Short title: Pollen galactose sensitivity via SWEET5 and GALK

Corresponding author: Li-Qing Chen

Tel: +1 217 265 9459

E-mail: lqchen77@illinois.edu

The hexose transporter SWEET5 confers galactose sensitivity to Arabidopsis pollen

via the galactokinase GALK

Jiang Wang[1,2], Ya-Chi Yu[1], Ye Li[1,3,4] and Li-Qing Chen[1,2*]

1. Department of Plant Biology, University of Illinois at Urbana-Champaign, Urbana, IL, 61801,

USA

2. Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign,

Urbana, IL, 61801, USA

3. Biotechnology Research Institute, Chinese Academy of Agricultural Sciences, 12

ZhongGuanCun South Street, Beijing, 100081, China

4. Current address: Key Laboratory of Agricultural Products Chemical and Biological Processing

Technology, Zhejiang University of Science and Technology, Hangzhou 310023, China

*Author for correspondence

Author Contributions

J.W. and L.-Q.C. conceived and designed the experiments. W.J., Y.-C.Y., and Y.L. conducted

experiments. J.W., and L.-Q.C wrote the manuscript.

One-sentence summary

SWEET5 mediates pollen galactose sensitivity via GALK that is required for efficient galactose

uptake in pollen during pollen germination.

Abstract

Galactose is an abundant and essential sugar used for the biosynthesis of many macromolecules in

different organisms, including plants. Galactose metabolism is tightly and finely controlled since

excess galactose and derivatives are inhibitory. In Arabidopsis, root growth and pollen germination

were strongly inhibited upon excess galactose. However, the mechanism of galactose induced

inhibition during pollen germination remains obscure. In this study, we characterized AtSWEET5

as a glucose and galactose plasma-membrane transporter localized in the pollen. SWEET5 protein

level start to accumulate since tricellular stage of pollen development and peaked in mature pollen

before rapidly declining after pollen was germinated. SWEET5 levels are responsible for the

dosage-dependent sensitivity of galactose and GALK is essential for the inhibitory effects of

galactose during pollen germination. The unexpected observation that GALK is required for

efficient galactose uptake in pollen may reveal an unknown regulatory mechanism for galactose

transporters. Overall, SWEET5 and GALK contribute to the maintenance of galactose metabolic

homeostasis during pollen germination, and galactose transport is positively regulated by GALK.

The study of SWEET5 upon galactose condition also suggests SWEET5 is a major low-affinity

hexose transporter at the early stage of pollen germination.

Introduction

Carbon flux, which is dependent on carbon supply and demand, needs to be well-controlled for plant growth and development (Hofmeyr and Cornish-Bowden, 2000; Bezrutczyk et al., 2018). Plants have developed sophisticated mechanisms to sense the sugar levels and adjust itself in response to developmental or environmental cues (Ruan, 2014). Plant sensitivity to various concentrations of sugar varies depending on the specific sugar involved. For example, in *Arabidopsis thaliana,* a low 10 mM concentration of galactose can trigger an inhibitory effect on Col-0 root growth (Yamada et al., 2011; Egert et al., 2012), while the much higher 167 mM (presented as 3%) concentration of glucose or 100 mM of sucrose still promote root growth (Kircher and Schopfer, 2012; Singh et al., 2014). Additionally, during *in vitro* pollen germination, 60 mM of galactose inhibits pollen germination (Hirsche et al., 2017), while 100 mM of glucose does not affect the pollen germination rate (Rottmann et al., 2018a) and high levels of sucrose (250-578 mM) are required for successful pollen germination (Wang et al., 2008; Rottmann et al., 2018a). However, the mechanism by which galactose inhibits plant growth has not been well-characterized. Sugar transporters and enzymes are key components in the supply-demand system to maintain carbon flux controlled (Ruan, 2014; Julius et al., 2017). The mutants of the two hexose transporter STPs (Sugar Transport Protein), AtSTP1 and AtSTP13, has shown a galactose-tolerant phenotype for root growth (Sherson et al., 2000, Yamada et al., 2011). The transporter that is involved in the process of galactose-suppressed pollen germination, however, remains unexplored.

STPs are important during early stage gametophyte development (Truernit et al. 1999) and pollen tube growth but not in the initial phase of pollen germination (Scholz-Starke et al. 2003, Buttner, 2007). Facilitator SWEETs (Sugar Will Eventually be Exported Transporter) have been shown to transport different sugars including galactose. OsSWEET5 (Zhou et al., 2014) and CsSWEET7a (Li et al., 2021) have been demonstrated to transport galactose. So far, all plant SWEETs with measured Km, e.g. AtSWEET1 (Chen *et al.*, 2010), AtSWEET12 (Chen *et al.*,2012), AtSWEET17 (Guo *et al.*, 2014), SlSWEET1a (Ho *et al.*, 2019) and CsSWEET7a (Li et al., 2021), have been found to function as low-affinity sugar transporters, ranging from ~10 to ~120 mM. AtSWEET5/VEX, belonging to SWEET family Clade II that primarily transports hexose (Chen et al., 2010; Eom et al., 2015), is strongly expressed in the pollen vegetative cell (Engel et al., 2005),

but no other details have been reported. We hypothesized that SWEET5 may be responsible for

the galactose-suppressed pollen germination.

Once galactose enters a cell, it is activated by galactokinase (GALK), a cytosolic enzyme, which

phosphorylates α-D-Gal into α-D-Gal-1-P at the C-1 position (Cardini and Leloir, 1953). Gal-1-P

can be further converted into UDP-Gal through a reversible reaction catalyzed either by Gal-1-P

uridylyltransferase (GALT) in the presence of UDP-Glc (mainly used in non-plant species) (Leloir,

1951) or by UDP-sugar pyrophosphorylase (USP) using UTP as a substrate (mainly found in plants)

(Feusi et al., 1999; Kotake et al., 2004). The activated UDP-Gal can be converted reversibly into

UDP-Glc by UDP-Glc epimerase (GALE/UGE) (Maxwell et al., 1960; Seifert et al., 2002). Only

a single copy of the GALK gene was reported to exist in Arabidopsis (Sherson et al., 1999; Yang

et al., 2009; Egert et al., 2012), and a galactose-insensitive phenotype in root growth was observed

in *galk* (Egert et al., 2012). As GALK is the first enzyme to catalyze galactose, we intended to

analyze its role together with SWEET5 transporter during pollen germination.

We show the plasma membrane localized SWEET5 transports both glucose and galactose. Data

from the current study indicates SWEET5 and GALK play critical roles in control galactose flux

during pollen germination. SWEET5 mediates galactose sensitivity in a GALK-dependent manner

during pollen germination and GALK positively regulates galactose uptake in the pollen, which

illuminates study of sugar flux control.

Results

SWEET5 transports galactose and contributes to galactose inhibition of Arabidopsis pollen

germination

To determine which SWEET in the mature pollen stage is able to transport galactose, we surveyed

the expression and protein accumulation of all SWEETs in Arabidopsis (Mergner et al., 2020).

Among all 17 members of the SWEET family, several were found to be expressed in mature pollen,

but only SWEET5, SWEET11 and SWEET12 proteins were detectable there (Figure 1A).

SWEET11 and SWEET12 have been reported to be transporters of sucrose (Chen et al., 2012),

while SWEET5 is the most likely transporter of galactose, since its homolog OsSWEET5 has been

found to transport galactose (Zhou et al., 2014). To test whether SWEET5 transports galactose,

we conducted growth assays with a yeast strain EBY.VW4000 lacking 17 hexose transporters

(Wieczorke et al., 1999). Yeast cells expressing SWEET5 grew on medium containing glucose

and galactose, but not fructose (Figure 1B). The K_m for galactose uptake of SWEET5 was

calculated as 671 ± 58.4 mM (Figure 1C), however, caution should be taken in interpreting this

K_m due to uncertain physiological effects on yeast when cultured with a high concentration (>500

mM). The pollen germination rate of Col-0 was dramatically reduced from 57% to 11% when

pollen germination medium (PGM) was supplemented with 60 mM Galactose (Figure 1D and 1E),

while remained unchanged when PGM was supplemented with either 60 mM glucose or fructose

(Supplemental Figure S1). The galactose suppression was barely detectable in *SWEET5* loss-of-

function mutants created from T-DNA insertion (*sweet5*) or CRISPR/CAS9 editing (Figure 1D

and 1E; Supplemental Figure S2). The phenotypes of the mutants were fully complemented by

transformation with a construct carrying the *SWEET5* promoter driving either endogenous

SWEET5 or a synonymous m*SWEET5* that carries mutations to avoid gRNA recognition in

CRISPR mutant lines (p*SWEET5*:*mSWEET5/sweet5_crispr*; Figure 1D and 1E; Supplemental

Figure S2D). Accordingly, the SWEET5 overexpression lines (p*SWEET5*:*SWEET5*/Col-0) were

more sensitive to galactose than Col-0 (Figure 1D and 1E). In pollen of *sweet5* mutant, there were

no observable morphological phenotypes under normal conditions, which may be due to functional

redundancy with other plasma membrane-localized hexose transporters (STPs, PMTs) detected at

either the RNA or protein level in pollen (Supplemental Figure S3). Among those hexose

transporters that were detected at the protein level, many can transport glucose and galactose

(Klepek et al., 2005; Klepek et al., 2010; Poschet et al., 2010; Rottmann et al., 2018b).

SWEET5 is mainly expressed at the late stages of pollen development

To determine the timing of the initiation of SWEET5-associated galactose inhibition. we examined

the spatial and temporal accumulation patterns of SWEET5 using transgenic lines harboring a

SWEET5 translational fusion with β-glucuronidase (GUS) or yellow fluorescent protein (YFP)

driven by its native SWEET5 promoter (p*SWEET5*:*gSWEET5-GUS* or p*SWEET5*:*gSWEET5-YFP*).

SWEET5 was almost exclusively found in the anthers and mature pollen at late stages of flower

development (Figure 2A), although GUS staining was detected at the early seedling stage with a

preference in the vein. The SWEET5 protein accumulation pattern was carefully assessed in the

flowers. Flowers at different developmental stages (Bowman, 1994) were imaged using
fluorescence microscopy to detect SWEET5 protein levels using the reporter YFP. As shown in
Figure 2C, SWEET5 started to accumulate in immature pollen grains from flower stage 11, in
which pollen mitosis I and II occur (Cecchetti et al., 2008) and peaked at flower stage 13 (anthesis).
The YFP signal was also observed over the course of male gametophyte development by isolating
developing pollen grains. The SWEET5-YFP protein was first detected at the tricellular pollen
(TCP) stage and peaked at the mature pollen grain (MPG) stage, while it was undetectable in both
uninucleate microspores (UNM) and bicellular pollen (BCP), as delineated by PI staining of the
cell wall and nucleus (Figure 2B). Additionally, SWEET5 protein level substantially reduced in
germinated pollen compared with non-germinated pollen grains after *in vitro* pollen germination
(Figure 2D). The subcellular localization of SWEET5 was determined using pollen from
transgenic plants harboring a SWEET5 (CDS) translational fusion with green fluorescent protein
(GFP) driven by its native SWEET5 promoter (p*SWEET5:cSWEET5-GFP*). SWEET5 is localized
to the plasma membrane (PM), and also endomembrane compartments as delineated by Nile red
staining for the lipid droplets (Figure 2E), which has been similarly observed for other plasma
membrane localized proteins in pollen (Hamilton et al., 2015). SWEET5 localization was further
transiently examined using the reporter mVenus in *N. benthamiana* leaves (Gookin and Assmann,
2014). SWEET5 protein was localized to the pavement cell periphery enclosing both chloroplast
and Golgi apparatus (Figure 2F), which suggests SWEET5 localized to the plasma membrane,
consistent with published Arabidopsis SWEETs except for SWEET2, SWEET16, and SWEET17
that are localized to the tonoplast (Klemens et al., 2013; Guo et al., 2014; Chen et al., 2015a).
161
Galactose sensitivity is GALK-dependent
To further elucidate whether GALK is involved in the galactose sensitivity during pollen
germination, we first surveyed whether GALK is accumulated in the pollen. The protein level of
GALK is relatively consistent across all tissues, including mature pollen grains (Supplemental
Figure S4A). Therefore, we tested pollen germination of the loss-of-function *galk* mutant (Egert
et al., 2012) on PGM containing galactose. The *galk* mutant showed a galactose-tolerant phenotype
similar to that of *sweet5* (Figure 3A; Supplemental Figure S5A), although in mature pollen it is a
knock-down mutant with 80% reduction in RNA transcripts (Supplemental Figure S4B and S4C).
The *galk* galactose-tolerant phenotype was fully complemented by *GALK* expressed under control

of the pollen-specific *LAT52* promoter (Figure 3A; Supplemental Figure S5A). Notably, the *galk*
complementation line showed a much stronger galactose sensitivity than Col-0, which is likely
due to the over-expression of GALK in the complementation line (Supplemental Figure S4C). As
expected, the GALK and SWEET5 double mutant *sweet5 × galk* showed a similar galactose-
tolerant phenotype to their individual single mutants, which indicates that GALK and SWEET5
function in the same pathway way. As Gal-1-P is suggested to exerts cellular toxicity from many
studies in animals (Lai et al., 2009), it is reasonable to speculate that pollen sensitivity to galactose
may be due to the accumulation of potential cytotoxic Gal-1-P, which depends on GALK activity
(de Jongh et al., 2008). To test this possibility, we compared the effects Gal-1-P on pollen
germination, 6 mM Gal-1-P was sufficient to mimic the suppression effect produced by 60 mM
galactose in Col-0 (Figure 3B; Supplemental Figure S5B). Gal-1-P can be converted into UDP-
Gal by USP in plants (Feusi et al., 1999). To test whether the USP-catalyzed reaction could
alleviate galactose suppression, we generated over-expression lines of AtUSP driven by the pollen-
specific *pLAT52* promoter in Col-0 background. However, USP overexpression lines failed to
improve the pollen germination rate under 60 mM galactose condition (Supplemental Figure S6).
186

Galactose inhibition is dosage-dependent

To determine the range of concentration at which galactose inhibits pollen germination, pollen
from different genotypes were germinated on PGM with galactose concentrations ranging from 0
to 600 mM. In Col-0, the pollen germination rate was not affected under 0.6 mM galactose, started
to decline significantly at 6 mM, followed by a substantial decrease at 60 mM and further reduced
to almost zero at 600 mM (Figure 4A, Supplemental Figure S5C). A time-course pollen
germination experiments of Col-0 further confirmed that the pollen germination rate was
significantly affected in the presence of 6 mM galactose starting from four hours after germination
(Supplemental Figure S5D). However, no significant difference was observed between Col-0 and
sweet5 mutants at 6 mM (Figure 5A, Supplemental Figure S5C). *sweet5* mutants were generally
tolerant to galactose up to 60 mM. Consistent with the observation, the SWEET5 overexpression
lines were more sensitive to ≥ 6 mM galactose than Col-0, with pollen germination rates reduced
to near zero by 60 mM galactose. These results suggest that SWEET5 is responsible for galactose
flux at the higher concentration range.

Galactose at 10 mM was reported to strongly inhibit root growth of Col-0 (medium containing no
sucrose, Yamada et al., 2011; Egert et al., 2012), while higher galactose (60 mM) is needed to
strongly inhibit Col-0 pollen germination (medium containing 15% (~ 439 mM) or 19.8 % (~578
mM) sucrose, Hirsche et al., 2017 or current study). To test whether the sucrose in the medium
may differentially affect observed galactose response, which may lead to biased comparison, we
conducted a galactose dosage-dependent assay for root growth using Col-0. Root growth was not
affected at 1 mM galactose but was strongly inhibited at 6 mM or higher without sucrose in the
medium (Supplemental Figure S7). By contrast, a significant root reduction was only observed at
60 mM galactose when 0.5% or 1% sucrose was supplemented in the medium. Therefore, the level
of inhibitory galactose concentration should be considered in the context of sucrose presence or
absence.
212
GALK positively regulates galactose uptake in pollen independent of SWEET5 expression
level
To assess the differences in galactose uptake among Col-0, *sweet5*, *galk*, and SWEET5
overexpression line, we conducted ^{14}C-galactose tracer uptake assays in pollen under various
galactose concentrations. Pollen grains were germinated in a liquid pollen germination system for
45 mins (the hydration period, Wang *et al*., 2008) to minimize the interference from transporters
expressed on pollen tubes. Less ^{14}C-galactose was taken up by *sweet5* pollen than by Col-0 pollen
at 60 mM galactose concentration; relative to the ~200 pmol galactose taken up by Col-0 pollen,
sweet5 utilized only ~50 pmol when 60 mM of galactose was externally supplied (Figure 4B).
Differences were not prominent at lower amounts of galactose (Figure 4B); however, the SWEET5
overexpression lines had significantly higher ^{14}C-galactose uptake than Col-0 under these
conditions (Figure 4C). Further, *sweet5* and *sweet5_crispr* pollen did not have significantly
reduced ^{14}C-galactose uptake under lower galactose conditions (Figure 4C), which is consistent
with involvement of other galactose transporters, that have high-affinity. Finally, ^{14}C-galactose
uptake was dramatically decreased in *galk* as well as *galk×sweet5* under all three tested galactose
conditions (Figure 4C), suggesting a positive feedback regulation of GALK on galactose
transporters, including SWEET5. The fact that the double *galk×sweet5* mutant phenocopies the
galk single mutant in both galactose uptake and *in vitro* pollen germination indicates that GALK
is epistatic to SWEET5 in controlling galactose uptake by pollen. However, the level of *SWEET5*

expression cannot explain the decrease of galactose uptake in *galk*, since expression of *SWEET5*

as well as other hexose transporters (STP 6 and 8) were significantly upregulated in *galk* (Figure

4D; Supplemental Figure S8).

Discussion

Pollen is a simple system to study the control of galactose flux

Over the past several decades, extensive efforts have been made to study galactose toxicity in

humans (galactosemia) and yeast; however, the mechanism underlying the toxicity is far from

understood (Lai et al., 2009). Considerably less attention has been paid to galactose inhibition by

plant scientists. In the current study, we found that the PM-localized hexose transporter

AtSWEET5 is responsible for galactose sensitivity during pollen germination. A high

concentration of sucrose at 19.8% is needed in the medium for in vitro pollen germination, which

may be partially cleaved into glucose and fructose by the Arabidopsis cell-wall invertase 2 before

imported into pollen (Hirsche et al., 2009). As SWEET5 also transports glucose (Figure 1B),

glucose may competitively suppress galactose uptake, resulting in much less galactose transported

in pollen. Thus, a high concentration of galactose, such as 60 mM, is needed to result in a severe

inhibitory effect on pollen germination. This concept was supported by the reduced galactose

response during root growth of wild type plant in presence of 0.5 % sucrose (Supplemental Figure

S7). Even when 60 mM external galactose was used, the estimated concentration was only about

135 µM in Col-0 pollen after incubation with 60 mM galactose for 45 min. This was calculated

based on the uptake of 200 pmol galactose (Figure 4B) determined from 20 flowers that yielded a

combined 40,000 pollen (2000 pollen per flower), and the mean pollen volume deduced from the

Arabidopsis pollen diameter of 20.72 µm (De Storme et al., 2013). While pollen has a fundamental

role in the sexual reproduction of flowering plants, pollen germination together with pollen tube

growth have served as a model system to study single cell growth and morphogenesis (Feijó et al.,

2004). In Arabidopsis, pollen grains are highly homogeneous and have strikingly low

transcriptional complexity (on average, 6044 genes expressed in mature pollen; Rutley and Twell,

2015) compared to sporophytic tissues or purified sporophytic cells (e.g. 13,222 genes expressed

in stomatal guard cells, and 11,696 genes expressed in root hair cells; Bates et al., 2012, Becker et

al., 2014). Our work to understand relationships among galactose transport, GALK, and galactose

sensitivity sheds light on developing alternative strategies to attenuate galactose suppression by
fine-tuning their relationships.

Model of galactose transport and metabolism during pollen *in vitro* germination

From the data presented in this study and reported from the publication, a schematic model of

galactose sensitivity in pollen was illustrated in Figure 5: SWEET5, as a uniporter, transports

galactose across pollen membrane across concentration gradient. If galactose is not sufficiently

transported into pollen grains due to a lack of enough SWEET5 protein, a galactose-tolerant

phenotype can be observed. Similarly, the lack of AtSTP1 and AtSTP13 have shown a galactose-

tolerant phenotype in conditions of up to 100 mM galactose (Sherson et al., 2000, Yamada et al.,

2011). Once galactose is taken up by cells, it will first be phosphorylated by GALK to produce the

potentially toxic Gal-1-P. A galactose detoxification pathway that stores excess galactose in the

vacuoles was proposed in the *galk* mutant (Egert et al., 2012). This pathway of detoxification

cannot explain the galactose-tolerant phenotype of *galk* pollen, because galactose uptake is

significantly decreased in *galk* pollen (Figure 4C) and vacuoles are nearly absent from mature

pollen grains (Pacini et al., 2011). Both decreased galactose uptake and the disrupted Gal-1-P

production may therefore be responsible for the galactose tolerance of the *galk* mutant during

pollen germination. Though possible free galactose accumulation by *galk* pollen, similar to *galk*

leaves (Egert et al., 2012), may negatively impact galactose influx during pollen germination and

obscure our understanding; the positive relationship between GALK and galactose uptake

activities suggests a yet-unknown mechanism associated with post-transcriptional and/or

translational regulation of PM-localized transporters.

Our study suggested that Gal-1-P is responsible for the galactose sensitivity observed during pollen

germination, which is supported by the results of Gal-1-P effect on *in vitro* pollen germination.

However, caution should be taken when interpreting these results, because it is still unclear how

Gal-1-P can be taken up by pollen as no such transporters have been identified. Gal-1-P showed a

significant increase as the external galactose concentration rises beyond the inhibitory levels in

corn and barley coleoptiles (Roberts et al., 1971). In animals or yeast, Gal-1-P has been reported

to potentially inhibit enzymes such as phosphoglucomutase (de Jongh et al., 2008), UDP-glucose

pyrophosphorylase (UGPase, catalyzing the reversible conversion from Glc-1-P to UDP-Glc) (Lai

et al., 2003), and inositol monophosphatase (Bhat, 2003) *in vitro,* however, no *in vivo* targets of

Gal-1-P have been identified. In limited studies in plants, Gal-1-P has only been reported to inhibit

UGPase activity in the crude extract from oat coleoptile, but not from the azuki bean epicotyl,

suggesting that galactose sensitivity appears to vary depending on the species (Yamamoto et al.,

1988). The double mutant of AtUGP1 and AtUGP2 exhibited dramatic growth defects and male

sterility (Park et al., 2010). Similarly, strong male sterility was also observed in the mutants that

lack USP to catalyzes the reversible conversion of monosaccharide-1-P and UDP-monosaccharide

(Schnurr et al., 2006; Kotake et al., 2007; Geserick and Tenhaken., 2013). However, the limited

production of UDP-Glc, UDP-GlcA, and UDP-Ara are not likely to be responsible for the *usp*

phenotype (Geserick and Tenhaken, 2013), since overexpressing UGPase for UDP-Glc synthesis

and overexpressing UDP-glucose dehydrogenase for UDP-GlcA synthesis have failed to rescue

the *usp* mutant (Geserick and Tenhaken, 2013), and the mutant of UDP-xylose-4-epimerse, which

contributes mostly to UDP-Ara production, does not differ from the wild type (Burget et al., 2003).

Therefore, the conversion between Gal-1-P and UDP-Gal might be important for the *usp*

phenotype. We tested this possibility by overexpressing USP; however, it failed to alleviate

galactose suppression during pollen germination. The pool of UDP-Gal may result not only from

the reaction catalyzed by USP, but also the reversible conversion of UDP-Gal and UDP-Glc

catalyzed by the epimerase of UGE. The lack of both UGE2 and UGE3 is responsible for a male

sterile phenotype resembling the *usp* phenotype (Rosti et al., 2007). Additionally, low endogenous

UGE1 activity has been suggested to be responsible for the more severe galactose inhibition, along

with an increase in the UDP-Gal content (Dormann and Benning, 1998). Thus, the dynamically

quantitative analysis of metabolites, including Gal-1-P, Glc-1-P, UDP-Gal and UDP-Glc, may help

to elucidate the role of USP or UGE in galactose sensitivity. We attempted to measure these

galactose-associated metabolites upon galactose treatment in *sweet5*, however, no clear

conclusions can be drawn due to the difficulties of separating glucose and galactose conjugated

phosphates and nucleotide sugars using a LC-MS/MS system.

SWEET5 orthologs may play important roles in hexose-utilizing pollen

Compared to the proton sugar symporters, SWEET5 likely functions as a uniporter, facilitating

substrate movement along a substrate concentration gradient (Chen et al., 2015c). The

physiological importance of SWEET family members was underscored mainly by their

involvement in multiple processes requiring sugar transport capability, for instances, SWEET11 and SWEET12 in apoplasmic phloem loading (Chen *et al*., 2012); SWEET9 in nectar secretion (Lin *et al*., 2014); and SWEET11, SWEET12, and SWEET15 in seed filling (Chen *et al*., 2015b). In the current work, we showed that SWEET5 is highly expressed in the mature pollen but dramatic declines during the transition from germinated pollen to pollen tubes, suggesting that SWEET5 functions during the early stages of pollen germination. By contrast, several STPs are primarily expressed in pollen tubes but are almost absent in mature pollen grains (Rottmann et al., 2018a). For examples, STP10 is a pollen tube-specific hexose transporter (Rottmann et al., 2016), the glucose transporter STP9 is weakly expressed in mature pollen grains and is primarily expressed in growing pollen tubes (Schneidereit et al., 2003). Therefore, SWEET5 and STPs may have complementary roles during pollen germination. As reported, high concentrations of sugars (ranging from 2% to 19.8%) are needed for successful germination *in vitro* in various species (Ylstra et al., 1998; Stadler et al., 1999; Wang et al., 2008) since pollen germination is an energy-demanding process (Reinders, 2016). However, the effect of different sugars on *in vitro* pollen germination depends on the species. For example, sucrose and hexose differentially affect *in vitro* pollen germination, ranging from growing equally well on glucose, fructose, or sucrose for petunia (Ylstra et al., 1998), cucumber (Cheng et al., 2015), and pearl millet (Reger et al., 1992), to a nearly complete inhibition by glucose or fructose for Arabidopsis (Hirsche et al., 2017). But for some species, such as tobacco, either sucrose or glucose can support pollen germination, while only sucrose can promote pollen tube growth (Goetz et al., 2017). Galactose alone can also promote pollen germination for example in *Picea wilsonii* (Zhou et al., 2020) and cucumber (Cheng et al., 2015) to a similar extent as sucrose. Therefore, the lack of a phenotype in *sweet5* under normal conditions may be due to the preference of Arabidopsis pollen during the initial phase of germination for sucrose. Considering SWEET5 also transports glucose, it is rational to speculate that SWEET5 orthologues, in the species that prefer to use hexose as a carbon source during pollen germination or pollen tube growth, may be essential for pollen development, and therefore seed production.

Conclusions

We have presented SWEET5 as a PM-localized hexose transporter that is highly expressed in

mature pollen grains. SWEET5 in synergy with GALK is responsible for the galactose sensitivity

observed during pollen germination in Arabidopsis. Pollen germination studies served as a tool to

elucidate SWEET5 biological functions and provide a model system that facilitates research to

uncover mechanisms underlying galactose sensitivity. The future comparison of the effect of

galactose or glucose on pollen germination, associated with either SWEET5 or its homologs from

species that favor different carbon sources for pollen germination, will offer new insights into the

carbon skeleton and/or energy supply mechanisms responsible for successful reproduction.

Materials and Methods

Constructs for yeast expression

366 The CDSs of *AtSWEET5* were amplified by RT–PCR using gene specific primers named by gene

367 (Supplemental Table S2, P1 and P2) and were subsequently cloned into the donor vector

368 pDONR221-f1 by BP clonase (Invitrogen) before transfer to pDRf1-GW (Loque et al., 2007) by

369 LR clonase (Invitrogen).

GUS and YFP fusion constructs under native promoters

To construct *SWEET5* fused with GUS (p*SWEET5*:g*SWEET5-GUS*), the fragment including the

SWEET5 native promoter sequence (1801 bp upstream of ATG) and genomic sequence of

SWEET5 was amplified (P3 and P4) and cloned into pDONR221-f1 by BP reactions followed by

recombination into the vector pMDC163 carrying the GUS gene (Curtis and Grossniklaus, 2003)

by LR reaction. For the SWEET5 overexpression construct (p*SWEET5*:g*SWEET5-YFP*), the entry

clone from above was transferred into pEarleyGateTW1 (Wang et al., 2016) by LR reactions.

eGFP fusion constructs under different promoters for strength comparisons and mutant complementation

The *SWEET5* (1216 bp upstream of ATG) native promoter, and *LAT52* promoter (602 bp;

Muschietti et al., 1994) were amplified with specific forward primers (P5-P8) containing a *BamH*I

restriction site and the reverse primers containing an *Xba*I restriction site and sub-cloned into the

eGFP-containing vector pGTKan3 (Kasaras and Kunze, 2010) via *BamH*I and *Xba*I restriction

sites. The CDS of *SWEET5, GALK,* and *USP* by their own specific primers (P9-P14) were

seamlessly subcloned into corresponding pGTKan3_promoter constructs by In-Fusion® (Takara)

following the linearization at *Xba*I and *Pst*I restriction sites. For the genome editing construct of

SWEET5, gRNA that targets the first exon of SWEET5 (Supplemental Figure S1D) was cloned

into p*YAO*:*hSpCas9* (Yan et al., 2015). For the synonymous complementation construct of Cas9

containing *sweet5_crispr,* synonymous *SWEET5* driven by pSWEET5 was amplified by primer

P15 and P16 based on pDONR221_pSWEET5-*SWEET5* with gRNA targeting the region modified

through site-directed mutagenesis before transferring into pEarleyGate301 (Earley et al., 2006) by

LR reactions.

Construct for subcellular localization

For subcellular localization of SWEET5, after the vector pDOE-13 (Gookin and Assmann, 2014)

was linearized by *SanD*I + *Aat*I restriction enzymes at MCS3, the CDS of SWEET5 was amplified

by primers P17 and P18 and subcloned into it by In-Fusion® (Takara). The agroinfiltration-based

assays were performed as previously described (Gookin and Assmann, 2014). The *N. benthamiana*

leaf disks were collected 72 h post infiltration for confocal imaging. Three independent trials were

conducted.

Plant materials and growth conditions

The Arabidopsis Col-0 plants were grown under controlled temperature (22°C) with a 16-h light

(100-150 $\mu Em^{-2}s^{-1}$)/ 8-h dark photoperiod. The flowers at stage 13 (fully opened) were collected

at around 5 hours into the light period for different experiments. T-DNA mutants of *galk* (GABI-

Kat 489D10), and *sweet5* (CS853155) were obtained from NASC or ABRC. Homozygous lines

were genotyped using primers of P19-P24 and used in related experiments. The floral dip method

(Clough and Bent, 1998) was used to generate all the transgenic lines used in this study. At least

15 T_1 lines were generated for each construct and at least 3 randomly selected lines were

propagated to generate T_3 homozygous seeds.

Yeast complementation assay and ^{14}C uptake assay

Yeast complementation and yeast ^{14}C uptake assays were conducted following the previously

described method (Chen et al., 2010). ^{14}C-galactose (0.5 μCi D-[1-^{14}C] galactose; 56.2 mCi/mmol)

was added per sample and four independent transformants were used for each uptake experiment.

For concentration- and time-dependent uptake of [14C] galactose for SWEET5, 5 galactose

concentrations (25, 50, 100, 250, 500 mM), and 4 time points (10 sec, 2.5, 5, and 10 min) for each

concentration were used to collect yeast cell samples and measure galactose uptake. Sugar uptake

at 10 sec was normalized as 0.

***In vitro* pollen germination**

The *in vitro* pollen germination assay was conducted according to a previously described method

(Wang et al., 2008) with minor modifications. The sucrose only basic pollen germination medium

(PGM) was composed of 19.8% (w/v) Suc (15% (w/v) Suc in liquid PGM), 1.5 mM boric acid,

0.8 mM $MgSO_4$, 1 mM KCl, 5 mM MES, 0.05% (w/v) lactalbumin hydrolysate, 10 µM myo-

inositol, 5 mM $CaCl_2$ and 1.5% (w/v) agarose. The pH was adjusted to 5.8 using 1 mM Tris (pH

= 8). The medium solution was heated to boiling on a heat plate, and then 500 µl was spread evenly

onto 75 × 25 mm glass slides. Slides were cooled down at room temperature before placing in an

opaque, sealed slide box. The medium was made freshly for each germination experiment. Six

flowers at stage 13 were collected from each genotype to spread pollen gently onto a ~1 cm^2 area

of the germination medium. The slide box was kept at room temperature for 8 hours in the dark

before samples on slides were imaged with a compound microscope. Pollen with pollen tubes

longer than the diameter of the pollen grain (about 20 µm) was considered as germinated pollen.

Pollen galactose uptake

The liquid pollen germination method was adopted from a previously described method (Wang et

al., 2008) with minor modifications for ^{14}C-galactose uptake. For each trial, 20 flowers (at least

2000 pollen grains collected per flower) at stage 13 for each genotype were collected into a 2 ml

tube, and then 1 ml of liquid PGM was added in and vortexed vigorously for 60 s to release mature

pollen grains into the solution. Subsequently, the pollen grains were counted using a

hemocytometer. Equal amounts of pollen for all tested genotypes were collected by centrifuging

at 15,000 g for 1 min. The pollen pellet was resuspended in 30 µl of liquid PGM containing ^{14}C-

galactose (0.3 µCi hot galactose) as well as different concentrations of cold galactose (ranging

from 0 to 60 mM), and subsequently cultured in a petri dish (35 mm in diameter). The petri dish

was covered and placed in the dark for 45 min before pollen was washed using 1 ml of ice-cold

liquid PGM and collected into a 1.5 ml tube. The pollen was precipitated at 15,000 g for 1 min and

the pellet was washed by 1 ml ice-cold pollen isolation buffer (PIB, composed of 100 mM NaPO$_4$,

pH 7.5, 1 mM EDTA, and 0.1% (v/v) Triton X-100) 5 more times. The pellet was resuspended in

100 µl PIB and transferred to a scintillation vial containing 5 ml Ultima Gold XR Scintillation

liquid (Perkinelmer). Radioactivity for each vial was measured by a liquid scintillation

spectrometry.

GUS histochemical analysis

GUS staining was performed as previously described (Chen et al., 2012). Twelve-day old seedlings

were used, and inflorescences and siliques were collected for histochemical GUS staining.

Microscopy imaging

A Zeiss Apotome.2 (Carl Zeiss, Thornwood, NY, USA) was used for fluorescence acquisition.

Different FL filter sets (YFP, CFP, RFP and GFP) were used to image samples as needed. A Zeiss

LSM 710 confocal microscope (Carl Zeiss, Thornwood, NY, USA) was used to image samples.

Argon laser excitation wavelength and emission bandwidth were 458 nm and 480–520 nm for

mTq2 (cyan), 488 nm and 500–550 nm for mVenus (yellow), and 633 nm and 633-740 nm for

chlorophyll autofluorescence (red) respectively. Image acquisition parameters were held

consistent. Raw data from each channel were not altered beyond equal signal increases.

RNA isolation and RT-qPCR

RNA isolation from pollen was performed as previously described with modifications (Lu, 2011).

For each independent sample, pollen grains from 50 flowers at stage 13 were collected. For

samples under galactose treatment, pollen grains from 50 flowers were germinated in liquid PGM

for 45 mins as described above in pollen galactose uptake. RNA was isolated using Trizol

(Invitrogen) as instructed by the manufacturer. First-strand cDNA was synthesized using oligo(dT)

and M-MuLV reverse transcriptase (NEB). Real-time qPCR was performed using PowerUp™

SYBR™ master mix (Applied Biosystems) according to the manufacturer's instructions on a

CFX96 Real-Time PCR Detection System (Bio-Rad) using gene specific primers (P25-P34). The

expression values were normalized to *ACTIN8* expression values in each repeat, and subsequently

normalized to Col-0 using $2^{-\Delta\Delta CT}$ method (Livak and Schmittgen, 2001).

Statistical analysis

The differences between the two subjects were determined using the two-tailed Student's t-test with equal variance. The differences among multiple subjects were assessed using one-way ANOVA followed by multiple comparison tests (Fisher's LSD method). All statistical analysis was performed using SPSS 26 statistical software (SPSS Inc, Chicago, IL, USA).

Accession numbers

Sequence information from this article can be found in the Arabidopsis Genome Initiative or GenBank/EMBL databases under the following accession numbers: SWEET5 (AT5G62850), GALK (AT3G06580), USP (AT5G52560).

Acknowledgments

We thank Dr. Yaxin Li for help with sample collection and thank Dr. Qi Xie for providing us with p*YAO:hSpCas9* vector. We also thank Mrs. Anita K. Snyder for editing. This work is supported by startup funds from the University of Illinois at Urbana-Champaign to Dr. Chen (to J.W., Y.-C. Y., and L.-Q.C). Y. L. was supported by China Scholarship Council (No. 201603250070).

References

Bates G, Rosenthal D, Sun J, Chattopadhyay M, Peffer E, Yang J, Ort D, Jones A (2012) A comparative study of the *Arabidopsis thaliana* guard-cell transcriptome and its modulation by sucrose. Plos One **7**

Becker J, Takeda S, Borges F, Dolan L, Feijo J (2014) Transcriptional profiling of Arabidopsis root hairs and pollen defines an apical cell growth signature. BMC Plant Biology **14**

Bezrutczyk M, Yang J, Eom J, Prior M, Sosso D, Hartwig T, Szurek B, Oliva R, Vera-Cruz C, White F, Yang B, Frommer W (2018) Sugar flux and signaling in plant-microbe interactions. Plant Journal **93:** 675-685

Bhat P (2003) Galactose-1-phosphate is a regulator of inositol monophosphatase: a fact or a fiction? Medical Hypotheses **60:** 123-128

Bowman J (1994) Arabidopsis: an atlas of morphology and development. Springer-Verlag New

York

Burget E, Verma R, Molhoj M, Reiter W (2003) The biosynthesis of L-arabinose in plants: molecular cloning and characterization of a golgi-localized UDP-D-xylose 4-epimerase encoded by the MUR4 gene of Arabidopsis. Plant Cell **15:** 523-531

Buttner M (2007) The monosaccharide transporter(-like) gene family in Arabidopsis. Febs Letters **581:** 2318-2324

Cardini C, Leloir L (1953) Enzymic phosphorylation of galactosamine and galactose. Archives of Biochemistry and Biophysics **45:** 55-64

Cecchetti V, Altamura M, Falasca G, Costantino P, Cardarelli M (2008) Auxin regulates Arabidopsis anther dehiscence, pollen maturation, and filament elongation. Plant Cell **20:** 1760-1774

Chen H, Huh J, Yu Y, Ho L, Chen L, Tholl D, Frommer W, Guo W (2015) The Arabidopsis vacuolar sugar transporter SWEET2 limits carbon sequestration from roots and restricts Pythium infection. Plant Journal **83:** 1046-1058

Chen L, Cheung L, Feng L, Tanner W, Frommer W, Kornberg R (2015) Transport of sugars. Annual Review of Biochemistry, Vol 84 **84:** 865-894

Chen L, Hou B, Lalonde S, Takanaga H, Hartung M, Qu X, Guo W, Kim J, Underwood W, Chaudhuri B, Chermak D, Antony G, White F, Somerville S, Mudgett M, Frommer W (2010) Sugar transporters for intercellular exchange and nutrition of pathogens. Nature **468:** 527-U199

Chen L, Lin I, Qu X, Sosso D, McFarlane H, Londono A, Samuels A, Frommer W (2015) A cascade of sequentially expressed sucrose transporters in the seed coat and endosperm provides nutrition for the Arabidopsis embryo. Plant Cell **27:** 607-619

Chen L, Qu X, Hou B, Sosso D, Osorio S, Fernie A, Frommer W (2012) Sucrose efflux mediated by SWEET proteins as a key step for phloem transport. Science **335:** 207-211

Cheng J, Wang Z, Yao F, Gao L, Ma S, Sui X, Zhang Z (2015) Down-regulating CsHT1, a cucumber pollen-specific hexose transporter, inhibits pollen germination, tube growth, and seed development. Plant Physiology **168:** 635-647

Clough S, Bent A (1998) Floral dip: a simplified method for Agrobacterium-mediated transformation of *Arabidopsis thaliana*. Plant Journal **16:** 735-743

Curtis M, Grossniklaus U (2003) A gateway cloning vector set for high-throughput functional

analysis of genes in planta. Plant Physiology **133:** 462-469

de Jongh W, Bro C, Ostergaard S, Regenberg B, Olsson L, Nielsen J (2008) The roles of galactitol, galactose-1-phosphate, and phosphoglucomutase in galactose-induced toxicity in *Saccharomyces cerevisiae*. Biotechnology and Bioengineering **101:** 317-326

De Storme N, Zamariola L, Mau M, Sharbel T, Geelen D (2013) Volume-based pollen size analysis: an advanced method to assess somatic and gametophytic ploidy in flowering plants. Plant Reproduction **26:** 65-81

Dormann P, Benning C (1998) The role of UDP-glucose epimerase in carbohydrate metabolism of Arabidopsis. Plant Journal **13:** 641-652

Earley K, Haag J, Pontes O, Opper K, Juehne T, Song K, Pikaard C (2006) Gateway-compatible vectors for plant functional genomics and proteomics. Plant Journal **45:** 616-629

Egert A, Peters S, Guyot C, Stieger B, Keller F (2012) An Arabidopsis T-DNA insertion mutant for galactokinase (AtGALK, At3g06580) hyperaccumulates free galactose and is insensitive to exogenous galactose. Plant and Cell Physiology **53:** 921-929

Engel M, Davis R, McCormick S (2005) Green sperm. Identification of male gamete promoters in Arabidopsis. Plant Physiology **138:** 2124-2133

Eom JS, Chen LQ, Sosso D, Julius BT, Lin IW, Qu XQ, Braun DM, Frommer WB (2015) SWEETs, transporters for intracellular and intercellular sugar translocation. Current Opinion in Plant Biology **25:** 53-62

Feijo J, Costa S, Prado A, Becker J, Certal A (2004) Signalling by tips. Current Opinion in Plant Biology **7:** 589-598

Feusi MES, Burton JD, Williamson JD, Pharr DM (1999) Galactosyl-sucrose metabolism and UDP-galactose pyrophosphorylase from *Cucumis melo* L-fruit. Physiologia Plantarum **106:** 9-16

Geserick C, Tenhaken R (2013) UDP-sugar pyrophosphorylase is essential for arabinose and xylose recycling, and is required during vegetative and reproductive growth in Arabidopsis. Plant Journal **74:** 239-247

Goetz M, Guivarch A, Hirsche J, Bauerfeind M, Gonzalez M, Hyun T, Eom S, Chriqui D, Engelke T, Grosskinsky D, Roitsch T (2017) Metabolic control of tobacco pollination by sugars and invertases. Plant Physiology **173:** 984-997

571   **Gookin T, Assmann S** (2014) Significant reduction of BiFC non-specific assembly facilitates in
572   planta assessment of heterotrimeric G-protein interactors. Plant Journal **80:** 553-567

573   **Guo W, Nagy R, Chen H, Pfrunder S, Yu Y, Santelia D, Frommer W, Martinoia E** (2014)
574   SWEET17, a facilitative transporter, mediates fructose transport across the tonoplast of
575   Arabidopsis roots and leaves. Plant Physiology **164:** 777-789

576   **Hamilton E, Jensen G, Maksaev G, Katims A, Sherp A, Haswell E** (2015) Mechanosensitive
577   channel MSL8 regulates osmotic forces during pollen hydration and germination. Science
578   **350:** 438-441

579   **Hirsche J, Engelke T, Voller D, Gotz M, Roitsch T** (2009) Interspecies compatibility of the
580   anther specific cell wall invertase promoters from Arabidopsis and tobacco for generating
581   male sterile plants. Theoretical and Applied Genetics **118:** 235-245

582   **Ho LH, Klemens PAW, Neuhaus HE, Ko HY, Hsieh SY, Guo WJ** (2019) SlSWEET1a is
583   involved in glucose import to young leaves in tomato plants. Journal of Experimental
584   Botany **70:** 3241-3254

585   **Hofmeyr J, Cornish-Bowden A** (2000) Regulating the cellular economy of supply and demand.
586   Febs Letters **476:** 47-51

587   **Julius B, Leach K, Tran T, Mertz R, Braun D** (2017) Sugar Transporters in Plants: New Insights
588   and Discoveries. Plant and Cell Physiology **58:** 1442-1460

589   **Kasaras A, Kunze R** (2010) Expression, localisation and phylogeny of a novel family of plant-
590   specific membrane proteins. Plant Biology **12:** 140-152

591   **Kircher S, Schopfer P** (2012) Photosynthetic sucrose acts as cotyledon-derived long-distance
592   signal to control root growth during early seedling development in Arabidopsis.
593   Proceedings of the National Academy of Sciences of the United States of America **109:**
594   11217-11221

595   **Klemens P, Patzke K, Deitmer J, Spinner L, Le Hir R, Bellini C, Bedu M, Chardon F, Krapp**
596   **A, Neuhaus H** (2013) Overexpression of the vacuolar sugar carrier AtSWEET16 modifies
597   germination, growth, and stress tolerance in Arabidopsis. Plant Physiology **163:** 1338-1352

598   **Klepek Y, Geiger D, Stadler R, Klebl F, Landouar-Arsivaud L, Lemoine R, Hedrich R, Sauer**
599   **N** (2005) Arabidopsis polyol transporter 5, a new member of the monosaccharide
600   transporter-like superfamily, mediates H^+-symport of numerous substrates, including myo-
601   inositol, glycerol, and ribosele. Plant Cell **17:** 204-218

Klepek Y, Volke M, Konrad K, Wippel K, Hoth S, Hedrich R, Sauer N (2010) *Arabidopsis*
thaliana polyol/mnosaccharide transporters 1 and 2: fructose and xylitol/H$^+$ symporters in
pollen and young xylem cells. Journal of Experimental Botany **61:** 537-550

Kotake T, Hojo S, Yamaguchi D, Aohara T, Konishi T, Tsumuraya Y (2007) Properties and
physiological functions of UDP-sugar pyrophosphorylase in Arabidopsis. Bioscience
Biotechnology and Biochemistry **71:** 761-771

Kotake T, Yamaguchi D, Ohzono H, Hojo S, Kaneko S, Ishida H, Tsumuraya Y (2004) UDP-
sugar pyrophosphorylase with broad substrate specificity toward various monosaccharide
1-phosphates from pea sprouts. Journal of Biological Chemistry **279:** 45728-45736

Lai K, Elsas L, Wierenga K (2009) Galactose toxicity in animals. Iubmb Life **61:** 1063-1074

Lai K, Langley S, Khwaja F, Schmitt E, Elsas L (2003) GALT deficiency causes UDP-hexose
deficit in human galactosemic cells. Glycobiology **13:** 285-294

Leloir L (1951) The enzymatic transformation of uridine diphosphate glucose into a galactose
derivative. Archives of Biochemistry and Biophysics **33:** 186-190

Li Y, Liu H, Yao X, Wang J, Feng S, Sun L, Ma S, Xu K, Chen L-Q, Sui X (2021) Hexose
transporter CsSWEET7a in cucumber mediates phloem unloading in companion cells for
fruit development. Plant Physiology

Lin I, Sosso D, Chen L, Gase K, Kim S, Kessler D, Klinkenberg P, Gorder M, Hou B, Qu X,
Carter C, Baldwin I, Frommer W (2014) Nectar secretion requires sucrose phosphate
synthases and the sugar transporter SWEET9. Nature **508:** 546-549

Livak K, Schmittgen T (2001) Analysis of relative gene expression data using real-time
quantitative PCR and the 2(T)(-Delta Delta C) method. Methods **25:** 402-408

Loque D, Lalonde S, Looger LL, von Wiren N, Frommer WB (2007) A cytosolic trans-
activation domain essential for ammonium uptake. Nature **446:** 195-198

Lu Y (2011) RNA isolation from Arabidopsis pollen grains. Bio-protocol **1:** e67

Maxwell E, Derobichonszulmajster H (1960) Purification of uridine diphosphate galactose-4-
epimerase from yeast, and the identification of protein-bound diphosphopyridine
nucleotide. Journal of Biological Chemistry **235:** 308-312

Mergner J, Frejno M, List M, Papacek M, Chen X, Chaudhary A, Samaras P, Richter S,
Shikata H, Messerer M, Lang D, Altmann S, Cyprys P, Zolg D, Mathieson T,
Bantscheff M, Hazarika R, Schmidt T, Dawid C, Dunkel A, Hofmann T, Sprunck S,

Falter-Braun P, Johannes F, Mayer K, Jurgens G, Wilhelm M, Baumbach J, Grill E, Schneitz K, Schwechheimer C, Kuster B (2020) Mass-spectrometry-based draft of the Arabidopsis proteome. Nature **579:** 409-414

Muschietti J, Dircks L, Vancanneyt G, Mccormick S (1994) LAT52 protein is essential for tomato pollen development - pollen expressing antisense LAT52 RNA hydrates and germinates abnormally and cannot achieve fertilization. Plant Journal **6:** 321-338

Pacini E, Jacquard C, Clement C (2011) Pollen vacuoles and their significance. Planta **234:** 217-227

Park J, Ishimizu T, Suwabe K, Sudo K, Masuko H, Hakozaki H, Nou I, Suzuki G, Watanabe M (2010) UDP-glucose pyrophosphorylase is rate limiting in vegetative and reproductive phases in *Arabidopsis thaliana*. Plant and Cell Physiology **51:** 981-996

Poschet G, Hannich B, Buttner M (2010) Identification and characterization of AtSTP14, a novel galactose transporter from Arabidopsis. Plant and Cell Physiology **51:** 1571-1580

Reger B, Pressey R, Chaubal R (1992) In vitro chemotropism of pearl-millet pollen tubes to stigma tissue - a response to glucose produced in the medium by tissue-bound invertase. Sexual Plant Reproduction **5:** 201-205

Reinders A (2016) Fuel for the road - sugar transport and pollen tube growth. Journal of Experimental Botany **67:** 2121-2123

Roberts R, Heishman A, Wicklin C (1971) Growth inhibition and metabolite pool levels in plant tissues fed D-glucosamine and D-galactose. Plant Physiology **48:** 36-42

Rosti J, Barton C, Albrecht S, Dupree P, Pauly M, Findlay K, Roberts K, Seifert G (2007) UDP-glucose 4-epimerase isoforms UGE2 and UGE4 cooperate in providing UDP-galactose for cell wall biosynthesis and growth of *Arabidopsis thaliana*. Plant Cell **19:** 1565-1579

Rottmann T, Fritz C, Sauer N, Stadler R (2018) Glucose uptake via STP transporters inhibits in vitro pollen tube growth in a HEXOKINASE1-dependent manner in *Arabidopsis thaliana*. Plant Cell **30:** 2057-2081

Rottmann T, Klebl F, Schneider S, Kischka D, Ruscher D, Sauer N, Stadler R (2018) Sugar transporter STP7 specificity for L-arabinose and D-xylose contrasts with the typical hexose transporters STP8 and STP12. Plant Physiology **176:** 2330-2350

Rottmann T, Zierer W, Subert C, Sauer N, Stadler R (2016) STP10 encodes a high-affinity

monosaccharide transporter and is induced under low-glucose conditions in pollen tubes

of Arabidopsis. Journal of Experimental Botany **67:** 2387-2399

Ruan Y, Merchant S (2014) Sucrose Metabolism: Gateway to Diverse Carbon Use and Sugar

Signaling. Annual Review of Plant Biology, Vol 65 **65:** 33-67

Rutley N, Twell D (2015) A decade of pollen transcriptomics. Plant Reproduction **28:** 73-89

Schneidereit A, Scholz-Starke J, Buttner M (2003) Functional characterization and expression

analyses of the glucose-specific AtSTP9 monosaccharide transporter in pollen of

Arabidopsis. Plant Physiology **133:** 182-190

Schnurr J, Storey K, Jung H, Somers D, Gronwald J (2006) UDP-sugar pyrophosphorylase is

essential for pollen development in Arabidopsis. Planta **224:** 520-532

Scholz-Starke J, Buttner M, Sauer N (2003) AtSTP6, a new pollen-specific H$^+$-monosaccharide

symporter from Arabidopsis. Plant Physiology **131:** 70-77

Seifert G, Barber C, Wells B, Dolan L, Roberts K (2002) Galactose biosynthesis in Arabidopsis:

genetic evidence for substrate channeling from UDP-D-galactose into cell wall polymers.

Current Biology **12:** 1840-1845

Sherson S, Gy I, Medd J, Schmidt R, Dean C, Kreis M, Lecharny A, Cobbett C (1999) The

arabinose kinase, ARA1, gene of Arabidopsis is a novel member of the galactose kinase

gene family. Plant Molecular Biology **39:** 1003-1012

Sherson S, Hemmann G, Wallace G, Forbes S, Germain V, Stadler R, Bechtold N, Sauer N,

Smith S (2000) Monosaccharide/proton symporter AtSTP1 plays a major role in uptake

and response of Arabidopsis seeds and seedlings to sugars. Plant Journal **24:** 849-857

Singh M, Gupta A, Laxmi A (2014) Glucose control of root growth direction in Arabidopsis

thaliana. Journal of Experimental Botany **65:** 2981-2993

Stadler R, Truernit E, Gahrtz M, Sauer N (1999) The AtSUC1 sucrose carrier may represent

the osmotic driving force for anther dehiscence and pollen tube growth in Arabidopsis.

Plant Journal **19:** 269-278

Truernit E, Stadler R, Baier K, Sauer N (1999) A male gametophyte-specific monosaccharide

transporter in Arabidopsis. Plant Journal **17:** 191-201

Wang R, Liu M, Yuan M, Oses-Prieto J, Cai X, Sun Y, Burlingame A, Wang Z, Tang W

(2016) The brassinosteroid-activated BRI1 receptor kinase is switched off by

dephosphorylation mediated by cytoplasm-localized PP2A B ' subunits. Molecular Plant **9:**

148-157

Wang Y, Zhang W, Song L, Zou J, Su Z, Wu W (2008) Transcriptome analyses show changes
in gene expression to accompany pollen germination and tube growth in Arabidopsis. Plant
Physiology **148:** 1201-1211

Wieczorke R, Krampe S, Weierstall T, Freidel K, Hollenberg C, Boles E (1999) Concurrent
knock-out of at least 20 transporter genes is required to block uptake of hexoses in
Saccharomyces cerevisiae. FEBS Letters **464:** 123-128

Yamada K, Kanai M, Osakabe Y, Ohiraki H, Shinozaki K, Yamaguchi-Shinozaki K (2011)
Monosaccharide absorption activity of Arabidopsis roots depends on expression profiles
of transporter genes under high salinity conditions. Journal of Biological Chemistry **286:**
43577-43586

Yamamoto R, Inouhe M, Masuda Y (1988) Galactose inhibition of auxin-induced growth of
monocotyledonous and dicotyledonous plants. Plant Physiology **86:** 1223-1227

Yan L, Wei S, Wu Y, Hu R, Li H, Yang W, Xie Q (2015) High-efficiency genome editing in
Arabidopsis using YAO promoter-driven CRISPR/Cas9 system. Molecular Plant **8:** 1820-
1823

Yang T, Bar-Peled L, Gebhart L, Lee S, Bar-Peled M (2009) Identification of galacturonic
acid-1-phosphate kinase, a new member of the GHMP kinase duperfamily in plants, and
comparison with galactose-1-phosphate kinase. Journal of Biological Chemistry **284:**
21526-21535

Ylstra B, Garrido D, Busscher J, van Tunen A (1998) Hexose transport in growing petunia
pollen tubes and characterization of a pollen-specific, putative monosaccharide transporter.
Plant Physiology **118:** 297-304

Zhou Y, Cui X, Hu A, Miao Y, Zhang L (2020) Characterization and functional analysis of
pollen-specific PwSWEET1 in *Picea wilsonii*. Journal of Forestry Research **31:** 1913-1922

Zhou Y, Liu L, Huang W, Yuan M, Zhou F, Li X, Lin Y (2014) Overexpression of OsSWEET5
in rice causes growth retardation and precocious senescence. Plos One **9:** e94210

Figure 1. Characterization of Arabidopsis SWEET5.

(A) 17 SWEET transcripts and protein levels in Arabidopsis mature pollen. Data were retrieved

from athena.proteomics.wzw.tum.de.

(B) Functional analysis of SWEET5 by EBY.VW4000 yeast complementation assay. The empty

vector (EV) as a negative control, or HXT5 as a positive control.

(C) Concentration-dependent uptake of ^{14}C-Galactose over time by yeast transformed with

SWEET5 or an empty vector as a negative control (mean ± SE, n = 4). The negative control is

subtracted.

(D) *In vitro* pollen germination assay for various genotypes on sucrose-based medium with or

without 60 mM galactose. The pictures were taken 8 hours post-germination.

(E) Statistical analysis of pollen germination rates for various genotypes. The means were

calculated from multiple repeats (± SE, n ≥ 9), with over 450 pollen grains/tubes counted in total.

The statistically significant differences among different samples were determined using one-way

ANOVA followed by multiple comparison tests and were represented by different letters (P <0.05).

Figure 2. Tissue-specific accumulation and cellular localization of SWEET5.

(A) SWEET5 tissue-specific expression was evaluated. Twelve-day old seedlings, inflorescences

and siliques from about one-month old Arabidopsis carrying p*SWEET5*:*gSWEET5-GUS* were

histochemically stained 4 h for GUS activity.

(B) SWEET5 localization in pollen grain was examined using fluorescence microscopy. Images

were captured using the YFP filter for YFP, the RFP filter for propidium iodine (PI) staining, and

under bright field. SWEET5 was detected in the tricellular pollen grain (TCP) and mature pollen

grain (MPG) but absent in early stages of uninucleate microspores (UNM) and bicellular pollen

(BCP).

(C) SWEET5 accumulation in different floral stages of Arabidopsis was examined using

fluorescence microscopy. The signals started to be detected from stage 11.

(D) SWEET5 accumulation was detected after pollen germination in p*SWEET5*:*gSWEET5-YFP*

lines. Pictures were taken 8 hours post-germination using a fluorescence microscope.

(E) SWEET5 cellular localization was examined in mature pollen carrying p*SWEET5*:*cSWEET5-*

GFP.

(F) SWEET5 cellular localization was examined 72 hours after infiltration of *Nicotiana*

benthamiana leaves with *pUBI10*:*cSWEET5-mVenus*. Pictures were taken using confocal

microscopy.

Figure 3. Involvement of GALK in galactose-inhibited pollen germination

(A) Examination of *in vitro* pollen germination on sucrose-based medium with 0 mM or 60 mM

galactose for various genotypes. (means ± SE, n ≥ 6 (over 300 pollen grains/tubes counted) and *P*

<0.05).

(B) Effects of galactose-1-phosphate (6 mM) on *in vitro* pollen germination (means ± SE, n ≥ 6

(over 300 pollen grain/tubes counted) and *P* <0.05).

Figure 4. Concentration-dependent galactose response.

(A) *In vitro* pollen germination of various genotypes on medium supplemented with different concentrations of galactose (ranging from 0 to 600 mM). The means (± SE) of at least 6 repeats were plotted (over 300 pollen grains/tubes tested) ($P < 0.05$).

(B) Concentration-dependent galactose uptake in pollen of Col-0 and *sweet5* (means ± SE, n=3 and $P < 0.05$). Pollen germinated on liquid medium containing 0 mM, 0.6 mM, 6 mM, or 60 mM galactose in addition to 0.3 μCi [14]C-galactose per sample for 45 min.

(C) Relative [14]C-galactose uptake by mature pollen grains of various genotypes under different galactose concentrations (means ± SE, n=3). Pollen germinated on liquid medium containing 0.6 mM, 6 mM, or 60 mM galactose in addition to 0.3 μCi [14]C-galactose per sample for 45 min. The [14]C-galactose uptake rate for individual samples was normalized to Col-0, whose uptake rate was converted to 100%, for each concentration.

(D) Expression of *SWEET5* of Col-0 and the *galk* mutant tested by RT-qPCR with or without galactose treatment (means ± SE, n=3 and $P < 0.05$). Pollen germinated on liquid medium containing 0 mM, or 60 mM galactose for 45 min. *ACT8* was used as an internal control gene.

The gene expression level of Col-0 under sucrose only condition was normalized to 1. Different

letters represent significant difference ($P < 0.05$).

Figure 5. Schematic representation of galactose transport and metabolism in Arabidopsis pollen.

A simplified pollen structure was drawn for clarity. Arrows represent positive regulation and bar-

headed lines indicate negative regulations. The black lines represent galactose metabolic flux. The

thick black line means higher flux. The green line indicates regulation and the green dotted line is

for hypothetical regulation. Galacto-lipids are predominantly found in plastid membrane. RFOs

(Raffinose Oligosaccharides) mainly include raffinose and stachyose containing one and two

Galactose moieties, respectively. Galactans represents the cell wall-bound galactose.

792