1   Immunosuppression broadens evolutionary pathways to treatment failure during *Acinetobacter*

2   *baumannii* pneumonia

6   Wenwen Huo[1,§], Lindsay M. Busch [1,2,§], Efrat Hamami[1], Juan Hernandez-Bird[1], Christopher W.
7   Marshall[3,4], Edward Geisinger[5], Vaughn S. Cooper[3], Tim van Opijnen[6], Jason W. Rosch[7] and
8   Ralph R. Isberg[1,*]

16   [1]Department of Molecular Biology and Microbiology, Tufts University School of Medicine, 150
17   Harrison Ave., Boston, MA 02111, USA.

18
19   [2]Current address: Division of Infectious Diseases, Emory University School of Medicine

20
21   [3]Department of Microbiology and Molecular Genetics and Center for Evolutionary Biology and
22   Medicine, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA.

23
24   [4]Current Address: Dept. of Biological Sciences, Marquette University, 1428 W. Clybourn St.,
25   Milwaukee, WI. 53233.

26
27   [5]Department of Biology, Northeastern University, Boston, MA, 02115, USA

28   [6]Department of Biology, Boston College, Boston, MA, USA.

29
30   [7]Department of Infectious Diseases, St Jude Children's Research Hospital, Memphis, TN, USA.

31
32   [§]Equal contributions.
33   [*]Corresponding Author

Abstract

Acinetobacter baumannii is increasingly refractory to antibiotic treatment in healthcare settings. As is true of most human pathogens, the genetic path to antimicrobial resistance (AMR) and the role that the immune system plays in modulating AMR during disease are poorly understood. Here we reproduced several routes to fluoroquinolone resistance, performing evolution experiments using sequential lung infections in mice that are replete or depleted of neutrophils, providing two key insights into the evolution of drug resistance. First, neutropenic hosts were demonstrated to act as reservoirs for the accumulation of drug resistance. Selection for variants with altered drug sensitivity profiles arose readily in the absence of neutrophils, while immunocompetent animals restricted the appearance of these variants. Secondly, antibiotic treatment failure was shown to occur without clinically defined resistance, an unexpected result that provides a model for how antibiotic failure occurs clinically in the absence of AMR. The genetic mechanism underlying both these results is initiated by mutations activating the drug egress pump regulator AdeL, which drives persistence in the presence of the antibiotic. Therefore, antibiotic persistence mutations are demonstrated to present a two-pronged risk during disease, causing drug treatment failure in the immunocompromised host while simultaneously increasing the likelihood of high-level AMR acquisition.

Introduction

53 *Acinetobacter baumannii* is an emerging Gram-negative pathogen, one of the high-

54 priority ESKAPE organisms that is increasingly refractory to multiple antibiotic treatment

55 regimens[1]. A significant proportion of healthcare-associated diseases that are associated with

56 this group of pathogens, such as ventilator associated pneumonia, is linked to their documented

57 multi-drug resistance (MDR) [2-5]. Of particular importance are patients in intensive care units

58 (ICU) who are critically ill and have depressed immunological clearance mechanisms that

59 increase the risk of infection by MDR pathogens [6,7]. As a consequence, the healthcare

60 environment with its immunologically compromised patients could provide a unique niche for

61 selection of MDR isolates [8]. Overlaying these issues is the fact that for many patients in

62 healthcare settings, antibiotic treatment failure is common but is often unexplained, as resistant

63 organisms cannot be identified[9].

64 *A. baumannii* clinical isolates have demonstrated a remarkable ability to successfully

65 battle antibiotic treatment in the clinic, resulting from high intrinsic resistance to antimicrobials

66 and the acquisition of drug resistance elements by the organism.[10,11,12, 13] A critical missing link

67 is a detailed roadmap for the stepwise evolution of antibiotic resistance in the clinic, particularly

68 in identifying *A. baumannii* subpopulations most likely to give rise to drug treatment failure.

69 Furthermore, whether a patient group exists that provides the reservoir for antimicrobial

70 resistance (AMR) acquisition is largely unknown. Particularly for healthcare-associated diseases,

71 patient groups susceptible to *A. baumannii* are by their nature compromised in a number of

72 fashions, with the potential for providing reservoirs for AMR evolution. The range of individuals

73 with altered immune function in these settings may allow for a diversity of host targets that can

74 act as primary amplifiers of resistance, with eventual spread to individuals with different sets of

susceptibilities. Therefore, as a model for healthcare-associated pneumonia we aimed to explore

how depletion of a single arm of innate immunity in the mouses often seen in hospitalized

patients, could help shape the antibiotic treatment outcome and support evolution of resistant

organisms that can spread to others in healthcare settings.

Fluoroquinolones (FQ) such as ciprofloxacin initially showed excellent activity against *A.*

baumannii infections[14]. Members of this drug class inhibit bacterial cell growth by covalently

linking to DNA gyrase (*gyrA*) and topoisomerase IV (*parC*), leading to double stranded DNA

breaks and cell death[15]. Drug resistant mutants arise fairly frequently in the clinic, with over 80%

of clinical isolates of *A. baumannii* currently being FQ resistant[16]. The most commonly reported

mechanisms of ciprofloxacin resistance in the clinic are target protein alterations and the

overexpression of efflux pumps. In *A. baumannii*, alterations in target proteins usually evolve in

a stepwise fashion, starting with *gyrA* (usually Ser81Leu) followed by *parC* (Ser83Leu).

Interestingly, in addition to the patient's underlying condition and hospitalization status, prior

exposure to fluoroquinolone is also a risk factor for *A. baumannii* colonization and infection[17],

indicating that resistance to this antibiotic class is linked to either the pathogenic potential of the

isolate or is highly associated with acquisition of MDR. Unknown is whether there exist early

adaptive mutations that enable precursor populations of *A. baumannii* to act as ancestors to drug

resistance.

Given the poor understanding of how antibiotic resistance evolves in the clinic, and the

lack of a detailed interrogation of the role played by the immune response in controlling

selection of AMR, we sought to identify the steps that lead to resistance in the presence or

absence of a single arm of innate immunity.   The rationale behind this approach is that clinical

antibiotic resistance is associated with mutations located outside well-characterized drug targets,

and these are difficult to identify bioinformatically or predict based on culture studies [18-20]. This

approach provides evidence that absence of neutrophil function allows outgrowth of drug

persisters and the appearance of fluoroquinolone resistance. In so doing, we tie drug resistance

evolution to the problem of unexplained antibiotic treatment failure.

Results

Recapitulation of *Acinetobacter baumannii* evolved drug resistance during

pneumonic disease. We serially passaged *A. baumannii* 15 times within a mouse pneumonia

model to analyze the dynamics and genetic trajectories of resistance to the FQ antibiotic

ciprofloxacin (CIP), with the purpose of determining whether neutrophils play a role in

suppressing drug resistance (Fig. 1A). The CIP^S ATCC reference strain 17978 (AB17978) was

passaged by oropharyngeal inoculation in either immunocompetent animals or those depleted of

neutrophils by pretreatment with two doses of cyclophosphamide. Mice in three parallel lines

were CIP treated at 7- and 19- hours post infection (hpi) and bacteria were collected from the

lungs of animals euthanized at 27 hpi. The enriched bacterial pools from each independent

infection were used as inocula for the next round of infection under the two immune status

conditions. The dynamics of bacterial yield was assessed from lung homogenates, and the CIP

minimum inhibitory concentration (MIC) of each population was determined followed by whole

genome sequencing of the heterogenous pool (mean genome coverage depth: 368.4).

Persistent neutropenia during successive passages quickly results in antibiotic

treatment failure. The most striking immediate result from the passage experiments is that the

protocol allowed drug resistant mutants to overgrow during passage in neutrophil-depleted

animals, while such variants still represented a small population in immunocompetent animals.

After each mouse passage in the presence of CIP, the efficiency of bacterial colonization of the

lungs was monitored by quantifying total bacterial colony forming units in the absence of drug

(CFU). Bacterial lung yields increased by 1000-fold in CIP-treated neutrophil-depleted mice

over the course of 15 passages, as compared to a 10-fold increase in immunocompetent mice

(Fig. 1B). After each passage, we observed 10-fold higher colonization in the neutrophil-

depleted compared to immunocompetent mice. By passages 5-7, the lung bacterial load in the

neutrophil-depleted mice reached the size of the initial inoculum (10^8 CFU), and animals showed

significant signs of disease such as lacrimation, piloerection, and decreased mobility in spite of

being CIP treated. To determine levels of CIP resistance after each passage, the harvested

bacteria were quantified on solid medium in the absence or presence of increasing amounts of

drug, plotting the fraction of surviving CFU as a function of CIP concentration (Material and

Methods; Fig. 2A and 2C). While there were differences among the three immunocompetent

lineages, clinically resistant bacteria (CIP $\geq$2ug/ml) only started to arise after the 10^{th} passage,

with the fraction of these isolates generally comprising less than 1% of the population after 15

passages. Therefore, in immunocompetent mice, 15 passages were insufficient to allow

outgrowth and fixation of a resistant population.

The evolutionary trajectories through neutrophil-depleted mice contrasted strongly with

immunocompetent mice, as the whole population became CIP resistant over the course of 15

passages. Bacteria resistant to at least 2 μg/ml CIP were observed as early as the 3^{rd} passage (Fig.

2C). After the 9^{th} passage, approximately 1×10^{-5} of the bacteria isolated from the neutrophil-

depleted mice showed resistance to 8 μg/mL CIP, four times higher than the clinically-defined

resistance breakpoint. By 15 passages, almost all bacteria from the neutrophil-depleted lineages

were clinically defined as CIP resistant (at concentration of at least 2 μg/mL), with a large

fraction showing resistance to CIP =8 μg/ml. Therefore, resistance acquisition during disease is

amplified in the absence of neutrophils.

Clinical failure results from a bacterial population that is clinically defined as drug

sensitive. By the 6th passage in neutrophil-depleted mice, yields of bacteria began to approach

those observed in the absence of antibiotic (Fig. 1B). Yields remained high in successive

passages (Fig. 2C). Therefore, the population that caused drug treatment failure at passage 6 was

dominated by bacteria that showed low viability at 2μg/ml, the clinical breakpoint, while more

resistant isolates arose from this population during subsequent passages.

To link specific mutational changes over time to these observed phenotypes, we

sequenced the pools of genomic DNA isolated from the original inoculum, the parent AB17978

strain, and the pools harvested after the 3rd, 6th, 9th, 12th and 15th passages. Mutation

frequencies were then plotted for each passage in the two immune conditions (Fig. 2B and 2D).

Based on the predicted sensitivity from the depth of pooled whole genome sequencing, we

expected that only nucleotide changes in greater than 5% of the population could be detected

(Materials and Methods).

The results from the pooled genomic sequencing allowed us to identify changes

associated with either the development of drug resistance, or the lack thereof, in these

populations. As predicted from Fig. 2A and the sensitivity limits of the analysis, there was no

evidence for mutations driving drug resistance at greater than 5% abundance in the three

populations evolved in immunocompetent mice. The only significant alteration, a mutation in a

gene encoding a putative solanesyl diphosphate synthase (ACX60_03850; *ispB*) was unrelated to

resistance, as single colony isolates showed no increased survival in the presence of CIP (Fig.

2E). It was clear, however, that drug resistance variants were present at low levels in these

populations and rose to about 0.1% by the 15th passage. Two of six randomly selected colonies

from the 15th passage that formed colonies on 2µg/ml had the canonical S81L mutation in GyrA.

The rest of the strains isolated on CIP plates harbored a combination of four mutations that were

similarly found in the neutrophil-depleted lineages, which were subjected to further analysis

(Fig. 2E).

Consistent with the dynamic nature of resistance acquisition within the neutrophil-

depleted lineages (Fig. 2B), analysis of the population identified the cause of drug treatment

failure and uncovered the mutations that drove the stepwise trajectory to CIP^R (Fig. 2D). In all

three lineages, a first step mutation occurred within *adeL* between passages 3 and 6 in

neutrophil-depleted mice in all three lines. The initial *adeL* mutations were different in each line

and disrupted the 3' end of the gene. Previously, mutations in this region of the gene have been

shown to activate the AdeFGH pump regulated by AdeL [21,22], and our data are consistent with

this interpretation (below, Fig. 3C). Mutations in various other locations arose as second step

mutations between passage 9 and 12. For instance, double mutants appeared to overgrow

Lineages 1 and 2, while a quadruple mutant appeared to overgrow Lineage 3.

To identify the various genotypes linked with increased CIP resistance, we isolated single

colonies after 15 passages with or without CIP (1µg/mL) selection and sequenced two individual

purified clones per condition. Indeed, among all single colonies harvested from the neutrophil-

depleted lineages, the majority of individual clones accumulated more than one mutation (Fig.

2E). In two lineages of the neutrophil-depleted mouse infections, alleles of DNA gyrase

observed infrequently in the clinic, were identified after 15 passages (*gyrA*(A117E) and

gyrB(Q447H), respectively). Having *gyrA*(A117E) alone was able to raise the CIP MIC by 16-

fold (Fig 2E). Clones having the *gyrB*(Q447H) allele were only observed linked to *adeL*(C312*),

resulting in CIP MIC that was 8-fold higher than that of the parent strain (Fig 2E). Besides the

mutations in target proteins, other single colonies that resulted in a similar MIC increase were

found to have a combination of four mutations (Fig. 2E). When comparing the single colonies

isolated that had a CIP of 1 µg/mL to those isolated without antibiotic, the colonies presented the

same genomic changes, indicating that the mutants described here became predominant without

requiring CIP selection *ex vivo*.

Mutations in *adeL* allow for persistence in the presence of high levels of

ciprofloxacin. To deconvolute the function of the various mutations observed, we backcrossed

mutations into the parent strain and assessed their relative contributions to CIP resistance. Of

particular interest were nucleotide changes identified in populations after the 6[th] passage in

neutrophil-depleted mouse lines, as CIP failed to efficiently restrict growth in the lung in these

passages, and this failure was associated with the outgrowth of single *adeL* mutations (Figs. 1A,

2D). The *adeL* gene (ACX60_06025)[22] has two predicted domains often associated with LysR

type transcriptional regulators: a helix-turn-helix domain and a substrate binding domain that are

responsible for regulation of the AdeEFG pump (Fig. 3A). Of the *adeL* mutations that arose

during the *in vivo* passages, one is located within the predicted substrate binding domain while

the others are at the C terminal end comprising in-frame deletions or early termination codons

(Fig. 3A and 3B). We constructed strains with each of these mutations and observed 100-1000X

increased *adeG* transcription levels compared to the parental strain, consistent with AdeFGH

efflux pump overproduction (Fig. 3C).

While these single *adeL* mutations resulted in an increase in MIC to ~1 µg/mL CIP (Fig.

4C), this level was below the recognized clinical breakpoint[23] and such strains would be

indicated as susceptible. As these are first step mutations, we hypothesize that variants harboring

these changes may facilitate the outgrowth of more resistant isolates by promoting tolerance or

persistence in the presence of antibiotic. To test this model, we evaluated the survival of *adeL*

mutants during exposure to high levels of CIP [24-28]. Exposure to 10 μg/mL CIP, roughly 20X

MIC of the parental strain, led to the majority of the *adeL* mutants (> 99%; MDK99) being

rapidly killed (within 1 hour), mimicking what was observed for the parental strain (Fig. 3D).

However, unlike the parental strain, a subpopulation was able to persist through 24 hours of drug

exposure (Fig. 3D). This phenotype was dependent on pump overproduction, as an *adeL* mutant

strain deleted for the *adeFGH* operon was indistinguishable from the parental strain (Fig. 3D).

These data argue that *adeL* mutations drive persistence in the presence of CIP *via* increased

expression of *adeFGH*, providing a reservoir for the outgrowth of strains with increased

resistance.

Resistance of evolved mutants in neutrophil-depleted mice can be explained by

combining *adeL* with *mfsA* mutations. We continued backcross analysis to identify the

minimal determinants necessary to confer clinical resistance during passage in the mouse. Many

of the single colony isolates from lineage 3 at passage 15 of the immunocompromised mice

possessed a combination of mutations in four genomic locations: *adeL*, *acfR*, *lpxD*, and the

intergenic *rpmL-mfsA* region, the latter of which is located between a ribosomal protein gene and

a predicted coding region for a putative MFS transporter (ACX60_15150, annotated as *nreB*;

Fig. 2D and 2E). To determine the role in drug susceptibility of each of these mutations, we

constructed separate strains carrying single mutations. All single mutants showed various MICs

well below 2 μg/ml CIP, with each of the *adeL* mutations as well as *mfsA*(-9) showing increased

MICs above the parental strain (Table S1; Fig. 4C). The *mfsA*(-9) mutation is located 9 bp

upstream of *mfsA*, within 2 bases of a previously reported mutation selected during evolution of

CIP resistance in *A. baumannii* in planktonic conditions[29]. Consistent with MFS transporter over-

production, we found that the *mfsA*(-9) mutation increased the *mfsA* mRNA levels by ~9-fold

relative to the parental strain, based on q-rtPCR analysis (Fig. 4B). Significantly, when the

mfsA(-9) mutation was combined with the *adeL*P131H mutation, the CIP MIC increased to

~4ug/ml and became indistinguishable from the quadruple mutant strain isolated from *in vivo*

passage 15 (Fig. 4C; Table S1). Therefore, combining two mutations that resulted in

upregulation of both the AdeFGH pump and a putative MFS transporter was sufficient to

reconstruct CIP resistance.

A hotspot for mutations within *lpxD* is tightly linked to fluoroquinolone resistance

in clinical isolates. The mutation *lpxD* T118_A119insA (locus tag: ACX60_07955; GenBank:

AKQ26661.1) was consistently found in CIP^R isolates from mice. This alteration within an

acyltransferase involved in lipooligosaccharide (LOS) biosynthesis [30-32] showed no effect on

MIC (Table S1). To test for clinical linkage of *lpxD* mutations to drug resistance, we analyzed

the genomes of 4671 clinical isolates [33] and identified fluoroquinolone resistant strains based on

the *gyrA*S81L resistance allele. The LpxD sequences were identified among this group, and

linkage to *gyrA*S81L was tested by plotting z-scores as a function of residue number (Fig. 5B;

Material and Methods). Of the CIP resistant isolates, 2762/4062 had alterations at residue E117

(z-score: 8.98; p-value, < 0.00001), which is adjacent to the T118 allele identified in our mouse

experiments, with both the E117K and T118_A119insA variants predicted to be in a turn

between two beta sheets (Fig. 5C)[34]. Although the significance of this linkage is unclear, the fact

that the *lpxD* T118_A119insA has a fitness defect in broth (Fig. 5D), raises the possibility that it

could contribute to a persistence phenomenon in tissues leading to evolution of drug resistance.

Evolutionary replay experiments reconstruct the pathway to drug resistance. The

observation that a resistant organism outcompeted a persistent strain (Fig. 2D) could be due to a

coincidence that occurred during serial passage or due to antibiotic-driven selection for strains

with increased MIC over persistent strains. To distinguish between these possibilities for the

neutrophil-depleted Lineage 3 variants, we performed two evolutionary replay experiments [35]. In

the first (Fig. 6A, B), bacteria were harvested after passage 9 and cycled 3 times in duplicate in

neutrophil-depleted animals to test if the *adeL* I335A336 allele would again be outcompeted by

initially rare drug-resistant mutants (Fig. 2D). The second approach was to mix a colony-purified

drug persistent *adeL* I335A336 single mutant (P9Cy3.1; Table S1) with a colony-purified drug

resistant quadruple mutant (Cy31; Table S1) in the approximate ratio present at passage 9 (95:5),

before passaging three times (Material and Methods; Fig. 6C).

In both experiments, we observed an increase in bacterial burden and fraction of bacteria

showing CIP MIC >2 after passaging the pools in neutrophil-depleted CIP-treated mice (Supp.

Fig. 1). Whole genome sequencing of the bacterial populations harvested after each passage

revealed a near identical evolutionary trajectory as seen from the original experiment, with the

single mutant associated with drug persistence eventually outcompeted by drug resistant mutants

(Fig. 6). Remarkably, in one of our replicate lineages (Fig. 6A), a new mutation arose in the

intergenic region between *rpmL* and *mfsA* (*mfsA*(-19)), arguing for the contribution of this

regulatory region in resistance evolution. Analyzed in isolation from other mutations, strains

harboring *mfsA*(-19) behaved similarly to single mutants having the previously described *mfsA*(-

9) mutation, as it results in a similar increase in MIC values (Fig. 4C), and is associated with up-

regulation of *mfsA* transcript levels (Fig. 4B). Therefore, this demonstrates that the evolutionary

trajectory is reproducible, and can be regenerated by phenotypically similar mutations that arise

spontaneously.

Ciprofloxacin treatment within the host simultaneously selects for increased

resistance and enhanced fitness. Antibiotic resistance is often associated with a fitness cost[36]

that may be compensated over time by continued selection on secondary genotypes that

overcome these costs. As a consequence, mutants with high resistance could maintain their

selective advantage over non or low-resistance mutants even in the absence of drug treatment. To

test this model, the relative fitness of the drug resistant mutants derived from neutrophil-depleted

lineages was compared to the parent strain AB17978. In the absence of CIP, variants isolated

after passage in neutrophil-depleted mice showed varying degrees of subtle growth impairment

in broth relative to the parental strain (Fig. 7A; Cy11, $p < 0.02$). Similar results were observed

when the mutants were challenged in competition with the WT strain in immunocompetent

animals, although the differences did not consistently rise to the level of statistical significance.

(Fig. 7B).

The lowered fitness of mutants, however, did not directly test the model that selection in

neutropenic mice increased fitness relative to first-step CIP^R variants. In the absence of CIP

treatment, the parent strain outcompeted the Lineage 3 quadruple mutant isolate Cy31 (Table S1)

under neutrophil-depleted conditions, similar to results observed in the presence of immune

function (Fig. 7C; Cy31 against WT). Nonetheless, the Cy31 strain had a clear fitness advantage

over its predecessor Lineage 3 *adeL* mutant when competed in the absence of antibiotics (Fig.

7C; Cy31 against P9Cy3.1). Therefore, in competition with the persister mutant, ciprofloxacin

simultaneously selected for an isolate with increased CIP resistance and enhanced fitness relative

to mutant competitors in the absence of drug. This predicts that drug-resistant mutants are able to

evolve to heightened relative fitness and replace persister mutants. As fitness increases, resistant

mutants should be able to compete with sensitive strains even after cessation of antibiotic

treatment, potentially escalating treatment failure and providing fertile ground for the emergence

of resistant strains with high fitness even in immunocompetent hosts.

Discussion

Comparative analyses of bacterial pathogen databases indicate that clinical antibiotic

resistance is associated with mutations located outside well-characterized drug targets [18-20].

Analysis of clinical resistance is often retrospective, with a few exceptional studies that have

allowed identification of a timeline of bacterial resistance in human populations[24-28,37-39]. Our

study bridges an important gap in understanding how resistance evolves, by following

experimental evolution of fluoroquinolone resistance in a pneumonia disease model. Strikingly,

in the absence of one arm of immune pressure, *A. baumannii* resistance arose after few passages

in murine lungs during CIP treatment (Fig. 1B), resulting in treatment failure. Conversely, while

there was a small subpopulation having clinically-defined resistance to CIP in healthy

immunocompetent mice, the overall frequency remained low and this did not lead to treatment

failure (Fig. 1B) (Fig. 2A).

Importantly, the spectrum of selected mutations was altered by the absence of

neutrophils. In contrast to the canonical target site *gyrA*S81L mutation observed in

immunocompetent animals, the *gyrA*A117E and *gyrB*Q447H alleles rarely observed in clinical

strains predominated in two of the lines from neutrophil-depleted animals (Fig 2E; 6 out of 12

CIP resistant clones). These unusual isolates showed clear fitness defects during disease and in

culture (Fig. 6). These non-canonical *gyrA* and *gyrB* mutations with low fitness may be unlikely

to arise in the presence of intact immune functions, which is consistent with evolutionary

trajectories being determined by the immunological state of the host. Therefore, we propose that

immunocompromised hosts are incubators for the generation of unique drug resistant variants,

with uncertain outcomes after pathogen exposure to the community at large. Although the

variants we described have reduced fitness relative to WT, continued passage demonstrated that

stepwise increase in drug resistance was associated with stepwise increase in fitness (Fig. 7).

Similar fitness effects have been obtained during continued passage in culture[29]. The recent

demonstration of enhanced viral evolution of SARS-CoV-2 in an immunocompromised patient

undergoing therapeutic antibody treatment is a graphic example of the potential interplay

between immunity, antimicrobials and evolution in the clinic, with potential largescale

community effects [40]. In fact, persistence mutants have been demonstrated to arise during

continued antibiotic therapy in an immunocompromised patient treated for vancomycin-resistant

enterococcus[41], further tying the results presented here with documented clinical outcomes.

Antibiotics primarily act by targeting essential cellular functions or pathways [42,43]. In

response to transient antibiotic exposure, a proportion of bacteria typically persist or tolerate this

treatment with mutations arising that can increase the fraction of the surviving population[44-46].

The mechanisms for allowing persistence/tolerance likely differ between bacterial species and

vary among antibiotics[47], but it is generally agreed that persistence/tolerance is caused by cell

dormancy[48-50] or transient expression of efflux pumps and stress response pathways[24,51].

Furthermore, persistence/tolerance mechanisms may play an important role in the relapse of

bacterial infections [52-54]. In our study, we found that persistence in *A. baumannii* can be

promoted by upregulation of a drug efflux pump, AdeFGH [22]. It is known that many clinical

isolates are found to overproduce efflux pumps that remove multiple antibiotics from the

bacterial cytoplasm[22,55], and that antibiotic persistence is associated with such pump

upregulation[56,57]. Therefore, it is likely that this is a common motif for initiating the

evolutionary path to drug resistance. Of particular note, mutations upregulating this pump are

rarely isolated in the laboratory through simple antibiotic single-step selections on solid agar,

indicating there may be a special environment in the lung that allows outgrowth of these mutants.

There are three important repercussions from the analysis of the AdeL efflux

overproducer mutations. First, overproduction results in drug treatment failure during

pneumonic disease in the mouse, in spite of the fact that these bacterial strains are CIP-sensitive

based on the international standard for clinical breakpoints[23]. Therefore, clinical drug treatment

failure in the absence of identified antibiotic resistance[9] could be explained by the outgrowth of

drug persistent mutants, taking advantage of fluctuations in drug availability to the infection site

between doses. Second, the mutation provides a molecular basis for *A. baumannii* to develop

second-step variants that lead to CIP resistance above the clinical breakpoint. Of interest,

inhibitors of efflux pumps have been suggested to re-sensitize bacteria to multiple antibiotics[58],

but perhaps such a regimen may also delay progression to resistance. Finally, isolates from drug

treatment failure occurring in immunocompromised patients could generate a pool of precursor

mutants giving rise to resistant isolates that eventually infect a broad range of patients.

We here propose a model of bacterial resistance progression in immunodepleted hosts

(Fig. 8). When infecting a host, the antibiotic susceptible bacteria colonize at a relatively low

rate. With active transmission and continuous CIP treatment, bacterial persisters arise that have

MICs below the clinical breakpoint, increasing bacterial load at the infection site. On continued

treatment, or transfer to a similar host undergoing CIP treatment, fully resistant mutants outgrow

the population. Although the presence of intact immune function may delay outgrowth of such

mutants, the fact that we could identify a mutant with a persister allele in the presence of immune

function indicates that such alleles could act as enablers of drug resistance in all hosts.

Genomic analyses can identify allelic variants linked to drug resistance, but verifying the

functional importance of these alleles is hindered by founder effects that are often difficult to

discount. The *lpxD* E117K mutation is one such allele associated with resistant *gyrA* S81L-

containing genomes. The fact that during multiple passages in immune-depleted mice the

adjacent *lpxD* T118 insA allele was isolated argues for a role of these altered residues in

supporting the outgrowth of drug resistant organisms. The *lpxD* T118 insA mutation likely has

an effect on envelope function. This raises the possibility that the mutation causes subtle

changes in permeability that slow drug access to target, or else the slowed growth of strains

harboring this allele increases drug tolerance during growth in tissues[47]. Our inability to

demonstrate increased tolerance to CIP and the fact that the clinical *lpxD* E117K did not show a

fitness defect when grown in LB broth (Fig. 5D) argues that the critical phenotypes may be

observed exclusively during growth in tissues, making them difficult to evaluate.

In summary, we hypothesize that resistance progression in clinical isolates follows a

similar trend witnessed in our experiments, albeit perhaps with greater complexity in the human.

Drug sensitive bacteria may colonize both healthy and immune system-compromised patients,

but resistance evolution occurs more rapidly within hosts having impaired neutrophil function.

The canonical resistance mutations that arise within healthy individuals may compete efficiently

with host-evolved mutations in patients having impaired neutrophil function. Hence, the

transmission from individuals having intact immune function may pose risks to

immunocompromised patients. Adding to the risks subjected to vulnerable patients, we found

that *A. baumannii* persistence mutants with MIC levels below the clinical breakpoint were

associated with treatment failure, emphasizing the difficulty in treating these patients (Fig. 1B)

[9,59-62]. While heterogeneity and variability between infected patients is inevitable, this work

provides insight into some genetic factors that could predispose individuals to treatment failures

in specific clinical contexts. Future work should be focused on how this evolutionary trend

towards drug resistance can be controlled, such as tackling the molecular basis for selection of

persister or other mutations that lead to treatment failure.

Material and Methods

Bacterial Strains. Bacterial strains used in this study are listed in Table S1.

Acinetobacter baumannii strains are derivatives of ATCC 17978. Bacteria were grown in

Lysogeny Broth (LB) (10 g/L tryptone, 5 g/L yeast extract, 10 g/L NaCl) or on LB agar plates

(LB supplemented with 15g/L agarose). Broth cultures were grown at 37°C in flasks with orbital

shaking at 200 rpm or in tubes with rotation on a roller drum at 56 rpm. Growth was monitored

by measuring absorbance spectrophotometrically at 600 nm (A_{600nm}). Plates were incubated at

37°C. Antibiotics were used at the following concentrations: Gentamicin: 10 μg/mL (*A.*

baumannii) and 50 μg/mL (*E. coli*); Carbenicillin: 100 μg/mL.

Murine experimental evolution. The *in vivo* passaging experiments were performed

identically in either immunocompetent or neutrophil-depleted 6-8 weeks-old female BALB/C

mice (Fig. 1A). Neutrophil depletion was induced via cyclophosphamide pretreatment [63]. Briefly,

150 mg/kg and 100 mg/kg of cyclophosphamide monohydrate (Sigma-Aldrich: C7397) were

administered 4 days and 1 day prior to infection, respectively. At the time of infection, mice

were transiently anesthetized by inhalation of isoflurane and lung infections were established

using ~10^8 colony forming units (CFU) of *A. baumannii* from mid-log phase cultures (A_{600nm}

~0.5) via the oropharyngeal aspiration [64]. At 7 and 19 hours post infection (hpi), 100 mg/kg of

ciprofloxacin (Sigma-Aldrich: PHR1044) was administered via subcutaneous injection. At 27

hpi, mice were euthanized, and lungs were removed aseptically and homogenized in 1 mL of ice-

cold 1x phosphate buffered saline (PBS). Afterwards, the homogenate was plated on large (150 x

15 mm) LB agar plates and incubated for ~17 hours at 37°C. The total number of bacteria was

quantified using CFU. The enriched bacteria were scraped off the LB plates and resuspended in

1x PBS. The resuspension was used for DNA isolation as well as the inoculum for the next

passage/round of infection (stored at -80°C in 20% glycerol prior to use). For each round of

mouse infection/passage, bacterial frozen stocks were revived by diluting them in LB broth and

growing them to exponential phase. Afterwards, ~10^8 CFU of the pool was used to establish

infection in each mouse. Three separate lineages of murine infections were maintained in parallel

in each condition (immunocompetent or neutrophil depleted), and within each lineage each

starting inoculum was passaged 15 times. Validation experiments involving the reconstruction of

identified mutants and re-passaging of these strains in mice were similarly performed with the

only difference being that the infection was terminated at 24 hpi instead of at 27 hpi.

Determining the fraction of resistance to set ciprofloxacin concentrations in each

pool. To evaluate the fraction of bacteria acquired from each passage that are viable on culturing

in various concentration of ciprofloxacin, the glycerol stock of each pool was revived in fresh LB

broth and grown to exponential phase. A total of ~10^7 CFU were used for serial dilutions in 1x

PBS and 10 µl from each diluted culture was spotted on LB agar plates containing the following

concentrations of ciprofloxacin: 0, 0.25, 0.5, 1, 2, 4 and 8 µg/ml. After an overnight incubation at

37°C, colonies were counted, and CFU/ml was calculated. A detection limit of 100 CFU/ml was

used. The fraction bacteria resistant to a certain concentration (C_0; $C_0 > 0$) of ciprofloxacin is

calculated as (CFU/ml at C_0 - CFU/ml at all concentrations above C_0) / Total CFU. A stacked bar

plot of the data was generated using Prism GraphPad (Fig. 2).

Whole Genome Sequencing. Genomic DNA (gDNA) was extracted from bacteria using

the DNeasy Blood and Tissue kit (Qiagen). Library preparation was performed using Illumina

Nextera with previously described modifications [65]. Libraries were enriched with 8 cycles of

PCR and sequenced using HiSeq2500 at Tufts University Core Facility using single-end 100 bp

reads. Reads were aligned to the *A. baumannii* ATCC 17978 genome and its plasmids (GenBank

Accession: CP012004; pAB1: CP000522; pAB2: CP00523; pAB3: CP012005), and variants

were identified at 5% cutoff using breseq 0.32 [66]. The predicted functional impact of substitution

variants was determined by using PROVEAN [67].

Competition Assays. Competition experiments were performed as described

previously[64]. Briefly, designated strains were grown to exponential phase ($A_{600nm} \sim 0.5\text{-}1.0$) and

mixed at a 1:1 ratio for the inoculum. Afterwards, $\sim 10^8$ CFU of the bacterial mixture was used to

challenge WT or cyclophosphamide-treated BALB/C mice via the oropharyngeal route as

described for the mouse passage experiments. 24 hr post inoculation, lungs were removed

aseptically, homogenized and extracts were spread on LB agar plates. To quantify the input and

output ratio of each designated strain, both the inoculum and the lung homogenate were serially

diluted in 1x PBS and plated on LB agar as well as LB agar supplemented with 2 µg/mL of

ciprofloxacin to differentiate between the WT and CIP resistant test strains. Competitive index

(CI) was calculated as CI = output (unmarked/marked) / input (unmarked/marked).

Persistence Assay. *A. baumannii* was grown in 3 individual LB broth cultures overnight

from separate colonies. The cultures were diluted 1000-fold into 8 ml LB broth and incubated by

rotation at 37°C for 2-3 hours until reaching mid or late exponential growth ($A_{600nm} \sim 0.3\text{-}0.8$).

Ciprofloxacin was added to reach a final concentration of 10 µg/ml, the concentration regarded

as 20x the MIC for the parent ATCC 17978 *A. baumannii* strain. These cultures were then

incubated at 37°C for up to 24 hours. After 0, 1, 3, 6, and 24 hours, 500 μl of culture was

removed, washed twice in 1x PBS, resuspended in 500 μl PBS, and sequentially diluted in 1x

PBS. Serial dilutions were used for CFU quantification by spot plating on LB agar plates.

Dilution factors harboring quantities ranging from 3-35 CFUs were used to calculate the

CFU/mL. The limit of detection by this assay was determined to be 375 CFUs. The CFU/mL

across the biological replicates at each time point throughout the drug challenge were plotted as

mean $\pm$ Standard Error of Mean.

qRT-PCR gene expression analysis. Bacteria were grown to early stationary phase in

LB broth, RNA was harvested and purified using RNeasy kit (Qiagen) followed by cDNA

synthesis using SuperScript VILO cDNA kit (Thermo Fisher). The cDNA was then amplified

with PowerUp SYBR Green Master Mix (Applied Biosystems) via a StepOnePlus Real-Time

PCR system (Applied Biosystems) per manufacturer's instructions, and target amplification

efficiency was evaluated by generating a standard curve with dilutions of cDNA (>95%

amplification efficiency for each primer pair). Primers were designed to amplify regions of

around 150 bp internal to genes (Table S1). Triple technical replicates were examined per

biological sample and at least three biological replicates per strain were tested, with controls

lacking reverse-transcriptase included to verify a lack of contaminant genomic DNA. Transcript

levels of specific targets from each strain were evaluated by the comparative $2^{-\Delta\Delta Ct}$ method to the

parental strain, normalizing to that of the endogenous control 16s. The transcript level for each

target across biological replicates was plotted as mean $\pm$ Standard Error of Mean (Fig. 3&4).

MIC determination. Minimum inhibitory concentrations (MICs) were determined by

broth microdilution. Overnight cultures of strains of interest were diluted 1000X in fresh LB

broth and grown to mid-logarithmic phase (A_{600} ~0.5). Afterwards, cultures were diluted to a

final $A_{600} = 0.003$ and tested in the presence of two-fold dilutions of antibiotics. 200 µl of

culture-antibiotic mixture was then aliquoted to a 96 well plate (COSTAR) and technical

duplicates were performed in a Biotek plate reader with rotation. Growth was monitored by

measuring A_{600} at 15 min. intervals for 16 hr, and the MIC was determined as the lowest

concentration of drug that prevented growth, using at least three biological replicates for each

strain. The MICs across biological replicates were plotted as mean $\pm$ Standard Error of Mean

(Fig. 4).

Molecular cloning and mutant construction. Plasmids and primers used in this study

are listed in Table S1. The mutant strains were constructed through sequential cloning into

pUC18[68] then pJB4648[69] as previously described[13]. Briefly, the mutant allele from each strain of

interest was amplified alongside upstream and downstream segments to generate a PCR product

~1500 bp in length flanked by appropriate restriction sites. The PCR product was then ligated

into pUC18 and propagated in *E. coli* DH5α. The PCR product was then subcloned from pUC18

into pJB4648 and propagated in *E. coli* DH5α λpir. After sequence confirmation, the pJB4648

plasmid construct containing the desired mutation was introduced into *A. baumannii* via

electroporation. Markerless, in-frame mutations were isolated via homologous recombination as

described [13].

Comparative sequence analysis. The LpxD protein sequence from ATCC 17978 was

compared to genomes downloaded from PATRIC database using tblastn to identify the amino

acid location of mismatches and gaps. In total, 4671 genomes were split into three groups: those

with a GyrA S81L mutation; those with the same GyrA sequence as in AB17978; and the rest.

Within each group, the mismatches/gaps at each amino acid location within LpxD were

displayed as number of genomes having mismatches/gaps at that location. A table of the data

was constructed showing the specific amino acid in the rows, the genome groups harboring these

mutations in the columns, and the total number of mismatches/gaps in the cells. The results in

each cell were then normalized using z-scores with the mean and standard deviation from

individual genome groups. The z-score was plotted as function of each residue.

Calculation of doubling time in broth culture. Overnight cultures of bacteria were

diluted 1000-fold in fresh LB broth and grown for 8 hours. The A_{600} values were measured every

hour using a spectrophotometer, and the log2 transformed values were plotted as a function of

time (hr). To calculate the doubling time, a modified sigmoid function ($y = \frac{L1}{e^{-k(x-x0)}+1} - L2$)

was fitted to the plotted curve(y: log2 transformed A_{600} value, x: time in hrs, x_0: timepoint when

the growth rate was the fastest, k, L1, L2: rate constants), with the fastest growth rate for each

culture determined as the time in which second derivative of the fitted curve approached 0 (or

when $x = x0$), at which point the doubling time was represented as the inverse of the slope (Δt =

$^4/_{k * L1}$) (Code available at: https://github.com/huoww07/calulate_bacteria_doubling_time). To

evaluate the goodness-of-fit, the R^2 value was calculated for each growth curve assay (the R^2

ranges from 0 to 1, representing the worst to the best fit). At least three biological replicates were

performed for each test strain (Table S2, Table S3) and the average doubling time and standard

deviation was calculated. Significance was determined using Student two-tailed t-test.

Accession numbers.

The datasets generated during the current study are available in the Sequence Read Archive (PRJNA485355).

Acknowledgements

This work is supported by NIAID U01AI12430 to TvO, JWR, VSC and RI, NIH R21AI128328 to RI and T32AI07422 support for JH-B. We thank the colleagues in all of our labs for support and constant critiquing of the experimental strategies used here.

Author Contributions

RI, JR and TvO devised the study. LB and JH performed the mice experiments. WH, EH and JH performed the wet-lab experiments. WH and LB performed the whole genome sequencing and bioinformatics analyses. CM, VC, EG and TvO collaborated on the bioinformatics analyses and performed training. WH, LM, EH, JH and RI interpreted the results and wrote the manuscript. WH, LB, EH, JH, CM, EG, VC, TvO, JR and RI approved the final manuscript.

Figure legends

Figure 1. Continuous passaging of *A. baumannii* in a lung infection model results in early antibiotic failure in neutrophil-depleted mice. A). The 15-passage strategy using the pneumonia model in both immunocompetent and neutrophil-depleted mice. Neutrophil depletion was induced by two doses of cyclophosphamide at 72- and 24-hrs prior of infection. At the time of infection, *A. baumannii* ATCC 17978 was oropharyngeally inoculated into mice at 10^8 CFU, and two doses of CIP were administered at 9- and 19-hrs post infection. At 27-hrs post infection, the mice were euthanized, lungs were homogenized, and bacteria were plated on LB agar plates . After overnight growth, the bacteria were collected and 10^8 CFU used for the next round of infection/passage. B). Rapid development of antibiotic failure as a consequence of passage in the absence of neutrophils. After lungs were homogenized, total CFU on LB agar in the absence of antibiotic were determined and plotted as function of passage number. Shown are three lines (L1, L2, L3) for immunocompetent and neutrophil-depleted animals treated with CIP as described in panel (A).

Figure 2. High level ciprofloxacin resistance evolves in a stepwise fashion during passage in neutrophil-depleted mice. (A,C). After each passage, bacteria from lung homogenates were serially diluted onto LB agar containing noted concentrations of CIP (μg/ml). The fraction of CFU on the graded series of CIP plates was determined for each pool and displayed in stacked bar plots for immunocompetent (A) and neutrophil-depleted (C) mouse infection lineages. The limit of detection was 2.7E-6. (B,D) The gDNA from each bacterial pool was isolated and subject to whole genome sequencing to identify the genomic mutations acquired throughout passaging (after passage 3, 6, 9, 12 and 15). The mutations were detected using a 5% abundance

cutoff and filtered against the parent strain AB17978. The relative abundance of each mutation

was plotted as a function of passage number from (B) immunocompetent lineages and (D)

neutrophil-depleted lineages (D). (E) After 15 passages, the bacterial pool from each lineage

was grown on LB with or without CIP (1 µg/mL) to select for single colonies. Colonies were

purified and subject to whole genome sequencing and MIC assays. In (E), each column

represents a single colony and the rows represent selection criteria, individual mutations and

MIC values. The detected mutations with >90% frequency were aggregated for all colonies and

the presence of each mutation in a certain colony was highlighted in purple. MICs were labeled

and color-coded for all colonies.

Figure 3. *adeL* mutations drive bacterial persistence through over-expression of the

AdeFGH pump. (A). Sites of nonsynonymous changes during *in vivo* passaging and domain

structure of AdeL. (B). Locations of nonsynonymous changes on predicted functional domains of

AdeL. (C). Mutations in *adeL* selected during mouse passage result in specific overproduction of

an AdeL-regulated gene *adeG*. Data were determined by qRT-PCR analysis of pump-encoding

components *adeG* (regulated by AdeL) or *adeB* and *adeJ* (components of other pumps) in noted

adeL mutant backgrounds. (D). Overproduction of AdeL-regulated pump components results in

increased persistence in presence of CIP. AB17978 derivatives described in legend were

exposed to CIP at 20X MIC for noted times and titered for viability by quantitating CFUs.

Figure 4. The presence of mutations in *adeL* and *mfsA* is sufficient to explain CIP[R] in

strains derived from mouse passage experiments. A) Two mutations located directly upstream

of *mfsA* selected during mouse passaging experiments. B) Transcription levels of *mfsA* and *rpmL*

were determined by qRT-PCR in strains harboring the two mutations upstream of *mfsA*. ****, p

< 0.0001. C) Double mutants containing *adeL* and *mfsA* mutations show MICs of CIP that are

indistinguishable from evolvedquadruple mutant strain. Data are mean MICs ± SEM.

Figure 5. The LpxD E117K allele is tightly linked to clinical fluoroquinolone resistance.

Assembled genomes of 4671 clinical isolates were obtained from PATRIC database[33]. A).

Fraction of genomes having the *gyrA*S81L resistance allele. Among the downloaded genomes,

4062 genomes presented the GyrA S81L allele (shown in red), 386 genomes had the same GyrA

allele as ATCC 17978 (shown in green) and the rest had other alleles (shown in black). B) LpxD

sequences from all genomes were aligned and compared to ATCC 17978 and the total number of

variants per defined amino acid was calculated for each group of genomes. The total number of

amino acid changes were normalized and presented as z-scores for each of the three groups, plotted

against the residue numbers. C). Structure of *Chlamydia trachomatis* LpxD protein [34] showing the

presumed site of the *A. baumannii* E117 based on sequence similarity. D). Broth growth of

AB17978 strains differing by single changes in *lpxD*. Shown are strains harboring either the WT,

lpxD T118A119_insA or *lpxD* E117K alleles.

Figure 6. The evolutionary trajectories of isolates acquired throughout passaging in

neutrophil-depleted mice is due to fitness advantage in presence of drug. (A,B) A single

adeL mutant is outcompeted during infection of CIP-treated mice by a quadruple mutant having

drug resistant *adeLmfsA* alleles. The passage 9 pool was oropharyngeally inoculated into

neutrophil-depleted mice in duplicate and passaged three times, subjecting each passage to deep

sequencing. Displayed are relative abundance of all mutations identified in deep sequencing

pools of the duplicate (panels A,B) passaging experiments as a function of the passage number.

C) Competition of purfied *adeL* and quadruple mutant having *adeLmfsA* alleles in presence of

CIP results in predominance of the quadruple mutant. The two strains were mixed at a ratio 95:5

(single:quadruple mutant), and oropharyngeally inoculated into neutrophil-depleted mice in the

presence of CIP through three successive passages. Deep sequencing of the initial inoculum and

each subsequent passage was performed to determine the ratio of the two strains, and the relative

abundance of each mutation is displayed as a function of passage.

Figure 7. Simultaneous selection for increased resistance and fitness. A) Growth rates in

broth culture of noted isolates from passage in mice. B) Neutrophil-replete and C) Neutrophil

depleted (cyclophosphamide treated) BALB/C mice were challenged with noted strain

combination to determine relative competition index of each pair of strains. C.I. values were

determined by C.I. = (strain1/strain2 output ratio)/(strain1/strain2 input ratio) (Materials and

Methods). Data are mean $\pm$ SEM, with significance determine by one-way ANOVA followed by

Multiple Comparison. *p<0.05; ****p<0.0001.

Figure 8. Model of bacterial resistance progression. The antibiotic susceptible inoculum

(blue) colonizes poorly in presence of antibiotic. During continued passage with CIP, in the

absence of neutrophils, the persister mutants (yellow) arise as first step mutations showing

intermediate MICs. With continued passage under the same antibiotic treatment regime, second

step mutations arise that have increased fitness in the presence of antibiotic (red) and dominate

the population.

Supplemental Figure 1. Bacterial load and resistance levels to CIP after re-passaging. A)

The bacterial load harvested from the lungs after each passage is plotted against the passage

number for each lineage. B) CIP resistance level of each pool (after each passage) was assayed

via quantification of CFU on LB agar supplemented with CIP.

Competing Interests Statement

The authors declare that they have no conflict of interest.

Reference

1   Boucher, Helen W. *et al.* Bad Bugs, No Drugs: No ESKAPE! An Update from the Infectious
Diseases Society of America. *Clin. Infect. Dis.* **48**, 1-12, doi:10.1086/595011 (2009).

2   Valentine, S. C. *et al.* Phenotypic and Molecular Characterization of *Acinetobacter baumannii*
Clinical Isolates from Nosocomial Outbreaks in Los Angeles County, California. *J. Clin. Microbiol.* **46**,
2499-2507, doi:10.1128/jcm.00367-08 (2008).

3   Sievert, D. M. *et al.* Antimicrobial-Resistant Pathogens Associated with Healthcare-Associated
Infections Summary of Data Reported to the National Healthcare Safety Network at the Centers for
Disease Control and Prevention, 2009–2010. *Infect. Control Hosp. Epidemiol.* **34**, 1-14,
doi:10.1086/668770 (2015).

4   Karlowsky, J. A., Hoban, D. J., DeCorby, M. R., Laing, N. M. & Zhanel, G. G. Fluoroquinolone-
Resistant Urinary Isolates of *Escherichia coli* from Outpatients Are Frequently Multidrug Resistant:
Results from the North American Urinary Tract Infection Collaborative Alliance-Quinolone Resistance
Study. *Antimicrob. Agents Chemother.* **50**, 2251-2254, doi:10.1128/aac.00123-06 (2006).

5   Azim, A. *et al.* Epidemiology of bacterial colonization at intensive care unit admission with
emphasis on extended-spectrum β-lactamase- and metallo-β-lactamase-producing Gram-negative bacteria
– an Indian experience. *J. Med. Microbiol.* **59**, 955-960, doi:10.1099/jmm.0.018085-0 (2010).

6   Montefour, K. *et al. Acinetobacter baumannii*: an emerging multidrug-resistant pathogen in
critical care. *Crit. Care Nurse* **28**, 15-25; quiz 26 (2008).

7   Zeighami, H., Valadkhani, F., Shapouri, R., Samadi, E. & Haghi, F. Virulence characteristics of
multidrug resistant biofilm forming *Acinetobacter baumannii* isolated from intensive care unit patients.
BMC Infect. Dis. **19**, doi:10.1186/s12879-019-4272-0 (2019).

8   Dijkshoorn, L., Nemec, A. & Seifert, H. An increasing threat in hospitals: multidrug-resistant
Acinetobacter baumannii. Nat. Rev. Microbiol. **5**, 939-951, doi:10.1038/nrmicro1789 (2007).

9   Ryan, K. *et al.* The impact of initial antibiotic treatment failure: Real-world insights in
healthcare-associated or nosocomial pneumonia. *J. Infect.* **77**, 9-17, doi:10.1016/j.jinf.2018.04.002
(2018).

10   Wong, D. *et al.* Clinical and Pathophysiological Overview of *Acinetobacter* Infections: a Century
of Challenges. *Clin. Microbiol. Rev.* **30**, 409-447, doi:10.1128/CMR.00058-16 (2017).

11   Fernandez-Cuenca, F. Relationship between beta-lactamase production, outer membrane protein
and penicillin-binding protein profiles on the activity of carbapenems against clinical isolates of
Acinetobacter baumannii. J. Antimicrob. Chemother. **51**, 565-574, doi:10.1093/jac/dkg097 (2003).

12   Lolans, K., Rice, T. W., Munoz-Price, L. S. & Quinn, J. P. Multicity outbreak of carbapenem-
resistant *Acinetobacter baumannii* isolates producing the carbapenemase OXA-40. *Antimicrob. Agents*
Chemother. **50**, 2941-2945, doi:10.1128/AAC.00116-06 (2006).

13   Geisinger, E. & Isberg, R. R. Antibiotic modulation of capsular exopolysaccharide and virulence
in *Acinetobacter baumannii. PLoS Pathog.* **11**, e1004691, doi:10.1371/journal.ppat.1004691 (2015).

14   Aldred, K. J., Kerns, R. J. & Osheroff, N. Mechanism of quinolone action and resistance.
Biochemistry **53**, 1565-1574, doi:10.1021/bi5000564 (2014).

15   Redgrave, L. S., Sutton, S. B., Webber, M. A. & Piddock, L. J. V. Fluoroquinolone resistance:
mechanisms, impact on bacteria, and role in evolutionary success. *Trends Microbiol.* **22**, 438-445,
doi:10.1016/j.tim.2014.04.007 (2014).

16   Hujer, K. M. *et al.* Rapid Determination of Quinolone Resistance in *Acinetobacter spp. J. Clin.*
Microbiol. **47**, 1436-1442, doi:10.1128/jcm.02380-08 (2009).

17   Munoz-Price, L. S. & Weinstein, R. A. *Acinetobacter* infection. *N. Engl. J. Med.* **358**, 1271-1281,
doi:10.1056/NEJMra070741 (2008).

18   Saroj, S. D., Clemmer, K. M., Bonomo, R. A. & Rather, P. N. Novel Mechanism for
Fluoroquinolone Resistance in *Acinetobacter baumannii. Antimicrob. Agents Chemother.* **56**, 4955-4957,
doi:10.1128/aac.00739-12 (2012).

19   Osińska, A., Harnisz, M. & Korzeniewska, E. Prevalence of plasmid-mediated multidrug
resistance determinants in fluoroquinolone-resistant bacteria isolated from sewage and surface water.
Environ. Sci. Pollut. Res. **23**, 10818-10831, doi:10.1007/s11356-016-6221-4 (2016).

20   Elhosseiny, N. M., Elhezawy, N. B. & Attia, A. S. Comparative proteomics analyses of
Acinetobacter baumannii strains ATCC 17978 and AB5075 reveal the differential role of type II secretion
system secretomes in lung colonization and ciprofloxacin resistance. *Microb. Pathog.* **128**, 20-27,
doi:10.1016/j.micpath.2018.12.039 (2019).

21   Yoon, E. J. *et al.* Contribution of resistance-nodulation-cell division efflux systems to antibiotic
resistance and biofilm formation in *Acinetobacter baumannii. mBio* **6**, doi:10.1128/mBio.00309-15
(2015).

22   Coyne, S., Rosenfeld, N., Lambert, T., Courvalin, P. & Perichon, B. Overexpression of
resistance-nodulation-cell division pump AdeFGH confers multidrug resistance in *Acinetobacter*
baumannii. Antimicrob. Agents Chemother. **54**, 4389-4393, doi:10.1128/AAC.00155-10 (2010).

23   USCAST. The United States Committee on Antimicrobial Susceptibility Testing (USCAST).
Breakpoint tables for interpretation of MIC and zone diameterresults. Version 6.0. (2020).

24   Brauner, A., Fridman, O., Gefen, O. & Balaban, N. Q. Distinguishing between resistance,
tolerance and persistence to antibiotic treatment. *Nat. Rev. Microbiol.* **14**, 320-330,
doi:10.1038/nrmicro.2016.34 (2016).

25   Kussell, E., Kishony, R., Balaban, N. Q. & Leibler, S. Bacterial persistence: a model of survival
in changing environments. *Genetics* **169**, 1807-1814, doi:10.1534/genetics.104.035352 (2005).

26   Balaban, N. Q. Persistence: mechanisms for triggering and enhancing phenotypic variability.
Curr. Opin. Genet. Dev. **21**, 768-775, doi:10.1016/j.gde.2011.10.001 (2011).

27   Balaban, N. Q. *et al.* Definitions and guidelines for research on antibiotic persistence. *Nat. Rev.*
Microbiol. **17**, 441-448, doi:10.1038/s41579-019-0196-3 (2019).

28   Liu, J., Gefen, O., Ronin, I., Bar-Meir, M. & Balaban, N. Q. Effect of tolerance on the evolution
of antibiotic resistance under drug combinations. *Science* **367**, 200-204, doi:10.1126/science.aay3041
(2020).

29   Santos-Lopez, A., Marshall, C. W., Scribner, M. R., Snyder, D. J. & Cooper, V. S. Evolutionary
pathways to antibiotic resistance are dependent upon environmental structure and bacterial lifestyle. *eLife*
8, doi:10.7554/eLife.47612 (2019).

30   Raetz, C. R. H., Reynolds, C. M., Trent, M. S. & Bishop, R. E. Lipid A Modification Systems in
Gram-Negative Bacteria. *Annu. Rev. Biochem.* **76**, 295-329,
doi:10.1146/annurev.biochem.76.010307.145803 (2007).

31   Lathe, R. & Lecocq, J. P. The firA gene, a locus involved in the expression of rifampicin
resistance in *Escherichia coli*. *Mol. Gen. Genet.* **154**, 53-60, doi:10.1007/bf00265576 (1977).

32   Kelly, T. M., Stachula, S. A., Raetz, C. R. & Anderson, M. S. The firA gene of *Escherichia coli*
encodes UDP-3-O-(R-3-hydroxymyristoyl)-glucosamine N-acyltransferase. The third step of endotoxin
biosynthesis. *J. Biol. Chem.* **268**, 19866-19874 (1993).

33   Davis, J. J. *et al.* The PATRIC Bioinformatics Resource Center: expanding data and analysis
capabilities. *Nucleic Acids Res.* doi:10.1093/nar/gkz943 (2019).

34   Buetow, L., Smith, T. K., Dawson, A., Fyffe, S. & Hunter, W. N. Structure and reactivity of
LpxD, the N-acyltransferase of lipid A biosynthesis. *Proc. Nat. Acad. Sci.* **104**, 4321-4326,
doi:10.1073/pnas.0606356104 (2007).

35   Blount, Z. D., Lenski, R. E. & Losos, J. B. Contingency and determinism in evolution: Replaying
life's tape. *Science* **362**, doi:10.1126/science.aam5979 (2018).

36   Melnyk, A. H., Wong, A. & Kassen, R. The fitness costs of antibiotic resistance mutations. *Evol.*
Appl. **8**, 273-283, doi:10.1111/eva.12196 (2014).

37   Subedi, D., Vijay, A. K., Kohli, G. S., Rice, S. A. & Willcox, M. Comparative genomics of
clinical strains of *Pseudomonas aeruginosa* strains isolated from different geographic sites. *Sci. Rep.* **8**,
doi:10.1038/s41598-018-34020-7 (2018).

38   Lieberman, T. D. *et al.* Genetic variation of a bacterial pathogen within individuals with cystic
fibrosis provides a record of selective pressures. *Nat. Genet.* **46**, 82-87, doi:10.1038/ng.2848 (2013).

39   Lieberman, T. D. *et al.* Parallel bacterial evolution within multiple patients identifies candidate
pathogenicity genes. *Nat. Genet.* **43**, 1275-1280, doi:10.1038/ng.997 (2011).

40   Choi, B. *et al.* Persistence and Evolution of SARS-CoV-2 in an Immunocompromised Host. *N.*
Engl. J. Med. **383**, 2291-2293, doi:10.1056/NEJMc2031364 (2020).

41   Honsa, E. S. *et al.* RelA Mutant Enterococcus faecium with Multiantibiotic Tolerance Arising in
an Immunocompromised Host. *mBio* **8**, doi:10.1128/mBio.02124-16 (2017).

42   Ma, C., Yang, X. & Lewis, P. J. Bacterial Transcription as a Target for Antibacterial Drug
Development. *Microbiol. Mol. Biol. Rev.* **80**, 139-160, doi:10.1128/mmbr.00055-15 (2016).

43   van Hoek, A. H. A. M. *et al.* Acquired Antibiotic Resistance Genes: An Overview. *Front.*
Microbiol. **2**, doi:10.3389/fmicb.2011.00203 (2011).

44   Moyed, H. S. & Bertrand, K. P. hipA, a newly recognized gene of *Escherichia coli* K-12 that
affects frequency of persistence after inhibition of murein synthesis. *J. Bacteriol.* **155**, 768-775,
doi:10.1128/JB.155.2.768-775.1983 (1983).

45   Schumacher, M. A. *et al.* HipBA-promoter structures reveal the basis of heritable multidrug
tolerance. *Nature* **524**, 59-64, doi:10.1038/nature14662 (2015).

46   Pontes, M. H., Groisman, E. A. & Garsin, D. A. A Physiological Basis for Nonheritable
Antibiotic Resistance. *mBio* **11**, doi:10.1128/mBio.00817-20 (2020).

47   Trastoy, R. *et al.* Mechanisms of Bacterial Tolerance and Persistence in the Gastrointestinal and
Respiratory Environments. *Clin. Microbiol. Rev.* **31**, doi:10.1128/cmr.00023-18 (2018).

48   Balaban, N. Q., Merrin, J., Chait, R., Kowalik, L. & Leibler, S. Bacterial persistence as a
phenotypic switch. *Science* **305**, 1622-1625, doi:10.1126/science.1099390 (2004).

49   Lewis, K. Persister cells, dormancy and infectious disease. *Nat. Rev. Microbiol.* **5**, 48-56,
doi:10.1038/nrmicro1557 (2007).

50   Shah, D. *et al.* Persisters: a distinct physiological state of *E. coli*. *BMC Microbiol.* **6**, 53,
doi:10.1186/1471-2180-6-53 (2006).

51   Nandakumar, M., Nathan, C. & Rhee, K. Y. Isocitrate lyase mediates broad antibiotic tolerance in
Mycobacterium tuberculosis. *Nat. Commun.* **5**, doi:10.1038/ncomms5306 (2014).

52   Michiels, J. E., Van den Bergh, B., Verstraeten, N. & Michiels, J. Molecular mechanisms and
clinical implications of bacterial persistence. *Drug Resist. Updat.* **29**, 76-89,
doi:10.1016/j.drup.2016.10.002 (2016).

53   Fisher, R. A., Gollan, B. & Helaine, S. Persistent bacterial infections and persister cells. *Nat. Rev.*
Microbiol. **15**, 453-464, doi:10.1038/nrmicro.2017.42 (2017).

54   Van den Bergh, B., Fauvart, M. & Michiels, J. Formation, physiology, ecology, evolution and
clinical importance of bacterial persisters. *FEMS Microbiol. Rev.* **41**, 219-251, doi:10.1093/femsre/fux001
(2017).

55   Yoon, E. J., Courvalin, P. & Grillot-Courvalin, C. RND-type efflux pumps in multidrug-resistant
clinical isolates of *Acinetobacter baumannii*: major role for AdeABC overexpression and AdeRS
mutations. *Antimicrob. Agents Chemother.* **57**, 2989-2995, doi:10.1128/AAC.02556-12 (2013).

56   Pu, Y. *et al.* Enhanced Efflux Activity Facilitates Drug Tolerance in Dormant Bacterial Cells.
Mol. Cell **62**, 284-294, doi:10.1016/j.molcel.2016.03.035 (2016).

57   Harms, A., Maisonneuve, E. & Gerdes, K. Mechanisms of bacterial persistence during stress and
antibiotic exposure. *Science* **354**, doi:10.1126/science.aaf4268 (2016).

58   Frimodt-Møller, J. & Løbner-Olesen, A. Efflux-Pump Upregulation: From Tolerance to High-
level Antibiotic Resistance? *Trends Microbiol.* **27**, 291-293, doi:10.1016/j.tim.2019.01.005 (2019).

59   Peeters, P. *et al.* The impact of initial antibiotic treatment failure: real-world insights in patients
with complicated, health care-associated intra-abdominal infection. *Infect. Drug Resist.* **12**, 329-343,
doi:10.2147/idr.S184116 (2019).

60   Soriano, A., Mensa, J., Meylan, S., Morata, L. & Kuehl, R. When antibiotics fail: a clinical and
microbiological perspective on antibiotic tolerance and persistence of *Staphylococcus aureus*. *J.*
Antimicrob. Chemother. **75**, 1071-1086, doi:10.1093/jac/dkz559 (2020).

61   Meylan, S., Andrews, I. W. & Collins, J. J. Targeting Antibiotic Tolerance, Pathogen by
Pathogen. *Cell* **172**, 1228-1238, doi:10.1016/j.cell.2018.01.037 (2018).

62   Karve, S. *et al.* The impact of initial antibiotic treatment failure: Real-world insights in patients
with complicated urinary tract infection. *J. Infect.* **76**, 121-131, doi:10.1016/j.jinf.2017.11.001 (2018).

63   Wang, E. *et al.* Pathogenesis of Pneumococcal Pneumonia in Cyclophosphamide-Induced
Leukopenia in Mice. *Infect. Immun.* **70**, 4226-4238, doi:10.1128/iai.70.8.4226-4238.2002 (2002).

64   Nielsen, T. B., Yan, J., Luna, B. & Spellberg, B. Murine Oropharyngeal Aspiration Model of
Ventilator-associated and Hospital-acquired Bacterial Pneumonia. *J. Vis. Exp.* doi:10.3791/57672 (2018).

65   Geisinger, E., Mortman, N. J., Vargas-Cuebas, G., Tai, A. K. & Isberg, R. R. A global regulatory
system links virulence and antibiotic resistance to envelope homeostasis in *Acinetobacter baumannii*.
PLoS Pathog. **14**, e1007030, doi:10.1371/journal.ppat.1007030 (2018).

66   Deatherage, D. E. & Barrick, J. E. Identification of mutations in laboratory-evolved microbes
from next-generation sequencing data using breseq. *Methods Mol. Biol.* **1151**, 165-188, doi:10.1007/978-
1-4939-0554-6_12 (2014).

67   Choi, Y., Sims, G. E., Murphy, S., Miller, J. R. & Chan, A. P. Predicting the functional effect of
amino acid substitutions and indels. *PLoS One* **7**, e46688, doi:10.1371/journal.pone.0046688 (2012).

68   Yanisch-Perron, C., Vieira, J. & Messing, J. Improved M13 phage cloning vectors and host
strains: nucleotide sequences of the M13mpl8 and pUC19 vectors. *Gene* **33**, 103-119, doi:10.1016/0378-
1119(85)90120-9 (1985).

69   Andrews, H. L., Vogel, J. P. & Isberg, R. R. Identification of linked *Legionella pneumophila*
genes essential for intracellular growth and evasion of the endocytic pathway. *Infect. Immun.* **66**, 950-958,
doi:10.1128/IAI.66.3.950-958.1998 (1998).

70   Hanahan, D., Jessee, J. & Bloom, F. R. Plasmid transformation of *Escherichia coli* and other
bacteria. *Methods Enzymol.* **204**, 63-113, doi:10.1016/0076-6879(91)04006-a (1991).

71   Kolter, R., Inuzuka, M. & Helinski, D. R. Trans-complementation-dependent replication of a low
molecular weight origin fragment from plasmid R6K. *Cell* **15**, 1199-1208, doi:10.1016/0092-
8674(78)90046-6 (1978).

72   Bouvet, P. J. M. & Grimont, P. A. D. Taxonomy of the Genus *Acinetobacter* with the
Recognition of *Acinetobacter baumannii sp. nov., Acinetobacter haemolyticus sp. nov., Acinetobacter*
johnsonii sp. nov., and *Acinetobacter junii sp. nov*. and Emended Descriptions of *Acinetobacter*
calcoaceticus and *Acinetobacter lwoffii*. *Int. J. Syst. Bacteriol.* **36**, 228-240, doi:10.1099/00207713-36-2-
228 (1986).

Figure 1. Continuous passaging of *A. baumannii* in a lung infection model results in early antibiotic failure in neutrophil-depleted mice. A). The 15-passage strategy using the pneumonia model in both immunocompetent and neutrophil-depleted mice. Neutrophil depletion was induced by two doses of cyclophosphamide at 72- and 24-hrs prior of infection. At the time of infection, *A. baumannii* ATCC 17978 was oropharyngeally inoculated into mice at 10^8 CFU,

and two doses of CIP were administered at 9- and 19-hrs post infection. At 27-hrs post infection,

the mice were euthanized, lungs were homogenized, and bacteria were plated on LB agar plates .

After overnight growth, the bacteria were collected and 10^8 CFU used for the next round of

infection/passage. B). Rapid development of antibiotic failure as a consequence of passage in the

absence of neutrophils. After lungs were homogenized, total CFU on LB agar in the absence of

antibiotic were determined and plotted as function of passage number. Shown are three lines

(L1, L2, L3) for immunocompetent and neutrophil-depleted animals treated with CIP as

described in panel (A).

Figure 2. High level ciprofloxacin resistance evolves in a stepwise fashion during passage in neutrophil-depleted mice. (A,C). After each passage, bacteria from lung homogenates were

serially diluted onto LB agar containing noted concentrations of CIP (μg/ml). The fraction of

CFU on the graded series of CIP plates was determined for each pool and displayed in stacked

bar plots for immunocompetent (A) and neutrophil-depleted (C) mouse infection lineages. The

limit of detection was 2.7E-6. (B,D) The gDNA from each bacterial pool was isolated and

subject to whole genome sequencing to identify the genomic mutations acquired throughout

passaging (after passage 3, 6, 9, 12 and 15). The mutations were detected using a 5% abundance

cutoff and filtered against the parent strain AB17978. The relative abundance of each mutation

was plotted as a function of passage number from (B) immunocompetent lineages and (D)

neutrophil-depleted lineages (D). (E) After 15 passages, the bacterial pool from each lineage

was grown on LB with or without CIP (1 μg/mL) to select for single colonies. Colonies were

purified and subject to whole genome sequencing and MIC assays. In (E), each column

represents a single colony and the rows represent selection criteria, individual mutations and

MIC values. The detected mutations with >90% frequency were aggregated for all colonies and

the presence of each mutation in a certain colony was highlighted in purple. MICs were labeled

and color-coded for all colonies.

Figure 3. *adeL* **mutations drive bacterial persistence through over-expression of the**

AdeFGH pump. (A). Sites of nonsynonymous changes during *in vivo* passaging and domain

structure of AdeL. (B). Locations of nonsynonymous changes on predicted functional domains of

AdeL. (C). Mutations in *adeL* selected during mouse passage result in specific overproduction of

an AdeL-regulated gene *adeG*. Data were determined by qRT-PCR analysis of pump-encoding

components *adeG* (regulated by AdeL) or *adeB* and *adeJ* (components of other pumps) in noted

adeL mutant backgrounds. (D). Overproduction of AdeL-regulated pump components results in

increased persistence in presence of CIP. AB17978 derivatives described in legend were

exposed to CIP at 20X MIC for noted times and titered for viability by quantitating CFUs.

Figure 4. The presence of mutations in *adeL* and *mfsA* is sufficient to explain CIP[R] in

strains derived from mouse passage experiments. A) Two mutations located directly upstream

of *mfsA* selected during mouse passaging experiments. B) Transcription levels of *mfsA* and *rpmL*

were determined by qRT-PCR in strains harboring the two mutations upstream of *mfsA*. ****, p

< 0.0001. C) Double mutants containing *adeL* and *mfsA* mutations show MICs of CIP that are

indistinguishable from evolvedquadruple mutant strain. Data are mean MICs $\pm$ SEM.

Figure 5. The LpxD E117K allele is tightly linked to clinical fluoroquinolone resistance.

Assembled genomes of 4671 clinical isolates were obtained from PATRIC database[33]. A).

Fraction of genomes having the *gyrA*S81L resistance allele. Among the downloaded genomes,

4062 genomes presented the GyrA S81L allele (shown in red), 386 genomes had the same GyrA

allele as ATCC 17978 (shown in green) and the rest had other alleles (shown in black). B) LpxD

sequences from all genomes were aligned and compared to ATCC 17978 and the total number of

variants per defined amino acid was calculated for each group of genomes. The total number of

amino acid changes were normalized and presented as z-scores for each of the three groups, plotted

against the residue numbers. C). Structure of *Chlamydia trachomatis* LpxD protein [34] showing the

presumed site of the *A. baumannii* E117 based on sequence similarity. D). Broth growth of

AB17978 strains differing by single changes in *lpxD*. Shown are strains harboring either the WT,

lpxD T118A119_insA or *lpxD* E117K alleles.

Figure 6. The evolutionary trajectories of isolates acquired throughout passaging in

neutrophil-depleted mice is due to fitness advantage in presence of drug. (A,B) A single

adeL mutant is outcompeted during infection of CIP-treated mice by a quadruple mutant having

drug resistant *adeLmfsA* alleles. The passage 9 pool was oropharyngeally inoculated into

neutrophil-depleted mice in duplicate and passaged three times, subjecting each passage to deep

sequencing. Displayed are relative abundance of all mutations identified in deep sequencing

pools of the duplicate (panels A,B) passaging experiments as a function of the passage number.

C) Competition of purfied *adeL* and quadruple mutant having *adeLmfsA* alleles in presence of

CIP results in predominance of the quadruple mutant. The two strains were mixed at a ratio 95:5

(single:quadruple mutant), and oropharyngeally inoculated into neutrophil-depleted mice in the

presence of CIP through three successive passages. Deep sequencing of the initial inoculum and

each subsequent passage was performed to determine the ratio of the two strains, and the relative

abundance of each mutation is displayed as a function of passage.

Figure 7. Simultaneous selection for increased resistance and fitness. A) Growth rates in

broth culture of noted isolates from passage in mice. B) Neutrophil-replete and C) Neutrophil

depleted (cyclophosphamide treated) BALB/C mice were challenged with noted strain

combination to determine relative competition index of each pair of strains. C.I. values were

determined by C.I. = (strain1/strain2 output ratio)/(strain1/strain2 input ratio) (Materials and

Methods). Data are mean $\pm$ SEM, with significance determine by one-way ANOVA followed by

Multiple Comparison. *p<0.05; ****p<0.0001.

Figure 8. Model of bacterial resistance progression. The antibiotic susceptible inoculum (blue) colonizes poorly in presence of antibiotic. During continued passage with CIP, in the absence of neutrophils, the persister mutants (yellow) arise as first step mutations showing intermediate MICs. With continued passage under the same antibiotic treatment regime, second step mutations arise that have increased fitness in the presence of antibiotic (red) and dominate the population.

925 **Supplemental Figure 1. Bacterial load and resistance levels to CIP after re-passaging.** A)
926 The bacterial load harvested from the lungs after each passage is plotted against the passage
927 number for each lineage. B) CIP resistance level of each pool (after each passage) was assayed
928 via quantification of CFU on LB agar supplemented with CIP.

929 **Table S1. Strains used in this study.**

Clone name	Description	Cipro MIC	reference	Relevant alleles
DH5α	*E. coli:* F⁻ *endA1 glnV44 thi-1 recA1 relA1 gyrA96 deoR nupG purB20* φ80d*lacZ*ΔM15 Δ(*lacZYA-argF*)U169, hsdR17(rK⁻mK⁺), λ⁻	N/A	[68]	N/A
DH5α λ*pir*	DH5α encoding π protein	N/A	[69]	N/A
WT/AB17978	*A. baumannii* ATCC 17978	0.5	[70]	N/A
Ab12	Single colonies selected without CIP from immunocompetent L1: 15 passages	0.5	This study	*ispB* R49H; ΔCP012005
Ab11		0.5	This study	*ispB* R49H; *acx60_02425* V138I
Ab21	Single colonies selected without CIP from immunocompetent L2: 15 passages	0.5	This study	*ispB* R49H; soxF V374*fs*
Ab22		0.5	This study	*adeS* F114L
Ab31	Single colonies selected without CIP from immunocompetent L3: 15 passages	0.5	This study	*alaA* H391fs
Ab33		0.5	This study	None detected
CIPAb11	Single colonies CIP 1 μg/mL selection from immunocompetent L1: 15 passages	4	This study	*acfR* V136D; *adeL* P131H; *lpxD* insA; *mfsA*(-9)
CIPAb12		4	This study	*acfR* V136D; *adeL* P131H; *lpxD* insA; *mfsA*(-9)
CIPAb23	Single colonies CIP 1 μg/mL selection from immunocompetent L2: 15 passages	4	This study	*acfR* V136D; *adeL* P131H; *lpxD* insA; *mfsA*(-9)
CIPAb24		8	This study	*acfR* V136D; *adeL* P131H; *lpxD* insA; *mfsA*(-9)
CIPAb32	Single colonies CIP 1 μg/mL selection from immunocompetent L3: 15 passages	8	This study	*gyrA* S81L
CIPAb34		4	This study	*gyrA* S81L
Cy11	Single colonies selected without CIP from neutrophil-depleted mice lineage 1: 15 passages	4	This study	*gyrB* Q447H; *adeL* C312*
Cy12		4	This study	*gyrB* Q447H; *adeL* C312*
Cy21	Single colonies selected without CIP from neutrophil-depleted mice lineage 2: 15 passages	8	This study	*gyrA* A117E
Cy22		8	This study	*gyrA* A117E

Strain	Description	MIC	Source	Mutations
Cy31	Single colonies selected without CIP from neutrophil-depleted mice lineage 3: 15 passages	4	This study	*acfR* V136D; *adeL* P131H; *lpxD* insA; *mfsA*(-9)
Cy32		4	This study	*acfR* V136D; *adeL* P131H; *lpxD* insA; *mfsA*(-9)
CIPCy11	Single colonies selected with CIP 1 µg/mL from neutrophil-depleted mice lineage 1: 15 passages	4	This study	*gyrB* Q447H; *adeL* C312*
CIPCy13		4	This study	*gyrB* Q447H; *adeL* C312*
CIPCy21	Single colonies selected with CIP 1 µg/mL from neutrophil-depleted mice lineage 2: 15 passages	8	This study	*acfR* V136D; *adeL* P131H; *lpxD* insA; *mfsA*(-9)
CIPCy22		8	This study	*acfR* V136D; *adeL* P131H; *lpxD* insA; *mfsA*(-9)
CIPCy32	Single colonies selected with CIP 1 µg/mL from neutrophil-depleted mice lineage 3: 15 passages	4	This study	*acfR* V136D; *adeL* P131H; *lpxD* insA; *mfsA*(-9)
CIPCy33		2	This study	*acfR* V136D; *adeL* P131H; *lpxD* insA; *mfsA*(-9)
P9Cy2.1	Single colony selected from neutrophil-depleted mice lineage 2: 9 passages	3	This study	*adeL* Y324*
P9Cy3.1	Single colony selected from neutrophil-depleted mice lineage 3: 9 passages	1	This study	*adeL* ΔI335A336
WH_B3	AB17978 *gyrB* Q447H	0.5	This study	*gyrB* Q447H
WH113	AB17978 *adeL* C312*	1	This study	*adeL* C312*
WH114	AB17978 *lpxD* T118_A119insA	0.25	This study	*lpxD* T118_A119insA
WH309	AB17978 *lpxD* E117K	0.25	This study	*lpxD* E117K
WH115	AB17978 *acfR* V136D	0.25	This study	*acfR* V136D
WH_mfsA_48	AB17978 *mfsA*(-9)	0.75	This study	*mfsA*(-9) A->G
WH_mfsA_12	AB17978 *mfsA*(-19)	0.75	This study	*mfsA*(-19) T->A
WH319	AB17978 *mfs*(-9) *adeL* P131H	4	This study	*mfs*(-9); *adeL* P131H
WH316	AB17978 *mfs*(-19) *adeL* P131H	2	This study	*mfs*(-19); *adeL* P131H

JH1	AB17978 *mfsA*(-19) *adeL* ΔI335A336	3	This study	*mfsA*(-19); *adeL* ΔI335A336
WH124	AB17978 *lpxD* insA *acfR* V136D *adeL* P131H	0.75	This study	*lpxD* insA; *acfR* V136D; *adeL* P131H
EHA8	AB17978 *adeL* ΔIA	1	This study	*adeL* ΔI335A336
EHA15	AB17978 *adeL* P131H	1	This study	*adeL* P131H
EHA123	AB17978 *adeL* ΔIA Δ*adeFGH*	1	This study	*adeL* ΔI335A336; Δ*adeFGH*

931 **Table S2. *in vitro* growth rates of LpxD mutants compared to AB17978.**

Time (hr)		AB17978				AB17978::LpxD T118_A119insA				AB17978::LpxD E117K			
Optical density at 600nm	0	0.003	0.004	0.005	0.004	0.003	0.004	0.004	0.004	0.004	0.004	0.004	0.004
	1	0.015	0.015	0.015	0.013	0.009	0.011	0.011	0.011	0.014	0.014	0.014	0.013
	2	0.144	0.138	0.143	0.131	0.054	0.059	0.064	0.067	0.124	0.132	0.129	0.12
	3	1.085	1.01	1.01	0.97	0.335	0.37	0.395	0.43	0.935	0.975	0.985	0.825
	4	2.47	2.68	2.445	2.38	1.16	1.27	1.355	1.39	2.36	2.475	2.475	2.27
	5	3.79	3.82	3.78	3.8	2.81	2.98	2.92	3.12	3.64	3.79	3.64	3.76
	6	4.26	4.23	4.3	4.29	3.56	3.52	3.57	3.81	4.41	4.31	4.33	4.12
	7	4.5	4.59	4.67	4.64	3.95	3.97	3.99	4.09	4.69	4.4	4.47	4.55
	8	4.3	4.29	4.58	4.64	4.34	4.36	4.23	4.28	4.44	4.44	4.49	4.32
Doubling time(min)		**17.325**	**17.479**	**17.667**	**17.258**	**21.579**	**21.904**	**21.563**	**21.165**	**17.927**	**17.511**	**17.546**	**18.005**
R^2		**0.999**	**1**	**0.995**	**0.996**	**1**	**1**	**1**	**0.999**	**0.999**	**0.999**	**0.999**	**0.998**

933 **Table S3. *in vitro* growth rates of the isolates acquired from the *in vivo* passaging experiments.**

Time (hr)		AB17978			AB17978::*adeL* ΔIA			Cy11			Cy21			Cy31		
	0	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001
	1	0.004	0.002	0.004	0.003	0.003	0.003	0.003	0.003	0.003	0.003	0.003	0.003	0.003	0.004	0.003
	2	0.02	0.021	0.025	0.013	0.015	0.014	0.012	0.011	0.011	0.014	0.012	0.013	0.017	0.015	0.015
Optical	3	0.106	0.182	0.201	0.074	0.077	0.055	0.034	0.042	0.03	0.059	0.042	0.049	0.109	0.052	0.094
density	4	0.566	0.712	0.747	0.417	0.416	0.284	0.111	0.235	0.084	0.33	0.22	0.222	0.532	0.272	0.488
at	5	1.6	1.65	1.8	1.22	1.2	0.99	0.65	0.9	0.51	1.02	0.93	1.95	1.23	1.03	1.24
600nm	6	2.03	2.04	2.12	1.44	1.38	1.3	1.06	1.23	0.94	1.35	1.23	1.22	1.55	1.36	1.55
	7	2.3	2.29	2.5	1.69	1.76	1.63	1.38	1.54	1.1	1.57	1.5	1.39	1.88	1.74	1.78
	8	2.56	2.55	2.7	1.85	1.83	1.89	1.51	1.73	1.37	1.78	1.61	1.57	2.01	1.91	2
Doubling time(min)		21.463	17.147	19.693	20.828	21.57	23.843	27.629	23.331	28.985	22.828	23.691	21.038	20.606	24.404	20.682
R^2		0.993	0.986	0.999	0.996	0.997	0.996	0.987	0.992	0.984	0.995	0.99	0.975	0.999	0.984	0.998