An inventory of early branch points in microbial phosphonate biosynthesis

Siwei Li and Geoff P. Horsman*

Department of Chemistry & Biochemistry, Wilfrid Laurier University, Waterloo, ON, Canada N2L 3C5

*Correspondence: ghorsman@wlu.ca

Keywords: phosphonates, microbial, biosynthesis

Abstract

Microbial phosphonate biosynthetic machinery has been identified in ~5% of bacterial genomes and encodes natural products like fosfomycin as well as cell surface decorations. Almost all biological phosphonates originate from the rearrangement of phosphoenolpyruvate (PEP) to phosphonopyruvate (PnPy) catalyzed by PEP mutase (Ppm), and PnPy is often converted to phosphonoacetaldehyde (PnAA) by PnPy decarboxylase (Ppd). Seven enzymes are known or likely to act on either PnPy or PnAA as early branch points en route to diverse biosynthetic outcomes, and these enzymes may be broadly classified into three reaction types: hydride transfer, aminotransfer, and carbon-carbon bond formation. However, the relative abundance of these branch points in microbial phosphonate biosynthesis is unknown. Also unknown is the proportion of *ppm*-containing gene neighborhoods encoding new branch point enzymes and potentially novel phosphonates. In this study we computationally sorted 434 *ppm*-containing gene neighborhoods based on these seven branch point enzymes. Unsurprisingly, the majority (56%) of these pathways encode for production of the common naturally occurring compound 2-aminoethylphosphonate (AEP) or a hydroxylated derivative. The next most abundant genetically encoded intermediates were phosphonoalanine (PnAla, 9.2%), 2-hydroxyethylphosphonate (HEP, 8.5%), and phosphonoacetate (PnAc, 6%). Significantly, about 13% of the gene neighborhoods could not be assigned to any of the seven branch points and may encode novel phosphonates. Sequence similarity network analysis revealed families of unusual gene neighborhoods including possible production of phosphonoacrylate and phosphonofructose, the apparent biosynthetic use of the C-P lyase operon, and a virus-encoded phosphonate. Overall, these results highlight the utility of branch point inventories to identify novel gene neighborhoods and guide future phosphonate discovery efforts.

IMPACT STATEMENT

Microbially-produced phosphonates are relatively rare and underexplored but include medically and agriculturally important molecules like fosfomycin and phosphinothricin, respectively. Because a single enzyme called phosphoenolpyruvate mutase (Ppm) initiates almost all phosphonate production, the composition of the 'gene neighborhood' surrounding a Ppm-encoding gene can inform hypotheses regarding the chemical output of this chromosomal region. After the initial Ppm-catalyzed reaction there are only a limited set of subsequently acting enzymes, or 'branch points', to direct these early-stage phosphonates to alternate chemical fates. However, the relative abundance of different branch points – or the existence of new ones – has not been evaluated. This study provides just such a 'branch point inventory' to determine relative proportions of known branch points and assess the diversity within each branch point. Significantly, this study suggests that a significant proportion (~13%) of gene neighborhoods do not fit into known branch points and therefore may be fertile hunting grounds for new phosphonate biochemistry.

Data Summary

INTRODUCTION

Phosphonic and phosphinic acid metabolites contain a carbon-phosphorus bond and represent a commercially successful but underexplored class of biological molecules exemplified by the antibiotic fosfomycin and the herbicide phosphinothricin.[1] In addition to diffusible small molecules, about half of all microbial phosphonate biosynthetic pathways are thought to produce phosphonoglycan and phosphonolipid cell wall constituents, yet almost nothing is known about their biological significance.[2] Moreover, phosphonate metabolism continues to unveil new and surprising catalytic transformations with strong potential to impact biocatalytic production of fine chemicals.[3, 4]

Almost all known biological phosphonates originate from rearrangement of phosphoenolpyruvate (PEP) to phosphonopyruvate (PnPy) catalyzed by phosphoenolpyruvate mutase (Ppm) (**Figure 1**).[5] Detecting Ppm activity long eluded researchers because PEP is heavily favored at equilibrium, and most phosphonate biosynthetic pathways couple the activity of Ppm with that of a second irreversible decarboxylation to phosphonoacetaldehyde (PnAA) catalyzed by PnPy decarboxylase (Ppd).[6] A typical third step involves formation of the most common phosphonate metabolite 2-aminoethylphosphonate (AEP) by the action of AEP transaminase (AEPT).[7]

Since the 1959[8] discovery of AEP as a ubiquitous biological phosphonate and subsequent identification of its three-enzyme pathway (Ppm-Ppd-AEPT), several additional enzymatic branch points emanating from either PnPy or PnAA have been characterized. In addition to AEPT, three other post-Ppd enzymes have been biochemically verified (**Figure 1, right side**): (i) an aldolase RhiG in the rhizocticin biosynthetic pathway catalyzes decarboxylative aldol condensation of oxaloacetate and PnAA to afford 2-keto-4-hydroxy-5-phosphonopentanoate (KHPnPA);[9] (ii)

Fe-containing alcohol dehydrogenases like PhpC reduce PnAA to 2-hydroxyethylphosphonate
(HEP);[10] and (iii) aldehyde dehydrogenase-catalyzed oxidation of PnAA to phosphonoacetate
(PnAc) has been biochemically characterized for PhnY of the phosphonoalanine (PnAla)
degradation pathway,[11] but is also predicted for the biosynthesis of *O*-phosphonoacetic acid
serine and some phosphonolipids.[2, 12] Although a PhnY-like reaction is also catalyzed by the
α-ketoglutarate-dependent dioxygenase FzmG en route to fosfazinomycins, FzmG and homologs
like PhnY* more commonly catalyze hydroxylation of a phosphonate α-carbon.[13, 14] Three
enzymes are thought to act directly on PnPy and therefore do not require the action of Ppd (**Figure**
1, left side). Of these three, only the aldol-like condensation with acetyl-CoA to generate
phosphonomethylmalate (PnMM) has been biochemically verified for the homocitrate synthase
homologs FrbC and Pfs2.[15, 16] In contrast, aspartate aminotransferase homologs like PalB have
yet to be characterized in vitro.[17, 18] Similarly, activity of the dehydrogenase VlpB has only
been inferred from studies of the valinophos gene cluster in which phosphonolactate (PnLac) was
isolated from spent media.[19] Although the ubiquitous and well-characterized enzymes PalA and
PhnX can also act on PnPy and PnAA respectively, these enzymes are catabolic rather than
biosynthetic. Overall, these seven biosynthetic branch points may be broadly classified as hydride
transfer, amino transfer, or carbon-carbon bond formation.
105
For several decades genome mining has played an increasing role in metabolite discovery, with
particular impact in microbial natural products.[20] New and more accessible tools like
antiSMASH,[21] PRISM,[22] and BiG-SCAPE[23] are available for identifying and classifying
biosynthetic gene clusters (BGCs). These tools are also more generally useful for classifying gene
neighborhoods to glean insights from massive amounts of sequence data, as shown by the recent
application of BiG-SCAPE to the computational characterization of Fe-S flavoenzymes.[24] The
apparently near-universal origin of biological phosphonates from the action of the Ppm enzyme
offers an opportunity for such classification. A landmark investigation of Ppm sequence data
indicated that about 5% of bacterial genomes encode Ppm, and about half are likely to be involved
in phosphonolipid of phosphonoglycan biosynthesis.[2] This study also revealed a strong
correlation between gene neighborhood and Ppm sequence. Ppm sequence diversity has since been
used to guide the discovery of several new phosphonates including valinophos and
phosphonoalamides, illustrating the utility of genome mining for phosphonate discovery.[18, 19]
119
Despite the success of Ppm sequence-based discovery, several general features of early-stage
phosphonate biosynthesis remain unclear. Specifically, the relative abundance of each biosynthetic
branch point remains unknown. Moreover, *ppm*-containing gene neighborhoods that do not encode
any of the seven branch points may represent fertile hunting grounds for new phosphonate
chemistry. Here we employ computational tools to inventory these seven phosphonate biosynthetic
branch points in *ppm*-containing gene neighborhoods from NCBI RefSeq complete bacterial
genomes, as well as archaea and viruses. The most common predicted intermediates include AEP
(56%), PnAla (9.2%), HEP (8.5%), and PnAc (6.2%). Significantly, ~13% of genomes examined
encode none of these seven gateway enzymes and may generate novel phosphonates like
phosphonofructose and phosphonoacrylate. Other surprises include the apparent biosynthetic use
of C-P lyase genes and a virus-encoded AEP pathway.
131
132
133 **METHODS**

Identification of Ppm-encoding genes

Genomes were downloaded from the National Center for Biotechnology Information (NCBI) RefSeq database in June 2020, including complete bacteria (17,258), archaea (1,077), fungi (326), protists (97), plants (124), and animals (556) (**Table S1**). In addition we scanned 364 'huge phage' genomes[25] and 501 draft contig metagenome-assembled genomes (MAGs) of Nucleo-Cytoplasmic Large DNA viruses (NCLDVs)[26]. Phosphoenolpyruvate mutase (Ppm) protein sequences were identified by hmmsearch (HMMER 3.1b2, February 2015; http://hmmer.org) for the Pfam family PEP_mutase (PF13714.7). Hits that aligned with at least 70% of profile HMMs were included. Sequences were then filtered for the presence of the Ppm-specific $EDKX_5NS$ motif [27] using the Perl script *ps_scan.pl* version 1.86 [28].

Collection and network analysis of phosphonate biosynthetic gene neighborhoods

Protein accession numbers were used to extract the corresponding RefSeq genome assembly accession numbers from feature tables and then download the full RefSeq assembly files in Genbank format for each identified assembly. Genome coordinates were extracted from each feature table to include *ppm* and five flanking genes on each side, and corresponding nucleotide accession numbers were used to download genbank files via batch entrez. A Python script was used to take the identified coordinates and extract each gene neighborhood from full GenBank files and output a collection of gene neighborhood GenBank files as input for BiG-SCAPE.[23] The files used for BiG-SCAPE analysis comprised *ppm* gene neighborhoods of 869 complete bacterial genomes, 12 archaea, and 1 virus, for a total of 882 gene neighborhoods.

Phylogenetic tree construction

The 424 bacterial Ppm protein sequences, the 12 Ppm sequences from the archaeal gene neighborhoods, and the single viral Ppm sequence (**Table S1**) were aligned against the PF13714 HMM using hmmalign. After removing unaligned and indel regions we were left with a 91 amino acid alignment of 432 Ppm sequences. A phylogenetic tree was constructed with FastTree [29] and the midpoint rooted tree was constructed using the phytools package in R [30]. The phylogenetic tree was visualized and annotated using the interactive Tree of Life (iTOL) [31].

Classifying Ppm-encoding neighborhoods based on early biosynthetic branch point

The 882 gene neighborhoods were also classified according to their predicted early biosynthetic branch points following the Ppm-catalyzed formation of PnPy common to all neighborhoods. This classification was performed based on the presence of the key enzymes outlined in **Figure 1**, and PFAM domains for each were identified by hmmscan (HMMER 3.1b2, February 2015; http://hmmer.org). Three of these enzymes act on PnPy and therefore only need Ppm co-encoded in the gene neighborhood (PFAM in parentheses): aminotransferase PalB (Aminotran_1_2 PF00155.22) to afford PnAla[17, 18], phosphonomethylmalate synthase FrbC (HMGL-like PF00682.20)[15], and short-chain dehydrogenase VlpB (2-Hacid_dh_C PF02826.2) to generate PnLac[19]. The remaining four enzymes act after PnPy decarboxylase (Ppd, PF002775 or PF002776)[6]: AEP transaminase PhnW (Aminotran_5 PF00266.20)[32], aldolase RhiG (HMGL-like PF00682.20)[9], phosphonoacetaldehyde dehydrogenase PhnY (Aldedh PF00171.23)[11], and phosphonoacetaldehyde reductase PhpC (Fe-ADH PF00465.20)[33].

Each pathway was classified according to the presence and absence of encoded enzymes (in addition to Ppm) required for each of the seven metabolic outcomes shown in **Figure 1**. For

example, a gene neighborhood was classified as 'AEPT' if it encoded (in addition to Ppm) Ppd
and AEPT but did not encode the competing RhiG, PhpC, and PhnY homologs. A series of Python
scripts were written to break each gene into individual protein sequences, use hmmscan to find all
PFAM domains and assign them to each gene neighbourhood, identify the presence/absence of
defined enzyme domains, and match each gene neighborhood to its corresponding BiG-SCAPE
gene cluster family (GCF) number. The network was visualized in Cytoscape and manually
curated for accurate branch point assignment of GCFs. The branch point inventory shown in Figure
2B was constructed by treating all BGCs possessing the same Ppm sequence as a single data point,
such that a total of 434 rather than 822 BGCs are represented.

189
190

RESULTS AND DISCUSSION
The microbial Ppm sequence inventory is dominated by *Burkholderia*
Consistent with previous work[2] we identified at least one Ppm-encoding gene in 4.2% of
complete bacterial genomes, 1.6% of archaea, 12% of protists, and 3.8% of animals (**Table S1**).
Although fungal phosphonates have been reported[34], we could not find Ppm sequences in fungal
genomes, nor could we identify Ppm encoded by plant or viral genomes in the NCBI RefSeq
database. Surprisingly, we detected Ppm encoded by one contig of a metagenomically-assembled
genome of a Nucleo-Cytoplasmic Large DNA virus (NCLDV).[26] The vast majority of Ppm
sequences (424 of 490) were bacterial, and many occurred in multiple genomes for a total of 869
Ppm-encoding gene neighborhoods in 723 bacterial genomes. Although most Ppm sequences were
found in a single genome, several were found in multiple genomes. In particular, two Ppm
sequences (WP_004523200 and WP_004525540) together accounted for over 200 gene
neighborhoods in different genome assemblies of *Burkholderia pseudomallei*; however, the
phylogenetic separation of these two sequences in distinct clades implies different phosphonate
products (**Figure 2A**).[2] We also noted the previously-described abundance of Proteobacteria and
in particular the genus *Burkholderia*, which comprised almost a quarter of the 424 bacterial Ppm
sequences analyzed (**Figure 2A**).

Gene neighborhoods predominantly encode transamination and hydride transfer
We classified 882 *ppm*-containing gene neighborhoods (*ppm* plus five genes on each side)
according to the presence of the known early-stage phosphonate biosynthetic genes shown in
Figure 1. After treating gene neighborhoods possessing the same Ppm sequence as a single data
point (e.g. all 103 neighborhoods possessing the Ppm sequence WP_004523200 were counted as
a single pathway leading to PnAc), we were left with 434 gene neighborhoods. The results of this
classification are summarized in **Figure 2B**. As expected, *ppd* is found in the vast majority of the
gene neighborhoods studied, indicating that ~83% of phosphonate biosynthetic pathways proceed
through the intermediacy of PnAA. The next most common transformation of PnPy (in 9.2% of
gene neighborhoods) is predicted to be catalyzed by PalB-like aminotransferases (PF00155
'Aminotran_1_2') to afford PnAla. Although PalB belongs to a degradative pathway,[17] a family
of homologous aminotransferases were recently implicated in phosphonoalamide
biosynthesis.[18] The citrate synthase-like aldol reaction catalyzed by FrbC homologs as well as
the VlpB-like reduction to PnLac together accounted for only 5.3% of the gene neighborhoods.
Also as expected, the vast majority of post-Ppd reactions involve transamination catalyzed by
AEPT, which is an aminotransferase (PF00266 'Aminotran_5') distinct from PalB homologs.
Overall, AEP is predicted to arise as an intermediate in more than half of all phosphonate

biosynthetic pathways. The next most common known intermediate appears to be HEP, which
accounts for 8.5% of neighborhoods. PhnY-generated PnAc was found in 6.2% of gene
neighborhoods and putative RhiG-like aldol reactions in only 1.6%. The relative scarcity of HEP
was surprising and may reflect an underrepresentation of marine sequences; HEP occurs in several
natural product biosynthetic pathways (fosfomycin, phosphinothricin, dehydrophos, argolaphos),
as a major polysaccharide component of marine dissolved organic matter [35], and a precursor to
methylphosphonate en route to methane production in the ocean.[36] In summary, most gene
neighborhoods encoded for early-stage amino (~65%) or hydride (~16%) transfer, with minor
contributions from aldol-type carbon-carbon bond-forming reactions (~5%). The remaining ~13%
of gene neighborhoods did not possess any of the known genes depicted in **Figure 1**, suggesting
possible novel transformations and phosphonate products.
237
Network analysis reveals several large gene cluster families
Although the above analysis classified key early branch points in phosphonate biosynthetic gene
neighborhoods, it did not capture diversity within each group. For example, how varied are the
gene neighborhoods comprising the 56% that appear capable of making AEP? To address this
question we used BiG-SCAPE network analysis to classify the 882 gene neighborhoods into 85
different gene cluster families (GCFs, **Figure 3**).
244
Half of the neighborhoods evaluated (444 of 882) belonged to AEPT- or PhnY-encoding
Burkholderia-dominated gene neighborhoods, represented respectively by the
'Burkholderiales_CDP-OH' and 'Burkholderia_TauD' groups of GCFs (boxes in **Figure 3**). These
larger groups respectively correspond to the Group 1 and 2 phosphonolipids described by Yu et
al.[2] and are predicted to produce the 1-hydroxy substituted phosphonates shown in **Figure 4**.
This predominance of *Burkholderia* may reflect database abundance due to medical importance
rather than natural occurrence; for example, most of the 126 neighborhoods of GCFs 877 and 817
appear to be nearly identical strains of *Burkholderia pseudomallei*, the causative pathogen of
meliodosis.[37] Furthermore, these two abundant groups contained the two Ppm sequences found
in over 200 mostly *Burkholderia pseudomallei* gene neighborhoods (large bars in **Figure 2A**):
WP_004523200 occurred in the majority of nearly identical PhnY-containing gene neighborhoods
of GCFs 817 and 877 that also encoded a TauD homolog (**Figure 4 inset**); WP_004525540
occurred in the majority of near-identical AEPT-encoding neighborhoods of GCFs 878 and 818.
The latter two neighborhoods belong to a larger 'Burkholderiales_CDP-OH' group of twelve
GCFs totaling 318 (36%) slightly more diverse neighborhoods. As shown in **Figure 4**, this group
has a common seven-gene core that encodes, in addition to the three AEP-producing enzymes
Ppm, Ppd, and AEPT: (i) a putative α-ketoglutarate-dependent iron(II) oxygenase, (ii) two
phosphonyl tailoring cytidylyltransferases (PntCs), one of which is fused to Ppm as often observed
in phosphonate biosynthetic gene clusters,[38] and (iii) two genes associated with phospholipid
biosynthesis. Interestingly, despite their prevalence and detection of *ppm* gene expression,[39]
phosphonolipids have not been characterized from either of these large groups; however, 1-
hydroxy-AEP is attached to a phosphonosphingolipid from *Bacteriovorax stolpii* (formerly
Bdellovibrio stolpii), but the gene cluster belongs to the more distantly related AEPT-encoding
GCF 761 (**Figures 3 and 5**).[40]
269
270 **The large group of AEPT-encoding gene neighborhoods is diverse**

Apart from the Group 1 'Burkholderiales_CDP-OH' phosphonolipids, the AEPT-containing gene neighborhoods are diverse (**Figure 5**). These neighborhoods include those encoding two structurally characterized phosphonates: the above-mentioned *B. stolpii* phosphonosphingolipid [40] and the phosphonoglycan component of the polysaccharide B virulence factor from *Bacteroides fragilis*.[41] Also included are the *Selenomonas* species of GCF 400 that encode multiple sulfatases and correspond to the previously assigned Group 4 phosphonolipids.[2] As evident from the similarity of aligned genes in **Figure 5**, several gene cluster families are related. For example, GCFs 841, 837, 805, and 453 are all Paenibacillaceae encoding two ATP-grasp amino acid ligases flanking the AEP biosynthetic core and therefore may produce a peptide. The ATP-grasp and Mur ligase domains of GCFs 859 and 840 suggest phosphonylated peptidoglycan. Although GCFs 864 and 857, exemplified by the human pathogen *Salmonella enterica*, possess the classical three-gene AEP core, they also encode an aminotransferase (PF00155) associated with PnAla production.[18] Squalene synthase and cyclases encoded by *Nitrosococcus* species of GCF 731 hint at a possible phosphonoterpene product as previously speculated.[42] This comparison of AEPT-encoding gene clusters also showcases diverse gene fusions. For example, Ppm can be encoded alone (GCFs 764 and 753) or as a fusion to a PntC cytidylyltransferase (e.g. PF12804 in GCF 761, or PF01467 in GCF 743) or Ppd (GCFs 333 and 881). Curiously, GCF 881 is a single gene neighborhood from a metagenomically assembled genome of a Nucleo-Cytoplasmic Large DNA Virus (NCLDV).[26] Immediately downstream of the AEP core are genes implicated in aminosugar synthesis consistent with production of a viral phosphonoglycan.

The unexplored diversity of PhnY-, PalB-, and PhpC-encoding pathways

Phosphonoacetate (PnAc) is an intermediate in only two characterized biosynthetic pathways: the fosfazinomycins [13] and *O*-phosphonoacetic acid serine[12]. In fosfazinomycin biosynthesis the transformation of PnAA to PnAc is probably due to the activity of FzmG, an a-ketoglutarate Fe(II)-dependent oxygenase whose primary role in this pathway is hydroxylation at C-1 of methylated PnAc.[13] The more common route to PnAc is via PhnY-like aldehyde dehydrogenases that have been characterized from phosphonate degradation pathways.[11] Evidence for biosynthetic roles for PhnY-like enzymes has been found in: (i) the PhpJ transformation of phosphonoformaldehyde to phosphonoformate in phosphintothricin biosynthesis[33], and (ii) PnAc formation in *O*-phosphonoacetic acid serine biosynthesis in *Streptomyces* sp. NRRL F-525.[12] The latter gene neighborhood is shown in **Figure S1A** along with other GCFs encoding PhnY-like aldehyde dehydrogenases, highlighting diversity and some interesting features. For example, the actinomycete-dominated GCF 846 and *Bradyrhizobium* species in GCF 829 encode non-ribosomal peptide synthetase (NRPS) systems within 15 genes of *ppm*, and the former encodes putative adenylation and thiolation domains within two genes of the encoded aldehyde dehydrogenase. Plant pathogens like *Xanthomonas vasicola* dominate GCFs 820 and 861, each of which possess a seven-gene core encoding an AurF-like *N*-oxygenase (PF11583), glutamine amidotransferase and asparagine synthase.

PalB-encoding families also very diverse and include GCF 870 of *Pseudomonas protegens* strains encoding NRPS systems within two genes of *ppm* (**Figure S1B**). Notably, none of these families represent the previously discovered group of phosphonalamides produced by various *Streptomyces* species and isolated from *Streptomyces* sp. NRRL B-2790, which are the only known phosphonate biosynthetic products of PalB-like activity.[18] The absence of the known phosphonoalamide gene clusters highlights the limitations of the current study using only bacterial complete genomes.

The formation of HEP catalyzed by PhpC-like iron-containing alcohol dehydrogenases appears in diverse families of gene neighborhoods that include pathogenic bacteria like *Helicobacter cinaedi* and *Campylobacter jejuni* (**Figure S1C**). Although this transformation is important in several *Streptomyces* natural products like fosfomycin, phosphinothricin, dehydrophos, and argolaphos,[1] none of the encoding gene clusters were found in the NCBI complete genome database. Of the ten gene cluster families identified, only two have yielded isolated phosphonate products: the methylphosphonate adduct from the marine archaeon *Nitrosopumilus maritimus*,[36] and the HEP conjugate purified from exopolysaccharide of *Stackebrandtia nassauensis*.[43] Interestingly, the two *Streptomyces rimosus* strains comprising GCF 782 encode not only a PhpC homolog, but also VlpB and PhnY homologs. This is one example highlighting the ambiguities resulting from our classification system because this cluster could be attributed to any number of groups and therefore can benefit from manual curation. Specifically, the presence of a possible 2-hydroxyethylphosphonate dioxygenase (HEPD) homolog of PhpD implies intermediacy of HEP in a manner reminiscent of phosphinothricin biosynthesis in *Streptomyces viridochromogenes*. In this case HEP would be oxidized by PhpC to phosphonoformaldehyde followed by PhnY-catalyzed formation of phosphonoformate analogous to the proposed action of PhpJ.[33]

VlpB-like dehyrogenases: beyond valinophos

To date PnLac has only been isolated in spent media of the valinophos producer *Streptomyces durhamensis* B-3309, and the dehydrogenase VlpB was proposed to catalyze its formation from PnPy.[19] Very similar gene clusters are found in two other *Streptomyces* strains of GCF 838: *S. atratus* strain SCSIO_ZH16 and *Streptomyces* sp. WAC 01529. Both of these clusters possess the *vlpABCDE* core proposed to generate 2,3-dihydroxypropylphosphonic acid (DHPPA) (**Figure S2**). Three other members of this family, all Proteobacteria, expand upon the five-gene valinophos core. The lone Firmicutes, *Tumebacillus algifaecus*, possesses a unique gene cluster that may encode the novel compound phosphonofructose, an analog of the sulfoglycolytic intermediate sulfofructose (**Figure S3**).[44]

The aldolases: FrbC- and RhiG-encoding pathways

The GCFs encoding Ppm, Ppd, and FrbC en route to the intermediate PnMM diverge from the standard FR-900098-like pathway logic of GCF 867 (**Figure 6A**) to include more unusual gene combinations in GCFs 352 and 630 (**Figure S4**). Like the FR-900098 gene cluster, GCF 867 encodes aconitase and isocitrate dehydrogenase homologs en route to the α-ketoglutarate analog 2-oxo-4-phosphonobutyrate.[15] Most GCF 867 gene clusters do not encode an aminotransferase, but like FR-900098 biosynthesis may nonetheless generate the glutamate analog 2-amino-4-phosphonobutyrate via the action of a promiscuous enzyme encoded elsewhere in the genome. However, we identified a plasmid of *Agrobacterium tumefaciens* strain CFBP6625 that also encodes a PalB-like aminotransferase and therefore may facilitate transamination to 2-amino-4-phosphonobutyrate (**Figure 6A**). The *Burkholderia*-dominated GCF 352 is the largest of the four FrbC GCFs and possesses an unusual gene combination including two dioxygenases, lumazine synthase, and a member of the beta-keto acid cleavage enzyme (BKACE) family.[45] The *Photorhabdus* GCF 630 neighborhoods encode aconitase and luciferase-like monooxygenase, an FMN-dependent enzyme found in natural product biosynthetic pathways such as mensacarcin and neoabyssomicin.[46, 47]

GCFs 764 and 753 (**Figure S4**) are very similar and were assigned as RhiG aldolase groups due
to the tight clustering of genes in the order *ppd-rhiG-ppm* and the inclusion of the known
rhizocticins producer *Bacillus subtilis* ATCC6633 in this group.[9] One distinguishing feature of
these rhizocticin-like gene clusters is the apparent separation of Ppd domains (PF2775 and
PF2776) onto individual proteins, which is occasionally observed in other gene clusters such as
the *O*-phosphonoacetic acid serine cluster from PnAc-producing *Streptomyces* sp. strain NRRL F-
525 (**Figure S1A**) [12]. The presence of an HMGL-like domain in the *Streptococcus pneumoniae*
clusters of GCF 872 led to assignment as possible RhiG-encoding pathways, with an unusual tRNA
synthetase and phosphopantetheine attachment site suggesting a peptidyl product (**Figure S5**).
Notably this neighborhood is conserved across only ten of 81 complete NCBI RefSeq *S.*
pneumoniae genomes, and the flanking recombinase/relaxase genes define potential gene cluster
boundaries.
375
Discovery potential in unclassified neighborhoods
Fifteen gene cluster families could not be assigned because they did not encode any of the seven
branch point enzymes acting on either PnPy or PnAA shown in **Figure 1**. Most of these GCFs
encoded Ppd ('post-Ppd other' in **Figure 3**) but several did not ('pre-Ppd other' in **Figure 3**).
Surveying these unclassified families revealed several unusual features. For example, GCF 842
neighborhoods of *Paenibacillus larvae* encode Ppm flanked by malate dehydrogenase and
fumarase (**Figure 6B**). Because PnPy is an analog of the citric acid cycle intermediate
oxaloacetate, we predict that malate dehydrogenase will catalyze a VlpB-like reduction to PnLac
followed by fumarase-catalyzed dehydration to the fumarate analog phosphonoacrylate. The
adjacent ATP-grasp domains imply peptide bond formation with an amino acid as seen for the
phosphonoalamides.[18] Interestingly, directing PnPy in the fumarase direction of the TCA cycle
would contrast with the citrate synthase direction observed in FR-900098 biosynthesis and in
FrbC-encoding families such as GCF 867 (**Figure 6**). GCF 540 encodes Ppm that is unusually
flanked by thymidylate kinase and an uncharacterized nucleotidyltransferase (PF14907), while
actinomycete-dominated GCF 699 has several gene neighborhoods encoding a PLP-dependent
enzyme (PF00291) that may also catalyze a PalB-like transamination to PnAla (**Figure S4**).
392
Most of the uncharacterized diversity occurs in gene neighborhoods that encode Ppm and Ppd and
therefore likely proceed through the intermediacy of PnAA (**Figure S4**). This includes the unusual
occurrence of the C-P lyase *phn* operon[48]: GCFs 573, 708, and 849 encode Ppm, Ppd, and a
possible dehydrogenase followed on the same strand by PhnGHIJKLMNP (**Figure S6**). The
absence of both the PhnCDE transporter and the PhnO *N*-acetylation enzyme imply a function
other than AEP import and degradation. The presence of an NAD(P)-dependent oxidoreductase
after Ppm-Ppd implies HEP or PnAc formation using atypical enzymes. Several gene clusters in
GCF 537 encode a hotdog-fold enzyme associated with A-factor biosynthesis, a
phosphopantetheine attachment site, and asparagine synthase. Neighborhoods dominated by
Clostridium botulinum and *C. tetani* in GCF 599 encode a methyltransferase and sugar
biosynthetic enzymes. In *Campylobacter* strains of GCF 385, the *ppm-ppd* core is flanked by genes
predicted to encode radical SAM and adenylsulfate kinase enzymes. *Fibrobacter succinogenes* of
GCF 60 is known to produce an exopolysaccharide possessing a novel *N*-hydroxyethyl derivative
of AEP.[49] However, the absence of AEPT suggests that AEP is not an intermediate but rather
favors the reductive amination of PnPy with ethanolamine. The *Streptomyces* of GCF 742 possess
a unique combination of polyprenyl synthetase, cytochrome P450, methyltransferase, and a TauD-

like oxygenase. Archaea of GCFs 869 and 685 have diverse genes surrounding *ppm* and *ppd*,
including calcineurin-like phosphoesterase (PF12850) and GDP-mannose 4,6-dehydratase
(PF16383) domains in common with *Fibrobacter succinogenes*.
412
Genomes with multiple Ppm-encoding neighborhoods
The vast majority of the 726 *ppm*-containing bacterial genomes evaluated possess either one
(~80% of genomes) or two (~18% of genomes) *ppm*-containing gene neighborhoods, but several
maintained three or four (SI file SI_multiply_ppm.xls), as previously noted[2]. Apart from a single
Streptomyces genome (*S. griseochromogenes* ATCC 14511) with three neighborhoods, all
genomes encoding more than two Ppm were *Burkholderia*, and four of the five genomes with four
ppm genes were strains of *B. oklahomensis*. An analysis of these *Burkholderia* Ppm gene
neighborhoods revealed 151 genomes with a single *ppm*; 119, 3, and 5 genomes housed two, three,
and four *ppm*, respectively (**Figure S7**). All genomes encoded AEPT-containing GCFs belonging
to the large 'Burkholderiales CDP-OH family' dominated by GCFs 818 and 878, but a clear
combinatorial progression emerges with additional clusters. Specifically, the second cluster is
always PhnY-containing GCFs 877 or 817, the third cluster belongs to PalB-like GCF 362, and
the fourth cluster is GCF 352 that we assigned as FrbC-like (**Figure S7**). The significance of this
expanded phosphonate biosynthetic capacity in *Burkholderia* is unclear, but the consistent
accumulation of the same four classes of phosphonate biosynthetic gene cluster is intriguing.
428
Conclusion
Phosphonates are an important but niche class of natural products and increasingly appreciated as
important cell surface modifications in the form of phosphonolipids and phosphonoglycans.
Because most biological phosphonates originate via Ppm catalysis, and Ppm sequence phylogeny
correlates with biosynthetic product[2], discovery of new phosphonates has largely focused on
ppm-guided actinomycete strain prioritization[19]. However, less is known about the diversity of
ppm gene neighborhoods and how they might be classified in order to identify and prioritize
overlooked gene clusters. Specifically, little is understood about the relative abundance of different
branch points in the early steps of phosphonate biosynthesis. This study provides an inventory of
these early branch points and highlights potential avenues of discovery. These highlights include:
(i) a possible phosphonoacrylate intermediate, (ii) integration of the degradative C-P lyase (*phn*
operon) components into biosynthetic gene clusters, (iii) a possible phosphonofructose
biosynthetic pathway, and (iv) a virus-encoded phosphonate, which hints at the broader possibility
of virus-encoded natural products. In summary, these results help contextualize current knowledge
of early branch points of phosphonate biosynthesis and provide a framework to generate
hypotheses and triage discovery efforts.
445
446

447 **Acknowledgements**
448 We thank Monica Papinski for providing Python functions that we modified and incorporated into
449 final scripts.

450
451 **Conflicts of Interest**
452 The authors declare that there are no conflicts of interest.
453

454 **Funding Information**

455   This work was supported by the Natural Sciences and Engineering Research Council of Canada
456   (NSERC) in the form of a Discovery Grant (G.P.H.) and an Undergraduate Student Research
457   Award (S.L.).

REFERENCES

1. **Horsman GP, Zechel DL**. Phosphonate Biochemistry. *Chem Rev* 2017;117:5704–5783.

2. **Yu X, Doroghazi JR, Janga SC, Zhang JK, Circello B, et al.** Diversity and abundance of phosphonate biosynthetic genes in nature. *Proc Natl Acad Sci USA* 2013;110:20759–20764.

3. **Metcalf WW, van der Donk WA**. Biosynthesis of phosphonic and phosphinic acid natural products. *Annu Rev Biochem* 2009;78:65–94.

4. **Parkinson EI, Erb A, Eliot AC, Ju K-S, Metcalf WW**. Fosmidomycin biosynthesis diverges from related phosphonate natural products. *Nat Chem Biol* 2019;15:1049–1056.

5. **Bowman Elise, McQueney Michael, Barry RJ, Dunaway-Mariano Debra**. Catalysis and thermodynamics of the phosphoenolpyruvate/phosphonopyruvate rearrangement. Entry into the phosphonate class of naturally occurring organophosphorus compounds. *J Am Chem Soc* 1988;110:5575–5576.

6. **Johnen S, Sprenger GA**. Characterization of recombinant thiamine diphosphate-dependent phosphonopyruvate decarboxylase from Streptomyces viridochromogenes Tü494. *Journal of Molecular Catalysis B: Enzymatic* 2009;61:39–46.

7. **Chen CCH, Zhang H, Kim AD, Howard A, Sheldrick GM, et al.** Degradation pathway of the phosphonate ciliatine: crystal structure of 2-aminoethylphosphonate transaminase. *Biochemistry* 2002;41:13162–13169.

8. **Horiguchi M, Kandatstu M**. Isolation of 2-Aminoethane Phosphonic Acid from Rumen Protozoa. *Nature* 1959;184:901–902.

9. **Borisova SA, Circello BT, Zhang JK, van der Donk WA, Metcalf WW**. Biosynthesis of Rhizocticins, Antifungal Phosphonate Oligopeptides Produced by Bacillus subtilis ATCC6633. *Chem Biol* 2010;17:28–37.

10. **Shao Z, Blodgett JAV, Circello BT, Eliot AC, Woodyer R, et al.** Biosynthesis of 2-hydroxyethylphosphonate, an unexpected intermediate common to multiple phosphonate biosynthetic pathways. *J Biol Chem* 2008;283:23161–23168.

11. **Agarwal V, Peck SC, Chen J-H, Borisova SA, Chekan JR, et al.** Structure and function of phosphonoacetaldehyde dehydrogenase: the missing link in phosphonoacetate formation. *Chem Biol* 2014;21:125–135.

12. **Freestone TS, Ju K-S, Wang B, Zhao H**. Discovery of a Phosphonoacetic Acid Derived Natural Product by Pathway Refactoring. *ACS Synth Biol* 2017;6:217–223.

13. **Huang Z, Wang K-KA, Lee J, van der Donk WA**. Biosynthesis of fosfazinomycin is a convergent process. *Chem Sci* 2015;6:1282–1287.

14.   **McSorley FR, Wyatt PB, Martinez A, DeLong EF, Hove-Jensen B, *et al.*** PhnY and
PhnZ Comprise a New Oxidative Pathway for Enzymatic Cleavage of a Carbon–Phosphorus
Bond. *J Am Chem Soc* 2012;134:8364–8367.

15.   **Eliot AC, Griffin BM, Thomas PM, Johannes TW, Kelleher NL, *et al.*** Cloning,
Expression, and Biochemical Characterization of Streptomyces rubellomurinus Genes
Required for Biosynthesis of Antimalarial Compound FR900098. *Chem Biol* 2008;15:765–
770.

16.   **Kim SY, Ju K-S, Metcalf WW, Evans BS, Kuzuyama T, *et al.*** Different biosynthetic
pathways to fosfomycin in Pseudomonas syringae and Streptomyces species. *Antimicrob
Agents Chemother* 2012;56:4175–4183.

17.   **Kulakova AN, Kulakov LA, Villarreal-Chiu JF, Gilbert JA, McGrath JW, *et al.***
Expression of the phosphonoalanine-degradative gene cluster from Variovorax sp. Pal2 is
induced by growth on phosphonoalanine and phosphonopyruvate. *FEMS Microbiol Lett*
2009;292:100–106.

18.   **Kayrouz CM, Zhang Y, Pham TM, Ju K-S**. Genome Mining Reveals the
Phosphonoalamide Natural Products and a New Route in Phosphonic Acid Biosynthesis. *ACS
Chem Biol* 2020;15:1921–1929.

19.   **Ju K-S, Gao J, Doroghazi JR, Wang K-KA, Thibodeaux CJ, *et al.*** Discovery of
phosphonic acid natural products by mining the genomes of 10,000 actinomycetes. *Proc Natl
Acad Sci U S A* 2015;112:12175–12180.

20.   **Ziemert N, Weber T, Medema MH**. 6.03 - Genome Mining Approaches to Bacterial
Natural Product Discovery. In: Liu H-W (Ben), Begley TP (editors). *Comprehensive Natural
Products III*. Oxford: Elsevier. pp. 19–33.

21.   **Blin K, Shaw S, Steinke K, Villebro R, Ziemert N, *et al.*** antiSMASH 5.0: updates to
the secondary metabolite genome mining pipeline. *Nucleic Acids Research* 2019;47:W81–
W87.

22.   **Skinnider MA, Dejong CA, Rees PN, Johnston CW, Li H, *et al.*** Genomes to natural
products PRediction Informatics for Secondary Metabolomes (PRISM). *Nucleic Acids
Research* 2015;43:9645–9662.

23.   **Navarro-Muñoz JC, Selem-Mojica N, Mullowney MW, Kautsar SA, Tryon JH, *et al.***
A computational framework to explore large-scale biosynthetic diversity. *Nature Chemical
Biology* 2020;16:60–68.

24.   **Pascal Andreu V, Fischbach MA, Medema MH**. Computational genomic discovery of
diverse gene clusters harbouring Fe-S flavoenzymes in anaerobic gut microbiota. *Microb
Genom*;6. Epub ahead of print May 2020. DOI: 10.1099/mgen.0.000373.

25.   **Al-Shayeb B, Sachdeva R, Chen L-X, Ward F, Munk P, *et al.*** Clades of huge phages
from across Earth's ecosystems. *Nature* 2020;578:425–431.

26.   **Moniruzzaman M, Martinez-Gutierrez CA, Weinheimer AR, Aylward FO**. Dynamic
genome evolution and complex virocell metabolism of globally-distributed giant viruses.
Nature Communications 2020;11:1710.

27.   **Chen CCH, Han Y, Niu W, Kulakova AN, Howard A, et al.** Structure and kinetics of
phosphonopyruvate hydrolase from Voriovorax sp. Pal2: New insight into the divergence of
catalysis within the pep mutase/isocitrate lyase superfamily. *Biochemistry* 2006;45:11491–
11504.

28.   **Gattiker A, Gasteiger E, Bairoch A**. ScanProsite: a reference implementation of a
PROSITE scanning tool. *Applied bioinformatics* 2002;1:107–108.

29.   **Price MN, Dehal PS, Arkin AP**. FastTree 2 - Approximately maximum-likelihood trees
for large alignments. *PLoS ONE*;5. Epub ahead of print 2010. DOI:
10.1371/journal.pone.0009490.

30.   **Revell LJ**. phytools: An R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution* 2012;3:217–223.

31.   **Letunic I, Bork P**. Interactive Tree of Life (iTOL) v4: Recent updates and new
developments. *Nucleic Acids Research* 2019;47:256–259.

32.   **Kim AD, Baker AS, Dunaway-Mariano D, Metcalf WW, Wanner BL, et al.** The 2-
aminoethylphosphonate-specific transaminase of the 2-aminoethylphosphonate degradation
pathway. *J Bacteriol* 2002;184:4134–4140.

33.   **Blodgett JAV, Thomas PM, Li G, Velasquez JE, van der Donk WA, et al.** Unusual
transformations in the biosynthesis of the antibiotic phosphinothricin tripeptide. *Nat Chem
Biol* 2007;3:480–485.

34.   **Wassef MK, Hendrix JW**. Ceramide aminoethylphosphonate in the fungus Pythium
prolatum. *Biochimica et Biophysica Acta (BBA) - Lipids and Lipid Metabolism* 1977;486:172–
178.

35.   **Repeta DJ, Ferrón S, Sosa OA, Johnson CG, Repeta LD, et al.** Marine methane
paradox explained by bacterial degradation of dissolved organic matter. *Nature Geoscience*
2016;9:884–887.

36.   **Metcalf WW, Griffin BM, Cicchillo RM, Gao J, Janga SC, et al.** Synthesis of
methylphosphonic acid by marine microbes: a source for methane in the aerobic ocean.
Science 2012;337:1104–1107.

37.   **Limmathurotsakul D, Golding N, Dance DA, Messina JP, Pigott DM, et al.** Predicted
global distribution of Burkholderia pseudomallei and burden of melioidosis. *Nat Microbiol*;1.
Epub ahead of print 1 January 2016. DOI: 10.1038/nmicrobiol.2015.8.

38.   **Rice K, Batul K, Whiteside J, Kelso J, Papinski M, et al.** The predominance of
nucleotidyl activation in bacterial phosphonate biosynthesis. *Nat Commun* 2019;10:1–12.

39.   **Bartlett C, Bansal S, Burnett A, Suits MD, Schaefer J,** *et al.* Whole-Cell Detection of
C–P Bonds in Bacteria. *Biochemistry* 2017;56:5870–5873.

40.   **Watanabe Y, Nakajima M, Hoshino T, Jayasimhulu K, Brooks EE,** *et al.* A novel
sphingophosphonolipid head group 1-hydroxy-2-aminoethyl phosphonate in Bdellovibrio
stolpii. *Lipids* 2001;36:513–519.

41.   **Baumann H, Tzianabos AO, Brisson JR, Kasper DL, Jennings HJ**. Structural
elucidation of two capsular polysaccharides from one strain of Bacteroides fragilis using high-
resolution NMR spectroscopy. *Biochemistry* 1992;31:4081–4089.

42.   **Wang L, Lim CK, Klotz MG**. High Synteny and Sequence Identity between Genomes
of Nitrosococcus oceani Strains Isolated from Different Oceanic Gyres Reveals Genome
Economization and Autochthonous Clonal Evolution. *Microorganisms*;8. Epub ahead of print
8 May 2020. DOI: 10.3390/microorganisms8050693.

43.   **Yu X, Price NPJ, Evans BS, Metcalf WW**. Purification and Characterization of
Phosphonoglycans from Glycomyces sp. Strain NRRL B-16210 and Stackebrandtia
nassauensis NRRL B-16338. *J Bacteriol* 2014;196:1768–1779.

44.   **Goddard-Borger ED, Williams SJ**. Sulfoquinovose in the biosphere: occurrence,
metabolism and functions. *Biochemical Journal* 2017;474:827–849.

45.   **Bastard K, Smith AAT, Vergne-Vaxelaire C, Perret A, Zaparucha A,** *et al.*
Revealing the hidden functional diversity of an enzyme family. *Nature Chemical Biology*
2014;10:42–49.

46.   **Ji X, Tu J, Song Y, Zhang C, Wang L,** *et al.* A Luciferase-Like Monooxygenase and
Flavin Reductase Pair AbmE2/AbmZ Catalyzes Baeyer–Villiger Oxidation in
Neoabyssomicin Biosynthesis. *ACS Catal* 2020;10:2591–2595.

47.   **Maier S, Heitzler T, Asmus K, Brötz E, Hardter U,** *et al.* Functional characterization
of different ORFs including luciferase-like monooxygenase genes from the mensacarcin gene
cluster. *Chembiochem* 2015;16:1175–1182.

48.   **Kamat SS, Singh S, Rajendran A, Gama SR, Zechel DL**. 4.16 - Enzymatic Strategies
for the Catabolism of Organophosphonates. In: Liu H-W (Ben), Begley TP (editors).
Comprehensive Natural Products III. Oxford: Elsevier. pp. 399–429.

49.   **Vinogradov E, Egbosimba EE, Perry MB, Lam JS, Forsberg CW**. Structural analysis
of the carbohydrate components of the outer membrane of the lipopolysaccharide-lacking
cellulolytic ruminal bacterium Fibrobacter succinogenes S85. *European Journal of*
Biochemistry 2001;268:3566–3576.

50.   **Gilchrist CLM, Chooi Y-H**. clinker & clustermap.js: automatic generation of gene
cluster comparison figures. *Bioinformatics*. Epub ahead of print 18 January 2021. DOI:
10.1093/bioinformatics/btab007.

Figure 1. Overview of early branch points in phosphonate biosynthesis. **Left side:** Pre-Ppd
transformations. Phosphonate biosynthesis starts from Ppm-catalyzed rearrangement of PEP to PnPy,
which is followed by either aldolase (e.g. FrbC)-catalyzed condensation to PnMM, aminotransferase (e.g.
PalB)-catalyzed formation of PnAla, dehydrogenase (e.g. VlpB)-catalyzed reduction to PnLac, degradative
reactions (PalA or PhnX, shown in grey), or Ppd-catalyzed decarboxylation to afford PnAA. **Right side:**
Post-Ppd transformations include aldolase (e.g. RhiG), reductase (e.g. PhpC)-catalyzed formation of HEP,
AEPT-catalyzed formation of AEP, and aldehyde reductase (e.g. PhnY)-catalyzed PnAc formation. Note
that the PnAA-to-PnAc transformation can also be catalyzed by the α-ketoglutarate-dependent
dioxygenase and PnhY* homolog FzmG. PFAM abbreviations and identification numbers of each enzyme
are in square brackets. Dotted arrows represent reactions not biochemically characterized with purified
enzyme. **Abbreviations:** PEP, phosphoenolpyruvate; PnPy, phosphonopyruvate; PnAA,
phosphonoacetaldehyde; PnLac, phosphonolactate; PnMM, phosphonomethylmalate; PnAla,
phosphonoalanine; PnAc, phosphonoacetate; AEP, 2-aminoethylphosphonate; HEP, 2-
hydroxyethylphosphonate; KHPnPA, 2-keto-4-hydroxy-5-phosphonopentanoate; Ppm, PEP mutase; Ppd,
PnPy decarboxylase; AEPT, AEP transaminase.

622
623
Figure 2. Classification of phosphonate biosynthetic genes. A) Phylogram of 424 unique Ppm sequences
from RefSeq bacterial complete genomes, plus 12 archaeal (black dots) and a single viral Ppm sequence
(purple star). Inner coloring represents the presence of neighboring genes encoding early-stage branch
point transformations; the colors in the outer ring represent bacterial phylum or class. The outmost ring of
bars represents the number of times a given sequence was found in RefSeq genomes, with the dotted line
marking a count of 10. The two largest bars represent sequences WP_004523200 and WP_004525540.
B) Sunburst chart of early branch point phosphonate biosynthetic steps based on 434 gene neighborhoods;
BGCs possessing the same Ppm sequence (e.g. WP_004523200) are treated as a single data point.
Starting from Ppm-generated PnPy, the inner ring illustrates that most PnPy (~83%, dark green 'Ppd') is
likely transformed to PnAA based on the presence of a gene encoding Ppd. The outer ring illustrates that
post-Ppd transformations are dominated by AEPT (red). The color scheme of B matches the inner ring of
A.
636

637
638
639 **Figure 3.** Network of 882 *ppm*-containing gene neighborhoods grouped by BiG-SCAPE into 85 gene cluster
640 families (GCFs). Each dot represents a gene neighborhood colored according to organism type (e.g.
641 phylum, genus). GCFs are organized into labeled boxes and colored by biosynthetic branch point. Each
642 GCF is accompanied by a number (for identifying in BiG-SCAPE) and a descriptive name.

643
644
Figure 4. The two most abundant families of *ppm*-containing gene neighborhoods in the NCBI RefSeq
database. Gene neighborhoods representative of the 318 'Burkholderiales-CDP-OH family' members are
shown aligned (with arbitrary color scheme) using clinker[50] to highlight the conserved seven-gene core,
which is expanded below to show annotations and putative phosphonate product. The accession numbers
WP_004525540 and WP_004523200 represent the most common Ppm in AEPT- and PhnY-containing
clusters, respectively. Gene cluster family numbers corresponding to those in Figure 3 are provided to the
left of each neighborhood. Inset illustrates the four-gene core representative of gene cluster families 877
and 817 with the putative phosphonate product. NCBI RefSeq accession numbers along with the
corresponding Ppm protein accession numbers for each gene neighborhood are as follows: *Cupriavidus*
metallidurans strain FDAARGOS_675, NZ_CP046331, WP_011516537; *Pandoraea apista* strain AU2161,
NZ_CP011501, WP_042112012; *Burkholderia glumae* strain 257sh-1 chromosome 2, NZ_CP035901,
WP_017423874; *Burkholderia gladioli* strain ATCC 10248 chromosome 2, NZ_CP009322,
WP_036031650; *Burkholderia pseudomallei* isolate UKMPMC2000 chromosome 2, NZ_LR595895,
WP_004525540; *Burkholderia pseudomallei* B03 chromosome 2, NZ_CP009150, WP_004525540;
Burkholderia thailandensis E254 chromosome 2, NZ_CP004382, WP_043296855; *Burkholderia cepacia*
ATCC 25416 chromosome 2, NZ_CP007748, WP_021162084; *Burkholderia cenocepacia* strain FL-5-3-
30-S1-D7 chromosome 2, NZ_CP013396, WP_023477253; *Burkholderia vietnamiensis* strain AU1233
chromosome 1, NZ_CP013433, WP_060044431; *Burkholderia territorii* strain RF8-non-BP5 chromosome
2, NZ_CP013365, WP_059507240.
664

Bacteroides fragilis strain NCTC 9343, NC_003228, WP_011202610; 321, *Bacteroides thetaiotaomicron*
strain 7330, NZ_CP012937, WP_008767814; 863, *Treponema denticola* ATCC 35405, NZ_002967,
WP_002679012; 866, *Clostridium beijerinckii* isolate C. beijerinckii DSM 6423, NZ_LN908213,
WP_077842376; 476, *Clostridium saccharobutylicum* strain NCP 195, NZ_CP016092, WP_022744173;
856, *Clostridium butyricum* strain 4-1, NZ_CP039705, WP_002579294; 400, *Selenomonas* sp. oral taxon
478 strain F0592, NZ_CP012071, WP_050342316.
689
690

691
692
693
694 **Figure 6.** Possible citric acid cycle homologs in **A)** FrbC-like pathway of GCF 867 and **B)** unclassified GCF
695 842. The transformations mimicking those of the citric acid cycle are in the dotted box region, and the
696 names of non-phosphonate analogs are italicized in grey. Abbreviations: 2-PnMM (2-
697 phosphonomethylmalate), 3-PnMM (3-phosphonomethylmalate), OPB (2-oxo-4-phosphonobutyrate), APB
698 (2-amino-4-phosphonobutyrate), PnLac (phosphonolactate), PnPy (phosphonopyruvate).
699