1

The methyl phosphate capping enzyme Bin3 is a stable component of the fission yeast

telomerase holoenzyme

Jennifer Porat[1], Moaine El Baidouri[2,3], Jorg Grigull[4], Jean-Marc Deragon[2,3,5], & Mark A.

Bayfield[1*]

1. Department of Biology, York University, Toronto, Canada

2. LGDP-UMR5096, Université de Perpignan Via Domitia, Perpignan, France

3. CNRS LGDP-UMR5096, Perpignan, France

4. Department of Mathematics and Statistics, York University, Toronto, Canada

5. Institut Universitaire de France, Paris, France

*Correspondence: bayfield@yorku.ca (M.A.B.)

Abstract:

The telomerase holoenzyme is critical for maintaining eukaryotic genome integrity. In addition

to a reverse transcriptase and an RNA template, telomerase contains additional proteins that

protect the telomerase RNA and promote holoenzyme assembly. Here we report that the

methyl phosphate capping enzyme (MePCE) Bin3 is a stable component of the *S. pombe*

telomerase holoenzyme. Bin3 associates with the telomerase holoenzyme and U6 snRNA

through an interaction with the recently described LARP7 family member Pof8, and we

demonstrate that these two factors are evolutionarily linked in fungi. Our data suggest that the

association of Bin3 with telomerase is independent of its methyltransferase activity, but rather

that Bin3 functions in telomerase holoenzyme assembly by promoting TER1 accumulation and

Pof8 recruitment to TER1. Taken together, this work yields new insight into the composition,

assembly, and regulation of the telomerase holoenzyme in fission yeast as well as the breadth

of its evolutionary conservation.

Introduction:

To ensure complete DNA replication, the termini of eukaryotic chromosomes contain tandem

repeats, or telomeres, that can be continually extended as the ends of linear chromosomes are

lost through DNA replication[1]. Telomeres serve as protection from DNA degradation, end-to-

end fusions, chromosomal rearrangements, and chromosome loss [1,2]. With very few exceptions,

eukaryotes extend telomeric DNA sequences through the telomerase holoenzyme, a complex

containing the telomerase reverse transcriptase (Trt1 in the fission yeast *Schizosaccharomyces*

pombe), an RNA template (TER1 in *S. pombe*), and accessory proteins that promote complex

assembly and tethering to telomeric DNA [1–5]. While the reverse transcriptase/RNA template

core of telomerase is generally well-conserved among eukaryotes, major differences exist,

including in the sequence and structure of the RNA template. In contrast to ciliate telomerase

RNAs, which are small RNA Polymerase III transcripts [6], yeast and metazoan telomerase RNAs

are longer and transcribed by RNA Polymerase II as precursors[7,8]. These precursors then

undergo an extensive maturation process to yield the mature form that integrates into the

telomerase holoenzyme [3,4,7,8].

In *S. pombe,* the mature form of TER1 is generated from an intron-containing precursor through

a spliceosome-catalyzed reaction involving release of the 5' exon prior to exon ligation [9]. TER1

maturation then proceeds with the sequential binding of the Sm and Lsm complexes. Sm

proteins, well-characterized for their role in splicing, bind TER1 immediately upstream the 5'

splice site and promote 3' maturation and the addition of a 5' trimethylguanosine (TMG) cap by

the methyltransferase Tgs1 [10]. The Sm complex is then replaced by the Lsm2-8 complex, which

serves to protect the mature 3' end of TER1 from exonucleolytic degradation and promotes the

interaction between TER1 and Trt1 in the active telomerase holoenzyme. The switch from the

Sm to Lsm complexes correlates with distinct, non-overlapping substrates: TER1 precursors are

exclusively bound by the Sm complex, while the mature form of TER1, ending in a polyuridylate

stretch upstream the spliceosomal cleavage site, is only bound by the Lsm2-8 complex [10].

More recently, a role has been proposed for Pof8, a La-related protein 7 (LARP7) homolog, in

telomerase assembly and telomere maintenance in *S. pombe*. Ciliate LARP7 family proteins,

including p65 from *Tetrahymena thermophila*, have previously been characterized for their

functions in telomerase assembly [11]. Binding of p65 to stem IV of the *T. thermophila* telomerase

RNA TER results in a conformational change to the RNA and subsequent binding of the reverse

transcriptase TERT, suggesting a p65-dependent hierarchical assembly of the telomerase

holoenzyme [12]. Pof8 is hypothesized to function similarly in *S. pombe*, with Pof8 binding to TER1

promoting recruitment of the Lsm2-8 complex and enhancing the interaction between TER1

and Trt1. Accordingly, Pof8 deletion strains possess shorter telomeres indicative of a defect in

telomerase function [13–15]. Pof8 also functions in telomerase RNA quality control through

recognition of the correctly folded pseudoknot region and subsequent recruitment of the Lsm2-

8 complex to protect the 3' end from degradation [16]. Telomerase-related functions in LARP7

family proteins, particularly Pof8, map to a conserved C-terminal domain characteristic of

LARP7 family members, the extended RNA recognition motif (xRRM)[17,18]. The Pof8 xRRM is a

major determinant for Pof8-mediated telomerase activity, contributing to TER1 binding and

complex assembly [14,19]. As other fungal Pof8 homologs also possess an xRRM, it is anticipated

that Pof8 may have a conserved role in telomere maintenance beyond fission yeast [14,20].

Conversely, LARP7 homologs in higher eukaryotes are best characterized in the transcriptional

regulatory 7SK snRNP, which includes the RNA polymerase III transcribed 7SK snRNA, the

methyl phosphate capping enzyme (MePCE), and hexamethylene bisacetamide-inducible

protein (HEXIM1/2). The 7SK snRNP inhibits transcription by sequestering positive transcription

elongation factor (P-TEFb) and preventing it from phosphorylating the C-terminal domain of the

largest subunit of RNA Polymerase II, which is associated with the transition into productive

transcription elongation [21–23]. The La module and xRRM of LARP7 bind 7SK through its

polyuridylate trailer and the 3' hairpin, respectively, and promote MePCE recruitment to the

complex [24,25]. In addition to adding a γ-monomethyl phosphate cap (i.e. CH_3-pppN) to 7SK

snRNA as a means of protecting it from 5' exonucleolytic degradation, MePCE remains stably

bound to 7SK snRNA to stabilize the complex [24]. MePCE, a homolog of the *Drosophila*

melanogaster Bin3/BCDIN3 protein, also catalyzes the addition of this atypical cap structure to

U6 snRNA [26]. Bin3/MePCE homologs are present in many eukaryotes, but to date little has been

studied outside of humans and *Drosophila* [27]. More recent work has identified BCIND3D, a

related Bin3 family protein overexpressed in breast cancer cells, as the enzyme responsible for

methylating the 5' monophosphate of histidine transfer RNA and pre-miR-145 [28,29]. Despite the

many insights into the role of Bin3/MePCE in RNA processing in higher eukaryotes, the function

of the *S. pombe* Bin3 homolog has yet to be determined.

In this work, we have explored the RNA targets of the *S. pombe* Bin3 homolog in an effort to

uncover its function. In addition to an expected association of Bin3 with U6 snRNA, we present

data showing that Bin3 is a stable component of the *S. pombe* telomerase RNP. Our results also

provide evidence for an evolutionarily conserved interaction between Bin3 and LARP7

homologs. Importantly, we show that the Bin3-Pof8 interaction is required for the recruitment

of Bin3 to both U6 snRNA and the active telomerase holoenzyme. Additionally, we provide data

that Bin3 does not catalyze the addition of a γ-monomethyl phosphate cap on TER1, suggesting

a catalytic independent function. Rather, we present evidence that Bin3 functions in concert

with Pof8 to promote telomerase assembly and TER1 accumulation. Taken together, our results

identify Bin3 as a novel component of the *S. pombe* telomerase holoenzyme, thus adding a new

layer of complexity to telomerase assembly and telomere maintenance.

Results:

***S. pombe* Bin3 interacts with U6 snRNA and the telomerase RNA TER1**

While Bin3/MePCE is conserved in many eukaryotes, its presence in the fission yeast *S. pombe*,

an organism lacking 7SK snRNA [27,30], is not well understood. Unlike the longer Bin3/MePCE

homologs from higher eukaryotes, *S. pombe* Bin3 only contains a conserved

methyltransferase/SAM-binding domain (Figure S1A). Alignment of the methyltransferase

domain of *S. pombe* Bin3 to that of *H. sapiens* MePCE and *D. melanogaster* Bin3 reveals that

previously identified residues critical for SAM and SAH binding and nucleotide binding [31] are

highly conserved in *S.* pombe (Figure S1B,C).

We first sought to determine the RNA substrates of *S. pombe* Bin3 to better understand its role

in the processing of fission yeast non-coding RNA(s). Following integration of a protein A (PrA)

tag into the Bin3 genomic locus, we performed RIP-Seq, followed by validation of potential

candidate substrates with northern blotting and semi-quantitative RT-PCR (Figure 1, Table S1).

Consistent with previous reports of MePCE interacting with U6 snRNA [26], U6 emerged as one of

the most highly enriched Bin3-interacting RNA transcripts in our Seq dataset (Figure 1A, Table

S1). *S. pombe* U6 is an unusual transcript in that it is transcribed by RNA Polymerase III yet

contains an mRNA-type intron removed by the spliceosome [32,33]. Alignment of RIP-Seq reads to

the *S. pombe* genome suggested that Bin3 interacts exclusively with the spliced form of U6. We

confirmed this with northern blotting, demonstrating robust immunoprecipitation of mature

U6 and no immunoprecipitation of the intron-containing precursor (Figure 1B). Chromatin

immunoprecipitation studies suggest that 7SK and U6 snRNA may be co-transcriptionally

modified by the human MePCE homolog [34], however our data are consistent with a model

wherein *S. pombe* Bin3 interacts with U6 post-splicing.

Surprisingly, the *S. pombe* telomerase RNA TER1 was the most highly enriched hit in the Bin3

immunoprecipitates (Figure 1A, C). Given the previously established link between the human

Bin3 homolog MePCE and human LARP7, [23] as well as the established link between the *S.*

pombe LARP7 homolog Pof8 and the telomerase holoenzyme [13–15], we considered the

possibility that Bin3 may also be part of the telomerase holoenzyme through an evolutionarily

conserved interaction with LARP7 family members.

Bin3 interacts with the mature form of TER1

Since TER1 processing and maturation involves a spliceosome-catalyzed reaction [9], we next set

out to determine the processing state of Bin3-associated TER1 by sequencing the 5' and 3' ends

through circularized rapid amplification of cDNA ends (cRACE) (Figure 2A). We also performed

cRACE on TER1 immunoprecipitated by Pof8, the telomerase reverse transcriptase Trt1, and

Lsm3, all known components of the mature TER1-containing active telomerase holoenzyme

[3,4,10,13–15]. All sequenced candidates had a discrete 5' end consistent with the reported 5' end [3,4]

and a similar distribution of 3' ends ending immediately upstream the 5' splice site [9], suggesting

that Bin3 interacts with the full-length, 3' end-matured TER1 associated with the active

telomerase holoenzyme. The heterogeneity observed at the 3' end of all candidates is likely the

result of exonucleolytic nibbling prior to binding of the Lsm2-8 complex [10]. To investigate this

further, we subjected TER1 RNA immunoprecipitated by Bin3, Trt1, Pof8 and Lsm3 to RNase H

cleavage to generate shorter 5' and 3' fragments and compared fragment size by northern blot

(Figure 2B). The sizes of the 5' and 3' ends of TER1 were similar across immunoprecipitates,

further substantiating that Bin3 binds the mature form of TER1, rather than the intron-

containing precursor.

Further TER1 processing in *S. pombe* involves the addition of a 5' trimethylguanosine (TMG) cap

by Tgs1 [10]. Since Bin3 homologs are normally associated with the formation of a γ-monomethyl

phosphate cap, and to further understand when Bin3 interacts with TER1 with respect to TER1

processing, we examined the cap structure of Bin3-associated TER1 by immunoprecipitating

RNA associated with Bin3 and subsequently re-immunoprecipitating this RNA with an anti-TMG

antibody. The specificity of the anti-TMG antibody was confirmed by showing that it effectively

enriched the TMG cap-containing U5 snRNA over the TMG cap-lacking U6 snRNA or an

unrelated tRNA (Figure 2C). While Bin3-associated U6 snRNA was not immunoprecipitated by

the anti-TMG antibody, in agreement with data demonstrating the presence of a γ-monomethyl

phosphate cap on U6 *in S. pombe* [35], Bin3-associated TER1 RNA was effectively enriched by anti-

TMG immunoprecipitation (Figure 2D). These data are consistent with previous work

demonstrating a TMG cap on TER1 in *S. pombe,* and also with Bin3 associating with TER1

following spliceosomeal cleavage and 5' TMG capping, similar to what has been posited for

Pof8 [13]. This suggests that Bin3 interacts with the primary cohort of TMG capped TER1

transcripts, and that TER1 is not a substrate for Bin3-catalyzed γ-monomethyl phosphate

capping.

Bin3 interacts with components of the mature telomerase holoenzyme

Since our results indicate that Bin3 interacts with the mature form of TER1, we set out to

confirm the presence of Bin3 in the *S. pombe* telomerase holoenzyme. We immunoprecipitated

Bin3 and identified interacting proteins through mass spectrometry, followed by validation

using co-immunoprecipitation (Figure 3). We identified Pof8, the Lsm2-8 complex, and Trt1 in

Bin3 immunoprecipitates, all of which make up the catalytically active core telomerase

holoenzyme [5,10,13–15,36,37] (Table S2). We also detected the RNase P and RNase MRP subunit

Pop100 in Bin3 immunoprecipitations. Pop1, Pop6, and Pop7, subunits of RNase P and RNase

MRP in budding yeast, have recently been identified as components of the budding yeast

telomerase complex [38–40], suggesting that the involvement of RNase P and RNase MRP subunits

in telomere maintenance is evolutionarily conserved. The finding that Bin3 interacts with core

components of the telomerase holoenzyme, and not the Sm complex nor the TMG capping

enzyme Tgs1, suggests Bin3 does indeed associate with the active telomerase holoenzyme

rather than a precursor. The involvement of Bin3 in the telomerase holoenzyme was further

substantiated by gene ontology analysis to determine overrepresented biological processes and

cellular components among the top 50 Bin3 interactors. Telomerase holoenzyme complex

assembly and telomere maintenance via telomerase emerged as the top overrepresented

biological processes (Figure 3A). Similarly, overrepresented cellular components include the

Lsm2-8 complex, the telomerase holoenzyme, and spliceosomal snRNPs (Table S3). We also

repeated immunoprecipitations in the presence of benzonase to determine if these interactions

are direct or mediated through a nucleic acid intermediate. With the exception of Pof8 and an

uncharacterized protein (SPCC18B5.09c), interactions between Bin3 and other components of

the telomerase holoenzyme were lost, suggestive of an interaction mediated by TER1.

We validated the interactions between Bin3 and Pof8, as well as Bin3 and Trt1, through co-

immunoprecipitation (Figures 3B, 3C, S2). We also performed co-immunoprecipitations with

benzonase, demonstrating that the interaction between Bin3 and Pof8 persists with benzonase

treatment, while the Bin3-Trt1 interaction is lost (Figures 3B and 3C). To confirm that the Bin3-

Trt1 interaction is mediated by TER1, we repeated co-immunoprecipitations in a TER1 knockout

strain. The direct interaction between Bin3 and Pof8 remained intact in the absence of TER1,

while the Bin3-Trt1 interaction was completely lost, indicative of Bin3 assembly in the

telomerase holoenzyme nucleated, in large part, by the telomerase RNA itself. The RNA-

dependence of the interaction between Bin3 and Trt1 is reminiscent both of the budding yeast

telomerase RNA TLC1, which acts as a scaffold for telomerase holoenzyme assembly [41], as well

as the MePCE/Bin3-containing 7SK snRNP [42]. The finding that Bin3 and Pof8 interact directly is

in agreement with previous reports demonstrating a protein-protein interaction between

MePCE and LARP7 in the context of the vertebrate 7SK snRNP [34,42], suggesting the direct

interaction may be evolutionarily conserved.

Bin3 is recruited to the active telomerase holoenzyme through the LARP7 family protein Pof8

We then tested whether Bin3 was recruited to the telomerase holoenzyme through its

interaction with Pof8, much like the human LARP7 homolog promotes MePCE recruitment to

7SK snRNA [24]. Immunoprecipitation of Bin3 in the context of a *pof8Δ* strain resulted in a

complete loss of TER1 binding, providing evidence that the interaction between Bin3 and TER1

is dependent on an interaction between Bin3 and Pof8 (Figure 4A,B). Unexpectedly, the

interaction between Bin3 and U6 was also lost in the *pof8Δ* strain. Recent findings indicate that

mammalian LARP7 binds U6 to guide post-transcriptional modification [43,44], so it is tempting to

speculate that Pof8 may also mediate Bin3 binding to U6 in fission yeast. To address this, we

immunoprecipitated Pof8 alongside Bin3 and other components of the telomerase holoenzyme

and looked for U6 enrichment. We observed slight enrichment of U6 in Pof8-myc

immunoprecipitates relative to immunoprecipitation using an untagged strain or the Trt1-myc

tagged strain (Figure 4C). We did not see enrichment for the intron-containing precursor,

suggesting that like Bin3, Pof8 interacts with the spliced form of U6. The slight enrichment of

U6 with Pof8 immunoprecipitation, compared to the more robust immunoprecipitation seen

with Bin3 and Lsm3, suggests that Pof8 may transiently bind U6, perhaps serving to load Bin3

on U6. Since LARP7 binding has been reported to disrupt the catalytic activity of MePCE [34], a

transient Pof8-U6 interaction may be consistent with U6 receiving a γ-monomethyl phosphate

cap, compared to TER1, where stable Pof8 binding may prevent Bin3 catalytic activity.

To further investigate how the presence of Bin3 in the TER1-containing telomerase holoenzyme

is reliant on Pof8, we fractionated lysates from wild type and Pof8 knockout strains on a

glycerol gradient and analyzed the sedimentation patterns of Bin3, Pof8, TER1, and U6 (Figure

S3). We found that Bin3 and Pof8 co-sedimented with TER1 in higher molecular weight

fractions in a wild-type strain, as well as in lighter molecular weight fractions that contained U6

(U6 present in higher molecular weight fractions likely corresponds to U6 within the

spliceosome). In contrast, Bin3 was depleted from higher molecular weight fractions containing

TER1 in the absence of Pof8. We also noted a slight shift in TER1 towards lighter fractions in the

Pof8 deletion strain, which could be due to a lighter-migrating telomerase holoenzyme lacking

Bin3 and Pof8. These data are also consistent with Bin3 stably associating with the telomerase

holoenzyme in a manner that is dependent on Pof8.

Knowing that Bin3 interacts with TER1 and components of the telomerase holoenzyme, we next

set out to confirm whether Bin3 is part of the catalytically active telomerase holoenzyme. We

performed a previously described *in vitro* telomerase assay that relies on the presence of the

TER1- and Trt1-containing telomerase holoenzyme to extend an oligonucleotide resembling

telomeric DNA [13]. Bin3 immunoprecipitates extended the oligonucleotide in a similar manner

previously demonstrated for Pof8 ([13] and see Figure 5D), as well as showed the same loss of

activity upon RNase A treatment, supporting the idea that Bin3, much like Pof8, is a component

of the active telomerase holoenzyme (Figure 4D). Consistent with previous results[13,16], we also

observed a loss of activity for Bin3 immunoprecipitated from a *pof8Δ* strain, which can largely

be attributed to the loss of functional, correctly assembled telomerase occurring in the absence

of Pof8. Since Trt1 is only recruited to the telomerase holoenzyme following Pof8 and Lsm2-8

binding, coupled with the idea that the assay relies on the presence of a reverse transcriptase,

these results are consistent with a model in which Bin3 is recruited to TER1 through its

interaction with Pof8 and remains bound through subsequent holoenzyme assembly and the

catalytic cycle (Figure 4E).

To further investigate the link between Bin3 and Pof8 in telomerase holoenzyme assembly, we

tested whether overexpression of Bin3 could rescue the short telomere phenotype of a *pof8Δ*

strain [13–15]. Consistent with a model in which Bin3 requires Pof8 for its assembly into the active

telomerase holoenzyme, Bin3 overexpression in a *pof8Δ* strain was insufficient to rescue

telomere shortening, as measured by Southern blot, as well chromosome fusions occurring

after telomeres reach a critically short length previously associated with *the pof8Δ* strain [13]

(Figure S4).

Bin3 promotes TER1 accumulation and Pof8 recruitment to telomerase

We then interrogated the functional role of Bin3 in telomerase by creating a Bin3 knockout

strain where we replaced the Bin3 open reading frame with a phleomycin resistance cassette

which was confirmed by PCR and sequencing, as well as qRT-PCR to confirm a lack of Bin3

mRNA (Figure S5A, B). To address conflicting reports in the literature as to the nature of Bin3's

essentiality in *S. pombe*[45,46], we back-crossed our Bin3 deletion strain with a wild type strain.

Sporulation and tetrad dissections yielded viable haploid colonies possessing phleomycin

resistance (Figure S5C), confirming that Bin3 is not essential and enabling us to conduct further

mechanistic studies.

Consistent with its strong dependence on Pof8 for recruitment to telomerase, Bin3 deletion

appears to decrease steady-state TER1 levels, much like a Pof8 deletion strain, with no

reduction in U6 abundance (Figure 5A). Importantly, re-introduction of plasmid expressed Bin3

to *bin3Δ* cells was sufficient to restore wild type levels of TER1 (Figure S5D). To understand the

mechanism by which Bin3 appears to promote TER1 accumulation, we combined our Bin3

deletion strain with deletion of Rrp6, an exonuclease component of the nuclear exosome that

has been previously implicated in TER1 degradation [13,15,16,47,48]. We observed a restoration of

TER1 levels in the *bin3Δ rrp6Δ* strain (Figure 5A), similar to what has been observed for Pof8,

where Rrp6 deletion rescues decreased TER1 levels[15]. From this, we hypothesize that Bin3

promotes TER1 accumulation by preventing 3' decay by the exosome to maintain steady state

TER1 levels.

In line with decreased TER1 levels, Bin3 deletion resulted in shorter telomere length compared

to a wild-type strain, which became more evident when genomic DNA was digested with ApaI

to yield shorter telomere fragments (Figure 5B). We continued passaging strains to get to the

point of crisis (>100 generations), where telomeres are lost and cells either die or circularize

their chromosomes, leading to a loss of an observable telomere signal by southern blot [49]. As

expected, a strain lacking TER1 showed a complete loss of telomeric DNA, whereas deletion of

Pof8 and Bin3 only resulted in shorter telomeres, suggesting that while the two proteins are

important for telomere maintenance, they are not strictly required like TER1 (Figure S6).

To better understand how Bin3 affects telomere length and TER1 accumulation, we examined

how the Pof8-TER1 interaction changes upon Bin3 deletion. Somewhat surprisingly, considering

that Bin3 completely relies on Pof8 for recruitment to telomerase, we found that Bin3 deletion

also impaired Pof8 binding to TER1 (Figure 5C), much like the cooperative binding of Pof8 and

Lsm2-8 to TER1 [16]. Similarly, we observed decreased telomerase activity immunoprecipitated

by Pof8 in the Bin3 knockout strain, with normalization of the intensity of the extension bands

to TER1 levels revealing that the compromised activity cannot only be attributed to the loss of

TER1 following Bin3 deletion (Figure 5D, E). Conversely, normalization of telomere extension

activity to the amount of TER1 immunoprecipitated by Pof8 revealed that although less Pof8-

containing telomerase complexes form in the absence of Bin3, the ones that do form are fully

functional (Figure 5D, F). Thus it appears that not only does Bin3 affect TER1 accumulation but

also contributes to Pof8 association with TER1, suggesting a further role for Bin3 in ensuring

holoenzyme functionality.

Pof8-like proteins are associated with Bin3-like proteins in diverse fungal lineages

The interaction between Bin3 and Pof8, coupled with the conservation of such an interaction in

other examined eukaryotes, led us to wonder at the extent of this interaction on a phylogenetic

scale. While Bin3/MePCE and LARP7/Pof8 family members are ubiquitous in higher eukaryotes,

their absence or presence is more varied in fungal lineages. We queried hundreds of

representative fungal species for the presence of Bin3- and Pof8-like proteins and identified

many species with a) both a Bin3 and Pof8 homolog, b) only a Bin3 homolog, or c) neither a

Bin3 nor a Pof8 homolog (Figure 6, S7). Conversely, only four out of 472 examined species

contained a Pof8-like protein but no Bin3-like protein, and in these species the Pof8 homolog

was noted to have a shortened N-terminal domain that would lack the La module observed in

other LARP7 family members, raising the possibility that members of this rare cohort may not

be bona-fide LARP7 family members, at least as these are currently appreciated [20]. Based on

the observed distribution, it seems likely that both genes were present in a fungi common

ancestor, but Pof8 or both Bin3 and Pof8 were lost in certain lineages. Examination of the

distribution of Bin3 and LARP7-family members in basal eukaryotic lineages reveals a similar

pattern: Bin3 can be present without LARP7, but there are no instances where a LARP7-family

member exists in a lineage lacking a Bin3 homolog (Table S8). Similarly, LARP7-family proteins

are represented in Alveolates, Stramenopiles, Amoebozoa, Fungi, and Metazoans, while Bin3-

family proteins are present in all eukaryotic lineages. Since the presence of Pof8/LARP7 is very

highly linked to the presence of a Bin3 homolog, these data suggest that the function of LARP7

family members may be more intimately associated with the function of Bin3/MePCE than has

been previously appreciated.

Discussion:

In this study, we have identified Bin3 as a novel component of the *S. pombe* telomerase

holoenzyme. In addition to showing an interaction between Bin3 and U6 snRNA, we

demonstrate that Bin3 interacts with the telomerase RNA TER1 and components of the active

telomerase holoenzyme in a manner that is dependent on the presence of the LARP7 family

protein Pof8. Together, our results indicating that Bin3 interacts with the mature, spliced form

of TER1 and that Bin3 immunoprecipitates possess telomerase activity *in vitro* strongly suggest

that Bin3 is a constitutive component of the telomerase holoenzyme containing Trt1, Pof8,

Lsm2-8, and Est1 scaffolded on TER1 (Figure 4E). Our results are in agreement with recent

findings from the Baumann lab reporting the presence of Bin3/Bmc1 in the telomerase

holoenzyme [50]. The idea of Pof8 and Bin3 scaffolded on TER1 is reminiscent of LARP7 and

MePCE in the 7SK snRNP, with Pof8/LARP7 recruiting Bin3/MePCE to the RNA substrate. Similar

to our data showing Pof8 recruiting Bin3 to TER1 and the telomerase holoenzyme, LARP7 has

also been demonstrated to recruit MePCE to the 7SK snRNP in human cells [24]. Thus, the

protein-protein interaction and subsequent recruitment to RNA substrates by LARP7 appears

evolutionarily conserved in fission yeast and higher eukaryotes. We also present data

demonstrating that the presence of Pof8-like proteins is nearly universally correlated with the

presence of Bin3-like proteins in fungi, suggesting that the Bin3-Pof8 interaction is highly

conserved and more prevalent than previously anticipated. Since Pof8 is required to load Bin3

onto the telomerase RNA and U6 snRNA in fission yeast, this phylogenetic distribution raises

the question of whether Bin3 homologs in fungi lacking a LARP7 homolog have evolved a

mechanism to bind RNA targets in a Pof8/LARP7 independent fashion.

Since TER1 undergoes several processing steps before assembly into the telomerase

holoenzyme, we were able to assess TER1 processing state to determine when Bin3 interacts

with TER1 with respect to the timing of TER1 maturation. We show that Bin3-associated TER1

has a mature 3' end, indicative of spliceosome-catalyzed end processing, and a 5' TMG cap. This

is consistent with previously reported data on the timing of Pof8 binding; since our data

indicates that the interaction between Bin3 and TER1 is dependent on Pof8, the most

parsimonious explanation is that Pof8, and thus Bin3, bind TER1 following 3' maturation and

TMG capping, preceding the binding of the Lsm2-8 complex [13]. We show that the interaction

between Bin3 and TER1 is dependent on Pof8, leading to the conclusion that Bin3 interacts with

TER1 at the same stage as Pof8, resulting in the hierarchical assembly of the active telomerase

holoenzyme.

The finding that Bin3-associated TER1 is immunoprecipitated by a TMG antibody argues against

a catalytic role for Bin3 in TER1 processing. Our data, coupled with previous work characterizing

the cap structure on the mature TER1 bound to Trt1 [3,4], supports the idea that Bin3 has a

capping-independent function in the telomerase holoenzyme. LARP7 binding to MePCE in the

7SK snRNP inhibits its capping activity; MePCE, in turn, promotes the interaction between 7SK

and LARP7, thereby stabilizing the complex [34]. In the *S. pombe* telomerase holoenzyme and U6

snRNP, we can hypothesize that Pof8 may also inhibit the methyltransferase activity of Bin3 by

binding to and occluding the active site. This is consistent with our results suggesting a transient

interaction between Pof8 and U6 that may function to load Bin3 on U6. In such a model, an

interaction between Bin3 and Pof8 is necessary to recruit Bin3 to U6 and following dissociation

of Pof8, the catalytic site of Bin3 would become available to cap U6. It is curious to note that

Bin3 deletion has no effect on U6 levels (Figure 5B), although such a finding is consistent with

what has been observed for human U6 upon MePCE depletion [26]. Since the lack of an MePCE-

catalyzed cap on human 7SK is associated with an increase in exonucleolytic degradation[26], this

therefore raises the question as to the function of the γ-monomethyl phosphate cap on U6, if

not to protect the 5' end from exonucleolytic degradation.

Additionally, our work identifying Bin3 as a telomerase-associated protein brings to mind work

identifying a box H/ACA motif at the 3' end of vertebrate telomerase RNA [51]. Box H/ACA RNAs

form an RNP complex with Dyskerin/Cbf5, Nop10, Nhp2, and Gar1 to guide noncoding RNA

pseudouridylation [52,53]. Although Dyskerin is a pseudouridine synthase, to date no evidence has

pointed to a modification function for Dyskerin in the context of telomerase RNA. Rather, the

Dyskerin-containing H/ACA complex has a structural role in telomerase biogenesis, with binding

of the H/ACA RNP to telomerase RNA leading to a conformational change in the RNA that

promotes 3' processing and maturation over exosome-mediated decay [54]. This is reminiscent of

the apparent catalytic independent role we observed for Bin3 in fission yeast telomerase,

although our data point towards Bin3 having a role promoting Pof8 binding to TER1, perhaps

also through Bin3-mediated conformational changes in TER1. The decreased TER1 levels

observed upon Bin3 deletion (Figure 5A) can likely be attributed to the decreased interaction

between Pof8 and TER1 in the absence of Bin3. We therefore propose a model in which the

direct interaction between Bin3 and Pof8 both recruits Bin3 to the telomerase holoenzyme, as

well as promotes Pof8 binding to TER1. This in turn promotes Lsm2-8 binding to the exposed 3'

end of TER1, which protects TER1 from exosome-mediated degradation [13,15,16]. The resulting

TER1-containing RNP then recruits Trt1, forming a stable complex capable of extending

telomeres.

Another facet of this work builds on the idea that components of the telomerase holoenzyme

are more conserved than previously appreciated. In addition to the ubiquitous telomerase

reverse transcriptase, previous work has identified LARP7 family proteins in both ciliates and

yeast [11,13–15], arguing a level of evolutionary conservation. We now also provide evidence

indicating the presence of RNase P and RNase MRP subunits in fission yeast, much like the

RNase P and RNase MRP subunits found in the budding yeast telomerase complex [38–40]. The

presence of RNase P and RNase MRP subunits in two divergent yeast species supports the

importance of these factors in the function of a breadth of telomerase holoenzymes.

Our results indicating that Bin3 is part of the telomerase holoenzyme were quite unexpected,

as Bin3 has been well-characterized in higher eukaryotes for its role in the processing,

maturation, and protection of RNA Polymerase III transcripts [26,34]. An intriguing idea involves

polymerase switching through telomerase RNA evolution. Ciliate telomerase RNA is transcribed

by RNA Polymerase III and subsequently binds the LARP7 homolog p65 through the terminal

polyuridylate stretch common to RNA Polymerase III transcripts [11,17]. Since MePCE and LARP7

also associate with RNA Polymerase III transcripts, it is possible that the Pof8-bound RNA

Polymerase II transcribed TER1, and the *S. pombe* telomerase RNP as a whole, represent an

intermediate step in the evolution of telomerase RNA between ciliates and the RNA Polymerase

II transcribed telomerase RNA of higher eukaryotes. Future work should investigate the

presence of Bin3/MePCE homologs in the telomerase holoenzymes of other species,

particularly in ciliates, which possess both an RNA Polymerase III transcribed telomerase RNA

and a LARP7 homolog. Such findings will continue to develop emerging ideas regarding

conservation in telomerase RNA processing and the composition of telomerase RNPs.

Acknowledgements:

We thank Toru Nakamura and Virginia Zakian for sharing published yeast strains and Marc

Fabian and Raymund Wellinger for helpful comments on the manuscript. J.P. is supported by a

Canada Graduate Scholarship (Doctoral) from the National Sciences and Engineering Research

Council of Canada. M.A.B. is supported by a Discovery Grant from NSERC ("Impact of chemical

modification of non-coding RNAs on gene expression in *S. pombe*"). JMD is supported by the

Institut Universitaire de France (IUF), JMD and MEB by the CNRS and the University of

Perpignan (UPVD) and their study is set within the framework of the "Laboratoires

d'Excellences (LABEX)" TULIP (ANR-10-LABX-41) and of the "École Universitaire de Recherche

(EUR)" TULIP-GS (ANR-18-EURE-0019).

Author contributions:

J.P. conceived of the study, designed, performed, and analyzed all experiments, and wrote the

first draft of the manuscript with input from all authors. M.E.B. and J-M.D. performed

phylogenetic analyses. J.P. analyzed RNA-Seq data with help from J.G.. M.A.B. conceived of the

study, designed experiments, supervised the project, and edited the manuscript.

Materials and Methods:

Strains, constructs, and growth media

Strains were grown in yeast extract with supplements (YES) or Edinburgh Minimal Media

(EMM), as indicated. Tag integration and knockouts were generated with a previously described

PCR-based strategy and verified by PCR and western blotting [55] (primer sequences in Table S5).

Protein A-tagged strains were generated according to [56]. The Bin3 knockout strain was

constructed by replacing the Bin3 open reading frame with the phleomycin resistance cassette

and flanking primers containing 750 nucleotides of homology to the Bin3 genomic locus.

Correct genotypes were selected on YES plates with the corresponding antibiotic (200 μg/mL

G418, Sigma; 100 μg/mL Nourseothricin, GoldBio; 100 μg/mL phleomycin, Invivogen). Other

strains were created by mating and antibiotic selection. A list of strains is provided in Table S4.

Native protein extract and immunoprecipitation

S. pombe cells were grown in YES at 32°C to mid-log phase, harvested, and subject to cryogenic

disruption using a mortar and pestle. Cell powder was lysed in 50 mM NaCl, 20 mM Hepes pH

7.4, 55 mM KOAc, 0.5% Triton X, 0.1% Tween-20, 0.2 mM PMSF, 1:100 Protease Inhibitor

Cocktail (Thermo, 78430), and 0.004 U/μL RNase inhibitor (Invitrogen, AM2694). For Protein A-

tagged strains, immunoprecipitation was carried out with Rabbit IgG-conjugated (MP-

Biomedicals, SKU 085594) Dynabeads (Invitrogen, 14301) as described [57]. Myc-tagged proteins

were immunoprecipitated with Protein G Dynabeads (Invitrogen, 10003D) coated with anti-myc

antibody (Cell Signaling, 2276S). Beads were washed 4 times with 400 μL lysis buffer. For RNA

immunoprecipitations, bound RNA was isolated by treatment of beads with 0.1% SDS and 0.2

mg/mL Proteinase K (Sigma, P2308) at 37°C for 30 minutes, followed by extraction with phenol:

chloroform:isoamyl alcohol (25:24:1) and ethanol precipitation.

<u>RNA preparation and Northern blotting</u>

Total RNA was isolated from 1% of native protein extracts by incubation with 0.5% SDS, 0.2

mg/mL proteinase K (Sigma, P2308), 20 mM Tris HCl pH 7.4, and 10 mM EDTA pH 8.0 for 15

minutes at 50°C, followed by phenol: chloroform extraction and ethanol precipitation. Northern

blot analysis was performed as described using 8% TBE-urea polyacrylamide gels [58]. Briefly,

electrophoresed RNA was transferred to positively charged nylon membranes (Perkin Elmer,

NEF988001) with the iBlot 2 system (Thermo, IB21001) and probed with ^{32}P γ-ATP-labeled DNA

probes (probe sequences provided in Table S6). For RNase H digestion, RNA was incubated with

25 pmol RNase H probe (provided in Table S6) for 5 minutes at 65°C and slow cooled to 37°C,

followed by digestion with 5 U RNase H (NEB, M02975) for 30 minutes at 37°C. The reaction

was stopped with the addition of 25 μM EDTA pH 8.0 and phenol: chloroform extraction and

ethanol precipitation. TMG-capped RNAs were isolated from RNase H-treated

immunoprecipitates with Protein G Dynabeads coated with an α-TMG antibody (Sigma,

MABE302), according to [59].

<u>qRT-PCR and semi-quantitative RT-PCR</u>

1 μg Turbo DNase-treated RNA was reverse transcribed with the iScript cDNA synthesis kit

(BioRad, 1708890), treated with 0.5 μL RNase cocktail (Invitrogen, AM2286), and diluted 1:10.

cDNA was quantified using the SensiFAST SYBR No-Rox kit (Bioline, BIO-98005) and 1 μM of

each primer (primer sequences provided in Table S6). qPCR settings were as follows: 95°C for

10 minutes and 40 cycles consisting of 10 seconds at 95°C, 20 seconds at 60°C, and 20 seconds

at 72°C, followed by melting curve analysis. TER1 and U6 levels were normalized to act1 mRNA

levels and subsequently subject to paired two-tailed student's *t*-tests and, where applicable,

one-way ANOVA followed by a Tukey posthoc test with α set to 0.05 (Table S10).

For semi-quantitative RT-PCR, DNase-treated RNA, 10 nmol dNTP mix, and 10 pmol gene-

specific reverse primers (Table S6) were heated to 65°C and slow-cooled to 37°C before reverse

transcription with 5 U AMV-RT (NEB, M0277L) at 42°C for 1 hour. cDNA was amplified with Taq

Polymerase (NEB, MO273L) using standard protocols and the following cycling conditions: 5

minute initial denaturation at 94°C, 22 (TER1) or 17 (U6) cycles of 30 seconds at 94°C, 30

seconds at 57°C, and 1 minute at 72°C, and a final 10 minute extension at 72°C.

<u>RIP Seq</u>

Libraries were constructed from immunoprecipitated RNA samples by the RNomics Platform at

the Université de Sherbrooke in Sherbrooke, Quebec. RNA quality was assessed with a

Bioanalyzer small RNA chip (Agilent, 5067-1548). Libraries were constructed with the NEBNext

Ultra II Directional Kit (NEB, E7760S) and amplified with 10 PCR cycles. cDNA libraries were

sequenced on an Illumina NextSeq 500 with 2 runs per sample, each for 50-bp single-end reads.

Following fastp processing (Version 0.20.1), reads from the first sequencing run were aligned to

the fission yeast genome (ASM294v2) with Bowtie 2 [60,61] and counted with featurecounts [62]

using the EF2 build of the ENSEMBL fission yeast genome. Differential expression analysis was

performed using edgeR, with reads filtered to include transcripts with at least 1 CPM in each

sample and libraries were normalized by TMM[63,64].

<u>Circularized Rapid Amplification of cDNA Ends (cRACE)</u>

Immunoprecipitated RNA was treated with 2 U TURBO DNase (Invitrogen, AM2239) and

dephosphorylated with 1 U calf intestinal alkaline phosphatase (NEB, M0290), followed by

decapping and circularization with RNA 5' Pyrophosphohydrolase (NEB, M0356S) and T4 RNA

Ligase 1 (NEB, M0204S). TER1 was amplified from circularized RNA with the OneStep RT-PCR Kit

(Qiagen, 210210) (primer sequences in Table S6). cRACE products were cloned into the pGEM-T

vector (Promega, A1360) and sequenced by the TCAG DNA Sequencing Facility at the Hospital

for Sick Children in Toronto.

<u>Mass spectrometry and GO analysis</u>

Half of the native protein extract was pre-treated with 0.625 U/μL Benzonase (Millipore, E1014)

at 37°C for 30 minutes. The reaction was stopped with 5 mM EDTA pH 8.0 and Protein A

immunoprecipitation was performed in the same manner as for RNA immunoprecipitations.

After washes with lysis buffer, beads were washed with 0.1 M NH_4OAc and 0.1 mM $MgCl_2$ and

eluted with 0.5 M NH_4OH for 20 minutes at room temperature. Eluates were lyophilized and

subject to in-solution tryptic digestion. LC-MS/MS analysis was performed by the IRCM

Proteomics Discovery Platform on a Q Exactive HF.

Raw data was processed and analyzed using the MaxQuant software package 1.5.1.2 [65] and the

fission yeast UP000002485 reference proteome (24/11/2019). Settings used for MaxQuant

analysis were as reported [66]. Proteins present in immunoprecipitations from both biological

replicates and absent from control immunoprecipitations were considered genuine Bin3

interacting partners and ranked by peak intensity. PANTHER overrepresentation tests (GO

Ontology database released 2019-12-09) were conducted on the top 50 Bin3 interacting

partners. Fisher's exact tests with FDR correction were performed using the GO biological

process complete and GO cellular component complete data sets.

<u>Co-immunoprecipitation and western blotting</u>

S. pombe cells were grown in YES at 32°C to mid-log phase, harvested, and subject to cryogenic

disruption using a mortar and pestle. Cell power was lysed in Co-IP lysis buffer (10 mM Tris HCl

pH 7.5, 150 mM NaCl, 0.5% NP40, 1:100 Protease Inhibitor Cocktail). Immunoprecipitations

were carried out as described and proteins were eluted by resuspending beads in 2X SDS

loading buffer (100 mM Tris HCl pH 6.8, 4% SDS, 20% glycerol, 0.2% bromophenol blue) and

boiling at 95°C for 5 minutes. Western blot analyses were performed using monoclonal anti-

myc (Cell Signaling, 2276S) at 1:5000 and monoclonal anti-beta actin (abcam, ab8226) at 1:2500

for primary antibodies and HRP-conjugated anti-mouse at 1:5000 for secondary antibodies.

Protein A-tagged proteins were detected with HRP-conjugated polyclonal anti-Protein A

(Invitrogen, PA1-26853) at 1:5000.

<u>Glycerol gradient sedimentation</u>

Native protein extracts were separated on a 10 mL 20-50% glycerol gradient (20 mM HEPES pH

7.6, 1 mM MgOAc, 1 mM DTT, 300 mM KOAc) and spun in an SW41 rotor for 20 hours at 30,000

rpm according to [67]. Individual fractions were divided in 2 for protein extraction by TCA

precipitation and RNA extraction with phenol: chloroform.

<u>Telomerase activity assay</u>

The telomerase activity assay was performed as described [13]. Briefly, bead slurries (see

immunoprecipitation) were incubated at 30°C for 90 minutes in a 10 μL reaction containing 50

mM Tris HCl pH 8.0, 1 mM MgOAc, 5% glycerol, 1 mM spermidine, 1 mM DTT, 100 mM KOAc,

0.2 mM dATP, dCTP, and dTTP, 5 μM telomerase assay primer (Table S6), and 0.3 μM 3000

Ci/mmol [α-^{32}P] dGTP (Perkin Elmer, BLU512H250UC). The reaction was stopped with the

addition of 0.5% SDS, 10 mM EDTA pH 8.0, 20 mM Tris HCl pH 7.5, 2 μg/μL Proteinase K

(Sigma), and 1000 cpm [γ-^{32}P] ATP-labeled 100-mer oligonucleotide and incubation at 42°C for

15 minutes, followed by phenol: chloroform extraction and ethanol precipitation. RNase A-

treated samples were pre-incubated with 20 ng RNase A (Invitrogen, AM2271) at 30°C for 10

minutes. Extension products were separated on a 10% urea-TBE polyacrylamide sequencing gel

at 60 W for 90 minutes, dried and exposed to a PhosphorScreen, and imaged with a Typhoon

imager.

<u>Genomic DNA extraction, southern blotting, and chromosome fusion PCR</u>

Genomic DNA was extracted from logarithmically growing cells according to [14]. For southern

blotting, 15 μg genomic DNA was digested overnight with EcoRI or ApaI and separated on a 1%

agarose gel, transferred to positively charged nylon membranes (Perkin Elmer, NEF988001) by

capillary transfer in 10X SSC, and probed with a ^{32}P γ-ATP-labeled telomere probe (probe

sequence provided in Table S6). The chromosome fusion PCR was adapted from [13]. Briefly, 50

μL PCR reactions contained 1 μL genomic DNA, 0.4 μM forward and reverse primers (sequences

provided in Table S6), 200 μM dNTPs, 1X Taq ThermoPol Buffer (NEB), and 1.25 U Taq

Polymerase (NEB, M0273). The Trt1 gene was amplified as a loading control in an identical PCR

reaction containing the corresponding primers. The PCR reaction consisted of an initial

denaturation at 95°C for 5 minutes, followed by 32 cycles of 95°C for 15 seconds, 55°C for 30

seconds, and 68°C for 3 minutes, with a final extension at 68°C for 10 minutes.

<u>Structure prediction</u>

The structure of *S. pombe* Bin3 was predicted using Phyre2 [68] and aligned with the co-crystal

structure of the *H. sapiens* MePCE methyltransferase domain bound to 7SK snRNA and SAH

(6DCB [31]) in PyMol [69]. Primary sequences were aligned with the Clustal Omega Multiple

Sequence Alignment tool [70].

<u>Phylogenetic tree construction and search for Bin3-like and Pof8-like sequences</u>

Protein sequences (GeneCatalogs) of 472 fungal species were downloaded from the JGI website

(https://mycocosm.jgi.doe.gov/) in fasta format. 5614520 proteins were downloaded and

merged into a single multifasta file. A blast (Blastp) against all proteins was performed using

diamond blastp [71] with a minimum e-value of 1e-5 and outfmt 6 (for tabular output). In order to

identify homologous proteins, proteins were clustered together if they shared at least 50%

similarity over 70% of coverage using SiLiX [72]. When a cluster contained two or more paralogous

genes derived from duplications, the species was removed from the cluster. Finally, only

clusters with a minimum of 100 species were retained for further analysis. Following these

cleaning steps, we obtained 5328 clusters containing one gene per species. Proteins sequences

of each cluster were aligned using MAFFT (v7.271) with default parameters

(https://mafft.cbrc.jp/alignment/server/). Multiple alignments were cleaned using trimA1

(v1.4.rev22 build [2015-05-21]) with -st 0.01 option. Fastree (version 2.1.10 SSE3) with default

parameters were used to build phylogenetic trees [73]. The consensus fungi tree was obtained

using ASTRAL (astral.5.6.3) software with default parameters [74]. To detect the presence of

genes coding for Bin3-like and Pof8-like proteins, blast searches (blastp and tbalstn) were

performed on each studied fungus genome, using initially either the full-length *S. pombe* Bin3

protein sequence or the *S. pombe* Pof8 xRRM conserved motif. This procedure was repeated in

each fungal clade using clade-specific Bin3-like and Pof8-like sequences detected by the initial

search using *S. pombe* sequences. Finally, new blast searches were performed all over again,

this time using Bin3-like or Pof8-like "probes" made of a collection of sequences representing

the diversity of these two proteins in fungi.

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange

Consortium via the PRIDE [75] partner repository with the dataset identifier PDX023356 and

10.6019/PDX023356 (reviewer account details: username: reviewer_pxd023356@ebi.ac.ul

password: M2zPpwXx). RNA Seq data have been deposited in NCBI's Sequence Read Archive

(SRA) database under BioProject number PRJNA776661

(reviewer link:

https://dataview.ncbi.nlm.nih.gov/object/PRJNA776661?reviewer=af5gv098pam3mdt9rsb2rtsl

ci).

References:

1.   Blackburn, E. H. & Collins, K. Telomerase: An RNP enzyme synthesizes DNA. *Cold Spring*

630   *Harb. Perspect. Biol.* (2011) doi:10.1101/cshperspect.a003558.

631   2.   Collins, K. The biogenesis and regulation of telomerase holoenzymes. *Nature Reviews*

632   *Molecular Cell Biology* (2006) doi:10.1038/nrm1961.

633   3.   Leonardi, J., Box, J. A., Bunch, J. T. & Baumann, P. TER1, the RNA subunit of fission yeast

634   telomerase. *Nat. Struct. Mol. Biol.* (2008) doi:10.1038/nsmb1343.

635   4.   Webb, C. J. & Zakian, V. A. Identification and characterization of the

636    Schizosaccharomyces pombe TER1 telomerase RNA. *Nat. Struct. Mol. Biol.* (2008)

637    doi:10.1038/nsmb1354.

638 5.  Nakamura, T. M. *et al.* Telomerase catalytic subunit homologs from fission yeast and

639    human. *Science (80-.).* (1997) doi:10.1126/science.277.5328.955.

640 6.  Greider, C. W. & Blackburn, E. H. A telomeric sequence in the RNA of Tetrahymena

641    telomerase required for telomere repeat synthesis. *Nature* (1989)

642    doi:10.1038/337331a0.

643 7.  Chapon, C., Cech, T. R. & Zaug, A. J. Polyadenylation of telomerase RNA in budding yeast.

644    *RNA* (1997).

645 8.  Feng, J. *et al.* The RNA component of human telomerase. *Science (80-.).* (1995)

646    doi:10.1126/science.7544491.

647 9.  Box, J. A., Bunch, J. T., Tang, W. & Baumann, P. Spliceosomal cleavage generates the 3'

648    end of telomerase RNA. *Nature* (2008) doi:10.1038/nature07584.

649 10. Tang, W., Kannan, R., Blanchette, M. & Baumann, P. Telomerase RNA biogenesis involves

650    sequential binding by Sm and Lsm complexes. *Nature* (2012) doi:10.1038/nature10924.

651 11. Witkin, K. L. & Collins, K. Holoenzyme proteins required for the physiological assembly

652    and activity of telomerase. *Genes Dev.* (2004) doi:10.1101/gad.1201704.

653 12. O'Connor, C. M. & Collins, K. A Novel RNA Binding Domain in Tetrahymena Telomerase

654    p65 Initiates Hierarchical Assembly of Telomerase Holoenzyme. *Mol. Cell. Biol.* (2006)

655    doi:10.1128/mcb.26.6.2029-2036.2006.

656 13. Páez-Moscoso, D. J. *et al.* Pof8 is a La-related protein and a constitutive component of

657    telomerase in fission yeast. *Nat. Commun.* (2018) doi:10.1038/s41467-017-02284-8.

14.   Mennie, A. K., Moser, B. A. & Nakamura, T. M. LARP7-like protein Pof8 regulates

telomerase assembly and poly(A)+TERRA expression in fission yeast. *Nat. Commun.*

(2018) doi:10.1038/s41467-018-02874-0.

15.   Collopy, L. C. *et al.* LARP7 family proteins have conserved function in telomerase

assembly. *Nat. Commun.* (2018) doi:10.1038/s41467-017-02296-4.

16.   Hu, X. *et al.* Quality-Control Mechanism for Telomerase RNA Folding in the Cell. *Cell Rep.*

(2020) doi:10.1016/j.celrep.2020.108568.

17.   Singh, M. *et al.* Structural Basis for Telomerase RNA Recognition and RNP Assembly by

the Holoenzyme La Family Protein p65. *Mol. Cell* (2012)

doi:10.1016/j.molcel.2012.05.018.

18.   Singh, M., Choi, C. P. & Feigon, J. xRRM: A new class of RRM found in the telomerase La

family protein p65. *RNA Biol.* (2013) doi:10.4161/rna.23608.

19.   Basu, R., Eichhorn, C. D., Cheng, R., Peterson, R. D. & Feigon, J. Structure of S. pombe

telomerase protein Pof8 C-terminal domain is an xRRM conserved among LARP7

proteins. *RNA Biol.* (2020) doi:10.1080/15476286.2020.1836891.

20.   Deragon, J.-M. M. Distribution, organization an evolutionary history of La and LARPs in

eukaryotes. *RNA Biol.* 1–9 (2020) doi:10.1080/15476286.2020.1739930.

21.   Nguyen, V. T., Kiss, T., Michels, A. A. & Bensaude, O. 7SK small nuclear RNA binds to and

inhibits the activity of CDK9/cyclin T complexes. *Nature* (2001) doi:10.1038/35104581.

22.   Yang, Z., Zhu, Q., Luo, K. & Zhou, Q. The 7SK small nuclear RNA inhibits the CDK9/cyclin

T1 kinase to control transcription. *Nature* (2001) doi:10.1038/35104575.

23.   Krueger, B. J. *et al.* LARP7 is a stable component of the 7SK snRNP while P-TEFb, HEXIM1

and hnRNP A1 are reversibly associated. *Nucleic Acids Res.* (2008)

doi:10.1093/nar/gkn061.

24. Muniz, L., Egloff, S. & Kiss, T. RNA elements directing in vivo assembly of the

7SK/MePCE/Larp7 transcriptional regulatory snRNP. *Nucleic Acids Res.* (2013)

doi:10.1093/nar/gkt159.

25. Eichhorn, C. D., Chug, R. & Feigon, J. hLARP7 C-terminal domain contains an xRRM that

binds the 3' hairpin of 7SK RNA. *Nucleic Acids Res.* (2016) doi:10.1093/nar/gkw833.

26. Jeronimo, C. *et al.* Systematic Analysis of the Protein Interaction Network for the Human

Transcription Machinery Reveals the Identity of the 7SK Capping Enzyme. *Mol. Cell*

(2007) doi:10.1016/j.molcel.2007.06.027.

27. Cosgrove, M. S., Ding, Y., Rennie, W. A., Lane, M. J. & Hanes, S. D. The bin3 RNA

methyltransferase targets 7SK RNA to control transcription and translation. *Wiley*

Interdisciplinary Reviews: RNA (2012) doi:10.1080/09540121.2017.1344767.

28. Xhemalce, B., Robson, S. C. & Kouzarides, T. Human RNA methyltransferase BCDIN3D

regulates MicroRNA processing. *Cell* (2012) doi:10.1016/j.cell.2012.08.041.

29. Martinez, A. *et al.* Human BCDIN3D monomethylates cytoplasmic histidine transfer RNA.

Nucleic Acids Res. (2017) doi:10.1093/nar/gkx051.

30. Marz, M. *et al.* Evolution of 7SK RNA and its protein partners in metazoa. *Mol. Biol. Evol.*

(2009) doi:10.1093/molbev/msp198.

31. Yang, Y., Eichhorn, C. D., Wang, Y., Cascio, D. & Feigon, J. Structural basis of 7SK RNA 5'-γ-

phosphate methylation and retention by MePCE. *Nat. Chem. Biol.* (2019)

doi:10.1038/s41589-018-0188-z.

32.   Tani, T. & Ohshima, Y. The gene for the U6 small nuclear RNA in fission yeast has an

703   intron. *Nature* (1989) doi:10.1038/337087a0.

704   33.   Tani, T. & Ohshima, Y. mRNA-type introns in U6 small nuclear RNA genes: Implications for

705   the catalysis in pre-mRNA splicing. *Genes Dev.* (1991) doi:10.1101/gad.5.6.1022.

706   34.   Xue, Y., Yang, Z., Chen, R. & Zhou, Q. A capping-independent function of MePCE in

707   stabilizing 7SK snRNA and facilitating the assembly of 7SK snRNP. *Nucleic Acids Res.*

708   (2009) doi:10.1093/nar/gkp977.

709   35.   Gu, J., Patton, J. R., Shimba, S. & Reddy, R. Localization of modified nucleotides in

710   Schizosaccharomyces pombe spliceosomal small nuclear RNAs: Modified nucleotides are

711   clustered in functionally important regions. *RNA* (1996).

712   36.   Beernink, H. T. H., Miller, K., Deshpande, A., Bucher, P. & Cooper, J. P. Telomere

713   maintenance in fission yeast requires an est1 ortholog. *Curr. Biol.* (2003)

714   doi:10.1016/S0960-9822(03)00169-6.

715   37.   Webb, C. J. & Zakian, V. A. Schizosaccharomyces pombe Ccq1 and TER1 bind the 14-3-3-

716   like domain of Est1, which promotes and stabilizes telomerase-telomere association.

717   *Genes Dev.* (2012) doi:10.1101/gad.181826.111.

718   38.   Lemieux, B. *et al.* Active Yeast Telomerase Shares Subunits with Ribonucleoproteins

719   RNase P and RNase MRP. *Cell* (2016) doi:10.1016/j.cell.2016.04.018.

720   39.   Garcia, P. D. *et al.* Stability and nuclear localization of yeast telomerase depend on

721   protein components of RNase P/MRP. *Nat. Commun.* (2020) doi:10.1038/s41467-020-

722   15875-9.

40.   Laterreur, N. *et al.* The yeast telomerase module for telomere recruitment requires a

specific RNA architecture. *RNA* **24**, (2018).

41.   Zappulla, D. C. & Cech, T. R. Yeast telomerase RNA: A flexible scaffold for protein

subunits. *Proc. Natl. Acad. Sci. U. S. A.* (2004) doi:10.1073/pnas.0403641101.

42.   Barboric, M. *et al.* 7SK snRNP/P-TEFb couples transcription elongation with alternative

splicing and is essential for vertebrate development. *Proc. Natl. Acad. Sci. U. S. A.* (2009)

doi:10.1073/pnas.0903188106.

43.   Hasler, D. *et al.* The Alazami Syndrome-Associated Protein LARP7 Guides U6 Small

Nuclear RNA Modification and Contributes to Splicing Robustness. *Mol. Cell* (2020)

doi:10.1016/j.molcel.2020.01.001.

44.   Wang, X. *et al.* LARP7-Mediated U6 snRNA Modification Ensures Splicing Fidelity and

Spermatogenesis in Mice. *Mol. Cell* (2020) doi:10.1016/j.molcel.2020.01.002.

45.   Hayles, J. *et al.* A genome-wide resource of cell cycle and cell shape genes of fission

yeast. *Open Biol.* (2013) doi:10.1098/rsob.130053.

46.   Kim, D. U. *et al.* Analysis of a genome-wide set of gene deletions in the fission yeast

Schizosaccharomyces pombe. *Nat. Biotechnol.* (2010) doi:10.1038/nbt.1628.

47.   Lee, N. N. *et al.* XMtr4-like protein coordinates nuclear RNA processing for

heterochromatin assembly and for telomere maintenance. *Cell* **155**, (2013).

48.   Coy, S., Volanakis, A., Shah, S. & Vasiljeva, L. The Sm Complex Is Required for the

Processing of Non-Coding RNAs by the Exosome. *PLoS One* **8**, (2013).

49.   Nakamura, T. M., Cooper, J. P. & Cech, T. R. Two modes of survival of fission yeast

without telomerase. *Science* vol. 282 (1998).

50.   Paez-Moscoso, D. J. *et al.* A putative cap binding protein and the methyl phosphate

capping enzyme Bin3/MePCE function in telomerase biogenesis. *bioRxiv*

2021.12.15.472819 (2021) doi:10.1101/2021.12.15.472819.

51.   Mitchell, J. R., Cheng, J. & Collins, K. A Box H/ACA Small Nucleolar RNA-Like Domain at

the Human Telomerase RNA 3′ End. *Mol. Cell. Biol.* (1999) doi:10.1128/mcb.19.1.567.

52.   Matera, A. G., Terns, R. M. & Terns, M. P. Non-coding RNAs: Lessons from the small

nuclear and small nucleolar RNAs. *Nature Reviews Molecular Cell Biology* (2007)

doi:10.1038/nrm2124.

53.   Lafontaine, D. L. J., Bousquet-Antonelli, C., Henry, Y., Caizergues-Ferrer, M. & Tollervey,

D. The box H + ACA snoRNAs carry Cbf5p, the putative rRNA pseudouridine synthase.

Genes Dev. (1998) doi:10.1101/gad.12.4.527.

54.   Tseng, C. K., Wang, H. F., Schroeder, M. R. & Baumann, P. The H/ACA complex disrupts

triplex in hTR precursor to permit processing by RRP6 and PARN. *Nat. Commun.* (2018)

doi:10.1038/s41467-018-07822-6.

55.   Bähler, J. *et al.* Heterologous modules for efficient and versatile PCR-based gene

targeting in Schizosaccharomyces pombe. *Yeast* (1998) doi:10.1002/(SICI)1097-

0061(199807)14:10<943::AID-YEA292>3.0.CO;2-Y.

56.   Oeffinger, M. *et al.* Comprehensive analysis of diverse ribonucleoprotein complexes. *Nat.*

Methods (2007) doi:10.1038/nmeth1101.

57.   Vakiloroayaei, A., Shah, N. S., Oeffinger, M. & Bayfield, M. A. The RNA chaperone La

promotes pre-tRNA maturation via indiscriminate binding of both native and misfolded

targets. *Nucleic Acids Res.* **45**, 11341–11355 (2017).

58.   Porat, J. & Bayfield, M. A. Use of tRNA-Mediated Suppression to Assess RNA Chaperone

768   Function. in *RNA Chaperones: Methods and Protocols* (ed. Heise, T.) 107–120 (Springer

769   US, 2020). doi:10.1007/978-1-0716-0231-7_6.

770 59. Hayes, K. E., Barr, J. A., Xie, M., Steitz, J. A. & Martinez, I. Immunoprecipitation of Tri-

771   methylated Capped RNA. *Bio-protocol* **8**, e2717 (2018).

772 60. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods*

773   **9**, (2012).

774 61. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient

775   alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, (2009).

776 62. Liao, Y., Smyth, G. K. & Shi, W. FeatureCounts: An efficient general purpose program for

777   assigning sequence reads to genomic features. *Bioinformatics* **30**, (2014).

778 63. Liu, R. *et al.* Why weight? Modelling sample and observational level variability improves

779   power in RNA-seq analyses. *Nucleic Acids Res.* **43**, (2015).

780 64. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: A Bioconductor package for

781   differential expression analysis of digital gene expression data. *Bioinformatics* **26**, (2009).

782 65. Cox, J. & Mann, M. MaxQuant enables high peptide identification rates, individualized

783   p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat. Biotechnol.*

784   (2008) doi:10.1038/nbt.1511.

785 66. Drissi, R., Dubois, M. L., Douziech, M. & Boisvert, F. M. Quantitative proteomics reveals

786   dynamic interactions of the minichromosome maintenance complex (MCM) in the

787   cellular response to etoposide induced DNA damage. *Mol. Cell. Proteomics* (2015)

788   doi:10.1074/mcp.M115.048991.

789 67. Lue, N. F. & Peng, Y. Identification and characterization of a telomerase activity from

Schizosaccharomyces pombe. *Nucleic Acids Res.* **25**, 4331–4337 (1997).

68.   Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N. & Sternberg, M. J. E. The Phyre2 web

portal for protein modeling, prediction and analysis. *Nat. Protoc.* (2015)

doi:10.1038/nprot.2015.053.

69.   DeLano, W. L. The PyMOL Molecular Graphics System, Version 1.8. *Schrödinger LLC*

(2014) doi:10.1038/hr.2014.17.

70.   Sievers, F. & Higgins, D. G. Clustal Omega. *Curr. Protoc. Bioinforma.* (2014)

doi:10.1002/0471250953.bi0313s48.

71.   Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using DIAMOND.

Nature Methods (2014) doi:10.1038/nmeth.3176.

72.   Miele, V., Penel, S. & Duret, L. Ultra-fast sequence clustering from similarity networks

with SiLiX. *BMC Bioinformatics* (2011) doi:10.1186/1471-2105-12-116.

73.   Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 - Approximately maximum-likelihood

trees for large alignments. *PLoS One* (2010) doi:10.1371/journal.pone.0009490.

74.   Mirarab, S. *et al.* ASTRAL: Genome-scale coalescent-based species tree estimation. in

Bioinformatics (2014). doi:10.1093/bioinformatics/btu462.

75.   Perez-Riverol, Y. *et al.* The PRIDE database and related tools and resources in 2019:

Improving support for quantification data. *Nucleic Acids Res.* (2019)

doi:10.1093/nar/gky1106.

Figure Legends:

Figure 1: Bin3 interacts with U6 snRNA and the telomerase RNA TER1

A) Enrichment of Bin3 PrA immunoprecipitated transcripts compared to an untagged control

(biological replicates = 3). Axes represent log2 of fold change and negative log of false discovery

rate.

B) Northern blot analysis of the mature and intron-containing U6 from total RNA and PrA

immunoprecipitates from untagged (wild type, wt) and PrA-tagged Bin3 strains.

C) Northern blot of the telomerase RNA TER1, U6, and U5 from total RNA and PrA

immunoprecipitates from untagged and PrA-tagged Bin3 strains, and semi-quantitative RT-PCR

analysis of TER1.

Figure 2: Bin3 interacts with the same TER1 species as well-established components of the

telomerase holoenzyme

A) The 5' and 3' ends of Bin3-, Trt1-, Pof8-, and Lsm3-associated TER1 were identified by cRACE.

The results of 10 independent clones per immunoprecipitation are shown below a schematic of

the architecture of TER1.

B) RNase H northern blots of RNase H-generated 5' and 3' ends of TER1 immunoprecipitated by

various telomerase components. The same blot was stripped and reprobed for U6.

C-D) Northern blot of α-TMG flow through (FT) and immunoprecipitated (IP) transcripts from

total RNA (C) and Bin3-associated RNA (D).

Figure 3: Bin3 interacts with components of the mature telomerase holoenzyme

A) Gene ontology analysis (biological process) of top 50 Bin3 protein interactors.

B-C) Examination and nucleic acid-dependence of interactions between Bin3 and Pof8 (B) and

Bin3 and Trt1 (C) by co-immunoprecipitation and western blotting. Blots were reprobed for

beta actin as a loading control.

Figure 4: Bin3 is recruited to the active telomerase holoenzyme by Pof8

A) Northern blot and semi-quantitative RT-PCR of TER1 and U6 in PrA immunoprecipitates for

an untagged wild type (wt), PrA-tagged (Bin3 PrA), and PrA-tagged Pof8 knockout strain (Bin3

PrA ΔPof8). Bin3-PrA was detected in input and immunoprecipitated samples by western blots

probing for PrA (bottom panel). Possible cleavage products are indicated with an asterisk.

B) qRT-PCR of TER1 and U6 in Bin3 PrA immunoprecipitates from a wild type and *pof8Δ* strain

normalized to input RNA. Relative TER1 and U6 IP was calculated by comparing percent

immunoprecipitation of TER1 or U6 to immunoprecipitation from an untagged strain (mean $\pm$

standard error, paired *t* test * at $p < 0.05$, ** at $p < 0.01$, *** at $p < 0.001$, and **** at p<0.0001)

(n=2 biological replicates).

C) Northern blot of mature and intron-containing U6 in PrA- and myc-tagged

immunoprecipitated RNA.

D) Telomerase assay of PrA-tagged Bin3 in a wild type and *pof8Δ* strain. A ^{32}P-labeled 100-mer

oligonucleotide was used as a loading control. Telomerase extension products were compared

to a terminal transferase ladder, with +1 and +4 extension products indicated. Western blot

probing for PrA following PrA immunoprecipitation is shown in the panel below.

E) Proposed model of the fission yeast telomerase holoenzyme. TER1 structure and binding

locations are based on models constructed by Hu and colleagues[16] and Mennie and

colleagues[14].

Figure 5: Bin3 promotes TER1 accumulation and Pof8 recruitment to telomerase

A) Quantitation of TER1 and U6 in total RNA by qRT-PCR, normalized to act1 mRNA. P-values

over bars represent comparison to a wild type strain (mean $\pm$ standard error, paired t test *

at $p < 0.05$, ** at $p < 0.01$, *** at $p < 0.001$, and **** at $p<0.0001$) (n=3 biological replicates).

B) Southern blot comparing telomere length in following 3 restreaks on rich media (one

restreak= 20-25 generations). Genomic DNA was digested with EcoRI (left) or ApaI (right) to

yield different sized telomere restriction fragments.

C) qRT-PCR of TER1 in Pof8 myc immunoprecipitates from a wild type and *bin3Δ* strain

normalized to input RNA. Relative TER1 IP was calculated by comparing percent

immunoprecipitation of TER1 to immunoprecipitation from an untagged strain (mean $\pm$

standard error, paired t test * at $p < 0.05$, ** at $p < 0.01$, *** at $p < 0.001$, and **** at $p<0.0001$)

(n=2 biological replicates). P-values over each bar represent results of a paired t-test with Pof8

myc.

D) Telomerase assay of myc-tagged Pof8 in a wild type and *bin3Δ* strain. A [32]P-labeled 100-mer

oligonucleotide was used as a loading control. Telomerase extension products were compared

to a terminal transferase ladder, with +1 and +4 extension products indicated. Western blot

probing for myc following myc immunoprecipitation is shown in the panel below.

E-F) Relative telomerase extension activity for myc-tagged Pof8 immunoprecipitates in a wild

type and *bin3Δ* strain. The intensity of the +4 extension product was normalized to a

precipitation loading control, then further normalized to TER1 expression (E) or the amount of

TER1 immunoprecipitated with Pof8 (F). P-values over each bar represent results of a paired t-

test with Pof8 myc (n=4).

Figure 6: Phylogenic distribution of Bin3 and Pof8 in fungi

Consensus cladogram describing the phylogenic relationships of 472 species representative of

fungi phylum and classes and highlighting (using a color code) the distribution of Bin3 and Pof8

in these species. The cladogram is a consensus tree of 5328 distinct protein coding gene trees

resulting from a genome-wide, against all, protein comparison (see Materials and Methods).

Only posterior probabilities inferior to 1 are shown. The Bin3 and Pof8 distribution is

recapitulated in Supplemental Table 8 with corresponding protein sequences. A cartoon

presenting structural domains of Pof8 is presented. Pof8 La module and RRM1 are only inferred

from secondary structure predictions and are represented by *La** and *RRM1** in the

corresponding cartoon. In four species (colored in orange in the cladogram) only an N-terminal

truncated version of Pof8 (Pof8*) that cannot accommodate the La module is present. Full

cladogram with species names and statistical supports of the different nodes is presented in

Figure S7.

Supplementary figure captions:

Supplementary figure 1: Bin3 is the *S. pombe* methyl phosphate capping enzyme (MePCE)

homolog

A) Schematic of the domain architecture of Bin3 homologs from *S. pombe*, *H. sapiens,* and *D.*

melanogaster.

B) Clustal alignment of the methyltransferase/SAM-binding domains of *S. pombe* Bin3, *H.*

sapiens MePCE, and *D. melanogaster* Bin3. Predicted SAM-binding residues are indicated in red

and nucleotide-binding residues are in blue.

C) Phyre2[68] structure prediction of *S. pombe* Bin3 bound to a 5' terminal guanosine (orange)

and SAH (green). Predicted SAM-binding residues are in red and nucleotide-binding residues

are in blue.

Supplementary figure 2: Reciprocal co-immunoprecipitations validate an interaction between

Bin3 and Pof8

The interaction between Bin3 and Pof8 was confirmed by α-myc co-immunoprecipitation.

Supplementary figure 3: Glycerol gradient sedimentation of Bin3, Pof8, TER1, and U6 in a wild

type and *pof8Δ* strain

Western blot and semi-quantitative RT-PCR of Bin3 PrA, Pof8 myc, TER1, and U6 in gradient

fractions.

Supplementary figure 4: Bin3 overexpression fails to rescue *pof8Δ*-dependent telomere

defects

A) Chromosome fusion PCR (top) following 2 restreaks on minimal media (one re-streak= 20-25

generations) from wild type and *pof8Δ* strains overexpressing an empty vector (pRep4x) or

Bin3. The Trt1 gene was amplified from genomic DNA as a loading control (bottom).

B) Southern blot comparing telomere length from the same genomic DNA as (A). Genomic DNA

was digested with EcoRI.

Supplementary figure 5: Validation of a Bin3 knockout strain confirms its viability

A) Schematic of Bin3 knockout (top) and PCR validation of the resultant strain using the

indicated primers.

B) qRT-PCR of bin3 mRNA normalized to act1 mRNA for wild type and *bin3Δ* strains (mean $\pm$

standard error, paired *t* test * at $p < 0.05$, ** at $p < 0.01$, *** at $p < 0.001$, and **** at p<0.0001)

(n=3 biological replicates). P-values over each bar represent results of a paired t-test with a wild

type strain.

C) Backcrossing of *bin3Δ* and a wild type strain, followed by tetrad dissection, confirms viability

of the *bin3Δ* strain on rich media. Tetrads from the same spore are arranged in columns.

Resulting haploid colonies possessing phleomycin resistance are circled in red and colonies that

didn't grow after tetrad dissection are circled in black.

D) qRT-PCR of TER1 normalized to act1 mRNA for wild type and *bin3Δ* strains transformed with

an empty vector (3x) or Bin3 (mean $\pm$ standard error, paired *t* test * at $p < 0.05$, ** at $p < 0.01$,

*** at $p < 0.001$, and **** at p<0.0001) (n=3 biological replicates). P-values over each bar

represent results of a paired t-test with a wild type strain transformed with an empty vector

(wild type + pRep3x).

Supplementary figure 6: Bin3 and Pof8 are not strictly required for telomere maintenance

Southern blot comparing ApaI-digested genomic DNA isolated after 5 restreaks on rich media. A

lack of signal with a telomeric probe represents survivors that have lost telomeric DNA and

circularized their chromosomes (*TER1Δ*).

Supplementary figure 7: Phylogenic distribution of Bin3 and Pof8 in fungi

Complete consensus cladogram describing the phylogenic relationships of 472 species

representative of fungi phylum and classes and highlighting (using a color code) the distribution

of Bin3 and Pof8 in these species. Only posterior probabilities inferior to 1 are shown.

Resulting haploid colonies possessing phleomycin resistance are circled in red and colonies that

didn't grow after tetrad dissection are circled in black.

D) qRT-PCR of TER1 normalized to act1 mRNA for wild type and *bin3Δ* strains transformed with

an empty vector (3x) or Bin3 (mean $\pm$ standard error, paired *t* test * at $p > 0.05$, ** at $p < 0.01$,

*** at $p < 0.001$, and **** at p<0.0001) (n=3 biological replicates). P-values over each bar

represent results of a paired t-test with a wild type strain transformed with an empty vector

(wild type + pRep3x).

Supplementary figure 6: Bin3 and Pof8 are not strictly required for telomere maintenance

Southern blot comparing ApaI-digested genomic DNA isolated after 5 restreaks on rich media. A

lack of signal with a telomeric probe represents survivors that have lost telomeric DNA and

circularized their chromosomes (*TER1Δ*).

Supplementary figure 7: Phylogenic distribution of Bin3 and Pof8 in fungi

Complete consensus cladogram describing the phylogenic relationships of 472 species

representative of fungi phylum and classes and highlighting (using a color code) the distribution

of Bin3 and Pof8 in these species. Only posterior probabilities inferior to 1 are shown.

Figure 1: Bin3 interacts with U6 snRNA and the telomerase RNA TER1
A) Enrichment of Bin3 PrA immunoprecipitated transcripts compared to an untagged control (biological replicates = 3). Axes represent log2 of fold change and negative log of false discovery rate.
B) Northern blot analysis of the mature and intron-containing U6 from total RNA and PrA immunoprecipitates from untagged (wild type, wt) and PrA-tagged Bin3 strains.
C) Northern blot of the telomerase RNA TER1, U6, and U5 from total RNA and PrA immunoprecipitates from untagged and PrA-tagged Bin3 strains, and semi-quantitative RT-PCR analysis of TER1.

Figure 2: Bin3 interacts with the same TER1 species as well-established components of the telomerase holoenzyme

A) The 5' and 3' ends of Bin3-, Trt1-, Pof8-, and Lsm3-associated TER1 were identified by cRACE. The results of 10 independent clones per immunoprecipitation are shown below a schematic of the architecture of TER1.

B) RNase H northern blots of RNase H-generated 5' and 3' ends of TER1 immunoprecipitated by various telomerase components. The same blot was stripped and reprobed for U6.

C-D) Northern blot of a-TMG flow through (FT) and immunoprecipitated (IP) transcripts from total RNA (C) and Bin3-associated RNA (D).

GO biological process	Fold enrichment	Raw P value	False discovery rate
Telomerase holoenzyme complex assembly	49.39	2.04E-08	5.33E-05
Telomere maintenance via telomerase	31.75	2.54E-04	4.74E-02
RNA-dependent DNA biosynthetic process	31.75	2.54E-04	4.58E-02
Ribonucleoprotein complex assembly	5.47	6.70E-06	2.50E-03
Ribonucleoprotein complex subunit organization	5.29	9.01E-06	2.77E-03
mRNA cis splicing, via spliceosome	5.25	6.76E-05	1.54E-02
mRNA splicing, via spliceosome	4.78	5.48E-05	1.43E-02
RNA splicing, via transesterification reactions with bulged adenosine as nucleophile	4.78	5.48E-05	1.43E-02
RNA splicing, via transesterification reactions	4.72	6.08E-05	1.45E-02
RNA splicing	4.52	8.60E-05	1.80E-02

Figure 3: Bin3 interacts with components of the mature telomerase holoenzyme
A) Gene ontology analysis (biological process) of top 50 Bin3 protein interactors.
B-C) Examination and nucleic acid-dependence of interactions between Bin3 and Pof8 (B) and Bin3 and Trt1 (C) by co-immunoprecipitation and western blotting. Blots were reprobed for beta actin as a loading control.

Figure 4: Bin3 is recruited to the active telomerase holoenzyme by Pof8

A) Northern blot and semi-quantitative RT-PCR of TER1 and U6 in PrA immunoprecipitates for an untagged wild type (wt), PrA-tagged (Bin3 PrA), and PrA-tagged Pof8 knockout strain (Bin3 PrA ΔPof8). Bin3-PrA was detected in input and immunoprecipitated samples by western blots probing for PrA (bottom panel). Possible cleavage products are indicated with an asterisk.

B) qRT-PCR of TER1 and U6 in Bin3 PrA immunoprecipitates from a wild type and pof8Δ strain normalized to input RNA. Relative TER1 and U6 IP was calculated by comparing percent immunoprecipitation of TER1 or U6 to immunoprecipitation from an untagged strain (mean ± standard error, paired t test * at p < 0.05, ** at p < 0.01, *** at p < 0.001, and **** at p<0.0001) (n=2 biological replicates).

C) Northern blot of mature and intron-containing U6 in PrA- and myc-tagged immunoprecipitated RNA.

D) Telomerase assay of PrA-tagged Bin3 in a wild type and pof8Δ strain. A 32P-labeled 100-mer oligonucleotide was used as a loading control. Telomerase extension products were compared to a terminal transferase ladder, with +1 and +4 extension products indicated. Western blot probing for PrA following PrA immunoprecipitation is shown in the panel below.

E) Proposed model of the fission yeast telomerase holoenzyme. TER1 structure and binding locations are based on models constructed by Hu and colleagues16 and Mennie and colleagues14.

Figure 5: Bin3 promotes TER1 accumulation and Pof8 recruitment to telomerase

A) Quantitation of TER1 and U6 in total RNA by qRT-PCR, normalized to act1 mRNA. P-values over bars represent comparison to a wild type strain (mean ± standard error, paired t test * at p < 0.05, ** at p < 0.01, *** at p < 0.001, and **** at p<0.0001) (n=3 biological replicates).

B) Southern blot comparing telomere length in following 3 restreaks on rich media (one restreak= 20-25 generations). Genomic DNA was digested with EcoRI (left) or ApaI (right) to yield different sized telomere restriction fragments.

C) qRT-PCR of TER1 in Pof8 myc immunoprecipitates from a wild type and bin3Δ strain normalized to input RNA. Relative TER1 IP was calculated by comparing percent immunoprecipitation of TER1 to immunoprecipitation from an untagged strain (mean ± standard error, paired t test * at p < 0.05, ** at p < 0.01, *** at p < 0.001, and **** at p<0.0001) (n=2 biological replicates). P-values over each bar represent results of a paired t-test with Pof8 myc.

D) Telomerase assay of myc-tagged Pof8 in a wild type and bin3Δ strain. A 32P-labeled 100-mer oligonucleotide was used as a loading control. Telomerase extension products were compared to a terminal transferase ladder, with +1 and +4 extension products indicated. Western blot probing for myc following myc immunoprecipitation is shown in the panel below.

E-F) Relative telomerase extension activity for myc-tagged Pof8 immunoprecipitates in a wild type and bin3Δ strain. The intensity of the +4 extension product was normalized to a precipitation loading control, then further normalized to TER1 expression (E) or the amount of TER1 immunoprecipitated with Pof8 (F). P-values over each bar represent results of a paired t-test with Pof8 myc (n=4).

Figure 6: Phylogenic distribution of Bin3 and Pof8 in fungi
Consensus cladogram describing the phylogenic relationships of 472 species representative of fungi phylum and classes and highlighting (using a color code) the distribution of Bin3 and Pof8 in these species. The cladogram is a consensus tree of 5328 distinct protein coding gene trees resulting from a genome-wide, against all, protein comparison (see Materials and Methods). Only posterior probabilities inferior to 1 are shown. The Bin3 and Pof8 distribution is recapitulated in Supplemental Table 8 with corresponding protein sequences. A cartoon presenting structural domains of Pof8 is presented. Pof8 La module and RRM1 are only inferred from secondary structure predictions and are represented by *La** and *RRM1** in the corresponding cartoon. In four species (colored in orange in the cladogram) only an N-terminal truncated version of Pof8 (Pof8*) that cannot accommodate the La module is present. Full cladogram with species names and statistical supports of the different nodes is presented in Figure S7.

Supplementary figure 1: Bin3 is the *S. pombe* methyl phosphate capping enzyme (MePCE) homolog
A) Schematic of the domain architecture of Bin3 homologs from *S. pombe*, *H. sapiens,* and *D. melanogaster*.
B) Clustal alignment of the methyltransferase/SAM-binding domains of *S. pombe* Bin3, *H. sapiens* MePCE, and *D. melanogaster* Bin3. Predicted SAM-binding residues are indicated in red and nucleotide-binding residues are in blue.
C) Phyre2 structure prediction 67 of *S. pombe* Bin3 bound to a 5' terminal guanosine (orange) and SAH (green). Predicted SAM-binding residues are in red and nucleotide-binding residues are in blue.

Supplementary figure 2: Reciprocal co-immunoprecipitations validate an interaction between Bin3 and Pof8

The interaction between Bin3 and Pof8 was confirmed by a-myc co-immunoprecipitation.

Supplementary figure 3: Glycerol gradient sedimentation of Bin3, Pof8, TER1, and U6 in a wild type and pof8Δ strain
Western blot and semi-quantitative RT-PCR of Bin3 PrA, Pof8 myc, TER1, and U6 in gradient fractions.

Supplementary figure 4: Bin3 overexpression fails to rescue *pof8Δ*-dependent telomere defects

A) Chromosome fusion PCR (top) following 2 restreaks on minimal media (one re-streak= 20-25 generations) from wild type and *pof8Δ* strains overexpressing an empty vector (pRep4x) or Bin3. The Trt1 gene was amplified from genomic DNA as a loading control (bottom).

B) Southern blot comparing telomere length from the same genomic DNA as (A). Genomic DNA was digested with EcoRI.

Supplementary figure 5: Validation of a Bin3 knockout strain confirms its viability

A) Schematic of Bin3 knockout (top) and PCR validation of the resultant strain using the indicated primers.
B) qRT-PCR of bin3 mRNA normalized to act1 mRNA for wild type and bin3Δ strains (mean ± standard error, paired t test * at p < 0.05, ** at p < 0.01, *** at p < 0.001, and **** at p<0.0001) (n=3 biological replicates). P-values over each bar represent results of a paired t-test with a wild type strain.
C) Backcrossing of bin3Δ and a wild type strain, followed by tetrad dissection, confirms viability of the bin3Δ strain on rich media. Tetrads from the same spore are arranged in columns. Resulting haploid colonies possessing phleomycin resistance are circled in red and colonies that didn't grow after tetrad dissection are circled in black.
D) qRT-PCR of TER1 normalized to act1 mRNA for wild type and bin3Δ strains transformed with an empty vector (3x) or Bin3 (mean ± standard error, paired t test * at p < 0.05, ** at p < 0.01, *** at p < 0.001, and **** at p<0.0001) (n=3 biological replicates). P-values over each bar represent results of a paired t-test with a wild type strain transformed with an empty vector (wild type + pRep3x).

wild type   TER1Δ   pof8Δ   bin3Δ

0.5 kB
0.4 kB
0.3 kB
0.2 kB
0.1 kB

Supplementary figure 6: Bin3 and Pof8 are not strictly required for telomere maintenance
Southern blot comparing ApaI-digested genomic DNA isolated after 5 restreaks on rich media. A lack of signal with a telomeric probe represents survivors that have lost telomeric DNA and circularized their chromosomes (*TER1Δ*).

Supplementary figure 7: Phylogenic distribution of Bin3 and Pof8 in fungi
Complete consensus cladogram describing the phylogenic relationships of 472 species representative of fungi phylum and classes and highlighting (using a color code) the distribution of Bin3 and Pof8 in these species. Only posterior probabilities inferior to 1 are shown.