1

Testing for pollinator recognition in multiple species of *Heliconia*

Dustin G. Gannon[1*], Adam S. Hadley[2,3], Urs G. Kormann[2,4], F. Andrew Jones[1,5], Matthew G.

Betts[2,6]

[1] Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon, USA

[2] Forest Biodiversity Research Network, Corvallis, Oregon, USA

[3] Department of Natural Resources and Energy Development, Government of New Brunswick,

Frederiction, Canada

[4] Ecological Research Group, Swiss Ornithological Institute, Sempach, Switzerland

[5] Smithsonian Tropical Research Institute, Apartado Postál 0843-03092, Panamá, Panamá

[6] Department of Forest Ecosystems and Society, Oregon State University, Corvallis, Oregon,

USA

20 *Corresponding author: gannondu@oregonstate.edu

Abstract

Many plants have evolved floral traits that, in effect, filter pollinator communities and promote pollination by efficient pollinators. Most documented pollinator filter traits act to deter or reduce visitation rates by a subsect of the community of floral visitors. However, a recently described pollinator filter termed 'pollinator recognition' (PR) acts at a stage after a pollinator visit. PR was first documented experimentally in *Heliconia tortuosa* whereby pollen tube germination – a proxy for reproduction – was enhanced following visits from morphologically specialized pollinators, but not generalists. This is thought to promote outcrossing among plants by preferentially investing in reproduction following visits by long-ranging hummingbirds with specialized bill shapes. To date, this plant behavior has only been described in *H. tortuosa*, but, if widespread, could have important ecological implications; given declines in abundances of specialist pollinators, visits by generalists would not buffer the loss of pollination services to plants with PR. We therefore tested for PR in four taxa spread widely across the Heliconiaceae.

We corroborated previous results that visits by long-billed, but not short-billed hummingbirds increased pollen tubes in *H. tortuosa* with aviary experiments that standardized pollen quality and minimized variation in pollen quantity. Across species, we found great variation in pollen tube responses to experimental treatments. For one species (*H. rostrata*), we found increased numbers of pollen tubes in those visited by hummingbirds compared to hand pollination alone, regardless of the visiting bird's bill morphology, indicating recognition of hummingbirds in general. In other cases, hummingbird visits decreased pollen tube counts compared to hand pollinations alone. Furthermore, our results could not substantiate any specific mechanism for pollinator recognition and highlight the need for further work on the complexities of and variability in reproductive strategies across plant taxa.

Introduction

Pollinator filters are floral traits that, in general, manipulate animal visitation patterns

presumably to promote conspecific pollen transfer or limit access to floral rewards (e.g., nectar)

for poor pollinators. For example, nectar that is distasteful to some pollinators will deter them

from visiting [1] and exploitation barriers, such as long corolla tubes that limit access to nectaries

[2,3], may make visitation unprofitable for some animals [4,5]. These filters reduce the richness

of pollinator species (or types within species) that visit a given flower species or type. However,

previous work with *Heliconia tortuosa* (Heliconiaceae) documented a cryptic pollinator filter

that could act at a stage after the pollinator visit to promote pollen germination and pollen tube

growth based on the identity, behavior, and bill morphology of hummingbird floral visitors [6].

In single-visit aviary experiments that controlled for variation in pollen deposition and

visitation rates by different pollinator species [6], the average number of pollen tubes that

germinated in a style (henceforth 'pollen tube rate') was nearly six times greater in flowers

visited by hummingbirds with bill shapes that are morphologically matched to the flowers (i.e.,

long, decurved bills) than in flowers visited by hummingbirds with mismatched bill shapes.

Furthermore, in a separate experiment, manual nectar removal showed higher pollen tube rates

than hand pollination alone. Betts et al. coined this behavior 'pollinator recognition' and posited

that nectar removal and pollen deposition by long-billed hummingbirds provides a cue for pollen

grain germination and pollen tube growth, thus reducing pollination efficiency by

morphologically mismatched hummingbirds that visit and transfer pollen but cannot access the

full volume of nectar at the base of the flower [6].

Betts et al. (18) speculated that pollinator recognition may be adaptive if it allows plants

to invest in reproduction following visits from high-quality pollinators (those more likely to

carry high-quality, outcrossed or unrelated pollen) and limit reproduction with the pollen

deposited by poor pollinators (those more likely to carry low-quality pollen). Despite receiving

visits from at least six hummingbird species, *H. tortuosa* specializes on long-billed

hummingbirds that are highly mobile [6,7] compared to the short-billed hummingbirds which

tend to defend territories and therefore move less. The mobile foraging behaviors of these birds

may make them more likely to carry high-quality pollen from geographically and genetically

distant sources [8], thus promoting outcrossing and genetic diversity in *H. tortuosa* populations.

We postulated that pollinator recognition may occur in other plant taxa, particularly in

relatively stable tropical systems with high pollinator diversity. Determining whether this is the

case is important for two reasons. First, we agree with Betts et al. [18] that filtering the short-

billed, territorial hummingbirds could promote outcrossing and enhance the genetic diversity of

pollen grains that reach the ovules, a hypothesis that is supported by landscape genetic studies of

H. tortuosa populations in Costa Rica [9,10]. Given the potential for fitness benefits, pollinator

recognition could be present in many related taxa and could be one means through which tight

morphological matching evolves despite apparently generalized interaction networks. Indeed,

hummingbird-*Heliconia* systems are often used as outstanding examples of morphological

matching (e.g., [11]).

Second, if the capacity for plants to actively filter floral visitors based on morphological

trait matching is widespread, this would have implications for the robustness of plant-pollinator

communities under climate and anthropogenic change [12,13]. Given local extinction or reduced

densities of morphologically matched pollinators, mismatched pollinators may alter their

foraging behaviors to exploit newly available resources [14–17]; however, visits from

mismatched pollinators would not compensate for the pollination services lost to a plant with a

pollinator recognition mechanism, even if they deposit pollen at the stigma. This could increase

the likelihood of coextinctions [12,18]. Thus, we sought to test for pollinator recognition in four

species distributed widely across the Heliconiaceae phylogeny as a first step in assessing the

generality of this cryptic pollinator filter.

Materials and Methods

Ethics statement

103 All experimental methods involving hummingbirds were approved by the Oregon State

104 University Animal Care and Use Committee (Animal Care and Use Permit 5020) and all

105 international research guidelines and practices were followed.

Study species

Heliconiaceae is a monogeneric family consisting of an estimated 200-250 species which

radiated rapidly c.a. 39-24 million years ago [19]. *Heliconia* species are rhizomatous perennial

herbs distributed widely throughout the Neotropics and on some South Pacific islands. Flowers

are situated within showy bracts and composed of six tepals, five of which are fused to create a

cylindrical perianth, the sixth peels back upon anthesis. A defining feature of the Heliconiaceae

is a staminode (modified stamen) that partially covers the opening to the nectar chamber at the

base of the perianth, which may need to be moved by a visiting animal when they extract the

nectar reward (though the mechanics of this have not be studied in detail). Flowers of the

Heliconiaceae last a single day from anthesis to dehiscence.

We targeted species that were common in the living collection at the Organization of

Tropical Studies Las Cruces Biological Station, Puntarenas Province, Coto Brus, Costa Rica, (8°

47′ 7″ N, 82° 57′ 32″ W) and could be found naturally or in ornamental gardens in the area. We

required that plants were setting seed when left unmanipulated, indicating that a viable pollen

source existed in the area, since previous work on mating systems in *Heliconia* suggests that the

hermaphroditic flowers of many species are self-incompatible to partially self-compatible, but

largely not selfing [6,9,20–23]. Furthermore, we required that wild, native hummingbirds could

be seen visiting the flowers of each target species in camera trap data [24] or during observation,

indicating that wild-caught hummingbirds would visit and drink from the flowers inside aviaries

despite the fact that many plant species in the collection are not native to Costa Rica. The plant

species that met these criteria included *H. hirsuta*, which is native to South America and

Trinidad [25], *H. rostrata*, native to western South America [25] but a common ornamental

throughout the tropics, and *H. wagneriana*, native to Costa Rica and Panama [26]. Furthermore,

because so little is known of this unusual plant behavior, we also sought to replicate the results of

the original study in the native *H. tortuosa,* an exercise rarely undertaken in experimental

ecology [27].

We selected two hummingbird species with different bill morphologies and foraging

behaviors as "treatments" in order to accentuate differences in morphological matching to and

nectar depletion from the range of flower shapes exemplified by the four *Heliconia* species (Fig

1). Green Hermit Hummingbirds (*Phaethornis guy*; GREH) are common traplining

hummingbirds in the region with long ($\bar{x} = 41.90$ mm, $s_x = 1.52$ mm), moderately decurved bills

($\bar{k} = 0.022$ mm^{-1}, $s_k = 0.004$ mm^{-1}, $n = 27$ birds of mixed sex, where k_i is the curvature of the

i^{th} bill measured as the inverse of the radius of the arc of the bill – see Temeles et al. [3]).

Rufous-tailed Hummingbirds (*Amazilia tzacatl*; RTAH) are common territorial hummingbirds

with short ($\bar{x} = 21.60$ mm, $s_x = 1.55$ mm, $n = 14$ birds of mixed sex), slightly decurved bills ($\bar{k} =$

0.016 mm^{-1}, $s_k = 0.002$ mm^{-1}; Fig 1).

Fig 1. Morphological mismatch between the hummingbird and *Heliconia* species used in

experiments. The average mismatch ($\widehat{\Delta}$) between a plant and hummingbird species was measured

as the Euclidean distance between a flower and a bird's bill in the $\mathbb{R}^2$ trait space, where one axis

was the total length of a bill or flower (mm) and the other was the radius of the arc along the

outside edge of the flower or bill (mm). We then computed the mean ($\widehat{\Delta}$) and standard deviation

(*s*) of the distances between each bill-flower pair, where *n* is the number of pairwise

comparisons.

Assuming that morphologically matched floral visitors increase the numbers of

successful pollen tubes for all *Heliconia* species, we predicted the following: 1) For *H.*

wagneriana and *H. tortuosa*, we would predict greater numbers of pollen tubes in flowers visited

by green hermit hummingbirds compared to rufous-tailed hummingbirds due to long and curved

flowers (Fig 1). 2) Because both *H. hirsuta* and *H. rostrata* have shorter, straighter flowers and

both hummingbird bill shapes approximate the shape of the flowers well (Fig 1), we would not

predict a large difference in the number of pollen tubes between flowers visited by green hermits

and those visited by rufous-tailed hummingbirds. We therefore used hand pollinations as a

control treatment in all experiments since hand pollinations do not replicate the physical

characteristics of a visit by a morphologically matched pollinator aside from pollen deposition.

Hence, we would predict the fewest pollen tubes in flowers pollinated by hand for all plant

species. Furthermore, this helped us control for potentially low genetic diversity in the pollen

pool since the control flowers (hand pollination only) and the treatment flowers (hand pollination

followed by a visit from a pollen-free hummingbird) both received pollen by hand from the same

donor.

Aviary experiments

To test whether pollen germination and tube growth is dependent on interactions with

morphologically matched floral visitors, we conducted 110 single-visit experiments ($n = 214$

flowers from 54 plants; see Table 1 for the number of replicates per treatment) with captive

hummingbirds inside portable aviaries. The aviaries measured 2 meters tall and one meter on a

side and could be quickly assembled around live plants (S1 File). In these experiments, we used

only virgin flowers that had been covered with mesh bags prior to anthesis in order to preclude

pollination by free-ranging pollinators. Flowers were not emasculated, however, due to low

numbers of pollen tubes in emasculated flowers in earlier experiments with plants in aviaries and

in natural settings (M. G. Betts and A. S. Hadley, *unpublished data*).

Table 1: Sample sizes for individual plants (grouping factor) and flowers (experimental units)

for each species × treatment combination.

	Hand pollination (HP)[*]	Hand pollination + short-billed bird (SB)	Hand pollination + long-billed bird (LB)	Hand pollination (HP)[**]	Hand pollination + pipette (BM)	Hand pollination + nectar extraction (HPNE)	Nectar extraction + hand pollination (NEHP)
H. hirsuta							
Plants	7	4	5	0	0	0	0
Flowers	11	5	7	0	0	0	0

H. rostrata							
Plants	19	13	10	13	7	15	5
Flowers	39	25	19	24	8	38	11
H. tortuosa							
Plants	10	8	6	10	7	10	4
Flowers	16	10	11	27	7	36	5
H. wagneriana							
Plants	12	6	11	11	7	8	4
Flowers	31	17	23	17	7	12	10

*Hand pollination controls for aviary experiments

**Hand pollination controls for nectar removal experiments

We selected inflorescences based on the availability of two virgin flowers and erected the aviary around the whole plant. We then randomly assigned one of the flowers as a control flower that received hand-pollination but no visit from a bird (HP treatment). The remaining flower was hand-pollinated with pollen from the same donor flower, then allowed a visit by either a pollen-free short-billed hummingbird (SB treatment; $n = 14$ _A. tzacatl_ individuals used in experiments) or a pollen free long-billed hummingbird (LB treatment; $n = 12$ _P. guy_ individuals used in experiments). To ensure the birds were free of pollen before the visit to the focal flower, we cleaned them using a soft paint brush and damp cotton swab under 20x magnification prior to releasing them into the aviary. Thus, flowers were the experimental units and individual plants were treated as a blocking effect to account for potential dependence among measurements on flowers from the same plant. Where possible, plants received all treatments (often on separate days).

By hand-pollinating all flowers using pollen from an arbitrarily selected pollen donor, we were able to control for differences in the quality of pollen delivered by the different pollinator

species. Indeed, we could not perfectly standardize the quantity of pollen grains at the stigmatic

surface because the size of *Heliconia* pollen grains makes it impractical to quantify the number

of grains in the field; however, we attempted to reduce variation in the quantity of pollen

available to the flowers by having the same experimenter apply pollen in an even layer across the

stigmatic surface with a toothpick under 20x magnification for every flower.

After the hummingbird visited the treatment flower (evidenced by bill insertion and a

clear attempt to feed from the flower), we terminated the experiment and checked the stigma

again to ensure that pollen was still present and in an even layer on the stigmatic surface before

again covering the flowers with mesh bags. All flowers were collected the following day, the

styles removed and preserved in formalin acetyl-acid, and scored for pollen tubes using epi-

florescence microscopy [6,20] (see S1 File for more information). All aviary experiments were

conducted during the 2018 and 2019 dry seasons (Feb-Mar).

Tests for a mechanism

We conducted additional experiments to test hypotheses of the mechanism of pollinator

recognition. Betts et al. [6] found increased pollen tube rates in flowers from which nectar was

removed compared to hand pollination alone. As an independent test of whether nectar removal

provides a cue to which plants respond, we manually extracted nectar from flowers of three of

the four species (*H. hirsuta* did not produce flowers regularly enough to conduct the full suite of

experiments) and compared pollen tube rates to control flowers that were hand-pollinated on the

same day. Alternatively, it is possible that the long-billed hummingbirds trigger a mechanical

cue [28] when they insert their bills into the flower. To test whether we could induce an increase

in pollen tube success rates using a mechanical stimulus, we molded a pipette tip to match the

curvature of the focal flower. We then inserted the pipette tip as a hummingbird would insert its

bill but did not remove any nectar. Finally, because we were unable to perfectly replicate the

timing of events in a natural pollinator visit in which nectar removal and pollen deposition

happen concurrently, we conducted some experiments in which we hand pollinated before

manually removing nectar and some in which we hand pollinated after removing nectar.

Differences in these pollen tube rates may indicate the importance of the timing of pollen

transfer and nectar removal or bill insertion (see S1 File for more detail).

Statistical methods

We separately analyzed pollen tube count data from each set of experiments (i.e., aviary

experiments as one dataset and nectar removal experiments as a second dataset). We fit

generalized linear mixed models (GLMMs) to both sets of data using the R package lme4 [29,30]

(see S1 File for full model specification). Our experiments were factorial with each plant species

receiving each treatment (4 plant species × 3 treatments = 12 means to estimate for the aviary

data; 3 plant species × 4 treatments = 12 means to estimate for the nectar removal data).

Furthermore, we included random effects for each plant (nested within plant species) to account

for potentially correlated observations that could arise from scoring pollen tubes in multiple

flowers from the same plant (since individual plants received more than one treatment).

Below, we report mean pollen tube counts per style estimated from the model fit for a

given treatment and plant species as $\hat{\lambda}$ with a subscript indicating the treatment and plant species.

We use *HP*, *SB*, *LB*, *HPNE*, *NEHP,* and *BM* to indicate the treatment (Table 1). Treatment codes

are as follows: *HP* identifies the hand-pollinated control flowers; *SB* indicates the treatment in

which we hand pollinated flowers, then allowed a clean, rufous-tailed hummingbird (short,

straight bill) to visit; *LB* indicates the treatment in which we hand pollinated flowers, then

allowed a pollen-free green hermit hummingbird (long bill) to visit; *HPNE* identifies the

treatment in which we hand pollinated the flowers then manually removed the nectar; *NEHP*

identifies the treatment in which we hand pollinated the flowers after removing nectar; and *BM*

identifies the treatment in which we inserted a pipette tip but did not attempt to remove nectar.

We use the letters *h*, *r*, *t*, and *w* to identify *H. hirstuta*, *H. rostrata*, *H. tortuosa*, and *H.*

wagneriana in the subscripts (respectively). We additionally report differences between

treatments as the fold change in pollen tube rates and use the notation $\hat{\lambda}_{i,j}/\hat{\lambda}_{k,j}$ for the fold

change between treatments i and k for plant species j. This is a natural way in which to report

differences between treatments given the log link function. Finally, we estimated ninety-five

percent confidence intervals from 1,000 bootstrapped mean predictions [29]. Confidence

intervals for the true parameters follow each estimate and are presented in square brackets.

Results

When we compared pollen tube counts in hand-pollinated control flowers (HP) and those

that were visited by a pollen-free, morphologically matched hummingbird, we found evidence

that a visit by a matched hummingbird increased pollen tube rates for *H. tortuosa* over hand

pollination alone (Fig 2). Pollen tube rates were 6.91 [3.56, 43.49] (estimate and 95% confidence

interval) times greater following visits from long-billed hummingbirds compared to the control

treatments with only hand pollination ($\hat{\lambda}_{HP,t} = 0.17 \, [7.21 \times 10^{-5}, 0.41]$ tubes per style;

$\hat{\lambda}_{LB,t} = 1.21 \, [0.56, 2.11]$ tubes per style). However, short-billed hummingbird visits did not

boost pollen tube rates for *H. tortuosa* much above the hand-pollinated controls ($\hat{\lambda}_{SB,t} =$

$0.36 \, [0.09, 0.80]$ tubes per style; Fig 2). Thus, the pollen tube rate in flowers visited by

morphologically matched hummingbirds was greater ($\hat{\lambda}_{LB,t}/\hat{\lambda}_{SB,t}$ = 3.32 [1.06, 10.33] times

greater) than the pollen tube rate in flowers visited by morphologically mismatched

hummingbirds (Figs 1&2).

Fig 2. Pollen tube rates in flowers that received hand pollination only (HP) and those visited by a

pollen-free hummingbird following hand pollination. Flowers were visited by either short-billed

(SB) rufous-tailed hummingbirds (*Amazilia tzacatl*) or long-billed (LB) green hermit

hummingbirds (*Phaethornis guy*). Error bars show the estimate plus or minus bootstrapped

standard errors. The grey, horizontal dashed lines show estimates ($\pm SE$) of the pollen tube rates

in flowers left open to free ranging pollinators.

In *H. rostrata* flowers, pollen tube rates in those visited by hummingbirds were greater

than hand pollination alone regardless of the bird species used in experiments. The estimated

rates were similar in flowers visited by long-billed hummingbirds and those visited by short-

billed hummingbirds, but were approximately double the rate in hand pollinated controls

($\hat{\lambda}_{HP,r}$ = 0.41 [0.22, 0.67] tubes per style; $\hat{\lambda}_{SB,r}$ = 0.75 [0.40, 1.16] tubes per style; $\hat{\lambda}_{LB,r}$ =

0.85 [0.45, 1.40] tubes per style; Fig 2).

For *H. hirsuta* and *H. wagneriana,* single visits from cleaned hummingbirds did not

enhance pollen tube success rates above hand pollination alone (Fig 2), and pollen tube rates in

H. wagneriana flowers that were visited by clean birds were actually reduced. The number of

pollen tubes per style in *H. wagneriana* flowers visited by green hermits were a little more than

half those of hand pollinations and short-billed hummingbird visits yielded pollen tube rates less

than half of hand pollination treatments ($\hat{\lambda}_{HP,w} = 1.39$ [0.91, 2.00] tubes per style; $\hat{\lambda}_{LB,w} = $

0.86 [0.51, 1.39] tubes per style; $\hat{\lambda}_{SB,w} = 0.56$ [0.23, 1.01] tubes per style; Fig 2).

When we experimentally removed nectar using pipette tips to test the hypothesis that

differential nectar removal may be the mechanism for pollinator recognition, the effect of nectar

removal on pollen tube rates for *H. rostrata* and *H. tortuosa* was negligible ($\hat{\lambda}_{HPNE,r} = $

0.25 [0.09, 0.47] tubes per style; $\hat{\lambda}_{HPNE,t} = 0.38$ [0.15, 0.75] tubes per style; Fig 3). We did find

weak evidence that our nectar removal treatments had a positive effect on pollen tube

germination relative to hand-pollinations alone for *H. wagneriana* ($\hat{\lambda}_{HP,w} = 0.90$, [0.46, 1.55]

tubes per style in the nectar removal data; $\hat{\lambda}_{HPNE,w} = 1.47$ [0.67, 2.70] tubes per style; Fig 3),

but this effect is driven in part by a small number of influential observations (>5 pollen tubes

found in 2 styles), and disappears after removing them, so we caution readers in their

interpretation of this result. Furthermore, we achieved higher pollen tube rates with hand

pollination alone during the aviary experiments ($\hat{\lambda}_{HP,w} = 1.39$ [0.91, 1.99] tubes per style from

aviary experiments) than in the nectar removal experiments.

Fig 3. Results from experimental tests of the effect of nectar depletion on pollen tube rates. We

used hand-pollination (HP) as a control treatment and compared pollen tube rates in flowers that

received the control treatment to those in flowers that received out-cross pollen by hand either

before (HP + nectar extraction) or after (Nectar extraction + HP) manual removal of the nectar in

the flower. As a test of whether pollen germination success could be boosted after the

mechanical stimulus of a hummingbird inserting its bill to drink from the flower, we tested for an

effect of pipette insertion without removing any nectar (HP + pipette). Error bars show estimates

of the mean plus or minus bootstrapped standard errors.

Inserting a hummingbird bill mimic (i.e., pipette tip) into flowers as a mechanical signal

without removing nectar also did not induce substantially higher pollen tube rates in any of the

tested species ($\hat{\lambda}_{BM,r} = 0.18$ [4.32×10^{-5}, 0.50] tubes per style; $\hat{\lambda}_{BM,t} = 0.34$ [$1.92 \times$

10^{-5}, 0.88] tubes per style; $\hat{\lambda}_{BM,w} = 1.25$ [$0.50, 2.39$] tubes per style; Fig 3), and, in all cases,

hand pollinating flowers after removing nectar resulted in the fewest pollen tubes per style out of

all treatments, generally about half of hand pollination alone ($\hat{\lambda}_{NEHP,r} = 0.08$ [$2.73 \times$

10^{-5}, 0.29] tubes per style; $\hat{\lambda}_{NEHP,t} = 0.20$ [1.12×10^{-6}, 0.74] tubes per style; $\hat{\lambda}_{NEHP,w} =$

0.33 [4.94×10^{-5}, 0.78] tubes per style; Fig 3).

Discussion

We tested whether pollinator recognition, a plant behavior first described in *Heliconia*

toruosa [6] in which plants may discriminate among floral visitors based on morphological

matching to the flower, may be present in more than one species of *Heliconia*. Our results

substantiate the findings of Betts et al. [6] that the number of successful pollen tubes in *H.*

tortuosa flowers is greater in those visited by morphologically matched hummingbirds compared

to those visited by mismatched birds after experimentally standardizing pollen quality, visit

frequency, and minimizing variation in pollen quantity. However, we did not find strong support

for any of the hypothesized mechanisms of recognition and found different responses to our

aviary treatments in each of the four *Heliconia* species we tested, highlighting the poorly

understood complexities of pollination in *Heliconia* plants.

The results from our aviary experiments with *H. tortuosa* are highly congruent with those

of Betts et al. [6] and support that plants specialize on long-billed, traplining hummingbirds.

Pollinator recognition in *H. tortuosa* is likely an additional pollinator filter acting in conjunction

with morphological barriers that often result in only imperfect resource partitioning by floral

visitors [4,31–33]. However, this could have strong ecological implications if disturbances

reduce the abundance of morphologically matched hummingbirds and the rewiring of the

remaining community of floral visitors fails to buffer the loss of pollination services to *H.*

tortuosa plants. In forest fragments around Coto Brus, mismatched hummingbirds account for

c.a. 10% of honest visits (those in which the visitor contacts the reproductive organs of the

flower) to *H. tortuosa* (K. Leimburger, *unpublished data*), but this proportion likely increases in

isolated fragments where morphologically matched hummingbirds are less common [34–36].

The combination of fragmentation and pollinator recognition therefore has a strong ecological

impact on *H. tortuosa* populations, which show reduced seed sets in fragmented forests

compared to those in continuous forest [37], presumably due to a paucity of morphologically

matched hummingbirds.

In *H. rostrata* styles, visits from clean hummingbirds to hand-pollinated flowers also

increased pollen tube rates, but regardless of the bird species. The estimated pollen tube rate in

H. rostrata flowers that were visited by long-billed hummingbirds was nearly identical to that

estimated for flowers visited by short-billed hummingbirds (Fig 1). However, given the relatively

short, straight corolla of *H. rostrata* (Fig 1), both hummingbird species we used for experiments

were able to achieve strong morphological matches and might not be expected to differ in their

visitation characteristics, such as nectar consumption. We did not destructively sample flowers

after hummingbird visits to measure the nectar remaining, but both species of birds can be seen

drinking from *H. rostrata* flowers in recorded videos (S1 and S2 videos).

In Peru, seven hummingbird species of various sizes and with various bill shapes have

been observed visiting *H. rostrata*, but nothing is known of their pollination efficiencies [38].

Based on our results showing increased pollen tube rates in bird-visited flowers compared to

hand pollination, we posit that *H. rostrata* could filter visits from animals without

complementary morphologies, such as insects. This idea is supported by data from Janeček et al.

[23] who recorded olive sunbirds (*Cyanomitra olivacea*) and Camaroon sunbirds (*Cyanomitra*

oritis) visiting *H. rostrata* flowers in South Africa where it has been introduced. These authors

found that *H. rostrata* flowers left open to visits from sunbirds had extremely low pollen tube

rates, as did hand-pollinated flowers. Further work in the native range of *H. rostrata* is necessary

to determine if *H. rostrata* cryptically filters the community of floral visitors.

We did not detect strong differences among treatments in *H. hirsuta* flowers, which all

showed pollen tube rates in the range of open pollination pollen tube rates (grey lines in Fig 2).

Interestingly, *H. hirsuta* flowers visited by long-billed hummingbirds following hand pollination

showed the lowest pollen tube rate. Similarly, we found reduced pollen tube rates in *H.*

wagneriana flowers that were visited by hummingbirds relative to hand-pollinations alone (Fig

2). While the mechanisms underlying these results in *H. wagneriana* and *H. hirsuta* flowers

remain unclear, we identified one way in which these species differ from the others that could

produce this result. Gannon et al. [24] discovered that *H. wagneriana* plants have a mechanism

for keeping the anthers protected within the perianth and then rapidly extending them as a

hummingbird visits. This is thought to protect pollen from desiccation and/or increase pollen

transfer to pollinators during the first visit. Once exposed, however, pollen grains desiccate

relatively quickly, and often fail to adhere to the stigmatic surface. While *H. hirsuta* anthers

protrude from the flower upon anthesis, pollen also appeared to dry quickly and often failed to

cling to the stigmatic surface. This may make the pollen grains of *H. wagneriana* and *H. hirsuta*

easy to dislodge. While we checked that pollen was still present on the stigma after a bird visited,

the size of *Heliconia* pollen makes exact quantification in the field infeasible. Thus, it is possible

that reduced outcross pollen loads after the birds visited resulted in reduced pollen tube counts

relative to hand pollination alone.

Using camera traps, Gannon et al. [24] found that c.a. 97% of the visits to open *H.*

wagneriana flowers around Las Cruces were by traplining species with morphologically

matched bill shapes. Similarly, Snow and Snow [39] report only green hermit and rufous-

breasted hermit (*Glaucis hirsutus*) visitors at *H. hirsuta* flowers in Trinidad (part of its native

range), both of which have well-matched bills. Given a relatively specialized set of floral visitors

to these two species, it seems unlikely that any cryptic recognition mechanism would adaptively

filter the community of floral visitors.

393 The mechanism of pollinator recognition

Previous work demonstrated increased pollen tube rates with manual nectar extraction

treatments compared to hand pollinations alone in *H. tortuosa* [6]. Betts et al. [18] hypothesized

that, because birds with well-matched bill morphologies can drain the nectar chamber but those

with mismatched bills often cannot [5,6], nectar removal could provide a cue to which plants

respond to promote successful pollen tube growth. We were unable to reproduce an increase in

pollen tube rates using manual nectar removal techniques, despite the use of identical nectar

removal methods (aside from the experimenter). While we found increased pollen tube rates in

H. wagneriana flowers following manual nectar removal (Fig 3), this result is tenuous because it

is in part driven by a few influential observations. Furthermore, the pollen tube counts in the

flowers from which we removed nectar are approximately equal to the pollen tube counts in our

hand pollinated flowers from the aviary experiments (Figs 2 and 3).

In summary, the evidence that nectar removal provides the cue for pollinator recognition

in *H. tortuosa* is equivocal and further experiments are necessary to verify nectar removal or

establish a new mechanism. We refrain from speculating on alternative mechanisms of

recognition, but instead suggest follow-up experiments that could help to disentangle the

apparent complexities of this system: 1) continued aviary experiments with Heliconia species in

in their native ranges with native floral visitors; 2) comparative studies of the mechanics of visits

to *H. tortuosa* flowers by different hummingbird species using high-speed videography to

identify potential mechanisms for higher pollen tube rates by morphologically matched species;

3) enhanced experimental techniques that can increase the consistency of manual nectar

extraction and test alternative potential mechanisms linked to morphological

matching (including, but not limited to, nectar removal, pollen deposition/removal, and other

aspects of visit mechanics).

Conclusions

Our results help to highlight the complexities of pollination and pollen germination

success. Despite our detailed and manipulative experiments, we found great variability in the

number of germinated pollen tubes depending on the floral visitor and plant species. We

therefore add to calls for more detailed and manipulative experiments to assess realized

pollination network structure and vulnerability to disturbance that take into account single-visit

pollination efficiency [40,41]. Furthermore, our results highlight the necessity and opportunity

for further study of pollinator recognition in *Heliconia*. Further research into the mechanism of

recognition in *H. tortuosa* and the generality of pollinator recognition in the Heliconiaceae is

warranted. Notably, Pedersen and Kress [21] report a c.a. four-fold increase in pollen tube rates

in *Heliconia paka* flowers that were visited by honeyeaters compared to those pollinated by

hand. These results would be consistent with what we would predict for *H. paka* given a

pollinator recognition mechanism. More generally, Young and Young [42] report that hand-

pollinated flowers had reduced reproductive output compared to open-pollinated flowers for 17

of 52 plant species from highly divergent lineages. We know of no follow-up experiments with

these or related taxa, but we urge others to conduct similar experiments to those presented here

to examine the potential for cryptic specialization in other pollination systems.

Acknowledgements

437   We thank C. Dowd, E. Sandi, M. Atencio, C. Tortorelli, and G. Doyle for invaluable

438 assistance with field experiments, and the staff of the Las Cruces Biological Station for

439 maintaining an excellent location in which to conduct research. We thank N. Waser, R. Sargent,

440 J. Lau, and seven anonymous reviewers for comments on previous versions of the manuscript.

Data accessibility

443   Data from pollination experiments and all R code necessary to reproduce the results can

444 be found on a public Github repository (https://github.com/Dusty-Gannon/PR-in-Heliconia).

Author contributions

M.G.B., A.S.H., and D.G.G. designed experiments. D.G.G. analyzed the data and wrote

the original version of the manuscript. All authors contributed to data collection and critical

review of the manuscript.

References

1. Stevenson PC, Nicolson SW, Wright GA. Plant secondary metabolites in nectar: impacts on
pollinators and ecological functions. Funct Ecol. 2017;31(1):65–75.

2. Gill FB. Ecological Fitting: Use of Floral Nectar in Heliconia stilesii Daniels by Three
Species of Hermit Hummingbirds. Condor. 1987;89(4):779–87.

3. Temeles EJ, Koulouris CR, Sander SE, Kress WJ. Effect of flower shape and size on
foraging performance and trade-offs in a tropical hummingbird. Ecology. 2009;90(5):1147–
61.

4. Rodríguez-Gironés MA, Sun S, Santamaría L. Passive partner choice through exploitation
barriers. Evol Ecol. 2015;29(3):323–40.

5. Temeles EJ, Liang J, Levy MC, Fan Y-L. Floral isolation and pollination in two
hummingbird-pollinated plants: the roles of exploitation barriers and pollinator competition.
Evol Ecol. 2019;33:481–97.

6. Betts MG, Hadley AS, Kress WJ. Pollinator recognition by a keystone tropical plant. PNAS.
2015;112(11):3433–8.

7. Stiles FG, Wolf LL. Hummingbird Territoriality at a Tropical Flowering Tree. Auk.
1970;87(3):467–91.

8. Wright S. Isolation by Distance. Genetics. 1943;28(2):114–38.

9. Torres-Vanegas F, Hadley AS, Kormann UG, Jones FA, Betts MG, Wagner HH. The
Landscape Genetic Signature of Pollination by Trapliners: Evidence From the Tropical
Herb, Heliconia tortuosa. Front Genet. 2019;10:1206.

10. Torres☐Vanegas F, Hadley AS, Kormann UG, Jones FA, Betts MG, Wagner HH. Tropical
deforestation reduces plant mating quality by shifting the functional composition of
pollinator communities. J Ecol. 2020;109(4):1730–46.

11. Temeles EJ, Kress WJ. Adaptation in a Plant-Hummingbird Association. Science.
2003;300(5619):630–3.

12. Kaiser☐Bunbury CN, Muff S, Memmott J, Müller CB, Caflisch A. The robustness of
pollination networks to the loss of species and interactions: a quantitative approach
incorporating pollinator behaviour. Ecol Lett. 2010;13(4):442–52.

13. Thébault E, Fontaine C. Stability of Ecological Communities and the Architecture of
Mutualistic and Trophic Networks. Science. 2010;329(5993):853–6.

14. Brosi BJ, Briggs HM. Single pollinator species losses reduce floral fidelity and plant
reproductive function. PNAS. 2013;110(32):13044–8.

15. Temeles EJ, Newman JT, Newman JH, Cho SY, Mazzotta AR, Kress WJ. Pollinator
Competition as a Driver of Floral Divergence: An Experimental Test. PLOS ONE. 2016
Jan 27;11(1):e0146431.

16. Bezemer N, Hopper SD, Krauss SL, Phillips RD, Roberts DG. Primary pollinator exclusion
has divergent consequences for pollen dispersal and mating in different populations of a
bird-pollinated tree. Mol Ecol. 2019;28(22):4883–98.

17. Temeles EJ, Bishop GA. A hurricane alters pollinator relationships and natural selection on
an introduced island plant. Biotropica. 2019;51(2):129–38.

18. Rezende EL, Lavabre JE, Guimarães PR, Jordano P, Bascompte J. Non-random
coextinctions in phylogenetically structured mutualistic networks. Nature.
2007;448(7156):925–8.

19. Iles WJ, Sass C, Lagomarsino L, Benson-Martin G, Driscoll H, Specht CD. The phylogeny
of Heliconia (Heliconiaceae) and the evolution of floral presentation. Mol Phylogenet Evol.
2016;117:150–67.

20. Kress WJ. Self-Incompatibility in Central American Heliconia. Evolution. 1983;37(4):735–
44.

21. Pedersen LB, Kress WJ. Honeyeater (Meliphagidae) pollination and the floral biology of
Polynesian Heliconia (Heliconiaceae). Pl Syst Evol. 1999;216(1):1–21.

22. Schleuning M, Templin M, Huamán V, Vadillo GP, Becker T, Durka W, et al. Effects of
Inbreeding, Outbreeding, and Supplemental Pollen on the Reproduction of a Hummingbird-
pollinated Clonal Amazonian Herb. Biotropica. 2011;43(2):183–91.

23. Janeček Š, Chmel K, Gómez GU, Janečková P, Chmelová E, Sejfová Z, et al. Ecological
fitting is a sufficient driver of tight interactions between sunbirds and ornithophilous plants.
Ecol Evol. 2020;10(4):1784–93.

24. Gannon DG, Kormann UG, Hadley AS, Betts MG, Jones FA. The "jack-in-the-box" stamens
of Heliconia wagneriana (Heliconiaceae). Ecology. 2018;99(2):488–90.

25. Bernal R, Gradstein S, Celis M. Catálogo de plantas y líquenes de Colombia. Bogotá:
nstituto de Ciencias Naturales, Universidad Nacional de Colombia; 2016.

26. Stiles G. Notes on the natural history of Heliconia (Musaceae) in Costa Rica. Brenesia.
1979;15:151–80.

27. Fraser H, Barnett A, Parker TH, Fidler F. The role of replication studies in ecology. Ecol
Evol. 2020;10(12):5197–207.

28. Braam J. In touch: plant responses to mechanical stimuli. New Phytol. 2005;165(2):373–89.

29. Bates D, Machler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4.
J Stat Softw. 2015;67(1):1–48.

30. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna,
Austria: R Foundation for Statistical Computing; 2020. Available from: https://www.R-
project.org/

31. Feinsinger P. Organization of a Tropical Guild of Nectarivorous Birds. Ecol Monogr.
1976;46(3):257–91.

32. Vlašánková A, Padyšáková E, Bartoš M, Mengual X, Janečková P, Janeček Š. The nectar
spur is not only a simple specialization for long-proboscid pollinators. New Phytol.
2017;215(4):1574–81.

33. Metelmann S, Sakai S, Kondoh M, Telschow A. Evolutionary stability of plant–pollinator
networks: efficient communities and a pollination dilemma. Ecol Lett. 2020;23(12):1747–
55.

34. Kormann U, Scherber C, Tscharntke T, Klein N, Larbig M, Valente JJ, et al. Corridors
restore animal-mediated pollination in fragmented tropical forest landscapes. Proc Royal
Soc B. 2016;283(1823):20152347.

35. Hadley AS, Frey SJK, Robinson WD, Betts MG. Forest fragmentation and loss reduce
richness, availability, and specialization in tropical hummingbird communities. Biotropica.
2018;50(1):74–83.

36. Morrison BML, Mendenhall CD. Hummingbird–Plant Interactions Are More Specialized in
Forest Compared to Coffee Plantations. Diversity. 2020;12(4):126.

37. Hadley AS, Frey SJ, Robinson WD, Kress WJ, Betts MG. Tropical forest fragmentation
limits pollination of a keystone understory herb. Ecology. 2014;95(8):2202–12.

38. Dalsgaard B, Maruyama PK, Sonne J, Hansen K, Zanata TB, Abrahamczyk S, et al. The
influence of biogeographical and evolutionary histories on morphological trait-matching
and resource specialization in mutualistic hummingbird-plant networks. Funct Ecol.
2021;35:1120–33.

39. Snow BK, Snow DW. Feeding Niches of Hummingbirds in a Trinidad Valley. J Anim Ecol.
1972;41(2):471–85.

40. Ballantyne G, Baldock KCR, Willmer PG. Constructing more informative plant–pollinator
networks: visitation and pollen deposition networks in a heathland plant community. Proc
Royal Soc B. 2015;282(1814):20151130.

41. King C, Ballantyne G, Willmer PG. Why flower visitation is a poor proxy for pollination:
measuring single-visit pollen deposition, with implications for pollination networks and
conservation. Methods Ecol Evol. 2013;4(9):811–8.

42. Young HJ, Young TP. Alternative Outcomes of Natural and Experimental High Pollen
Loads. Ecology. 1992;73(2):639–47.

Supporting information

S1 video. Green hermit visit to *H. rostrata.* Green hermit hummingbird (*Phaethornis guy*)

visiting and drinking nectar from a *Heliconia rostrata* flower.

S2 video. Rufous-tailed hummingbird visit to *H. rostrata*. Rufous-tailed hummingbird

(*Amazilia tzacatl*) visiting and drinking nectar from a *Heliconia rostrata* flower.

S1 File. Supplementary methods. This file contains additional information on the field,

laboratory, and statistical methods.

SB - *Amazilia tzacatl* ♂

$\hat{\Delta} = 45.75$ mm
$s = 7.25$ mm
$n = 490$

$\hat{\Delta} = 32.70$ mm
$s = 4.80$ mm
$n = 182$

$\hat{\Delta} = 35.19$ mm
$s = 5.47$ mm
$n = 140$

$\hat{\Delta} = 50.52$ mm
$s = 4.39$ mm
$n = 126$

H. tortuosa

H. rostrata

H. hirsuta

H. wagneriana

$\hat{\Delta} = 21.75$ mm
$s = 5.84$ mm
$n = 945$

$\hat{\Delta} = 16.09$ mm
$s = 7.25$ mm
$n = 351$

$\hat{\Delta} = 27.09$ mm
$s = 13.79$ mm
$n = 270$

$\hat{\Delta} = 34.07$ mm
$s = 6.44$ mm
$n = 243$

1 cm

LB - *Phaethornis guy* ♂