Naïve individuals promote collective exploration in homing pigeons

Gabriele Valentini[1,2], Theodore P. Pavlic[2,3,4,5,6], Sara Imari Walker[1,3,6], Stephen C. Pratt[2,6], Dora Biro[7,8] & Takao Sasaki[9]

[1] Arizona State University, School of Earth and Space Exploration, Tempe, AZ 85287, USA
[2] Arizona State University, School of Life Sciences, Tempe, AZ 85287, USA
[3] Arizona State University, Beyond Center for Fundamental Concepts in Science, Tempe, AZ 85287, USA
[4] Arizona State University, School of Computing, Informatics, and Decision Systems Engineering, Tempe, AZ 85287, USA
[5] Arizona State University, School of Sustainability, Tempe, AZ 85287, USA
[6] Arizona State University, ASU–SFI Center for Biosocial Complex Systems, Tempe, AZ 85287, USA
[7] University of Oxford, Department of Zoology, Oxford, OX1 3PS, UK
[8] University of Rochester, Department of Brain and Cognitive Sciences, Rochester, NY 14627, USA
[9] University of Georgia, Odum School of Ecology, Athens, GA 30602, USA

Abstract. Group-living animals that rely on stable foraging or migratory routes can develop behavioural traditions to pass route information down to inexperienced individuals. Striking a balance between exploitation of social information and exploration for better alternatives is essential to prevent the spread of maladaptive traditions. We investigated this balance during cumulative route development in the homing pigeon *Columba livia*. We quantified causal interactions within pairs of birds in a transmission-chain experiment and determined how birds with different levels of experience contributed to the exploration–exploitation trade-off. Newly introduced naïve individuals were initially more likely to initiate exploration than experienced birds, but the pair soon settled into a pattern of alternating leadership with both birds contributing equally. Experimental pairs showed an oscillating pattern of exploration over generations that might facilitate the discovery of more efficient routes. Our results introduce a new perspective on the roles of leadership and information pooling in the context of collective learning.

Introduction

The coordinated motion of groups is a widespread phenomenon observed in multiple taxa (Vicsek and Zafeiris 2012). Among other adaptive advantages, such as increased energetic efficiency and decreased odds of predation (Krause and Ruxton 2002), collective motion also allows group members to increase their sensory and cognitive capacity (Berdahl et al. 2013; Gelblum et al. 2020) and to acquire valuable social information for navigation (Couzin 2009; Couzin et al. 2011). In many animals, this social information concerns well-established foraging or migratory routes that can, in some species, persist over successive generations (Helfman and Schultz 1984; Sasaki and Biro 2017; Jesmer et al. 2018). Knowledge and skills that accumulate over generations can provide groups with an enhanced ability to solve difficult problems (Biro, Sasaki, and Portugal 2016). Not only can later generations build on the success of earlier ones, but

the introduction of new members, even those with no prior knowledge, adds diversity that can enhance the

group's behavioural solutions (Mehlhorn et al. 2015). As is often the case (Hills et al. 2015), behavioural

patterns that lead to a search for improvement, whether individually, socially, or over multiple generations,

involve an exploration–exploitation trade-off. In navigation problems, both solitary individuals and groups

have to balance between exploiting previously acquired information necessary to navigate a known route

and exploring for additional information that might allow them to approach the optimal route (Fu and Gray

2006). However, how moving collectives compromise between these tasks has received limited attention.

Understanding the exploration–exploitation trade-off is complicated by ambiguity about group

leadership (Couzin et al. 2005; Garland et al. 2018). Although some collectives (*e.g.*, ants, honeybees) can

allocate certain individuals to spatial exploration while others continue to exploit accumulated information

(Hills et al. 2015), individuals in cohesively moving groups are highly coupled and can only benefit from

compromising between exploring and exploiting if they do so in unison. If they are to stay together, the

group must reach consensus between following a known route or departing from it to find better routes,

foraging patches, or temporary resting locations. Elucidating whether different group members contribute

differently to this process is crucial to understanding how groups compromise between exploration and

exploitation.

We investigate this question in the context of navigation through natural landscapes using the homing

pigeon *Columba livia* as our model system. After successive homing journeys from a given release site,

pigeons develop stable idiosyncratic routes that are followed with high fidelity (Meade, Biro, and Guilford

2005; Guilford and Biro 2014). These birds rely on sequences of localized visual landmarks to recapitulate

familiar yet individually distinct routes (Biro, Meade, and Guilford 2004; Meade, Biro, and Guilford 2005).

Each route is learned in a gradual process starting with an exploration phase that samples new landmarks,

from which the bird eventually converges upon a stable sequence of landmarks (Biro, Meade, and Guilford

2004). Experiments with paired birds show that route information can be passed from experienced birds to

naïve individuals through social learning (Pettit, Flack, et al. 2013) and can be modified through information

pooling when individuals with different idiosyncratic routes share information to reach a compromise

between their routes (Biro et al. 2006). Although learning generally improves route efficiency, both social

learning and information pooling tend to reach a plateau beyond which further improvement in efficiency is

not seen. However, for birds flying together, the introduction of a naive individual in place of an experienced

one effectively leads to the resumption of exploratory behaviour and further route improvement (Sasaki and

Biro 2017). Yet, it remains unclear to what extent a bird's prior experience influences the balance between

exploration and exploitation and how birds with potentially different route preferences jointly shape a route.

Indeed, the mechanisms underlying how different individual preferences are combined into a collective

outcome is one of the key foci in studies of collective animal behaviour. Broadly, group decisions can range

from despotic with a single leader to democratic in which input from different individuals is aggregated to

reach consensus (Conradt and Roper 2003). Evidence of both despotic and democratic decisions exists in

homing pigeons (Biro et al. 2006; Nagy et al. 2010; Jorge and Marques 2012). When leadership is defined

as disproportionate input into collective navigational decisions, either through spatial position (Pettit, Perna, et al. 2013), route similarity (Flack et al. 2012), or directional correlation delay (Nagy et al. 2010), a number of different factors have been shown to play a role in it. Leadership dynamics are influenced by individual differences among birds in fidelity to their own routes (Freeman et al. 2011), their typical flight speed (Pettit et al. 2015), their personality (Sasaki et al. 2018), as well as their level of experience (Flack et al. 2012). Moreover, equally experienced birds are known to come to a compromise by averaging their idiosyncratic routes so long as the pair's route remains within a threshold distance from each bird's favoured one – a low level of conflict. Higher levels of conflict lead instead to a splitting of the pair or to the emergence of a single leader (Biro et al. 2006). Nonetheless, experience alone is unable to fully recover the leadership structure characteristic of larger flocks (Watts et al. 2016). Spatial position offers some insight into leadership: on average, birds flying closer to the front of the flock have a stronger influence on the flock's directional choices than birds flying at the back (Nagy et al. 2010; Pettit, Perna, et al. 2013). Even so, the moment-to-moment relationship between leadership and level of experience remains unclear.

Leader–follower interactions of this sort can be accurately captured using information-theoretic measures that quantify causal relations in terms of predictive information (Butail, Mwaffo, and Porfiri 2016; Kim et al. 2018; Crosato et al. 2018; Ray et al. 2019; Valentini et al. 2020). One of these measures, *transfer entropy,* quantifies information about the future behaviour of a focal individual that can be obtained exclusively from knowledge of the present behaviour of another subject (Schreiber 2000). Transfer entropy measures information transferred from the present of the sender to the future of the receiver (Lizier and Prokopenko 2010). It explicitly accounts for autocorrelations characteristic of individual birds' trajectories (Mitchell et al. 2019) by discounting predictive information available from the sender's present that is already included in the receiver's past. Furthermore, it does not require a model of how sender and receiver interact, and it is well suited to study social interactions both over space and over time (Lizier, Prokopenko, and Zomaya 2008). Consequently, transfer entropy can capture causal interactions due not only to alignment forces (Nagy et al. 2010) but also to attraction and repulsion forces that result in temporarily unaligned states (Pettit, Perna, et al. 2013).

We study collective decision making and the exploration–exploitation trade-off using an experimental analysis of cumulative route development in homing pigeons (Sasaki and Biro 2017). In these experiments, pairs consisting of a naïve and an experienced bird were required to successively solve the same homing task a fixed number of times. This set of paired flights allowed the naïve bird to acquire knowledge of localized visual landmarks necessary for homing. The more experienced bird was then replaced with a new naïve individual and the learning process was repeated through five generations of replacement in a transmission-chain design. Route efficiency was measured as the ratio of the beeline distance between the release site and the home loft (*i.e.*, the ideal optimum) and the actual distance travelled by birds. The results showed that, although homing efficiency dropped considerably every time a new naïve bird was introduced, transmission-chain pairs continued to improve within and over generations, eventually outperforming both solo and fixed-pair controls (respectively, 0.92 efficiency versus 0.83 and 0.85). In contrast, the efficiency

of solo and fixed pairs plateaued after they had first established their idiosyncratic routes (around the 9^{th}–
10^{th} release for the former and the 7^{th}–8^{th} release for the latter).

The continued improvement seen in transmission chains might result from a variety of decision-making
mechanisms ranging from fully despotic to increasingly democratic. A simplified perspective of this
continuum allows us to consider four alternative hypotheses. In two of these alternatives, a single despotic
leader, either the naïve or the experienced bird, determines the entire homing route. Whereas evidence of
social learning (Sasaki and Biro 2017) suffices to dismiss the possibility of leadership by the naïve
individual, leadership by the experienced individual could still be the only process in place if social learning
is unidirectional and the naïve individual merely triggers the experienced bird to resume and lead
exploration. Under the other two hypotheses, birds could pool their personal information by means of
democratic processes based on moment-by-moment integration of individual preferences or transient,
alternating leadership (Conradt 2012). The third hypothesis entails the experienced bird contributing only
its past route information and relying instead on the naïve individual for the discovery of route innovations.
If this hypothesis holds, we expect the naïve bird to disproportionally lead phases of exploration. Otherwise
(fourth hypothesis), both experienced and naïve birds might contribute through exploration to the discovery
of new information.

We discriminated between these alternative hypotheses by using transfer entropy to reveal the extent
to which birds influence each other and to investigate if relative spatial position can accurately predict
leader–follower dynamics. On this basis, we studied the contribution of each bird to the exploration–
exploitation trade-off over different stages of route development. This exploration–exploitation perspective
of homing route development allowed us to characterize the efficiency of choices made by birds over the
course of the experiment and to shed light on the superior performance of experimental pairs with respect
to solo and fixed pairs controls.

Results

Birds pool information. The social-learning hypothesis under which the naïve bird passively copies the
idiosyncratic route of the experienced one (*i.e.*, the despotic leader) entails a transfer of information that is
unidirectional – from the experienced to the naïve bird. Instead, under the two alternative hypotheses based
on democratic decision-making, the two birds rely on bidirectional information transfer to pool information
and increase the efficiency of their route (Pettit, Flack, et al. 2013; Sasaki and Biro 2017). We rejected the
unidirectional social-learning hypothesis by finding causal evidence of information pooling: the naïve bird
actively influenced the behaviour of the experienced one for a large portion of the parameter space
(Figure S1). As is common practice with these measures (Porfiri 2018), we selected the parameter
configuration that maximized the total transfer of information between the two birds (one sample every 0.2
seconds, $k = 10$) . This was maximal for the shortest sampling period (*i.e.*, prediction interval) of 0.2
seconds and progressively decreased towards 0 for larger periods up to 4 seconds, indicating that the effect
of an interaction between birds was transitory and lasted for a limited period of time. Using this configuration,

we compared measurements of information transfer against those of a surrogate dataset created by pairing

trajectories of birds that were not flown together. We found that levels of mutual influence between birds

that flew together were significantly higher than those observed in the surrogate dataset both overall and

for each generation separately (Mann–Whitney–Wilcoxon, columns 2 and 3 of Table S1).

(a)   (b)

Figure 2. Panel (a) shows the net predictive power of the two birds, $\tilde{T}_{E \to N} - \tilde{T}_{N \to E}$, over generations, where $\tilde{T}_{X \to Y} =$

$\frac{T_{X \to Y}}{H_Y} \cdot 100$ *with $X, Y \in \{E, N\}$, highlighting which bird is more informative (purple for the experienced bird, green for the*

naïve bird). The red line corresponds to a linear fit over generations using the Theil–Sen estimator. Panel (b) shows

the predictive power of naïve and experienced birds over generations measured as the percentage $\tilde{T}_{X \to Y}$ of uncertainty

reduction about one bird given by the other.

During the first two generations of paired flights (Figure 2a, paired analysis), when there was a large margin

to improve the efficiency of the pair's trajectory, the naïve bird was more informative than the experienced

one, evidenced by a stronger influence over the latter. At generation 4 there was a balance between the

two birds whereas the experienced bird eventually became the better source of predictive information in

the last generation. A linear fit over generations of the paired comparison (Figure 2a, red line) showed an

increasing influence of the experienced bird over the naïve one (Theil–Sen estimator, slope 0.534, $p <$

0.001). Additionally, a non-paired comparison of the same results revealed that, although the naïve bird

had, on average, a marginally higher predictive power than that of the experienced one (18.7–21.4% versus

17.7–20.5%), both birds explained a large portion of each other's behaviour (Figure 2b) suggesting non-

trivial leadership dynamics. These results do not show whether different levels of experience within the pair

led to asymmetric contributions of birds to route development, with the experienced bird providing only its

past route information and the naïve bird in charge of discovering route innovations, or if both birds

contributed to the exploration for possible route alternatives. To discriminate between these remaining

hypotheses, we first developed the means to evaluate leadership on a moment-to-moment basis.

(a)   **(b)**

Figure 3. Panel (a) shows sample flight trajectories for a number of different releases, r, of the same pair of birds. Colors highlight which bird is ahead of the other during different segments of the route. Panel (b) shows the local transfer of information (mean and 0.95 confidence interval) between the experienced bird and the naïve one as a function of their relative distance (colors represent the direction of information transfer) estimated using smoothed conditional means.

Relative position determines temporary leadership. Consistent with information sharing within each pair, we found that experienced and naïve birds repeatedly switched their positions at the front and back of the pair (Figure 3a). Previous studies found evidence that birds that spent, on average, more time at the front of the flock had a tendency to assume leadership roles (Nagy et al. 2010; Pettit, Perna, et al. 2013). To see whether this average relationship between leadership and position holds at each point in time, we investigated the spatiotemporal dynamics of information transfer (Lizier, Prokopenko, and Zomaya 2008). We did so by considering the amount of predictive information obtained by each bird as a function of the distance from the experienced to the naïve bird projected over their mean direction of motion (Figure 3b). We found that within a distance of up to 30 meters, the bird flying ahead was consistently more informative than that flying in the back. This is not only further evidence that the bird flying ahead acts as the leader, influencing the path of the follower behind it, but, because of its finer grain, it also enables relative distance between birds to be used as a (more parsimonious) moment-to-moment measure of causal influence within the pair.

The experienced and the naïve bird alternated leading segments of the route whose duration was stochastic and resembled a log-normal distribution (see Supplementary Material). Although the naïve bird flew at the front of the pair for longer segments (Whitney–Mann–Wilcoxon, $p < .047, W = 56865638$), the difference was very small (0.3 seconds) and largely driven by the flights of one generation. Indeed, for all generations but the third ($p < .001, W = 6588674$), the distribution of consecutive time spent at the front of the pair by the experienced bird cannot be distinguished from that of the naïve individual (Table S2). The tails of these distributions approach that of an exponential distribution and suggest that temporary leadership might be decided on the basis of stochastic processes (Biro et al. 2006) instead of certain rules such as fixed periods of time. Moreover, with the exception of generation 3 where 54% of the route was led

by the naïve bird (Wilcoxon signed rank-test, $p = .03, V = 1851$), there was no significant difference in the proportion of a flight spent by each bird at the front of the pair (Table S3) suggesting a relatively egalitarian relation between birds despite differing levels of experience.

(a) (b) (c)

Figure 4. Panel (a) shows the proportion of exploration over releases for the experimental group (the red dotted vertical line separates solo flights at generation 1 from paired flights at generations 2-5), the solo control, and the fixed-pairs control. Smoothed lines are computed using generalized additive models using shrinkage cubic regression splines (mean and standard error); points represent averages for individual releases. Panel (b) shows the proportion of a flight led by each bird during phases of exploration and exploitation in the experimental treatment. Darker colors correspond to exploitation, lighter colors to exploration; purple represents the experienced bird, green the naïve one; red lines represent linear fits to data pooled from both birds using the Theil–Sen estimator (slopes and p-values: respectively, 0.0203, $p = 0$ for exploitation and -0.0171, $p = 0$ for exploration). Panel (c) shows the probability for the naïve individual to initiate phases of exploration over releases (exact binomial test, $p < .01$ for release 1).

Exploration–exploitation dynamics explain flight performance. Sasaki and Biro (2017) previously showed that flight efficiency varies across treatments with experimental pairs eventually outperforming both fixed pairs of birds and solo individuals. The discovery of route innovations and, in particular, how birds with different levels of experience contribute to this task, is the key to the superior performance of experimental pairs. To understand this phenomenon and thus shed light on the pair's information pooling mechanism, we investigated how pigeons balance between exploitation of known information, closely following (< 300 meters) their most recent route, and exploration for possible route improvements. To do so, we labelled segments of flight trajectories as a function of the point-to-point distance of a focal trajectory from the immediately preceding one (*i.e.*, baseline) and compared the exploration–exploitation dynamics both across treatments and between experienced and naïve birds.

During the initial part of the experiment (Figure 4a, first 12 releases), exploration decreased steadily in all conditions with birds that flew individually in the experimental group (*i.e.*, generation 1) performing similarly to those of the solo control (respectively, 36.7% and 34.2%) whereas fixed pairs of birds explored significantly more (51.7%, Whitney–Mann–Wilcoxon, $p < .001$, Table S4). However, while exploration steadily decreased for solo and fixed pairs of birds in the successive 48 releases, experimental pairs showed a markedly different pattern of exploration oscillating over generations (Figure 4a, releases 13–60). Each time a new naïve individual was paired with an experienced one (dotted vertical lines), exploration increased for about 5 to 6 releases, reaching values well beyond those of both solo birds and fixed pairs; then exploration decreased within a few releases (2 to 4) to the same levels as those of fixed pairs. On average during generations 2 to 5, experimental pairs explored (32.9%) significantly more than both solo

(15.7%, $p < .001$) and fixed pairs of birds (29.3%, $p = .0456$). These results also held when exploration and

exploitation were defined with respect to the last release of the previous generation. Under this model,

differences between experimental and fixed pairs were even more pronounced, with the former

characterized by 46.6% exploration and the latter by only 32.4% ($p < .001$, Table S5 and Figure S6). The

inferior flight efficiency of solo and fixed pairs of birds might thus be explained, at least in some measure,

by a lower likelihood to discover route improvements due to limited exploration.

The superior homing performance of experimental pairs is suggested to be rooted in their ability to

select novel portions of a route introduced by the naïve individual that are more efficient while discarding

inefficient ones (Sasaki and Biro 2017). Under this hypothesis, we expect to observe, not only increasing

homing efficiency over generations, but also an asymmetric pattern of leadership in which the naïve

individual leads periods of exploration and the experienced one leads periods of exploitation. We found

instead no significant difference between the contributions of the experienced bird and those of the naïve

one both overall and within generations (Figure 4b and Table S7). The sole exception is represented by

generation 3 during which the naïve bird contributed more than the experienced one to exploitation

(Wilcoxon signed-rank test, $p = .035, V = 1871$). Experienced and naïve birds led the pair with

approximately the same frequency in both exploration and exploitation, suggesting that deviations from

established routes were not caused only, or even mainly, by the naïve bird (see also Figure S4, inset). We

did find evidence of behavioural asymmetries, in that transitions from exploitation to exploration were

marginally more likely to be initiated by naive birds (exact binomial test, $p = .042, n = 964$, Table S8);

however, this result was driven by those of generation 3 ($p = .02, n = 301$) whereas no difference was

detected in other generations. Transitions from exploration to exploitation were equally likely to be initiated

by the two birds both overall and within each generation. However, when transitions are considered over

the 12 releases composing each generation (Figure 4c), the naïve individual was more likely to initiate

phases of exploration during the first release ($p < .01, n = 44$, Table S9) doing so 70.5% of the time

compared to 29.5% for the experienced bird. After the first release, transitions that initiate phases of

exploration were about as likely to be initiated by either of the two birds independently of their level of

experience.

Discussion

For many group-living animals, searching for optimal travel routes can be a complex task as social

information about routes can persist over generations regardless of its quality (Helfman and Schultz 1984;

Sasaki and Biro 2017; Jesmer et al. 2018; Laland and Williams 1998). This search is inherently subject to

a trade-off between the exploitation of well-established route information accumulated over time and

exploration for innovations that constitute potential improvements (Hills et al. 2015). Striking a balance is

fundamental as a pronounced reliance on exploitation of learned information can hinder innovations

(Davies, Krebs, and West 2012) and thus promote the maintenance of potentially suboptimal behaviour

and even of maladaptive behavioural traditions (Laland and Williams 1998). Equally, an over-reliance on
exploration without exploiting the rewards of beneficial innovations eventually impedes improvements in
performance over time (Fu and Gray 2006; Mehlhorn et al. 2015).

We studied the causal structure of this process in flights of the homing pigeon *C. livia*, as this species
is capable of both social learning and information pooling (Biro et al. 2006; Pettit, Flack, et al. 2013). Of
particular interest for our study is the increase in route efficiency that results from the pairing of naïve
individuals with experienced ones (Pettit, Flack, et al. 2013), including when this happens iteratively over
multiple generations (Sasaki and Biro 2017). Previous work has proposed information pooling as the
underlying mechanism driving this increase in flight performance (Sasaki and Biro 2017). Using transfer
entropy to measure predictive information (Schreiber 2000; Pilkiewicz et al. 2020), we found quantitative
evidence that supports the information-pooling hypothesis in the strength of causal interactions within pairs
of birds. Experienced and naïve birds influence each other's behaviour; about 20% of the future directional
choices of any individual in a pair is explained by the behaviour of the other individual. These results
contrast with our expectations for unidirectional social learning that entails an asymmetric pattern of
leadership with a pronounced role for experienced individuals.

Our analysis showed that, in a multi-generational transmission-chain design, the naïve bird has a higher
influence than the experienced one during the early generations. In later generations, as flight efficiency
increases, the experienced bird becomes the better source of predictive information. We can further
hypothesize that, over generations, as birds explore an increasing portion of the search space and exhaust
remaining alternatives, a newly introduced bird should become less likely to contribute productive
innovations, and an increasing proportion of innovations should lead to errors instead of improvement. This
theoretical reasoning is analogous to the diminishing marginal value from returning to a previous location
when searching for an object in space (Stone 1976). From an information-foraging perspective (Stephens
and Krebs 1986; Pirolli 2007), the time invested in harvesting innovations on the introduction of a naïve bird
corresponds to the time invested in attending to a newly discovered patch; at some point, the opportunity
cost of further harvesting becomes too high to justify remaining in the patch. Thus, after the introduction of
a naïve bird, the information-foraging pair shifts from information-harvesting exploration back toward
information-preserving exploitation, as would be expected in an optimal search problem (Stone 1976).

How do experimental pairs improve their homing routes over generations? Previous studies where
leadership was defined on the basis of route similarity showed that, in pairs with a large difference in
experience between birds, experienced individuals were more likely to assume leadership (Flack et al.
2012). Still using route similarity, Sasaki and Biro (2017) found evidence of social learning with naïve
individuals learning routes from their experienced partners. Moreover, because newly formed pairs in the
transmission-chain design also improved performance generation after generation, they proposed that
naïve individuals could contribute innovations that pigeons evaluate in terms of route efficiency and prune
away when inefficient. However, defining leadership in terms of causal interactions instead of route
similarity allowed us to show that there is an asymmetric relation between innovators and exploiters only

during the initial flight of a newly formed pair. Although leadership is ephemeral and equally shared between birds during exploration and exploitation independently of their level of experience, over the course of this first flight the naïve individual disproportionally initiates phases of exploration, attracting the experienced bird to unfamiliar areas and triggering it to also resume the search. After that, both birds are equally likely to initiate transitions between exploration and exploitation. Moreover, as it is unlikely that experimental pairs were merely better than control groups at evaluating efficiency, we believe that their superior performance is rooted instead in their complex exploration–exploitation dynamics that allowed them to better cover the search space.

Personally acquired information allows solo individuals to improve flight performance (Meade, Biro, and Guilford 2005) but only to discover routes moderately efficient (consistently within 0.8–0.85 efficiency across a large number of experiments, reviewed in Guilford and Biro 2014) because solo individuals rapidly reduce their exploration efforts to seek out novel information. Experimental and fixed pairs of birds on the other hand explore more and thus outperform solo individuals. Together, two birds have superior sensory and cognitive capacities compared to single birds (Krause, Ruxton, and Krause 2010) which facilitates the discovery of better routes that are then learned collectively (Biro, Sasaki, and Portugal 2016; Kao et al. 2014). The reasons why pairs explore more than solo individuals might lie partly with the conflicts characteristic of newly formed pairs (Biro et al. 2006) if the resolution of conflict, *e.g.*, through averaging individual inputs, indirectly prompts pairs to explore more and discover route innovations. This hypothesis could also explain why experimental pairs outperform fixed pairs. Differently from fixed pairs that undergo a process of mutual habituation as they develop a stable route that likely reduces conflicts, the introduction of a naïve individual at the start of each generation repeatedly creates an experience imbalance in the newly formed pair. This experience imbalance could be the source of new conflicts possibly explaining why experimental pairs reach levels of exploration generally higher than those of fixed pairs. The process of gradually settling on a joint route over the course of a generation following this initial perturbation is also reminiscent of the transient effects observed when an ant with outside information joins a group of nestmates transporting an object: the new information temporarily steers the collective in the right direction but its effects on collective motion vanish quickly (Gelblum et al. 2015, 2016, 2020).

Adopting an explicit exploration–exploitation perspective to study search strategies and doing so through the use of predictive information to quantify causal interactions has the potential to advance our understanding of collective navigation in larger flocks. The conceptual framework of exploration and exploitation as well as the methods we proposed can also benefit researchers studying other taxa that move in groups with the potential to learn from previous experiences, such as shoaling fish or certain primates. In principle, this information-theoretic approach could also be applied to the study of information transfer between the environment and the individuals within a group. The ability to quantify causal interactions of this sort could shed light on broader questions in ecology involving animals moving in a group and their environment such as the impact of visual landmarks on navigation or the effects of terrestrial migration on the environment (Bracis and Mueller 2017; de Guinea et al. 2021).

Materials and Methods

Data and source code are available in (Valentini et al. 2021).

Experimental subjects and procedure. Data were taken from a previous study on cumulative route
development in the homing pigeon *Columba livia* (Sasaki and Biro 2017). Pairs of birds composed of an
experienced and a *naïve* bird were released together from the same site and allowed to fly back to the
home loft. Pairs were created and released over 5 successive generations of a transmission chain, each
generation lasting 12 consecutive releases of the same pair, according to the following procedure: initially,
at generation 1, a naïve bird was released alone 12 times, allowing it to develop its own idiosyncratic route
to the home loft. This bird, now experienced, was then paired with a new (naïve) bird at generation 2, and
together they performed another 12 flights. This process was then repeated in the next generation with a
new pair of birds composed of the former naïve bird and of a new naïve one. Data were gathered for a total
of 10 independent transmission chains, each lasting 5 generations (see Sasaki and Biro 2017 for details).
Birds flying at an average linear distance greater than 250 meters from each other were not considered as
pairs and were thus excluded from the analysis, leaving 343 flights with a mean ± SD flight duration of
8.65 ± 1.33 minutes. Additionally, in two control conditions, nine solo birds and six pairs (all initially naïve)
were released from the same site for a total of 60 releases (the equivalent of 5×12 releases for the
transmission chains).

375
Figure 1. Illustration of the methodological approach. The spatial trajectories of an experienced (E) and a naïve (N) bird
(point 1) are encoded as clockwise and counter-clockwise rotations (point 2) which we represent as discrete time series
(point 3). The combination of rotations encoded in both series (point 4) is used to estimate the probabilities required to
compute transfer entropy (point 5) and to determine the influence of one individual over the future behavior of the other
(point 6). This example illustrates transfer entropy from experienced to naïve, but we also computed it for the opposite
direction.

Data collection and pre-processing. Flight trajectories of birds were sampled at a frequency of 5 Hz
using GPS loggers, converted from the geographic coordinate system to the metric system, and projected
over the 2-dimensional plane (see Sasaki and Biro 2017). Each trajectory consisted of a time-ordered
series of positions in space, $(x_i : (x^1, x^2)_i, i \geq 1)$, see Figure 1 (point 1). We encoded the pattern of rotations
of each flight using a binary symbolic representation where symbols 0 and 1 represent, respectively, a

clockwise and a *counter-clockwise* rotation (point 2). The direction of rotation was computed as the cross
product $\overrightarrow{x_{i-1}x_i} \times \overrightarrow{x_i x_{i+1}}$ between the motion vector at time i and that at time $i+1$. The rotation is clockwise
when the product is negative and counter-clockwise when it is positive. We also measured the distance
$d_{EN}(i)$ of the experienced bird from the naïve one projected over the current direction of motion of the pair
(cf. Nagy et al. 2010 and Supplementary Material). Using this distance, we then determined the relative
position of birds over time: when $d_{EN} > 0$, the experienced bird was flying ahead of the naïve bird; when
$d_{EN} < 0$, it was flying behind. Previous tests using the same GPS loggers showed that these devices have
a sufficient level of accuracy with a small normally distributed spatial error (SD of 0.05 meters) affecting
the tracking accuracy of the direction of motion and a relatively larger error (median of 1.69 meters)
affecting that of the relative position (Pettit, Perna, et al. 2013).

Measuring information transfer. We quantified the amount of information transferred between birds
using information-theoretic measures (Cover and Thomas 2005; Pilkiewicz et al. 2020) estimated from the
series of rotations of the experienced, $E = (e_i, i \geq 1)$, and of the naïve, $N = (n_i, i \geq 1)$, birds (Figure 1,
point 3). We aimed to quantify causal interactions between birds in a Wiener–Granger sense by measuring
the extent to which the current behaviour of one bird allows us to predict the future behaviour of the other
(Bossomaier et al. 2016). Here we describe the process for predicting the naïve bird's behaviour from that
of the experienced one, but we also used the same method for the opposite direction. The average amount
of information necessary to fully predict the next rotation of the naïve bird is quantified by the marginal
entropy of its series of rotations $H(N^{i+1}) = -\sum_{n_i} p(n_i) \log_2 p(n_i)$ (Figure 1, Venn diagram, lower-left set).
This is equal to 1 bit if the flight of the naïve bird is maximally uncertain (i.e., clockwise and counter-
clockwise rotations are equally likely) and to 0 bits if the flight is fully deterministic (i.e., rotations are either
all clockwise or all counter clockwise). As a result of temporal autocorrelation (Mitchell et al. 2019), part of
this information might be contained in the recent history of rotations of the naïve bird, $n_i^{(k)} =$
$\{n_{i-k+1}, \dots, n_{i-1}, n_i\}$ for history length k (Figure 1, point 4, dark green entries in the naïve time series). The
remaining predictive information, which is not explained by the past behaviour of the naïve bird, is
quantified by the marginal entropy,

$$H(N^{i+1}|N^{(k)}) = -\sum_{n_i^{(k)}, n_{i+1}} p(n_i^{(k)}, n_{i+1}) \log_2 \frac{p(n_i^{(k)}, n_{i+1})}{p(n_i^{(k)})},$$

of its future rotations, N^{i+1}, conditioned on the outcome of the past k rotations, $N^{(k)}$ (Figure 1, Venn
diagram, overlap of the white and light purple areas).

Of interest to us was how much of this remaining information (necessary to predict the future
direction of rotation of the naïve bird) can be obtained by the current behaviour of the experienced bird
(Figure 1, points 4–6). This is given by the transfer entropy, which estimates the time-delayed effects on
the naïve bird of its interaction with the experienced one: $T_{E \to N} = H(N^{i+1}|N^{(k)}) - H(N^{i+1}|N^{(k)}, E)$
(Schreiber 2000). Transfer entropy is time directional, from the present of one bird to the future of the other,

and considered for this reason a measure of information transfer (Lizier and Prokopenko 2010). It is defined
as

$$T_{E \to N} = \sum_{n_{i+1}, n_i^{(k)}, e_i} p(n_{i+1}, n_i^{(k)}, e_i) \log_2 \frac{p(n_{i+1} \mid n_i^{(k)}, e_i)}{p(n_{i+1} \mid n_i^{(k)})}$$

and formally measures the reduction of uncertainty of the future rotation N^{i+1} of the naïve bird given by
knowledge of the current rotation E of the experienced bird at the net of possible autocorrelations in the
naïve bird's past $N^{(k)}$ (Figure 1, Venn diagram, area with red border). The logarithmic part of the above
equation is known as local transfer entropy and measures over time whether the interaction at time i
(Figure 1, point 4, dark green and dark purple entries) was informative (positive) or misinformative
(negative) (Lizier, Prokopenko, and Zomaya 2008).
The above information-theoretic measures were computed in R 3.6.1 using the rinform-1.0.2
package (Moore et al. 2018) by estimating probabilities separately for each flight. To test whether causal
interactions were significant, we also evaluated a surrogate dataset artificially created by pairing
trajectories of birds that were not flown together: the trajectory of each experienced (or naïve) bird was
paired with that of the naïve (or experienced) one from every other pair of birds not containing the same
subjects.
437
Measuring exploration and exploitation. Pairwise analysis of successive routes from the last three flights
of the experienced bird during training (generation 1) showed that, once established, the bird largely
remained within a point-to-point distance of 300 meters from its idiosyncratic route (Figure S3a). Therefore,
we used 300 meters as a threshold to differentiate flight segments between those exploring new solutions
and those exploiting known ones. We then compared the trajectories of consecutive flights for the same
subjects to label each segment in the experimental and control datasets. For each focal trajectory that we
aimed to label, we considered the trajectory of the previous release as a baseline trajectory for the
comparison. In the case of paired birds (i.e., both experimental and fixed-pairs control), we considered the
trajectory of the pair defined by the mean position of the two birds over time. For the first release of each
generation in the experimental group, we used as baseline trajectory the last release of the previous
generation because in this case there was no previous flight of the same pair to compare with. This
approach to define exploration and exploitation led to a model of exploitation (*i.e.*, the baseline trajectory)
that varied over successive releases. Because the introduction of a naïve bird at each generation was likely
to affect the baseline model of exploitation in a more pronounced manner than that of solo and fixed pairs
of birds, this model might have been susceptible to differences between the experimental design of
transmission chain experiments with respect to those of the two controls. To control for this scenario, we
also explored an alternative approach where exploitation was defined on the basis of only information
available to the experienced bird at the beginning of a new generation. In this case, the last release of the
previous generation was used as the baseline trajectory for all releases within a given generation (see

Supplementary Methods). In both cases, we also measured the distance $d_{EN}(i)$ between experienced and

naïve birds to determine which bird was flying at the front of the pair for a given route segment.

References

Berdahl, Andrew M, Colin J Torney, Christos C Ioannou, Jolyon J Faria, and Iain D Couzin. 2013.
    "Emergent Sensing of Complex Environments by Mobile Animal Groups." *Science* 339 (6119): 574–
    76. https://doi.org/10.1126/science.1225883.

Biro, Dora, Jessica Meade, and Tim Guilford. 2004. "Familiar Route Loyalty Implies Visual Pilotage in the
    Homing Pigeon." *Proceedings of the National Academy of Sciences* 101 (50): 17440–43.
    https://doi.org/10.1073/pnas.0406984101.

Biro, Dora, Takao Sasaki, and Steven J Portugal. 2016. "Bringing a Time–Depth Perspective to Collective
    Animal Behaviour." *Trends in Ecology & Evolution* 31 (7): 550–62.
    https://doi.org/10.1016/j.tree.2016.03.018.

Biro, Dora, David J T Sumpter, Jessica Meade, and Tim Guilford. 2006. "From Compromise to Leadership
    in Pigeon Homing." *Current Biology* 16 (21): 2123–28. https://doi.org/10.1016/j.cub.2006.08.087.

Bossomaier, Terry, Lionel Barnett, Michael Harré, and Joseph T Lizier. 2016. *An Introduction to Transfer
    Entropy*. Springer. https://doi.org/10.1007/978-3-319-43222-9.

Bracis, Chloe, and Thomas Mueller. 2017. "Memory, Not Just Perception, Plays an Important Role in
    Terrestrial Mammalian Migration." *Proceedings of the Royal Society B: Biological Sciences* 284
    (1855): 20170449. https://doi.org/10.1098/rspb.2017.0449.

Butail, Sachit, Violet Mwaffo, and Maurizio Porfiri. 2016. "Model-Free Information-Theoretic Approach to
    Infer Leadership in Pairs of Zebrafish." *Phys. Rev. E* 93 (4): 42411.
    https://doi.org/10.1103/PhysRevE.93.042411.

Conradt, Larissa. 2012. "Models in Animal Collective Decision-Making: Information Uncertainty and
    Conflicting Preferences." *Interface Focus* 2 (2): 226–40. https://doi.org/10.1098/rsfs.2011.0090.

Conradt, Larissa, and T J Roper. 2003. "Group Decision-Making in Animals." *Nature* 421 (6919): 155–58.
    https://doi.org/10.1038/nature01294.

Couzin, Iain D. 2009. "Collective Cognition in Animal Groups." *Trends in Cognitive Sciences* 13 (1): 36–
    43. https://doi.org/10.1016/j.tics.2008.10.002.

Couzin, Iain D, Christos C Ioannou, Güven Demirel, Thilo Gross, Colin J Torney, Andrew Hartnett,
    Larissa Conradt, Simon A Levin, and Naomi E Leonard. 2011. "Uninformed Individuals Promote
    Democratic Consensus in Animal Groups." *Science* 334 (6062): 1578 LP – 1580.
    https://doi.org/10.1126/science.1210280.

Couzin, Iain D, Jens Krause, Nigel R Franks, and Simon A Levin. 2005. "Effective Leadership and
    Decision-Making in Animal Groups on the Move." *Nature* 433 (7025): 513–16.
    https://doi.org/10.1038/nature03236.

Cover, Thomas M, and Joy A Thomas. 2005. *Elements of Information Theory. Elements of Information*
Theory. Hoboken, NJ, USA: John Wiley & Sons, Inc. https://doi.org/10.1002/047174882X.

Crosato, Emanuele, Li Jiang, Valentin Lecheval, Joseph T Lizier, X Rosalind Wang, Pierre Tichit, Guy
Theraulaz, and Mikhail Prokopenko. 2018. "Informative and Misinformative Interactions in a School
of Fish." *Swarm Intelligence*. https://doi.org/10.1007/s11721-018-0157-x.

Davies, Nicholas B, John R Krebs, and Stuart A West. 2012. *An Introduction to Behavioural Ecology*.
John Wiley & Sons.

Flack, Andrea, Benjamin Pettit, Robin Freeman, Tim Guilford, and Dora Biro. 2012. "What Are Leaders
Made of? The Role of Individual Experience in Determining Leader–Follower Relations in Homing
Pigeons." *Animal Behaviour* 83 (3): 703–9. https://doi.org/10.1016/j.anbehav.2011.12.018.

Freeman, Robin, Richard Mann, Tim Guilford, and Dora Biro. 2011. "Group Decisions and Individual
Differences: Route Fidelity Predicts Flight Leadership in Homing Pigeons (*Columba Livia*)." *Biology*
Letters 7 (1): 63–66. https://doi.org/10.1098/rsbl.2010.0627.

Fu, Wai-Tat, and Wayne D Gray. 2006. "Suboptimal Tradeoffs in Information Seeking." *Cognitive*
Psychology 52 (3): 195–242. https://doi.org/10.1016/j.cogpsych.2005.08.002.

Garland, Joshua, Andrew M Berdahl, Jie Sun, and Erik M Bollt. 2018. "Anatomy of Leadership in
Collective Behaviour." *Chaos: An Interdisciplinary Journal of Nonlinear Science* 28 (7): 75308.
https://doi.org/10.1063/1.5024395.

Gelblum, Aviram, Ehud Fonio, Yoav Rodeh, Amos Korman, and Ofer Feinerman. 2020. "Ant Collective
Cognition Allows for Efficient Navigation through Disordered Environments." Edited by Gordon J
Berman. *ELife* 9 (May): e55195. https://doi.org/10.7554/eLife.55195.

Gelblum, Aviram, Itai Pinkoviezky, Ehud Fonio, Abhijit Ghosh, Nir Gov, and Ofer Feinerman. 2015. "Ant
Groups Optimally Amplify the Effect of Transiently Informed Individuals." *Nature Communications* 6.
https://doi.org/10.1038/ncomms8729.

Gelblum, Aviram, Itai Pinkoviezky, Ehud Fonio, Nir S Gov, and Ofer Feinerman. 2016. "Emergent
Oscillations Assist Obstacle Negotiation during Ant Cooperative Transport." *Proceedings of the*
National Academy of Sciences 113 (51): 14615–20. https://doi.org/10.1073/pnas.1611509113.

Guilford, Tim, and Dora Biro. 2014. "Route Following and the Pigeon's Familiar Area Map." *Journal of*
Experimental Biology 217 (2): 169–79. https://doi.org/10.1242/jeb.092908.

Guinea, Miguel de, Alejandro Estrada, Karline R M Janmaat, K Anne-Isola Nekaris, and Sarie Van Belle.
2021. "Disentangling the Importance of Social and Ecological Information in Goal-Directed
Movements in a Wild Primate." *Animal Behaviour* 173: 41–51.
https://doi.org/10.1016/j.anbehav.2020.12.017.

Helfman, Gene S, and Eric T Schultz. 1984. "Social Transmission of Behavioural Traditions in a Coral
Reef Fish." *Animal Behaviour* 32 (2): 379–84. https://doi.org/10.1016/S0003-3472(84)80272-9.

Hills, Thomas T, Peter M Todd, David Lazer, A David Redish, and Iain D Couzin. 2015. "Exploration
versus Exploitation in Space, Mind, and Society." *Trends in Cognitive Sciences* 19 (1): 46–54.

https://doi.org/10.1016/j.tics.2014.10.004.

Jesmer, Brett R, Jerod A Merkle, Jacob R Goheen, Ellen O Aikens, Jeffrey L Beck, Alyson B Courtemanch, Mark A Hurley, et al. 2018. "Is Ungulate Migration Culturally Transmitted? Evidence of Social Learning from Translocated Animals." *Science* 361 (6406): 1023–25. https://doi.org/10.1126/science.aat0985.

Jorge, Paulo E, and Paulo A M Marques. 2012. "Decision-Making in Pigeon Flocks: A Democratic View of Leadership." *Journal of Experimental Biology* 215 (14): 2414–17. https://doi.org/10.1242/jeb.070375.

Kao, Albert B, Noam Miller, Colin Torney, Andrew Hartnett, and Iain D Couzin. 2014. "Collective Learning and Optimal Consensus Decisions in Social Animal Groups." *PLOS Computational Biology* 10 (8): 1–11. https://doi.org/10.1371/journal.pcbi.1003762.

Kim, Changsu, Tommaso Ruberto, Paul Phamduy, and Maurizio Porfiri. 2018. "Closed-Loop Control of Zebrafish Behaviour in Three Dimensions Using a Robotic Stimulus." *Scientific Reports* 8 (1): 657. https://doi.org/10.1038/s41598-017-19083-2.

Krause, Jens, and Graeme D Ruxton. 2002. *Living in Groups*. Oxford University Press.

Krause, Jens, Graeme D Ruxton, and Stefan Krause. 2010. "Swarm Intelligence in Animals and Humans." *Trends in Ecology & Evolution* 25 (1): 28–34. https://doi.org/https://doi.org/10.1016/j.tree.2009.06.016.

Laland, Kevin N, and Kerry Williams. 1998. "Social Transmission of Maladaptive Information in the Guppy." *Behavioral Ecology* 9 (5): 493–99. https://doi.org/10.1093/beheco/9.5.493.

Lizier, Joseph T, and Mikhail Prokopenko. 2010. "Differentiating Information Transfer and Causal Effect." *The European Physical Journal B* 73 (4): 605–15. https://doi.org/10.1140/epjb/e2010-00034-5.

Lizier, Joseph T, Mikhail Prokopenko, and Albert Y Zomaya. 2008. "Local Information Transfer as a Spatiotemporal Filter for Complex Systems." *Phys. Rev. E* 77 (2): 26110. https://doi.org/10.1103/PhysRevE.77.026110.

Meade, Jessica, Dora Biro, and Tim Guilford. 2005. "Homing Pigeons Develop Local Route Stereotypy." *Proceedings of the Royal Society B: Biological Sciences* 272 (1558): 17–23. https://doi.org/10.1098/rspb.2004.2873.

Mehlhorn, Katja, Ben R Newell, Peter M Todd, Michael D Lee, Kate Morgan, Victoria A Braithwaite, Daniel Hausmann, Klaus Fiedler, and Cleotilde Gonzalez. 2015. "Unpacking the Exploration–Exploitation Tradeoff: A Synthesis of Human and Animal Literatures." *Decision* 2 (3): 191–215. https://doi.org/10.1037/dec0000033.

Mitchell, David J, Antoine M Dujon, Christa Beckmann, and Peter A Biro. 2019. "Temporal Autocorrelation: A Neglected Factor in the Study of Behavioral Repeatability and Plasticity." *Behavioral Ecology*. https://doi.org/10.1093/beheco/arz180.

Moore, Douglas G, Gabriele Valentini, Sara Imari Walker, and Michael Levin. 2018. "Inform: Efficient Information-Theoretic Analysis of Collective Behaviors." *Frontiers in Robotics and AI* 5: 60. https://doi.org/10.3389/frobt.2018.00060.

Nagy, Máté, Zsuzsa Ákos, Dora Biro, and Tamás Vicsek. 2010. "Hierarchical Group Dynamics in Pigeon
Flocks." *Nature* 464 (7290): 890–93. https://doi.org/10.1038/nature08891.
Pettit, Benjamin, Zsuzsa Ákos, Tamás Vicsek, and Dora Biro. 2015. "Speed Determines Leadership and
Leadership Determines Learning during Pigeon Flocking." *Current Biology* 25 (23): 3132–37.
https://doi.org/10.1016/j.cub.2015.10.044.
Pettit, Benjamin, Andrea Flack, Robin Freeman, Tim Guilford, and Dora Biro. 2013. "Not Just Passengers:
Pigeons, *Columba Livia*, Can Learn Homing Routes While Flying with a More Experienced
Conspecific." *Proceedings of the Royal Society B: Biological Sciences* 280 (1750): 20122160.
https://doi.org/10.1098/rspb.2012.2160.
Pettit, Benjamin, Andrea Perna, Dora Biro, and David J T Sumpter. 2013. "Interaction Rules Underlying
Group Decisions in Homing Pigeons." *Journal of The Royal Society Interface* 10 (89): 20130529.
https://doi.org/10.1098/rsif.2013.0529.
Pilkiewicz, K R, B H Lemasson, M A Rowland, A Hein, J Sun, Andrew M Berdahl, M L Mayo, et al. 2020.
"Decoding Collective Communications Using Information Theory Tools." *Journal of The Royal
Society Interface* 17 (164): 20190563. https://doi.org/10.1098/rsif.2019.0563.
Pirolli, Peter. 2007. *Information Foraging Theory: Adaptive Interaction with Information*. New York, NY,
USA: Oxford University Press.
Porfiri, Maurizio. 2018. "Inferring Causal Relationships in Zebrafish-Robot Interactions through Transfer
Entropy: A Small Lure to Catch a Big Fish." *Animal Behavior and Cognition* 5 (4): 341–67.
https://doi.org/10.26451/abc.05.04.03.2018.
Ray, Subash K, Gabriele Valentini, Purva Shah, Abid Haque, Chris R Reid, Gregory F Weber, and Simon
Garnier. 2019. "Information Transfer During Food Choice in the Slime Mold Physarum
Polycephalum." *Frontiers in Ecology and Evolution* 7: 67. https://doi.org/10.3389/fevo.2019.00067.
Sasaki, Takao, and Dora Biro. 2017. "Cumulative Culture Can Emerge from Collective Intelligence in
Animal Groups." *Nature Communications* 8 (1): 15049. https://doi.org/10.1038/ncomms15049.
Sasaki, Takao, Richard P Mann, Katherine N Warren, Tristian Herbert, Tara Wilson, and Dora Biro. 2018.
"Personality and the Collective: Bold Homing Pigeons Occupy Higher Leadership Ranks in Flocks."
Philosophical Transactions of the Royal Society B: Biological Sciences 373 (1746): 20170038.
https://doi.org/10.1098/rstb.2017.0038.
Schreiber, Thomas. 2000. "Measuring Information Transfer." *Physical Review Letters* 85 (2): 461–64.
https://doi.org/10.1103/PhysRevLett.85.461.
Stephens, David W, and John R Krebs. 1986. *Foraging Theory*. Vol. 1. Princeton, New Jersey, USA:
Princeton University Press.
Stone, Lawrence D. 1976. *Theory of Optimal Search*. New York, NY, USA: Academic Press.
Valentini, Gabriele, Nobuaki Mizumoto, Stephen C Pratt, Theodore P Pavlic, and Sara Imari Walker.
2020. "Revealing the Structure of Information Flows Discriminates Similar Animal Social Behaviors."
ELife 9: e55395. https://doi.org/10.7554/eLife.55395.

Valentini, Gabriele, Theodore P Pavlic, Sara Imari Walker, Stephen C Pratt, Dora Biro, and Takao Sasaki.

2021. "Data and Code from: Naïve Individuals Promote Collective Exploration in Homing Pigeons."

Figshare. https://doi.org/10.6084/m9.figshare.14043362.

Vicsek, Tamás, and Anna Zafeiris. 2012. "Collective Motion." *Physics Reports* 517 (3): 71–140.

https://doi.org/10.1016/j.physrep.2012.03.004.

Watts, Isobel, Benjamin Pettit, Máté Nagy, Theresa Burt de Perera, and Dora Biro. 2016. "Lack of

Experience-Based Stratification in Homing Pigeon Leadership Hierarchies." *Royal Society Open*

Science 3 (1): 150518. https://doi.org/10.1098/rsos.150518.

Acknowledgements

615 GV, TPP, SIW, and SCP were supported by NSF grant No. PHY-1505048 awarded to SIW, TPP, and SCP.
616 GV was also supported by research funds from Arizona State University to Prof. Bert Hölldobler. DB was
617 supported by grant No. TWCF0316 from the Templeton World Charity Foundation's "Diverse Intelligences"
618 scheme. The authors would like to thank Dr. Albert B. Kao and Prof. Andrew Berdahl for helpful discussions.

619
620

Author contributions statement

622 GV and TS designed the study with input from all authors. TS and DB provided data. GV analysed data
623 with feedback from all authors. All authors contributed to the writing of the manuscript.
624
625

Additional information

627 **Competing interests:** The authors declare no competing interest.

Supplementary Information

Naïve individuals promote collective exploration in homing pigeons

Gabriele Valentini, Theodore P. Pavlic, Sara Imari Walker, **Stephen C. Pratt, Dora Biro &**
Takao Sasaki

Methods

Relative position of birds over time. We determined the relative position of each bird within the pair using the distance d_{EF} of the experienced bird from the naïve one projected onto the direction of motion of the flock. We modified the method proposed in (Nagy et al. 2010), which gives the distance of the experienced bird from the centre of the flock, to also include the segment from the centre of the flock to the naïve bird. This is given by

$$d_{EN}(i) = \left(\overrightarrow{x_E}(i) - \overrightarrow{x_N}(i) \right) \cdot \overrightarrow{v_{pair}}(i) \cdot 2,$$

where $\overrightarrow{x_E}(i)$ and $\overrightarrow{x_N}(i)$ are the positions of the experienced and the naïve bird and $\overrightarrow{v_{pair}}(i)$ is the normalized velocity of the pair. The normalized velocity is computed as

$$\overrightarrow{v_{pair}}(i) = \frac{\langle \overrightarrow{\dot{x}_k}(i) \rangle_k}{\left| \langle \overrightarrow{\dot{x}_k}(i) \rangle_k \right|}.$$

As the flock is composed of two birds only, the projected distance of the experienced bird from the naïve one projected onto the direction of motion of the pair is positive, when the experienced bird is flying *ahead* of the naïve one, and it is negative when the experienced bird is flying *behind*.

Exploration–exploitation with respect to the previous generation. Our primary approach to define exploration and exploitation is based on comparing pairs of successive releases using the trajectory of the previous release as our baseline model of exploitation and then labelling newer portions of a focal route as exploration. The model of exploitation (i.e., baseline trajectory) thus varies for each focal release similarly to a moving-average window over successive releases. An alternative approach to define exploration and exploitation is to consider a constant model of exploitation for each release within a given generation. This can be obtained by setting the baseline trajectory to equal the last trajectory of the previous generation. As a consequence, exploitation is defined on the basis of the information available only to the experienced bird at the beginning of a new generation; every portion of the route within that generation that is more than 300 meters away from the baseline is considered exploration.

At the first generation of the experimental group, when birds are trained individually for the successive transmission chain experiment and there is no previous generation available to provide us with

39 a baseline trajectory, we consider the first release as the baseline trajectory for the remainder of the
40 generation. In the case of solo and fixed pairs controls, for which there is no obvious definition of a
41 generation, we considered the 60 releases in each of these experiments as formed by 5 generations, each
42 lasting 12 releases, and used an equivalent approach to that of the experimental group to define exploration
43 and exploitation.
44
45 **Results**
46

(a)   **(b)**

Figure S1. Landscape of information transfer as a function of the history length, $k \in \{1, ...,17\}$, and of the sampling period, $\{0.2, 0.4, ..., 4.0\}$ seconds. Panel (a) shows the total transfer of information between the pair of birds, $T_{E \to N} + T_{N \to E}$, averaged over all releases and generations. Panel (b) shows the net transfer of information between the pair of birds, $T_{E \to N} - T_{N \to E}$, averaged over all releases and generations. Positive (respectively, negative) values represent configurations where the experienced (naïve) bird is more informative than the naïve (experienced) one. The triangle represents the configuration with maximum total transfer of information.

Landscape of information transfer and choice of parameters. We first explored how information
transfer between the birds varied as a function of two parameters: the sampling period (seconds) and the
history length used in the computation of transfer entropy. Our original GPS data are sampled at a frequency
of 1 sample every 0.2 seconds (5 Hz); we can further subsample these data by dropping samples, for
example, using one sample every 0.4, 0.6, ..., 4.0 seconds as the sampling period. The history length
represents the number of past rotations by the bird we want to predict that we consider when computing
transfer entropy from the other bird in the pair.
The total transfer of information between the birds varies as a function of these parameters (Figure
S1a). It peaks in the region delimited by history lengths of 8–10 time-steps and a sampling period between
0.2–1.2 seconds, whereas it vanishes otherwise. The total transfer of information, which is generally
adopted as a measure to choose study parameters (Porfiri 2018), reaches its maximum for a history length
of $k = 10$ at one sample every 0.2 seconds (black triangle in Figure S1a). We use this parameter
configuration for the rest of our information-theoretic analysis. For this and for most other parameter
configurations, the naïve bird is more informative about the future behaviour of the experienced one than

the other way around (Figure S1b). Only in two regions, both far from the point maximizing the total transfer

of information, is the experienced bird more informative than the naïve one; however, the overall amount

of information transferred between birds in these regions is negligible.

Comparison of information transfer with the surrogate dataset. To verify their significance, we

compared our estimates of information transfer between the two birds with equivalent measurements taken

from the surrogate dataset. We expect causal effects measured in the original dataset to be stronger than

those found in the surrogate one. As shown in Table S1, our expectations are fully met: the original dataset

shows values of transfer entropy significantly higher than those observed in the surrogate dataset, both for

the entire dataset as well as for each separate generation.

Table S1. Statistical comparison of information transfer between the original and the surrogate dataset over all
generations and over separate generations. Column 1 reports the generation and sample sizes. Columns 2 and 4 report
the differences between the mean value of transfer entropy of the original dataset and that of the surrogate dataset.
Columns 3 and 5 report the results of one-sided two-sample Whitney–Mann–Wilcoxon rank-sum tests with continuity
correction (p-value and W statistic) testing if the original dataset has significantly higher transfer entropy than the
surrogate one. Significant p-values are reported in bold.

Original vs surrogate dataset

Generation	$T_{E \to N} - T_{E \to N}^s$	$H_1: T_{E \to N} > T_{E \to N}^s$	$T_{N \to E} - T_{N \to E}^s$	$H_1: T_{N \to E} > T_{N \to E}^s$
All ($n = 343, n^s = 29035$)	$\mu = 0.0089$	$\boldsymbol{p < .001}$ ($W = 6145522$)	$\mu = 0.0088$	$\boldsymbol{p < .001}$ ($W = 6126284$)
2 ($n = 94, n^s = 7912$)	$\mu = 0.0062$	$\boldsymbol{p < .001}$ ($W = 445262$)	$\mu = 0.0074$	$\boldsymbol{p < .001}$ ($W = 452733$)
3 ($n = 99, n^s = 9801$)	$\mu = 0.0094$	$\boldsymbol{p < .001}$ ($W = 615721.5$)	$\mu = 0.0119$	$\boldsymbol{p < .001}$ ($W = 645665.5$)
4 ($n = 81, n^s = 6561$)	$\mu = 0.0071$	$\boldsymbol{p = .006}$ ($W = 308433.5$)	$\mu = 0.0057$	$\boldsymbol{p = .015}$ ($W = 303085.5$)
5 ($n = 69, n^s = 4761$)	$\mu = 0.0111$	$\boldsymbol{p < .001}$ ($W = 214258.5$)	$\mu = 0.0075$	$\boldsymbol{p = .002}$ ($W = 197618.5$)

Analysis of time spent by each bird at the front of the pair. We divided each flight into different

segments, with each segment representing a consecutive portion of the route with either the experienced

or the naïve bird at the front of the pair, and then measured the segment durations (Figure S2). The

distribution of segment durations resembles a log-normal distribution for both the entire dataset and for

individual generations. Experienced and naïve birds are characterized by very similar distributions closely

overlapping each other. Overall, the naïve bird spent significantly longer periods of time at the front of the

pair (Table S2), but the difference is small and largely driven by that of generation 3 (the only generation

showing significant differences). With the exception of generation 3, the experienced and the naïve bird

spent, on average, an approximately equal portion of the route at the front of the pair (Table S3). At

generation 3, the naïve bird was at the front of the pair for a significantly larger portion of the route with

respect to the experienced bird (respectively, 54% versus 46%).

(a)   **(b)**

Figure S2. Probability density function of the duration of flight segments with either the experienced or the naïve bird at the front of the pair. Panel (a) shows the results aggregated over all generations. Panel (b) shows the results separately for each generation.

Table S2. Statistics about the duration of flight segments with either the experienced or the naïve bird at the front of the pair. Column 1 reports the generation and sample sizes. Columns 2 and 3 give the mean duration and the standard deviation of segments for the experienced and for the naïve bird. Column 4 reports the results of two-sided two-sample Whitney–Mann–Wilcoxon rank-sum tests with continuity correction (p-value and W statistic) testing differences between the distribution of the duration D of flight segments for the two birds. Significant p-values are reported in bold.

Generation	Experienced	Naïve	$H_1: D_E \neq D_N$
All ($n^E = 10725, n^N = 10773$)	$\mu = 7.93, \sigma = 12.92$	$\mu = 8.23, \sigma = 14.12$	$\boldsymbol{p = .047}$ ($W = 56865638$)
2 ($n^E = 2512, n^N = 2492$)	$\mu = 8.89, \sigma = 17.51$	$\mu = 8.81, \sigma = 13.55$	$p = .13$ ($W = 3052262$)
3 ($n^E = 3690, n^N = 3737$)	$\mu = 6.83, \sigma = 9.08$	$\mu = 7.96, \sigma = 14.51$	$\boldsymbol{p < .001}$ ($W = 6588674$)
4 ($n^E = 2297\ n^N = 2283$)	$\mu = 8.63, \sigma = 13.65$	$\mu = 8.55, \sigma = 14.23$	$p = .86$ ($W = 2630132$)
5 ($n^E = 2226, n^N = 2261$)	$\mu = 7.98, \sigma = 11.23$	$\mu = 7.73, \sigma = 13.93$	$p = .23$ ($W = 2568854$)

Table S3. Statistics about the proportion of a flight with either the experienced or the naïve bird at the front of the pair. Column 1 reports the generation and sample size. Columns 2 and 3 give the mean and the standard deviation of the proportion of a flight with either the experienced or the naïve bird at the front of the pair. Column 4 reports the results of two-sided paired Wilcoxon signed-rank tests with continuity correction (p-value and V statistic) testing differences between the distribution of the proportion P of a flight for the two birds. Significant p-values are reported in bold.

Generation	Experienced	Naïve	$H_1: P_E \neq P_N$
All ($n = 341$)	$\mu = 0.49, \sigma = 0.2$	$\mu = 0.51, \sigma = 0.2$	$p = .46$ ($V = 27817.5$)
2 ($n = 92$)	$\mu = 0.51, \sigma = 0.23$	$\mu = 0.49, \sigma = 0.23$	$p = .69$ ($V = 2243$)
3 ($n = 99$)	$\mu = 0.46, \sigma = 0.17$	$\mu = 0.54, \sigma = 0.17$	$\boldsymbol{p = .03}$ ($V = 1851$)
4 ($n = 81$)	$\mu = 0.5, \sigma = 0.2$	$\mu = 0.5, \sigma = 0.2$	$p = .84$ ($V = 1705$)
5 ($n = 69$)	$\mu = 0.49, \sigma = 0.2$	$\mu = 0.51, \sigma = 0.2$	$p = .95$ ($V = 1219$)

Exploration versus exploitation with respect to successive releases. We used a distance-based mechanism to label portions of a focal route as either exploration or exploitation depending on their point-to-point distance from the baseline trajectory at the previous release. To determine a suitable threshold, we compared successive trajectories flown by the experienced bird towards the end of training (generation 1, last three flights) and studied the distribution of distances between successive trajectories (Figure S3a). The distribution of distances is right-skewed and approximately exponential. A threshold of 300 meters captures a large portion of the probability mass, about 70.5%, whereas larger distances are progressively less likely. On this basis, we set a threshold of 300 meters to distinguish between phases of exploitation (< 300 meters) and phases of exploration (≥ 300 meters). Figure S3b shows the distribution of distances between consecutive flights observed during each generation whereas Figure S4a shows the same results aggregated over all generations as well as the proportion of time (inset) that each bird is leading as a function of the same distance. Figures S4b and S4c show similar results for fixed pairs of birds and solo individuals. Whereas fixed pairs are characterized by a distribution similar to that of experimental pairs, solo birds have a markedly shifted distribution towards exploitation at the expense of exploration.

(a)   (b)

Figure S3. Distribution of minimum distances between the pairs of consecutive flights. Panel (a) shows the results for the trained birds during the first generation of the experiment. Panel (b) shows the results for the pair of birds during all remaining generations of the experiments. Vertical lines highlight the division of the probability mass between exploitation (left) and exploration (right) defined by a 300 meters threshold.

(a)   (b)   (c)

Figure S4. Illustration of the distribution of point-to-point distances between pairs of consecutive flights highlighting the 300 meter threshold that demarks the end of exploitation and the beginning on exploration. Panel (a) reports the results for experimental pairs over generations 2–5, panel (b) reports those for the fixed pairs control, and panel (c) those for

We compared the proportion of exploration and exploitation across experimental conditions before the

beginning (*i.e.*, generation 1) and during transmission chain experiments (*i.e.*, generation 2 to 5). During

the first 12 releases (Table S4), individuals from the experimental group that flew solo during generation 1

could not be distinguished from birds in the solo control whereas fixed pairs of birds showed levels of

exploration significantly higher than those of solo birds. During releases 13 to 60, that is, when naïve

individuals are iteratively introduced in the transmission chains at the beginning of each generation,

experimental pairs showed instead significantly higher levels of exploration than both solo and fixed pairs

of birds, with these former exploring much less than birds that flew in pairs.

140
Table S4. Statistical comparison of mean proportions of a flight spent exploring versus exploiting across treatments
(experimental pairs, solo and fixed pairs controls) for the first 12 releases (generation 1) and for releases 13 to 60
(generation 2 to 5). Entries report the proportion of exploration vs exploitation for pairs of treatments as well as the
results of two-sided two-sample Whitney–Mann–Wilcoxon rank-sum tests with continuity correction (p-value and W
statistic) for differences in proportion of exploration. Significant p-values are reported in bold. Results of testing for the
proportion of exploitation are equivalent and not repeated below.

Releases	*Dataset*	Solo control	Fixed pairs control
1 to 12	Experimental (gen. 1)	Row: 36.7% vs 63.3% Col: 34.2% vs 65.8% $p = .55$ ($W = 5288$)	Row: 36.7% vs 63.3% Col: 51.7% vs 48.3% $\boldsymbol{p <.001}$ ($W = 2230$)
	Solo control	–	Row: 36.7% vs 63.3% Col: 34.2% vs 65.8% $\boldsymbol{p <.001}$ ($W = 1837$)
13 to 60	Experimental (gen. 2-5)	Row: 32.9% vs 67.1% Col: 15.7% vs 84.3% $\boldsymbol{p <.001}$ ($W = 94108.5$)	Row: 32.9% vs 67.1% Col: 29.3% vs 70.7% $\boldsymbol{p =.0456}$ ($W = 50472$)
	Solo control	–	Row: 15.7% vs 84.3% Col: 29.3% vs 70.7% $\boldsymbol{p <.001}$ ($W = 31517$)

Exploration versus exploitation with respect to the previous generation. In addition to the exploration–

exploitation analysis with respect to successive releases, we also performed a similar analysis where

exploitation is defined with respect to the previous generation (see supplementary Methods above) in order

to validate the robustness of the main results obtained with our primary approach. The distributions of point-

to-point distances between baseline and focal trajectories (Figure S5) resembled those observed when

comparing successive releases (Figure S4). Differences across experimental and control treatments (Table

S5) were much more pronounced under this model but in line with the results obtained above (Table S4).

Moreover, the overall exploration trends reported in Figure S6 showed the same signatures of the analysis

over successive releases (Figure 5a). Exploration decreased over generations in all experimental

conditions with experimental pairs exploring more than both fixed pairs of birds and solo individuals.

Figure S5. Illustration of the distribution of point-to-point distances between each flight of a generation (focal trajectories) and the last flight of the previous generation (baseline trajectory). Colors and vertical lines highlight the 300 meter threshold that demarks the end of exploitation and the beginning of exploration. Panel (a) reports the results for experimental pairs over generations 2–5, panel (b) reports those for the fixed pairs control, and panel (c) those for the solo control.

Table S5. Statistical comparison of mean proportions of a flight spent exploring versus exploiting across treatments (experimental pairs, solo and fixed pairs controls) for the first 12 releases (generation 1) and for releases 13 to 60 (generation 2 to 5) when considering the last release at the previous generation as the baseline trajectory. Entries report the proportion of exploration vs exploitation for pairs of treatments as well as the results of two-sided two-sample Whitney–Mann–Wilcoxon rank-sum tests with continuity correction (p-value and W statistic) for differences in proportion of exploration. Significant p-values are reported in bold. Results of testing for the proportion of exploitation are equivalent and not repeated below.

Releases	*Dataset*	Solo control	Fixed pairs control
1 to 12	Experimental (gen. 1)	Row: 61.5% vs 38.5% Col: 55.8% vs 44.2% $p = .11$ ($W = 5592$)	Row: 61.5% vs 38.5% Col: 82.9% vs 17.1% $\boldsymbol{p <.001}$ ($W = 1721$)
	Solo control	–	Row: 55.8% vs 44.2% Col: 82.9% vs 17.1% $\boldsymbol{p <.001}$ ($W = 1163$)
13 to 60	Experimental (gen. 2-5)	Row: 46.6% vs 53.4% Col: 18.9% vs 81.1% $\boldsymbol{p <.001}$ ($W = 103095$)	Row: 46.6% vs 53.4% Col: 32.4% vs 67.6% $\boldsymbol{p <.001}$ ($W = 60502$)
	Solo control	–	Row: 18.9% vs 81.1% Col: 32.4% vs 67.6% $\boldsymbol{p <.001}$ ($W = 30744$)

Figure S6. Illustration of the proportion of exploration (respectively, one minus the proportion of exploitation) over releases when considering the last release at the previous generation as the baseline trajectory. Results are shown for

the experimental group, the solo control, and the fixed pairs control. The red dotted vertical line delineates the end of
the experimental group's training phase. Smoothed lines are computed with generalized additive models using
shrinkage cubic regression splines (mean and standard error); points represent averages for individual releases.

The role of leadership during exploration and exploitation. Using this distance-based mechanism, we
label each segment of a flight as either exploration or exploitation. For each segment, we keep track of
which bird, experienced or naïve, is leading the pair. We then compute the proportion of time spent by the
experienced bird and by the naïve one leading the pair during either an exploration or exploitation phase
(Table S6). The combined proportion of time of both birds in either exploration or exploitation gives instead
the corresponding contribution of the pair. Overall, each bird spends approximately 16.5% of the time
leading the pair during an exploration phase and 33.5% of the time leading the pair during an exploitation
phase. With the exception of exploitation phases in generation 3, there is no significant difference between
the proportion of time spent by the two bird in an exploration or exploitation phase (Table S7).

Table S6. Proportion of a flight led by each of the two birds, calculated separately for exploration and exploitation
phases. Column 1 reports the generation and sample size. Columns 2 and 3 give the mean and the standard deviation
of the proportion of a flight led, respectively, by the experienced and the naïve bird for the case of exploration. Columns
4 and 5 give the mean and the standard deviation of the proportion of a flight led, respectively, by the experienced and
the naïve bird for the case of exploitation.

	Exploration		Exploitation	
Generation	Experienced	Naïve	Experienced	Naïve
All $(n = 341)$	$\mu = 0.17, \sigma = 0.15$	$\mu = 0.16, \sigma = 0.13$	$\mu = 0.32, \sigma = 0.18$	$\mu = 0.35, \sigma = 0.2$
2 $(n = 92)$	$\mu = 0.2, \sigma = 0.17$	$\mu = 0.17, \sigma = 0.14$	$\mu = 0.31, \sigma = 0.2$	$\mu = 0.32, \sigma = 0.21$
3 $(n = 99)$	$\mu = 0.16, \sigma = 0.13$	$\mu = 0.19, \sigma = 0.13$	$\mu = 0.3, \sigma = 0.16$	$\mu = 0.36, \sigma = 0.17$
4 $(n = 81)$	$\mu = 0.18, \sigma = 0.16$	$\mu = 0.16, \sigma = 0.13$	$\mu = 0.32, \sigma = 0.18$	$\mu = 0.34, \sigma = 0.22$
5 $(n = 69)$	$\mu = 0.13, \sigma = 0.13$	$\mu = 0.1, \sigma = 0.1$	$\mu = 0.37, \sigma = 0.17$	$\mu = 0.4, \sigma = 0.21$

198 *Table S7. Statistical comparison of leadership by experienced vs. naïve birds, tested separately for exploration and*
199 *exploitation over all generations and over separate generations. Column 1 reports the generation and sample size.*
200 *Columns 2 and 3 report the results of two-sided paired Wilcoxon signed-rank tests with continuity correction (p-value*
201 *and V statistic) for differences in proportion of flight led between experienced and naïve birds for exploration and*
202 *exploitation, respectively. Significant p-values are reported in bold.*

	Experienced vs Naïve	
Generation	Exploration	Exploitation
All $(n = 341)$	$p = .44 \ (V = 28820)$	$p = .13 \ (V = 26240)$
2 $(n = 92)$	$p = .17 \ (V = 2490)$	$p = .71 \ (V = 2048)$
3 $(n = 99)$	$p = .08 \ (V = 1891)$	$\boldsymbol{p = .035} \ (V = 1871)$
4 $(n = 81)$	$p = .36 \ (V = 1767)$	$p = .79 \ (V = 1603)$
5 $(n = 69)$	$p = .22 \ (V = 1187)$	$p = .8 \ (V = 1131)$

Transitions between phases of exploration and exploitation. Although once initiated, phases of exploration and phases of exploitation are led in equal manner by the experienced and the naïve bird, the experience unbalance within the pair might affect the likelihood of a bird to initiate transitions from one phase of the exploration–exploitation process to other. To investigate this question, we measured transition probabilities for both birds over generations (Table S8) and over releases (Table S9). Transitions from exploration to exploitation are not significantly different between experienced and naïve birds, but transitions from exploitation to exploration are significantly (albeit marginally) more likely to be initiated by naïve birds. However, this result seems driven by the data of generation 3, where the naïve individual is marginally but significantly more likely than the experienced one to initiate exploration phases, while in all other generations both birds are equally likely to initiate changes in either direction. When looking at transition probabilities over releases, we found a similar trend except for the first release where the naïve bird is much more likely to initiate exploration phases.

Table S8. Statistical comparison of the proportion of transitions from exploitation to exploration and from exploration to exploitation led by the experienced and by the naïve bird over generations. Column 1 reports the generation number. Columns 2 and 4 report the estimated probabilities that transitions are led by the experienced bird, P_E, and by the naïve bird, P_N, for transitions, respectively, from exploitation to exploration and from exploration to exploitation. Columns 3 and 5 give the results of exact binomial tests (p-value and sample size n) of the null hypothesis that the probability P_E that transitions are led by the experienced bird equals 0.5. Significant p-values are reported in bold.

	Exploitation → Exploration		Exploration → Exploitation	
Generation	P_E versus P_N	$H_1: P_E \neq 0.5$	P_E versus P_N	$H_1: P_E \neq 0.5$
All	$P_E = 0.467, P_N = 0.533$	$\boldsymbol{p =.042}$ $(n = 964)$	$P_E = 0.513, P_N = 0.487$	$p = .42$ $(n = 966)$
2	$P_E = 0.494, P_N = 0.506$	$p = .9$ $(n = 247)$	$P_E = 0.52, P_N = 0.48$	$p = .56$ $(n = 244)$
3	$P_E = 0.432, P_N = 0.568$	$\boldsymbol{p =.02}$ $(n = 301)$	$P_E = 0.483, P_N = 0.517$	$p = .6$ $(n = 300)$
4	$P_E = 0.5, \quad P_N = 0.5$	$p = 1$ $(n = 216)$	$P_E = 0.539, P_N = 0.461$	$p = .28$ $(n = 219)$
5	$P_E = 0.45, P_N = 0.55$	$p = .18$ $(n = 200)$	$P_E = 0.522, P_N = 0.478$	$p = .57$ $(n = 203)$

Table S9. Statistical comparison of the proportion of transitions from exploitation to exploration and from exploration to exploitation led by the experienced and by the naïve bird over releases. Column 1 reports the release number. Columns 2 and 4 report the estimated probabilities that transitions are led by the experienced bird, P_E, and by the naïve bird, P_N, for transitions, respectively, from exploitation to exploration and from exploration to exploitation. Columns 3 and 5 give the results of exact binomial tests (p-value and sample size n) of the null hypothesis that the probability P_E that transitions are led by the experienced bird equals 0.5. Significant p-values are reported in bold.

	Exploitation → Exploration		Exploration → Exploitation	
Release	P_E versus P_N	$H_1: P_E \neq 0.5$	P_E versus P_N	$H_1: P_E \neq 0.5$
1	$P_E = 0.295, P_N = 0.705$	$\boldsymbol{p <.01}$ $(n = 44)$	$P_E = 0.477, P_N = 0.523$	$p = .88$ $(n = 44)$
2	$P_E = 0.478, P_N = 0.522$	$p = .81$ $(n = 69)$	$P_E = 0.522, P_N = 0.478$	$p = .81$ $(n = 67)$
3	$P_E = 0.519, P_N = 0.481$	$p = .83$ $(n = 81)$	$P_E = 0.575, P_N = 0.425$	$p = .22$ $(n = 80)$
4	$P_E = 0.576, P_N = 0.424$	$p = .27$ $(n = 66)$	$P_E = 0.6, P_N = 0.4$	$p = .14$ $(n = 65)$
5	$P_E = 0.487, P_N = 0.513$	$p = .91$ $(n = 76)$	$P_E = 0.519, P_N = 0.481$	$p = .82$ $(n = 77)$
6	$P_E = 0.54, P_N = 0.46$	$p = .48$ $(n = 100)$	$P_E = 0.465, P_N = 0.535$	$p = .55$ $(n = 99)$

7	$P_E = 0.443, P_N = 0.557$	$p = .34$ $(n = 88)$	$P_E = 0.494, P_N = 0.506$	$p = 1.0$ $(n = 89)$
8	$P_E = 0.409, P_N = 0.591$	$p = .18$ $(n = 66)$	$P_E = 0.597, P_N = 0.403$	$p = .14$ $(n = 67)$
9	$P_E = 0.438, P_N = 0.562$	$p = .22$ $(n = 112)$	$P_E = 0.496, P_N = 0.504$	$p = 1$ $(n = 115)$
10	$P_E = 0.422, P_N = 0.578$	$p = .17$ $(n = 90)$	$P_E = 0.522, P_N = 0.478$	$p = .75$ $(n = 92)$
11	$P_E = 0.391, P_N = 0.609$	$p = .053$ $(n = 87)$	$P_E = 0.453, P_N = 0.547$	$p = .45$ $(n = 86)$
12	$P_E = 0.541, P_N = 0.459$	$p = .52$ $(n = 85)$	$P_E = 0.482, P_N = 0.518$	$p = .83$ $(n = 85)$

References

Nagy, Máté, Zsuzsa Ákos, Dora Biro, and Tamás Vicsek. 2010. "Hierarchical Group Dynamics in Pigeon Flocks." *Nature* 464 (7290): 890–93. https://doi.org/10.1038/nature08891.

Porfiri, Maurizio. 2018. "Inferring Causal Relationships in Zebrafish-Robot Interactions through Transfer Entropy: A Small Lure to Catch a Big Fish." *Animal Behavior and Cognition* 5 (4): 341–67. https://doi.org/10.26451/abc.05.04.03.2018.