Title

Mass-sensitive particle tracking (MSPT) to elucidate the membrane-associated MinDE reac-

tion cycle

Authors

Tamara Heermann[1,#], Frederik Steiert[1,2,#], Beatrice Ramm[1,3], Nikolas Hundt[4,#,*] and Petra

Schwille[1,*]

# equal contribution

Affiliations

[1]Department of Cellular and Molecular Biophysics, Max Planck Institute of Biochemistry, Am

Klopferspitz 18, 82152 Planegg, Germany

[2]Physics Department, Technical University Munich, Garching, Germany.

[3]current affiliation: Department of Physics, Princeton University, 08544 Princeton, NJ, USA

[4]Department of Cellular Physiology, Biomedical Center (BMC), Ludwig-Maximilians-Universi-

tät München, Großhaderner Str. 9, 82152 Planegg, Germany

* corresponding authors:

Prof. Petra Schwille

ORCID: 0000-0002-6106-4847

schwille@biochem.mpg.de

Phone: +49 89 8578-2900

Fax: +49 89 8578-2903

Dr. Nikolas Hundt

ORCID: 0000-0001-8217-671X

nikolas.hundt@med.uni-muenchen.de

Phone: +49 (0) 89 2180 71563

Fax: +49 (0) 89 2180 71732

Keywords

Self-organisation; single-particle tracking; Min system; iSCAT; mass photometry

Abbreviations

adenosine-5'-triphosphate (ATP), adenosine-5'-diphosphate (ADP), bovine serum albumin

(BSA), 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine-N-cap biotinyl (18:1 Biotinyl Cap PE),

dioleoyl-sn-glycero-3-phosphocholine (DOPC), dioleoyl-sn-glycero-3-phosphoglycerol

(DOPG), horseradish peroxidase (HRP), interferometric scattering (iSCAT), phosphate binding

loop (P-loop), point spread function (PSF), single-particle tracking (SPT), small unilamellar

vesicle (SUV), supported lipid bilayer (SLB), mass photometry (MP), mass-sensitive particle

tracking (MSPT), membrane targeting sequence (MTS)

Abstract

In spite of their great importance in biology, methods providing access to spontaneous molec-

ular interactions with and on biological membranes have been sparse. So far, it has been

consensus that their observation with sufficient sensitivity and time resolution requires the in-

troduction of - predominantly fluorescent- labels to the system. However, the recent advent of

mass photometry to quantify mass distributions of unlabelled biomolecules landing on surfaces

raised hopes that this approach could be transferred to membranes. Here, we introduce mass-

sensitive particle tracking (MSPT), enabling simultaneous label-free tracking and monitoring

of molecular masses of single biomolecules diffusing on lipid membranes. We applied this

approach to the highly non-linear reaction cycles underlying MinDE protein self-organisation.

MSPT allowed us to determine the stoichiometry and turnover of individual membrane-bound

MinD/MinDE protein complexes and to quantify their size-dependent diffusion. We found that

MinD assembles into complexes larger than the commonly postulated dimer, through lateral

interactions of membrane-bound complexes and subunit recruitment from solution. Further-

more, the ATPase-activating protein MinE interconnects MinD into high-molecular-weight het-

eromeric complexes and affects their subunit turnover and concerted membrane release. This

study demonstrates the potential of MSPT to enhance our quantitative understanding of both

prokaryotic and eukaryotic membrane-associated biological systems.

Introduction

The recruitment of proteins to lipid interfaces is crucial for various cell biological processes,

such as the regulation of membrane trafficking[1], mediation of signalling cascades[2], and the

establishment of cell polarity. These membrane-associated reactions often rely on short-lived

complexes that coexist in a dynamic equilibrium with their respective cytosolic forms[3]. Transi-

ent interactions on the membrane eventually serve as nucleation sites for the assembly of

larger, more stable complexes. Additionally, the spatial distribution, stoichiometry and temporal

dynamics of membrane-associated complexes are often heterogeneous[4]. This combination of

fast dynamics and compositional heterogeneity makes membrane-associated reactions diffi-

cult targets for conventional analytical techniques, which determine the composition and follow

the dynamics of molecular systems based on ensemble-averaged measures[5]. In recent dec-

ades, single-particle tracking (SPT) has revolutionised the analysis of membrane-associated

systems by following the dynamics of individual molecules at nanometre precision and milli-
second time resolution[6–8], using fluorophores as labels in combination with highly sensitive
microscopy. More recently, scattering-based detection using gold nanoparticles as labels has
pushed the spatiotemporal resolution down to the sub-nanometre and microsecond range[9–12].
The single-molecule nature of these approaches has provided detailed mechanistic insight into
the dynamics of biomolecular systems[13]. However, as label-based technique, SPT also suffers
from label-induced artifacts: large particles, but also small fluorescent tags[14] may perturb na-
tive protein function, and fluorescence-specific phenomena like photo-bleaching and -blinking
hinder continuous particle tracking and limit the attainable spatiotemporal resolution[15]. More
importantly, although fluorescence-based SPT may provide access to interaction dynamics
between tagged molecules through brightness changes, it has proven extremely complicated
to extract the molecular composition of the tracked particles. In general, relating the signal of
an external marker to the molecular stoichiometry of the labelled particle requires careful char-
acterisation of labelling efficiencies and imaging conditions, which is very often hampered by
quenching effects. Especially for multicomponent systems, access to the molecular composi-
tion of single particles has been prohibitive for standard biological applications.

Recent advances in interferometric scattering (iSCAT) microscopy[11,16] made it possible to de-
tect individual biomolecules label-free based on light scattering. The linear relationship be-
tween iSCAT contrast and molecular mass of a biomolecule led to the development of mass
photometry (MP)[17,18]. This technique allows the determination of size distributions of biomole-
cules in solution by measuring the individual masses of molecules landing on a glass surface
and counting their relative abundances, to determine both molecular identity and composition.
For purely shot noise-limited detection of a single molecule landing on a glass surface on top
of the static scattering pattern, MP applies a sophisticated background removal strategy opti-
mised for this type of experiment, which is not compatible with the detection of mobile mole-
cules diffusing on lipid membranes. Here, by introducing a new iSCAT image processing and
analysis strategy, we enable mass-sensitive particle tracking (MSPT) of single unlabelled bio-
molecules on a supported lipid bilayer (SLB). We show that the iSCAT signal-to-mass linearity
holds true for membrane-associated proteins and that we can relate molecular composition,
accessible via mass, to diffusive behaviour. Moreover, we can follow the mass time course
along individual trajectories, making it possible to observe the (dis-)assembly of biomolecular
complexes in real-time. The approach is fast and provides high particle statistics within
minutes, all without need for protein labelling and its caveats.

We showcase the abilities of this method for the detailed analysis of complex biological sys-
tems by analysing the membrane-associated reaction cycle of the *Escherichia coli* Min system.

This system consists of three proteins - MinC, MinD and MinE - and is essential for the spati-
otemporal regulation of the division site in *E. coli*[19]. To perform this task, the ATPase MinD and
the ATPase-activating protein MinE oscillate between the cell poles, forming a concentration
gradient of the passenger protein MinC. This gradient supposedly enables MinC to inhibit FtsZ
polymerisation at the poles and directs Z-ring formation to the mid-cell[20,21]. To this end, the
phospholipid bilayer acts as a catalytic interface and membrane interaction of MinD and MinE
is mediated by amphipathic helices, i.e. membrane targeting sequences (MTS)[22–25]. The inter-
action of MinD and MinE with the membrane decreases their diffusion rates and facilitates
lateral molecular interactions enabling the system to self-organise[26]. Despite its compositional
simplicity the system exhibits surprisingly complex, non-linear dynamics. Its self-organisation
can be reconstituted *in vitro*[26] and the underlying mechanism has been probed by various
techniques such as SPT[13], high-speed atomic force microscopy[27], nuclear-magnetic reso-
nance[28], electron microscopy[29], plasmonic nano sensors[30] and mutational analysis[31,32]. De-
spite this intense characterisation, molecular details about MinDE self-assembly, such as the
presumed cooperativity in bilayer-attachment of MinD, have remained poorly understood.
Thus, the system was set to benefit from the unique ability of MSPT to characterise dynamics
as well as molecular composition of membrane-attached protein complexes. Using MSPT, we
dissected the membrane-associated MinDE reaction cycle by determining the stoichiometry,
turnover, and diffusion of individual membrane-bound MinD/MinDE protein complexes. Our
results indicate that MinD, in contrast to the classical model, assembles not only into dimers,
but also forms larger oligomers, confirming recent findings[27]. We furthermore show that MinE
promotes MinD self-assembly on the lipid bilayer due to its ability to interconnect MinD into so
far unresolved higher-order heteromeric complexes. We believe our experiments on the Min
system demonstrate that MSPT is a powerful, widely applicable tool for the mechanistic anal-
ysis of both pro- and eukaryotic membrane-associated systems.
137
138   **Results**
139
Mass-sensitive dynamic imaging of membrane-associated protein complexes
The major obstacle for detecting single macromolecules with iSCAT microscopy (Fig. 1a) is
separating their comparatively small signal from the dominant scattering background. In a
standard mass photometry experiment, molecules landing on a glass surface are detected
through continuous comparison of an averaged image with the average of its preceding frames
as background (Supplementary Fig. 1a)[17,18]. In this image processing approach, a molecule
appears as a dark spot at the moment of landing and disappears when it becomes part of the
static background[17]. The mass of the particle can then be determined by fitting the detected
peak signal with a model point spread function (PSF). However, for a moving molecule, this

strategy produces distorted images of the molecule's PSF. The missing particle density at its

previous location produces a bright spot, while the added density at the new position generates

a dark spot. The spatial overlap of these patterns causes moving objects to appear as dark

fronts carrying bright tails, thus hampering the determination of their mass and location (Sup-

plementary Fig. 1b, Supplementary Movie 1). To address this issue, we employed the temporal

median of an image sequence as background estimate[10,33,34]. If during the median period mov-

ing molecules only occasionally cross a surface location, the median signal at that location will

be a good estimate of the empty background. To retain shot-noise limited detection, which is

strongly affected by sample drift, we calculated the pixel-wise median of an image sequence

as background estimate for its central frame and moved this median window from frame to

frame throughout the video (Supplementary Fig. 1c). Due to the clear separation of static back-

ground from moving objects, background corrected movies showed clear, undistorted images

of moving PSFs (Fig. 1a, Supplementary Fig. 1c, Supplementary Table 1 and Supplementary

Movie 1), a prerequisite that allows mass-sensitive particle tracking (MSPT) of single unla-

belled biomolecules diffusing on lipid interfaces.

First, we set out to assess the quality of mass determination of mobile molecules compared to

landing particles in conventional mass photometry. To this end, we used a bilayer supple-

mented with biotinylated lipids and attached a set of biotinylated standard proteins with known

mass via divalent streptavidin[35]. This system has several advantages, such as the ability to

cover different protein size regimes, standardised membrane binding, and the added benefit

of simplified complex stoichiometries by using divalent streptavidin. For each molecule, we

determined its median contrast throughout its trajectory. Analogous to conventional mass pho-

tometry (Fig. 1b, left column), histograms of the contrasts of all molecules determined with

MSPT revealed the particle size distribution (Fig. 1b, right column, Supplementary Movie 2).

For our standard proteins, iSCAT contrast as a function of mass exhibited the expected linear

relationship[17]. The calibration line obtained from standards diffusing on SLBs was in fact indis-

tinguishable from the one determined with molecules landing on glass (Fig. 1c). This result

suggests that mass calibrations performed with landing assays can be transferred to particles

diffusing on membranes. However, this strictly needs to be verified for any new lipid/buffer

combination, protein system and imaging condition.

Besides mass determination, MSPT also enables the analysis of the diffusive behaviour of

membrane-bound molecules. For this purpose, it is important to choose the median window

size for background estimation such that particles travel sufficient distances during the median

period. Hence, we first systematically tested the minimum median window sizes required to

extract the correct diffusion coefficients at particle diffusion speeds expected for our SLB sys-
tem. We generated artificial movies of randomly diffusing particles at varying speeds and com-
pared the input diffusion coefficients with diffusion coefficients extracted using different median
window sizes and extraction methods (see methods, Supplementary Figures 2 and 3 as well
as Supplementary Discussion for details). Based on the results of our simulation, we chose a
jump-distance analysis[36] to extract diffusion coefficients from our experimental videos. As an
experimental verification of the diffusion coefficients obtained in this manner, we again made
use of streptavidin attached to a membrane via biotinylated lipids. In line with literature values
ranging from 0.8 to 2.0 μm^2 s^{-1} [37–39], the lateral diffusion coefficient of streptavidin was found
to be 1.3 ± 0.1 μm^2 s^{-1} (Fig. 1d). Similar diffusion coefficients were obtained for all standard
proteins attached via divalent streptavidin (Supplementary Fig. 4). To highlight a unique ad-
vantage of MSPT, we plotted the diffusion coefficient versus the respective molecular mass
obtained from individual trajectories, enabling an unprecedented direct connection of these
two parameters. As displayed in Figure 1d, the unimodal distribution of membrane-bound
streptavidin indicates a distinct population of tetramers undergoing Brownian motion. Having
validated the method with membrane-attached streptavidin, we thought the method offers the
potential of detailed insight into the dynamics of molecular interactions within more complex
membrane-bound systems. To further explore the method's capabilities, we turned to the
membrane-associated *E. coli* Min system.

Figure 1 - Principle of mass-sensitive particle tracking (MSPT). (a) Schematic displaying the inter-
ferometric scattering (iSCAT)-based measurement principle of MSPT. In this setup, proteins freely dif-
fuse on a supported lipid bilayer and can be tracked and identified according to the linear relationship
between their interferometric scattering contrast and molecular weight. Exemplary structures of three
aldolase oligomer states (PDB: 4S1F[40]) are shown in the top panel, and their respective iSCAT images
at the bottom. Scale bar: 1 µm. **(b)** Probability density distributions of standard proteins determined
using the conventional MP landing assay (left panel) or using MSPT (right panel). All data represent

pooled distributions of three independent experiments per condition: alcohol dehydrogenase (ADH; par-

ticle number n = 9828), bovine serum albumin (BSA; n = 11,408), TEV protease (TEV; n = 1,705), β-

amylase (bAm; n = 10,043), protein A (prA; n = 12,720), divalent streptavidin (Strep; n = 16,699 trajec-

tories), divalent streptavidin with biotinylated aldolase (Strep-ALD; n = 16,727 trajectories), divalent

streptavidin with biotinylated bovine serum albumin (Strep-BSA; n = 8,842 trajectories) and divalent

streptavidin with biotinylated protein A (Strep-prA; n = 22,424 trajectories). Dashed lines mark peaks

not considered for mass calibration (left panel). Continuous lines represent oligomer states included in

the mass calibration. 2D plots of mass vs. diffusion coefficient for the four proteins measured with MSPT

(right panel) are shown in Supplementary Fig. 4. **(c)** Comparison of the contrast-to-mass calibration for

MP and MSPT, derived from peak contrasts in panel (b) and their assigned sequence masses (see

Supplementary Table 2, 3). Error bars represent the standard error of the peak locations estimated by

bootstrapping. **(d)** 2D kernel density estimation of 1.25 nM tetravalent streptavidin bound to biotinylated

lipids on a supported lipid bilayer (n = 73,901 trajectories of three independent replicates; particle den-

sity: 0.2 μm^{-2}). Marginal probability distributions of the molecular mass and the diffusion coefficient are

presented at the top and right, respectively.

Cooperative Membrane-Catalysed Association Dynamics of MinD

The MinDE system is known for its ability to self-organise into mesoscopic protein patterns on

lipid membranes[26]. To generate these patterns, MinD and MinE are generally assumed to un-

dergo a canonical membrane binding-unbinding cycle displayed in Figure 2a. In brief, upon

ATP-complexation cytosolic MinD dimerises and localises to the membrane interface[23]. After

homodimeric MinE binds to MinD, nucleotide hydrolysis is stimulated and MinD dissociates

from the lipid bilayer to return to its monomeric state[41]. Despite this established model, it has

remained rather enigmatic how ATP-dependent dimerisation and the resulting increase in

MinD membrane affinity alone can confer the non-linear attachment required for pattern for-

mation. In recent years, it has been proposed that the presence of higher MTS valences and

thus higher-order oligomer structures might contribute to the local self-enhancement of MinD

at the membrane[27,31,42]. Due to the fast diffusion and attachment/detachment dynamics, how-

ever, it has remained challenging to provide convincing evidence of their existence.

Taking advantage of the ability to monitor the mass of membrane-attached and diffusing pro-

tein complexes, we thus applied our MSPT approach to investigate potential oligomeric spe-

cies formed by the MinDE system. We started by reconstituting the membrane recruitment of

MinD and compared the mass distribution of MinD in solution, measured in an MP landing

assay, with the mass distribution of MinD on a lipid bilayer (Fig. 2b, Supplementary Fig. 5). In

the presence of ATP, MinD monomers were detected as the predominant species in solution.

In contrast, the major species on the membrane was a MinD dimer already at sparse densities

corresponding to one particle per frame in the field of view (31.9 μm^2). However, it should be

noted that at the chosen imaging conditions for MSPT, the signal-to-noise ratio of monomers

(33 kDa) was too low for their quantitative detection (Supplementary Figures 9 and 10). Hence,

an adequate estimate of their membrane abundance cannot be stated.

One of the major advantages of iSCAT-based imaging is that it provides a direct estimate of

the molecular density on a bilayer from the number of detected particles, as compared to sin-

gle-molecule fluorescence where factors like labelling efficiency and photo bleaching have to

be taken into account. Accordingly, we could classify video sections of membrane-bound MinD

into conditions of different particle densities and observed the resulting change in the oligo-

meric distribution (Fig. 2c, Supplementary Fig. 6, 7 and Supplementary Movie 3). While MinD

was mainly present in the dimer state at low particle densities (0.1 μm^{-2}), the MinD population

shifted towards a broad distribution with higher-order complexes on crowded bilayers (0.8 μm^{-2}

). This data directly demonstrates that MinD indeed assembles into complexes larger than a

dimer as previously suggested[27,31]. To further investigate the structural determinants for

higher-order MinD assemblies, we performed the same MSPT experiment with a MinD mutant

(D40A) that is reported to predominantly reside in the dimeric state due to its impaired ability

to hydrolyse ATP[43]. According to the prevailing mechanism, it should not be possible for a

locked dimer mutant to participate in MinD self-assembly, due to its inability to switch between

the monomeric and dimeric state[23,41,44]. Nevertheless, we found a distinct population of MinD

D40A tetramers in our contour plot (Fig. 2d, Supplementary Fig. 8), suggesting that MinD is

capable of forming higher-order oligomers using an interface distinct from the canonical dimer-

isation site.

Another striking aspect of the D40A mutant was its clear separation into two distinct popula-

tions (dimer and tetramer), whereas the distribution of the WT protein appeared unresolved.

This result suggested that the WT was able to also recruit MinD monomers forming trimeric

species as intermediates, which were not as abundant for the D40A mutant. In order to provide

a more quantitative measure for the comparison between WT and D40A mutant, we decon-

volved their mass distributions into the underlying components to determine the relative abun-

dance of each species as a function of particle density on the bilayer. To estimate the shapes

of the individual components, we used our video simulation routine and determined separate

mass distributions for particles representing only one type of MinD oligomer at a range of par-

ticle densities (Supplementary Figures 9, 10 and Supplementary Discussion). Next, we fit the

mass distributions of MinD WT and D40A using a linear combination of simulated distributions

from six components corresponding to monomers – hexamers and extracted the relative abun-

dance of each oligomer as a function of particle density (Fig. 2e, g, Supplementary Fig. 11).

These graphs show a sequential appearance of increasingly larger oligomers of MinD for

higher molecule densities (Fig. 2f). Compared to the WT, the D40A mutant had a higher ten-
dency to populate stoichiometries with even numbers of subunits and to transform its dimer
state into the tetramer state, likely due to the increased stability of the dimer (Fig. 2h).

Furthermore, MSPT confers the unique possibility to determine oligomer-specific lateral diffu-
sion coefficients. This knowledge can be used to deduce structural information about the ob-
served molecules when considering the theory of Evans-Sackmann[45], which postulates the
relation of an object's membrane inclusion size with its respective diffusion coefficient. Assum-
ing a similar membrane viscosity as for pure DOPC membranes[46], we can thus estimate that
a MinD D40A dimer with a diffusion coefficient of 0.85 μm²/s has an inclusion size of 5 nm,
and a tetramer with 0.34 μm²/s an inclusion size of 9 nm (Supplementary Fig. 13). These esti-
mates imply that all monomers[43] of the tetramer are able to insert their MTS into the bilayer,
suggesting certain geometrical constraints for subunit orientation.

Figure 2 - Lateral MinD–MinD interactions lead to self-assembly into large homo-oligomers. (a)
Schematic of the canonical membrane binding-unbinding cycle of MinDE. Upon ATP-complexation,
MinD dimerises (1) and attaches to the membrane interface (2). In the event of MinE binding (3), MinE
stimulates the intrinsic ability of MinD to hydrolyse ATP, which leads to the release of MinD from the
membrane in its monomeric form (4). **(b)** MinD mass distribution in solution (grey line; n = 16,101 tra-
jectories) and upon attachment to the supported lipid bilayer (blue line; n = 13,917 trajectories). For
solution experiments, 175 nM MinD with 0.5 μM ATP were measured in the conventional MP landing
assay. The membrane mass distribution of MinD was determined using MSPT at a particle density of
0.03 μm⁻². **(c)** 2D kernel density estimation of membrane-attached MinD and **(d)** MinD D40A at particle
densities of 0.1 μm⁻² (light blue, n = 117,086 trajectories) and 0.8 μm⁻² (dark blue, n = 152,685 trajecto-
ries) and 0.1 μm⁻² (light green, n = 7,831 trajectories) and 0.8 μm⁻² (dark green, n = 3,150 trajectories),
respectively. Marginal probability distributions of both molecular mass and diffusion coefficient are pre-
sented at the top and right, respectively. **(e)** Representative mass distributions (grey) of MinD and MinD

D40A **(g)** and estimation (black line, coloured lines highlight underlying components) of its six compo-

nents for three different particle densities. MinD: 0.1 µm^{-2} – light blue, 0.4 µm^{-2} – blue, 0.7 µm^{-2} – dark

blue; MinD D40A: 0.1 µm^{-2} – light green, 0.3 µm^{-2} – green, 0.6 µm^{-2} – dark green. **(f,h)** Relative oligomer

abundance as a function of particle density. Error bars are the standard deviation of fitting results from

three data subsets. The oligomer analysis is based on a total of n = 1,102,940 trajectories for MinD and

n = 194,545 for MinD D40A.

Time-resolved mass analysis of single MinD trajectories

Aside from measuring a particle's location frame by frame, MSPT also allows to determine its

respective mass in a time-resolved fashion, thus enabling the detection of attachment and

detachment events along the trajectory of a single particle (Fig. 3a, Supplementary Fig. 14).

For MinD, the minimal expected mass increment (33 kDa) for monomer-wise turnover was

close to the measurement uncertainty (28 kDa standard deviation) of the mass for a single

frame. To minimise user bias, we employed a step-finding algorithm that locates mass change

points along trajectories based on statistical criteria[47].

Figure 3b displays examples of mass time courses for individual particles and the mass steps

detected by the algorithm. Note that in a majority of cases, MinD complexes retained their size

throughout their entire trajectory (Fig. 3b, upper panel). However, in ~10% of trajectories, the

mass of a tracked particle changed during its trajectory (Figure 3b, bottom panel), suggesting

the attachment or detachment of MinD subunits to and from the membrane-bound complex.

By analysing the sizes of mass steps along all trajectories, it was possible to obtain a distribu-

tion of subunit sizes attaching and detaching from membrane-bound MinD complexes (Fig.

3c), which appeared to mostly (dis-)assemble in one- or two-subunit increments at low particle

densities. However, for higher MinD particle densities, when larger oligomers had accumu-

lated, these complexes often turned-over greater subunits, as indicated by the shift of the dis-

tribution towards higher mass steps[27] (Fig. 3c, dark blue profile). Moreover, the combined in-

formation of a mass plateau level and its dwell time as annotated in Fig. 3b could be used to

extract the subunit turnover rates of each oligomer species (Fig. 3d). Here, the dwell time

preceding a mass increase could be used to deduce subunit attachment rates (Fig. 3d – upper

panel), whereas the dwell time followed by a mass decrease provided an estimate of detach-

ment rates (Fig. 3d – lower panel). The resulting lifetime plots suggested that membrane-at-

tached dimers had a faster subunit turnover than tetramers, implying higher stability of these

larger complexes.

Plateaus at the end of trajectories (Fig. 3b, bottom panel, last plateau), and trajectories without

any mass change at all (Fig. 3b, top panel), could be used to identify the molecular weight of

particles completely released (rl) from the membrane (Fig. 3e). Notably, this time-resolved

mass analysis of the individual trajectories significantly improved the mass resolution, as com-
pared to the median-based particle mass estimates used in Figure 2. Hence, MinD dimers,
trimers and tetramers were now fully resolved as separate peaks (Fig. 3e). Accordingly, one
could now recognize that the major species released from the membrane was a dimer at low
particle densities and a tetramer at high particle densities. The corresponding dwell time plot
showed that tetramers stayed associated to the bilayer significantly longer than dimers, in line
with a higher avidity in membrane binding conferred by additional MTS (Fig. 3f). For compari-
son, the dimer-arrested mutant MinD D40A almost exclusively dissociated as dimers or te-
tramers (Supplementary Fig. 15, 16). Taken together, our detailed trajectory analysis confirms
our previous observations suggesting that MinD WT assembles into species of higher order
with an intermediate trimer state. This indicates that subunits bind at a location different from
the canonical dimerisation site.

Figure 3 - Subunit (dis-)assembly of MinD particles diffusing on membranes. (a) Schematic rep-
resentation of the time-resolved mass analysis of single MinD trajectories, which reveals attachment
(at) and detachment (dt) events along the trajectory as well as a particle's full membrane release (rl) at
the end of its trajectory. **(b)** Representative mass time traces of MinD trajectories (grey line) and corre-
sponding step fits (black line) determined by a step-finding algorithm that locates mass change points
within a trajectory. **(c)** Mass step size distribution derived from step fits as depicted in (b) revealing MinD
subunit sizes for attachment (at) and detachment (dt) events at particle densities: 0.1 µm^{-2} – pale blue
(n = 20,796 plateaus), 0.3 µm^{-2} – light blue (n = 26,177 plateaus), 0.6 µm^{-2} –blue (n = 25,864 plateaus),
0.8 µm^{-2} –dark blue (n = 7,506 plateaus). **(d)** Dwell time plots for MinD particles before attachment (at)
events (top plot), representing the lengths of mass plateaus preceding a mass increase, and before
detachment (dt) events (bottom plot), representing the lengths of mass plateaus preceding a mass de-
crease. Dwell times are shown for the MinD dimer (66 kDa, light blue line) and the MinD tetramer state
(132 kDa, dark blue line). Plateau numbers for (at): dimer - n = 23,782; tetramer - n = 5,088; for (dt):

dimer - n = 3,143; tetramer - n = 10,406. Insets: Mean dwell times of the two oligomer states. Error bars

represent the standard deviation of the dwell times estimated by bootstrapping. **(e)** MinD mass distribu-

tion for membrane release (rl) events. MinD particle densities: 0.1 μm^{-2} – pale blue (n = 117,086 plat-

eaus), 0.3 μm^{-2} – light blue (n = 169,957 plateaus), 0.6 μm^{-2} – blue (n = 284,916 plateaus), 0.8 μm^{-2} –

dark blue (n = 120,654 plateaus). **(f)** Plot of the dwell times before membrane release (rl) for the MinD

dimer and tetramer state. Plateau numbers: dimer - n = 562,011; tetramer - n = 73,037. Inset: Mean

dwell times of the two oligomer states. Error bars represent the standard error of dwell times estimated

by bootstrapping.

MinE-induced formation of large heteromeric complexes

In the past decade, several different mechanistic models have been proposed to explain the

role of the ATPase-activating protein MinE for MinDE detachment dynamics[13,48,49]. Some of

these models are based on the cooperation of both MinE and MinD dimer to prompt mem-

brane-release through MinD ATPase activity stimulation[44,48]. However, this effect alone cannot

explain the recently observed cooperative membrane-detachment of MinDE filaments[27]. To

address this issue, we used MSPT to determine the stoichiometry of the membrane-bound

MinDE complex and followed its membrane dynamics on a molecular level.

In accordance with previous structural studies that suggest a conformational switch of MinE

allowing MinD binding only upon the encounter of membrane-bound MinD[48,50], we found no

indication for MinDE interaction in solution (Supplementary Fig. 17). In the presence of a sup-

ported lipid bilayer, however, the MinDE complex existed predominantly in a stable double-

dimeric state (Fig. 4a – light pink, Supplementary Fig. 18). Furthermore, if the MinDE complex

encountered more proteins on a crowded bilayer, MinE promoted the interconnection into very

large heteromeric MinDE complexes, a behaviour unexpected considering the common mod-

els (Fig. 4a – magenta). One possible explanation for this behaviour is the ability of a MinE

dimer to symmetrically bind to both sides of a MinD dimer, thus effectively acting as a bridge

between MinD assemblies[49]. Accordingly, our time-resolved mass step analysis revealed that

during subunit turnover on membrane-bound particles, predominantly dimeric and tetrameric

subunits attached (at) and detached (dt) at high particle densities of 0.6 and 0.8 μm^{-2} (Fig. 4b,

Supplementary Fig. 19). This effect required a critical minimum density on the bilayer, since

MinDE complexes in sparsely populated environments (0.1 and 0.3 μm^{-2}) mainly exhibited

conversion in their minimum subunit increments. Hence, their final oligomeric state during

membrane release (rl) resembled a MinDE dimer (Fig. 4c, light pink). Strikingly, on bilayers

with high protein density, MinDE complex sizes released from the membrane increased be-

yond the sizes observed for MinD alone and reached masses of >350 kDa (Fig. 4c, purple).

Based on our mass dwell time analysis, we found that the presence of MinE generally reduced
the diffusion coefficients (Fig. 4d) of MinDE oligomers and slowed down turnover rates (Fig.
4e, pink), when compared to their respective MinD versions (Fig. 4e, blue). In addition, MinDE
complexes were found to reside significantly longer on the membrane before full release (Fig.
4f). This suggests that MinE can effectively stabilise membrane-bound MinD when present in
equimolar amounts as previously suggested[13,49].

Figure 4 - MinE interconnects MinD oligomers into large complexes with a prolonged membrane
dwell time. (a) 2D kernel density estimation of membrane-attached MinDE complexes at particle den-
sities of 0.1 µm⁻² (light pink, n = 200,436 trajectories) and 0.8 µm⁻² (purple, n = 30,082 trajectories).
Marginal probability distributions of both molecular mass and diffusion coefficient are presented at the
top and right, respectively. **(b)** Mass step size distribution revealing MinDE subunit turnover (at and dt
events) on membrane-bound particles at particle densities of 0.1 µm⁻² – pale pink (n = 105,438 plat-
eaus), 0.3 µm⁻² – light pink (n = 73,471 plateaus), 0.6 µm⁻² – pink (n = 9,247 plateaus), 0.8 µm⁻² – dark
purple (n = 4,040 plateaus). **(c)** MinDE mass distribution for membrane release (rl) at MinDE membrane
particle densities of: 0.1 µm⁻² – pale blue (n = 200,436 plateaus), 0.3 µm⁻² – light blue (n = 158,660
plateaus), 0.6 µm⁻² – blue (n = 35,501 plateaus), 0.8 µm⁻² – dark blue (n = 26,708 plateaus). **(d)** Analysis
of oligomer specific diffusion coefficients for MinD (blue lines) and MinDE complexes (pink lines). light
blue/pink – dimer (MinD/MinDE n = 439,568/206,422 trajectories); dark blue/purple – tetramer dimer
(MinD/MinDE n = 118,922/47,136 trajectories). Inset: Mean diffusion coefficients of each oligomer state.
Error bars represent the standard deviation of the dwell times estimated by bootstrapping. **(e)** Dwell time
plots for MinD and MinDE attachment (at) events (top plot) as well as for MinD and MinDE detachment
(dt) events (bottom plot). Dwell times are shown for the MinD dimer (light blue line) and MinD tetramer
(dark blue line) as well as for their respective MinDE versions (hetero-dimer state – light pink, hetero-

tetramer state – purple). Plateau numbers for (at): MinD dimer – n = 23,782, MinDE dimer – n = 37,278,

MinD tetramer – n = 5,088, MinDE tetramer – n = 11,698; (dt): MinD dimer – n = 3,143, MinDE dimer –

n = 3,974, MinD tetramer – n = 10,406, MinDE tetramer – n = 22,501. Insets: Mean dwell times of the

two oligomer states. Error bars represent the standard deviation of the dwell times estimated by boot-

strapping. **(f)** Plot of the dwell times before membrane release (rl) for the MinDE dimer and tetramer

state and the respective MinD versions for comparison. Plateau numbers: MinD dimer – n = 562,011,

MinDE dimer – n = 277,782, MinD tetramer – n = 73,037, MinDE tetramer – n = 60,114. Inset: Mean

dwell times of the two oligomer states. Error bars represent the standard error of the dwell times esti-

mated by bootstrapping.

Conclusions

In this work, we presented a simple and versatile single-molecule-based method for the deter-

mination of membrane-associated oligomer distributions, subunit turnover and mass-resolved

residence times in the context of the *E. coli* Min system. Based on our results, we propose the

extension of the established membrane binding-unbinding models for MinDE self-organisa-

tion[44,48] (Fig. 5). Our data support the original model regarding nucleotide exchange from ADP

to ATP triggering membrane-dependent dimerisation of MinD, thus coupling the process to

energy dissipation. On the membrane, we find that lateral MinD interactions as well as recruit-

ment of MinD subunits from solution lead to the formation of a dynamic mixture of MinD oligo-

meric states that assemble through attachment and detachment of subunits at a location dif-

ferent from the canonical dimerisation site[31], a behaviour unexpected by common models. We

assume that the ability of MinD to assemble into these complexes is generally required for its

local self-accumulation and in combination with the initial nucleotide-dependent membrane

recruitment explains the observed attachment cooperativity during MinDE self-organisation. At

elevated membrane densities, a small percentage of MinD self-assembles into tetramers,

which either fall apart or interact with MinE. In contrast to the previously assumed cooperation

of the MinE and MinD dimer to prompt membrane-release[44,48], we found that MinE promotes

the interconnection[49] of heteromeric MinDE complexes. We assume that this behaviour is

based on the ability of two MinE dimers to symmetrically bind to both sides of a MinD dimer,

thus effectively acting as a bridge between MinD assemblies[49]. The existence of multivalent

membrane-bound structures would likewise explain the prolonged residence times of MinDE

complexes compared to their respective MinD oligomer variants. Notably, opposed to the pre-

viously assumed monomer detachment[51], we found MinDE to detach from the membrane in-

terface in complexes with sizes beyond 350 kDa, which could correspond to eight MinDE sub-

units. This corroborates that MinE could induce nucleotide-conversion of MinD subunits and

thereby weaken the overall membrane avidity of the MinDE complex prior to its full dissociation

from the membrane interface[27]. Upon membrane release, larger complexes might remain tem-

porarily stable, stay in the vicinity of the bilayer, and can potentially rebind close to their disso-

ciation spot, once MinD subunits have exchanged their nucleotide. An indication of such be-

haviour is the gradual shift of MinD/MinDE mass distributions, for both membrane recruitment

and release events, towards higher masses over the course of a video (Supplementary Fig.

20). To conclude, we believe that MinDE self-organisation arises from an interplay of cooper-

ative MinD membrane attachment into higher-order oligomers, anisotropy of the local MinD

concentration through quick re-binding of oligomers to the membrane, and ATP-dependent

membrane release of MinD assemblies coordinated by the ATPase-stimulating activity of

MinE.

The mechanistic dissection of the MinDE membrane-cycle described in this study constitutes

a detailed technical demonstration of the capabilities of mass-sensitive particle tracking

(MSPT). By combining the advantages of mass photometry[17,18], where biomolecular complex

stoichiometries are determined from the mass of individual particles, with single-particle track-

ing, MSPT provides invaluable insights into the complex dynamics of biomolecules on lipid

membranes. When exploring the size range for quantitative characterisation of transient mem-

brane interactions and oligomer stoichiometries, we found that >50 kDa proteins on supported

lipid bilayers could be reliably resolved with our system at relatively low particle densities (see

Supplementary Discussion). Although the application to even smaller proteins will require im-

proved methodology with regard to detection and data processing, the ability of label-free de-

tection of membrane complexes provides powerful advantages over conventional fluores-

cence-based single-particle tracking approaches: 1) The possibility to simultaneously deter-

mine particle densities, i.e. local concentrations on the bilayer; 2) The correlation of mass and

diffusion coefficient, which, in combination with a derived membrane inclusion size, provides

information about the stoichiometry and arrangement of a membrane-bound complex; 3) The

analysis of time-resolved subunit turnover and its kinetics by analysing mass changes along a

particle's trajectory; 4) Extended observation times due to the absence of photo-bleaching,

thus enabling the collection of high particle statistics in a short period of time. Ultimately, we

believe that mass-sensitive particle tracking will make a strong contribution to the quantitative

understanding of both prokaryotic and eukaryotic membrane-associated biological systems.

Figure 5 - Schematic of the proposed membrane-associated MinDE reaction cycle. (1) Upon nu-
cleotide exchange from ADP to ATP, membrane-dependent dimerisation of MinD is triggered. **(2)** On
the membrane, lateral MinD interactions and recruitment of MinD subunits from solution lead to MinD
higher-order structures that assemble through attachment of subunits at a location different from the
canonical dimerisation site. MinD assemblies then either dissociate from the membrane **(3)**, or encoun-
ter MinE **(4)**. MinE promotes the interconnection of very large heteromeric MinDE complexes that, due
to their multivalent MTS structure, reside significantly longer on the membrane interface **(5)**. **(6)** How-
ever, MinE also induces nucleotide-conversion of MinD subunits, thereby weakening the overall mem-
brane avidity of the MinDE complex, prior to its full release from the membrane interface in complexes
>350 kDa.

Methods

ADP/ATP stock solution
Both ADP and ATP stocks were prepared from their respective salt hydrates (#A2754 and
#A2383, Sigma Aldrich, St. Louis, USA), supplemented with an equal molar amount of $MgCl_2$
and adjusted to pH 7.5 with 1 M Tris-HCl. Final nucleotide concentration was spectroscopically
determined (λ = 259 nm; V-650, Jasco, Pfungstadt, Germany) using an extinction coefficient
of 15,400 M^{-1} cm^1.

Protein Purification and Modification
Purifications of MinD, MinD D40A and MinE were performed as previously described [52] and all
used plasmid constructs are indicated in Supplementary Table 4. Divalent streptavidin was
assembled as outlined in Howarth *et al.* [53], both pET21a-Streptavidin-Alive (Addgene plasmid
#20860) and pET21a-Streptavidin-Dead (Addgene plasmid #20859) were a gift from Alice Ting
[53]. Biotin-modification of aldolase (#28403842, Cytiva, Marlborough, USA) with EZ-Link™ Ma-
leimide-PEG2-Biotin (#A39261, Thermo Fisher Scientific, Waltham, USA) was achieved
through incubation at RT for 1 h and subsequent size-exclusion chromatography on a 16/600

Superdex 200 pg column (GE Healthcare, Pittsburgh, USA), equilibrated in storage buffer (50

mM HEPES pH 7.25, 150 mM KCl, 0.1 mM EDTA, 10% glycerol, 0.4 mM TCEP), using an

Äkta Pure chromatography system (GE Healthcare, Pittsburgh, USA). LC-MS and SDS-PAGE

was performed to assess purity and integrity of all purified or modified proteins. A customized

Bradford assay (#5000006, Bio-Rad Protein Assay; Bio-Rad Laboratories Inc., Hercules, USA)

was used to determine protein concentrations and single-use aliquots were flash frozen in

liquid nitrogen and stored at -80 °C.

Small Unilamellar Vesicles (SUVs) and Supported Lipid Bilayer (SLB) Formation

For the formation of small unilamellar vesicles (SUV), dioleoyl-sn-glycero-3-phosphocholine

(DOPC; #850375, Avanti Polar Lipids, Alabaster, USA), dioleoyl-sn-glycero-3-phosphoglycerol

(DOPG; #840475, Avanti Polar Lipids, Alabaster, USA) and 1,2-dioleoyl-sn-glycero-3-phos-

phoethanolamine-N-cap biotinyl (18:1 Biotinyl Cap PE; #870273, Avanti Polar Lipids, Alabas-

ter, USA) were dissolved in chloroform (Sigma Aldrich, St. Louis, USA) and mixed in a ratio of

70 mol % DOPC to 30 mol % DOPG or 70 mol % DOPC with 29.99 mol % DOPG and 0.01

mol % Biotinyl Cap PE. After solvent evaporation through nitrogen, residual chloroform was

removed for 1 h in a vacuum desiccator. Lipid film hydration was achieved with Min buffer (25

mM Tris/HCl pH 7.5, 150 mM KCl, 5 mM $MgCl_2$) and SUVs were formed through consecutive

freeze-thaw cycles (8-10) using liquid nitrogen and a 90 °C water bath. For monodisperse

vesicle distribution, lipid mixtures were extruded across a Whatman nucleopore membrane

(#110603, GE Healthcare, Chicago, USA) with a pore size of 50 nm for 37 passes.

SLBs were formed by fusion of SUVs on cleaned glass cover slides (Paul Marienfeld GmbH &

Co. KG, Lauda-Königshofen, Germany) that were assembled into a flow chamber through

double-sided sticky tape (Scotch, Conrad Electronic SE, Germany). Prior to their assembly,

cover slides (#0102242, #1.5, 24 x 60 mm; #0102062, #1.5, 24 x 24 mm) were cleaned by

sequential sonication in Milli-Q water, isopropanol, and Milli-Q water (15 min each), and sub-

sequently dried under a nitrogen stream. Slides were then activated for 30 s (30% power, 0.3

mbar) in a Zepto plasma cleaner (Diener electronic GmbH, Ebhausen, Germany) using oxygen

as the process gas. After flow chamber assembly, SUVs were added to each reaction chamber

at a final concentration of 0.4 mg/mL in Min buffer with additional 2 mM $CaCl_2$ to promote

vesicle rupture. Unfused SUVs were removed through subsequent washing with Min buffer.

Mass Calibration Curves and Diffusion Control

To convert interferometric scattering contrast into the respective protein mass, the contrast of

a set of mass standards was measured for each experimental setup – MP landing assays and

MSPT on the lipid interface. In line with Young *et al.* [17], mass calibration for landing assays

was performed in flow chambers filled with filtered (sterile syringe filters 0.45 μm cellulose

acetate membrane, VWR International, Radnor, USA) SLB buffer (25 mM Tris/HCl pH 7.5, 150

mM KCl). As mass standards, either 50 nM TEV protease (MPIB Core Facility), 50 nM Pierce™

recombinant protein A (#21184, Thermo Fisher Scientific, Waltham, USA) 50 nM bovine serum

albumin (#A4612, Sigma Aldrich, St. Louis, USA), 20 nM alcohol dehydrogenase (#A8656,

Sigma Aldrich, St. Louis, USA), or 20 nM β-amylase (#A8781, Sigma Aldrich, St. Louis, USA)

were injected and landing events recorded. To also enable mass calibration on supported lipid

bilayers, membranes containing 0.01 mol % Biotinyl Cap PE were used. As linker for the at-

tachment of biotinylated standard proteins, 2.5 nM divalent streptavidin were incubated for 10

min (RT) prior to the addition of 100 nM biotin labeled bovine albumin (#A8549, Sigma Aldrich,

St. Louis, USA), biotinylated Pierce™ protein A (#29989, Thermo Fisher Scientific, Waltham,

USA) or custom biotinylated aldolase (#28403842, Cytiva, Marlborough, USA). These concen-

trations correspond to 3-6 particles (median value) in the field of view for each standard protein,

equivalent to a particle density of 0.09 to 0.19 μm^{-2}.

For diffusion coefficient verification, 1.25 nM unconjugated tetravalent Streptavidin

(#SNN1001, Thermo Fisher Scientific, Waltham, USA) were incubated for 10 min (RT) on a

supported lipid membrane containing biotinylated lipids.

Mass Photometry Landing Assay

To determine the nucleotide-dependent solution-state of MinD (175 nM) and MinDE (175 nM),

all proteins were diluted in filtered (sterile syringe filters 0.45 μm cellulose acetate membrane,

VWR International, Radnor, USA) Min buffer in the presence of either 0.5 mM ADP or ATP.

Measurements were performed in flow chambers and 50 μL protein solution were flushed in

five times consecutively, to collect sufficient landing events.

***In Vitro* Reconstitution of Min Complexes on a Lipid Bilayer**

MinD

For MSPT of single MinD complexes on the lipid interface, we added increasing protein con-

centrations (50, 75, 100, 125, 150, 175 and 200 nM, n = 3 flow chambers each) to a flow

chamber with a supported lipid bilayer in the presence of 0.5 mM ATP in filtered Min buffer.

The same experiments were performed likewise for MinD D40A. Videos of 0.5 mM ATP in Min

buffer without protein were recorded as image background control (0.8 % of particles detected

as compared to videos containing MinD).

MinDE

Co-reconstitution of MinDE was performed in a similar manner as for MinD, except for the
addition of equimolar amounts of both reactants (50, 75, 100, 125, 150, 175 and 200 nM, n =
3 flow chambers each) to the sample chamber. For MinD D40AE, experiments were performed
with 200 nM of each protein.

Microscopy
Imaging of MinDE landing assays was performed on a custom-built interferometric scattering
microscope described in [17,54] with a 445 nm laser diode for illumination and 635 nm for focus-
stabilization. Image acquisition was controlled using custom-written software in Labview de-
scribed in [17]. Landing events were recorded at a frame rate of 1 kHz for 60 s per video. Videos
were saved 5-fold frame-averaged (200 Hz effective frame rate) and 3-fold pixel-binned (70.2
nm effective pixel size).

SLB experiments were carried out on a commercial Refeyn One^{MP} mass photometer (Refeyn
Ltd., Oxford, UK). Movies were acquired for either 45 s (landing assay of mass standards, Fig.
1) or 350 s (SLB assays) with the Aquire^{MP} (Refeyn Ltd., v2.3) software at a frame rate of 1
kHz. Movies were saved 5-fold frame-averaged (200 Hz effective frame rate) and 4-fold pixel-
binned (84.4 nm effective pixel size).

Data analysis
Image Processing of Conventional MP Landing Assays
Videos of the proteins MinD, MinE and their equimolar mixture MinDE landing on glass cover
slides were analysed with the software Discover^{MP} (version 2.1.0, Refeyn Ltd.), using a rolling
background removal strategy[17].

MinD monomers have a molecular mass of 33 kDa, which approaches the lower detection limit
of the mass photometer. We have therefore systematically determined the optimal frame av-
eraging factor *navg* and filter thresholds *T1* and *T2* for the detection of MinD monomers and
dimers. To this end, we generated a semi-synthetic movie that used frames from a video rec-
orded in a chamber with Min buffer alone to reconstruct the experimental background and
added simulated PSFs (same as fitting model PSF) as landing events that had the expected
scattering contrast of MinD monomers (contrast = 1.9×10^{-3}) or MinD dimers (contrast = $3.5 \times$
10^{-3}). We then varied *navg* as well as *T1* and *T2*, ran the analysis procedure, and evaluated
the number of true positive and false positive detections. To determine the maximum number
of true positive detections possible at the respective signal-to-noise ratio, we simulated 1000
frames with 100 landing events that were not allowed to spatially overlap closer than 12 pixels
and 26 frames temporally. Based on the simulation, we chose *navg* = 12, *T1* = 1.1 and *T2* =

0.2 to process the experimental videos. Using these parameters, the number of true positive

detections of monomers was 68.8 ± 6.6% (mean ± STD, n = 5 simulated videos), while the

total of false positive detections was 9.4 ± 2.1%. For dimers, a simulation with the same pa-

rameters gave 95.8 ± 0.8% true positive detections and 4.2 ± 1.1% false positive ones.

Mass Calibration for Landing Assays

Landing events imaged in three independent experiments per standard protein were pooled

and kernel density estimations were calculated. For each standard protein, the contrast asso-

ciated with a probability density maximum was determined and plotted against its nominal

mass (see Supplementary Table 2). In case of alcohol dehydrogenase and β-amylase, the

peak with higher contrast was used.

Error bars in Fig 1c represent the standard error of the peak position calculated from 10,000

bootstrap resamples.

Image Processing and Analysis Procedure for Single-Particle Tracking and Mass Determina-

tion of Biomolecules on SLBs

To detect and analyse diffusing biomolecules on a lipid bilayer, we applied a new image pro-

cessing strategy that removed the dominant static scattering background, while conserving the

shape of mobile features and displaying them on top of a shot-noise limited background.

Around each frame, a pixel-wise temporal median image was calculated that only contains the

features which did not move in the median period. This static background, calculated for each

frame, was then removed from the respective frame with index i according to equation (1) and

with n denoting the median half-size.

(1) $$mobile\ features\ frame_i = \left(\frac{frame_i}{median(frame_{i-n}:frame_{i+n})}\right) - 1$$

Through this process, moving objects in the mobile features frame appeared as undistorted

PSFs and displayed iSCAT contrasts similar to those obtained for molecules in the conven-

tional landing assay. We kindly received a custom-written Python script from Philipp Kukura

and Gavin Young which automatised this image analysis procedure and we modified it to load

and process videos in the Refeyn format (.mp files). To track individual proteins diffusing on

supported lipid bilayers in the background-corrected movies, the Python script also included a

single-particle tracking routine. For particle detection, a Laplace filter was applied to each

frame (*scipy.ndimage.filters.gaussian_laplace* function), to suppress shot noise and highlight

potential particles. The Laplace-filtered images were thresholded, such that only objects with

at least the size of 53 kDa remained, a threshold that would preserve streptavidin as our small-

est standard protein and MinD dimers. Local maxima were then found (*scipy.ndimage.fil-*
ters.maximum_filter function) as candidate pixels that contained a particle. Around each of
these candidate pixels in the non-Laplace-filtered image, a region of interest (ROI) of 13 by 13
pixels (84.4 nm/pixel) was excised and fit by the model PSF used in the Discover[MP] software
to extract particle contrast and location at sub-pixel resolution. Particle locations were linked
into trajectories using the *Trackpy* package (version 0.4.2, *link_df* function) [55]. Only trajectories
of particles with a lifetime of at least 5 frames (25 ms) were considered for subsequent anal-
yses. Supplementary Table 1 summarises all parameters used for particle detection, fitting and
trajectory linking.

Determination of Diffusion Coefficients
To determine translational diffusion coefficients from individual trajectories, a jump-distance
analysis was performed as described in [36]. Cumulative frequency distributions of jump dis-
tances for time lags from Δ1 to Δ4 frames were fitted globally (*scipy.optimize.least_squares*,
Trust Region Reflective algorithm) with either one or two species depending on which model
resulted in a reduced chi-squared statistic.

(2)
$$P_{1c}(r^2, \Delta t) = 1 - e^{-\frac{r^2}{4D_1\Delta t + \sigma}}$$

(3)
$$P_{2c}(r^2, \Delta t) = 1 - \sum_{i=1}^{2} a_i \, e^{-\frac{r^2}{4D_i\Delta t + \sigma}}, \qquad \sum_{i=1}^{2} a_i = 1$$

701 D_i and a_i are the diffusion coefficient and abundance of component *i*, *Δt* is the time lag and *σ*
702 is an offset parameter taking into account localisation uncertainty and motion blur [56]. In the
703 case of a two-component model (~10% of all trajectories), an effective diffusion coefficient was
704 calculated according to

(4)
$$D_t = a_1 D_1 + a_2 D_2,$$

For comparison with the jump distance approach in the diffusion simulation (Supplementary
Figure 2), diffusion coefficients were also estimated based on the mean squared displacement
(MSD). Here, the first three or four non-trivial time-averaged MSD points were included in the
linear regression depending on the trajectory length [56].
In both fit routines, the diffusion coefficient was restricted to values greater than 0.0001 μm^2/s
while the offset parameter σ was unconstrained. Trajectories where the diffusion coefficient
returned by the fit hit the lower boundary were excluded from the analysis.

Kernel Density Estimation in 1D

For univariate distributions, kernel density estimations (KDEs) were computed with the Python package *KDEpy* (version 1.0.10, *FFTKDE* function) [https://github.com/tommyod/KDEpy], using a Gaussian kernel and the Improved Sheather-Jones plug-in bandwidth selector[57].

Kernel Density Estimation in 2D

2D maps of the diffusion coefficient and mass were computed with the Python package *fastkde* (version 1.0.14, *fastkde.pdf* function) [58]. The probability density for the diffusion coefficient was calculated in base 10 logarithmic space. Marginal distributions displayed on top and on the right of the graph were calculated by summing up the bivariate probability densities along the y- and x-axis, respectively.

The resulting 2D kernel densities were plotted as filled contours with either six (if an overlay of two conditions is shown) or eight linearly spaced levels. The opacity of the levels was reduced linearly from 100% for the highest to 0% for the lowest level, rendering the lowest level fully transparent.

Mass Calibration for MSPT

Analogous to the mass calibration for landing assays, peak contrasts were related to the nominal mass of streptavidin (Strep) or to the mass of streptavidin and the standard protein (Strep-ALD, Strep-BSA, Strep-prA). For the streptavidin-protein complexes, the peak with lowest contrast was excluded from analysis as it resembles streptavidin without an attached standard protein (blue crosses, Supplementary Fig. 4). The two other peak(s) with higher contrast were assigned to dimeric and tetrameric aldolase, monomeric and dimeric BSA and monomeric protein A (Supplementary Table 3).

Error bars in Fig 1c represent the standard error of the peak position calculated from 10,000 bootstrap resamples.

Simulated Videos of Diffusing Particles

To investigate the influence of extreme diffusion coefficients and high particle densities on the fidelity of our MSPT analysis, we implemented a Python routine that adds artificial diffusing particles to an experimental video of an empty SLB. Thereby, it was possible to control the number of particles in the field of view (FOV) and to individually set both the iSCAT contrast and the particles' diffusion coefficient. In a first step, the trajectories of all particles were generated by choosing a random starting location within an area that is 4 times the video FOV size (x- and y-axes extended 2-fold). The travelled per-frame distance Δs in the x and y directions was drawn randomly from normal distributions with standard deviation $2D\Delta t$ (equations 5).

(5) $$p(\Delta s) = \frac{1}{\sqrt{2\pi(2D\Delta t)^2}}e^{-\frac{s^2}{2\cdot(2D\Delta t)^2}}$$

D is the diffusion coefficient converted into pixel dimensions and *Δt* the frame time. In order to

keep the number of particles in the simulated area constant, we added periodic boundaries.

Whenever a particle would cross a boundary of the simulated area, its location was shifted to

the corresponding location at the opposite boundary. Trajectory coordinates that entered the

FOV of the experimental video, i.e., 128 x 35 pixels, were populated with PSFs of a defined

ratiometric contrast according to the Refeyn fitting model. Finally, the background-free rati-

ometric video of simulated PSFs was merged with a raw experimental iSCAT video of an empty

SLB via pixel-wise calculation of

(6) $$contrast_{simulated} = (contrast_{ratiometric} + 1) \cdot camera\ count_{experimental}$$

The resulting simulated videos were then analysed by our MSPT analysis routine in the same

way as an experimental video with real particles.

Analysis of MinD and MinD40A Oligomerisation

Each trajectory was assigned to an apparent membrane protein density determined as the

median of all trajectories detected during the trajectory`s lifetime, divided by the area of the

FOV (31.9 μm^2). For trajectory densities ranging from 0.03 to 0.8 μm^{-2} (corresponding to 1 to

27 particles in the FOV), the KDE of each mass distribution was calculated. The densities were

binned in an overlapping manner, such that all trajectories were selected that had trajectory

numbers within a range of plus or minus two (e.g. at a density of 0.1 μm^{-2}, corresponding to 4

trajectories/FOV, trajectories with membrane coverage between 2 and 6 trajectories/FOV were

pooled). Each KDE was fit with a linear combination of monomer to hexamer mass distributions

that were inferred from single component simulations of MinD or MinD D40A at varying particle

densities (Supplementary Fig. 9). The mass distributions for each oligomer and density were

selected based on the number of localised particles in the experiment as a certain fraction of

particles (~40%) is lost in the trajectory linking and filtering (minimum length of 5 frames) pro-

cess (Supplementary Fig. 12). Additionally, the sum of contributions of the six components

was constrained to add up to unity in the least-squares sense. Prior to fitting, the simulated

mass distributions were smoothed using a moving average with a window length of 4.5 kDa.

To estimate the uncertainty of the extracted abundances of the components, the dataset at

each trajectory density was split randomly into three samples before fitting each subset indi-

vidually. This procedure was repeated 10 times and the average standard deviation of the per-

split results was calculated. Unless otherwise stated, the reported particle densities throughout

the manuscript correspond to trajectory densities (i.e. linked particle densities) as a more ro-
bust means to estimate the current membrane coverage (in contrast to localised particle den-
sities; see Supplementary Figs. 7, 12).
789
Step Detection
To detect mass change points during trajectories, we employed a MATLAB (R2020a, The
MathWorks, Natick, MA) implementation of the Kalafut-Visscher algorithm[59] described in[47].
While the algorithm has no parameters except the time series itself, we noticed that the results
depended on the length of the time series and the position of potential change points if they
were located close to the beginning or the end of a trajectory. Thus, change points were more
likely to be inserted, the shorter a time series was and the closer a point was to the boundaries
of the time series. To address this ambiguity, we concatenated the trajectories of a specific
imaging condition (e.g. of all videos containing MinD, Supplementary Fig. 14a) and divided this
concatenated series into n subsets of equal length l, which were then analysed with the step
detection algorithm (Supplementary Fig. 14b). This procedure enabled an equal treatment of
trajectories of different lengths and avoided bias in change point detection at the beginning
and the end of trajectories. Additionally, step identification was repeated $l-1$ times shifting the
start point of the linked time series circularly by one increment in each iteration. This second
procedure ensured that the detected steps would not depend on the start points of a subset.
From the l outputs generated in total, the relative significance of a step was reflected by the
fraction f of iterations, in which a change point was found at a particular location, ranging from
0 (never) to 1 (in every iteration, Supplementary Fig. 14c).
The choice of fraction f as well as the length l is somewhat arbitrary yet interconnected. In-
creasing l leads to fewer detected steps which can be compensated for by reducing f. Since
the longest trajectory in all datasets has a length of 558 frames and an increasing length leads
to a finer grating of the tunable parameter f, we set l to 1000 frames. By visual inspection, we
chose a fraction f of 0.25 as an appropriate tradeoff between under- and over-identification of
steps considering the noise level in our trajectories (Supplementary Fig. 14d).
814
Subunit Attachment/Detachment During Membrane Diffusion and Release of Complexes
The concatenated mass traces and step-fitting results were split into their original trajectories
(Supplementary Fig. 14e). For each trajectory, step heights (equivalent to mass changes) and
dwell lengths (equivalent to time intervals without a change of mass) were extracted and cat-
egorised into three classes (Fig. 3b). Depending on whether the sign of the succeeding mass
change is positive or negative, the preceding level is classified as an attachment (at) or as a
detachment plateau (dt), respectively. The last plateau in a trajectory prior to the dissociation
of the whole particle from the membrane is regarded separately as release (rl). If no step is

detected in the mass time series as it was the case in ~90% of all trajectories, the single plat-
eau is also classified as release (rl). Particles leaving the FOV have been excluded from the
analysis. Therefore, it can be assumed that the end of a trajectory represents the particle's
release from the membrane. Generally, the mass which is initially binding to the membrane is
not included in the list of steps for attachment or detachment events.

The distribution of mass changes during the trajectory as well as the distribution of complex
mass released from the membrane for different densities are displayed as univariate kernel
density estimates. To characterise the residence times of different oligomers, histograms of
the observed dwell lengths were generated for plateaus with masses that correspond to di-
meric (66 ± 17 kDa) or tetrameric (132 ± 17 kDa) MinD complexes. For each variant the aver-
age dwell times were calculated according to equation *(7)*

$$\bar{t} = \frac{\sum_{i=m}^{n} N_i t_i}{\sum_{i=m}^{n} N_i} - t_m,$$

(7)

835 where N_i is the total number of plateaus with dwell time t_i, and t_m represents the most frequent
836 dwell time. Due to the noise level of mass detections in our trajectories, short dwell times could
837 not be quantitatively detected. Hence, dwell times shorter than 5 frames (25 ms) were excluded
838 from the analysis.

839 The standard error of the mean dwell time was estimated from 10,000 bootstrap resamples.

Acknowledgments

We gratefully thank the group of Philipp Kukura at the University of Oxford for hosting early
experiments and Gavin Young for sharing image processing and particle tracking code. We
thank Max Felix Hantke, Josiah Kane, and the rest of the Refeyn software team for providing
mass photometry image analysis code and their support. We thank Daniel Bollschweiler (Cryo-
EM MPIB Core Facility) for the initial introduction to the commercial Refeyn One^{MP} mass pho-
tometer, the MPIB Biochemistry Core Facility (Recombinant Protein Production) and Kerstin
Andersson for assistance with protein purification, S. Bauer for assistance with vesicle prepa-
ration and Florian Stehr for helpful discussions. T.H. and P.S. acknowledge funding through
the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - Project-ID
201269156 – SFB 1032 (A09) and B.R. and P.S. through Project 269423233 – TRR 174. B.R.
was supported by a DFG fellowship through the Graduate School of Quantitative Biosciences
Munich (QBM). N.H. was supported by a DFG return grant HU 2462/3-1. P.S. acknowledges
support through the research network MaxSynBio via the joint funding initiative of the German
Federal Ministry of Education and Research (BMBF) and the Max Planck Society.

Author Contributions

T.H., F.S., N.H. contributed equally to this work. T.H., N.H., F.S. and P.S. conceived the study. T.H. designed and performed all experiments. B.R. designed and purified proteins. F.S. and N.H. analysed the data. T.H., F.S. and N.H. wrote the manuscript draft. All authors discussed and interpreted results. All authors revised, reviewed and approved the manuscript.

Competing Interests Statement

The authors declare no conflict of interest.

Data Availability Statement

The datasets generated and analysed during the current study are available from the corresponding authors on reasonable request.

Code Availability Statement

The custom-written Python script for particle detection and particle simulation (both as binary executable) as well as particle analysis (source code accessible) is available from the corresponding authors on reasonable request.

References

1. Gonzalez, L. & Scheller, R. H. Regulation of membrane trafficking: Structural insights from a Rab/effector complex. *Cell* vol. 96 755–758 (1999).
2. Cho, W. & Stahelin, R. V. Membrane-protein interactions in cell signaling and membrane trafficking. *Annual Review of Biophysics and Biomolecular Structure* vol. 34 119–151 (2005).
3. Bagheri, Y., Ali, A. A. & You, M. Current Methods for Detecting Cell Membrane Transient Interactions. *Frontiers in Chemistry* vol. 8 (2020).
4. Miller, H., Zhou, Z., Shepherd, J., Wollman, A. J. M. & Leake, M. C. Single-molecule techniques in biophysics: A review of the progress in methods and applications. *Reports Prog. Phys.* **81**, 024601 (2018).
5. Manzo, C. & Garcia-Parajo, M. F. A review of progress in single particle tracking: From methods to biophysical insights. *Reports on Progress in Physics* vol. 78 (2015).
6. Gelles, J., Schnapp, B. J. & Sheetz, M. P. Tracking kinesin-driven movements with nanometre-scale precision. *Nature* **331**, 450–453 (1988).
7. Funatsu, T., Harada, Y., Tokunaga, M., Saito, K. & Yanagida, T. Imaging of single fluorescent molecules and individual ATP turnovers by single myosin molecules in aqueous solution. *Nature* **374**, 555–559 (1995).

8.   Schmidt, T., Schütz, G. J., Baumgartner, W., Gruber, H. J. & Schindler, H. Imaging of
single molecule diffusion. *Proc. Natl. Acad. Sci. U. S. A.* **93**, 2926–2929 (1996).

9.   Taylor, R. W. *et al.* Interferometric scattering microscopy reveals microsecond
nanoscopic protein motion on a live cell membrane. *Nat. Photonics* **13**, 480–487
(2019).

10.   Kukura, P. *et al.* High-speed nanoscopic tracking of the position and orientation of a
single virus. *Nat. Methods* **6**, 923–927 (2009).

11.   Jacobsen, V., Stoller, P., Brunner, C., Vogel, V. & Sandoghdar, V. Interferometric
optical detection and tracking of very small gold nanoparticles at a water-glass
interface. *Opt. Express* **14**, 405 (2006).

12.   Ueno, H. *et al.* Simple dark-field microscopy with nanometer spatial precision and
microsecond temporal resolution. *Biophys. J.* **98**, (2010).

13.   Loose, M., Fischer-Friedrich, E., Herold, C., Kruse, K. & Schwille, P. Min protein
patterns emerge from rapid rebinding and membrane interaction of MinE. *Nat. Struct.*
Mol. Biol. **18**, 577–583 (2011).

14.   Ha, T. & Tinnefeld, P. Photophysics of fluorescent probes for single-molecule
biophysics and super-resolution imaging. *Annual Review of Physical Chemistry* vol. 63
595–617 (2012).

15.   Garcia-Parajo, M. F., Segers-Nolten, G. M. J., Veerman, J. A., Greve, J. & Van Hulst,
N. F. Real-time light-driven dynamics of the fluorescence emission in single green
fluorescent protein molecules. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 7237–7242 (2000).

16.   Lindfors, K., Kalkbrenner, T., Stoller, P. & Sandoghdar, V. Detection and spectroscopy
of gold nanoparticles using supercontinuum white light confocal microscopy. *Phys.*
Rev. Lett. **93**, (2004).

17.   Young, G. *et al.* Quantitative mass imaging of single biological macromolecules.
Science **360**, 423–427 (2018).

18.   Piliarik, M. & Sandoghdar, V. Direct optical sensing of single unlabelled proteins and
super-resolution imaging of their binding sites. *Nat. Commun.* **5**, (2014).

19.   de Boer, P. A. J., Crossley, R. E. & Rothfield, L. I. A division inhibitor and a topological
specificity factor coded for by the minicell locus determine proper placement of the
division septum in E. coli. *Cell* **56**, 641–649 (1989).

20.   Raskin, D. M. & de Boer, P. A. MinDE-dependent pole-to-pole oscillation of division
inhibitor MinC in Escherichia coli. *J. Bacteriol.* **181**, 6419–24 (1999).

21.   Raskin, D. M. & De Boer, P. A. J. Rapid pole-to-pole oscillation of a protein required
for directing division to the middle of Escherichia coli. *Proc. Natl. Acad. Sci. U. S. A.*
96, 4971–4976 (1999).

22.   Szeto, T. H., Rowland, S. L., Rothfield, L. I. & King, G. F. Membrane localization of

MinD is mediated by a C-terminal motif that is conserved across eubacteria, archaea,

and chloroplasts. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 15693–15698 (2002).

23. Hu, Z., Gogol, E. P. & Lutkenhaus, J. Dynamic assembly of MinD on phospholipid

vesicles regulated by ATP and MinE. *Proc. Natl. Acad. Sci.* **99**, 6761–6766 (2002).

24. Hsieh, C. W. *et al.* Direct MinE-membrane interaction contributes to the proper

localization of MinDE in E. coli. *Mol. Microbiol.* **75**, 499–512 (2010).

25. Shih, Y. L. *et al.* The N-terminal Amphipathic Helix of the topological specificity factor

mine is associated with shaping membrane curvature. *PLoS One* **6**, e21425 (2011).

26. Loose, M., Fischer-Friedrich, E., Ries, J., Kruse, K. & Schwille, P. Spatial regulators

for bacterial cell division self-organize into surface waves in vitro. *Science* **320**, 789–

792 (2008).

27. Miyagi, A., Ramm, B., Schwille, P. & Scheuring, S. High-speed atomic force

microscopy reveals the inner workings of the MinDE protein oscillator. *Nano Lett.* **18**,

288–296 (2018).

28. Ayed, S. H. *et al.* Dissecting the role of conformational change and membrane binding

by the bacterial cell division regulator MinE in the stimulation of MinD ATPase activity.

J. Biol. Chem. **292**, 20732–20743 (2017).

29. Ghosal, D., Trambaiolo, D., Amos, L. A. & Löwe, J. MinCD cell division proteins form

alternating co-polymeric cytomotive filaments. *Nat. Commun.* **5**, 5341 (2014).

30. Ye, W. *et al.* Plasmonic Nanosensors Reveal a Height Dependence of MinDE Protein

Oscillations on Membrane Features. *J. Am. Chem. Soc.* **140**, (2018).

31. Heermann, T., Ramm, B., Glaser, S. & Schwille, P. Local Self-Enhancement of MinD

Membrane Binding in Min Protein Pattern Formation. *J. Mol. Biol.* **432**, 3191–3204

(2020).

32. Kretschmer, S., Zieske, K. & Schwille, P. Large-scale modulation of reconstituted Min

protein patterns and gradients by defined mutations in MinE's membrane targeting

sequence. *PLoS One* **12**, 1–16 (2017).

33. Ortega Arroyo, J. *et al.* Label-free, all-optical detection, imaging, and tracking of a

single protein. *Nano Lett.* **14**, 2065–2070 (2014).

34. Mosby, L. S. *et al.* Myosin II Filament Dynamics in Actin Networks Revealed with

Interferometric Scattering Microscopy. *Biophys. J.* **118**, 1946–1957 (2020).

35. Howarth, M. *et al.* A monovalent streptavidin with a single femtomolar biotin binding

site. *Nat. Methods* **3**, 267–273 (2006).

36. Weimann, L. *et al.* A Quantitative Comparison of Single-Dye Tracking Analysis Tools

Using Monte Carlo Simulations. *PLoS One* **8**, (2013).

37. Johansson, B., Höök, F., Klenerman, D. & Jönsson, P. Label-free measurements of

the diffusivity of molecules in lipid membranes. *ChemPhysChem* **15**, 486–491 (2014).

38.   Horton, M. R., Höfling, F., Rädler, J. O. & Franosch, T. Development of anomalous
969   diffusion among crowding proteins. *Soft Matter* **6**, 2648–2656 (2010).

970   39.   Horton, M. R., Reich, C., Gast, A. P., Rädler, J. O. & Nickel, B. Structure and
971   dynamics of crystalline protein layers bound to supported lipid bilayers. *Langmuir* **23**,
972   6263–6269 (2007).

973   40.   Stellmacher, L. *et al.* Acid-Base Catalyst Discriminates between a Fructose 6-
974   Phosphate Aldolase and a Transaldolase. *ChemCatChem* **7**, 3140–3151 (2015).

975   41.   Hu, Z. & Lutkenhaus, J. Topological regulation of cell division in E. coli:
976   Spatiotemporal oscillation of MinD requires stimulation of its ATPase by MinE and
977   phospholipid. *Mol. Cell* **7**, 1337–1343 (2001).

978   42.   Ramm, B. *et al.* The MinDE system is a generic spatial cue for membrane protein
979   distribution in vitro. *Nat. Commun.* **9**, (2018).

980   43.   Wu, W., Park, K. T., Holyoak, T. & Lutkenhaus, J. Determination of the structure of the
981   MinD-ATP complex reveals the orientation of MinD on the membrane and the relative
982   location of the binding sites for MinE and MinC. *Mol. Microbiol.* **79**, 1515–1528 (2011).

983   44.   Park, K. T., Wu, W., Lovell, S. & Lutkenhaus, J. Mechanism of the asymmetric
984   activation of the MinD ATPase by MinE. *Mol. Microbiol.* **85**, 271–281 (2012).

985   45.   Evans, E. & Sackmann, E. Translational and rotational drag coefficients for a disk
986   moving in a liquid membrane associated with a rigid substrate. *J. Fluid Mech.* **194**,
987   553–561 (1988).

988   46.   Khmelinskaia, A., Mücksch, J., Petrov, E. P., Franquelim, H. G. & Schwille, P. Control
989   of Membrane Binding and Diffusion of Cholesteryl-Modified DNA Origami
990   Nanostructures by DNA Spacers. *Langmuir* **34**, 14921–14931 (2018).

991   47.   Little, M. A. *et al.* Steps and bumps: Precision extraction of discrete states of
992   molecular machines. *Biophys. J.* **101**, 477–485 (2011).

993   48.   Park, K. T. *et al.* The min oscillator uses MinD-dependent conformational changes in
994   MinE to spatially regulate cytokinesis. *Cell* **146**, 396–407 (2011).

995   49.   Vecchiarelli, A. G. *et al.* Membrane-bound MinDE complex acts as a toggle switch that
996   drives Min oscillation coupled to cytoplasmic depletion of MinD. *Proc. Natl. Acad. Sci.*
997   *U. S. A.* **113**, E1479–E1488 (2016).

998   50.   Ghasriani, H. & Goto, N. K. Regulation of symmetric bacterial cell division by MinE
999   What is the role of conformational dynamics? *Commun. Integr. Biol.* **4**, 101–103
1000   (2011).

1001   51.   Lackner, L. L., Raskin, D. M. & De Boer, P. A. J. ATP-dependent interactions between
1002   Escherichia coli Min proteins and the phospholipid membrane in vitro. *J. Bacteriol.*
1003   **185**, 735–749 (2003).

1004   52.   Ramm, B., Glock, P. & Schwille, P. In vitro reconstitution of self-organizing protein

1005    patterns on supported lipid bilayers. *J. Vis. Exp.* 1–13 (2018) doi:10.3791/58139.

1006 53. Howarth, M. & Ting, A. Monovalent streptavidin expression and purification. *Protoc.*

1007    *Exch.* (2008) doi:10.1038/nprot.2008.81.

1008 54. Cole, D., Young, G., Weigel, A., Sebesta, A. & Kukura, P. Label-Free Single-Molecule

1009    Imaging with Numerical-Aperture-Shaped Interferometric Scattering Microscopy. *ACS*

1010    *Photonics* **4**, 211–216 (2017).

1011 55. Allan, D. *et al.* soft-matter/trackpy: Trackpy v0.4.2. (2019)

1012    doi:10.5281/ZENODO.3492186.

1013 56. Michalet, X. Mean square displacement analysis of single-particle trajectories with

1014    localization error: Brownian motion in an isotropic medium. *Phys. Rev. E - Stat.*

1015    *Nonlinear, Soft Matter Phys.* **82**, 041914 (2010).

1016 57. Botev, Z. I., Grotowski, J. F. & Kroese, D. P. Kernel density estimation via diffusion.

1017    *Ann. Stat.* **38**, 2916–2957 (2010).

1018 58. O'Brien, T. A., Kashinath, K., Cavanaugh, N. R., Collins, W. D. & O'Brien, J. P. A fast

1019    and objective multidimensional kernel density estimation method: FastKDE. *Comput.*

1020    *Stat. Data Anal.* **101**, 148–160 (2016).

1021 59. Kalafut, B. & Visscher, K. An objective, model-independent method for detection of

1022    non-uniform steps in noisy signals. *Comput. Phys. Commun.* **179**, 716–723 (2008).

1023