<u>**Title:**</u>

Pathogen invasion-dependent tissue reservoirs and plasmid-encoded antibiotic degradation
promote within-host plasmid transfer

<u>**Authors:**</u>

Erik Bakkeren[1,2,3*], Joana A. Herter[1], Jana S. Huisman[4,5], Ersin Gül[1], Alexander O. Brachmann[1], Jörn
Piel[1], Roland R. Regoes[4], Sebastian Bonhoeffer[4], Médéric Diard[6,7], Wolf-Dietrich Hardt[1*]

<u>**Affiliations:**</u>

[1]Institute of Microbiology, Department of Biology, ETH Zurich, Zurich, Switzerland
[2]Present affiliation: Department of Zoology, University of Oxford, Oxford, UK
[3]Present affiliation: Department of Biochemistry, University of Oxford, Oxford, UK
[4]Institute of Integrative Biology, Department of Environmental Systems Science, ETH Zurich, Zurich,
Switzerland
[5]Swiss Institute of Bioinformatics, Lausanne, Switzerland
[6]Biozentrum, University of Basel, Basel, Switzerland
[7]Botnar Research Centre for Child Health, Basel, Switzerland

<u>*for correspondence:</u>
hardt@micro.biol.ethz.ch
erik.bakkeren@zoo.ox.ac.uk

<u>**Key words:**</u>

Plasmids, Salmonella, Antibiotic resistance, tissue reservoirs, persistence

Abstract:

Many plasmids encode antibiotic resistance genes. Through conjugation, such plasmids can rapidly disseminate between bacterial strains. Previous work has identified gut-luminal donor/recipient blooms and tissue-lodged plasmid-bearing persister cells of the enteric pathogen *Salmonella enterica* serovar Typhimurium (*S*.Tm) that survive antibiotic-therapy in host tissues, as factors promoting plasmid dissemination among *Enterobacteriaceae*. However, the buildup of tissue reservoirs and their contribution to plasmid spread await experimental demonstration. Here, we asked if reseeding-acquisition-invasion cycles by *S*.Tm could serve to diversify tissue-lodged plasmid reservoirs, and thereby promote plasmid spread. Starting with intraperitoneal mouse infections, we demonstrate that *S*.Tm reseeding the gut lumen from either growing populations or persisters initiates clonal expansion. ESBL plasmid-encoded gut luminal antibiotic degradation by donors can foster recipient survival under beta-lactam antibiotic treatment, enhancing transconjugant formation upon reseeding. *S*.Tm transconjugants can subsequently re-enter host tissues introducing the new plasmid into the tissue-lodged reservoir. Population dynamics analyses pinpoint recipient migration into the gut lumen as rate-limiting for plasmid transfer dynamics in our model. Priority effects may limit reservoir formation in host tissues. Once transconjugant reservoirs are formed, persisters and chronically infecting transconjugant populations may serve complementary functions. The former are more numerous and may be more efficient vehicles promoting plasmid spread, while persister subpopulations guarantee plasmid reservoir maintenance even if the reservoir lacks appropriate resistance genes. Overall, our proof of principle data indicates that luminal antibiotic degradation and shuttling between the gut lumen and tissue-resident reservoirs can promote the accumulation and spread of plasmids within a host over time.

Main text:

Introduction:

The accumulation of antibiotic resistance genes in pathogenic bacterial strains is an important cause of antibiotic treatment failure. Conjugative antibiotic resistance plasmids are key drivers of this accumulation, accelerating the emergence of new bacterial strains genetically resistant to antibiotics (1). The mechanisms that contribute to the spread of resistance plasmids within an infected host are still not fully established.

Recently, we showed that the spread of plasmids within hosts can be aided by reservoirs of cells phenotypically recalcitrant to antibiotic therapy (2). This process is called persistence and refers to a property of bacterial populations in which one subpopulation is killed by an antibiotic parallel to another subpopulation that is killed slowly ("persisters"), defined by phenotypic and not genotypic properties (3, 4). Persisters can not only lead to antibiotic treatment failure, but they also have important implications for the evolution of antibiotic resistance (5-7) or virulence (8, 9), and serve as long term reservoirs promoting the spread of resistance plasmids. We showed this using the model organism *Salmonella enterica* serovar Typhimurium (*S*.Tm) (2). Invasive *S*.Tm cells form reservoirs of recalcitrant cells inside of host tissues (i.e., tissue persister reservoirs) that are difficult to treat with antibiotics (2, 9-13). After treatment, these *S*.Tm cells can reseed the gut lumen from their reservoirs along with any plasmids they carry. In the gut lumen, this provides the plasmids with the opportunity to conjugate into new host bacteria. Thereby, plasmids can subvert persisters to ensure long-term association within a given host. Besides persisters, *S*.Tm tissue invasion also creates a susceptible subpopulation within the host tissues. In the absence of antibiotic therapy, the latter cells are generally more numerous than the persisters and are thought to drive chronic infections (12-15). Moreover, these cells are known to re-seed the gut lumen over time (16). However, the role of this second tissue-lodged pathogen population in plasmid dissemination remained to be formally established.

Although very specific characteristics define antibiotic persistence (i.e., biphasic killing curves defining the susceptible and tolerant subpopulations) (3), there are many similarities to persistent (aka long-term, chronic) infection. Numerous bacterial pathogens can survive or evade host immune defenses and antibiotics alike in a recalcitrant state (e.g. *Pseudomonas aeruginosa, Escherichia coli, Staphylococcus aureus,* or *Salmonella enterica*) (4, 8, 17). Using mouse models, tissue reservoirs of *S*.Tm antibiotic persisters located in the intestinal mucosa or systemic sites can reseed the gut lumen after the cessation of antibiotic therapy (2, 9, 12, 18, 19). In chronically infected hosts, the pathogen population tends to diversify forming non-growing and growing subpopulations which are kept in check by immune defenses, but can rise upon neutralization of cytokines (e.g. IFNγ or TNFα) (14, 20). During monoclonal infections characteristic of many experimental models, intraspecies priority effects from *S*.Tm colonizing a given site within the host reduce the chances of colonization by pathogen cells that are arriving later (16). However, gut colonization is rarely a monoclonal process. This has been demonstrated in longitudinal clinical studies monitoring fecal *Enterobacteriaceae* populations, but also shown in swine where different strains of *Salmonella* colonize the lymph nodes compared to the gut lumen (21-24). Based on these considerations, we hypothesized that repeated re-seeding and re-infection events during persistent infection may promote plasmid spread.

Finally, we reasoned that certain resistance plasmids should themselves expedite reseeding-transfer-invasion cycles within the infected host. In particular, this should pertain to plasmid-encoded resistance genes encoding enzymes degrading antibiotics with an extra-cytoplasmic target (e.g. beta-lactamases). Among these, the extended spectrum beta-lactamases (ESBL) capable of hydrolyzing a wide variety of beta-lactam antibiotics including ampicillin and cephalosporins, are of particular concern (25). These periplasmic enzymes may facilitate reseeding of sensitive recipients, as

resistance plasmids encoding beta-lactamases are able to reach high density under beta-lactam
treatment, and clear the local environment from antibiotics, as shown *in vitro* (26). However, it had
remained unclear if such donor-mediated reduction of local beta-lactam concentrations would
suffice for promoting survival of susceptible recipients *in vivo*, fostering plasmid transfer and
subsequent reservoir formation in the host's tissue.
To address this, we quantitatively assessed the link between within-host plasmid transfer,
invasion-reservoir formation-reseeding cycles and plasmid accumulation in the host's tissues using
S.Tm mouse models for both antibiotic persistence and chronic infection. To study these processes in
a proof of concept approach, we chose donor-recipient-plasmid combinations featuring a high
conjugation efficiency and relieved potential restrictions to conjugation or reservoir formation in the
mouse gut like colonization resistance conferred by the microbiota, and priority effects of strains that
already colonize tissue reservoirs. Thereby, we could demonstrate expansion of the plasmid diversity
in the gut tissue reservoir, a role for the chronically infecting *S*.Tm subpopulation and the effect of
gut luminal antibiotic degradation by beta-lactamase expressing donor strains.
118
119 **Results:**
120
121 **Plasmids that enter *S*.Tm in the gut lumen can be stored in intestinal tissue reservoirs**
122
To study plasmid transfer within a host and its subsequent storage in the host tissues, we
modified a mouse model from our previous work (2). For most plasmid transfer experiments, we
used the IncI1 plasmid P2 (a close relative of extended-spectrum beta-lactamase (ESBL) plasmids;
also known as pCol1b9), which naturally occurs in *S*.Tm SL1344 and spreads in the gut of antibiotic
pretreated mice and during diet-elicited *Enterobacteriaceae* blooms (2, 27-31).
For all experiments, we used streptomycin resistant derivatives of *S*.Tm ATCC 14028S that
naturally do not contain P2 as recipient strains. We infected 129/SvEv mice intraperitoneally (i.p.)
with recipients to allow the formation of tissue reservoirs of *S*.Tm (**Fig. 1A**; left side, green) (2). The
advantage of this approach is three-fold. First, it allows us to study plasmid transfer and subsequent
tissue storage in models for antibiotic persistence (the focus of (2)) and to extend the work to
persistent infection (this work). This is attributable to the capacity of the *Nramp1*-positive 129/SvEv
mice to limit systemic pathogen growth during chronic infections (14, 32). Second, i.p. infection
bypasses the need for gut colonization, ensuring that plasmid transfer observed later in the
experiment is truly a result of plasmid conjugation into recipients entering from tissue reservoirs into
the gut lumen. Third, the i.p. infection model mimics tissue reservoirs that are established after
systemic spread following gut colonization (and therefore allows us to study the role of persistent
infection in plasmid transfer dynamics) (2, 14, 16, 33). We would like to point out one caveat of this
i.p. model, in that it eliminates the contribution of priority effects attributable to the inoculation
route. As natural infections would occur via the oral colonization route, sites that are normally
occupied by invading *S*.Tm cells (e.g. the mesenteric lymph node; mLN) are less colonized after i.p.
infection. This means that invasion events from the gut luminal side later in the experiment are likely
over-estimated in this model. On the other hand, this provides us with an exquisitely sensitive system
to ask if tissue re-invasion can in principle expand the plasmid reservoirs within a host.
Next, we treated mice with a dose of streptomycin to suppress the microbiota and allow
colonization of P2^{cat} carrying donor strains (Sm^R) introduced orally (**Fig. 1A**, blue). Note that both the
donors and recipient strains used in this experimental model are streptomycin resistant. Therefore,
the systemic infection by the recipients remains unaffected by the antibiotic (= persistent systemic
infection model). We used a derivative of *S*.Tm SL1344 that lacks functional type 3 secretions systems
(TTSS)-1 and -2 (*invG* and *ssaV* mutant; SL1344^{Avir}) (34). This prevented invasion of the *S*.Tm donors
into tissues to exclude tissue-localized plasmid transfer. We used SL1344 derivatives to study

interspecies interactions, rather than isogenic strain interactions that would limit co-existence of
donors and recipients in the gut after reseeding (16, 35). Feces were monitored over time to
investigate plasmid transfer dynamics and mice were euthanized at day 10 post infection to assess
the formation of new intra-cellular plasmid reservoirs by recipients that had obtained a plasmid
(transconjugants; **Fig. 1A**, red).
One day after the addition of donors (SL1344^{Avir} P2^{cat}), recipients started reseeding into the
gut lumen and transconjugants were detected in the feces (**Fig. 1B, Fig. S1A**). By day 10 of the
experiment, transconjugants were detected in the majority of mice (8/13 mice; **Fig. 1B**). Reseeding
and plasmid acquisition depended on the TTSS-2 encoded on SPI-2, as recipients lacking a functional
TTSS-2 (14028S *ssaV* mutant; *S.*Tm^{SPI-2}) failed to reseed frequently, and consequently transconjugants
were rarely observed (1/8 mice; **Fig. 1B. Fig. S1B**). Their failure to establish substantial tissue
reservoir sizes (e.g. in the spleen, liver, or gall bladder) could explain the limited reseeding of *S.*Tm^{SPI-2}
(**Fig. 1C**). These results indicated that bacteria reseeding from tissue reservoirs can gain plasmids in
the gut. In contrast to previous work (2) this occurred without previous selection for persister sub-
populations. This allows us to assess the contribution of growing tissue-lodged pathogen
subpopulations. Importantly, we analyzed here if the gut luminal plasmid was carried back into host
tissues by transconjugants. Indeed, transconjugants were detected in the mLNs of mice where gut
luminal plasmid transfer was observed. Occasionally such transconjugants were also observed in
other tissue reservoirs such as the spleen and liver associated with further systemic spread after
tissue invasion (**Fig. 1C**).
Next, we tested if this process could be generalized to other donors colonizing the gut lumen.
For this, we used *E. coli* 536 carrying P2^{cat} as a donor. This strain is naturally streptomycin resistant
and colonizes the streptomycin-pretreated mouse gut for long periods of time (36, 37). Also in this
case, reseeding of *S.*Tm 14028S recipients occurred followed by the formation of transconjugants in
the majority of the mice (7/12 mice) (**Fig. 1D, Fig. S1C**). Once again, this was associated with the
formation of new plasmid tissue reservoirs by invading transconjugants, primarily in the mLN (**Fig.**
1E). To assess if this experimental model allowed the comparison between both persistent infection
and antibiotic persistence, we performed a control experiment in which we treated mice three times
with ceftriaxone 2 days after i.p. infection to select for persisters (**Fig. S2**), as done previously (2).
After the addition of *E. coli* 536 P2^{cat} donors, *S.*Tm^{WT} recipients began to reseed in some mice.
However, this occurred later and in a smaller number of the analyzed mice, as antibiotic persisters
are much less abundant (and need to re-enter growth) compared to the much larger population of
persistently infecting *S.*Tm^{WT} recipient cells. Regardless, this reseeding was also associated with
plasmid transfer and the occasional formation of tissue reservoirs of transconjugants in the mLN (**Fig.**
S2).
Altogether, this demonstrates that plasmids can subvert both *S.*Tm derived from tissue
lodged populations of antibiotic persisters, or from persistent infections. Cells from either type of
tissue lodged population can re-enter the gut lumen, and serve as recipients in interspecies
interactions. This process can be followed by reinvasion forming new tissue reservoirs, capturing a
record of conjugative plasmids that had previously been present in the gut lumen.
193
194 **Reseeding of the gut lumen, plasmid transfer, and the formation of new reservoirs is limited by the**
195 **gut luminal carrying capacity and the conjugation rate**
196
Next, we examined the dynamics of pathogen-assisted plasmid transfer and storage, and the
possible routes that recipients and transconjugants take to exit and re-enter host tissues in more
detail. First, we explored possible sources of persistently infecting recipients that reseed the gut
lumen by characterizing the localization of *S.*Tm in tissue reservoirs. We infected mice i.p. with *S.*Tm
14028S recipients and euthanized mice 2 days later, while keeping the gut luminal microbiota intact

(at the time point when normally the donors are added). As expected based on previous work (2, 14,
16, 38, 39), S.Tm were found in the spleen, liver, and kidney at high density (**Fig. S3**), which likely
occurred because of blood flow to these organs immediately after i.p. infection of S.Tm (39).
Although bacteria were not detected in the feces (our readout for reseeding), S.Tm was found in the
content of the small intestine (i.e. primarily the jejunum and ileum, but also the duodenum), and this
was correlated with S.Tm in the tissues of these sites (**Fig. S3**). Furthermore, we found high densities
of S.Tm in the lining of the small intestinal tissues that contains the gut-associated tissues such as
Peyer's patches (here called small intestinal gut associated lymphoid tissues; S.I. GALT). The gall
bladder contained some S.Tm, but the low densities suggest that the gall bladder is not the primary
route of reseeding in this model (as suggested in some experimental models (16)). Future work may
address if S.Tm reseeds from the small intestinal tissues after spread from blood or lymphatic supply.
Instead, we focused on the dynamics of reseeding followed by plasmid transfer into this
luminal pathogen population, and asked in which conditions this occurs. We mixed equal ratios of
recipient strains harbouring three different sequence-tagged barcodes at a neutral location in the
chromosome (14028S Sm^R *TAG1-3*; 1:1:1 ratio) and i.p. infected mice (same mice as in **Fig. 1B**). As
donor strains, we used SL1344^{Avir} containing P2 labelled with four additional unique sequence tags
(P2^{TAG4-7}; 1:1:1:1 ratio). All tags can be identified using quantitative PCR (39). At the end of the
experiment, we analyzed the ratio of each of these tags in the recipient population (enriching for
recipients = recipients + transconjugants) by qPCR. To visually correlate reseeding of S.Tm into the
gut lumen with the localization of these tags in tissue reservoirs, we sorted and color-coded the tags
according to the most abundant tag in the feces. While chromosomal tags were evenly distributed in
primary tissue reservoirs (e.g. spleen and liver), the feces were typically dominated by just one of the
three tagged strains (**Fig. 2A, Fig. S4**). Thus, reseeding from these tissue reservoirs is a rare process
that is followed by clonal expansion. We also analyzed the gall bladder as a possible source of
reseeding, and the mLN as a site where new tissue reservoirs are formed after reinvasion of
reseeding S.Tm. Although the chromosomal tags were slightly more skewed in the gall bladder
compared to the spleen and liver, there was no correlation between the rank of the tags found in the
feces and in the gall bladder (**Fig. 2A, Fig. S4A**; e.g. mouse 7 contains different most abundant tags in
the feces and gall bladder). Interestingly, the mLN and the feces typically shared the same most
abundant chromosomal tag (~10-fold median reduction of the proportion of the most and second
most abundant tag in the feces; **Fig. 2A**), suggesting that reseeding followed by reinvasion could be
skewing this distribution. Note that the mLN population likely reflects the population that has
invaded into the gut tissue. However, due to unavoidable background contamination from separating
the gut tissue from the gut lumen, we chose to examine the mLN instead. This correlation between
mLN and fecal chromosomal tags was not seen in one mouse (mouse 9 in **Fig. S4**), in which recipients
were not found at high densities in the feces, further supporting this hypothesis. We plotted the
plasmid tags found in the transconjugant population in the same manner (**Fig. 2B**). In mice where
plasmid transfer was detected (population sizes in **Fig. S1A**; raw tag data in **Fig. S4**), almost all
plasmid tags were found, although their distribution was skewed (**Fig. 2B**). Importantly, the plasmid
tags found in the feces were mostly also found in the mLN, implying that plasmids were carried into
the mLN by transconjugants that had formed in the gut lumen (**Fig. 2B**). In addition, the data
indicated that mLN entry of the plasmid-bearing transconjugants occurred more than once in most of
the mice.
Next, we developed a mathematical model (**Supplementary information**) to explore the
relative contribution of reseeding (migration rate; μ) and plasmid transfer (conjugation rate per
mating pair; donor or transconjugant to recipient; γ) to the bacterial dynamics. We fit the model to
the experimental data (evenness of tags in **Fig. 2A-B**; recipient and transconjugant population sizes in
Fig. S1A; **Supplementary information**) to reveal the most likely migration and conjugation rates (**Fig.
2C** in red; μ = 1.78 events per day; γ = 3.16×10^{-11} events per CFU/g feces per day; **Supplementary**

table S2). This identified recipient reseeding as the rate limiting step in our observed plasmid transfer
dynamics, although the conjugation rate influenced the evenness of the plasmid tags and the time
until recipients were converted into transconjugants (**Fig. S5**). In our experimental system, we used
streptomycin to decrease colonization resistance, effectively raising the carrying capacity
(parametrized at 10^9 CFU/g feces for **Fig. 2C, Fig. S5**). However, since conjugation is density
dependent, we reasoned that the level of colonization resistance should influence our results. To test
the range of *Enterobacteriaceae* densities that would allow appreciable densities of transconjugants
(e.g. sufficient to allow invasion into new tissue reservoirs), we fixed the migration and conjugation
rates to the most likely values from **Fig. 2C.** We then varied the carrying capacity as a proxy for
colonization resistance, using the population sizes of recipients and transconjugants at the end of the
experiment (day 10) as a readout. While recipient densities increased proportionally to the carrying
capacity, the transconjugant population size remained below 10^6 CFU/g feces until the carrying
capacity reached 10^7 - 10^8 CFU/g feces (**Fig. 2E;** grey dotted lines). To validate this with our
experimental data, we re-analyzed the experimental data from **Fig. 1B-E** (pooling data from both
*S.*Tm^{Avir} and *E. coli* donors), correlating the donor population size at day 10 of the experiment (as a
proxy for colonization resistance) with the transconjugant populations in both the feces and the mLN
at day 10 (**Fig. S6A-B**). As suggested by our model, transconjugant formation and reinvasion was
more likely in mice that harboured higher gut luminal donor densities, indicating a reduction in
colonization resistance (**Fig. S6A-B**). We confirmed this trend in simulations with both a higher and
lower conjugation rate than in our experimental system (**Fig. S6C-D**). Varying the conjugation rate
influenced the carrying capacity at which transconjugant population sizes reached appreciable
densities by the end of the simulation. This highlights a critical role for colonization resistance in
limiting tissue-reservoir-assisted plasmid spread dynamics, but also for the plasmid transfer rate
defined by the plasmid-strain association.
275
276 **Plasmids can use re-invading *S.*Tm to form a persistent tissue reservoir that survives antibiotics and**
277 **permits spread to further strains in the gut post-antibiotic treatment**
278
We then addressed the implications of newly formed plasmid tissue reservoirs for future
plasmid transfer dynamics in the gut. For this, in the first phase of the experiment, we performed the
same experiment as in **Fig. 1D** (using *S.*Tm 14028S Sm^R as a recipient and *E. coli* 536 P2^{cat} as a donor),
but on day 10 we gave mice three doses of ciprofloxacin instead of stopping the experiment (second
phase of the experiment; **Fig. 3A**). As before, *S.*Tm reseeded from tissue reservoirs, obtained a
plasmid in the gut lumen, and transconjugants carried the plasmid into host tissues (i.e. mLN) before
the ciprofloxacin treatment killed gut luminal populations and left only tissue-associated persisters
(including some of the newly formed *S.*Tm 14028S Sm^R P2^{cat} cells) to survive (**Fig. 3A, Fig. S7A**). We
used ampicillin in the drinking water to suppress reseeding of these tissue-associated reservoirs
before the third phase of the experiment. In this third phase, we introduced a secondary recipient
(SL1344 ΔP2) and monitored plasmid transfer (and thus the formation of secondary transconjugants)
to this strain. Strikingly, secondary transconjugants were formed in most mice (3/5 mice, numbered
1-3; **Fig. 3A, Fig. S7A**; black dots), correlating with both the presence of primary transconjugants in
newly formed tissue reservoirs in the mLN, and reseeding of primary transconjugants (**Fig. 3B; Fig.**
S7B). Importantly, *E. coli* donors were not detected in this experiment after the ciprofloxacin
treatment, suggesting that plasmid transfer was due to reseeding primary transconjugants from their
new reservoirs (**Fig. S7A-B**). A control experiment in which we euthanized a subset of mice at day 15
(when we normally added the secondary recipient) and investigated the tissue reservoirs verified this
(**Fig. S8**).
Next, we tested if multiple plasmids could accumulate in the gut lumen from tissue
reservoirs. We performed the first two phases of the experiment as above (**Fig. 3A-B**), but this time

in the third phase of the experiment, we added a second *E. coli* donor containing an extended-
spectrum beta-lactamase (ESBL) resistance plasmid (pESBL) shown to conjugate efficiently both in
antibiotic pre-treated mice and mice containing a reduced complexity microbiota (2, 40). As in the
previous experiment, P2^{cat} transconjugants were formed in the first phase of the experiment,
followed by the elimination of bacteria from the gut lumen and storage of P2^{cat} transconjugants in
tissue reservoirs in the second phase of the experiment (**Fig. 3C**; **Fig. S7C-D**). Following the
introduction of pESBL donors, pESBL transconjugants were detected in most mice (4/7 mice) as
additional recipients reseeded from their reservoirs (**Fig. 3C; Fig. S7C**). Importantly, in 2 of these 4
mice (labeled "a" and "b", P2^{cat} transconjugants were also detected in the feces, suggesting that P2^{cat}
transconjugants reseeded from their reservoirs (**Fig. 3C; Fig. S7C**). As in the other experiments, pESBL
transconjugants could also be detected in the mLN of one mouse (**Fig. 3D**). These data indicate that
invasive enteropathogens like *S.*Tm might allow different resistance plasmids to accumulate in tissue
reservoirs in the long run. However, it is important to reiterate that invasion into the mLN is likely
influenced by priority effects, making each subsequent invasion less likely. This may explain why both
types of transconjugants were not found in the same mLN in the tested mice (mouse "a" and "b"; **Fig.**
3D). Nevertheless, the accumulation of two plasmids in the gut lumen can be facilitated by tissue
reservoirs.
Altogether, this indicates that tissue reservoirs of *S.*Tm can influence plasmid spread in the
gut both by spreading plasmids to different recipients over time, or accumulating multiple plasmids
over time.
320
321 **ESBL resistance plasmids can facilitate reseeding of *S.*Tm from tissue reservoirs under beta-lactam**
322 **treatment, and promote their spread**
323
Plasmids are selfish genetic elements that can use persistent pathogen cells to form long
term-reservoirs in host tissues. We reasoned that some plasmids (e.g. those encoding for resistances
that can clear antibiotics nearby, such as beta-lactamases) may have another way to promote such
transconjugant-dependent reservoir formation by protecting potential recipients in their
surroundings. The survival of *S.*Tm in the presence of otherwise lethal concentrations of beta-lactam
antibiotics mediated by ampicillin-resistant *E. coli* has been described *in vitro*, based on the
extracellular acting mechanism of beta-lactamases (26). Furthermore, antibiotic treatment (of which
beta-lactams comprise a large fraction) is commonly used in the treatment of bacterial infections in
humans and livestock (25, 41, 42). Therefore, we asked if gut luminal *Enterobacteriaceae* harbouring
plasmids that confer resistance to the antibiotic used could facilitate survival of *S.*Tm recipients long
enough to obtain a plasmid and survive in the gut lumen. As a class of multi-drug resistance plasmids
that has drawn particular attention as a contributor to antibiotic resistance spread, and because of
their extracellular mechanism of action, we chose to study ESBL plasmids under beta-lactam
treatment (25, 43-45).
As previously, we infected *S.*Tm recipients (14028S Sm^R) i.p. into mice. One day post
infection, we introduced *E. coli* orally either bearing pESBL (same strains used in **Fig. 3C-D**; clinical
isolate characterized and sequenced in (2, 40)), or P2^{cat} as a control (isogenic *E coli* strains; the P2^{cat} *E.*
coli was cured of pESBL, followed by *in vitro* conjugation of P2^{cat}), and simultaneously gave the mice
ampicillin in the drinking water (**Fig. 4**). Treating mice with 0.5 g/L ampicillin in the drinking water
corresponds to a daily dose of 2-6g in a 70 kg human, close to the recommended treatment schedule
of oral ampicillin of 500mg every 6 hours (46). As expected, *E. coli* pESBL grew to high densities,
whereas in the control animals inoculated with ampicillin susceptible *E. coli* P2^{cat} (negative control
donors; chloramphenicol resistance), the *E. coli* could not survive (**Fig. S9A**). Over time, *S.*Tm began
reseeding from their tissue reservoirs in both groups of mice (**Fig. S9B**), but could only bloom in the
gut lumen in the presence of pESBL *E. coli* (**Fig. 4A**). This was associated with transfer of the pESBL,

followed by clonal expansion as *S*.Tm became resistant to ampicillin, leading to >99% of the *S*.Tm gut
luminal population obtaining a plasmid (**Fig. 4B**). Importantly, since *S*.Tm is an enteric pathogen that
utilizes inflammation to bloom in the gut (47), using an ELISA for fecal lipocalin-2, inflammation was
detected in mice with pESBL *E. coli*, and reached particularly high levels in mice where reseeding of
S.Tm occurred early during the experiment (i.e. >10^3 lipocalin-2 ng/g feces; **Fig. 4C**). Interestingly,
low-grade inflammation was detected in the absence of gut luminal *S*.Tm (**Fig. 4C**) and could suggest
that systemic *S*.Tm may contribute to low-grade gut inflammation, propagated by gut luminal *E. coli*.
To confirm that reseeding of *S*.Tm was dependent on lowering the concentration of fecal ampicillin
by pESBL *E. coli*, we measured the concentration of fecal ampicillin using mass spectrometry (**Fig.
4D**). In the presence of *E. coli* ESBL, fecal ampicillin concentrations plummeted below the minimum
inhibitory concentration (MIC; detected *in vitro* (2)) of *S*.Tm (**Fig. 4D**), likely explaining the reseeding
dynamics. Thus, donor mediated antibiotic degradation may protect the re-seeding recipients and
thereby promote pESBL transfer. Importantly, as in the previous experiments, *S*.Tm transconjugants
formed new plasmid reservoirs in host tissues, primarily in the mLN (**Fig. 4E**). This indicates that ESBL
plasmids promote their spread in host-associated *Enterobacteriaceae* in two different ways: by
lowering local antibiotic concentrations that ensure survival of potential recipients, and by subverting
tissue-lodged pathogens (persisters and growing populations alike) to form long-term reservoirs
within host tissues.
367
368 ### Discussion:
369
Our results suggest that plasmids can leverage facultative intracellular enteropathogenic
bacteria to promote both the formation of plasmid reservoirs in host tissues, and the spread of
plasmids into and among gut luminal microbes passing through a given host. In our experiments
using *S*.Tm infection in mice, repeated cycles of pathogen invasion and tissue reservoir formation
followed by reseeding the gut lumen allow plasmids to form long-term reservoirs in the host's tissues
(**Fig. S10, steps 1-5**). These reservoirs act as a record of gut luminal horizontal gene transfer, here
shown by conjugation (**Fig. 1-4**). By assessing the genetic element's tissue reservoir formation, our
current study significantly extends the ecological implications of our previous work where we had
focused on persisters, showing that these antibiotic persisters can promote the release of donors
into the gut lumen (2). Here we show that tissue reservoirs can influence plasmid dynamics also
without selection for persister survival in antibiotic treated tissues (**Fig. S10, step 1**). Most likely, it
will therefore happen during persistent infection involving phenotypically susceptible pathogen cells.
Since a natural ecological succession of *Enterobacteriaceae* has been documented in the gut (21, 24),
an overarching reseeding-conjugation-reinvasion cycle could result in the distribution of different
alleles in the various strains colonizing a host. The host tissue reservoirs would dramatically expand
the time scale of such transfer thereby enabling horizontal gene transfer even when the initial donor
and the later recipient have never "co-existed" within the host. This could pertain to resistance genes
or virulence determinants alike. In line with this hypothesis, different strains of commensal and
pathogenic Enterobacteriaceae that show vastly different plasmid, prophage and mutational profiles
are often observed within a given host population or even within the same host (21-23, 48, 49).
Conversely, the same plasmid has been detected in different enterobacterial strains over time within
the same patient (24). The role of the pathogen's reseeding-acquisition-reinvasion cycles in
resistance plasmid transfer may be particularly important in cases where different *S*.Tm are present
within the same host (e.g. the gut lumen vs. the host tissues), as observed in pig farms (23).
Which factors limit plasmid storage and transfer from host-tissue reservoirs? Through
population dynamic analysis, we showed that reseeding of recipient strains from tissue reservoirs is
the rate-limiting step for the formation of new transconjugant reservoirs (**Fig. 2; Fig. S10, step 2a**).
This is in line with our previous work (2) where we suggested that donor-reseeding from persister

reservoirs in host tissues is a rate limiting process, and reinforces the importance of understanding
the mechanisms of reseeding, host tissue invasion and the formation of new tissue reservoirs.
Previous studies have suggested the gall bladder as a route of reseeding for *Salmonella* via bile
secretions (16, 50, 51). In our experimental model, we found that sequence tags in the gall bladder
did not necessarily correlate with those found at highest density in the feces, suggesting, along with
others (16), that additional routes of reseeding may exist. Phagocytic cells such as macrophages and
dendritic cells serve as reservoirs for *S.*Tm (10-13, 20, 52) and can be reactivated by IFNγ depletion
(14). The high density of *S.*Tm in the tissues that line the small intestine including the gut associated
lymphoid tissues (here called S.I. GALT), correlating with *S.*Tm found in the gut lumen of the small
intestine, could suggest that trafficking of *S.*Tm-bearing immune cells to these sites (53) contributes
to gut luminal reseeding, but further work is needed for confirmation.
Interventions such as vaccination have shown success in reducing the formation of tissue
reservoirs and consequently plasmid transfer dynamics associated with reseeding (2, 28, 54). Here,
we showed that once reseeding occurred, the transfer of plasmids followed by reinvasion is likely,
given a sufficiently high carrying capacity (as a proxy for decreased colonization resistance) in the gut
lumen. This is because plasmid transfer is dependent on both the density of donors and recipients.
Invasion may also be density dependent (55); approximately 300 invasion events occur into the mLN
per day initially in streptomycin pretreated mouse models (56) and this likely decreases
proportionally with colonization. In a healthy microbiota community, colonization resistance
prohibits enterobacteriaceal blooms in >90% of the hosts (57, 58) and dramatically reduces the
efficiency of plasmid transfer (29, 30). However, gut perturbations such as diet shifts, inflammation,
antibiotic treatment, or a reduction in microbiota complexity (e.g. dysbiosis) can all lead to increases
in loads of *Enterobacteriaceae* and consequently plasmid transfer (**Fig. S10, step 3**) (2, 29, 30, 40, 47,
59, 60). Therefore, a healthy microbiota is integral in minimizing pathogen evolution in two
complementary ways, by reducing horizontal gene transfer and by reducing the likelihood of re-
invasion and subsequent formation of tissue reservoirs. Further work will be needed to validate how
such perturbations accelerate the dynamics of plasmid spread and reservoir formation.
Lastly, we showed that plasmids encoding secreted antibiotic degrading enzymes can
promote their own spread by an additional mechanism *in vivo*: protecting potential new recipients
sensitive to the antibiotic (**Fig. S10, step 2b**). Reseeding of *S.*Tm, initially susceptible to the beta-
lactam, can occur even in the presence of the antibiotic, dependent on the presence of *E. coli*
producing beta-lactamase (**Fig. 4**). The prevalence of ESBL-producing *E. coli* is rising in recent years,
and ESBL genes are frequently plasmid-encoded (25, 61, 62). The extracellularly acting beta-
lactamases are therefore public goods. *In vitro* this was shown to allow the survival of susceptible
*S.*Tm in the presence of otherwise lethal concentrations of ampicillin (26). In our experiments, we
showed that the same process occurs in the gut lumen, with reseeding *S.*Tm followed by conjugation,
clonal expansion, and tissue reinvasion as a consequence (**Fig. 4**). As one of the prodigal mechanisms
for blooms (of resistant bacteria), antibiotic treatment could serve to amplify reseeding followed by
plasmid exchange and tissue reservoir reformation, given sufficient protection from the antibiotic *in*
trans. This obviously depends on the presence of tissue-invasive enteropathogens in the hosts, which
is quite common in today's farming industry (23, 63), as well as the mechanism of action of the
antibiotic used. Conversely, pathogen intrinsic mechanisms transiently promoting survival may
further prolong recipient survival in an antibiotic loaded gut lumen and thereby further promote
plasmid spread. The transient overexpression of efflux pumps was shown to confer survival long
enough to obtain a plasmid in the presence of tetracycline above the MIC *in vitro* (64). Further work
will be necessary to determine if reseeding can be facilitated in the presence of antibiotics,
depending on the specific resistance genes (other than ESBLs) encoded on plasmids in strains
colonizing the gut lumen.

Overall, here we show that plasmids can subvert tissue-associated pathogen reservoirs to
promote long-term carriage within the host and subsequent spread. There seem to be two types of
tissue reservoirs with partially complementary features (**Fig. S10**). The tissue-associated reservoirs
formed by enteropathogens in chronic (persistent) infections are relatively large and facilitate re-
seeding of the gut lumen and subsequent plasmid transfer at relatively high rates. In contrast,
persistent pathogen cells are less frequent and need to re-start growth before engaging in any
luminal plasmid transfers, likely making them less efficient at luminal re-seeding. However, they
ensure survival of recipients and plasmid reservoir maintenance even in cases of therapy with
antibiotics to which these persisters are genetically susceptible. Although in our experimental model
plasmids can benefit strongly from tissue reservoirs, their actual transfer is dependent on the strain
densities in the gut lumen. Our study provides a proof of concept that tissue reservoirs can act as a
record of evolutionary events such as plasmid transfer in the gut lumen. It is conceivable that
mutational events or transfer of other mobile genetic elements, for example bacteriophages carrying
accessory genes, could also be stored in tissue reservoirs. Regardless, it remains to be seen how
often such cases occur in nature, since our work identifies these processes to be dependent on the
carrying capacity in the gut and conjugation rates of plasmid-strain pairs, and to what extent this
contributes to the dissemination of accessory genes on mobile genetic elements, such as antibiotic
resistance genes.

 <u>Figure captions:</u>

Figure 1. *S*.Tm in tissue reservoirs can reseed the gut lumen, obtain a plasmid from
Enterobacteriaceae in the gut lumen, and form new plasmid-bearing reservoirs. A) Proposed
hypothesis. *S*.Tm recipients (green) establish tissue reservoirs after intraperitoneal injections and
spread to organs, dependent on the Type-three secretions system-2 (TTSS-2; encoded on SPI-2). The
microbiota (grey) provides colonization resistance against colonization of the gut lumen. Donors
(blue) colonize the gut lumen by an oral infection following a streptomycin treatment to suppress the
microbiota. *S*.Tm recipients reseed from their reservoirs and obtain a plasmid from donors forming a
transconjugant (red). These transconjugants then form new tissue reservoirs. **B)** Transconjugants are
formed in the gut lumen. Mice were i.p. infected with 10^3 CFU of a 1:1:1 mix of 14028S Sm^R *TAG1-3*
(*S*.Tm^{WT}; n=13) or 10^3 CFU of 14028S^{SPI-2} SmR (*S*.Tm^{SPI-2}; *ssaV* mutant; n=8). On day 2 post infection,
20 mg of streptomycin was given 4 hours before 5×10^7 CFU of a 1:1:1:1 mix of *S*.Tm^{Avir} (*invG ssaV*
mutant) P2^{TAG4-7}. Selective plating was used to determine the fecal transconjugant loads. **C)** Tissue
reservoirs of *S*.Tm. Organs of mice in **panel B** were analyzed for recipients (green circles) or
transconjugants (red circles). **B-C)** Dotted line indicates the detection limit. A two-tailed Mann
Whitney-U test was used to compare *S*.Tm^{WT} (solid circles) to *S*.Tm^{SPI-2} (hollow circles) in each sample
(p>0.05 not significant (ns), p<0.05 (*), p<0.01 (**), p<0.001 (***), p<0.0001 (****)). **D)** *S*.Tm
recipients can obtain P2^{cat} from *E. coli*. Mice were i.p. infected with 10^3 CFU of 14028S Sm^R *TAG1*
(*S*.Tm^{WT}; n=12). On day 2 post infection, 20 mg of streptomycin was given 4 hours before 5×10^7 CFU
of *E. coli* 536 P2^{cat}. Selective plating and colony colour on MacConkey agar was used to determine the
fecal transconjugant loads. **E)** Tissue reservoirs of *S*.Tm after conjugation with *E. coli*. Organs in **panel**
B were analyzed for recipients (green squares) or transconjugants (red squares). **D-E)** Dotted line
indicates the detection limit. **B,D)** Bars indicate median. **C,E)** Lines indicate median. GB = gall bladder;
mLN = mesenteric lymph node. **B-E)** Fecal populations of donors, recipients, and transconjugants are
presented in **Fig. S1**.

Figure 2. Reseeding from tissue reservoirs is a rate limiting step that suffices in conjugation and
formation of new reservoirs given a sufficient carrying capacity. A-B) Mice in **Fig. 1** where reseeding
of recipients occurred (n=10) were analyzed for the distribution of sequence tags in enrichments of
recipients at day 10 of the experiment (including transconjugants) by qPCR (raw tag data shown in
Fig. S4). The dotted line indicates the conservative detection limit. Lines indicate the median. The
proportions of tags are plotted relative to all 7 tags (3 recipient and 4 plasmid tags). **A)** Recipient tags
and **B)** plasmid tags were sorted according the most abundant recipient or plasmid tag in the feces,
in each mouse. GB = gall bladder; mLN = mesenteric lymph node. **C)** Fitting of simulations to
experimental data. Individual summary statistics used to fit simulations are shown in **Fig. S5**.
Parameters and method for fitting is described in **Supplementary information.** The migration rate of
μ=1.78 events per day and the conjugation rate of γ=3.16×10^{-11} events per CFU/g feces per day
provide the best fit to the experimental data (red indicates rate pairs that fit the experimental data
most often). **D)** The migration and conjugation rates were fixed to the most likely values in **panel C**
and the carrying capacity was varied. Recipient (green) and transconjugant (red) populations at the
end of the simulation (day 10; n=100 simulations per carrying capacity) are shown as the mean with
the range of values. The lowest value was set to 1 CFU/g feces to allow visualization on a log scale.
The grey dotted lines correspond to the carrying capacity at which transconjugants are present at 10^6
CFU/g feces, as a density threshold that we speculate should facilitate some frequency of invasion
into tissue reservoirs.

Figure 3. Newly formed plasmid reservoirs can spread plasmids to recipients after antibiotic
treatment and plasmids can accumulate in the gut dependent on tissue reservoirs. A) *S*.Tm

transconjugants can conjugate to secondary recipients after survival of antibiotic treatment, by
survival in tissue reservoirs. Mice were i.p. infected with 10^3 CFU of 14028S Sm^R *TAG1* (Kan^R; n=5). On
day 2 post infection, 20 mg of streptomycin was given 4 hours before 5x10^7 CFU of *E. coli* 536 P2^{cat}
(CmR). Selective plating and colony colour on MacConkey agar was used to determine the fecal
transconjugant loads (red circles). On day 10, mice were given 3 mg ciprofloxacin for three days in a
row, along with 2 g/l ampicillin in the drinking water from day 11-15. On day 15, a secondary
recipient (SL1344 ΔP2; Amp^R) was added (5x10^7 CFU oral). Selective plating allowed the enumeration
of secondary transconjugants (black circles). Bars indicate the median. Dashed black lines connect
primary or secondary transconjugant populations of the same mice (mice are numbered 1-3; the
same numbering is used in **panel B**). **B)** Tissue reservoirs of *S*.Tm. Organs of mice in **panel A** were
analyzed for primary transconjugants and secondary transconjugants. Lines indicate the median. GB
= gall bladder; mLN = mesenteric lymph node. The numbers correspond to the same mouse
numbering as in **panel A**. **C)** *S*.Tm recipients can allow the formation of multiple transconjugants over
time dependent on multiple reseeding events. Mice (n=7) were i.p. infected with recipients 14028S
Sm^R *TAG1* (Kan^R) and donors *E. coli* 536 P2^{cat} (CmR) as in **panel A**. On day 10, mice were given
antibiotics as in **panel A**. On day 15, a second donor, (*E. coli* pESBL; Amp^R) was added (5x10^7 CFU
oral). Selective plating allowed the enumeration of both P2^{cat} (red squares) and pESBL
transconjugants (grey sqares). Bars indicate the median. **D)** Tissue reservoirs of *S*.Tm. Organs of mice
in **panel C** were analyzed for P2^{cat} and pESBL transconjugants. Lines indicate the median. GB = gall
bladder; mLN = mesenteric lymph node. Dashed lines connect the two transconjugant populations in
the same fecal sample (two mice harbor both types of transconjugants: mouse (a) and (b); these
mouse labels are also shown on the mLN population). **A-D)** Dotted lines indicate the detection limits.
Population sizes of all strains and subpopulations are presented in **Fig. S7**.
536
Figure 4. Reseeding of *S*.Tm followed by reception of a plasmid and storage in new tissue reservoirs
can occur under beta-lactam counter-selection, dependent on *E. coli* conjugative ESBL plasmids. A-
E) Mice were i.p. infected with 10^3 CFU of a 1:1:1:1:1:1:1 mix of 14028S Sm^R *TAG1-7* (Kan^R). On day 1
post infection, mice were given 0.5 g/l ampicillin in the drinking water and either 5x10^7 CFU of *E. coli*
P2^{cat} (Cm^R; black circles; n=10) or *E. coli* pESBL (Amp^R; orange circles; n=10). Dotted lines indicate
detection limits. Lines connect medians on each day. A two-tailed Mann Whitney-U test was used to
compare *S*.Tm^{WT} + *E. coli* P2^{cat} to *S*.Tm^{WT} + *E. coli* P2^{cat} in each sample (p>0.05 not significant (ns),
p<0.05 (*), p<0.01 (**), p<0.001 (***), p<0.0001 (****)). **A)** Reseeding *S*.Tm was analyzed by
selective plating on kanamycin-containing MacConkey agar. **B)** Conjugation was determined by
selective and/or replica plating. **C)** Inflammation was quantified using an ELISA for fecal lipocalin-2.
D) Fecal ampicillin was measured using mass spectrometry. The blue dashed line indicates the
minimum inhibitory concentration of *S*.Tm *in vitro* (2). The red dashed line indicates the
concentration of ampicillin given to the mice in the drinking water. **E)** Organs were analyzed for
transconjugant populations on day 10 of the experiment. GB = gall bladder; mLN = mesenteric lymph
node.
552
553

Materials and methods:

Strains, plasmids, and primers used in this study

Bacterial strains used in this study are derivatives of *S.*Tm SL1344 (65), *S.*Tm ATCC 14028S (66), *E. coli* 536 (36, 67), or *E. coli* ESBL15 (2, 68), and are listed in **Table 1**. For cultivation of bacterial strains, lysogeny broth (LB) medium was used containing the appropriate antibiotics (50 μg/ml streptomycin (AppliChem); 50 μg/ml kanamycin (AppliChem); 15 μg/ml chloramphenicol (AppliChem); 100 μg/ml ampicillin (AppliChem)) at 37°C (or 30°C if containing pCP20 or pCP20-IncI1). Genetic constructs (e.g. gene deletions, neutral isogenic sequence tags, or the P3 plasmid) were introduced into the desired background strain using P22 HT105/1 *int-201* phage transduction (69). Bacterial plasmids used to confer resistance or for construction of strains are listed in **Table 2** and were transformed into cells using electroporation.

E. coli ESBL 15 was cured of its plasmid, pESBL, using plasmid incompatibility. To create an incompatible plasmid, pCP20-IncI1 was cloned using the replicon of pESBL. A PCR amplicon of the replication initiation protein of pESBL with 831 and 420 bp upstream and downstream flanking regions, respectively (to ensure the entire replicon was cloned), was cloned into pCP20 digested with *Pst*I and *Eco*RV (removing the ampicillin resistance cassette) using Gibson Assembly (NEB; protocol as described by the manufacturer). The resulting plasmid, pCP20-IncI1, was electroporated into *E. coli* ESBL15 and grown at 30°C under chloramphenicol selection. Loss of resistance of ampicillin (and consequently loss of pESBL) was confirmed by streaking on LB with ampicillin. The resulting clones were restreaked on LB without chloramphenicol grown at 37°C to cure pCP20-IncI1.

Conjugative plasmids were transferred into the desired strain (e.g. P2^{cat} into *E. coli* ΔpESBL and *E. coli* 536) using *in vitro* conjugation assays. In brief, 10^5 CFU from an overnight culture of the donor strain (SL1344 P2^{cat}) was mixed with the desired recipient, allowing conjugation overnight at 37°C on a rotating wheel. Cells were plated on selective MacConkey agar to identify transconjugants by resistance phenotype and/or colour (i.e., *S.*Tm is *lac* negative and thus forms yellow colonies on MacConkey agar while *E. coli* is *lac* positive and forms red colonies).

All strains and plasmids were genotyped prior to use using the primers listed in **Table 3**.

Table 1. Strains used in this study

Strain name	Strain Number	Relevant genotype	Resistance*	Reference
SL1344	SB300	Wild-type	Sm	(65)
ATCC 14028S	14028S	Wild-type	None	(66)
SL1344 P2^{cat}	M995	*cat* on P2	Sm, Cm	(30)
SL1344 ΔP2	M1404	P2 cured	Sm	(30)
E. coli 536	Z2366	Wild-type	Sm	(36, 67)
E. coli 536 P2^{cat}	Z2124	*cat* on P2	Sm, Cm	This study
SL1344 ΔP2 pM975	Z2287	P2 cured; *bla*	Sm, Amp	This study
E. coli ESBL15	Z2115	*CTX-M1* on pESBL	Amp	(2, 68)
14028S Sm^R	T186	*strAB* on P3	Sm	This study
14028S Sm^R *TAG1*	Z2279	*WITS1-aphT*	Sm, Kan	This study
14028S Sm^R *TAG2*	Z2281	*WITS2-aphT*	Sm, Kan	This study
14028S Sm^R *TAG3*	Z2283	*WITS11-aphT*	Sm, Kan	This study
14028S Sm^R *TAG4*	T270	*WITS13-aphT*	Sm, Kan	This study
14028S Sm^R *TAG5*	T272	*WITS17-aphT*	Sm, Kan	This study
14028S Sm^R *TAG6*	T274	*WITS19-aphT*	Sm, Kan	This study
14028S Sm^R *TAG7*	T276	*WITS21-aphT*	Sm, Kan	This study
14028S^{SPI-2} Sm^R	T284	*ssaV::aphT*	Sm, Kan	This study
SL1344^{Avir} P2^{TAG4}	Z2292	*WITS13-cat* on P2; Δ*invG* Δ*ssaV*	Sm, Cm	This study

SL1344^{Avir} P2^{TAG5}	Z2293	*WITS17-cat* on P2; Δ*invG* Δ*ssaV*	Sm, Cm	This study
SL1344^{Avir} P2^{TAG6}	Z2294	*WITS19-cat* on P2; Δ*invG* Δ*ssaV*	Sm, Cm	This study
SL1344^{Avir} P2^{TAG7}	Z2295	*WITS21-cat* on P2; Δ*invG* Δ*ssaV*	Sm, Cm	This study
E. coli Δ pESBL	Z2156	pESBL cured	None	This study
E. coli Δ pESBL P2^{cat}	T305	pESBL cured; *cat* on P2	Cm	This study

585 * Relevant resistances only: Sm = ≥50 µg/ml streptomycin; Cm = ≥15 µg/ml chloramphenicol; Kan =
586 ≥50 µg/ml kanamycin; Amp = ≥100 µg/ml ampicillin.

588 **Table 2. Plasmids used in this study**

Plasmid name	Relevant genotype	Resistance	Reference
pM975	*bla*; used to confer ampicillin resistance	Amp	(34)
pCP20	FLP recombinase	Amp, Cm	(70)
pCP20-IncI1 (pZ2496)	FLP-*bla*::IncI1 replicon	Cm	This study
P3 (pRSF1010)	*strAB*	Sm	(49)
P2 (pCol1b9)	Wild-type	None	(30)
P2^{cat}	*cat*	Cm	(30)
P2^{TAG}	*WITS13, 17, 19, or 21-cat* on P2	Cm	(2)
pESBL (pESBL15)	*CTX-M-1*	Amp	(2, 68)

589
590

591 **Table 3. Primers used in this study**

Primer name	Sequence (5' to 3')	Purpose	Reference
ESBL15_repl_gibs_for	GCC AGT TAA TAG TTT GCG CAA CGT TGT TGC CAT TGC TGC ACT GAG CTA CCA TAG ATG AC	PCR amplification of IncI1 replicon from pEBSL for Gibson assembly	This study
ESBL15_repl_gibs_rev	TAC AAT TAT TCC TTA CTA TGG ACA AAA ACA TCA ATC TGA T GTG GTT TCA GAA CGG TGA G	PCR amplification of IncI1 replicon from pEBSL for Gibson assembly	This study
ESBL15_ori_ver_up	CCA GTT AAT AGT TTG CGC AAC	Validation of pCP20-IncI1 cloning	This study
ESBL15_ori_intl_ rev	CTT TCA GCG CTT TAT AGC G	Validation of pCP20-IncI1 cloning	This study
ESBL15_ori_intl_for	CTG TTC CGA TGA CCA TCT G	Validation of pCP20-IncI1 cloning	This study
ESBL15_ori_ver_dw	CTC CAG TTT AAA TAC AAG ACG	Validation of pCP20-IncI1 cloning	This study
ssaV-137F	GCAGCGTTCCAGGGTATTCC	Verification of Δ*ssaV* in the chromosome	(31)
ssaV+155R	CAGCAAGTTCTTCTCCAGGC	Verification of Δ*ssaV* in the chromosome	(31)
invG-134F	GAAGGCCACGAGAACATCAC	Verification of Δ*invG* in the chromosome	(31)
invG+112R	GCGGCCTGTTGTATTTCCGC	Verification of Δ*invG* in the chromosome	(31)
P3_SmR_for	CTA GTA TGA CGT CTG TCG C	Verification of P3	This study
P3_SmR_rev	CAC GTT TCG CAA CCT GTT C	Verification of P3	This study
WITS1	ACG ACA CCA CTC CAC ACC TA	qPCR for TAG1	(39)
WITS2	ACC CGC AAT ACC AAC AAC TC	qPCR for TAG2	(39)
WITS11	ATC CCA CAC ACT CGA TCT CA	qPCR for TAG11	(39)
WITS13	GCT AAA GAC ACC CCT CAC TCA	qPCR for TAG13	(39)
WITS17	TCA CCA GCC CAC CCC CTC A	qPCR for TAG17	(39)
WITS19	GCA CTA TCC AGC CCC ATA AC	qPCR for TAG19	(39)
WITS21	ACA ACC ACC GAT CAC TCT CC	qPCR for TAG21	(39)
ydgA	GGC TGT CCG CAA TGG GTC	qPCR for all tags	(39)

592
Infection experiments

All animal experiments were performed in 8-12 week-old specified opportunistic pathogen-
free (SOPF) 129/SvEv mice. These mice contain a functional *Nramp1* allele (also known as *Slc11a1*)

and are resistant to *S*.Tm and therefore allow for long-term infections (32, 33). All infection
experiments were approved by the responsible authorities (Tierversuchskommission, Kantonales
Veterinäramt Zürich, licenses 193/2016 and 158/2019). Sample size was not predetermined and mice
were randomly assigned to treatment groups.

For all experiments, overnight cultures of recipient *S*.Tm (14028S Sm^R derivatives) containing
the appropriate antibiotics were washed with sterile PBS 2x before being diluted to 10^4 CFU/ml in
PBS. Mixtures of the appropriate strains (in the case of tagged strains) or single strains were
intraperitoneally injected into mice as a 100 µl volume (~10^3 CFU per mouse). All mice used for
plasmid transfer experiments were housed in individual cages to ensure experimental independence.
For experiments that lasted longer than 10 days, mice were caged in pairs initially, but split to
individual cages for the final 10 days (due to ethical reasons, mice were not individually caged for
longer than 10 days).

On day 2 post infection, mice were given an oral dose of streptomycin (20 mg), as previously
described for the streptomycin pretreated mouse model (60). Overnight cultures of donor strains
(Sm^R; SL1344 or *E. coli* 536 derivatives) containing the appropriate antibiotics were subcultured for 4
hours 1:20 in 2 ml LB without antibiotics, and then washed in PBS. 4 hours after streptomycin
treatment, donors were given orally (~$5x10^7$ CFU per mouse). After 10 days, mice were either
euthanized or given doses of oral ciprofloxacin (3 mg per mouse; ciprofloxacin hydrochloride
monohydrate, Sigma-Aldrich, dissolved in 100 µl sterile dH$_2$O) for three consecutive days, while
ampicillin (2 g/L) was provided in the drinking water on days 11-15 to prevent reseeding before the
third strain was added. After each ciprofloxacin treatment, cages were changed to prevent
reinfection due to coprophagy. On day 15 of the experiment, mice were either euthanized, or given a
third strain (either a second recipient or a second donor; Amp^R). These mice were euthanized on day
22.

For mice used to enrich for persisters in tissue reservoirs (**Fig. S2**), ceftriaxone (1.5 mg
ceftriaxone disodium salt hemi(heptahydrate) dissolved in 100 µl PBS; Sigma-Aldrich) was injected
intraperitoneally for 3 consecutive days. Donors were added after a streptomycin treatment (as
above) on day 6 and again without streptomycin on day 8 to ensure robust colonization. These mice
were euthanized on day 15.

For mice used to investigate reseeding in the presence of ampicillin counterselection
dependent on ESBL-producing *E. coli* (**Fig. 4**), on day 1 post intraperitoneal infection of recipients, the
mice were given 0.5 g/L ampicillin in the drinking water and given ~$5x10^7$ CFU pESBL or P2^{cat} *E. coli*
donors orally (in PBS, following a 4 hours subculture as described above). An ELISA for mouse
lipocalin-2 was performed on feces (protocol according to the manufacturer; R&D Systems kit) to
determine the inflammatory state of the gut.

In all mouse experiments, feces were collected daily into pre-weighed Eppendorf tubes and
homogenized in PBS using a steel ball at 25 Hz for 1 minute. Populations of donors, recipients, and
transconjugants were diluted and enumerated using selective MacConkey agar. Replica plating was
used to accurately determine the ratio of recipients or donors to transconjugants if transconjugants
were on the same order of magnitude as the donors or recipients. When mice were euthanized (at
day 2, 10, 15, or 22; specified in the figure legends), the spleen, liver, gall bladder, and mesenteric
lymph node, were collected and homogenized in PBS. Populations of bacteria were enumerated as
for the feces. For mice euthanized at day 2 post infection (**Fig. S3**), additional organs were collected.
For the duodenum, jejunum, ileum, cecum, and colon, first the content was collected, and then 1cm
of each tissue was opened longitudinally and washed briefly in PBS before homogenization. For the
purposes of this study, the small intestinal and large intestinal gut-associated lymphoid tissue (S.I.
and L.I. GALT) was collected by removing the lining of the tissues (including the fat) for each of the
small intestine and large intestine. For analysis of bacterial loads in the blood, 100 µl of blood was
aspirated from the heart immediately after euthanasia and collected in PBS with 2% BSA and 1 mM

EDTA. Blood samples were also homogenized. Dissection tools were disinfected in 70% EtOH in
between each organ to minimize the chance of cross-contamination between organs.
648
649   **Analysis of population dynamics using neutral sequence tags**
650
Mice were infected with an equal ratio of S.Tm 14028S Sm[R] recipient strains (total ~10^3 CFU)
bearing sequence tags at a neutral locus in the chromosome (39). For analysis of reseeding and
plasmid transfer dynamics in **Fig. 2**, three recipient tagged strain were used (TAG1-3 Kan[R]; 1:1:1
ratio). On day 2 post infection, an equal ratio of donor tagged strains (S.Tm SL1344[Avir] P2[TAG] strains;
TAG4-7 Cm[R]; 1:1:1:1) were given at a total inoculum size of ~$5x10^7$ CFU per mouse orally. The inocula
were enriched in either LB+chloramphenicol (for donors) or LB+kanamycin (for recipients). At the end
of the experiment (i.e., on day 10 post infection), mice were euthanized and
recipient+transconjugant populations were enriched from 100 µl of the feces and organ
homogenates (in parallel to selective plating) in LB supplemented with kanamycin.
Enrichments were concentrated and genomic DNA was extracted using a QIAamp DNA Mini
Kit (Qiagen). qPCR analysis was performed according to temperature conditions as previously
described (39) using qPCR primers specific to the TAG and a universal second primer, *ydgA* (**Table 3**).
The relative proportion was calculated by dividing the DNA copy number (calculated from the C_T
value) of each tag detected, by the sum of all 7 tags in the sample. A dilution standard of purified
chromosomal DNA allowed for a correlation between DNA copy number and C_T value. For each qPCR
run, the detection limit was determined by the C_T value of the most-diluted DNA standard. The least
precise detection limit defines the conservative detection limit plotted on graphs and used for the
mathematical model ($8.9x10^{-5}$). Once the relative proportion was determined, recipient tags were
separated from plasmid tags (however, the proportion remains relative to all tags). This data is
presented in **Fig. S4**. Next, the tags were sorted according to the most abundant in the feces in each
given mouse (separately for recipient and plasmid tags). This ranking was conserved for analysis in
other organs to allow a correlation between the most abundant tag in the organs relative to the
feces. For example, if TAG2 appeared the most abundant in the feces but least abundant in the mLN,
it would ranked and coloured according to the most abundant tag, and it would still appear the same
colour in the mLN, despite it not being the most abundant tag in the mLN. This data is presented in
Fig. 2A-B.
677
678   **Measurement of fecal ampicillin**
679
Detection and absolute quantification of ampicillin was carried out using high-performance
liquid chromatography heated electrospray ionization high-resolution mass spectrometry (HPLC HESI
HRMS). Fecal samples were homogenized in PBS and immediately frozen after collection. Prior to
analysis, samples were thawed and centrifuged at high speed. The supernatants were analyzed on a
Dionex UltiMate 3000 HPLC coupled to a Q Exactive[TM] Hybrid Quadrupole-Orbitrap[TM] mass
spectrometer. Chromatographic separation was obtained on a Phenomenex Kinetex 2.6 µm XB-C18
150 × 4.6 mm column at 30°C. Water (A) and acetonitrile (B), each containing 0.1% formic acid, were
employed as mobile phases. A gradient of total 12.5 min was applied at a flowrate of 800 µL/min,
starting with 3% B for 2 min, 3% to 50% B in 3 min, 50% to 98% B in 5 min, 98% B for 1 min, 98% to
3% B in 1 min and 3% B for 0.5 min. MS-settings: spray voltage (+) 3.5 kV, capillary temperature 320
°C; sheath gas (57.50), auxiliary gas (16.25), sweep gas (3.25); probe heater 462.50°C; S-Lens RF (50),
resolution (70.000); AGC target (3e6), microscans (1), maximum IT 200 ms, scan range 250–750 *m/z*.
For quantification, an ampicillin standard curve was recorded using H_2O diluted concentrations of 10
ng/mL, 100 ng/mL, 500 ng/mL, 1 µg/mL, 10 µg/mL and 100 µg/mL from a 1 mg/mL ampicillin stock
solution. Samples were analyzed by retention time of ampicillin and the respective ion adduct [M+H]$^+$

= 350.1169 *m/z* with a mass tolerance of 5 ppm. 15 µL of each sample were injected and ampicillin
concentrations were calculated using the Thermo Xcalibur 2.2 Quan Browser software. The
concentration given in µg/ml of supernatant was normalized to the weight of feces collected and
converted to µg/ml fecal content using the average density of feces (71).
699
700 **Statistical analysis**
701
Statistical tests on experimental data were performed using GraphPad Prism 8 for Windows.
The test used is described in the figure caption. The mathematical model was fit to the experimental
measurements using an Approximate Bayesian Computation (ABC) approach (72). For the fit, we
considered three sets of summary statistics: the skew of the plasmid and chromosomal tag
distributions, the total size of the transconjugant and recipient population on day 10, and the time at
which the transconjugant and recipient populations first exceeded 10^6 CFU/g feces. A simulation was
called "successful" if all summary statistics were within three standard deviations of the
experimentally observed mean of these statistics. Migration and conjugation rates were varied on a
grid, all other parameters were kept fixed (**Supplementary table S1**). All R-code needed to simulate
the stochastic model, estimate the most likely parameters from the experimental data, and plot the
results, is included in the Github repository https://github.com/JSHuisman/Recorder .
713

714 <u>**Acknowledgements:**</u>
715
We would like to thank the staff at RCHCI and EPIC animal facilities, and members of the
Hardt, Sunagawa and Slack labs for helpful discussions. This work has been supported by grants from
the Swiss National Science Foundation (NRP 72 407240_167121, 310030B_173338 and
310030_192567), the Gebert Rüf Foundation (GRS-060/18) and the Monique Dornonville de la Cour
Foundation to WDH. MD is funded by an SNF professorship grant (PP00PP_176954) and a BRCCH
multi-investigator grant, JSH is funded by NRP72 grant 407240-167121 from the Swiss National
Science Foundation, and EB acknowledges a Boehringer Ingelheim Fonds PhD fellowship.

723
724
725 <u>**Contributions:**</u>
726
EB and WDH conceived the project and designed the experiments. EB, JAH, and EG carried
out the mouse experiments and analyzed data. JSH designed and performed the mathematical
modeling. RRR and SB supervised the mathematical modelling. AOB and JP measured fecal ampicillin
using mass spectrometry. EB wrote the initial draft of the manuscript. EB, MD, and WDH reviewed
and edited the manuscript to its final form. All authors read, commented on, and approved this
manuscript.

 Supplementary information:

Description of the mathematical model

We model the reseeding of Salmonella recipients from systemic sites into the gut, and the
subsequent creation of transconjugants through plasmid transfer from donors. We follow the
dynamics of Donor (D), Recipient (R), and Transconjugant (T) populations in the gut. These
populations are further divided into isogenic subpopulations based on their sequence tags. The
donors D_j carry a plasmid tag $j \in 1 \ldots N_j$, where N_j is the number of plasmid tags (N_j = 4). Recipients
R_i carry a chromosomal tag $i \in 1 \ldots N_i$, where N_i is the total number of chromosomal tags (N_i = 3).
Transconjugants T_{ij} carry both a chromosomal and a plasmid tag.

To describe the dynamics of these populations, we make several assumptions. Recipients are
introduced into the gut at an overall rate μ. Here we assume this rate is constant, although it likely
increases slightly during the experiment as a function of the recipient population size at systemic
sites. We assume the populations of recipients and donors in the gut are well-mixed, and plasmid
transfer is described by mass-action kinetics. Plasmids are thus transferred between donors and
recipients at a constant per-contact rate γ_D. Once transconjugants are formed, these can also transfer
their plasmid at a constant per-contact rate γ_T. All plasmids are isogenic; they transfer from donors or
transconjugants at the same rate and can be transferred only to naïve recipients.

Resources are limited in the gut, so we model bacterial population growth as a logistic
function of the population size. We assume the donor population is limited by a different resource
than the recipient and transconjugant populations, since they are not isogenic (35). The populations
thus reach separate carrying capacities, denoted by K_D and K_{RT} respectively. Otherwise, the growth
dynamics are assumed population independent: the bacteria are born at a birth-rate r, are cleared at
a clearance rate c (this includes both death and efflux from the gut), and similar to (2) we
parametrize this with a residual birth-rate at carrying capacity $r_K = c$ (to ensure some continued
population turnover even at high densities).

We simulate the population dynamics stochastically using the tau-leaping method from the R
package *adaptivetau* (73). The corresponding deterministic equations (in the limit of large numbers)
are:

$$\dot{D_j} = (r + r_K)D_j - D_j\left(c + \mathrm{r}\frac{\sum_j D_j}{K_D}\right)\#(1)$$

$$\dot{R_i} = (r + r_K)R_i + \frac{1}{N_i}\mu - \sum_j\left(\gamma_D D_j + \gamma_T \sum_i T_{ij}\right)R_i - R_i\left(c + \mathrm{r}\frac{\sum_i(R_i + \sum_j T_{ij})}{K_{RT}}\right)\#(2)$$

$$\dot{T_{ij}} = (r + r_K)T_{ij} + \left(\gamma_D D_j + \gamma_T \sum_i T_{ij}\right)R_i - T_{ij}\left(c + \mathrm{r}\frac{\sum_i(R_i + \sum_j T_{ij})}{K_{RT}}\right)\#(3)$$

Parametrizing the mathematical model

In the main simulations, we varied the migration rate μ and the conjugation rates $\gamma_{D/T}$ while the other
parameters were kept fixed (summarized in **Table S1**). The population-independent parameters
describing the bacterial growth dynamics (*r, c, r_K*), as well as the carrying capacities $K_{D/RT}$ and
inoculum density D_0, were set to the values from (2) . There they were parametrized based on
experimental data from the same mouse model system (28, 60). The number of distinguishable
chromosomal and plasmid tag populations match those used in this paper. We fixed the conjugation

rates from donors and transconjugants to the same value, and the carrying capacities of donors and
recipients+transconjugants were set to the same value.

Table S1. Parameter values used in the simulations

Parameter	Function	Value	Units
r	Birth-rate	$44\ln(2)$	Per day
c	Clearance-rate	$4\ln(2)$	Per day
r_K	Residual birth-rate at carrying capacity	$4\ln(2)$	Per day
$K = K_{D/RT}$	Carrying capacity	10^9	CFU/g feces
D_0	Size of the donor inoculum	10^7	CFU/g feces
N_i	Number of distinguishable chromosomal tags	3	Dimensionless
N_j	Number of distinguishable plasmid tags	4	Dimensionless
μ	Reseeding rate of recipients from systemic sites	**Uniform on the discrete grid** $10^{-6} - 10^3$ in 0.25 log increments	Per day
$\gamma = \gamma_{D/T}$	The rate of conjugation from donors or transconjugants per mating pair	**Uniform on the discrete grid** $10^{-12} - 10^3$ in 0.25 log increments	Per CFU/g feces per day

Parameter estimation in the mathematical model

We use Approximate Bayesian Computation (ABC) (72) to infer the most likely values of migration
rate μ and conjugation rate γ. We varied both parameters on a grid, running 100 simulations per
parameter set. For each parameter combination, we calculated summary statistics (see below) to
compare the simulations to the experimental data.

i. The Evenness index E of the plasmid and chromosomal tag distributions. E is defined as $1 - g$
where g is the Gini index (2, 74).
ii. The size of the recipient and transconjugant populations on day 10, respectively.
iii. The time at which the recipient and transconjugant populations first exceed 10^6 CFU/g feces,
respectively.

For **Fig. 2C** we calculated the likelihood of the parameter combination as the percentage of
simulations that return all summary statistics within three standard deviations of the experimentally
observed mean of these statistics. The experimental data used to calculate the mean of these
statistics is derived from **Fig. 1B, S1A,** and **Fig. 2A-B** (n=8). Due to the assumptions of our model, we
excluded mice where reseeding did not occur. Therefore, the migration rate calculated here may be
slightly overestimated. However, we do not expect the relative importance of reseeding or
conjugation to change due to the exclusion of these mice.

Table S2. Parameter estimates

Simulation	Maximum likelihood (μ,γ) pair	μ value from the marginal posterior distribution	γ value from the marginal posterior distribution
Main text (**Fig. 2D**)	μ=1.78 (per day) γ=3.16×10^{-11} (per CFU/g feces per day)	7.1 (per day) HPD: [0.03, 31.6]	9.36×10^{-10} (per CFU/g feces per day) HPD: [1×10^{-12}, 5.6×10^{-9}]

*HPD: Highest posterior density interval

Dependence of transconjugant emergence on the carrying capacity in the gut

In a separate analysis, we estimated the evenness of the plasmid and chromosomal tag distributions,
as well as the final recipient and transconjugant population sizes, as a function of the carrying
capacity. Here we fixed the migration rate μ and conjugation rate γ at their most likely values (μ =
1.78 per day, $\gamma = 3.16\times10^{-11}$ per CFU/g feces per day), and varied the carrying capacity K uniformly on
the grid $10^3 - 10^{10}$ in 0.25 log increments.

In addition, we repeated this set of simulations with a conjugation rate that was either 100-fold
higher, or 100-fold lower than the most likely value.

802
803
804
805

Supplementary figure captions

Figure S1. Fecal populations for mice in Figure 1. Donors (blue; Cm^R), recipients (green; Kan^R), and
transconjugants (red; Cm^R and Kan^R) were enumerated by selective and/or replica plating on
MacConkey agar. The black dotted lines represent the detection limits for transconjugants and
donors (and recipients in **panel B**). The green dotted lines represent the detection limits for
recipients. The detection limit is higher for recipients once the density of bacteria reaches >10^8 CFU/g
feces (because of the limit of colonies on a plate for replica plating). Before this happens, recipient
populations can appear below this line (then the black dotted line is the detection limit). Lines of
colour connect medians between days. **A-B)** Mice from **Fig. 1B-C** infected with SL1344^{Avir} P2^{cat} as
donors. **A)** Mice infected with S.Tm^{WT} recipients (n=13). **B)** Mice infected with S.Tm^{SPI-2} recipients
(n=8). **C)** Mice from **Fig. 1D-E** infected with S.Tm^{WT} recipients and *E. coli* 536 P2^{cat} as donors (n=12).

**Figure S2. Control for reseeding and obtaining a plasmid from persisters induced by antibiotic
treatment.** Mice were i.p. infected with 10^3 CFU of 14028S Sm^R *TAG1* (Kan^R; n=7). On day 2 post
infection, doses of ceftriaxone (1.5 mg i.p. per mouse per day) were given for three consecutive days.
On day 6, 20 mg of streptomycin was given 4 hours before 5x10^7 CFU of *E. coli* 536 P2^{cat} (Cm^R). *E. coli*
was given again on day 8 post infection to ensure colonization. **A)** Donors (blue; Cm^R), recipients
(green; Kan^R), and transconjugants (red; Cm^R and Kan^R) were enumerated by selective and/or replica
plating on MacConkey agar. The black dotted lines represent the detection limits for transconjugants
and donors. The green dotted lines represent the detection limits for recipients. The detection limit is
higher for recipients once the density of bacteria reaches >10^8 CFU/g feces (because of the limit of
colonies on a plate for replica plating). Before this happens, recipient populations can appear below
this line (then the black dotted line is the detection limit). Lines of colour connect medians between
days. **B)** Organs of mice in **panel A** were analyzed for population sizes of recipients and
transconjugants. Lines indicate medians. Dotted line indicates the detection limit. GB = gall bladder;
mLN = mesenteric lymph node.

Figure S3. Localization of tissue reservoirs of S.Tm prior to reseeding. Mice (n=7) were i.p. infected
with 10^3 CFU of a 1:1:1:1:1:1:1 mix of 14028S Sm^R *TAG1-7*. At day 2 post infection, S.Tm in the
organs were enumerated. For feces and the intestinal content, pre-weighed tubes were used to
determine S.Tm per g of content. Blood is reported per ml. Intestinal tissues are reported per cm.
The small intestinal (S.I.) and large intestinal (L.I.) gut-associated lymphoid tissues (GALT) contain the
lining of the intestinal tissues, which includes the GALT (but also fat). Here, we called them S.I. and
L.I. GALT, but they can contain more than just this region. Dotted lines indicate the detection limits.
Lines indicate medians.

Figure S4. Raw tag proportions before sorting and re-colouring for mice in Figure 2. Relative tag
proportions detected by qPCR in the inocula (R. ino. = recipient inoculum; D. ino. = donor inoculum),
the feces, and the organs at day 10 of the experiment. All samples are derived from an enrichment
in LB+kanamycin (to enrich for recipients+transconjugants). Each tag is given a unique colour. Mice
are separated by vertical dashed lines. Sample groups are separated by vertical solid lines. Dotted
line indicates the conservative detection limit. GB = gall bladder; mLN = mesenteric lymph node. **A)**
Recipient tags. **B)** Plasmid tags.

**Figure S5. Summary statistics used for fitting the mathematical model to the experimental data in
Fig. 2C. A-F)** The mathematical model was fit to the experimental data using 6 metrics (described in
the **Supplementary information**). The values of all 6 metrics were calculated on the grid of possible
migration and conjugation rate pairs. For each metric, the experimentally determined mean is shown

in red (an arrow and red text indicates the mean value), and values of simulations that are above the experimental mean are indicated in a gradient towards white, whereas values that are below the experimental mean are indicated in a gradient towards black. Grey fill indicates migration and conjugation rate pairs where the given metric cannot be calculated (e.g. evenness of tags cannot be calculated if reseeding of recipients into the feces does not occur; the recipient population never reaches appreciable densities if the conjugation rate is too high). **A)** Evenness of chromosomal tags in the recipient+transconjugant fecal population at day 10 (experimental mean = 0.0166; SD = 0.0508). **B)** Evenness of plasmid tags in the transconjugant fecal population at day 10 (experimental mean = 0.358; SD = 0.282). **C)** Days of the experiment (timeline as in **Fig. 1**) until the recipient population exceeds 10^6 CFU / g feces (experimental mean = 4.875; SD = 1.642). **D)** Days of the experiment (timeline as in **Fig. 1**) until the transconjugant population exceeds 10^6 CFU / g feces (experimental mean = 5.00; SD = 0.926). **E)** Recipient population in the feces at day 10 (experimental mean = 2.9×10^8 CFU / g feces; SD = 3.75×10^8 CFU / g feces). **F)** Transconjugant population in the feces at day 10 (experimental mean = 2.61×10^8 CFU / g feces; SD = 3.45×10^8 CFU / g feces).

Figure S6. Dependence of recipient and transconjugant population sizes on the carrying capacity with a higher or lower conjugation rate compared to Fig. 2D, and correlation with experimental data in Fig. 1. A-B) Transconjugant invasion increases proportional to the carrying capacity, as a proxy for colonization resistance provided by the microbiota. Correlation of the donor population size at day 10 (as a measure for carrying capacity; either SL1344 $S.Tm^{Avir}$ or *E. coli* 536; data from **Fig. S1A, C;** n=25) to the transconjugant population in the feces **(panel A;** data from **Fig. S1A, C)** or mesenteric lymph node (mLN; **panel B**; data from **Fig. 1C, E)**. Dotted lines indicate the detection limits by selective plating. A line of best fit is shown and a linear regression is performed: **A)** R^2 = 0.5604; F test determined non-zero slope (p<0.0001); **B)** R^2 = 0.4913; F test determined non-zero slope (p<0.0001). **C-D)** Simulation data. The migration rate was fixed to the most likely values in **Fig. 2C**, but the conjugation rate was fixed to 3.16×10^{-9}, 100-fold higher than in **Fig. 2D** to test the case of a plasmid with a higher conjugation rate **(panel C),** or fixed to 3.16×10^{-13}, 100-fold lower than in **Fig. 2D** to test the case of a plasmid with a lower conjugation rate **(panel D)**. The carrying capacity was varied. Recipient (green) and transconjugant (red) populations at the end of the simulation (day 10; n=100 simulations per carrying capacity) are shown as the mean with the range of values. The lowest value was set to 1 CFU/g feces to allow visualization on a log scale. The grey dotted lines correspond to the carrying capacity at which transconjugants are present at 10^6 CFU/g feces, as a density threshold that we speculate should facilitate some frequency of invasion into tissue reservoirs.

Figure S7. Additional fecal and organ populations for mice in Figure 3. A-B) Secondary conjugation experiment (circles); mice in **Fig. 3A-B**. Donors (blue; Cm^R), primary recipients (green; Kan^R), primary transconjugants (red; Cm^R and Kan^R), secondary recipients (beige; Amp^R), and secondary transconjugants (black; Amp^R and Cm^R) were enumerated by selective and/or replica plating on MacConkey agar. **B)** Organs of mice in **panel A** were analyzed for population sizes of donors, recipients and transconjugants. **C-D)** Multiple plasmids experiment (squares); mice in **Fig. 3C-D**. P2^{cat} donors (blue; Cm^R), recipients (green; Kan^R), P2^{cat} transconjugants (red; Cm^R and Kan^R), pESBL donors (purple; Amp^R), and pESBL transconjugants (grey; Amp^R and Kan^R) were enumerated by selective and/or replica plating on MacConkey agar. **D)** Organs of mice in **panel A** were analyzed for population sizes of donors, recipients and transconjugants. **A,C)** The black dotted lines represent the detection limits for transconjugants and donors. The green dotted lines represent the detection limits for recipients. The detection limit is higher for recipients once the density of bacteria reaches >10^8 CFU/g feces (because of the limit of colonies on a plate for replica plating). Before this happens, recipient populations can appear below this line (then the black dotted line is the detection limit).

903 Lines of colour connect medians between days. **B,D)** Lines indicate medians. Dotted line indicates the
904 detection limit. GB = gall bladder; mLN = mesenteric lymph node.
905
Figure S8. Control that *E. coli* donors are not detected in tissue reservoirs after antibiotic
treatment. Mice (n=5) were i.p. infected with 10^3 CFU of 14028S Sm^R *TAG1* recipients (Kan^R) and
5x10^7 CFU of *E. coli* 536 P2^{cat} donors (CmR) as in **Fig. 3**. Selective plating and colony colour on
MacConkey agar was used to determine population sizes of donors (blue), recipients (green), and
transconjugants (red; Cm^R and Kan^R). On day 10, mice were given 3 mg ciprofloxacin for three days in
a row, along with 2 g/l ampicillin in the drinking water from day 11-15. **A)** Fecal population sizes. The
black dotted lines represent the detection limits for transconjugants and donors. The green dotted
lines represent the detection limits for recipients. The detection limit is higher for recipients once the
density of bacteria reaches >10^8 CFU/g feces (because of the limit of colonies on a plate for replica
plating). Before this happens, recipient populations can appear below this line (then the black dotted
line is the detection limit). Lines of colour connect medians between days. **B)** Organs of mice in **panel**
A were analyzed for population sizes of donors, recipients and transconjugants. Lines indicate
medians. Dotted line indicates the detection limit. GB = gall bladder; mLN = mesenteric lymph node.
919
Figure S9. *E. coli* population sizes and total *S*.Tm population sizes in tissue reservoirs of mice in Fig.
4. A) *E. coli* P2^{cat} (Cm^R) and *E. coli* pESBL (Amp^R) populations were enumerated by selective plating
and colour on MacConkey agar. Coloured lined connect medians. **B)** Total *S*.Tm populations (Kan^R) in
organs were determined by selective plating on kanamycin MacConkey agar. GB = gall bladder; mLN
= mesenteric lymph node. Lines indicate the media. **A-B)** A two-tailed Mann Whitney-U test was used
to compare *S*.Tm^{WT} + *E. coli* P2^{cat} (black; n=10) to *S*.Tm^{WT} + *E. coli* P2^{cat} (orange; n=10) in each sample
($p>0.05$ not significant (ns), $p<0.05$ (*), $p<0.0001$ (****)). Dotted lines indicate the detection limits.
927
Figure S10. Working model for how the infection cycle of *S*.Tm may promote the spread and
accumulation of plasmids within a host. Invasive *S*.Tm (green for recipients, red for transconjugants)
can establish reservoirs within host tissues, storing their current plasmid complement in these
reservoirs (n plasmids). A sub-population of these cells can survive antibiotic treatment as persisters
(smaller green or red cells), that can regrow after the withdrawal of antibiotics. In extension to
previous work (2), here we show that reservoirs of *S*.Tm that chronically infect the host can reseed
from host tissues, even in the absence of selection for persisters (**step 1**). Exiting the host tissues into
the gut lumen is a rate-limiting process (**step 2a**; (2)). This process can be facilitated by
Enterobacteriaceae donors (blue; plasmid shown in black) that produce extracellularly acting
plasmid-encoded antibiotic resistance genes (such as beta-lactamases; blue colour radiating from the
cell indicates local degradation of beta-lactam antibiotics by beta-lactamases), that can allow the
local survival of reseeding recipients under antibiotic treatment (**step 2b**). Once both donors and
recipients co-occur in the gut lumen at sufficient density, plasmid transfer proceeds (**step 3**; (2, 27-
30, 40)). When colonization resistance is relaxed, invasive transconjugants (red) can reach sufficient
densities to invade back into host tissues and form new reservoirs that now contain n+1 plasmids
(**step 4**). In the tissue, transconjugant subpopulations can also survive antibiotic therapy as persisters
(smaller red cells). This process is repeatable in certain conditions. Either transconjugants can reseed
to donate the plasmid to further gut luminal recipients (**step 5a**), or recipients can reseed to receive
an additional plasmids (**step 5b**). Novel steps of this process, demonstrated here using the
streptomycin treatment mouse model, are indicated with bold numbers, whereas steps that have
been previously shown are indicated with non-bold numbers (and key references are listed in the
caption).
950

References:

1. G. D. Wright, The antibiotic resistome: the nexus of chemical and genetic diversity. *Nat Rev Microbiol* **5**, 175-186 (2007).
2. E. Bakkeren *et al.*, Salmonella persisters promote the spread of antibiotic resistance plasmids in the gut. *Nature* **573**, 276-280 (2019).
3. N. Q. Balaban *et al.*, Definitions and guidelines for research on antibiotic persistence. *Nat Rev Microbiol* 10.1038/s41579-019-0196-3 (2019).
4. B. Gollan, G. Grabe, C. Michaux, S. Helaine, Bacterial Persisters and Infection: Past, Present, and Progressing. *Annu Rev Microbiol* **73**, 359-385 (2019).
5. I. Levin-Reisman *et al.*, Antibiotic tolerance facilitates the evolution of resistance. *Science* **355**, 826-830 (2017).
6. J. Liu, O. Gefen, I. Ronin, M. Bar-Meir, N. Q. Balaban, Effect of tolerance on the evolution of antibiotic resistance under drug combinations. *Science* **367**, 200-204 (2020).
7. E. M. Windels *et al.*, Bacterial persistence promotes the evolution of antibiotic resistance by increasing survival and mutation rates. *ISME J* **13**, 1239-1251 (2019).
8. E. Bakkeren, M. Diard, W. D. Hardt, Evolutionary causes and consequences of bacterial antibiotic persistence. *Nat Rev Microbiol* 10.1038/s41579-020-0378-z (2020).
9. M. Diard *et al.*, Antibiotic treatment selects for cooperative virulence of Salmonella typhimurium. *Curr Biol* **24**, 2000-2005 (2014).
10. S. Helaine *et al.*, Internalization of Salmonella by macrophages induces formation of nonreplicating persisters. *Science* **343**, 204-208 (2014).
11. D. A. C. Stapels *et al.*, Salmonella persisters undermine host immune defenses during antibiotic treatment. *Science* **362**, 1156-1160 (2018).
12. P. Kaiser *et al.*, Cecum lymph node dendritic cells harbor slow-growing bacteria phenotypically tolerant to antibiotic treatment. *PLoS Biol* **12**, e1001793 (2014).
13. B. Claudi *et al.*, Phenotypic variation of Salmonella in host tissues delays eradication by antimicrobial chemotherapy. *Cell* **158**, 722-733 (2014).
14. D. M. Monack, D. M. Bouley, S. Falkow, Salmonella typhimurium persists within macrophages in the mesenteric lymph nodes of chronically infected Nramp1+/+ mice and can be reactivated by IFNgamma neutralization. *J Exp Med* **199**, 231-241 (2004).
15. T. D. Lawley *et al.*, Host transmission of Salmonella enterica serovar Typhimurium is controlled by virulence factors and indigenous intestinal microbiota. *Infect Immun* **76**, 403-416 (2008).
16. L. H. Lam, D. M. Monack, Intraspecies competition for niches in the distal gut dictate transmission during persistent Salmonella infection. *PLoS Pathog* **10**, e1004527 (2014).
17. S. S. Grant, D. T. Hung, Persistent bacterial infections, antibiotic tolerance, and the oxidative stress response. *Virulence* **4**, 273-283 (2013).
18. K. Endt *et al.*, Peroral ciprofloxacin therapy impairs the generation of a protective immune response in a mouse model for Salmonella enterica serovar Typhimurium diarrhea, while parenteral ceftriaxone therapy does not. *Antimicrob Agents Chemother* **56**, 2295-2304 (2012).
19. I. A. Onwuezobe, P. O. Oshun, C. C. Odigwe, Antimicrobials for treating symptomatic non-typhoidal Salmonella infection. *Cochrane Database Syst Rev* **11**, CD001167 (2012).
20. T. H. M. Pham *et al.*, Salmonella-Driven Polarization of Granuloma Macrophages Antagonizes TNF-Mediated Pathogen Restriction during Persistent Infection. *Cell Host Microbe* **27**, 54-67 e55 (2020).
21. J. N. V. Martinson *et al.*, Rethinking gut microbiome residency and the Enterobacteriaceae in healthy human adults. *ISME J* **13**, 2306-2318 (2019).
22. O. Tenaillon, D. Skurnik, B. Picard, E. Denamur, The population genetics of commensal Escherichia coli. *Nat Rev Microbiol* **8**, 207-217 (2010).
23. B. San Roman *et al.*, Relationship between Salmonella infection, shedding and serology in fattening pigs in low-moderate prevalence areas. *Zoonoses Public Health* **65**, 481-489 (2018).

24.   R. Leon-Sampedro *et al.*, Pervasive transmission of a carbapenem resistance plasmid in the
gut microbiota of hospitalized patients. *Nat Microbiol* 10.1038/s41564-021-00879-y (2021).

25.   WHO (2017) Global priority list of antibiotic-resistant bacteria to guide research, discovery,
and development of new antibiotics. (WHO (World Health Organization)).

26.   M. H. Perlin *et al.*, Protection of Salmonella by ampicillin-resistant Escherichia coli in the
presence of otherwise lethal drug concentrations. *Proc Biol Sci* **276**, 3759-3768 (2009).

27.   M. Diard *et al.*, Inflammation boosts bacteriophage transfer between Salmonella spp. *Science*
355, 1211-1215 (2017).

28.   K. Moor *et al.*, High-avidity IgA protects the intestine by enchaining growing bacteria. *Nature*
10.1038/nature22058 (2017).

29.   S. Y. Wotzka *et al.*, Escherichia coli limits Salmonella Typhimurium infections after diet shifts
and fat-mediated microbiota perturbation in mice. *Nat Microbiol* 10.1038/s41564-019-0568-
5 (2019).

30.   B. Stecher *et al.*, Gut inflammation can boost horizontal gene transfer between pathogenic
and commensal Enterobacteriaceae. *Proc Natl Acad Sci U S A* **109**, 1269-1274 (2012).

31.   E. Bakkeren *et al.*, Cooperative virulence can emerge via horizontal gene transfer but is
stabilized by transmission. *bioRxiv* 10.1101/2021.02.11.430745, 2021.2002.2011.430745
(2021).

32.   O. Cunrath, D. Bumann, Host resistance factor SLC11A1 restricts Salmonella growth through
magnesium deprivation. *Science* **366**, 995-999 (2019).

33.   B. Stecher *et al.*, Chronic Salmonella enterica serovar Typhimurium-induced colitis and
cholangitis in streptomycin-pretreated Nramp1+/+ mice. *Infect Immun* **74**, 5047-5057 (2006).

34.   S. Hapfelmeier *et al.*, The Salmonella pathogenicity island (SPI)-2 and SPI-1 type III secretion
systems allow Salmonella serovar typhimurium to trigger colitis via MyD88-dependent and
MyD88-independent mechanisms. *J Immunol* **174**, 1675-1685 (2005).

35.   S. M. Lee *et al.*, Bacterial colonization factors control specificity and stability of the gut
microbiota. *Nature* **501**, 426-429 (2013).

36.   E. Brzuszkiewicz *et al.*, How to become a uropathogen: comparative genomic analysis of
extraintestinal pathogenic Escherichia coli strains. *Proc Natl Acad Sci U S A* **103**, 12879-12884
(2006).

37.   M. Ghalayini *et al.*, Long-term evolution of the natural isolate of Escherichia coli 536 in the
mouse gut colonized after maternal transmission reveals convergence in the constitutive
expression of the lactose operon. *Mol Ecol* **28**, 4470-4485 (2019).

38.   M. Hensel *et al.*, Simultaneous identification of bacterial virulence genes by negative
selection. *Science* **269**, 400-403 (1995).

39.   A. J. Grant *et al.*, Modelling within-host spatiotemporal dynamics of invasive bacterial
disease. *PLoS Biol* **6**, e74 (2008).

40.   F. Benz *et al.*, Plasmid- and strain-specific factors drive variation in ESBL-plasmid spread in
vitro and in vivo. *ISME J* 10.1038/s41396-020-00819-4 (2020).

41.   WHO, *Antimicrobial resistance: global report on surveillance* (World Health Organization
(WHO), 2014).

42.   M. Cully, Public health: The politics of antibiotics. *Nature* **509**, S16-17 (2014).

43.   J. Davies, D. Davies, Origins and evolution of antibiotic resistance. *Microbiol Mol Biol Rev* **74**,
417-433 (2010).

44.   A. Carattoli, Resistance plasmid families in Enterobacteriaceae. *Antimicrob Agents*
Chemother **53**, 2227-2238 (2009).

45.   P. Bajaj, N. S. Singh, J. S. Virdi, Escherichia coli beta-Lactamases: What Really Matters. *Front*
Microbiol **7**, 417 (2016).

46.   M. J. Wirfs, *The APRN and PA's Complete Guide to Prescribing Drug Therapy 2020* (Springer
Publishing Company, 2019).

47.   B. Stecher *et al.*, Salmonella enterica serovar typhimurium exploits inflammation to compete
with the intestinal microbiota. *PLoS Biol* **5**, 2177-2189 (2007).

48.   P. T. Desai *et al.*, Evolutionary Genomics of Salmonella enterica Subspecies. *MBio* **4** (2013).

49. C. Kroger *et al.*, The transcriptional landscape and small RNAs of Salmonella enterica serovar
Typhimurium. *Proc Natl Acad Sci U S A* **109**, E1277-1286 (2012).
50. G. Gonzalez-Escobedo, J. M. Marshall, J. S. Gunn, Chronic and acute infection of the gall
bladder by Salmonella Typhi: understanding the carrier state. *Nat Rev Microbiol* **9**, 9-14
(2011).
51. P. Everest, J. Wain, M. Roberts, G. Rook, G. Dougan, The molecular mechanisms of severe
typhoid fever. *Trends Microbiol* **9**, 316-320 (2001).
52. G. Rosenberg *et al.*, Host succinate is an activation signal for Salmonella virulence during
intracellular infection. *Science* **371**, 400-405 (2021).
53. A. Habtezion, L. P. Nguyen, H. Hadeiba, E. C. Butcher, Leukocyte Trafficking to the Small
Intestine and Colon. *Gastroenterology* **150**, 340-354 (2016).
54. K. Moor *et al.*, Peracetic Acid Treatment Generates Potent Inactivated Oral Vaccines from a
Broad Range of Culturable Bacterial Species. *Front Immunol* **7**, 34 (2016).
55. M. Ackermann *et al.*, Self-destructive cooperation mediated by phenotypic noise. *Nature*
454, 987-990 (2008).
56. P. Kaiser, E. Slack, A. J. Grant, W. D. Hardt, R. R. Regoes, Lymph node colonization dynamics
after oral Salmonella Typhimurium infection in mice. *PLoS Pathog* **9**, e1003532 (2013).
57. B. Stecher *et al.*, Like will to like: abundances of closely related species can predict
susceptibility to intestinal colonization by pathogenic and commensal bacteria. *PLoS Pathog*
6, e1000711 (2010).
58. B. Stecher, W. D. Hardt, Mechanisms controlling pathogen colonization of the gut. *Curr Opin*
Microbiol **14**, 82-91 (2011).
59. M. Kreuzer, W.-D. Hardt, How Food Affects Colonization Resistance Against
Enteropathogenic Bacteria. *Annual Review of Microbiology* **74** (2020).
60. M. Barthel *et al.*, Pretreatment of mice with streptomycin provides a *Salmonella* enterica
serovar Typhimurium colitis model that allows analysis of both pathogen and host. *Infect*
Immun **71**, 2839-2858 (2003).
61. E. Temkin, A. Adler, A. Lerner, Y. Carmeli, Carbapenem-resistant Enterobacteriaceae: biology,
epidemiology, and management. *Ann N Y Acad Sci* **1323**, 22-42 (2014).
62. D. Yang, Y. Guo, Z. Zhang, Combined porin loss and extended spectrum beta-lactamase
production is associated with an increasing imipenem minimal inhibitory concentration in
clinical Klebsiella pneumoniae strains. *Curr Microbiol* **58**, 366-370 (2009).
63. T. P. Van Boeckel *et al.*, Global trends in antimicrobial resistance in animals in low- and
middle-income countries. *Science* **365** (2019).
64. S. Nolivos *et al.*, Role of AcrAB-TolC multidrug efflux pump in drug-resistance acquisition by
plasmid transfer. *Science* **364**, 778-782 (2019).
65. S. K. Hoiseth, B. A. Stocker, Aromatic-dependent *Salmonella* typhimurium are non-virulent
and effective as live vaccines. *Nature* **291**, 238-239 (1981).
66. T. Jarvik, C. Smillie, E. A. Groisman, H. Ochman, Short-term signatures of evolutionary change
in the Salmonella enterica serovar typhimurium 14028 genome. *J Bacteriol* **192**, 560-567
(2010).
67. H. Berger, J. Hacker, A. Juarez, C. Hughes, W. Goebel, Cloning of the chromosomal
determinants encoding hemolysin production and mannose-resistant hemagglutination in
Escherichia coli. *J Bacteriol* **152**, 1241-1247 (1982).
68. S. Tschudin-Sutter *et al.*, Prospective Validation of Cessation of Contact Precautions for
Extended-Spectrum beta-Lactamase-Producing Escherichia coli(1). *Emerg Infect Dis* **22**, 1094-
1097 (2016).
69. N. L. Sternberg, R. Maurer, Bacteriophage-mediated generalized transduction in Escherichia
coli and Salmonella typhimurium. *Methods Enzymol* **204**, 18-43 (1991).
70. K. A. Datsenko, B. L. Wanner, One-step inactivation of chromosomal genes in Escherichia coli
K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**, 6640-6645 (2000).
71. D. M. Brown, D. Butler, N. R. Orman, J. W. Davies, Gross solids transport in small diameter
sewers. *Water Science and Technology* **33**, 25-30 (1996).

1110 72. P. Marjoram, J. Molitor, V. Plagnol, S. Tavare, Markov chain Monte Carlo without likelihoods.
1111  *Proc Natl Acad Sci U S A* **100**, 15324-15328 (2003).
1112 73. P. Johnson (2016) adaptivetau: Tau-Leaping Stochastic Simulation. (R package version 2.2.1).
1113 74. L. Maier *et al.*, Granulocytes impose a tight bottleneck upon the gut luminal pathogen
1114  population during Salmonella typhimurium colitis. *PLoS Pathog* **10**, e1004557 (2014).

1115

Bakkeren et al. Figure 1

Bakkeren et al. Figure 2

Time-course of secondary conjugation experiment

A

E. coli donor (oral)

Cipro. (oral)

Secondary *S*.Tm^WT recipient (oral)

Amp. in water

B

Transconjugant tissue reservoirs at Day 22 (Secondary conjugation experiment)

● Primary transconjugants
● Secondary transconjugants

Time-course of multiple plasmids experiment

C

E. coli P2^cat donor (oral)

Cipro. (oral)

E. coli pESBL donor (oral)

Amp. in water

D

Transconjugant tissue reservoirs at Day 22 (Multiple plasmids experiment)

■ P2^cat transconjugants
■ pESBL transconjugants

Bakkeren et al. Figure 3

Bakkeren et al. Figure 4

Bakkeren et al. Figure S1

A

Sm (oral)
+
E. coli 536 donor (oral)

14028S *S*.Tm^WT recipient (i.p.)

Ceft. (i.p.)

E. coli 536 donor (oral)

CFU / g feces

Day of experiment

B

Tissue reservoirs at Day 15
(Ceftriaxone treatment; *E. coli* 536 donor)

CFU / organ (or g feces)

Feces   Spleen   Liver   GB   mLN

Donors (*E. coli* 536)
Recipients (14028S *S*.Tm^WT)
Transconjugants (14028S *S*.Tm^WT)

Bakkeren et al. Figure S2

Bakkeren et al. Figure S3

A

Recipient tags at Day 10

Mouse number

Recipient tag
- Tag 1
- Tag 2
- Tag 3

B

Plasmid tags in transconjugants at Day 10

Mouse number

Plasmid tag
- Tag 4
- Tag 5
- Tag 6
- Tag 7

Bakkeren et al. Figure S4

Bakkeren et al. Figure S5

A Experimental data from
Fig. S1A,C

B Experimental data from
Fig. 1C,E and Fig. S1A,C

C Simulation data
100-fold <u>higher</u> conjugation rate
compared to Fig. 2D

D Simulation data
100-fold <u>lower</u> conjugation rate
compared to Fig. 2D

● Recipients
● Transconjugants

Bakkeren et al. Figure S6

Bakkeren et al. Figure S7

A

Sm (oral)
+
14028S *E. coli*
S.Tm^WT   536
recipient donor
(i.p.)   (oral)

Cipro.
(oral)

Amp. in water

Day of experiment

CFU / g feces

B

Tissue reservoirs at Day 15

CFU / organ (or g feces)

Feces   Spleen   Liver   GB   mLN

Donors (*E. coli* 536)
Recipients (14028S *S*.Tm^WT)
Transconjugants (14028S *S*.Tm^WT)

Bakkeren et al. Figure S8

A *E. coli* colonization

E. coli pESBL or **P2**^{cat} donor (oral)

S.Tm^{WT} recip. (i.p.)

Amp. in water

ns **** **** **** **** **** **** **** ****

E. coli CFU / g feces

Day of experiment

B *S*.Tm tissue reservoirs at Day 10

S.Tm CFU / organ (or g feces)

*   *   *   ns   *

Feces Spleen Liver GB mLN

● *S*.Tm^{WT} + *E. coli* P2^{cat}
● *S*.Tm^{WT} + *E. coli* pESBL

Bakkeren et al. Figure S9

Bakkeren et al. Figure S10