PKR-mediated Stress Response Enhances Dengue and Zika Virus Replication

Taissa Ricciardi-Jorge[1*], Edroaldo Lummertz da Rocha[1], Edgar Gonzalez-Kozlova[1,2], Gabriela Flavia

Rodrigues-Luiz[1], Brian J Ferguson[2], Nerea Irigoyen[2], Daniel Santos Mansur[1*]

[1] Departamento de Microbiologia, Imunologia e Parasitologia, Centro de Ciências Biológicas (CCB),

Universidade Federal de Santa Catarina (UFSC), Florianópolis, SC, Brasil.

[2] Present address, Icahn School of Medicine, New York, USA.

[3] Department of Pathology, University of Cambridge, Tennis Court Road, Cambridge, UK.

* Corresponding authors: daniel.mansur@ufsc.br; t.ricciardi.jorge@gmail.com

ABSTRACT

The mechanisms by which *Flaviviruses* use non-canonical translation to support their

replication in host cells are largely unknown. Here we investigated how the integrated stress response

(ISR), which promotes translational arrest by eIF2α phosphorylation (p–eIF2α), regulates *Flavivirus*

replication. During Dengue virus (DENV) and Zika virus (ZIKV) infection, eIF2α activation peaked at

24 hours post infection and was dependent on PKR but not type I interferon. The ISR is activated

downstream of p-eIF2α during infection with either virus, but translation arrest only occurred following

DENV infection. Despite this difference, both DENV and ZIKV replication was impaired in cells

lacking PKR, independent of IFN-I/NF-kB signaling or cell viability. By using a ZIKV 5'UTR reporter

system as a model, we found that this region of the genome is sufficient to promote an enhancement of

viral mRNA translation in the presence of an active ISR. Together we provide evidence that

Flaviviruses escape ISR translational arrest and co-opt this response to increase viral replication.

INTRODUCTION

The Flavivirus genus is part of the *Flaviviridae* family and contains the causative agents of several human diseases of high global impact, such as: DENV, ZIKV, Japanese encephalitis virus (JEV), yellow fever virus (YFV), West Nile virus (WNV) and others. Flaviviruses are enveloped ssRNA(+)viruses of about 50 nm in diameter with 10-11 kb genome containing a single ORF flanked by two non-coding segments (UTRs) 5´- (capped) and 3'- (non-polyadenylated) [1]. Flavivirus replication is cytoplasmic in strong association with ER, forming viral replication complexes by convolution of ER membrane [2].

Despite the presence of a 5´-cap on its genome, DENV has been shown to use a cap-independent mechanism for protein synthesis under the suppression of cap-binding protein, eIF4E [3]. Using DENV and ZIKV, this mechanism was later shown to be mediated by an internal ribosomal entry site (IRES localized in viral -UTR [4]. A recent report has confirmed the existence of the IRES element [5], indicating, along with previous suggestions [3,6,7] that a cap-dependent translation would be used for viral polyprotein synthesis until the suppression of canonical translation initiation when a switch to cap-independent translation would take place. This alternative cap-independent mechanism, including cell conditions to promote the switch in translation mode, remains to be fully characterized.

A well-known cellular control over the translation machinery is coordinated by four kinases of the integrated stress response (ISR), that act as sensors of different stressors. They are: (i) PKR, activated by dsRNA; (ii) HRI, activated by oxidative stress and/or heme deficiency; (iii) PERK, activated by reticulum stress; and (iv) GCN2, activated by deprivation of amino acids [8–10]. Therefore, ISR is capable of detecting viral infections directly by ligating replication intermediates (dsRNA) or indirectly via structural or metabolic stresses. A shared target of these four kinases is the subunit alpha of the translation factor eIF2 (eIF2α), which is part of the ternary complex (TC, eIF2-GTP-tRNA$_i^{Met}$), necessary for translation initiation. Phosphorylation of eIF2α (p-eIF2α) prevents the exchange of the associated inactive GDT for the active GTP, blocking its recycling for participation in the formation of new 43S complexes, thus inhibiting general cell translation [11,12].

In the present work, we show that PKR is the main kinase leading to eIF2α phosphorylation and cellular protein translation arrest during DENV/ZIKV infection. This event, however, is linked to the establishment of an optimal cellular environment for virus replication. Using ZIKV as a model, we demonstrate that eIF2α phosphorylation requires viral 5´-UTR to promote enhanced viral translation. These results suggest that ISR may provide the conditions necessary for viral alternative translation boosting viral replication.

RESULTS

DENV4 and ZIKV can induce ISR activation during infection

Interferon stimulated genes are central for Flavivirus restriction, therefore we used human lung A549 cells, as they support DENV and ZIKV replication and also produce and respond to type one interferons (IFN-I) [4,13–17]. We evaluated eIF2α phosphorylation during DENV4 and ZIKV infection as a marker of ISR activation by these viruses. By immunoblot analysis, we show that both viruses induce p–eIF2α by 24 hours post infection (h p.i.) (Figure 1A). This time point coincides with PKR activation by autophosphorylation (Thr451), which suggests the participation of this kinase in ISR activation. Quantification by flow cytometry enables us to determine that eIF2α phosphorylation happens after 12 h p.i. reaching a peak at 24 h p.i. (Figure 1B). This time frame is surprisingly late, considering that from at least 12 h p.i. infectious virus particles can already be detected in cell supernatants under these experimental conditions (Supplementary Figure 1).

To further investigate the context of eIF2α phosphorylation, cells were co-stained with a monoclonal antibody directed against the flavivirus envelope protein (Figure 1C-D). At 24 h p.i. DENV4 infected cells exhibited a significant double-positive population, with about 90% of p–eIF2α+ cells being infected. However, during ZIKV infection, there is a lower proportion of double-positive cells, with about 40% of envelope positive cells in the p–eIF2α+ population. These cells may be infected with ZIKV, but they do not yet express enough envelope protein to allow detectable staining. The same profile of co-staining quantified by cytometry could also be visualized by immunofluorescence (Figure 1D).

In summary, although both infections led to different profiles of anti–p–eIF2α and anti–flavivirus co-staining, both viruses promote stress responses within the same time frame.

Figure 1 – DENV4 and ZIKV promote phosphorylation of eIF2α in A549 cells. A549 cells were
infected with DENV4 (MOI 2) or ZIKV (MOI 3) and harvested at the indicated times for analysis. (A)
Immunoblot analysis of cell extracts resolved in denaturing SDS-PAGE. (B) Flow cytometry analysis
for quantification of cells expressing p–eIF2α within the total population (C) Dot plot and Venn diagram
of representative results from co-staining with anti-Flavivirus E protein and anti-p–eIF2α at 24 h p.i.
analysed by flow cytometry and (D) immunofluorescence of infections under the same conditions,
representative images of three independent experiments. In the column chart, the bars represent the
means ± SEM from three independent experiments. Statistical analysis was performed by paired *t* test
comparing infected samples to the respective uninfected control. *: p ≤0.05; **: p ≤0.01; ***: p ≤0.001;
ns / no markup: no statistical difference; h p.i: hours post-infection.

Phosphorylation of eIF2α during DENV4/ZIKV infection is PKR-dependent and IFN-I independent

To investigate which ISR kinase promotes eIF2α phosphorylation in our model, gene deletion by CRISPR-Cas9 was used to produce A549 PKR and IFNAR knockout cell lines. In PKR[-/-] cells (Supplementary Figure 2A) infected with DENV4 or ZIKV p–eIF2α was abrogated indicating that ISR activation promoted by both viruses is mostly dependent on this kinase (Figure 2A,B). Since PKR is an IFN-stimulated gene (ISG), and considering the late ISR activation, the participation of IFN-I signalling in eIF2α phosphorylation was assessed. However, there was no change in the level of p–eIF2α in infected IFNAR[-/-] A549s compared to WT cells (Figure 2A,B). Likewise, experiments using a PKR/IFNAR double-knockout (Supplementary Figure 2B) lineage confirmed the requirement for constitutive PKR expression to drive eIF2α phosphorylation.

The persistence of a small population p–eIF2α+ in the PKR[-/-] cells infected with ZIKV, indicates the activation of other ISR kinase by this virus. To investigate PERK activity, a selective inhibitor, GSK2656157 [18], was used. GSK2656157 treatment did not alter the induction of p–eIF2α in DENV4 infected cells but promoted a small decrease in p–eIF2α population in ZIKV infected cells (Figure 2C). This demonstrates a minor participation of PERK in the ISR activation in the ZIKV model.

In summary, the phosphorylation of eIF2α during infection by DENV4 or ZIKV is mainly PKR dependent and IFN-I independent in this model.

Figure 2 – Phosphorylation of eIF2α during DENV4 or ZIKV infections is PKR-dependent and IFN-independent. (A) Flow cytometry analysis for quantification of cells expressing p–eIF2α within the total population of the indicated lineages of A549 cells infected with DENV4 (MOI 2) at 24 h p.i.; and (B) same experiment using ZIKV (MOI 3) (C) Flow cytometry analysis for quantification of cells expressing p–eIF2α within the total population A549 WT and PKR$^{-/-}$cells infected with DENV4 (MOI 2) or ZIKV (MOI 3) or treated with thapsigargin 2μM for 45 minutes in the presence or absence of the PERK inhibitor GSK2656157 (5 μM) for 24h; and (D) immunoblot analysis of cell extracts from the same experiment resolved in denaturing SDS-PAGE. Representative image of two independent experiments. DKO: double IFNAR$^{-/-}$/PKR$^{-/-}$ knockout. In column charts, bars represent the mean ± SEM from three independent experiments. Statistical analysis was performed by one-way ANOVA, followed by Tukey's test for multiple comparisons.*: $p \leq 0.05$; **: $p \leq 0.01$; ***: $p \leq 0.001$; ns / no markup: no statistical difference.

PKR is not blocked by DENV4 or ZIKV but shows delayed recognition dsRNA

The late PKR activation during infection with both viruses is not explained by the need of type I IFN signaling to boost PKR levels. Therefore, we tested whether PKR activity is blocked by infection at earlier time points. To test this, the dsRNA analogue, poly (I:C), was used to stimulate PKR. Poly (I:C) transfection was performed at 3 h p.i. and cells were analysed at 9 h p.i., that is, before virus-promoted eIF2α phosphorylation (Figure 3A). The comparable levels of p–eIF2α between mock and infected cells stimulated with poly (I:C) suggest that PKR is not blocked by DENV4 or ZIKV in the early stages of infection (Figure 3A). The absence of p–eIF2α in PKR$^{-/-}$ cells demonstrates that eIF2α phosphorylation induced by poly (I:C) is also dependent on this kinase.

If PKR is not blocked in the early stages of infection by DENV or ZIKV, the late ISR activation could be due to the inability of RNA sensors such as PKR to access viral dsRNA, as suggested previously [19–21]. To test this hypothesis, infected cells were incubated in a semisolid medium so viral spread happens from cell to cell by juxtaposition allowing temporal inferences of events based on cell location from the centre of the viral plaque. Immunofluorescence of DENV4 infected cells shows that cells at the edge of the viral plaque do not phosphorylate eIF2α despite the presence of dsRNA (Figure 3B). This result indicates that ISR activation is delayed, occurring in infected cells after the establishment of viral replication complexes.

Figure 3 – PKR is not blocked by DENV4 or ZIKV but shows delayed recognition dsRNA. (A)
Cells were infected with MOCK/DENV4/ZIKV and incubated for 1h30 before wash, then stimulated
with poly (I:C) at 3 hpi with 10 μg/ml for 6 hours. Analysis for quantification of cell population
expressing p–eIF2α after stimulation with poly (I:C) was performed by flow cytometry. Data obtained
from three independent experiments. (B) Immunofluorescence of A549 WT cells infected with DENV4
for 48 h p.i. in semisolid medium. Cells fixed, permeabilized and co-stained with DAPI, anti-dsRNA
and anti-p–eIF2α. Representative image of three independent experiments. In the column chart, bars
represent the means ± SEM. Statistical analysis was performed by one-way ANOVA, followed by
Tukey's test for multiple comparisons. *: p ≤0.05; **: p ≤0.01; ***: p ≤0.001; ns / no markup: no
statistical difference.

DENV4, but not ZIKV, induces translation arrest in infected cells

To investigate the consequences of eIF2α phosphorylation on cellular and viral protein
translation, viral infections were performed in semi-solid medium and a puromycin labelling protocol
was used to assess general translation activity [22]. By immunofluorescence analysis, it was possible to
observe that p–eIF2α in WT lineage promotes cellular translation arrest during DENV4 infection,

evidenced by the absence of puromycin incorporation in the centre of viral plaques (Figure 4A). At the

edges of the plaque, however, anti-puromycin and anti-flavivirus simultaneous staining reinforces data

presented previously (Fig 3B) that translation arrest happens relatively late during DENV4 infection.

In PKR$^{-/-}$ cells, however, co-staining is present throughout the viral plaque, suggesting cell translation

downregulation during DENV4 infection is caused by ISR activation via PKR.

Surprisingly for ZIKV, there is a different profile, with more intense incorporation of

puromycin by infected WT cells than bystander cells (Figure 4B). Bystander cells are here defined as

cells that were in the same environment of infected cells (i.e., same well or organism) but viral protein

was not detected. Anti-puromycin signal is stronger in the perinuclear region which coincides with viral

replication sites (Figure 4C). The phenotype of increased translation activity in infected cells is lost in

PKR$^{-/-}$ cell lines, suggesting that PKR activation allows increased viral translation.

Figure 4 - DENV4, but not ZIKV, induces PKR-dependent translation arrest in infected cells.

Immunofluorescence of A549 WT or PKR$^{-/-}$ cells infected with DENV4 (A) or ZIKV (B) for 48 h p.i. in semisolid medium. Cells labelled with puromycin before fixation and permeabilization. Co-staining with DAPI, anti-puromycin, anti-flavivirus envelope (E) and secondary antibodies. (C) Zoom on WT

cells infected with ZIKV for a better appreciation of the immunostaining pattern. Representative images of three different experiments.

PKR is required for expression of eIF2α-downstream genes and proteins during DENV4/ZIKV infections

Phosphorylation of eIF2α also leads to transcriptional up-regulation of genes from cellular stress resolution pathways [23,24]. To evaluate the impact of PKR deficiency in events downstream of eIF2α phosphorylation, the expression of *DDIT3* (CHOP) and *GADD34* was analysed. Gene transcription was evaluated by qPCR (Figure 5A) and both *DDIT3* and *GADD34* were less expressed in PKR$^{-/-}$ cells during DENV4/ZIKV infection than in WT cells. This downregulation is subtle, probably due to control of these genes by other pathways such as IRF3/7 [25,26] and STAT3-NFkB or NF-YA [27,28]. However, when GADD34 expression is analysed by immunoblot (Figure 5B) or flow cytometry (Figure 5C) the downregulation of this protein follows p–eIF2α profile (Fig. 2A-B). These results are in line with the literature since GADD34 mRNA translation requires p–eIF2α to suppress uORF2 reading, which in turn inhibits the CDS translation [29,30], demonstrating the transcriptional and post-transcriptional cellular control over eIF2α downstream signalling.

In summary, these results show that during infection by DENV4/ZIKV the activation of the stress response is not restricted to phosphorylation of PKR and eIF2α; and reinforces that the activation of this pathway is abolished in the PKR$^{-/-}$ cells.

Figure 5 – PKR is required for expression of eIF2α-downstream genes and proteins during DENV4/ZIKV infections. A549 WT or PKR-/- cells infected with DENV4 (MOI 2) or ZIKV (MOI 3) and harvested at 24 h p.i. for analysis. (A) Quantification by RT-qPCR of DDIT3 and GADD34 gene expression. Relative expression calculated by 2-ΔΔCt methods using mock WT cells as reference. (B) Immunoblot analysis of GADD34 protein expression. Representative image of two independent experiments. (C) Flow cytometry analysis for quantification of cells expressing GADD34 within the total population. In the column charts, bars represent the means ± SEM from three independent

experiments. Statistical analysis was performed by paired *t* test comparing the two cell lineages under the same conditions. *: p ≤0.05; **: p ≤0.01; ***: p ≤0.001; ns / no markup: no statistical difference.

Disruption of the PKR–eIF2α pathway impairs DENV4 and ZIKV replication

Considering the efficacy of p-eIF2α in blocking cellular translation and the dependency of PKR for this phosphorylation, we evaluated the impact of PKR deletion on virus replication. Surprisingly, replication of both viruses was impaired in PKR$^{-/-}$ cells, shown by lower virus titres (Figure 6A) and smaller plaque sizes and number of cells/plaque (Figure 6B-C). Viral genome replication, assessed by qPCR, does not show statistically significant difference, but a trend of lower replication in PKR$^{-/-}$ cells can be seen and may reflect impairment of other viral replication steps (Figure 6D).

The lower virus titres and diminished plaque sizes in PKR$^{-/-}$ cells are particularly counterintuitive since this cell line is defective in activating the whole antiviral pathway coordinated by PKR. This finding indicated the possibility that the PKR deletion has a negative impact on other antiviral pathways and/or these viruses could have co-opted the ISR to promote viral replication.

Figure 6 - Disruption of the PKR–eIF2α pathway impairs DENV4 and ZIKV replication. (A) Viral titration from A549 WT or PKR$^{-/-}$ cells infected with DENV4 (MOI 2) or ZIKV (MOI 3) and harvested at 24 h p.i. for analysis. Plaque size comparison of virus in A549 WT or PKR$^{-/-}$ cells infected with DENV4 and incubated for 8 days or infected with ZIKV and incubated for 5 days: measurements of (B) plaque areas and (C) number of cells per plaque from immunofluorescence. (D) Quantification by RT-

qPCR of DENV4 or ZIKV viral RNA from A549 WT or PKR[-/-] cells infected with DENV4 (MOI 2) or

ZIKV (MOI 3) and harvested at 24 h p.i. for analysis. Relative expression calculated by $2^{-\Delta\Delta Ct}$ method

using mock WT cells as a reference. PFU: plate forming units; FFU: focus forming units. In all charts,

bars represent the means ± SEM from three independent experiments. Statistical analysis was performed

by paired *t* test comparing the two cell lineages under the same conditions. *: p ≤0.05; **: p ≤0.01; ***:

p ≤0.001; ns / no markup: no statistical difference.

PKR deletion does not affect the innate immune response or cell viability

The phenotype of reduced viral replication in PKR[-/-] lineage could be explained by changes

in the cellular innate immune response, as several studies report the role of PKR in sustaining NF-kB

and IFN signalling [31–36] and promoting IFNγ and TNFα expression [37,38]. To investigate the

possibility of alterations in innate immune pathways as a result of PKR deletion, gene and protein

expression of elements of this response was evaluated.

Genes activated by IRF3/7, IFN-I or NF-κB did not show any significant difference in

transcription between WT and PKR[-/-] lineages during virus infections (Figure 7A). Immunoblotting

corroborated this transcriptional analysis, as no differences were observed in STAT1 activation or IFIT1

and ISG15 expression levels between the two cell lines, thus demonstrating virus-induced IFN-I

signalling is not affected by PKR deletion (Figure 7B).

As PKR can also play a role in cell survival during viral infections [39], cell viability was

analysed at 24 h p.i. by cytometry. Also in this assay, no difference between WT and PKR[-/-] lineages

was observed during virus infections (Figure 7C).

Together, this set of results allow us to conclude that the PKR deletion does not significantly

affect the innate immune response or cell survival during DENV4 or ZIKV infection. Therefore,

alterations in these pathways do not explain the lower viral replication seen in the PKR[-/-] cells.

Figure 7 - PKR deletion does not affect the innate immune response or cell viability. A549 WT or
PKR-/- cells infected with DENV4 (MOI 2) or ZIKV (MOI 3) and harvested at 24 h p.i. for analysis.
(A) Quantification by RT-qPCR of IFNB, IFNL, ISG15 and TNFA1 gene expression. Relative
expression calculated by 2-ΔΔCt methods using mock WT cells as a reference. (B) Immunoblot analysis
of cell extracts resolved in denaturing SDS-PAGE. Representative image of two independent
experiments. (C) Flow cytometry analysis for quantification of living cells by staining with Zombie-
NIR viability dye. Mock WT cells set as 100% reference. In the column charts, bars represent the means
± SEM from three independent experiments. Statistical analysis was performed by paired *t* test
comparing the two cell lineages under the same conditions. *: p ≤0.05; **: p ≤0.01; ***: p ≤0.001; ns /
no markup: no statistical difference.

ZIKV co-opts PKR–eIF2α pathway to increase viral translation through its 5′-UTR

ISR control over cell translation can promote the expression of certain mRNAs as a
consequence of their 5′-UTR sequence and/or structure [30,40]. Therefore, we next sought to
investigate whether viral 5′-UTR could subvert the ISR to promote replication. Hence, a translation
activity assay was performed based on the ZIKV genome using a luciferase reporter readout. For this
assay, a reporter plasmid was designed with the first 194 nucleotides of ZIKV, this includes the 5′-UTR
followed by the first 87 nt from the viral polyprotein fused in frame with the firefly luciferase gene
(Figure 8A). A control plasmid for expression of *Renilla* luciferase with the same backbone that the

firefly luciferase reporter was co-transfected for data normalization. Results are presented in Relative

Translation Activity (RTA), given in percentage, using values obtained with WT mock cells as

reference (100%).

Cells were infected with ZIKV for 12 hours before plasmid transfection, then analysed at 36

h p.i. to maximize the time of luciferase expression. The comparison between mock and infected

conditions for each cell line reveals that virus infection promotes a 2-fold increase in translation activity

by WT and IFNAR$^{-/-}$ cells (Figure 8B). In contrast, in PKR$^{-/-}$ and double knockout cells this effect is

suppressed as relative translation activity in these cells, under infected and uninfected conditions, is

similar or lower to WT uninfected cells. These results suggest that the ZIKV 5′-UTR is sufficient to

promote a relative advantage of mRNA translation in the presence of p–eIF2α.

Figure 8 - ZIKV 5′-UTR is sufficient to increase relative reporter translation in PKR–eIF2α

competent cells. (A) Diagram of reporter construct used for the assay. (B) Relative translation of ZIKV

reporter in A549 (WT/PKR$^{-/-}$/IFNAR$^{-/-}$/DKO) cells infected with ZIKV (MOI 3), transfected at 12 h p.i.

and analysed at 36 h p.i. C: first 87 nt of viral capsid protein. In the column chart, the bars represent the

means ± SEM from three independent experiments. *: p ≤0.05; **: p ≤0.01; ***: p ≤0.001; ns / no

markup: no statistical difference.

DISCUSSION

Given the *Flavivirus* genome structure, it is suggested that these viruses would use a cap-

dependent translation mechanism for viral polyprotein synthesis until the suppression of canonical

initiation machinery when a switch to cap-independent translation would take place [3,5]. Cell

conditions to promote the switch in translation mode and the required auxiliary factors (eIFs) for

alternative translation remain to be fully characterized.

In the present work, we investigated how ISR, known to promote translation arrest through

eIF2α phosphorylation, would play a role in this elusive non-canonical translation mechanism employed

by flaviviruses. Using A549 cells we verified that PKR promotes p-eIF2α during DENV4 and ZIKV

infections. In accordance, previous reports have also detected p-eIF2α in A549 cells during DENV

[6,41], ZIKV [42] and JEV [43] infections; as well as studies using other cell lines [44,45]. Further

characterization of p-eIF2α was performed by flow cytometry to allow quantification and co-staining

for other markers. Although DENV4 and ZIKV promote eIF2α phosphorylation within the same time

frame, these viruses displayed different profiles of anti–p–eIF2α and anti–flavivirus co-staining,

indicating distinct dynamics between viral replication and activation of stress response for each virus.

Therefore, cytometry is here proposed as a complementary method to investigate ISR activation, in

addition to traditionally used immunoblot and immunofluorescence, as it can provide more detailed

information.

It is worth noting that a few studies observe PERK/UPR activation during DENV and ZIKV

infections without excluding the participation of other ISR kinases especially regarding eIF2α

phosphorylation [46–50]. In the present model, PERK plays a minor role in ZIKV-induced p-eIF2α.

Here, PKR is clearly accountable for ISR activation during infection with both DENV4 and ZIKV. This

response was shown to be activated from PKR to p-eIF2α-downstream genes in WT lineage but

abolished in PKR$^{-/-}$ lineage. However, other flaviviruses such as JEV and WNV seem to phosphorylate

eIF2α at least partially dependent on PERK [46–50]. For JEV, this could be due to PKR blocking

promoted by viral NS2a [43].

An increased expression of p-eIF2α downstream genes was also found in Huh7 cells infected

with DENV [51]. In this study, single-cell RNA-seq analysis showed upregulation of *DDIT3* and

GADD34 (aka *PPP1R15A*) mRNAs, as well as of other ATF4-downstream genes (*ASNS, CTH,*

HERPUD1, SOD2, TRIB3) [52–56] within the infected population (Supplementary Figure 3A-B). A

second study using the same approach in PBMC [57] revealed an upregulation of *GADD34* mRNA in

DENV-infected monocytes and T cells (Supplementary Figure 3C-D). The upregulation of these

mRNAs suggests ISR activation after DENV infection in these cell models. However, as shown here,

the expression of these genes is also regulated by post-transcriptional mechanisms dependent on the

presence of p-eIF2α.

In our model, a PKR-dependent translation arrest was confirmed in DENV-infected cells as

expected, but curiously not in ZIKV-infected cells. For the latter, increased puromycin incorporation in

perinuclear areas indicates a possible viral escaping mechanism from ISR translation control. This

difference might be related to the distinct dynamics between viral replication and activation of stress

response for each virus seen in the co-staining assay. Therefore, it calls for further investigation using

methods such as live-cell imaging, to track down the sequence of events during infection.

Roth et al, reported a general translation arrest after infection of Huh7 cells with several

Flavivirus species and strains despite the absence of p-eIF2α [7]. This conflicting result could be due

to the different cell lineage used. Although we do not exclude the possibility that other translation

suppression mechanisms, such as RIDD [58], RNase L [59] and 4E-BP dephosphorylation [60,61], can also take place in our model in later times after infection.

Despite the differences in translational activity under activated ISR, both DENV4 and ZIKV presented an approximated 50% less viral replication in PKR$^{-/-}$ cells. This decrease suggests that ISR activation is not essential for viral replication but rather co-opted by both viruses to counter a major cellular antiviral pathway for their own benefit. Indeed, after discarding the hypothesis of PKR deletion disturbing common antiviral pathways in this model (e.g., type I IFN expression, NFκB activation and apoptosis), translation activity assay has shown that ZIKV 5′-UTR promotes a relative advantage of mRNA translation in the presence of p–eIF2α. This finding reveals that ZIKV not only escapes ISR translation arrest, but also uses this response to increase viral replication. Moreover, it also indicates that viral alternative translation does not require eIF2/TC for initiation. The independence of eIF2 for translation initiation under stress conditions has been previously reported within *Flaviviridae* family, since HCV (*Hepacivirus* genus) and *Pestivirus C*(formerly CSFV, *Pestivirus* genus) IRES present similar functioning [62–64].

Here we demonstrate that DENV and ZIKV induce eIF2α phosphorylation in a PKR dependent manner and that its absence is detrimental for optimal virus replication. Using ZIKV as a model we showed that this effect maps to the 5'UTR of the viral genome. Altogether these results suggest that the activation of the PKR/p-eIF2α axis may be necessary for the engagement of the non-canonical translation employed by Flaviviruses.

Supplementary Figure 1 – Viral titration from A549 WT or PKR$^{-/-}$ cells infected with (A) DENV4 (MOI 2) or (B) ZIKV (MOI 3) and harvested at 12 and 24 h p.i. for analysis. PFU: plate forming units; FFU: focus forming units. In all charts, points represent the means ± SEM from three independent experiments.

Supplementary Figure 2 – Characterization of knockout cell lines. (A) Clones of A549 PKR[-/-]:

Immunoblot of unstimulated and stimulated cells with 100 IU/ml of IFN-α2a for 12 hours

demonstrating successful PKR deletion. (B) Clones of A549 IFNAR[-/-]/PKR[-/-] after stimulation with 100

IU/ml of IFN-α2a for 12 hours demonstrating the deletion of PKR on the parental IFNAR[-/-] cells. DKO:

double-knockout.

Supplementary figure 3 – p–eIF2α-downstream genes are upregulated in DENV-infected Huh7

cells and PBMCs. (A) Dataset from [51], of DENV-infected Huh7 cells, analysed by single-cell RNA-

seq. Heatmap representing expression variability of stress-related genes in uninfected, bystander and

infected cells; and (B) violin plot representing the expression of p-eIF2α- and ATF4-downstream genes

in uninfected, bystander and infected cell populations. (C) Dataset from [57], of PBMCs from patients infected with DENV, analysed by single-cell RNA-seq. Dot plot representing the expression of PPP1R15A/GADD34 in different cell population and infection conditions; and (D) violin plot representing the expression of PPP1R15A/GADD34 the whole PBMC population under uninfected, bystander and infected conditions.

MATERIALS AND METHODS

Cells and viruses. VERO-E6 (ATCC® CRL-1586™) A549 (ATCC® CCL-185™) and derived cell lines were grown at 37° C/5% CO2 in DMEM-F12 medium supplemented with 1 U/ml penicillin /streptomycin and 5% fetal bovine serum. C6/36 cells (ATCC® CRL-1660 were grown at 28° C in Leibovitz's L-15 medium supplemented with 1 U/ml penicillin /streptomycin and 5% fetal bovine serum. IFNAR1 knockout cell line has been previously described [65,66]. PKR$^{-/-}$ and DKO (IFNAR$^{-/-}$/PKR$^{-/-}$) were produced with a pair of sgRNA guides (PKR1-FWD: CACCGATTATGAACAGTGTGCATCG; PKR1-REV: AAACCGATGCACACTGTTCATAATC; PKR2-FWD:CACCGAAACAGTTCTTCGTTGCTTA; PKR2-REV: AAACTAAGCAACGAAGAACTGTTTC) cloned into pSpCas9(BB)-2A-Puro (PX459) vector (Addgene) according to developer's protocol [67]. Clonal populations of new cell lines were isolated and tested to confirm gene deletion (Supplementary figure 2). Experiments were carried out with the 1B6 clone of PKR$^{-/-}$ cells and the 2B6 clone of DKO cells.

DENV-4 virus, TVP-360 strain (KUCZERA et al., 2016) and ZIKV, strain BR 2015/15261 (STROTTMANN et al., 2019) were kindly provided by Dr Claudia N. Duarte dos Santos (Fiocruz-PR, Brazil). Viral stocks were produced in C6/36 cells and all viral titrations were performed in VERO cells by plaque-forming units (PFU) or focus-forming units (FFU).

Poly (I:C)Transfection. Cell stimulation with poly (I:C) ("polyinosinic: polycytidylic acid") was performed with Lipofectamine 3000 (Invitrogen) following manufacturer's instructions for A459 cells at the proportion of 1 ug of poly(I:C) in 1 ml of medium in a 12-well plate.

Western-blots. Samples were lysed on ice with RIPA buffer (150 mM NaCl; 50 mM Tris-HCl pH8; 1% NP40; 0.1% SDS; 0.5% sodium deoxycholate) supplemented with cOmplete™ and PhosSTOP inhibitors™ (Roche) according to manufacturer's recommendations. Protein extracts were heat-denatured in the loading buffer and resolved in SDS-PAGE. Antibodies used for immunoblot reactions were: PKR antibody (Santa Cruz Biotechnology, Inc, sc-707, dilution 1:100); phospho-PKR (Thr451) polyclonal antibody, Invitrogen, 44-668G, dilution 1:100); phospho-eIF2α (Ser51) antibody (Cell Signaling Technology, Inc., #9721, dilution 1:1000); GADD34 antibody (Proteintech, 10449-1-AP, dilution 1:500); α-tubulin (DM1A) mAb (Cell Signaling Technology, Inc., #3873, dilution 1:1000);

STAT1 antibody (AbCam, ab3987, dilution 1:1000); phospho-STAT1(Y701) M135 (Abcam, ab29045, dilution 1:1000); IFIT1 polyclonal antibody (Thermo/Pierce, PA3-848, dilution 1:1000); ISG15 mouse mAb (R&D System, MAB4845, dilution 1:1000); anti-mouse IgG, HRP-linked Antibody (Cell Signaling Technology, Inc., #7076, dilution 1:5000); anti-rabbit IgG, HRP-linked Antibody, Cell Signaling Technology, Inc., #7074, dilution 1:5000). The imaging of blots was carried out on the Chemidoc MP (Bio-Rad).

Flow cytometry. Cells were fixed in BD-Phosflow Lyse/Fix buffer. Primary and secondary antibodies were diluted in permeabilization buffer (0.1% saponin; 1% fetal bovine serum, in PBS) for staining. Between each step, samples were washed in blocking buffer (1% fetal bovine serum in PBS). Antibodies and dyes used for flow cytometry were: phospho-eIF2α (Ser51) antibody (Cell Signaling Technology, Inc., #3597, dilution 1:300); human monoclonal antibody DV 18.4 [68](dilution 1:100); GADD34 antibody (Proteintech, 10449-1-AP, dilution 1:250 (IF/FACS); Zombie-NIR fixable viability dye (BioLegend, 423105, dilution 1:750); Goat anti-Rabbit IgG (H+L) Alexa Fluor 647 conjugated (Invitrogen, A-21245, dilution 1:500); Goat anti-Rabbit IgG (H+L) Alexa Fluor 488 conjugated (Invitrogen, A-11008, dilution 1:500); Goat anti-Human IgG (H+L) Alexa Fluor 488 conjugated (Invitrogen, A-11013, dilution 1:500). Data acquisition was performed with FACSVerse™ (BD) and data analysis with FlowJo software (BD).

Immunofluorescence: Cells were grown and infected on 13 mm cover slips in 24-wells plates and incubated in liquid or semisolid medium according to experiment design. Fixation in 3% paraformaldehyde for 20 minutes and permeabilization with Triton X-100 0,5% in PBS for 5 minutes. Primary and secondary antibodies were diluted in blocking buffer (2% bovine serum albumin in PBS) for staining. Between each step, samples were washed in PBS. Antibodies and dyes used for immunofluorescence were: DAPI (Invitrogen); anti-dsRNA MAb J2 (SCICONS, 10010500, dilution 1:200); human monoclonal antibody DV 18.4 [68](dilution 1:100); anti-puromycin 12D10 (Millipore, MABE343, dilution 1:20,000); phospho-eIF2α (Ser51) antibody (Cell Signaling Technology, Inc., #3597, dilution 1:300); Goat anti-Rabbit IgG (H+L) Alexa Fluor 647 conjugated (Invitrogen, A-21245, dilution 1:500); Goat anti-Rabbit IgG (H+L) Alexa Fluor 488 conjugated (Invitrogen, A-11008, dilution 1:500); Goat anti-Human IgG (H+L) Alexa Fluor 488 conjugated (Invitrogen, A-11013, dilution 1:500); Donkey Anti-Mouse IgG H&L Alexa Fluor 647 conjugated (Abcam, ab150107, dilution 1:500). Image acquisition was performed in the confocal microscope Leica DMI6000 B.

RT-qPCR:RNA was extracted and purified with HiYield TotalRNA mini kit (RBC Real Biotech Corporation, YRB50) according to manufacturer's instructions. Reverse transcription reaction was performed with MMLV "in house" kit from 1 μg of total RNA. Quantitative PCRs were performed with GoTaq ® qPCR Master mix (Promega) with the following primer sets: h18S (F: 5'-

TAGAGGGACAAGTGGCGTTC-3', R: 5'-CGCTGAGCCAGTCAGTGT-3'); hIFNL1 (F: 5'-

TTCCAAGCCCACCACAACTG-3', R: 5'-GAGTGACTCTTCCAAGGCGT-3'); hIFNB1 (F: 5'-

AAACTCATGAGCAGTCTGCA-3', R: 5'-AGGAGATCTTCAGTTTCGGAGG-3'); hTNFA (F: 5'-

TCTTCTCGAACCCCGAGTGA-3', R: 5'-CCTCTGATGGCACCACCAG-3'); hISG15 (F: 5'-

TCCTGGTGAGGAATAACAAGGG-3', R: 5'-TCAGCCAGAACAGGTCGTC-3'); hDDIT3 (F: 5'-

GGAGCATCAGTCCCCCACTT-3', R: 5'-TGTGGGATTGAGGGTCACATC-3'); hGADD34 (F: 5'-

CTGGCTGGTGGAAGCAGTAA-3', R: 5'-TATGGGGGATTGCCAGAGGA-3'); DENV4 vRNA (F:

5'-TTGTCCTAATGATGCTGGTCG-3', R: 5'-TCCACCTGAGACTCCTTCCA-3'); ZIKV vRNA (F:

5'-CTGTGGCATGAACCCAATAG-3', R: 5'-ATCCCATAGAGCACCACTCC-3'). Data acquisition

with StepOne Plus real time PCR system (Applied Biosystems) and relative mRNA expression was

calculated by $2^{-\Delta\Delta CT}$ method.

ZIKV Luciferase reporter vectors: The pSGDLuc vector previously described [69], was

used as a template to design the *Renilla* luciferase and the ZIKV firefly luciferase reporters. The *Renilla*

luciferase was amplified using the following oligonucleotides 5´-

TCCGCCCAGTTCCGCCCATTCTCCGC-3´ and 5´-

GCGCTCTAGATTATCTCGAGGTGTAGAAATAC- 3´ and then cloned back into the pSGDLuc

previously digested with *AvrII* and *XbaI* This will create a plasmid only expressing the *Renilla*

luciferase gene under the T7 or SV40 promoter.

The first 194 nucleotides of the ZIKV genome were cloned first in the pSGD plasmid using the set of

primers 5´-GCGCCTCGAGATAAGTTGTTGATCTGTGTGAATCAGAC-3´ and 5´-

GCGCAGATCTGCCCCCAAAGGGGCTCACACGGGCTAC-3´ in order to introduce this fragment

in frame with the 2A-firefly luciferase gene. Using this intermediate plasmid as a template, the ZIKV-

2A-firefly luciferase gene was later amplified with oligonucleotides 5´-

TCCGCCCAGTTCCGCCCATTCTCCGC-3´and 5´-

GATTCACACAGATCAACAACTccctatagtgagtc to introduce the T7 promoter upstream of the ZIKV

coding region; and 5´-GCTTTACTGGGGCTACGATCTTTTGC-3´ and 5´-

gactcactatagggAGTTGTTGATCTGTGTGAATC-3´ to amplify the ZIKV region. This was cloned

back into the pSGD plasmid previously digested with *AvrII* and *BglII* . This will create a plasmid

expressing the ZIKV-2A-firefly luciferase fused gene under the T7 or SV40 promoter.

Luciferase assays: Cells were transfected using Lipofectamine 3000 (Invitrogen) with firefly

and *Renilla Luciferase* plasmids at the proportion of 98 ng and 2 ng respectively for every 2×10^4 cells

in a 96-well plate. For cell lysis and luminescence reaction, Dual-Luciferase® Reporter Assay System

(Promega) was used following the manufacturer's instructions.

Single-cell RNA sequencing analysis: Processed, publicly available single-cell RNA-seq data are available through the GEO accession number GSE110496 and GSE116672. We downloaded processed single-cell data and metadata from the supplementary information from the respective publication [51,57].

Then, we used CellRouter [70] to perform quality control, normalization, analysis and visualize the expression of selected genes with log2 fold changes > 0.15.

AUTHOR CONTRIBUTIONS

Conceptualization, T.R.J. and D.S.M.; Methodology, T.R.J. and D.S.M.; Formal Analysis and Investigation, T.R.J.; *In silico* analysis, E.L.R., E.G.K., G.F.R.L.; Resources, D.S.M., B.F., N.I., Writing-Original Draft, T.R.J. and D.S.M., Writing-Review and editing, T.R.J., B.F., N.I. and D.S.M., Visualization, T.R.J. and D.S.M., Supervision, D.S.M., Project Administration and Funding Acquisition, D.S.M.

DECLARATION OF INTERESTS

The authors declare no competing interests.

ACKNOWLEDGEMENTS

This work was supported by Comissão de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) Computational Biology programme (23038.010048/2013-27), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) (1 311406/2017-3 and 407609/2018-0) and the Academy of Medical Sciences/U.K. (NAF004/1005) to D.S.M., Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) (41907/2017-7) to T.R.J., BBSRC grant BB/S001336/1 and Wellcome Trust grant 201946/Z/16/Z to B.J.F; and an Isaac Newton Trust Grant (18.40r), a Royal Society Research Grant (RGS\R1\191137) and an Isaac Newton Trust/Wellcome Trust ISSF/University of Cambridge Joint Research Grant to NI.

Multi-User Laboratory in Biological Studies (LAMEB) and LCME at UFSC.

REFERENCES

1. Chambers TJ, Hahn CS, Galler R, Rice CM (1990) Flavivirus genome organization, expression, and replication. *Annu Rev Microbiol* **44**: 649–688.
2. Gillespie LK, Hoenen A, Morgan G, Mackenzie JM (2010) The endoplasmic reticulum provides the membrane platform for biogenesis of the flavivirus replication complex. *J Virol* **84**: 10438–10447.
3. Edgil D, Polacek C, Harris E (2006) Dengue virus utilizes a novel strategy for translation initiation when cap-dependent translation is inhibited. *J Virol* **80**: 2976–2986.

4. Song Y, Mugavero J, Stauft CB, Wimmer E (2019) Dengue and Zika Virus 5' Untranslated
Regions Harbor Internal Ribosomal Entry Site Functions. *MBio* **10.**:
5. Fernández-García L, Angulo J, Ramos H, Barrera A, Pino K, Vera-Otarola J, López-Lastra M
(2020) The internal ribosome entry site of the Dengue virus mRNA is active when cap-
dependent translation initiation is inhibited. *J Virol*.
6. Hou J-N, Chen T-H, Chiang Y-H, Peng J-Y, Yang T-H, Cheng C-C, Sofiyatun E, Chiu C-H,
Chiang-Ni C, Chen W-J (2017) PERK Signal-Modulated Protein Translation Promotes the
Survivability of Dengue 2 Virus-Infected Mosquito Cells and Extends Viral Replication. *Viruses*
9.:
7. Roth H, Magg V, Uch F, Mutz P, Klein P, Haneke K, Lohmann V, Bartenschlager R, Fackler
OT, Locker N, et al. (2017) Flavivirus Infection Uncouples Translation Suppression from
Cellular Stress Responses. *MBio* **8.**:
8. Donnelly N, Gorman AM, Gupta S, Samali A (2013) The eIF2α kinases: their structures and
functions. *Cell Mol Life Sci* **70**: 3493–3511.
9. Pakos-Zebrucka K, Koryga I, Mnich K, Ljujic M, Samali A, Gorman AM (2016) The integrated
stress response. *EMBO Rep* **17**: 1374–1395.
10. Wek RC, Jiang H-Y, Anthony TG (2006) Coping with stress: eIF2 kinases and translational
control. *Biochem Soc Trans* **34**: 7–11.
11. Krishnamoorthy T, Pavitt GD, Zhang F (2001) Tight binding of the phosphorylated α subunit of
initiation factor 2 (eIF2α) to the regulatory subunits of guanine nucleotide exchange factor eIF2B
is required for …. *and cellular biology*.
12. Jackson RJ, Hellen CUT, Pestova TV (2010) The mechanism of eukaryotic translation initiation
and principles of its regulation. *Nat Rev Mol Cell Biol* **11**: 113–127.
13. Wang K, Zou C, Wang X, Huang C, Feng T, Pan W, Wu Q, Wang P, Dai J (2018) Interferon-
stimulated TRIM69 interrupts dengue virus replication by ubiquitinating viral nonstructural
protein 3. *PLoS Pathog* **14**: e1007287.
14. Whelan JN, Li Y, Silverman RH, Weiss SR (2019) Zika Virus Production Is Resistant to RNase
L Antiviral Activity. *J Virol* **93.**:
15. Gullberg RC, Steel JJ, Pujari V, Rovnak J, Crick DC, Perera R (2018) Stearoly-CoA desaturase 1
differentiates early and advanced dengue virus infections and determines virus particle
infectivity. *PLoS Pathog* **14**: e1007261.
16. Zhang J, Lan Y, Li MY, Lamers MM, Fusade-Boyer M, Klemm E, Thiele C, Ashour J, Sanyal S
(2018) Flaviviruses Exploit the Lipid Droplet Protein AUP1 to Trigger Lipophagy and Drive
Virus Production. *Cell Host Microbe* **23**: 819–831.e5.
17. Espada CE, da Rocha EL, dos Santos AA, Soares ZG, Malaquias G, Patrício DO, Kozlova EG,
dos Santos PF, Bordignon J, Sanford TJ, et al. (2019) ISG15/USP18/STAT2 is a molecular hub
regulating autocrine IFN I-mediated control of Dengue and Zika virus replication. *bioRxiv*.
18. Axten JM, Romeril SP, Shu A, Ralph J, Medina JR, Feng Y, Li WHH, Grant SW, Heerding DA,
Minthorn E, et al. (2013) Discovery of GSK2656157: An Optimized PERK Inhibitor Selected for
Preclinical Development. *ACS Med Chem Lett* **4**: 964–968.
19. Fredericksen BL, Gale M Jr (2006) West Nile virus evades activation of interferon regulatory
factor 3 through RIG-I-dependent and -independent pathways without antagonizing host defense
signaling. *J Virol* **80**: 2913–2923.
20. Espada-Murao LA, Morita K (2011) Delayed cytosolic exposure of Japanese encephalitis virus
double-stranded RNA impedes interferon activation and enhances viral dissemination in porcine
cells. *J Virol* **85**: 6736–6749.
21. Overby AK, Popov VL, Niedrig M, Weber F (2010) Tick-borne encephalitis virus delays
interferon induction and hides its double-stranded RNA in intracellular membrane vesicles. *J
Virol* **84**: 8470–8483.
22. Schmidt EK, Clavarino G, Ceppi M, Pierre P (2009) SUnSET, a nonradioactive method to
monitor protein synthesis. *Nat Methods* **6**: 275–277.
23. Harding HP, Novoa I, Zhang Y, Zeng H, Wek R, Schapira M, Ron D (2000) Regulated
translation initiation controls stress-induced gene expression in mammalian cells. *Mol Cell* **6**:
1099–1108.
24. Novoa I, Zeng H, Harding HP, Ron D (2001) Feedback Inhibition of the Unfolded Protein

568		Response by GADD34-Mediated Dephosphorylation of eIF2α. *J Cell Biol* **153**: 1011–1022.

Response by GADD34-Mediated Dephosphorylation of eIF2α. *J Cell Biol* **153**: 1011–1022.

25. Clavarino G, Cláudio N, Couderc T, Dalet A, Judith D, Camosseto V, Schmidt EK, Wenger T,
Lecuit M, Gatti E, et al. (2012) Induction of GADD34 is necessary for dsRNA-dependent
interferon-β production and participates in the control of Chikungunya virus infection. *PLoS*
Pathog **8**: e1002708.

26. Dalet A, Argüello RJ, Combes A, Spinelli L, Jaeger S, Fallet M, Vu Manh T-P, Mendes A,
Perego J, Reverendo M, et al. (2017) Protein synthesis inhibition and GADD34 control IFN-β
heterogeneous expression in response to dsRNA. *EMBO J* **36**: 761–782.

27. Huang Y, Chuang AY, Romano R-A, Liégeois NJ, Sinha S, Trink B, Ratovitski E, Sidransky D
(2010) Phospho-DeltaNp63alpha/NF-Y protein complex transcriptionally regulates DDIT3
expression in squamous cell carcinoma cells upon cisplatin exposure. *Cell Cycle* **9**: 328–338.

28. Canino C, Luo Y, Marcato P, Blandino G, Pass HI, Cioce M (2015) A STAT3-
NFkB/DDIT3/CEBPβ axis modulates ALDH1A3 expression in chemoresistant cell
subpopulations. *Oncotarget* **6**: 12637–12653.

29. Lee Y-Y, Cevallos RC, Jan E (2009) An upstream open reading frame regulates translation of
GADD34 during cellular stresses that induce eIF2alpha phosphorylation. *J Biol Chem* **284**:
6661–6673.

30. Young SK, Willy JA, Wu C, Sachs MS, Wek RC (2015) Ribosome Reinitiation Directs Gene-
specific Translation and Regulates the Integrated Stress Response. *J Biol Chem* **290**: 28257–
28271.

31. Gilfoy FD, Mason PW (2007) West Nile virus-induced interferon production is mediated by the
double-stranded RNA-dependent protein kinase PKR. *J Virol* **81**: 11148–11158.

32. Zamanian-Daryoush M, Mogensen TH, DiDonato JA, Williams BR (2000) NF-kappaB
activation by double-stranded-RNA-activated protein kinase (PKR) is mediated through NF-
kappaB-inducing kinase and IkappaB kinase. *Mol Cell Biol* **20**: 1278–1290.

33. Czerkies M, Korwek Z, Prus W, Kochańczyk M, Jaruszewicz-Błońska J, Tudelska K, Błoński S,
Kimmel M, Brasier AR, Lipniacki T (2018) Cell fate in antiviral response arises in the crosstalk
of IRF, NF-κB and JAK/STAT pathways. *Nat Commun* **9**: 493.

34. Deb A, Haque SJ, Mogensen T, Silverman RH, Williams BR (2001) RNA-dependent protein
kinase PKR is required for activation of NF-kappa B by IFN-gamma in a STAT1-independent
pathway. *J Immunol* **166**: 6170–6180.

35. Kårehed K, Dimberg A, Dahl S, Nilsson K, Oberg F (2007) IFN-gamma-induced upregulation of
Fcgamma-receptor-I during activation of monocytic cells requires the PKR and NFkappaB
pathways. *Mol Immunol* **44**: 615–624.

36. Li Y, Xie J, Wu S, Xia J, Zhang P, Liu C, Zhang P, Huang X (2013) Protein kinase regulated by
dsRNA downregulates the interferon production in dengue virus- and dsRNA-stimulated human
lung epithelial cells. *PLoS One* **8**: e55108.

37. Kaempfer R, Ilan L, Cohen-Chalamish S, Turgeman O, Namer LS, Osman F (2019) Control of
mRNA Splicing by Intragenic RNA Activators of Stress Signaling: Potential Implications for
Human Disease. *Front Genet* **10**: 464.

38. Ben-Asouli Y, Banai Y, Pel-Or Y, Shir A, Kaempfer R (2002) Human interferon-gamma mRNA
autoregulates its translation through a pseudoknot that activates the interferon-inducible protein
kinase PKR. *Cell* **108**: 221–232.

39. Donzé O, Deng J, Curran J, Sladek R, Picard D, Sonenberg N (2004) The protein kinase PKR: a
molecular clock that sequentially activates survival and death programs. *EMBO J* **23**: 564–571.

40. Young SK, Wek RC (2016) Upstream Open Reading Frames Differentially Regulate Gene-
specific Translation in the Integrated Stress Response. *J Biol Chem* **291**: 16927–16935.

41. Umareddy I, Pluquet O, Wang QY, Vasudevan SG, Chevet E, Gu F (2007) Dengue virus
serotype infection specifies the activation of the unfolded protein response. *Virol J* **4**: 91.

42. Hou S, Kumar A, Xu Z, Airo AM, Stryapunina I, Wong CP, Branton W, Tchesnokov E, Götte
M, Power C, et al. (2017) Zika Virus Hijacks Stress Granule Proteins and Modulates the Host
Stress Response. *J Virol* **91**.:

43. Tu Y-C, Yu C-Y, Liang J-J, Lin E, Liao C-L, Lin Y-L (2012) Blocking double-stranded RNA-
activated protein kinase PKR by Japanese encephalitis virus nonstructural protein 2A. *J Virol* **86**:
10347–10358.

44. Peña J, Harris E (2011) Dengue virus modulates the unfolded protein response in a time-
dependent manner. *J Biol Chem* **286**: 14226–14236.
45. Tan Z, Zhang W, Sun J, Fu Z, Ke X, Zheng C, Zhang Y, Li P, Liu Y, Hu Q, et al. (2018) ZIKV
infection activates the IRE1-XBP1 and ATF6 pathways of unfolded protein response in neural
cells. *J Neuroinflammation* **15**: 275.
46. Gladwyn-Ng I, Cordón-Barris L, Alfano C, Creppe C, Couderc T, Morelli G, Thelen N, America
M, Bessières B, Encha-Razavi F, et al. (2018) Stress-induced unfolded protein response
contributes to Zika virus-associated microcephaly. *Nat Neurosci* **21**: 63–71.
47. Datan E, Roy SG, Germain G, Zali N, McLean JE, Golshan G, Harbajan S, Lockshin RA, Zakeri
Z (2016) Dengue-induced autophagy, virus replication and protection from cell death require ER
stress (PERK) pathway activation. *Cell Death Dis* **7**: e2127.
48. Wang Q, Xin X, Wang T, Wan J, Ou Y, Yang Z, Yu Q, Zhu L, Guo Y, Wu Y, et al. (2019)
Japanese Encephalitis Virus Induces Apoptosis and Encephalitis by Activating the PERK
Pathway. *J Virol* **93**.:
49. Sharma M, Bhattacharyya S, Sharma KB, Chauhan S, Asthana S, Abdin MZ, Vrati S, Kalia M
(2017) Japanese encephalitis virus activates autophagy through XBP1 and ATF6 ER stress
sensors in neuronal cells. *J Gen Virol* **98**: 1027–1039.
50. Ambrose RL, Mackenzie JM (2011) West Nile virus differentially modulates the unfolded
protein response to facilitate replication and immune evasion. *J Virol* **85**: 2723–2732.
51. Zanini F, Pu S-Y, Bekerman E, Einav S, Quake SR (2018) Single-cell transcriptional dynamics
of flavivirus infection. *Elife* **7**.:
52. Gjymishka A, Su N, Kilberg MS (2009) Transcriptional induction of the human asparagine
synthetase gene during the unfolded protein response does not require the ATF6 and IRE1/XBP1
arms of the pathway. *Biochem J* **417**: 695–703.
53. Dickhout JG, Carlisle RE, Jerome DE, Mohammed-Ali Z, Jiang H, Yang G, Mani S, Garg SK,
Banerjee R, Kaufman RJ, et al. (2012) Integrated stress response modulates cellular redox state
via induction of cystathionine γ-lyase: cross-talk between integrated stress response and thiol
metabolism. *J Biol Chem* **287**: 7603–7614.
54. Ma Y, Hendershot LM (2004) Herp is dually regulated by both the endoplasmic reticulum stress-
specific branch of the unfolded protein response and a branch that is shared with other cellular
stress pathways. *J Biol Chem* **279**: 13792–13799.
55. Fusakio ME, Willy JA, Wang Y, Mirek ET, Al Baghdadi RJT, Adams CM, Anthony TG, Wek
RC (2016) Transcription factor ATF4 directs basal and stress-induced gene expression in the
unfolded protein response and cholesterol metabolism in the liver. *Mol Biol Cell* **27**: 1536–1551.
56. Ohoka N, Yoshii S, Hattori T, Onozaki K, Hayashi H (2005) TRB3, a novel ER stress-inducible
gene, is induced via ATF4-CHOP pathway and is involved in cell death. *EMBO J* **24**: 1243–
1255.
57. Zanini F, Robinson ML, Croote D, Sahoo MK, Sanz AM, Ortiz-Lasso E, Albornoz LL, Rosso F,
Montoya JG, Goo L, et al. (2018) Virus-inclusive single-cell RNA sequencing reveals the
molecular signature of progression to severe dengue. *Proc Natl Acad Sci U S A* **115**: E12363–
E12369.
58. Hollien J, Weissman JS (2006) Decay of endoplasmic reticulum-localized mRNAs during the
unfolded protein response. *Science* **313**: 104–107.
59. Donovan J, Rath S, Kolet-Mandrikov D, Korennykh A (2017) Rapid RNase L-driven arrest of
protein synthesis in the dsRNA response without degradation of translation machinery. *RNA* **23**:
1660–1671.
60. Pause A, Belsham GJ, Gingras AC, Donzé O, Lin TA, Lawrence JC Jr, Sonenberg N (1994)
Insulin-dependent stimulation of protein synthesis by phosphorylation of a regulator of 5'-cap
function. *Nature* **371**: 762–767.
61. Poulin F, Gingras AC, Olsen H, Chevalier S, Sonenberg N (1998) 4E-BP3, a new member of the
eukaryotic initiation factor 4E-binding protein family. *J Biol Chem* **273**: 14002–14007.
62. Pestova TV, de Breyne S, Pisarev AV, Abaeva IS, Hellen CUT (2008) eIF2-dependent and eIF2-
independent modes of initiation on the CSFV IRES: a common role of domain II. *EMBO J* **27**:
1060–1072.
63. Robert F, Kapp LD, Khan SN, Acker MG, Kolitz S, Kazemi S, Kaufman RJ, Merrick WC,

Koromilas AE, Lorsch JR, et al. (2006) Initiation of protein synthesis by hepatitis C virus is refractory to reduced eIF2.GTP.Met-tRNA(i)(Met) ternary complex availability. *Mol Biol Cell* **17**: 4632–4644.

64. Rivas-Estilla AM, Svitkin Y, Lopez Lastra M, Hatzoglou M, Sherker A, Koromilas AE (2002) PKR-dependent mechanisms of gene expression from a subgenomic hepatitis C virus clone. *J Virol* **76**: 10637–10653.

65. Antunes KH, Fachi JL, de Paula R, da Silva EF, Pral LP, Dos Santos AÁ, Dias GBM, Vargas JE, Puga R, Mayer FQ, et al. (2019) Microbiota-derived acetate protects against respiratory syncytial virus infection through a GPR43-type 1 interferon response. *Nat Commun* **10**: 3273.

66. Guimarães ES, Gomes MTR, Campos PC, Mansur DS, Dos Santos AA, Harms J, Splitter G, Smith JA, Barber GN, Oliveira SC (2019) Cyclic Dinucleotides Trigger STING-Dependent Unfolded Protein Response That Favors Bacterial Replication. *J Immunol* **202**: 2671–2681.

67. Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F (2013) Genome engineering using the CRISPR-Cas9 system. *Nat Protoc* **8**: 2281–2308.

68. Beltramello M, Williams KL, Simmons CP, Macagno A, Simonelli L, Quyen NTH, Sukupolvi-Petty S, Navarro-Sanchez E, Young PR, de Silva AM, et al. (2010) The human immune response to Dengue virus is dominated by highly cross-reactive antibodies endowed with neutralizing and enhancing activity. *Cell Host Microbe* **8**: 271–283.

69. Loughran G, Howard MT, Firth AE, Atkins JF (2017) Avoidance of reporter assay distortions from fused dual reporters. *RNA* **23**: 1285–1289.

70. Lummertz da Rocha E, Rowe RG, Lundin V, Malleshaiah M, Jha DK, Rambo CR, Li H, North TE, Collins JJ, Daley GQ (2018) Reconstruction of complex single-cell trajectories using CellRouter. *Nat Commun* **9**: 892.