Chemical-genetic interaction mapping links carbon metabolism and cell wall

structure to tuberculosis drug efficacy

Eun-Ik Koh[1], Nadine Ruecker[2], Megan K. Proulx[1], Vijay Soni[2], Kenan C. Murphy[1],

Kadamba G. Papavinasasundaram[1], Charlotte J. Reames[1], Carolina Trujillo[2], Anisha

Zaveri[2], Matthew D. Zimmerman[3], Roshanak Aslebagh[4,5], Richard E. Baker[1], Scott A.

Shaffer[4,5], Kristine M. Guinn[6], Michael Fitzgerald[7], Véronique A. Dartois[3,8], Sabine Ehrt[2],

Deborah T. Hung[7,9,10], Thomas R. Ioerger[11], Eric Rubin[6], Kyu Y. Rhee[2], Dirk

Schnappinger[2] and Christopher M. Sassetti[1]

[1] Department of Microbiology and Physiological Systems, University of Massachusetts

Medical School, Worcester, Massachusetts, 01655 USA

[2] Department of Microbiology and Immunology, Weill Cornell Medical College, New

York, New York, 10065 USA

[3] Center for Discovery and Innovation, Hackensack Meridian Health, Nutley, New

Jersey, 07110 USA

[4] Department of Biochemistry and Molecular Pharmacology, University of

Massachusetts Medical School, Worcester, Massachusetts, 01655 USA

[5] Mass Spectrometry Facility, University of Massachusetts Medical School, Shrewsbury,

Massachusetts, 01545 USA

[6] Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of

Public Health, Boston, Massachusetts, 02115 USA

[7] Broad Institute of MIT and Harvard, Cambridge, Massachusetts, 02142 USA

[8] Department of Medical Sciences, Hackensack School of Medicine, Nutley, New

Jersey, 07110 USA

[9] Department of Molecular Biology and Center for Computation and Integrative Biology,

Massachusetts General Hospital, Boston, Massachusetts, 02114 USA

[10] Department of Genetics, Harvard Medical School, Boston, Massachusetts, 02114

USA

[11] Department of Computer Science and Engineering, Texas A&M University, College

Station, Texas, 77840 USA

Corresponding author:

Christopher M. Sassetti

368 Plantation Street, AS8-2051, Worcester, MA 01605

508-856-3678

Christopher.sassetti@umassmed.edu

Abstract

Current chemotherapy against *Mycobacterium tuberculosis (Mtb)*, an important human pathogen, requires a multidrug regimen lasting several months. While efforts have been made to optimize therapy by exploiting drug-drug synergies, testing new drug combinations in relevant host environments remains arduous. In particular, host environments profoundly affect the bacterial metabolic state and drug efficacy, limiting the accuracy of predictions based on *in vitro* assays alone. In this study, we utilize conditional *Mtb* knockdown mutants of essential genes as an experimentally-tractable surrogate for drug treatment, and probe the relationship between *Mtb* carbon metabolism and chemical-genetic interactions (CGI). We examined the anti-tubercular drugs isoniazid, rifampicin and moxifloxacin, and found that CGI are differentially responsive to the metabolic state, defining both environment-independent and – dependent synergies. Specifically, growth on the *in vivo*-relevant carbon source, cholesterol, reduced rifampicin efficacy by altering mycobacterial cell surface lipid composition. We report that a variety of perturbations in cell wall synthesis pathways restore rifampicin efficacy during growth on cholesterol, and that both environment-independent and cholesterol-dependent *in vitro* CGI could be leveraged to enhance bacterial clearance in the mouse infection model. Our findings present an atlas of novel chemical-genetic-environmental synergies that can be used to optimize drug-drug interactions as well as provide a framework for understanding *in vitro* correlates of *in vivo* efficacy.

Significance

Efforts to improve tuberculosis therapy include optimizing multi-drug regimens to take advantage of drug-drug synergies. However, the complex host environment has a profound effect on bacterial metabolic state and drug activity, making predictions of optimal drug combinations difficult. In this study, we leverage a newly developed library of conditional knockdown *Mycobacterium tuberculosis* mutants in which genetic depletion of essential genes mimics the effect of drug therapy. This tractable system allowed us to assess the effect of growth condition on predicted drug-drug interactions. We found that these interactions can be differentially sensitive to the metabolic state and select *in vitro*-defined synergies can be leveraged to accelerate bacterial killing during infection. These findings suggest new strategies for optimizing tuberculosis therapy.

Introduction

The current chemotherapeutic regimen for tuberculosis (TB) is the product of many decades of basic and clinical research. Since the first trials of streptomycin monotherapy in 1948 were rapidly followed by the emergence of antibiotic resistant clones (1), multidrug regimens to both suppress resistance and accelerate bacterial killing have become standard. The current regimen used against drug-sensitive strains of *Mycobacterium tuberculosis* (*Mtb*) consists of four antibiotics, isoniazid (INH), rifampicin (RIF), pyrazinamide (PZA), and ethambutol (EMB), and was optimized in a series of clinical trials in the 1970s (2, 3). While this "short-course regimen" has been credited with curing over 50 million patients, its delivery is complicated by the need for 6-9 months of drug administration (4). Furthermore, even in clinical trial settings where the delivery of this extended regimen is assured, 5-10% of patients fail therapy (5). The frequent transmission of antibiotic resistant *Mtb* strains has further complicated TB treatment options, requiring the use of less optimized drug combinations that are administered for even longer periods (6).

The factors that necessitate prolonged therapy are complex and specific to the infection environment. While antibiotics such as INH and RIF cause rapid cell death *in vitro*, their antimicrobial activities are much slower during infection (7). Limited drug penetration into *Mtb*-containing tissue lesions may reduce the efficacy of some drugs. For example, the intralesional RIF concentrations are significantly lower than what is achieved in the plasma (8, 9), and clinical studies suggest that increased RIF dosing improves bacterial clearance (10, 11). However, drug penetration is unlikely to fully account for the reduced

antibiotic efficacy in host tissue, as this complex and stressful environment has also

been shown to alter the physiology of the pathogen to induce a drug-tolerant state (12–

14). While a number of host immune-related stresses may be involved in this process,

simple changes in macronutrient availability can have important consequences. *Mtb* has

access to a mixture of glycolytic carbon sources, fatty acids and cholesterol in host

tissue (15–17), and altering the availability of these carbon sources *in vitro* can change

the efficacy of anti-tubercular compounds (18–23). A role for differential carbon

catabolism in determining drug efficacy is also supported by the identification of natural

genetic variants in clinical *Mtb* isolates that enhance drug tolerance by altering either

glycerol, lipid, or sterol catabolism (24, 25). The profound effect of the host environment

on bacterial metabolic state and drug activity makes it difficult to predict the ultimate

efficacy of an antibiotic regimen based on *in vitro* assays alone.

The most advanced efforts to accelerate TB therapy involve optimizing multi-drug

regimens to take advantage of drug-drug synergies. These pharmacological

interactions can improve therapy by increasing drug exposure, decreasing MIC, or

enhancing the maximal effect of the treatment (26). However, despite the demonstrated

benefits of synergistic regimens in many therapeutic realms, the combinatorial burden of

testing all potential multi-drug combinations remains cumbersome. While recent

advances to more efficiently predict synergies *in vitro* have proven valuable (27, 28), it

remains unclear whether the environmental influences that so profoundly alter the

efficacy of individual drugs will also influence drug-drug interactions (29, 30).

In this study, we leveraged a newly developed genetic resource to understand the effect

of growth conditions on potential drug-drug synergies. This "hypomorph library" consists

of individually DNA barcoded strains in which individual essential genes are tagged with

the DAS+4 sequence that targets proteins for degradation upon inducible induction of

the SspB adapter (31). Graded *sspB* expression produces different degrees of depletion

to model the effect of partial chemical inhibition (32). This approach allowed for highly

parallel assessment of chemical-genetic interactions (CGI) between individual

hypomorphic alleles and TB drugs under different growth conditions. We report that

drugs of distinct classes produce synergies that are differentially sensitive to the

environment, and that both condition-independent and –dependent CGIs can be

identified for all drugs tested. In particular, RIF efficacy was impaired as a result of cell

surface lipid alterations that occur during growth in cholesterol, and the cell could be

resensitized through a condition-specific interaction with the cell wall synthetic

machinery both *in vitro* and in the mouse lung. These observations provide a

compendium of chemical-genetic synergies that can be exploited to enhance therapy

and begin to define *in vitro* correlates of *in vivo* efficacy.

Results

Genetic strategy to define essential bacterial functions that alter drug efficacy *in*

vitro.

Host environmental factors alter antibiotic efficacy against TB (12, 24, 33, 34). To

determine if carbon source availability specifically affects antibiotic efficacy, we used the

Mtb strain H37Rv in minimal medium with glycerol, acetate or cholesterol as sole

carbon sources in a growth inhibition assay with the TB drugs isoniazid (INH), rifampicin

(RIF) and moxifloxacin (MOX). As these carbon sources support different growth rates,

endpoint-based antibiotic activity measurements such as MIC assays can be difficult to

interpret. Therefore we quantified growth rates in each carbon source over multiple

concentrations of antibiotics to determine the concentration that decreased the growth

rate by 50% (GR_{50}) (24, 35). We found different carbon sources to alter the GR_{50} of

these drugs in distinct ways. For INH, glycerol-dependent growth increased efficacy,

relative to the non-glycolytic carbon sources. In contrast, growth in cholesterol was

found to decrease RIF efficacy, relative to the other conditions. Media composition had

a more modest effect on MOX, altering GR_{50} by only approximately 2-fold **(Fig1A)**.

These findings show that carbon metabolism plays an important role in antibiotic

efficacy and has distinct effects on different drugs.

To understand the mechanisms linking drug efficacy and metabolism, we utilized a

barcoded-hypomorph library consisting of 465 essential genes to identify CGI with these

drugs (32). The library was grown in minimal media with glycerol, acetate or cholesterol,

and antibiotics were added at concentrations ranging from 0.05X to 1X GR_{50} for each.

Bacteria were subjected to these conditions in 96-well plates for 14 days, at which point

the relative abundance of individual mutants was assessed through multiplex PCR and

Next-Generation Sequencing **(Fig1B, S.Table 1)**. Relative abundances of individual

mutants were calculated by normalizing the mutant's barcode count to the total barcode

count and comparing the relative abundance of each mutant in different drug

concentrations with the untreated control samples.

We initially investigated the validity of this dataset by determining if genetic inhibition of

each drug's target produced the expected CGI with that compound in standard glycerol-

dependent growth conditions. RIF and MOX interact with the RNA polymerase subunit,

RpoB or the DNA gyrase subunit, GyrA, respectively. As expected, *rpoB* and *gyrA*

hypomorphic mutants were hypersensitive to RIF or MOX, and the CGI increased with

drug concentration **(Fig1C)**. The major target of INH, InhA, was not present in our

mutant pool. However, the library did contain a *ndh* mutant. Deficiency in this NADH-

dehydrogenase is known to decrease INH efficacy, likely due to the inefficient formation

of the active INH-NAD adduct (36). Consistent with these observations, we found that

the *ndh* hypomorphic mutant showed an INH dose-dependent increase in abundance in

our dataset **(Fig1C)**.

The hypomorph library was constructed to contain up to five different versions of each

DAS+4 tagged mutant, which express different levels of *sspB* and therefore produce

graded levels of protein depletion (31, 37). To investigate the relationship between

target protein abundance and phenotype, we correlated the strength of *sspB* expression

with drug sensitivity for a number of genes that were expected to alter antibiotic efficacy

(Fig1D). For RIF, we concentrated on *rpoB*. For INH, we examined *ndh* as well as

genes associated with mycolate biosynthesis, and found *kasA* and *hadB* mutants to be

significantly hypersensitive to increasing INH concentrations. For MOX, in addition to

gyrA, depletion of proteins associated with DNA integrity or deoxynucleotide production,

dnaN, *lexA* and *ndrF2,* showed significantly decreased fitness over increasing MOX

concentrations. For some mutants, particularly those involved in DNA metabolism, we

noted a dose-dependent effect of *sspB* expression, with larger phenotypes

corresponding to greater degrees of depletion. However, this was not a universal

phenomenon, and we often found that independent mutants expressing different *sspB*

levels produced consistent phenotypic effects. The observed differences in sensitivity to

protein depletion likely reflect the unique biochemistry of each pathway (38). As a result,

we considered each mutant corresponding to a target protein independently in the

following analyses.

CGI profiles are determined by the combination of carbon source and drug

To understand the relative importance of carbon metabolism and drug on bacterial

physiology, we compared the "fitness profiles" of each condition, which represented the

relative fitness of each mutant strain. We first selected the fold change in abundance

(log2FC) values of every mutant at 0.65X GR_{50} drug concentration that showed a

statistically significant fitness difference when at least one treated sample was

compared to the untreated control (918 mutants, Q <0.05). 0.65X GR_{50} drug

concentration displayed the largest number of mutants with a statistically significant

fitness difference across most conditions. Upon hierarchical clustering of both genes

and conditions, we found three major condition clusters that corresponded to each

antibiotic **(Fig2A)**. These condition clusters were differentiated by two distinct clusters of

mutants; one large gene set differentiated MOX from the other drugs (Fig 2A bottom

cluster), and differential fitness effects on a second smaller gene set differentiated INH

and RIF (Fig2A top cluster). Within each condition cluster, fitness effects due to carbon

source were apparent.

To catalog the genes that differentiate these conditions, we compared each drug-

treated sample (0.65X GR_{50}) with the corresponding untreated control and chose genes

with log2FC < -1 and p < 0.05 for each carbon source. For all antibiotics tested, there

were genes that showed synergy across all carbon sources as well as those associated

with a single carbon source **(Fig2B)**. Notably, MOX produced the largest number of

synergies in a condition-independent manner, while INH and RIF produced the largest

number of synergies during acetate and cholesterol-dependent growth respectively. To

more quantitatively assess the relative contribution of carbon source and drug, we used

Principal Component Analysis (PCA) to examine the amount of variance contributed by

each, concentrating only on the INH and RIF conditions, which produced the most

condition-specific effects. PC1 aligned to distinguish the carbon sources, while PC2

differentiated among drug treatments **(Fig2C)**. The first and second principal

components accounted for 37.6% and 23.7% respectively, of the overall variance

across the 9 data sets, indicating that drug and carbon source played a relatively equal

role in shaping bacterial physiology.

Condition-independent CGI can be reproduced with individual mutants.

Genes found to alter antibiotic efficacy from the high-throughput hypomorph analysis

were validated using individual mutant strains. For initial validation studies, we

concentrated on mutants predicted to show drug-hypersensitivity under all media

conditions. The repression of a drug's target is thought to increase sensitivity to

inhibition by reducing the fraction of the target that needs to be inhibited to reduce

growth (39). As such, we anticipated that drug target inhibition would result in condition-

independent effects. This prediction was verified using a *rpoB* hypomorphic mutant,

which our screening data predicted to increase RIF efficacy in all carbon sources

(Fig3A). When the MIC of RIF was determined using a standard microplate growth

assay, we verified that this degree of RpoB depletion decreased the RIF MIC by 4 to 16-

fold in each of the three media conditions **(Fig3A)**. In addition to known drug targets, a

number of additional genes were predicted to enhance RIF efficacy in a carbon source-

independent manner, including the transcriptional regulator, Rv0472c. This gene has

been named, mycolic acid desaturase regulator (*madR*), based on its specific effect on

the mycolate-modifying *desA1* and *desA2* genes (40). As we found for RpoB, MadR

depletion sensitized the bacterium to RIF in all tested carbon sources in the single strain

assay **(Fig3B)**. Finally, to ensure the specificity of these CGI, we investigated whether

they could be reversed by specific metabolite complementation. As predicted by the

screen, depletion of the lysine biosynthetic enzyme, LysA, produced RIF

hypersensitivity in both glycerol and cholesterol media, and we found this phenotype

was largely reversed by supplementation with 3mM lysine **(Fig3C)**. These data support

the predictions of the primary screen, and demonstrate that certain CGI are relatively

unaffected by carbon source.

Depletion of cell-wall biosynthetic genes produces condition-specific RIF synergy

As cholesterol represents a primary carbon source for *Mtb* during infection (16, 41), and

RIF efficacy is thought to be limited by drug exposure (42–44), we probed the

hypomorph fitness profiles for clues to the mechanism underlying the cholesterol-

dependent increase in RIF MIC. We found that inhibition of genes associated with

different components of the mycobacterial cell wall, including arabinogalactan (*embC*

and *aftB*), and mycolic acid (*kasA*) (45, 46) specifically sensitized the bacterium to RIF

during cholesterol growth **(Fig4A)**. These mutants were significantly underrepresented

by up to 8-fold upon RIF treatment specifically in cholesterol growth conditions across

multiple *sspB* doses and drug concentrations. To further explore the effect of cell wall

inhibition, we tested individual mutants with defects in peptidoglycan (*dapE*),

arabinogalactan (*aftB*), and mycolic acid (*hadB*) synthesis (45, 47–49). In this format,

we found that perturbation in each of these cell wall layers produced a similar effect, as

they all preferentially sensitized the bacterium to RIF in cholesterol growth conditions

(Fig4BCD). These observations suggest that alterations in cell wall structure might

underlie the relationship between cholesterol catabolism and reduced RIF efficacy.

RIF efficacy correlates with propionate catabolism and branched chain lipid

abundance.

Mtb catabolizes cholesterol to pyruvate, propionyl-CoA and acetyl-CoA (17, 50).

Propionyl-CoA is a precursor of branched-chain lipid synthesis, and catabolism of either

cholesterol or propionate increases cellular propionyl-CoA levels causing an increase in

both the abundance and chain-length of cell wall lipids, such as diacyltrehalose (DAT),

polyacyltrehalose (PAT), and sulfolipid-1 (SL-1) (50–53). To understand whether these

alterations in cell wall lipids could be responsible for altering RIF efficacy, we compared

RIF GR_{50} values in cells grown in cholesterol to those grown in butyrate media

supplemented with increasing concentrations of propionate. We observed that GR_{50}

values increased up to 8-fold with increasing propionate levels, and the addition of 0.1X

propionate, reflecting a 1:10 ratio of propionate to butyrate (weight:weight), mimicked

the effect of cholesterol **(Fig5A)**. To more quantitatively associate branched-chain lipid

abundance with drug efficacy, we performed relative quantification of the abundance of

SL-1 during growth in butyrate, butyrate and propionate, and cholesterol using mass

spectrometry. As expected, we found that growth in either propionate or cholesterol

increased both the *m/z* range of SL-1 and its total abundance, and the effect of

propionate was dose-dependent in the tested range **(Fig5B)**. This dose-dependency

allowed us to correlate changes in sulfolipid abundance with changes in rifampicin

efficacy. At multiple propionate concentrations, we calculated the GR_{50} for RIF and the

total abundance of SL-1. We found a correlation between these values, as 0.003X

propionate supplementation had no effect on either SL-1 abundance or RIF GR_{50}, and

0.1X increased both values **(Fig5C)**. These findings implied that the propionyl-CoA

derived from cholesterol alters rifampicin efficacy through modification of mycobacterial

surface lipids.

The different branched chain lipid species of *Mtb* are produced by distinct biosynthetic

pathways, and inhibiting the synthesis of one lipid can produce a compensatory

increase in others (52). Therefore, to assess whether increased lipid abundance was

causally related to RIF efficacy, we employed an *Mtb* mutant lacking the PhoPR

regulatory system, which is required for the synthesis of multiple branched chain lipid

species (54–56). In contrast to wild type *Mtb,* a Δ*phoPR* deletion mutant showed little

change in rifampicin efficacy with propionate supplementation or cholesterol compared

to butyrate growth conditions, and possessed significantly decreased sulfolipid levels

compared to WT **(Fig6AB)**. Similar to the effects seen with the Δ*phoPR* mutant, cell wall

defective hypomorphs (*aftB, hadB, dapE*) also maintained a relatively consistent RIF

GR50 in all media conditions, compared to wild type **(Fig6CD)**. These findings suggest

that cholesterol catabolism reduces RIF efficacy via a propionyl-CoA driven increase in

synthesis of cell surface lipids, and that this effect can be reversed by perturbing the

structure of the cell envelope.

In vitro* CGI predict drug efficacy *in vivo

To determine whether synergies identified *in vitro* predict strategies to accelerate

bacterial killing in the lung, we infected C57BL/6J mice via the aerosol route with a

pooled culture consisting of 3 barcoded WT strains and a number of select hypomorph

mutants. We concentrated on genes that either displayed carbon source-independent

synergy with RIF (*rpoB*, *madR*), or mutants that disrupt the peptidoglycan (*dapE, murA*)

or arabinogalactan (*aftB*) layer of the cells wall and would be expected to reverse the

effect of host-derived cholesterol catabolism **(Fig3AB, Fig4BC, S.Fig1)**. To ensure that

the animals were treated with a relevant dose of antibiotic, we measured the plasma

concentration of RIF in mice and adjusted the dosing to match the 24-hour exposure

observed during clinical TB therapy **(Fig7A)** (57). The mice were fed doxycycline chow

starting 3 days before infection to repress *sspB* expression in the mutants upon

infection, and allow them to grow normally. At 3 weeks post-infection, doxycycline was

withdrawn to initiate protein degradation, and two weeks of RIF treatment was

administered **(Fig7B)** (58). The antibiotic regimen effectively decreased the total

number of lung CFU **(Fig7C)**. Lung homogenates were washed free of RIF and plated

on 7H10 plates with supplementation to recover viable clones. The relative abundance

of individual mutants was normalized to the average abundance of three barcoded WT

strains for each mouse and fitness changes were calculated by comparing to those from

untreated mice. Each of the mutants showed significant decrease in abundance in RIF

treated animals, relative to wild type. AftB and MurA depletion increased bacterial

clearance to the same degree as RpoB depletion. Depleting MadR had an even greater

effect, and the abundance of this mutant was reduced below our limit of detection in four

out of five animals **(Fig7D)**. DapE depletion mutant was below our limit of detection in

both untreated and treated animals. These findings show that CGI identified *in vitro* can

accelerate bacterial clearance during infection, and highlight the importance of the

environment for defining relevant interactions.

Discussion

Optimizing combination therapy is critical to improving TB treatment. While efficient

strategies for predicting and quantifying drug-drug interactions under *in vitro* conditions

have been developed, the impact of bacterial environment on these interactions has not

been determined (27, 28). As a result, it remains unclear how well these *in vitro* data

predict the effects of combination therapy in the infection environment. In this study, we

modeled drug-drug interactions using a newly developed genetic hypomorph library to

assess the impact of bacterial carbon metabolism on CGI. This work revealed that

synergies with distinct drugs are differentially sensitive to the environment, suggesting

tailored approaches to optimization.

While global analysis of our chemical-genetic-environmental dataset suggested that

drug mechanism and environment both play critical roles in shaping chemical-genetic

interactions, all drugs were not equally affected by condition. In particular, the individual

effect of the fluoroquinolone, MOX, was unaffected by carbon source, and this relative

indifference to the environment was also observed for CGI with MOX. These

observations contrast with RIF and INH, where carbon source plays an important role in

determining the MIC of the drug alone, as well as shaping chemical-genetic interaction

profiles. While we can only speculate on the mechanistic basis for these differences

between drugs, we found that the cholesterol-dependent decrease in RIF sensitivity is

due to the profound effect of carbon source on cell envelope structure that is likely to

alter RIF penetration. Similarly, the activation of the prodrug, INH, is affected by the

redox state of the cell (59), another process that is impacted by carbon catabolism. In

contrast, MOX is not a prodrug and we speculate that it may not be limited by cellular

penetration as strongly as RIF. More practically, these observations suggest that drug-

drug interactions with MOX may be particularly robust to changes in condition, and

therefore may translate well between *in vitro* and *in vivo* conditions.

While RIF synergies were complex and condition-dependent compared to MOX, we

found that the cholesterol-specific effects translate well to the infection environment and

may be particularly relevant to treatment. As reduced RIF exposure limits the efficacy of

this critical component of our standard regimen, even small increases in MIC during

infection could have a significant effect on bacterial killing. A number of observations

indicate that RIF efficacy is limited by similar mechanisms *in vivo* and in cholesterol-

containing media. Firstly, the increase in branched-chain lipid abundance that is

associated with reducing RIF efficacy in cholesterol- or propionate-containing media has

also been observed during infection, where similar carbon sources are utilized (16, 52).

Furthermore, genetic perturbation of the same cell wall synthetic enzymes sensitized

Mtb to RIF in both conditions. This effect may underlie the increased cell-associated RIF

upon simultaneous INH and ethambutol treatment *in vitro* (60). While the importance of

cell wall architecture and cell envelope lipid production suggests that cellular

permeability to RIF may be altered during growth on cholesterol, we note that lipid

synthesis also has important effects on central carbon metabolic pathways that could

influence antibiotic susceptibility (15). Additional enzymes identified in this study may

also be attractive targets for further drug development, such as AftB, a

arabinofuranosyltransferase distinct from EmbA and EmbB that are targets of

ethambutol (46). Similarly, enzymes in the DAP (diaminopimelic acid) biosynthesis

pathway produce precursors for the synthesis of peptidoglycan and lysine (61), and the

loss of either sensitizes the cell to RIF.

Using a chemical-genetic system, we determined that cellular environments can have a

profound effect on drug-drug interactions, and that even the use of relatively simple *in*

vitro culture conditions can identify synergies that are relevant during infection. These

studies provide a tractable system that can incorporate more complex culture systems,

or even cellular models, to identify additional *in vivo*-relevant synergies. Expanding this

CGI atlas promises to elucidate the processes limiting drug efficacy during infection and

guide drug development and regiment optimization efforts.

Material and methods

Bacterial cultures

M. tuberculosis (H37Rv) cells were cultured in Middlebrook 7H9 medium supplemented

with 10% oleic acid-albumin-dextrose-catalase (OADC), 0.5% glycerol, and 0.05%

Tween-80, or in minimal medium supplemented with 0.1% Tyloxapol and variable levels

of glycerol, acetate, butyrate, propionate, or cholesterol up to 0.1%. Minimal medium

was made as previously described but with ferric chloride (100uM) replacing ferric

ammonium citrate (50).

Cells were also cultured on 7H10 agar medium supplemented with 10% OADC and

0.5% glycerol. Streptomycin (20ug/mL) was supplemented when necessary.

Anhydrotetracycline (ATC, 500ng/mL) was supplemented periodically until cultures

reach exponential growth. ATC was removed from cultures prior to antibiotic growth

inhibition assays by washing in PBS supplemented with 0.1% Tyloxapol. *ΔphoPR*

mutant was generated as described previously (62).

Multiplexed library screening

Library pools of hypomorph mutants were prepared as described previously (32). Pools

were grown in minimal medium with 0.1% glycerol, 0.1% acetate, or 0.1% cholesterol

on 96-well plates for 2 weeks. Isoniazid and moxifloxacin were used at 1ug/mL and

rifampicin was used at 0.062 ug/mL. Antibiotics were subsequently serially diluted in a

2-fold manner. An untreated growth condition was included for each study. Samples

were inoculated at OD600 0.05 and growth was monitored using OD600. Upon

completion, 96-well plates were heat-inactivated at 85 degrees for 2 hours. Barcodes

were PCR amplified as described previously (32). Individual libraries were mixed with

1:1 20% DMSO and heated for 10 minutes at 98 degrees prior to PCR reaction.

Amplified barcodes were purified using SPRI-based purification methods and

sequenced using Next-generation sequencing methods. Sequence alignment and

analysis were conducted using Bowtie software package with index mismatch set to 2

bases and barcode mismatch set to 1 base. Relative abundance of every mutant was

calculated as mean ratio of barcode abundance of mutant relative to total barcode

abundance of library. Log2FC of a mutant was calculated as the log2 of the ratio of the

mean relative abundances in a given antibiotic condition relative to its untreated control.

Hierarchical clustering and principal component analysis were conducted using ClustVis

(63). Hierarchical clustering was applied to vectors of Log2FC of each gene across all

conditions. PCA was performed on relative abundances across all conditions. Statistical

significance was determined by unpaired *t*-test with Benjamini-Hochberg multiple testing

correction.

Antibiotic growth inhibition assay

Mtb cells were grown in minimal medium with 0.1% glycerol, 0.01% acetate, 0.01%

butyrate, 0.01% propionate, or 0.01% cholesterol on 96-well plates. Isoniazid and

moxifloxacin were used at 1ug/ml and rifampicin was used at 0.062 ug/mL. Antibiotics

were subsequently serially diluted in a 2-fold manner. Untreated condition was included

for each study. Cells were inoculated at OD600 0.05 and growth was monitored using

OD600. Antibiotic efficacy was determined using growth rate inhibition as done

previously (24). The exponential growth constant (k) value was determined for all

conditions. The k-value of each antibiotic concentration was normalized to the k-value of

the no-drug control. The GR_{50} value was defined as the concentration of antibiotic that

resulted in 50% decrease in growth rate. L-lysine (Sigma) was added to growth

conditions to final concentration of 3mM.

Lipid extraction and mass spectrometry

Mtb cells were grown in minimal medium with 0.01% butyrate, propionate or cholesterol.

Cells were inoculated at OD600 0.1 and grown to final OD600 of 0.7. Cells were

pelleted and heat-inactivated at 85 degrees for 45 minutes. Cells were washed in 10%

glycerol to remove residual detergents from growth media. Lipid extraction was

conducted with methyl tert-butyl ether (MTBE) as described previously (64). 1.5mL

methanol and 5mL MTBE were mixed with the cell pellet and placed into a glass tube

with a Teflon-lined cap. The mixture was incubated at room temperature for 5 hours

while shaking. Phase-separation was achieved by adding 0.75mL water to the tubes,

incubating for 10 minutes at room temperature, then centrifuging at 1000g for 10

minutes. The 2.5mL of the top phase was collected and transferred to a fresh glass tube

with Teflon-lined cap. Samples were dried using a nitrogen evaporator and stored in -20

degrees prior to injection to mass spectrometer. After evaporation of MTBE, 300 μL of

2:1 methanol:chloroform was added to each tube followed by vigorous vortex.

Solubilized lipids were then transferred to a new, pre-weighed tube, dried using a

nitrogen evaporator, and then tubes were re-weighed to determine the lipid mass in

each sample. Then 1.5 µL of 100 ng/µL N-omega-CD_3-octadecanoyl

monosialoganglioside GM_2 (Matreya, State College, PA) was added to each sample for

normalization and samples were reconstituted in 300 µL of 2:1 methanol: chloroform for

mass spectrometry analysis.

Samples were analyzed by direct infusion on a syringe pump to an Orbitrap Velos Pro

(Thermo Fisher Scientific, Waltham, MA) mass spectrometer operating in the negative

electrospray ionization mode. Mass spectra were acquired at a flow rate of 10 µL/min

for 2 minutes in two different mass ranges, *m/z* 200-2000 and *m/z* 300-3000, acquiring

two replicates for each range using a resolution of 30,000 (*m/z* 200), an AGC target

population of $5e^5$ and a maximum ion injection time of 100 ms. Data were analyzed in

Xcalibur 2.2 (Thermo Scientific). Briefly, mass spectra were averaged over the entire

acquisition range (0-2 minutes), producing one averaged spectrum for each analysis.

Peak lists were exported in Excel and intensity data was normalized to the intensity of

the GM2 ganglioside spike.

Plasma pharmacokinetic analysis

C57BL/6J mice were purchased from Jackson Laboratories. Housing and

experimentation were done in accordance with the guidelines set by the Department of

Animal Medicine of University of Massachusetts Medical School and Institutional Animal

Care and Use Committee and adhered to the laws of the United States and regulations

of the Department of Agriculture. Eight- to 12-week mice were administered 0.1g/L RIF

through drinking water for three days. Blood was collected during a 12-hour period 24

hours post RIF administration, kept on ice, centrifuged at approximately 2500 x g for 5

minutes. After centrifugation, plasma was collected and stored at -80°C until analysis.

Neat 1 mg/mL DMSO stocks for RIF were serial diluted in 50/50 acetonitrile/water to

create neat spiking stocks. Standards and quality control (QC) samples were created by

adding 10 µL of spiking stock to 90 µL of drug free plasma. Ten microliters of control,

standard, QC, or study sample were added to 100 µL of acetonitrile/methanol 50/50

protein precipitation solvent containing 20 ng/mL RIF-d8. Extracts were vortexed for 5

minutes and centrifuged at 4000 RPM for 5 minutes. 100 µL of supernatant was

combined with 5 µL of 75 mg/mL ascorbic acid to stabilize RIF. 100 µL of mixture was

combined with 100 µL of Milli-Q water prior to HPLC-MS/MS analysis. Mouse control

plasma (K2EDTA) was sourced from Bioreclamation. Mouse control lungs were

collected in house. RIF was sourced from Sigma Aldrich and RIF-d8 was purchased

from Toronto Research Chemicals.

LC-MS/MS analysis was performed on a Sciex Applied Biosystems Qtrap 6500+ triple-

quadrupole mass spectrometer coupled to a Shimadzu Nexera X2 UHPLC system to

quantify each drug in plasma. Chromatography was performed on a Agilent SB-C8

(2.1x30 mm; particle size, 3.5 µm) using a reverse phase gradient. Milli-Q deionized

water with 0.1% formic acid was used for the aqueous mobile phase and 0.1% formic

acid in acetonitrile for the organic mobile phase. Multiple-reaction monitoring of

precursor/product transitions in electrospray positive-ionization mode was used to

quantify the analytes. Sample analysis was accepted if the concentrations of the quality

control samples were within 20% of the nominal concentration. The compounds were

ionized using ESI positive mode ionization and monitored using masses RIF

(823.50/791.60) and RIF-d8 (831.50/799.60). Data processing was performed using

Analyst software (version 1.6.2; Applied Biosystems Sciex).

***In vivo* antibiotic susceptibility**

C57BL/6J mice were purchased from Jackson Laboratories. Housing and

experimentation were done in accordance with the guidelines set by the Department of

Animal Medicine of University of Massachusetts Medical School and Institutional Animal

Care and Use Committee and adhered to the laws of the United States and regulations

of the Department of Agriculture. Eight- to 12-week mice were infected with pools of

strains at equal ratios through the aerosol route (500 to 1000 CFU/mouse). Mice were

fed doxycycline-containing chow (Purina 5001 with 2000 ppm doxycycline, Research

Diets C11300-2000i) starting 3 days pre-infection to 21 days post-infection. At 21 days

post-infection, 0.1g/L rifampicin was administered through drinking water for 14 days. At

35 days post-infection, mice were sacrificed, spleen and lungs were isolated and

homogenized, and CFU was determined by plating dilutions on 7H10 agar with 50ug/mL

streptomycin and 500ng/mL ATC. For library recovery, approximately 1 million CFU per

mouse were plated on 7H10 agar with 50ug/mL streptomycin and 500ng/ml ATC.

Genomic DNA was extracted and normalized as done previously (34) and sequenced

using multiplex PCR methods. Sequence alignment and analysis were conducted using

Bowtie software package as described above. Relative abundance of every mutant was

calculated as mean ratio of barcode abundance of mutant relative to the average

barcode abundance of three barcoded H37Rv strains. Log2FC of a mutant was

calculated as the log2 of the ratio of the mean relative abundances of rifampicin treated

mice relative to the untreated mice. Statistical significance was determined by unpaired

t-test.

Figure 1. Genetic strategy to define essential bacterial functions that alter drug

efficacy *in vitro*. (A) Normalized growth inhibition of WT *Mtb* across increasing

concentrations of INH, RIF and MOX in minimal media with glycerol (black), acetate

(blue) and cholesterol (red) as sole carbon sources. Results shown as means from 3

biological replicates with standard deviations. **(B)** Barcoded hypomorph mutants were

pooled and grown in 96-well plates containing minimal media with glycerol, acetate or

cholesterol as the sole carbon source for 14 days. Antibiotics were added to individual

wells as well as untreated controls. Chromosomal barcodes were PCR amplified and

pooled for Illumina Next Generation Sequencing (NGS). Barcodes were analyzed to

quantify changes in fitness of individual strains from different conditions. **(C)** Boxplot

representing changes in relative abundances of *rpoB*, *gyrA* and *ndh* mutants during RIF,

MOX and INH treatment respectively. Data represents 5 biological replicates.

Significance was calculated using unpaired t-test and compared to untreated conditions,

554   *$p<0.05$, **$p<0.01$, ***$p<0.001$. **(D)** Heat map representing changes in fitness of

555   individual mutants, shown as Log2 fold change, during INH, RIF and MOX treatment in

556   glycerol growth conditions. Mutants were chosen from previous association with

557   respective antibiotic. sspB numbers denote *sspB* expression level of mutant. Results

558   shown as means from 5 biological replicates.

Figure 2. Carbon sources mediate antibiotic-genetic interactions of essential

bacterial functions. (A) Heat map of hypomorph mutants shown as Log2 fold change

of 0.65X GR_{50} of INH, RIF and MOX against untreated controls. Results shown as

means from 5 biological replicates. Each antibiotic (INH, RIF and MOX) in single carbon

source (glycerol, acetate and cholesterol) conditions were compared using hierarchical

clustering methods. **(B)** Venn diagrams showing the total number of genes with log2FC

> -1 and $p<0.05$ at 0.65X GR_{50} of INH, RIF and MOX, and the % of genes associated

with specific conditions. (**C**) Relative abundance datasets of the hypomorph libraries

from glycerol (black), acetate (blue) and cholesterol (red) growth conditions with 0.65X

GR_{50} INH (triangle) and RIF (square) treatment as well as untreated controls (circle)

572   were examined using principal component analysis. Units shown on axes are arbitrary

573   values in principal component space.

Figure 3. Condition-independent chemical-genetic synergies can be reproduced

with individual mutants. (A) RpoB depletion shows decreased fitness compared to the

library over increasing concentrations of RIF for glycerol (black), acetate (blue) and

cholesterol (red). Normalized growth inhibition assay shows *rpoB* hypomorph mutant

(green) has decreased RIF MIC compared to WT (black) across multiple carbon

conditions. **(B)** MadR depletion shows decreased fitness compared to the library over

increasing concentrations of rifampicin for glycerol (black), acetate (blue) and

cholesterol (red). Normalized growth inhibition assay shows *madR* hypomorph mutant

(brown) has decreased RIF MIC compared to WT (black) across multiple carbon

conditions. **(C)** LysA depletion shows decreased fitness compared to the library over

increasing concentrations of rifampicin for glycerol (black), acetate (blue) and

cholesterol (red). Normalized growth inhibition assay shows *lysA* hypomorph mutant

(green) has decreased RIF MIC compared to WT (black), which can be reversed by

3mM lysine supplementation (purple). Library results are shown as means from 5

biological replicates, and significance was calculated using unpaired t-test with

Benjamini-Hochberg multiple testing correction, $*q<0.05$, $**q<0.01$, $***q<0.001$.

Normalized growth inhibition results are shown as means from 3 biological replicates

with standard deviations.

Figure 4. Cell-wall biosynthesis hypomorph mutants show altered RIF efficacy

during cholesterol growth conditions. (A) Heat map showing hypomorph mutants of

select mycobacterial cell-wall biosynthesis genes and pathways with significant changes

in fitness during cholesterol growth, shown as Log2 fold change at 0.65X and 0.75X

GR_{50} of RIF against untreated controls. Results shown as means from 5 biological

replicates. Classification of mutants was done using Mycobrowser. **(B-D)** Normalized

growth inhibition of WT and select mutants across increasing concentrations of RIF in

minimal media with sole carbon sources. Depletion of AftB (blue), DapE (red) and HadB

(green) show decreased RIF MIC compared to WT (black) only during cholesterol

growth conditions. Results shown as means from 3 biological replicates with standard

deviations.

Figure 5. Propionate and cholesterol catabolism decreases RIF efficacy in

correlation with increased branched chain lipid length and abundance. (A)

Normalized growth inhibition of WT across increasing concentrations of RIF in minimal

media with 1X butyrate (BT) and corresponding propionate (PP) supplementations as

well as cholesterol. Results shown as means from 3 biological replicates with standard

deviations. **(B)** Normalized mass spectra of extracted sulfolipids from WT grown in

minimal media with 1X butyrate (BT) and corresponding propionate (PP)

supplementations as well as cholesterol. MS spectra intensities were normalized using

GM2 ganglioside internal standard and cell density. Results shown as representative

from 2 biological and 2 technical replicates. **(C)** Plotted GR_{50} values (black) and relative

622   sulfolipid abundances (red) across increasing propionate levels. Relative sulfolipid

623   abundances were determined by the combined normalized mass spectra intensities

624   from *m/z* 2000-2800 for each growth condition. Results shown as means with standard

625   deviations.

Figure 6. Disruption of the cell envelope reverses propionate and cholesterol

catabolism dependent decrease in RIF efficacy. (A) Normalized growth inhibition of

Δ*phoPR* mutant across increasing concentrations of RIF in minimal media compared to

WT. Results shown as means from 3 biological replicates. **(B)** Relative sulfolipid

abundances of WT and Δ*phoPR* mutant grown in minimal media with 1X butyrate and

0.1X propionate (1X BT+0.1X PP), and cholesterol. Relative sulfolipid abundances were

determined by the combined normalized mass spectra intensities from *m/z* 2000-2800

for each growth condition. MS spectra intensities were normalized using GM2

ganglioside internal standard and cell density. Results shown as means with standard

deviations from 2 biological and 2 technical replicates. Significance was calculated

using unpaired t-test, *$p<0.05$, **$p<0.01$. **(C)** Normalized growth inhibition of *aftB* mutant

641   across increasing concentrations of RIF in minimal media compared to WT. Results

642   shown as means from 3 biological replicates. **(D)** Calculated fold change in GR_{50} values

643   of 1X butyrate and 0.1X propionate (1X BT+0.1X PP) and cholesterol conditions against

644   1X butyrate alone for WT and mutant strains. Significance was calculated using

645   unpaired t-test, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$.

Figure 7. Essential bacterial functions that alter drug efficacy *in vivo*. (A) RIF

plasma concentrations were measured in mice over 12-hour period 24 hours post RIF

administration (0.1g/L). Results shown as Area Under the Concentration (AUC) time

profile. Dotted lines indicate RIF plasma range observed during clinical TB therapy (57).

(B) C57BL/6J mice were infected through the aerosol route with a pooled culture of

individual hypomorph mutants and barcoded WT strains. Mice were fed doxycycline

chow starting 3 days before infection to 3 weeks post infection. Mice were subsequently

switched to normal chow and water with or without RIF (0.1g/L) for 2 additional weeks.

Lungs were harvested, homogenized and plated for *Mtb* outgrowth. Upon DNA

extraction of grown colonies, chromosomal barcodes were PCR amplified and pooled

for Illumina Next Generation Sequencing (NGS). Barcode abundances of individual

mutants were normalized to WT strains and analyzed to quantify changes in fitness

during RIF treatment. **(C)** RIF treated mice show significant decrease in *Mtb* CFU in

lungs compared to untreated controls. Each dot represents a single mouse. Significance

was calculated using unpaired t-test, ***$p<0.001$. **(D)** Depletion of RpoB, AftB and MurA

show significant decrease in fitness, shown as Log2 fold change, during rifampicin

treatment compared to untreated controls. Each dot represents a single mouse and

empty dot indicates that the relative abundance was below the limit of detection (Dotted

line). Significance was calculated using unpaired t-test, *$p<0.05$, **$p<0.01$

Supplementary Figure 1. MurA hypomorph mutant shows altered RIF efficacy

during cholesterol growth conditions. (A) Normalized growth inhibition of WT and

MurA hypomorph mutant across increasing concentrations of RIF in minimal media with

cholesterol as the sole carbon source. Depletion of MurA (purple) show decreased RIF

MIC compared to WT (black) during cholesterol growth conditions. Results shown as

means from 3 biological replicates with standard deviations.

Acknowledgements

We are thankful to the members of the Sassetti lab for both technical assistance and

helpful discussion. We thank Sovie Lavalette-Levy and Curtis Engelhart for technical

help. This work was supported by the Office of the Assistant Secretary of Defense for

Health Affairs through the Peer Reviewed Medical Research Program, Focused

Program Award, under award no. W81XWH-17-1-0692. Opinions, interpretations,

conclusions, and recommendations are those of the author and are not necessarily

endorsed by the Department of Defense. The work was additionally supported by the

NIH (AI095208). Model figures were created with BioRender.com

1. ,   Streptomycin Treatment of Pulmonary Tuberculosis. *Br. Med. J.* **2**, 769–782 (1948).

2.   W. Fox, D. Mitchison, Short-course chemotherapy for pulmonarytuberculosis. *Am Rev*
Respir Dis **111**, 325–353 (1975).

3.   World Health Organization, *Guidelines for treatment of tuberculosis, 4th edition* (World
Health Organization, 2010).

4.   World Health Organization, *Global tuberculosis report 2020.* (World Health
Organization, 2020).

5.   S. H. Gillespie, *et al.*, Four-Month Moxifloxacin-Based Regimens for Drug-Sensitive
Tuberculosis. *http://dx.doi.org/10.1056/NEJMoa1407426* (2014)
https:/doi.org/10.1056/NEJMoa1407426 (November 4, 2020).

6.   P. Nahid, *et al.*, Treatment of Drug-Resistant Tuberculosis. An Official
ATS/CDC/ERS/IDSA Clinical Practice Guideline. *Am. J. Respir. Crit. Care Med.* **200**, e93–
e142 (2019).

7.   J. E. Gomez, J. D. McKinney, M. tuberculosis persistence, latency, and drug tolerance.
Tuberculosis **84**, 29–44 (2004).

8.   A. A. Ordonez, *et al.*, Dynamic imaging in patients with tuberculosis reveals
heterogeneous drug exposures in pulmonary lesions. *Nat. Med.* **26**, 529–534 (2020).

9.   M. C. Kjellsson, *et al.*, Pharmacokinetic Evaluation of the Penetration of
Antituberculosis Agents in Rabbit Pulmonary Lesions. *Antimicrob. Agents Chemother.*
56, 446–457 (2012).

10. R. J. Svensson, *et al.*, Greater Early Bactericidal Activity at Higher Rifampicin Doses
Revealed by Modeling and Clinical Trial Simulations. *J. Infect. Dis.* **218**, 991–999
(2018).

11. M. J. Boeree, *et al.*, High-dose rifampicin, moxifloxacin, and SQ109 for treating
tuberculosis: a multi-arm, multi-stage randomised controlled trial. *Lancet Infect. Dis.*
17, 39–49 (2017).

12. S.-H. Baek, A. H. Li, C. M. Sassetti, Metabolic regulation of mycobacterial growth and
antibiotic sensitivity. *PLoS Biol.* **9**, e1001065 (2011).

13. K. N. Adams, *et al.*, Drug Tolerance in Replicating Mycobacteria Mediated by a
Macrophage-Induced Efflux Mechanism. *Cell* **145**, 39–53 (2011).

14. Y. Liu, *et al.*, Immune activation of the host cell induces drug tolerance in
Mycobacterium tuberculosis both in vitro and in vivo. *J. Exp. Med.* **213**, 809–825
(2016).

15. R. R. Lovewell, C. M. Sassetti, B. C. VanderVen, Chewing the fat: lipid metabolism and
homeostasis during M. tuberculosis infection. *Curr. Opin. Microbiol.* **29**, 30–36 (2016).

16. A. K. Pandey, C. M. Sassetti, Mycobacterial persistence requires the utilization of host
cholesterol. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 4376–4380 (2008).

17. A. D. Baughn, K. Y. Rhee, Metabolomics of Central Carbon Metabolism in
Mycobacterium tuberculosis. *Microbiol. Spectr.* **2** (2014).

18. D. A. Aguilar-Ayala, *et al.*, Antimicrobial activity against Mycobacterium tuberculosis
under in vitro lipid-rich dormancy conditions. *J. Med. Microbiol.* **67**, 282–285 (2018).

19. J. J. Baker, R. B. Abramovitch, Genetic and metabolic regulation of Mycobacterium
tuberculosis acid growth arrest. *Sci. Rep.* **8**, 1–16 (2018).

20. S. G. Franzblau, *et al.*, Comprehensive analysis of methods used for the evaluation of
compounds against Mycobacterium tuberculosis. *Tuberculosis* **92**, 453–488 (2012).

21. K. Pethe, *et al.*, A chemical genetic screen in Mycobacterium tuberculosis identifies
carbon-source-dependent growth inhibitors devoid of in vivo efficacy. *Nat. Commun.* **1**,
1–8 (2010).

22. P. Gopal, *et al.*, Pyrazinamide Resistance Is Caused by Two Distinct Mechanisms:
Prevention of Coenzyme A Depletion and Loss of Virulence Factor Synthesis. *ACS*
Infect. Dis. **2**, 616–626 (2016).

23. N. P. Kalia, *et al.*, Carbon metabolism modulates the efficacy of drugs targeting the
cytochrome bc1:aa3 in Mycobacterium tuberculosis. *Sci. Rep.* **9** (2019).

24. M. M. Bellerose, *et al.*, Common Variants in the Glycerol Kinase Gene Reduce
Tuberculosis Drug Efficacy. *mBio* **10**, e00663-19 (2019).

25. N. D. Hicks, *et al.*, Clinically prevalent mutations in Mycobacterium tuberculosis alter
propionate metabolism and mediate multidrug tolerance. *Nat. Microbiol.* **3**, 1032–
1042 (2018).

26. G. R. Zimmermann, J. Lehár, C. T. Keith, Multi-target therapeutics: when the whole is
greater than the sum of the parts. *Drug Discov. Today* **12**, 34–42 (2007).

27. M. Cokol, N. Kuru, E. Bicak, J. Larkins-Ford, B. B. Aldridge, Efficient measurement and
factorization of high-order drug interactions in Mycobacterium tuberculosis. *Sci. Adv.*
3, e1701881 (2017).

28. S. Ma, *et al.*, Transcriptomic Signatures Predict Regulators of Drug Synergy and Clinical
Regimen Efficacy against Tuberculosis. *mBio* **10** (2019).

29. J. Pasipanodya, T. Gumbo, An Oracle: Antituberculosis Pharmacokinetics-
Pharmacodynamics, Clinical Correlation, and Clinical Trial Simulations To Predict the
Future. *Antimicrob. Agents Chemother.* **55**, 24–34 (2011).

30. E. Nuermberger, C. Sizemore, K. Romero, D. Hanna, Toward an Evidence-Based
Nonclinical Road Map for Evaluating the Efficacy of New Tuberculosis (TB) Drug
Regimens: Proceedings of a Critical Path to TB Drug Regimens-National Institute of
Allergy and Infectious Diseases In Vivo Pharmacology Workshop for TB Drug
Development. *Antimicrob. Agents Chemother.* **60**, 1177–1182 (2016).

31. J.-H. Kim, *et al.*, Protein inactivation in mycobacteria by controlled proteolysis and its
application to deplete the beta subunit of RNA polymerase. *Nucleic Acids Res.* **39**,
2210–2220 (2011).

32. E. O. Johnson, *et al.*, Large-scale chemical–genetics yields new M. tuberculosis inhibitor
classes. *Nature* **571**, 72–78 (2019).

33. J. M. Stokes, A. J. Lopatkin, M. A. Lobritz, J. J. Collins, Bacterial Metabolism and
Antibiotic Efficacy. *Cell Metab.* **30**, 251–259 (2019).

34. M. M. Bellerose, *et al.*, Distinct Bacterial Pathways Influence the Efficacy of Antibiotics
against Mycobacterium tuberculosis. *mSystems* **5** (2020).

35. M. Hafner, M. Niepel, M. Chung, P. K. Sorger, Growth rate inhibition metrics correct for
confounders in measuring sensitivity to cancer drugs. *Nat. Methods* **13**, 521–527
(2016).

36. A. S. G. Lee, A. S. M. Teo, S.-Y. Wong, Novel Mutations in ndh in Isoniazid-Resistant
Mycobacterium tuberculosis Isolates. *Antimicrob. Agents Chemother.* **45**, 2157–2159
(2001).

37. J.-H. Kim, *et al.*, A genetic strategy to identify targets for the development of drugs that
prevent bacterial persistence. *Proc. Natl. Acad. Sci.* **110**, 19095–19100 (2013).

38. J.-R. Wei, *et al.*, Depletion of antibiotic targets has widely varying effects on growth.
Proc. Natl. Acad. Sci. **108**, 4176–4181 (2011).

39. G. Giaever, *et al.*, Genomic profiling of drug sensitivities via induced haploinsufficiency.
Nat. Genet. **21**, 278–283 (1999).

40. E. J. Peterson, *et al.*, Path-seq identifies an essential mycolate remodeling program for
mycobacterial host adaptation. *Mol. Syst. Biol.* **15** (2019).

41. R. V. der Geize, *et al.*, A gene cluster encoding cholesterol catabolism in a soil
actinomycete provides insight into Mycobacterium tuberculosis survival in
macrophages. *Proc. Natl. Acad. Sci.* **104**, 1947–1952 (2007).

42.   A. H. Diacon, *et al.*, Early Bactericidal Activity of High-Dose Rifampin in Patients with
Pulmonary Tuberculosis Evidenced by Positive Sputum Smears. *Antimicrob. Agents*
Chemother. **51**, 2994–2996 (2007).

43.   E. Chigutsa, *et al.*, Impact of Nonlinear Interactions of Pharmacokinetics and MICs on
Sputum Bacillary Kill Rates as a Marker of Sterilizing Effect in Tuberculosis.
Antimicrob. Agents Chemother. **59**, 38–45 (2015).

44.   J. G. Pasipanodya, *et al.*, Serum Drug Concentrations Predictive of Pulmonary
Tuberculosis Outcomes. *J. Infect. Dis.* **208**, 1464–1473 (2013).

45.   M. Jankute, J. A. G. Cox, J. Harrison, G. S. Besra, Assembly of the Mycobacterial Cell Wall.
Annu. Rev. Microbiol. **69**, 405–423 (2015).

46.   M. Seidel, *et al.*, Identification of a Novel Arabinofuranosyltransferase AftB Involved in
a Terminal Step of Cell Wall Arabinan Biosynthesis in Corynebacterianeae, such as
Corynebacterium glutamicum and Mycobacterium tuberculosis. *J. Biol. Chem.* **282**,
14729–14740 (2007).

47.   V. Usha, *et al.*, Reconstruction of diaminopimelic acid biosynthesis allows
characterisation of Mycobacterium tuberculosis N -succinyl-L,L-diaminopimelic acid
desuccinylase. *Sci. Rep.* **6**, 1–10 (2016).

48.   M. Jankute, *et al.*, Disruption of Mycobacterial AftB Results in Complete Loss of
Terminal β(1 ⮕ 2) Arabinofuranose Residues of Lipoarabinomannan. *ACS Chem. Biol.*
12, 183–190 (2017).

49.   C. Lefebvre, *et al.*, HadD, a novel fatty acid synthase type II protein, is essential for
alpha- and epoxy-mycolic acid biosynthesis and mycobacterial fitness. *Sci. Rep.* **8**, 1–15
(2018).

50.   J. E. Griffin, *et al.*, Cholesterol Catabolism by Mycobacterium tuberculosis Requires
Transcriptional and Metabolic Adaptations. *Chem. Biol.* **19**, 218–227 (2012).

51.   M. Jackson, G. Stadthagen, B. Gicquel, Long-chain multiple methyl-branched fatty acid-
containing lipids of Mycobacterium tuberculosis: Biosynthesis, transport, regulation
and biological activities. *Tuberculosis* **87**, 78–86 (2007).

52.   M. Jain, *et al.*, Lipidomics reveals control of Mycobacterium tuberculosis virulence
lipids via metabolic coupling. *Proc. Natl. Acad. Sci.* **104**, 5133–5138 (2007).

53.   X. Yang, N. M. Nesbitt, E. Dubnau, I. Smith, N. S. Sampson, Cholesterol Metabolism
Increases the Metabolic Pool of Propionate in M. tuberculosis. *Biochemistry* **48**, 3819–
3821 (2009).

54.   S. B. Walters, *et al.*, The Mycobacterium tuberculosis PhoPR two-component system
regulates genes essential for virulence and complex lipid biosynthesis. *Mol. Microbiol.*
60, 312–330 (2006).

55.   J. G. Asensio, *et al.*, The Virulence-associated Two-component PhoP-PhoR System
Controls the Biosynthesis of Polyketide-derived Lipids in Mycobacterium tuberculosis.
J. Biol. Chem. **281**, 1313–1316 (2006).

56.   J. J. Baker, B. K. Johnson, R. B. Abramovitch, Slow growth of Mycobacterium
tuberculosis at acidic pH is regulated by phoPR and host-associated carbon sources.
Mol. Microbiol. **94**, 56–69 (2014).

57.   K. E. Stott, *et al.*, Pharmacokinetics of rifampicin in adult TB patients and healthy
volunteers: a systematic review and meta-analysis. *J. Antimicrob. Chemother.* **73**,
2305–2313 (2018).

58.   M. Gengenbacher, *et al.*, Tissue Distribution of Doxycycline in Animal Models of
Tuberculosis. *Antimicrob. Agents Chemother.* **64** (2020).

59.   N. L. Wengenack, H. M. Hoard, F. Rusnak, Isoniazid Oxidation by Mycobacterium
tuberculosis KatG: A Role for Superoxide Which Correlates with Isoniazid
Susceptibility. *J. Am. Chem. Soc.* **121**, 9748–9749 (1999).

60.   M. B. McNeil, S. Chettiar, D. Awasthi, T. Parish, Cell wall inhibitors increase the
accumulation of rifampicin in Mycobacterium tuberculosis. *Access Microbiol.* **1** (2019).

61.   M. S. Pavelka, W. R. Jacobs, Biosynthesis of diaminopimelate, the precursor of lysine
and a component of peptidoglycan, is an essential function of Mycobacterium
smegmatis. *J. Bacteriol.* **178**, 6496–6507 (1996).

62.   J. Lee, T. Repasy, K. Papavinasasundaram, C. Sassetti, H. Kornfeld, Mycobacterium
tuberculosis Induces an Atypical Cell Death Mode to Escape from Infected
Macrophages. *PLOS ONE* **6**, e18367 (2011).

63.   T. Metsalu, J. Vilo, ClustVis: a web tool for visualizing clustering of multivariate data
using Principal Component Analysis and heatmap. *Nucleic Acids Res.* **43**, W566–W570
(2015).

64.   V. Matyash, G. Liebisch, T. V. Kurzchalia, A. Shevchenko, D. Schwudke, Lipid extraction
by methyl-tert-butyl ether for high-throughput lipidomics. *J. Lipid Res.* **49**, 1137–1146
(2008).

849