Low mammalian species richness is associated with Kyasanur Forest Disease outbreak risk in deforested
landscapes in the Western Ghats, India

Michael G. Walsh[1,2,3,4] Rashmi Bhat[3,4], Venkatesh Nagarajan-Radha[5], Prakash Narayanan [3,4], Navya
Vyas[3,4], Shailendra Sawleshwarkar[3,4,6], Chiranjay Mukhopadhyay[7,8]

[1]The University of Sydney, Faculty of Medicine and Health, School of Public Health, Camperdown, New
South Wales, Australia, [2]The University of Sydney, Faculty of Medicine and Health, Sydney Institute for
Infectious Diseases, Westmead, New South Wales, Australia, [3]One Health Centre, The Prasanna School
of Public Health, Manipal Academy of Higher Education, Manipal, Karnataka, India, [4]The Prasanna School
of Public Health, Manipal Academy of Higher Education, Manipal, Karnataka, India, [5]Behaviour Ecology
and Evolution Lab, The University of Sydney, Faculty of Science, School of Life and Environmental
Sciences, Camperdown, New South Wales, Australia, [6]The University of Sydney, Faculty of Medicine and
Health, Westmead Clinical School, Westmead, New South Wales, Australia, [7]Department of
Microbiology, Kasturba Medical College, Manipal Academy of Higher Education, Manipal, Karnataka,
India, [8]Centre for Emerging and Tropical Diseases, Kasturba Medical College, Manipal Academy of Higher
Education, Manipal, Karnataka, India

*Address correspondence to:
Michael Walsh, PhD, MPH
Senior Lecturer, Infectious Diseases Epidemiology
Sydney Institute for Infectious Diseases
Sydney School of Public Health
The University of Sydney
176 Hawksbury Road
Westmead NSW 2145 Australia
thegowda@gmail.com
michael.walsh1@sydney.edu.au

Highlights

• Local mammalian species richness is estimated across the entire Western Ghats region
• Low species richness is associated with high KFDV risk in deforested landscapes
• This work identifies key landscapes for wildlife disease surveillance development

Abstract

Kyasanur forest disease virus (KFDV) is a rapidly expanding tick-borne zoonotic virus with natural foci in
the forested region of the Western Ghats of South India. The Western Ghats is one of the world's most
important biodiversity hotspots and, like many such areas of high biodiversity, is under significant
pressure from anthropogenic landscape change. The current study sought to quantify mammalian
species richness using ensemble models of the distributions of a sample of species extant in the Western
Ghats and to explore its association with KFDV outbreaks, as well as the modifying effects of
deforestation on this association. Species richness was quantified as a composite of individual species'
distributions, as derived from ensembles of boosted regression tree, random forest, and generalised
additive models. Species richness was further adjusted for the potential biotic constraints of sympatric
species. Both species richness and forest loss demonstrated strong positive associations with KFDV
outbreaks, however forest loss substantially modified the association between species richness and
outbreaks. High species richness was associated with increased KFDV risk but only in areas of low forest
loss. In contrast, lower species richness was associated with increased KFDV risk in areas of greater
forest loss. This relationship persisted when species richness was adjusted for biotic constraints at the
taluk-level. In addition, the taluk-level species abundances of three monkey species (*Macaca radiata*,
Semnopithecus hypoleucus, and *Semnopithecus priam*) were also associated with outbreaks. These
results suggest that increased monitoring of wildlife in areas of significant habitat fragmentation may
add considerably to critical knowledge gaps in KFDV epidemiology and infection ecology and should be
incorporated into novel One Health surveillance development for the region. In addition, the inclusion of
some primate species as sentinels of KFDV circulation into general wildlife surveillance architecture may
add further value.

Keywords: Kyasanur Forest Disease, biodiversity, deforestation, zoonoses, Western Ghats, tick-borne
disease

Introduction

The sharing of space between wildlife and humans can be disruptive to both animals and people

particularly in the context of altered habitat. Human pressure on wildlife influences species' community

composition, population movement and density, resource provisioning[1,2], and eco-immunology[3],

which can subsequently alter pathogen circulation among wildlife hosts as well as introduce conduits

that allow the movement of pathogens from reservoir hosts to novel human hosts[4], with potentially

catastrophic effects[5–7]. As such, the growing wildlife-human interface requires both improved

surveillance of pathogen circulation among wildlife and a greater understanding of the impact of

anthropogenic landscape change on wildlife communities, their pathogens, and mechanisms for

potential spillover of novel pathogens to humans. Importantly, as new wildlife-human interfaces emerge

in close proximity to major transportation hubs, the potential for high-impact spillovers with the

capacity for rapid regional or global dissemination is increasing[8].

88   Currently, there is a lack of zoonosis surveillance in wildlife globally, but the absence is

particularly noteworthy in the world's biodiversity hotspots that are undergoing rapid anthropogenic

change. The Western Ghats region of South India is one of the world's most important biodiversity

hotspots[9] and also experiences significant pressure in the form of landscape change due to

deforestation, commercial agricultural exploitation, and urbanisation[10]. The monsoon rainforests of

this region support approximately 500 bird species, 225 reptile species, 219 amphibian species, and 133

mammal species[11,12]. Nevertheless, only 10% of the region's 160,000 km^2 area is formally

protected[11]. As anthropogenic pressure increases in the region, conflict between many of these

species, especially mammals, and humans has grown [13,14], facilitating exposure to novel pathogens.

The general lack of wildlife pathogen surveillance across this rapidly changing region is compounded by

an even greater absence of work examining the influence of community ecology on the processes of

pathogen circulation and spillover events. The emerging tick-borne zoonotic virus, Kyasanur Forest

disease virus (KFDV), simultaneously represents 1) a critical public health priority in the Western Ghats

states because of its recent rapid expansion and the high mortality (2% - 10%) associated with its

neurological and haematological complications [15–18], and 2) a model system for understanding viral

spillover because of the extensive wildlife-human interface across the region[8]. Kyasanur Forest disease

virus is a Flavivirus transmitted by several tick species, while the primary vector is the forest tick,

Haemaphysalis spinigera. This species is found in high relative abundance in the region, has high viral

prevalence, and feeds on many taxa of mammalian hosts including humans[19–21]. Humans typically

are exposed to these ticks in the anthropogenic ecotones of forest fringe[22]. Outbreaks of KFDV have

expanded from a single district in the state of Karnataka in the decade following the virus' first

identification in 1957 to an extensive region now comprising five states across South India[15,22].

Recent work has demonstrated a strong direct association between the expansion of KFDV outbreaks

and the loss of native forest[23,24], which was further supported by phylogeographic analyses[25]. The

study that reported the KFDV outbreak association with forest loss also noted that areas of high

mammalian species richness presented greater risk[23]. While this previous study's assessment of forest

loss was of high quality, species richness was assessed more crudely using the IUCN-derived species'

range for each mammal species present across the region. As such, variation of species' distributions

within those ranges was not incorporated into the assessment of species richness, nor was there any

assessment of potential interaction between sympatric species. Additionally, the associations between

KFDV and individual species' abundance and relative abundance were not previously explored. It is

anticipated that further investigation of these aspects of mammalian biogeography and community

ecology in KFDV outbreak hotspots will add considerable insight into the epidemiology of KFDV spillover

while also providing an ecological evidence-base for developing municipal wildlife surveillance

infrastructure at the taluk level (sub-district) across the Western Ghats states. The incorporation of

wildlife monitoring and sampling into KFDV surveillance is particularly important since the investigation

of infection in wildlife hosts has fallen off dramatically in recent decades, prior to which some

mammalian susceptibility had been identified via serology but reservoir host competence was not

established[15]. This decline in wildlife surveillance has resulted in a considerable knowledge gap in the

fundamental infection ecology of KFDV, while also leaving any new efforts at building surveillance

infrastructure uninformed with respect to what species should be sampled and where they should be

sampled. Given the extensive forest loss in the region, delineating potential host communities across the

spectrum of habitat fragmentation will be important in understanding how risk of spillover is modulated

by wildlife communities. For example, as mentioned, the extent of viral competence is not known for

most mammals of the Western Ghats and therefore identifying key reservoir hosts, or distinguishing

between maintenance and amplification hosts, is not currently possible. As such, we remain largely

ignorant of the importance of generalist species, which typically are more resilient to anthropogenic

pressure, frequently dominate fragmented landscapes as biodiversity is lost, and often host zoonotic

pathogens [26–28]. Conversely, the extent to which greater species richness in less fragmented

landscapes may buffer against viral transmission (or against tick dispersal and feeding success) is also

unknown. Therefore, there is an urgent need for targeted surveillance mechanisms to sample wildlife

across heterogenous landscapes to fill these critical knowledge gaps and thereby ultimately develop a

more sound approach to the control and prevention of this rapidly expanding tick-borne arbovirus.

The specific aims of the current study were as follows. First, we sought to estimate the species

distributions of a sample of extant mammals of the Western Ghats (including endemic and non-endemic

species) and use these to construct a more representative metric of species richness across the region.

Second, we sought to adjust overall species richness, as well as individual species' abundance and

relative abundance for the biotic constraints of sympatry. Third, we sought to interrogate the

association between species richness and KFDV outbreaks and the modification of this association by

forest loss to develop much needed landscape epidemiology targets for improved surveillance of this

expanding zoonotic arbovirus.

Materials and Methods

Data Sources

The Global Biodiversity Information Facility was used to obtain all direct observations of extant

mammals in the Western Ghats[12] and Peninsular India between 1 January, 2010 and 1 November,

2020 (www.gbif.org). A total of 2826 mammals were observed over this period[29]. This comprised a

total of 99 species observed across the region. However, habitat suitability was estimated only for those

species with a sample size of at least 30 individuals, for a total of 24 species.

The KFDV outbreaks used in this investigation have been described in detail previously[23].

Briefly, two independent sources of outbreaks reported between 1 January, 2012 and 30 June, 2019

were used for model training and testing, respectively. The training data (47 outbreaks) were sourced

from the ProMED-mail[30] electronic surveillance system and the testing data (39 outbreaks) comprised

an independent set of laboratory-confirmed outbreaks described in the literature[23]. This allowed for

external validation of KFDV outbreak models rather than partitioning data from a single source into

training and testing.

Mean annual precipitation and mean annual temperature were obtained from the WorldClim

Global Climate database at a resolution of 30 arc seconds (~ 1 km) [31]. The rasters derived from the

WorldClim database were mean measurements between the period 1950 to 2000, and thus represent

estimates of climate rather than weather. Some regions represented by the WorldClim database have

sparse weather stations contributing to climate estimates. However, India's extensive network of

weather stations make its local contributions to decadal climate interpolation more representative than

many other large countries[32].

The Priestley-Taylor α coefficient (P-Tα) is the ratio of actual evapotranspiration to potential

evapotranspiration and was included here as a metric for water-soil balance[33,34]. In contrast to solar

energy input alone, the P-Tα represents water availability in the soil as a function of the local

vegetation's water requirements and is therefore a more robust estimate of soil-water balance. A 30 arc

seconds resolution raster of P-Tα was obtained from the Consultative Group for International

Agricultural Research (CGIAR) Consortium for Spatial Information. The ratio ranges from 0 (extreme

water stress) to 1 (no water stress)[35].

Forest cover was first acquired from the Global Land Cover Climatology MODIS-based data

product, which corrects temporally aggregated 10-year land cover types by weighting each land cover

type by their annual uncertainty and then validates the weighted composite against the System for

Terrestrial Ecosystem Parameterization[36]. The product used here for forest cover represents the

period 2001-2010 at a resolution of 15 arc seconds (~ 500 m).

Landsat data compiled at 1 arc second (~ 30 m) resolution by the Global Forest Change project

was used to quantify forest loss between 2000 and 2012[37]. Landsat imagery was processed using a

stratified random sampling validation procedure, with 99.6% accuracy in global settings and 99.5%

accuracy in tropical settings[37]. Individual tiles were merged across the region of the Western Ghats

and a new quantile raster constructed to represent deciles of forest loss.

An established validated metric for the human footprint (HFP) was used to control for spatial

background sampling as described below in the analysis section. The metric was acquired from the

Socioeconomic Data and Applications Center (SEDAC) repository[38] at a resolution of 30 arc seconds

and has been described in detail previously[39]. Briefly, HFP comprises 2 levels of classification. First, an

index of human influence was determined based on eight domains: 1) population density, distance to 2)

roads, 3) railway lines, 4) navigable rivers, and 5) coastlines, 6) degree of night-time artificial light, 7)

urban vs rural location, and 8) land cover. These domains were then scored to derive the human

influence index (HII), ranging from 0 (no human impact) to 64 (greatest human impact). The ratio of the

range of minimum and maximum HII in the local terrestrial biome to the range of minimum and

maximum HII across all biomes represents the final HFP metric and is expressed as a percentage[39].

Because KFDV outbreaks disproportionally affect marginalised communities with limited access

to health care, this investigation adjusted for potential reporting bias using a measure of the distribution

of health system performance as an indication of the local capacity to detect outbreaks (see modelling

description below). The infant mortality ratio (IMR) was chosen as a proxy for health system

performance because this has been verified as a robust indicator of health infrastructure and health

system performance and used extensively to compare health service delivery[40,41]. The IMR strongly

correlates with disability-adjusted life expectancy (DALE), the Human Development Index (HDI), and the

Inequality-Adjusted Human Development Index (IHDI). As such, the IMR is an important indicator of

structural issues that affect health care access and delivery, such as economic development, general

living conditions, social well-being, and the quality of the environment[40,42]. The raster of the IMR was

obtained from SEDAC[43].

Data Analysis

Species distribution modelling. The landscape suitability of each of the 24 mammal species was

estimated using an ensemble approach comprising two machine learning methods (boosted regression

trees (BRT) and random forests (RF)) and generalised additive models (GAM)). The machine learning

frameworks, BRT and RF, are powerful approaches to improving standard decision trees by iteratively

growing many trees and amalgamating the results. As the algorithms partition the data space according

to rules that optimise homogeneity among predictors and a response (e.g. species presence), many

decision trees are grown and combined, resulting in optimised decision trees that reduce overfitting and

can capture complex interactions between the predictors [44–47]. There are two key differences in

these machine learning approaches. First, for each decision tree partition in RF, only a random subset of

predictors is selected from the set of all predictors, which decorrelates the trees and reduces overfitting.

Second, rather than decorrelating trees based on the sampling of subsets as with RF, BRT instead

reduces overfitting by growing trees sequentially and learning from previously grown trees. In contrast,

the GAM framework allows for nonlinear relationships between outcomes and covariates by fitting

multiple basis functions for smoothed covariates [48,49]. Five-fold cross-validation was used to fit each

model under the three distinct modelling frameworks (BRT, RF, and GAM). Mean annual temperature, P-

T α, and forest cover were included as landscape features at 30 arc seconds resolution in all models. For

each species, each of the three models describing their landscape suitability was evaluated according to

performance, based on the area under the receiver operating characteristic curve (AUC), and fit, based

on the model deviance. Subsequently, an ensemble landscape suitability was computed from the three

model classes using their weighted mean, with weights based on AUC [50]. To adjust for potential spatial

sampling bias in the GBIF database, background points were sampled proportional to the human

footprint as a proxy for landscape accessibility. The estimates of each species' distribution as derived

from these ensembles were then summed across all species to estimate local species richness across the

Western Ghats region. This estimate as well as an additional estimate adjusted for the biotic constraints

of sympatric species (see description below) were subsequently used as more appropriate metrics for

mammalian species richness and the evaluation of its influence on KFDV outbreaks. As an additional

sensitivity analysis to test whether associations with species richness may be influenced by the limited

regional sample of mammalian species used to estimate species richness, an alternate India-wide

sample (number of observations = 5833, number of species = 30) was used to re-estimate individual

species habitat suitability and then species richness. The sdm package[50] in the R platform[51] was

used for fitting each model and the derivation of the three-model ensembles to each species.

Point process modelling. The KFDV outbreaks were fitted as an inhomogeneous Poisson point

process[52,53]. Under this model framework the spatial dependence of the outbreaks' distribution can

be determined and evaluated with respect to specific landscape features. The background points used in

these models were sampled proportional to IMR, as described above, to control for potential outbreak

reporting bias. Previous work demonstrated a strong association between KFDV outbreaks and both

forest loss and mammalian species richness[23]. However, the association with the latter was 1) based

on a crude representation of species richness that did not account for local heterogeneity of individual

species' distributions and 2) could not be used to evaluate the potential effect modification of species

richness by forest loss. The current investigation provides this critical missing component by

incorporating a more suitable metric of species richness based on the individual estimated species'

distributions described above and subsequently interrogating the interaction between species richness

and forest loss. In addition, to account for the important climate features that were incorporated in the

previous work, precipitation and temperature were also included in the current work. Finally, the

possibility of confounding of the associations between KFDV outbreaks and species richness and forest

loss by alternative forms of human influence in the landscape was assessed by including HFP as a

covariate in an additional model. Therefore, the suite of multiple inhomogeneous Poisson models

comprised the newly-quantified mammalian species richness metric described above, forest loss, an

interaction term for species richness and forest loss, mean annual precipitation, mean annual

temperature, and HFP all aggregated up to 2.5 arc minutes. The landscape features included in the

multiple inhomogeneous Poisson models were not highly correlated so multicollinearity in the models

was not of concern (all values of the Pearson's r were < 0.5). Model fit was assessed using the Akaike

information criterion (AIC) and model performance was tested against an independent, laboratory-

confirmed set of outbreak data, as described above. Performance was evaluated using the AUC. The

spatstat R package was used to fit the point process models[54].

Taluk-level modelling. Species richness was further adjusted for, and individual species abundance and

relative abundance were computed based on, the potential modulating effects of sympatric species at

the scale of the taluk. Taluks are 3rd-level, subdistrict administration units in India and are small enough

to reasonably represent shared space between sympatric species, while simultaneously large enough to

delineate the minimal municipal infrastructure (i.e. delineating the most local focus) used in the

Western Ghats states for the implementation of animal and human disease surveillance. Specifically,

biotic constraints were applied to the estimates of each species distribution according to the spatially-

explicit species assemblage modelling (SESAM) framework[55–57]. Under this framework, first the

landscape suitability for each species was estimated according to the ensemble method described

above. Second, species richness was calculated as the sum of the individual species distributions, again

using the method described above. Third, the biotic constraint was applied whereby each species is

evaluated with respect to all other species present within each taluk via the probability ranking

rule[56,58] to determine whether a given species should be retained within, or excluded from, each

taluk "community". The third step ranks each species from highest to lowest predicted habitat suitability

based on each species' suitability obtained in the first step. Then, species are selected for inclusion

beginning with the species with highest suitability and proceeding down the catalogue of ranked species

until the sum of selected species is equal to the expected species richness value for each location

calculated in the second step. This provides an estimate of each species' presence (or absence) at the

taluk level given the potential biotic constraints of the other species within the taluk. For those species

thusly identified as present within the taluk community, all 1 km^2 pixels exceeding the true skill statistic

(TSS)[59] of their ensemble landscape suitability estimate were designated present (1 = present, 0 =

otherwise) and summed across all pixels within the taluk. This yielded a taluk-level estimate of species

abundance for each species under consideration. Similarly, the taluk-level relative abundance for each

species was quantified by dividing each individual species abundance by the total abundance of all

species identified as present within each taluk under the SESAM framework. Finally, a new estimate of

species richness was computed by summing all the species identified as present within each taluk under

the SESAM framework. This systematic approach thus yielded taluk-level community estimates of each

species' abundance and relative abundance, as well as an updated estimate of species richness, all

adjusted for the biotic constraints of sympatric species. The SESAM framework, including the probability

ranking rule was implemented using the ecospat package in R[57].

Integrated nested Laplace approximation (INLA) models[60] were used to estimate the

association between KFDV outbreaks and the biotically constrained estimate of mammalian species

richness and its interaction with forest loss at the level of the taluk, as well as associations between

KFDV outbreaks and individual species abundance and relative abundance to identify potential species

that may be good candidates for initial wildlife surveillance. While the latter may represent species that

are important as reservoir hosts, we did not attempt to assess species' roles as hosts in the current

study since we did not measure their competence for KFDV. Instead, individual species associations

were investigated simply to identify whether certain species may be useful sentinels of KFDV outbreaks.

The binomial likelihood family was used for the INLA models with spatial autocorrelation estimated

using random effects with Besag–York–Mollie priors. Under the binomial family, taluks were modelled

as either having experienced a KFDV outbreak under the period of study (outbreak positive) or not

(outbreak negative). This approach was taken since the majority of taluks in the region did not

experience outbreaks, and, among those that did, most only experienced one or two outbreaks.

Nevertheless, to evaluate the effect of different distribution families, we also fitted these models with

the zero-inflated Poisson family as a sensitivity analysis. Since the KFDV outbreaks were necessarily

aggregated by taluk, to increase the outbreak sample size available for taluk-level modelling, the

independent datasets described above for the point process models (PPMs) were combined for the INLA

models. Since the primary objective was estimation rather than prediction so as to infer associations

between KFDV outbreaks and species richness, this was deemed an acceptable trade-off. The fit of INLA

models was evaluated using the Watanabe-Akaike information criterion (WAIC). The INLA models were

fit using the inla package in R (www.r-inla.org) [60].

Results

The distribution of mammalian species richness, estimated as the sum of the sample of individual

species' distributions, is presented in Figure 1 juxtaposed with the distribution of KFDV outbreaks. The

individual species distribution models generally performed well (Supplementary material, S1 Table 1).

The distribution of each species' landscape suitability is presented in S2 Figure 1 and S3 Figure 2. Both

forest loss and the new quantification of mammalian species richness based on individual species'

landscape suitability alone were very strongly associated with KFDV outbreaks at local scale (Table 1).

Strong associations were apparent whether these features were considered alone in the crude bivariate

PPMs (Table 1A), or together in the multiple PPMs (Table 1B, 1C, and 1D). Critically, while species

richness demonstrated a similar association with KFDV outbreaks to that shown previously[23], the

current investigation identified significant and substantial interaction between species richness and

forest loss. As such, increasing mammalian species richness was crudely associated with increasing KFDV

outbreak occurrence. However, when species richness was considered in concert with forest loss, the

KFDV association with the former was modified by the latter (coefficient = -0.41). This relationship was

maintained in the final model (Table 1C), which incorporated the climate features and was a markedly

better fit (AIC = -36.34 vs. -22.80) although performance was slightly diminished (AUC = 0.74 vs. 0.78).

Finally, confounding of these associations by HFP was not supported since the regression coefficients

were unaltered by the inclusion of HFP, HFP itself was not associated with outbreaks, and the model

overall was a poorer fit to the data (Table 1D). The distribution of KFDV outbreak risk across the Western

Ghats based on the best fitting model (Table 1C) is presented in Figure 2. The individual species

distribution models and projected suitability using the alternate country-wide sample are presented in

S4 Table 2 and S5 Figure3 and S6 Figure 4. There was no marked difference in the strength or direction

of the association between KFDV outbreak risk and species richness or its interaction with forest loss in

the sensitivity analysis using the India-wide sample, however the model based on the latter

demonstrated both poorer fit and performance (S7 Table 3, S8 Figure 5), so the original model based on

the regional sample was retained as the primary model for this analysis.

The newly quantified metric for taluk-level species richness based on the SESAM framework is presented

in Figure 3. In comparison to the best-fitting high-resolution PPM model (Table 1C), the taluk-level INLA

model showed the same strong interaction between forest loss and mammal species richness, even as

the latter was adjusted for biotic constraint under the SESAM framework and scaled up to taluk (Table 2,

Figure 4). Further sensitivity analysis showed that these associations all persisted when the distribution

family was specified as zero-inflated Poisson (S9 Table 4 and S10 Figure 6), although temperature was no

longer associated with outbreak occurrence. Finally, as with the PPMs described above, the associations

persisted (S11 Table 5) when species richness was estimated based on the alternate India-wide sample

(S12 Figure 7) although the model overall was a poorer fit. Based on taluk-level outbreak risk (Figure 4),

we identified the 50th percentile of risk where deforestation and species richness optimally converge for

targeted selection for a state-administered (e.g. Karnataka) animal surveillance program (Figure 5). In

addition to the broad metric of species richness, we also explored the associations between KFDV

outbreaks and each individual species' abundance and relative abundance to determine if any particular

species may prove useful as a sentinel surveillance target. Despite several species demonstrating

significant individual associations with KFDV occurrence at fine scale based on their estimated landscape

suitability (S13 Table 6), most species did not demonstrate associations with outbreaks at the level of

the taluk (S14 Table 7). However, there were a few notable exceptions. Species abundance of the

primates *Macaca radiata*, *Semnopithecus hypoleucos* and *Semnopithecus priam*, were all associated

with increased KFDV occurrence at the taluk level. Species relative abundance was also associated with

KFDV outbreaks for *S. hypoleucos*, but not for any other primate species. The distributions of these

species' abundances and relative abundances per taluk are presented in S15 Figure 8. Finally, both the

abundance and relative abundance of *Panthera pardus*, and the abundance of *Axis axis*, were also

associated with increased KFDV outbreak occurrence.

Discussion

This is the first study to attempt a general estimate of mammalian species richness across the Western

Ghats region based on ensemble modelling of the landscape suitability of several extant species, and to

extend this by adding biotic constraints of sympatric species to the estimates of species richness, as well

as individual species abundance and relative abundance. This is also the first study to demonstrate

negative effect modification of the association between mammalian species richness and KFDV outbreak

occurrence by forest loss such that higher species richness was associated with increased KFDV outbreak

risk where deforestation was minimal, but in areas of increasing forest loss lower species richness was

associated with increased risk. This association was identified at both fine scale and at the level of the

taluk. Finally, three monkey species were identified (*M. radiata*, *S. hypoleucus*, and *S. priam*) whose

taluk-level species abundances were indicators of outbreaks. Taken together, these findings have

identified important interaction between landscape degradation and mammalian species richness that

may influence the ecology of this tick-borne virus in its natural foci. The fact that this relationship held at

the taluk level even after controlling for biotic constraints indicates that enhanced monitoring of wildlife

and human populations at the level of the taluk, which is well-suited to the municipal infrastructure of

current human and animal health systems in the Western Ghats states, may offer promise as the focus

of novel One Health surveillance development. In addition, the incorporation of some primate species as

sentinels of KFDV circulation into general wildlife surveillance may add further value to such a One

Health program.

Previous work identified increased risk of KFDV outbreaks associated with both forest loss and

high mammalian species richness[23]. However, the latter association was difficult to interpret due to

the fairly crude measure of species richness, which was simply a composite of IUCN mammal species'

ranges across the region. Moreover, this work was not able to assess interaction between forest loss

and mammalian biodiversity. The positive association between KFDV outbreak risk and species richness

shown in that study was reasonable given the expectation of greater exposure to potential KFDV hosts

and their tick vectors following increasing incursion into areas of high species richness. Nevertheless, it

is also of particular interest to identify how habitat loss may affect the community ecology of

mammalian hosts in ways that could influence KFDV spillover to humans. Therefore, better metrics of

species richness were required across the region to more appropriately evaluate the modifying effect of

forest loss on the association between species richness and KFDV outbreaks. The results presented in

this current investigation indicated that the relationship between species richness and forest loss is

indeed more nuanced. As shown previously[23], increased species richness was associated with

increased KFDV outbreak risk but in the current study this relationship only held in landscapes with

minimal forest loss. In contrast, as forest loss increased, lower species richness was associated with

increased risk. While this finding allows no definitive assertion regarding the nature of KFDV circulation

among mammalian hosts, such as the relative importance of the dilution effect to transmission

dynamics[61], it does provide important evidence that KFDV circulation among reservoir hosts persists,

and spillover to humans continues, in human-altered landscapes exhibiting lower biodiversity, which is

consistent with previous evidence showing clear pathways from habitat fragmentation and biodiversity

loss to increased spillover[1,62]. Despite the current study's inability to make specific claims about the

role of any individual species in the viral maintenance and infection ecology of KFDV, the findings do

suggest that, in general, species resilient to anthropogenic landscape change may be most relevant to

the expansion of KFDV spillover across the region. This finding closely parallels recent work

demonstrating similar relationships more globally for other zoonotic pathogens[26,27]. Moreover, the

work may help to address a critical surveillance gap currently impeding a fuller understanding of KFDV

infection ecology. Following the initial identification of KFDV in 1957, there was a rapid but limited flurry

of research attempting to identify wildlife reservoirs for KFDV in the Bandipur Forest range[15]. Several

mammalian species were identified as hosts, but these studies were not definitive with respect to KFDV

infection ecology as many relied on serology, rather than measuring host competence, and many of the

studies employed limited sampling strategies[15]. Subsequently, wildlife investigations all but stopped

by the early 1980s before KFDV reservoirs could be established. Therefore, the current study could have

important implications for reinstating widespread wildlife surveillance of KFDV in the Western Ghats

with a focus on synanthropic species.

It is important to caution against interpretation of these findings to infer specific dynamics of

infection ecology at either the level of the community or individual species because infection

competence was not assessed in any of the mammals under investigation nor was interspecific

interaction among species directly observed at species' locations. However, the results do provide some

of the most actionable evidence to date for targeted, landscape-designed wildlife surveillance across the

region. Specifically, these findings can help to frame a landscape-based cluster sampling design for the

monitoring of wildlife distribution and movement in response to human induced landscape change in

concert with pathogen surveillance in animals and humans (Figure 5). Critically, the fact that this study's

findings were consistent at the level of the taluk indicates that the existing municipal infrastructure for

animal and human health can be used as the necessary scaffolding for novel One Health surveillance

architecture and service delivery. Moreover, the taluk-level abundance of bonnet macaques (*M. radiata*)

and grey langurs (*Semnopithecus spp.*), which have previously been identified as highly susceptible to

KFDV[63–65], were also associated with outbreaks. As mentioned above, the current study can make no

claims regarding the roles of individual species in the infection ecology of KFDV, but the associations

with these primate species' abundance within taluks indicates that these species may provide useful

sentinels under a One Health surveillance framework. In addition, leopard (*P. pardus*) abundance and

relative abundance and chital (*A. axis*) abundance were also associated with outbreak risk. While neither

of these species have been shown to be susceptible to KFDV, both are known to host KFDV-relevant

ticks and distribute these vectors in the landscape[66,67]. Therefore, their associations with increased

KFDV outbreak occurrence in the current study merits a closer look at the role these species could play

as tick distributors, particularly since both of these species are highly adaptive to anthropogenic

landscapes. As such leopards and chital could have repercussions for human exposure to KFDV vectors,

their unlikely role as important KFDV hosts notwithstanding.

440   Further to the cautionary interpretation around KFDV infection ecology encouraged above,

some further description of this study's limitations is provided. First, as previously mentioned direct

observation of interspecific interaction was not possible under the current investigation, which

precludes the ability to make specific claims about the true nature and influence of interaction among

communities at various scales across the region. The findings reported here will need to be verified by

systematically conducted field investigations of directly observed interspecific interaction at multiple

scales. Second, sufficient observations were not available for each extant mammal species of the

Western Ghats and so the new quantification of species richness presented here, although much

improved over previous estimates, is based on only a sample of Western Ghats mammals and therefore

remains a proxy for true species richness. Third, spatial biases may affect the distribution of species

observations, as well as the reporting of KFDV outbreaks, across the region. To control for these biases,

respectively, background points were not selected randomly but instead selected proportional to the

human footprint as a measure of accessibility and IMR as a measure of health system performance and

access. Finally, this study provides no description of the roles of individual species as reservoirs, whether

operating as maintenance, amplification, or bridging hosts, since neither infection competence nor

infection susceptibility was measured in these species in the current study. Rather, individual species'

landscape suitability was used to 1) calculate species richness and evaluate this as a landscape feature of

importance to the distribution of KFDV outbreak occurrence, particularly with respect to forest loss, and

2) identify individual species that may serve as useful sentinels in the development of wildlife

surveillance mechanisms. Therefore, rather than defining how mammalian community ecology

determines KFDV infection ecology, the current work delineates optimal landscape targets for

developing the surveillance instruments necessary to appropriately monitor and sample wildlife so that

the necessary community ecology and infection ecology data that are needed can be generated.

Conclusions

This study provides novel metrics of mammalian species richness in the Western Ghats, one of the

world's critical biodiversity hotspots, and demonstrates the influence of this landscape feature and its

interaction with deforestation on the expansion of one of India's most important emerging zoonotic

arboviruses. The insight gained can provide municipally-directed targets for landscape-based One Health

surveillance of animals and humans in the face of anthropogenic landscape change, which can begin to

provide a definitive understanding of KFDV infection ecology while simultaneously identifying the

wildlife-human interfaces most vulnerable to pathogen spillover and which may yield the highest benefit

from tailored intervention.

Declarations of interest: none.

Funding: This research did not receive any specific grant from funding agencies in the public,
commercial, or not-for-profit sectors.

Author contributions: Conceptualization: Michael Walsh, Rashmi Bhat, Venkatesh Nagarajan-Radha,
Shailendra Sawleshwarkar, Chiranjay Mukhopadhyay; Data curation: Michael Walsh; Formal analysis:
Michael Walsh; Investigation: Michael Walsh, Rashmi Bhat, Venkatesh Nagarajan-Radha, Prakash
Narayanan, Navya Vyas, Shailendra Sawleshwarkar, Chiranjay Mukhopadhyay; Methodology: Michael
Walsh, Rashmi Bhat, Venkatesh Nagarajan-Radha, Shailendra Sawleshwarkar; Supervision: Michael
Walsh, Chiranjay Mukhopadhyay; Validation: Michael Walsh, Rashmi Bhat, Venkatesh Nagarajan-Radha,
Shailendra Sawleshwarkar, Chiranjay Mukhopadhyay; Visualization: Michael Walsh; Writing - original

draft: Michael Walsh, Rashmi Bhat, Venkatesh Nagarajan-Radha, Prakash Narayanan, Navya Vyas,
Shailendra Sawleshwarkar, Chiranjay Mukhopadhyay; Writing - Michael Walsh, Rashmi Bhat, Venkatesh
Nagarajan-Radha, Prakash Narayanan, Navya Vyas, Shailendra Sawleshwarkar, Chiranjay
Mukhopadhyay.

512   References

513   [1]   C.L. Faust, H.I. McCallum, L.S.P. Bloomfield, N.L. Gottdenker, T.R. Gillespie, C.J. Torney, A.P.
514   Dobson, R.K. Plowright, Pathogen spillover during land conversion, Ecol. Lett. 21 (2018) 471–483.
515   https://doi.org/10.1111/ele.12904.

516   [2]   R.K. Plowright, C.R. Parrish, H. McCallum, P.J. Hudson, A.I. Ko, A.L. Graham, J.O. Lloyd-Smith,
517   Pathways to zoonotic spillover, Nat. Rev. Microbiol. 15 (2017) 502–510.
518   https://doi.org/10.1038/nrmicro.2017.45.

519   [3]   D.J. Becker, G.F. Albery, M.K. Kessler, T.J. Lunn, C.A. Falvo, G. Czirják, L.B. Martin, R.K. Plowright,
520   Macroimmunology: The drivers and consequences of spatial patterns in wildlife immune defence,
521   J. Anim. Ecol. 89 (2020) 972–995. https://doi.org/10.1111/1365-2656.13166.

522   [4]   B. Borremans, C. Faust, K.R. Manlove, S.H. Sokolow, J.O. Lloyd-Smith, Cross-species pathogen
523   spillover across ecosystem boundaries: Mechanisms and theory, Philos. Trans. R. Soc. B Biol. Sci.
524   374 (2019). https://doi.org/10.1098/rstb.2018.0344.

525   [5]   L.F. Wang, B.T. Eaton, Bats, civets and the emergence of SARS, Curr. Top. Microbiol. Immunol.
526   315 (2007) 325–344. https://doi.org/10.1007/978-3-540-70962-6_13.

527   [6]   C.E.M. Coltart, B. Lindsey, I. Ghinai, A.M. Johnson, D.L. Heymann, The Ebola outbreak, 2013–
528   2016: Old lessons for new epidemics, Philos. Trans. R. Soc. B Biol. Sci. 372 (2017).
529   https://doi.org/10.1098/rstb.2016.0297.

530   [7]   K.G. Andersen, A. Rambaut, W.I. Lipkin, E.C. Holmes, R.F. Garry, The proximal origin of SARS-CoV-
531   2, Nat. Med. 26 (2020) 450–452. https://doi.org/10.1038/s41591-020-0820-9.

532   [8]   M.G. Walsh, S. Sawleshwarkar, S. Hossain, S.M. Mor, Whence the next pandemic? The
533   intersecting global geography of the animal-human interface, poor health systems and air transit
534   centrality reveals conduits for high-impact spillover, One Heal. (2020).
535   https://doi.org/10.1016/j.onehlt.2020.100177.

536   [9]   N. Myers, R.A. Mittermeier, C.G. Mittermeier, G.A.B. da Fonseca, J. Kent, Biodiversity hotspots
537   for conservation priorities, Nature. 403 (2000) 853–858. https://doi.org/10.1038/35002501.

538   [10]   C.S. Jha, C.B.S. Dutt, K.S. Bawa, Deforestation and land use changes in Western Ghats, India, Curr.
539   Sci. 79 (2000) 231–243.

540   [11]   N.R. Gunawardene, A.E.D. Daniels, I.A.U.N. Gunatilleke, C.V.S. Gunatilleke, P. V Karunakaran, K.G.
541   Nayak, S. Prasad, P. Puyravaud, B.R. Ramesh, K.A. Subramanian, G. Vasanthy, A brief overview of
542   the Western Ghats – Sri Lanka biodiversity hotspot, Curr. Sci. 93 (2007) 1–6.

543   [12]   C. Venkataraman, M. Kamalakannan, D. Dam, An Update on the Mammals of Western Ghats, in:
544   Sivaperuman, Chandrakasan, Venkataraman, Krishnamoorthy (Eds.), Indian Hotspots Vertebr.
545   Faunal Divers. Conserv. Manag., 1st ed., Springer, Singapore, 2018: pp. 205–221.
546   https://doi.org/10.1007/978-981-10-6983-3.

547   [13]   M.D. MADHUSUDAN, Recovery of wild large herbivores following livestock decline in a tropical
548   Indian wildlife reserve, J. Appl. Ecol. 41 (2004) 858–869. https://doi.org/10.1111/j.0021-
549   8901.2004.00950.x.

[14]   B.B. Singh, A.A. Gajadhar, Role of India's wildlife in the emergence and re-emergence of zoonotic

pathogens, risk factors and public health implications, Acta Trop. 138 (2014) 67–77.
https://doi.org/10.1016/j.actatropica.2014.06.009.

[15]   P. Pattnaik, Kyasanur forest disease: an epidemiological view in India, Rev. Med. Virol. 16 (2006)
151–165.

[16]   D.T. Mourya, P.D. Yadav, Recent Scenario of Emergence of Kyasanur Forest Disease in India and
Public Health Importance, Curr. Trop. Med. Reports. 3 (2016) 7–13.
https://doi.org/10.1007/s40475-016-0067-1.

[17]   Y.K. Gurav, P.D. Yadav, M.D. Gokhale, T.R. Chiplunkar, R. Vishwanathan, D.Y. Patil, R. Jain, A.M.
Shete, S.L. Patil, G.D. Sarang, G.N. Sapkal, M.D. Andhare, Y.R. Sale, P.S. Awate, D.T. Mourya,
Kyasanur Forest Disease Prevalence in Western Ghats Proven and Confirmed by Recent Outbreak
in Maharashtra, India, 2016, Vector-Borne Zoonotic Dis. 18 (2018) 164–172.
https://doi.org/10.1089/vbz.2017.2129.

[18]   S.Z. Shah, B. Jabbar, Z. ur Rahman, S. Nadeem, I. Jabbar, S. Azam, N. Ahmed, H. Nasir, A. Rehman,
Epidemiology, Pathogenesis, and Control of a Tick-Borne Disease- Kyasanur Forest Disease:
Current Status and Future Directions, Front. Cell. Infect. Microbiol. 8 (2018).
https://doi.org/10.3389/fcimb.2018.00149.

[19]   N. Naren babu, A. Jayaram, H. Hemanth Kumar, P. Pareet, S. Pattanaik, A.M. Auti, J.
Abdulmajeed, H. Maity, S. Devadiga, Y. Bhandari, H. Agre Deepchand, M. Shakir, N. Kumar, G.
Arunkumar, Spatial distribution of Haemaphysalis species ticks and human Kyasanur Forest
Disease cases along the Western Ghats of India, 2017–2018, Exp. Appl. Acarol. 0 (2019) 0.
https://doi.org/10.1007/s10493-019-00345-9.

[20]   C. Sadanandane, M.D. Gokhale, A. Elango, P. Yadav, D.T. Mourya, P. Jambulingam, Prevalence
and spatial distribution of Ixodid tick populations in the forest fringes of Western Ghats reported
with human cases of Kyasanur forest disease and monkey deaths in South India, Exp. Appl.
Acarol. 75 (2018) 135–142. https://doi.org/10.1007/s10493-018-0223-5.

[21]   G. Geevarghese, A.C. Mishra, Haemaphysalis ticks of India, Elsevier, 2011.

[22]   K. Ajesh, B.K. Nagaraja, K. Sreejith, Kyasanur forest disease virus breaking the endemic barrier:
An investigation into ecological effects on disease emergence and future outlook, Zoonoses
Public Health. 64 (2017) e73–e80. https://doi.org/10.1111/zph.12349.

[23]   M.G. Walsh, S.M. Mor, H. Maity, S. Hossain, Forest loss shapes the landscape suitability of
Kyasanur Forest disease in the biodiversity hotspots of the Western Ghats, India, Int. J.
Epidemiol. (2019). https://doi.org/10.1093/ije/dyz232.

[24]   B. V. Purse, N. Darshan, G.S. Kasabi, F. Gerard, A. Samrat, C. George, A.T. Vanak, M. Oommen, M.
Rahman, S.J. Burthe, J.C. Young, P.N. Srinivas, S.M. Schäfer, P.A. Henrys, V.K. Sandhya, M.M.
Chanda, M. V. Murhekar, S.L. Hoti, S.K. Kiran, Predicting disease risk areas through co-production
of spatial models: The example of Kyasanur Forest Disease in India's forest landscapes, PLoS Negl.
Trop. Dis. 14 (2020) e0008179. https://doi.org/10.1371/journal.pntd.0008179.

[25]   P.D. Yadav, S. Patil, S.M. Jadhav, D.A. Nyayanit, V. Kumar, S. Jain, J. Sampath, D.T. Mourya, S.S.
Cherian, Phylogeography of Kyasanur Forest Disease virus in India (1957–2017) reveals evolution
and spread in the Western Ghats region, Sci. Rep. 10 (2020) 1–12.
https://doi.org/10.1038/s41598-020-58242-w.

[26]   R. Gibb, D.W. Redding, K.Q. Chin, C.A. Donnelly, T.M. Blackburn, T. Newbold, K.E. Jones, Zoonotic
host diversity increases in human-dominated ecosystems, Nature. 584 (2020) 398–402.
https://doi.org/10.1038/s41586-020-2562-8.

[27]   C.K. Johnson, P.L. Hitchens, P.S. Pandit, J. Rushmore, T.S. Evans, C.C.W. Young, M.M. Doyle,
Global shifts in mammalian population trends reveal key predictors of virus spillover risk,
Proceedings. Biol. Sci. 287 (2020) 20192736. https://doi.org/10.1098/rspb.2019.2736.

[28]   R.S. Ostfeld, T. Levi, A.E. Jolles, L.B. Martin, P.R. Hosseini, F. Keesing, Life History and
Demographic Drivers of Reservoir Competence for Three Tick-Borne Zoonotic Pathogens, PLoS
One. 9 (2014) e107387. https://doi.org/10.1371/journal.pone.0107387.

[29]   GBIF, GBIF Occurrence Download - Mammals, Glob. Biodivers. Inf. Facil. (2020).
https://www.gbif.org/occurrence/download/0103251-200613084148143.

[30]   International Society for Infectious Diseases, ProMED-mail, (n.d.). http://www.promedmail.org/.

[31]   WorldClim - Global Climate, Data for current conditions (~1950-2000) | WorldClim - Global
Climate Data, WorldClim - Glob. Clim. Data. (n.d.). http://www.worldclim.org/current (accessed
October 23, 2014).

[32]   R.J. Hijmans, S.E. Cameron, J.L. Parra, P.G. Jones, A. Jarvis, Very high resolution interpolated
climate surfaces for global land areas, Int. J. Climatol. 25 (2005) 1965–1978.
https://doi.org/10.1002/joc.1276.

[33]   C.H.B. Priestley, R.J. Taylor, On the assessment of surface heat flux and evaporation using large-
scale parameters, Mon. Weather Rev. 100 (1972) 81–92. https://doi.org/10.1175/1520-
0493(1972)100<0081:OTAOSH>2.3.CO;2.

[34]   a. Khaldi, A. Khaldi, A. Hamimed, Using the Priestley-Taylor expression for estimating actual
evapotranspiration from satellite Landsat ETM + data, Proc. Int. Assoc. Hydrol. Sci. 364 (2014)
398–403. https://doi.org/10.5194/piahs-364-398-2014.

[35]   A. Trabucco, R.J. Zomer, Global soil water balance geospatial database, CGIAR Consort. Spat. Inf.
(2010). http://www.cgiar-csi.org.

[36]   P.D. Broxton, X. Zeng, D. Sulla-Menashe, P.A. Troch, A global land cover climatology using MODIS
data, J. Appl. Meteorol. Climatol. 53 (2014) 1593–1605. https://doi.org/10.1175/JAMC-D-13-
0270.1.

[37]   M.C. Hansen, P. V. Potapov, R. Moore, M. Hancher, S.A. Turubanova, A. Tyukavina, D. Thau, S. V.
Stehman, S.J. Goetz, T.R. Loveland, A. Kommareddy, A. Egorov, L. Chini, C.O. Justice, J.R.G.
Townshend, High-Resolution Global Maps of 21st-Century Forest Cover Change, Science (80-.).
342 (2013) 850–853. https://doi.org/10.1126/science.1244693.

[38]   Socioeconomic Data and Applications Center | SEDAC, Methods » Last of the Wild, v2 | SEDAC,
(n.d.). http://sedac.ciesin.columbia.edu/data/collection/wildareas-v2/methods (accessed
December 23, 2014).

[39]   E.W. Sanderson, M. Jaiteh, M.A. Levy, K.H. Redford, A. V. Wannebo, G. Woolmer, The Human
Footprint and the Last of the Wild, Bioscience. 52 (2002) 891. https://doi.org/10.1641/0006-
3568(2002)052[0891:THFATL]2.0.CO;2.

[40] D.D. Reidpath, P. Allotey, Infant mortality rate as an indicator of population health, J. Epidemiol.
Community Health. 57 (2003) 344–346. https://doi.org/10.1136/jech.57.5.344.

[41] J. Choi, M. Ki, H.J. Kwon, B. Park, S. Bae, C.-M. Oh, B.C. Chun, G.-J. Oh, Y.H. Lee, T.-Y. Lee, H.K.
Cheong, B.Y. Choi, J.H. Park, S.K. Park, Health Indicators Related to Disease, Death, and
Reproduction, J. Prev. Med. Public Heal. 52 (2019) 14–20. https://doi.org/10.3961/jpmph.18.250.

[42] J. Ignacio Ruiz, K. Nuhu, J. Tyler McDaniel, F. Popoff, A. Izcovich, J. Martin Criniti, Inequality as a
powerful predictor of infant and maternal mortality around the world, PLoS One. 10 (2015).
https://doi.org/10.1371/journal.pone.0140796.

[43] Center for International Earth Science Information Network - CIESIN - Columbia University, Global
Subnational Infant Mortality Rates, Version 2, Palisades, 2019.
https://doi.org/10.7927/H4PN93JJ.

[44] J. Elith, J.R. Leathwick, T. Hastie, A working guide to boosted regression trees., J. Anim. Ecol. 77
(2008) 802–13. https://doi.org/10.1111/j.1365-2656.2008.01390.x.

[45] J. Friedman, Greedy function approximation: a gradient boosting machine, Ann. Stat. 29 (2001)
1189–1232.

[46] L. Breiman, Random forests, Mach. Learn. 45 (2001) 5–32.
https://doi.org/10.1023/A:1010933404324.

[47] G. James, D. Witten, T. Hastie, R. Tibshirani, An introduction to Statistical Learning, 2000.
https://doi.org/10.1007/978-1-4614-7138-7.

[48] S.N. Wood, Stable and Efficient Multiple Smoothing Parameter Estimation for Generalized
Additive Models, J. Am. Stat. Assoc. 99 (2004) 673–686.
https://doi.org/10.1198/016214504000000980.

[49] S.N. Wood, Generalized additive models : an introduction with R, 2nd ed., Chapman and
Hall/CRC, New York, 2017. https://books.google.com.au/books?id=HL-
PDwAAQBAJ&printsec=frontcover&dq=Wood,+S.N.+(2017)+Generalized+Additive+Models:+An+I
ntroduction+with+R+(2nd+edition).+Chapman+and+Hall/CRC.&hl=en&sa=X&ved=0ahUKEwiB_43
x9NvjAhUC4nMBHU7CDWQQ6AEIMDAB#v=onepage&q=Wood%2 (accessed July 30, 2019).

[50] B. Naimi, M.B. Araújo, sdm: a reproducible and extensible R platform for species distribution
modelling, Ecography (Cop.). 39 (2016) 368–375. https://doi.org/10.1111/ecog.01881.

[51] R Core Team, R: A language and environment for statistical computing, R Foundation for
Statistical Computing, Vienna, 2016. https://www.r-project.org/.

[52] A. Baddeley, R. Turner, Practical Maximum Pseudolikelihood for Spatial Point Patterns (with
Discussion), Aust. <html_ent Glyph="@amp;" ascii="&"/> New Zeal. J. Stat. 42 (2000) 283–322.
https://doi.org/10.1111/1467-842X.00128.

[53] A. Baddeley, E. Rubak, R. Turner, Spatial Point Patterns: Methodology and Applications with R,
CRC Press, 2015. https://books.google.com/books?id=rGbmCgAAQBAJ&pgis=1 (accessed
February 5, 2016).

[54] A. Baddeley, R. Turner, spatstat: An R Package for Analyzing Spatial Point Patterns, J. Stat. Softw.
12(6). (2005). http://www.jstatsoft.org/v12/i06/ (accessed October 23, 2014).

[55]   A. Guisan, C. Rahbek, SESAM - a new framework integrating macroecological and species
671   distribution models for predicting spatio-temporal patterns of species assemblages, J. Biogeogr.
672   38 (2011) 1433–1444. https://doi.org/10.1111/j.1365-2699.2011.02550.x.

673   [56]   M. D'Amen, A. Dubuis, R.F. Fernandes, J. Pottier, L. Pellissier, A. Guisan, Using species richness
674   and functional traits predictions to constrain assemblage predictions from stacked species
675   distribution models, J. Biogeogr. 42 (2015) 1255–1266. https://doi.org/10.1111/jbi.12485.

676   [57]   V. Di Cola, O. Broennimann, B. Petitpierre, F.T. Breiner, M. D'Amen, C. Randin, R. Engler, J.
677   Pottier, D. Pio, A. Dubuis, L. Pellissier, R.G. Mateo, W. Hordijk, N. Salamin, A. Guisan, ecospat: an
678   R package to support spatial analyses and modeling of species niches and distributions,
679   Ecography (Cop.). 40 (2017) 774–787. https://doi.org/10.1111/ecog.02671.

680   [58]   M. D'Amen, J.-N. Pradervand, A. Guisan, Predicting richness and composition in mountain insect
681   communities at high resolution: a new test of the SESAM framework, Glob. Ecol. Biogeogr. 24
682   (2015) 1443–1453. https://doi.org/10.1111/geb.12357.

683   [59]   O. ALLOUCHE, A. TSOAR, R. KADMON, Assessing the accuracy of species distribution models:
684   prevalence, kappa and the true skill statistic (TSS), J. Appl. Ecol. 43 (2006) 1223–1232.
685   https://doi.org/10.1111/j.1365-2664.2006.01214.x.

686   [60]   H. Rue, S. Martino, N. Chopin, Approximate Bayesian inference for latent Gaussian models by
687   using integrated nested Laplace approximations, J. R. Stat. Soc. B. 71 (2009) 319–392.
688   https://folk.ntnu.no/hrue/r-inla.org/papers/inla-rss.pdf (accessed November 8, 2017).

689   [61]   M. Begon, Effects of Host Diversity on Disease Dynamics, in: R.S. Ostfeld, F. Keesing, V.T. Eviner
690   (Eds.), Infect. Dis. Ecol. Eff. Ecosyst. Dis. Dis. Ecosyst., First, Princeton University Press, Princeton,
691   2008: pp. 12–29.

692   [62]   D.A. Wilkinson, J.C. Marshall, N.P. French, D.T.S. Hayman, Habitat fragmentation, biodiversity loss
693   and the risk of novel infectious disease emergence, J. R. Soc. Interface. 15 (2018) 20180403.
694   https://doi.org/10.1098/rsif.2018.0403.

695   [63]   M.K. Goverdhan, P.K. Rajagopalan, D.P. Narasimha Murthy, S. Upadhyaya, J. Boshell-M, H.
696   Trapido, T. Ramachandra Rao, Epizootiology of Kyasanur Forest Disease in wild monkeys of
697   Shimoga district, Mysore State (1957-1964)., Indian J. Med. Res. 62 (1974) 497–510.
698   http://www.ncbi.nlm.nih.gov/pubmed/4215749 (accessed March 6, 2019).

699   [64]   M.A. Sreenivasan, H.R. Bhat, P.K. Rajagopalan, The epizootics of Kyasanur Forest disease in wild
700   monkeys during 1964 to 1973., Trans. R. Soc. Trop. Med. Hyg. 80 (1986) 810–4.
701   http://www.ncbi.nlm.nih.gov/pubmed/3603621 (accessed October 30, 2014).

702   [65]   R.H. Kenyon, M.K. Rippy, K.T. McKee, P.M. Zack, C.J. Peters, Infection of Macaca radiata with
703   viruses of the tick-borne encephalitis group., Microb. Pathog. 13 (1992) 399–409.
704   http://www.ncbi.nlm.nih.gov/pubmed/1297916 (accessed March 6, 2019).

705   [66]   P. Rajagopalan, A. Patil, J. Boshell, Ixodid ticks on their mammalian hosts in the Kyasanur Forest
706   disease area of Mysore State, India, 1961-64 , Indian J. Med. Res. 56 (1968) 510–525.
707   https://pubmed.ncbi.nlm.nih.gov/4971268/ (accessed March 23, 2021).

708   [67]   K. Muraleedharan, Wildlife Arthropods of Karnataka with Special Reference to KFD Endemic Area
709   of Shivamogga District▯: Those Parasitic on Small and Large Mammals, Vet. Res. Int. 4 (2016)
710   114–123.

Table 1. Unadjusted (A) and adjusted (B, C, and D) regression coefficients and 95% confidence intervals
for the associations between Kyasanur Forest disease virus outbreaks and landscape features as derived
from inhomogeneous Poisson point process models (PPMs) of the outbreaks. Coefficients in A represent
crude associations from bivariate PPMs, whereas those in B, C, and D represent adjusted associations
wherein each landscape feature is adjusted for all other features included in the multiple PPMs. The
Akaike information criterion (AIC) and area under the receiver operating characteristic curve (AUC) show
model fit and performance, respectively.

Landscape features	Coefficient	95% confidence interval	AIC	AUC
A. Bivariate PPM				
Mammal richness	0.08	0.02 – 0.15	31.79	0.62
Forest loss (deciles)	1.22	0.85 – 1.59	27.14	0.76
Annual precipitation (100 mm)	0.06	0.04 - 0.08	28.64	0.63
Annual temperature (Celsius)	-0.02	-0.03 – -0.01	56.88	0.53
B. Multiple PPM – without climate				
Mammal richness	0.59	0.37 – 0.82	-22.80	0.78
Forest loss (deciles)	5.21	3.94 – 6.48		
Mammal richness:forest loss interaction	-0.41	-0.54 – -0.28		
C. Multiple PPM – with climate				
Mammal richness	0.39	0.12 – 0.65	-36.34	0.74
Forest loss (deciles)	4.82	3.46 – 6.19		
Mammal richness:forest loss interaction	-0.37	-0.51 – -0.23		
Annual precipitation (100 mm)	0.05	0.02 – 0.08		
Annual temperature (Celsius)	-0.03	-0.05 – -0.01		
D. Multiple PPM – with the human footprint				
Mammal richness	0.39	0.12 – 0.66	-35.34	0.75
Forest loss (deciles)	4.86	3.47 – 6.24		
Mammal richness:forest loss interaction	-0.37	-0.51 – -0.23		
Annual precipitation (100 mm)	0.05	0.02 – 0.08		
Annual temperature (Celsius)	-0.03	-0.05 – -0.01		
Human footprint (%)	-0.02	-0.05 – 0.02		

726 Table 2. Taluk-level integrated nested Laplace approximation model (binomial family) of Kyasanur Forest
727 Disease virus outbreaks. Mammal species richness was adjusted for the biotic constraints of sympatric
728 species using the SESAM framework. The Watanabe-Akaike information criterion (WAIC) was used to
729 assess model fit.

Host and landscape features	Coefficient	95% credible interval	WAIC
Mammal richness$_{SESAM}$	0.86	0.39 – 1.35	150.03
Forest loss (deciles)	5.62	2.67 – 8.79	
Mammal richness:forest loss interaction	-0.54	-0.85 – -0.26	
Annual precipitation (100 mm)	0.00	0.000 – 0.1	
Annual temperature (Celsius)	-0.02	-0.034 – -0.004	

Figure legends

Figure 1. The distribution of taluks (beige) across the general region of the Western Ghats comprises five
states (Maharashtra, Goa, Karnataka, Kerala, and Tamil Nadu) in Peninsular India (left panel). Kyasanur
Forest Disease virus outbreaks captured via ProMED-mail (red) and independently sourced (black) are
superimposed over taluks. The distribution of mammalian species richness as estimated from a sample
of Western Ghats species from 2010 to 2020 is presented in the right panel.

Figure 2. The distribution of Kyasanur Forest Disease virus outbreak risk derived from the
inhomogeneous Poisson model is presented in the centre panel with the lower and upper 95%
confidence limits to the left and right, respectively.

Figure 3. The adjusted distribution of mammalian species richness as estimated from a sample of
Western Ghats species. Species richness was adjusted using the SESAM framework to introduce biotic
constraints to adjust the presence of each species for the presence of other species within each taluk.

Figure 4. The distribution of Kyasanur Forest Disease virus outbreak risk derived from the integrated
nested Laplace approximation model (binomial family) is presented in the centre panel with the lower
and upper 95% credible limits to the left and right, respectively.

Figure 5. Taluks identified as priority targets (red) for implementing animal-human surveillance in
Karnataka state. Having located priority taluks, a sample of these can be taken either by random draw or
proportional to the distribution of key landscape features, such as species richness and/or forest loss.
Once the taluks are selected, One Health surveillance infrastructure can be developed by 1) working in
concert with the forest department to select sampling grids from within the sampled taluks for the
monitoring and sampling of wildlife, 2) working in concert with animal husbandry and veterinary
services to generate random samples of farms/livestock holders from animal censuses to monitor and
sample cattle, and 3) working in concert with primary healthcare centres to improve human case
detection and care delivery, conduct serosurveys, and implement prevention outreach.

Lower 95% C.L
□ under 0.011
□ 0.011 to 0.026
□ 0.026 to 0.052
■ 0.052 to 0.094
■ over 0.094

Fitted outbreaks
□ under 0.011
□ 0.011 to 0.026
□ 0.026 to 0.052
■ 0.052 to 0.094
■ over 0.094

Upper 95% C.L.
□ under 0.011
□ 0.011 to 0.026
□ 0.026 to 0.052
■ 0.052 to 0.094
■ over 0.094