Low mammalian species richness increases Kyasanur Forest Disease outbreak risk in deforested landscapes in the Western Ghats, India

Michael G. Walsh[1,2,3,4] Rashmi Bhat[3,4], Venkatesh Nagarajan-Radha[5], Prakash Narayanan [3,4], Navya Vyas[3,4], Shailendra Sawleshwarkar[3,4,6], Chiranjay Mukhopadhyay[7,8]

[1]The University of Sydney, Faculty of Medicine and Health, School of Public Health, Camperdown, New South Wales, Australia, [2]The University of Sydney, Faculty of Medicine and Health, Marie Bashir Institute for Infectious Diseases and Biosecurity, Westmead, New South Wales, Australia, [3]One Health Centre, The Prasanna School of Public Health, Manipal Academy of Higher Education, Manipal, Karnataka, India, [4]The Prasanna School of Public Health, Manipal Academy of Higher Education, Manipal, Karnataka, India, [5]Behaviour Ecology and Evolution Lab, The University of Sydney, Faculty of Science, School of Life and Environmental Sciences, Camperdown, New South Wales, Australia, [6]The University of Sydney, Faculty of Medicine and Health, Westmead Clinical School, Westmead, New South Wales, Australia, [7]Department of Microbiology, Kasturba Medical College, Manipal Academy of Higher Education, Manipal, Karnataka, India, [8]Centre for Emerging and Tropical Diseases, Kasturba Medical College, Manipal Academy of Higher Education, Manipal, Karnataka, India

*Address correspondence to:
Michael Walsh, PhD, MPH
Senior Lecturer, Infectious Diseases Epidemiology
Marie Bashir Institute for Infectious Diseases and Biosecurity
Sydney School of Public Health
The University of Sydney
176 Hawksbury Road
Westmead NSW 2145 Australia
thegowda@gmail.com
michael.walsh1@sydney.edu.au

Abstract

Kyasanur forest disease virus (KFDV) is a rapidly expanding tick-borne zoonotic virus with natural foci in the forested region of the Western Ghats of South India. The Western Ghats is one of the world's most important biodiversity hotspots and, like many such areas of high biodiversity, is under significant pressure from anthropogenic landscape change. The current study sought to quantify mammalian species richness using ensemble models of the distributions of a sample of species extant in the Western Ghats and to explore its association with KFDV outbreaks, as well as the modifying effects of deforestation on this association. Species richness was quantified as a composite of individual species' distributions, as derived from ensembles of boosted regression tree, random forest, and generalised additive models. Species richness was further adjusted for the potential biotic constraints of sympatric species. Both species richness and forest loss demonstrated strong positive associations with KFDV outbreaks, however forest loss substantially modified the association between species richness and outbreaks. High species richness was associated with increased KFDV risk but only in areas of low forest loss. In contrast, lower species richness was associated with increased KFDV risk in areas of greater forest loss. This relationship persisted when species richness was adjusted for biotic constraints at the taluk-level. In addition, the taluk-level species abundances of three monkey species (*Macaca radiata*, *Semnopithecus hypoleucus*, and *Semnopithecus priam*) were also associated with outbreaks. These results suggest that increased monitoring of wildlife in areas of significant habitat fragmentation may add considerably to critical knowledge gaps in KFDV epidemiology and infection ecology and should be incorporated into novel One Health surveillance development for the region. In addition, the inclusion of some primate species as sentinels of KFDV circulation into general wildlife surveillance architecture may add further value.

Keywords: Kyasanur Forest Disease, biodiversity, deforestation, zoonoses, Western Ghats, tick-borne disease

Introduction

The sharing of space between wildlife and humans can be disruptive to both animals and people particularly in the context of altered habitat. Human pressure on wildlife influences species' community composition, population movement and density, resource provisioning[1,2], and eco-immunology[3], which can subsequently alter pathogen circulation among wildlife hosts as well as introduce conduits that allow the movement of pathogens from reservoir hosts to novel human hosts[4], with potentially catastrophic effects[5–7]. As such, the growing wildlife-human interface requires both improved surveillance of pathogen circulation among wildlife and a greater understanding of the impact of anthropogenic landscape change on wildlife communities, their pathogens, and mechanisms for potential spillover of novel pathogens to humans. Importantly, as new wildlife-human interfaces emerge in close proximity to major transportation hubs, the potential for high-impact spillovers with the capacity for rapid regional or global dissemination is increasing[8].

Currently, there is a lack of zoonosis surveillance in wildlife globally, but the absence is particularly noteworthy in the world's biodiversity hotspots that are undergoing rapid anthropogenic change. The Western Ghats region of South India is one of the world's most important biodiversity hotspots[9] and also experiences significant pressure in the form of landscape change due to deforestation, commercial agricultural exploitation, and urbanisation[10]. In addition to the general lack of wildlife pathogen surveillance across this rapidly changing region, there is an even greater absence of work examining the influence of community ecology on the processes of pathogen circulation and spillover events. The emerging tick-borne zoonotic virus, Kyasanur Forest Disease virus (KFDV), simultaneously represents 1) a critical public health priority in the Western Ghats states because of its recent rapid expansion and the high mortality (2% - 10%) associated with its neurological and haematological complications [11–14], and 2) a model system for understanding viral spillover because of the extensive wildlife-human interface across the region[8]. Outbreak occurrence has expanded from

a single district in the state of Karnataka in the decade following the virus' first identification in 1957 to an extensive region now comprising five states[11,15]. Recent work has demonstrated a strong direct association between the expansion of KFDV outbreaks and the loss of native forest[16,17], which was further supported by phylogeographic analyses[18]. The study that identified the direct association with forest loss also noted that areas of high mammalian species richness presented greater risk[16]. While this previous study's assessment of forest loss was of high quality, species richness was assessed more crudely using the IUCN-derived species' range for each mammal species present across the region. As such, variation of species' distributions within those ranges was not incorporated into the assessment of species richness, nor was there any assessment of potential interaction between sympatric species. Additionally, the associations between KFDV and individual species' abundance and relative abundance were not previously explored. It is anticipated that further investigation of these aspects of mammalian biogeography and community ecology in KFDV outbreak hotspots will add considerable insight into the epidemiology of KFDV spillover while also providing an ecological evidence-base for developing municipal wildlife surveillance infrastructure at the taluk level (sub-district) across the Western Ghats states. The incorporation of wildlife monitoring and sampling into KFDV surveillance is particularly important since the investigation of infection in wildlife hosts has fallen off dramatically in recent decades[11].

The specific aims of the current study were as follows. First, we sought to estimate the species distributions of a sample of extant mammals of the Western Ghats (including endemic and non-endemic species) and use these to construct a more representative metric of species richness across the region. Second, we sought to adjust overall species richness, as well as individual species' abundance and relative abundance for the biotic constraints of sympatry. Third, we sought to interrogate the association between species richness and KFDV outbreaks and the modification of this association by

forest loss to develop much needed landscape epidemiology targets for improved surveillance of this expanding zoonotic arbovirus.

Methods

Data Sources

The Global Biodiversity Information Facility was used to obtain all direct observations of extant mammals in the Western Ghats[19] and Peninsular India between 1 January, 2010 and 31 December, 2020 (www.gbif.org). A total of 2826 mammals were observed over this period[20]. This comprised a total of 99 species observed across the region. However, habitat suitability was estimated only for those species with a sample size of at least 30 individuals, for a total of 24 species.

The KFDV outbreaks used in this investigation have been described in detail previously[16]. Briefly, two independent sources of outbreaks reported between 1 January, 2012 and 30 June, 2019 were used for model training and testing, respectively. The training data (47 outbreaks) were sourced from the ProMED-mail[21] electronic surveillance system and the testing data (39 outbreaks) comprised an independent set of laboratory-confirmed outbreaks described in the literature[16]. This allowed for external validation of KFDV outbreak models rather than partitioning data from a single source into training and testing.

Mean annual precipitation and mean annual temperature were obtained from the WorldClim Global Climate database at a resolution of 30 arc seconds (~ 1 km) [22]. The rasters derived from the WorldClim database were mean measurements between the period 1950 to 2000, and thus represent estimates of climate rather than weather. Some regions represented by the WorldClim database have a sparse representation of weather stations contributing to climate estimates. However, India's extensive network of weather stations make its local contributions to decadal climate interpolation more representative than many other large countries[23].

The Priestley-Taylor α coefficient (P-Tα) is the ratio of actual evapotranspiration to potential evapotranspiration and was included here as a metric for water-soil balance[24,25]. In contrast to solar energy input alone, the P-Tα represents water availability in the soil as a function of the local vegetation's water requirements and is therefore a more robust estimate of soil-water balance. A 30 arc seconds resolution raster of P-Tα was obtained from the Consultative Group for International Agricultural Research (CGIAR) Consortium for Spatial Information. The ratio ranges from 0 (extreme water stress) to 1 (no water stress)[26].

Forest cover was first acquired from the Global Land Cover Climatology MODIS-based data product, which corrects temporally aggregated 10-year land cover types by weighting each land cover type by their annual uncertainty and then validates the weighted composite against the System for Terrestrial Ecosystem Parameterization[27]. The product used here for forest cover represents the period 2001-2010 at a resolution of 15 arc seconds (~ 500 m).

Landsat data compiled at 1 arc second (~ 30 m) resolution by the Global Forest Change project was used to quantify forest loss between 2000 and 2012[28]. Landsat imagery was processed using a stratified random sampling validation procedure, with 99.6% accuracy in global settings and 99.5% accuracy in tropical settings[28]. Individual tiles were merged across the region of the Western Ghats and a new quantile raster constructed to represent deciles of forest loss.

An established validated metric for the human footprint (HFP) was used to control for spatial sampling and reporting biases as described below in the analysis section. The metric was acquired from the Socioeconomic Data and Applications Center (SEDAC) repository[29] at a resolution of 30 arc seconds and has been described in detail previously[30]. Briefly, HFP comprises 2 levels of classification. First, an index of human influence was determined based on eight domains: 1) population density, distance to 2) roads, 3) railway lines, 4) navigable rivers, and 5) coastlines, 6) degree of night-time

artificial light, 7) urban vs rural location, and 8) land cover. These domains were then scored to derive the human influence index (HII), ranging from 0 (no human impact) to 64 (greatest human impact). The ratio of the range of minimum and maximum HII in the local terrestrial biome to the range of minimum and maximum HII across all biomes represents the final HFP metric and is expressed as a percentage[30].

Data Analysis

Species distribution modelling. The landscape suitability of each of the 24 mammal species was estimated using an ensemble approach comprising two machine learning methods (boosted regression trees (BRT) and random forests (RF)) and generalised additive models (GAM)). The machine learning frameworks, BRT and RF, are powerful approaches to improving standard decision trees by iteratively growing many trees and amalgamating the results. As the algorithms partition the data space according to rules that optimise homogeneity among predictors and a response (e.g. species presence), many decision trees are grown and combined, resulting in optimised decision trees that reduce overfitting and can capture complex interactions between the predictors [31–34]. There are two key differences in these machine learning approaches. First, for each decision tree partition in RF, only a random subset of predictors is selected from the set of all predictors, which decorrelates the trees and reduces overfitting. Second, rather than decorrelating trees based on the sampling of subsets as with RF, BRT instead reduces overfitting by growing trees sequentially and learning from previously grown trees. In contrast, the GAM framework allows for nonlinear relationships between outcomes and covariates by fitting multiple basis functions for smoothed covariates [35,36]. Five-fold cross-validation was used to fit each model under the three distinct modelling frameworks (BRT, RF, and GAM). Mean annual temperature, P-T α, and forest cover were included as landscape features at 30 arc seconds resolution in all models. For each species, each of the three models describing their landscape suitability was evaluated according to performance, based on the area under the receiver operating characteristic curve (AUC), and fit, based

on the model deviance. Subsequently, an ensemble landscape suitability was computed from the three model classes using their weighted mean, with weights based on AUC [37]. To adjust for potential spatial sampling bias in the GBIF database, background points were sampled proportional to the human footprint as a proxy for landscape accessibility. The estimates of each species' distribution as derived from these ensembles were then summed across all species to estimate local species richness across the Western Ghats region. This estimate as well as an additional estimate adjusted for the biotic constraints of sympatric species (see description below) were subsequently used as more appropriate metrics for mammalian species richness and the evaluation of its influence on KFDV outbreaks. The sdm package[37] in the R platform[38] was used for fitting each model and the derivation of the three-model ensembles to each species.

Point process modelling. The KFDV outbreaks were fitted as an inhomogeneous Poisson point process[39,40]. Under this model framework the spatial dependence of the outbreaks' distribution can be determined and evaluated with respect to specific landscape features. The background points used in the model were sampled proportional to HFP to control for potential outbreak reporting bias. Previous work demonstrated a strong association between KFDV outbreaks and both forest loss and mammalian species richness[16]. However, the association with the latter was 1) based on a crude representation of species richness that did not account for local heterogeneity of individual species' distributions and 2) could not be used to evaluate the potential effect modification of species richness by forest loss. The current investigation provides this critical missing component by incorporating a more suitable metric of species richness based on the individual estimated species' distributions described above and subsequently interrogating the interaction between species richness and forest loss. In addition, to account for the important climate features that were incorporated in the previous work, precipitation and temperature were also included in the current work. Therefore, the full multiple inhomogeneous Poisson model comprised the newly-quantified mammalian species richness metric described above,

forest loss, an interaction term for species richness and forest loss, mean annual precipitation and mean annual temperature all aggregated up to 2.5 arc minutes. The landscape features included in the multiple inhomogeneous Poisson models were not highly correlated so multicollinearity in the models was not of concern (all values of the Pearson's r were < 0.5). Model fit was assessed using the Akaike information criterion (AIC) and model performance was tested against an independent, laboratory-confirmed set of outbreak data, as described above. Performance was evaluated using the AUC. The spatstat R package was used to fit the point process models[41].

Taluk-level modelling. Individual species abundance and relative abundance, as well as species richness, were further adjusted for the potential modulating effects of sympatric species at the scale of the taluk. Taluks are 3rd-level, subdistrict administration units in India and are small enough to reasonably represent shared space between sympatric species, while simultaneously large enough to delineate the minimal municipal infrastructure (i.e. delineating the most local focus) used in the Western Ghats states for the implementation of animal and human disease surveillance. Specifically, biotic constraints were applied to the estimates of each species distribution according to the spatially-explicit species assemblage modelling (SESAM) framework[42–44]. Under this framework, first the landscape suitability for each species was estimated according to the ensemble method described above. Second, species richness was calculated as the sum of the individual species distributions, again using the method described above. Third, the biotic constraint was applied whereby each species is evaluated with respect to all other species present within each taluk via the probability ranking rule[43,45] to determine whether a given species should be retained within, or excluded from, each taluk "community". This provides an estimate of each species' presence (or absence) at the taluk level given the potential biotic constraints of the other species within the taluk. For those species thusly identified as present within the taluk community, all 1 km^2 pixels exceeding the true skill statistic (TSS)[46] of their ensemble landscape suitability estimate were designated present (1 = present, 0 = otherwise) and summed across all pixels

within the taluk. This yielded a taluk-level estimate of species abundance for each species under consideration. Similarly, the taluk-level relative abundance for each species was quantified by dividing each individual species abundance by the total abundance of all species identified as present within each taluk under the SESAM framework. Finally, a new estimate of species richness was computed by summing all the species identified as present within each taluk under the SESAM framework. This systematic approach thus yielded taluk-level community estimates of each species' abundance and relative abundance, as well as an updated estimate of species richness, all adjusted for the biotic constraints of sympatric species. The SESAM framework, including the probability ranking rule was implemented using the ecospat package in R[44].

Integrated nested Laplace approximation (INLA) models[47] were used to estimate the association between KFDV outbreaks and the biotically constrained estimate of mammalian species richness and its interaction with forest loss at the level of the taluk, as well as associations between KFDV outbreaks and individual species abundance and relative abundance to identify potential species that may be good candidates for initial wildlife surveillance. While the latter may represent species that are important as reservoir hosts, we did not attempt to assess species' roles as hosts in the current study since we did not measure their competence for KFDV. Instead, individual species associations were investigated simply to identify whether certain species may be useful sentinels of KFDV outbreaks. The binomial likelihood family was used for the INLA models with spatial autocorrelation estimated using random effects with Besag–York–Mollie priors. Under the binomial family, taluks were modelled as either having experienced a KFDV outbreak under the period of study (outbreak positive) or not (outbreak negative). This approach was taken since the majority of taluks in the region did not experience outbreaks, and, among those that did, most only experienced one or two outbreaks. Nevertheless, to evaluate the effect of different distribution families, we also fitted these models with the zero-inflated Poisson family as a sensitivity analysis. Since the KFDV outbreaks were necessarily

aggregated by taluk, to increase the outbreak sample size available for taluk-level modelling, the independent datasets described above for the point process models (PPMs) were combined for the INLA models. Since the primary objective was estimation rather than prediction so as to infer associations between KFDV outbreaks and species richness, this was deemed an acceptable trade-off. The fit of INLA models was evaluated using the Watanabe-Akaike information criterion (WAIC). The INLA models were fit using the inla package in R (www.r-inla.org) [47].

Results

The distribution of mammalian species richness, estimated as the sum of the sample of individual species' distributions, is presented in Figure 1 juxtaposed with the distribution of KFDV outbreaks. The individual species distribution models generally performed well (Supplementary material, S1 Table 1). The distribution of each species' landscape suitability is presented in S2 Figure 1 and S3 Figure 2. Both forest loss and the new quantification of mammalian species richness based on individual species' landscape suitability alone were very strongly associated with KFDV outbreaks at local scale (Table 1). Strong associations were apparent whether these features were considered alone in the crude bivariate PPMs (Table 1A), or together in the multiple PPMs (Table 1B and 1C). Critically, while species richness demonstrated a similar association with KFDV outbreaks to that shown previously[16], the current investigation identified significant and substantial interaction between species richness and forest loss. As such, increasing mammalian species richness was crudely associated with increasing KFDV outbreak occurrence. However, when species richness was considered in concert with forest loss, the KFDV association with the former was modified by the latter (coefficient = -0.26). This relationship was maintained in the full model, which incorporated the climate features and was a markedly better fit (AIC = -30.19 vs. -9.71) although performance was slightly diminished (AUC = 0.74 vs. 0.78). The distribution of KFDV outbreak risk across the Western Ghats based on the full model is presented in Figure 2.

The newly quantified metric for taluk-level species richness based on the SESAM framework is presented in Figure 3. In comparison to the best-fitting high-resolution PPM model (Table 1C), the taluk-level INLA model showed the same strong interaction between forest loss and mammal species richness, even as the latter was adjusted for biotic constraint under the SESAM framework and scaled up to taluk (Table 2). The sensitivity analysis showed that these associations all persisted when the distribution family was specified as zero-inflated Poisson (S4 Table 2 and S5 Figure 3), although temperature was no longer associated with outbreak occurrence. Based on taluk-level outbreak risk (Figure 4), we identified the 50[th] percentile of risk where deforestation and species richness optimally converge for targeted selection for a state-administered (e.g. Karnataka) animal surveillance program (Figure 5). In addition to the broad metric of species richness, we also explored the associations between KFDV outbreaks and each individual species' abundance and relative abundance to determine if any particular species may prove useful as a sentinel surveillance target. Despite several species demonstrating significant individual associations with KFDV occurrence at fine scale based on their estimated landscape suitability (S6 Table 3), most species did not demonstrate associations with outbreaks at the level of the taluk (S7 Table 4). However, there were a few notable exceptions. Species abundance of the primates *Macaca radiata*, *Semnopithecus hypoleucos* and *Semnopithecus priam*, were all associated with increased KFDV occurrence at the taluk level. Species relative abundance was also associated with KFDV outbreaks for *S. hypoleucos*, but not for any other primate species. The distributions of these species' abundances and relative abundances per taluk are presented in S8 Figure 4. Finally, both the abundance and relative abundance of *Panthera pardus*, and the abundance of *Axis axis*, were also associated with increased KFDV outbreak occurrence.

Discussion

This is the first study to attempt a general estimate of mammalian species richness across the Western Ghats region based on ensemble modelling of the landscape suitability of several extant species, and to

extend this by adding biotic constraints of sympatric species to the estimates of species richness, as well as individual species abundance and relative abundance. This is also the first study to demonstrate negative effect modification of the association between mammalian species richness and KFDV outbreak occurrence by forest loss such that higher species richness was associated with increased KFDV outbreak risk where deforestation was minimal, but in areas of increasing forest loss lower species richness was associated with increased risk. This association was identified at both fine scale and at the level of the taluk. Finally, three monkey species were identified (*M. radiata*, *S. hypoleucus*, and *S. priam*) whose taluk-level species abundances were indicators of outbreaks. Taken together, these findings have identified important interaction between landscape degradation and mammalian species richness that may influence the ecology of this tick-borne virus in its natural foci. The fact that this relationship held at the taluk level even after controlling for biotic constraints indicates that enhanced monitoring of wildlife and human populations at the level of the taluk, which is well-suited to the municipal infrastructure of current human and animal health systems in the Western Ghats states, may offer promise as the focus of novel One Health surveillance development. In addition, the incorporation of some primate species as sentinels of KFDV circulation into general wildlife surveillance may add further value to such a One Health program.

Previous work identified increased risk of KFDV outbreaks associated with both forest loss and high mammalian species richness[16]. However, the latter association was difficult to interpret due to the fairly crude measure of species richness, which was simply a composite of IUCN mammal species' ranges across the region. Moreover, this work was not able to assess interaction between forest loss and mammalian biodiversity. The positive association between KFDV outbreak risk and species richness shown in that study was reasonable given the expectation of greater exposure to potential KFDV hosts and their tick vectors following increasing incursion into areas of high species richness. Nevertheless, it is also of particular interest to identify how habitat loss may affect the community ecology of

mammalian hosts in ways that could influence KFDV spillover to humans. Therefore, better metrics of species richness were required across the region to more appropriately evaluate the modifying effect of forest loss on the association between species richness and KFDV outbreaks. The results presented in this current investigation indicated that the relationship between species richness and forest loss is indeed more nuanced. As shown previously[16], increased species richness was associated with increased KFDV outbreak risk but in the current study this relationship only held in landscapes with minimal forest loss. In contrast, as forest loss increased, lower species richness was associated with increased risk. While this finding allows no definitive assertion regarding the nature of KFDV circulation among mammalian hosts, such as the relative importance of the dilution effect to transmission dynamics[48], it does provide important evidence that KFDV circulation among reservoir hosts persists, and spillover to humans continues, in human-altered landscapes exhibiting lower biodiversity, which is consistent with previous evidence showing clear pathways from habitat fragmentation and biodiversity loss to increased spillover[1,49]. Despite the current study's inability to make specific claims about the role of any individual species in the viral maintenance and infection ecology of KFDV, the findings do suggest that, in general, species resilient to anthropogenic landscape change may be most relevant to the expansion of KFDV spillover across the region. This finding closely parallels recent work demonstrating similar relationships more globally for other zoonotic pathogens[50,51]. Moreover, the work may help to address a critical surveillance gap currently impeding a fuller understanding of KFDV infection ecology. Following the initial identification of KFDV in 1957, there was a rapid but limited flurry of research attempting to identify wildlife reservoirs for KFDV in the Bandipur Forest range[11]. Several mammalian species were identified as hosts, but these studies were not definitive with respect to KFDV infection ecology as many relied on serology, rather than measuring host competence, and many of the studies employed limited sampling strategies[11]. Subsequently, wildlife investigations all but stopped by the early 1980s before KFDV reservoirs could be established. Therefore, the current study could have

important implications for reinstating widespread wildlife surveillance of KFDV in the Western Ghats with a focus on synanthropic species.

It is important to caution against interpretation of these findings to infer specific dynamics of infection ecology at either the level of the community or individual species because infection competence was not assessed in any of the mammals under investigation nor was interspecific interaction among species directly observed at observed species' locations. However, the results do provide some of the most actionable evidence to date for targeted, landscape-designed wildlife surveillance across the region. Specifically, these findings can help to frame a landscape-based cluster sampling design for the monitoring of wildlife distribution and movement in response to human induced landscape change in concert with pathogen surveillance in animals and humans (Figure 5). Critically, the fact that this study's findings were consistent at the level of the taluk indicates that the existing municipal infrastructure for animal and human health can be used as the necessary scaffolding for novel One Health surveillance architecture and service delivery. Moreover, the taluk-level abundance of bonnet macaques (*M. radiata*) and grey langurs (*Semnopithecus spp.*), which have previously been identified as highly susceptible to KFDV[52–54], were also associated with outbreaks. As mentioned above, the current study can make no claims regarding the roles of individual species in the infection ecology of KFDV, but the associations with these primate species' abundance within taluks indicates that these species may provide useful sentinels under a One Health surveillance framework. In addition, leopard (*P. pardus*) abundance and relative abundance and chital (*A. axis*) abundance were also associated with outbreak risk. While neither of these species have been shown to be susceptible to KFDV, both are known to host KFDV-relevant ticks and distribute these vectors in the landscape[55,56]. Therefore, their associations with increased KFDV outbreak occurrence in the current study merits a closer look at the role these species could play as tick distributors, particularly since both of these species are highly adaptive to anthropogenic landscapes. As such leopards and chital could have

repercussions for human exposure to KFDV vectors, their unlikely role as important KFDV hosts notwithstanding.

Further to the cautionary interpretation around KFDV infection ecology encouraged above, some further description of this study's limitations is provided. First, as previously mentioned direct observation of interspecific interaction was not possible under the current investigation, which precludes the ability to make specific claims about the true nature and influence of interaction among communities at various scales across the region. The findings reported here will need to be verified by systematically conducted field investigations of directly observed interspecific interaction at multiple scales. Second, sufficient observations were not available for each extant mammal species of the Western Ghats and so the new quantification of species richness presented here, although much improved over previous estimates, is only a sample of Western Ghats mammals and therefore remains a proxy for true species richness. Third, spatial sampling bias may affect the distribution of species observations, as well as the reporting of KFDV outbreaks, across the region. To control for this sampling bias, background points were not selected randomly but instead selected proportional to the human footprint as a measure of accessibility. Finally, this study provides no description of the roles of individual species as reservoirs, whether operating as maintenance, amplification, or bridging hosts, since neither infection competence nor infection susceptibility was measured in these species in the current study. Rather, individual species' landscape suitability was used to 1) calculate species richness and evaluate this as a landscape feature of importance to the distribution of KFDV outbreak occurrence, particularly with respect to forest loss, and 2) identify individual species that may serve as useful sentinels in the development of wildlife surveillance mechanisms. Therefore, rather than defining how mammalian community ecology determines KFDV infection ecology, the current work delineates optimal landscape targets for developing the surveillance instruments necessary to appropriately

monitor and sample wildlife so that the necessary community ecology and infection ecology data that are needed can be generated.

In conclusion, this study provides novel metrics of mammalian species richness in the Western Ghats, one of the world's critical biodiversity hotspots, and demonstrates the influence of this landscape feature and its interaction with deforestation on the expansion of one of India's most important emerging zoonotic arboviruses. The insight gained can provide municipally-directed targets for landscape-based One Health surveillance of animals and humans in the face of anthropogenic landscape change, which can begin to provide a definitive understanding of KFDV infection ecology while simultaneously identifying the wildlife-human interfaces most vulnerable to pathogen spillover and which may yield the highest benefit from tailored intervention.

References

[1] C.L. Faust, H.I. McCallum, L.S.P. Bloomfield, N.L. Gottdenker, T.R. Gillespie, C.J. Torney, A.P. Dobson, R.K. Plowright, Pathogen spillover during land conversion, Ecol. Lett. 21 (2018) 471–483. https://doi.org/10.1111/ele.12904.

[2] R.K. Plowright, C.R. Parrish, H. McCallum, P.J. Hudson, A.I. Ko, A.L. Graham, J.O. Lloyd-Smith, Pathways to zoonotic spillover, Nat. Rev. Microbiol. 15 (2017) 502–510. https://doi.org/10.1038/nrmicro.2017.45.

[3] D.J. Becker, G.F. Albery, M.K. Kessler, T.J. Lunn, C.A. Falvo, G. Czirják, L.B. Martin, R.K. Plowright, Macroimmunology: The drivers and consequences of spatial patterns in wildlife immune defence, J. Anim. Ecol. 89 (2020) 972–995. https://doi.org/10.1111/1365-2656.13166.

[4] B. Borremans, C. Faust, K.R. Manlove, S.H. Sokolow, J.O. Lloyd-Smith, Cross-species pathogen spillover across ecosystem boundaries: Mechanisms and theory, Philos. Trans. R. Soc. B Biol. Sci. 374 (2019). https://doi.org/10.1098/rstb.2018.0344.

[5] L.F. Wang, B.T. Eaton, Bats, civets and the emergence of SARS, Curr. Top. Microbiol. Immunol. 315 (2007) 325–344. https://doi.org/10.1007/978-3-540-70962-6_13.

[6] C.E.M. Coltart, B. Lindsey, I. Ghinai, A.M. Johnson, D.L. Heymann, The Ebola outbreak, 2013–2016: Old lessons for new epidemics, Philos. Trans. R. Soc. B Biol. Sci. 372 (2017). https://doi.org/10.1098/rstb.2016.0297.

[7] K.G. Andersen, A. Rambaut, W.I. Lipkin, E.C. Holmes, R.F. Garry, The proximal origin of SARS-CoV-2, Nat. Med. 26 (2020) 450–452. https://doi.org/10.1038/s41591-020-0820-9.

[8] M.G. Walsh, S. Sawleshwarkar, S. Hossain, S.M. Mor, Whence the next pandemic? The intersecting global geography of the animal-human interface, poor health systems and air transit centrality reveals conduits for high-impact spillover, One Heal. (2020). https://doi.org/10.1016/j.onehlt.2020.100177.

[9] N. Myers, R.A. Mittermeier, C.G. Mittermeier, G.A.B. da Fonseca, J. Kent, Biodiversity hotspots for conservation priorities, Nature. 403 (2000) 853–858. https://doi.org/10.1038/35002501.

[10] C.S. Jha, C.B.S. Dutt, K.S. Bawa, Deforestation and land use changes in Western Ghats, India, Curr. Sci. 79 (2000) 231–243.

[11] P. Pattnaik, Kyasanur forest disease: an epidemiological view in India, Rev. Med. Virol. 16 (2006) 151–165.

[12] D.T. Mourya, P.D. Yadav, Recent Scenario of Emergence of Kyasanur Forest Disease in India and Public Health Importance, Curr. Trop. Med. Reports. 3 (2016) 7–13. https://doi.org/10.1007/s40475-016-0067-1.

[13] Y.K. Gurav, P.D. Yadav, M.D. Gokhale, T.R. Chiplunkar, R. Vishwanathan, D.Y. Patil, R. Jain, A.M. Shete, S.L. Patil, G.D. Sarang, G.N. Sapkal, M.D. Andhare, Y.R. Sale, P.S. Awate, D.T. Mourya, Kyasanur Forest Disease Prevalence in Western Ghats Proven and Confirmed by Recent Outbreak in Maharashtra, India, 2016, Vector-Borne Zoonotic Dis. 18 (2018) 164–172. https://doi.org/10.1089/vbz.2017.2129.

[14] S.Z. Shah, B. Jabbar, Z. ur Rahman, S. Nadeem, I. Jabbar, S. Azam, N. Ahmed, H. Nasir, A. Rehman,

Epidemiology, Pathogenesis, and Control of a Tick-Borne Disease- Kyasanur Forest Disease: Current Status and Future Directions, Front. Cell. Infect. Microbiol. 8 (2018). https://doi.org/10.3389/fcimb.2018.00149.

[15] K. Ajesh, B.K. Nagaraja, K. Sreejith, Kyasanur forest disease virus breaking the endemic barrier: An investigation into ecological effects on disease emergence and future outlook, Zoonoses Public Health. 64 (2017) e73–e80. https://doi.org/10.1111/zph.12349.

[16] M.G. Walsh, S.M. Mor, H. Maity, S. Hossain, Forest loss shapes the landscape suitability of Kyasanur Forest disease in the biodiversity hotspots of the Western Ghats, India, Int. J. Epidemiol. (2019). https://doi.org/10.1093/ije/dyz232.

[17] B. V. Purse, N. Darshan, G.S. Kasabi, F. Gerard, A. Samrat, C. George, A.T. Vanak, M. Oommen, M. Rahman, S.J. Burthe, J.C. Young, P.N. Srinivas, S.M. Schäfer, P.A. Henrys, V.K. Sandhya, M.M. Chanda, M. V. Murhekar, S.L. Hoti, S.K. Kiran, Predicting disease risk areas through co-production of spatial models: The example of Kyasanur Forest Disease in India's forest landscapes, PLoS Negl. Trop. Dis. 14 (2020) e0008179. https://doi.org/10.1371/journal.pntd.0008179.

[18] P.D. Yadav, S. Patil, S.M. Jadhav, D.A. Nyayanit, V. Kumar, S. Jain, J. Sampath, D.T. Mourya, S.S. Cherian, Phylogeography of Kyasanur Forest Disease virus in India (1957–2017) reveals evolution and spread in the Western Ghats region, Sci. Rep. 10 (2020) 1–12. https://doi.org/10.1038/s41598-020-58242-w.

[19] C. Venkataraman, M. Kamalakannan, D. Dam, An Update on the Mammals of Western Ghats, in: Sivaperuman, Chandrakasan, Venkataraman, Krishnamoorthy (Eds.), Indian Hotspots Vertebr. Faunal Divers. Conserv. Manag., 1st ed., Springer, Singapore, 2018: pp. 205–221. https://doi.org/10.1007/978-981-10-6983-3.

[20] GBIF, GBIF Occurrence Download, Glob. Biodivers. Inf. Facil. (n.d.). https://www.gbif.org/occurrence/download/0109657-200613084148143 (accessed February 12, 2021).

[21] International Society for Infectious Diseases, ProMED-mail, (n.d.). http://www.promedmail.org/.

[22] WorldClim - Global Climate, Data for current conditions (~1950-2000) | WorldClim - Global Climate Data, WorldClim - Glob. Clim. Data. (n.d.). http://www.worldclim.org/current (accessed October 23, 2014).

[23] R.J. Hijmans, S.E. Cameron, J.L. Parra, P.G. Jones, A. Jarvis, Very high resolution interpolated climate surfaces for global land areas, Int. J. Climatol. 25 (2005) 1965–1978. https://doi.org/10.1002/joc.1276.

[24] C.H.B. Priestley, R.J. Taylor, On the assessment of surface heat flux and evaporation using large-scale parameters, Mon. Weather Rev. 100 (1972) 81–92. https://doi.org/10.1175/1520-0493(1972)100<0081:OTAOSH>2.3.CO;2.

[25] a. Khaldi, A. Khaldi, A. Hamimed, Using the Priestley-Taylor expression for estimating actual evapotranspiration from satellite Landsat ETM + data, Proc. Int. Assoc. Hydrol. Sci. 364 (2014) 398–403. https://doi.org/10.5194/piahs-364-398-2014.

[26] A. Trabucco, R.J. Zomer, Global soil water balance geospatial database, CGIAR Consort. Spat. Inf. (2010). http://www.cgiar-csi.org.

[27]   P.D. Broxton, X. Zeng, D. Sulla-Menashe, P.A. Troch, A global land cover climatology using MODIS data, J. Appl. Meteorol. Climatol. 53 (2014) 1593–1605. https://doi.org/10.1175/JAMC-D-13-0270.1.

[28]   M.C. Hansen, P. V. Potapov, R. Moore, M. Hancher, S.A. Turubanova, A. Tyukavina, D. Thau, S. V. Stehman, S.J. Goetz, T.R. Loveland, A. Kommareddy, A. Egorov, L. Chini, C.O. Justice, J.R.G. Townshend, High-Resolution Global Maps of 21st-Century Forest Cover Change, Science (80-.). 342 (2013) 850–853. https://doi.org/10.1126/science.1244693.

[29]   Socioeconomic Data and Applications Center | SEDAC, Methods » Last of the Wild, v2 | SEDAC, (n.d.). http://sedac.ciesin.columbia.edu/data/collection/wildareas-v2/methods (accessed December 23, 2014).

[30]   E.W. Sanderson, M. Jaiteh, M.A. Levy, K.H. Redford, A. V. Wannebo, G. Woolmer, The Human Footprint and the Last of the Wild, Bioscience. 52 (2002) 891. https://doi.org/10.1641/0006-3568(2002)052[0891:THFATL]2.0.CO;2.

[31]   J. Elith, J.R. Leathwick, T. Hastie, A working guide to boosted regression trees., J. Anim. Ecol. 77 (2008) 802–13. https://doi.org/10.1111/j.1365-2656.2008.01390.x.

[32]   J. Friedman, Greedy function approximation: a gradient boosting machine, Ann. Stat. 29 (2001) 1189–1232.

[33]   L. Breiman, Random forests, Mach. Learn. 45 (2001) 5–32. https://doi.org/10.1023/A:1010933404324.

[34]   G. James, D. Witten, T. Hastie, R. Tibshirani, An introduction to Statistical Learning, 2000. https://doi.org/10.1007/978-1-4614-7138-7.

[35]   S.N. Wood, Stable and Efficient Multiple Smoothing Parameter Estimation for Generalized Additive Models, J. Am. Stat. Assoc. 99 (2004) 673–686. https://doi.org/10.1198/016214504000000980.

[36]   S.N. Wood, Generalized additive models : an introduction with R, 2nd ed., Chapman and Hall/CRC, New York, 2017. https://books.google.com.au/books?id=HL-PDwAAQBAJ&printsec=frontcover&dq=Wood,+S.N.+(2017)+Generalized+Additive+Models:+An+Introduction+with+R+(2nd+edition).+Chapman+and+Hall/CRC.&hl=en&sa=X&ved=0ahUKEwiB_43x9NvjAhUC4nMBHU7CDWQQ6AEIMDAB#v=onepage&q=Wood%2 (accessed July 30, 2019).

[37]   B. Naimi, M.B. Araújo, sdm: a reproducible and extensible R platform for species distribution modelling, Ecography (Cop.). 39 (2016) 368–375. https://doi.org/10.1111/ecog.01881.

[38]   R Core Team, R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, 2016. https://www.r-project.org/.

[39]   A. Baddeley, R. Turner, Practical Maximum Pseudolikelihood for Spatial Point Patterns (with Discussion), Aust. <html_ent Glyph="@amp;" ascii="&"/> New Zeal. J. Stat. 42 (2000) 283–322. https://doi.org/10.1111/1467-842X.00128.

[40]   A. Baddeley, E. Rubak, R. Turner, Spatial Point Patterns: Methodology and Applications with R, CRC Press, 2015. https://books.google.com/books?id=rGbmCgAAQBAJ&pgis=1 (accessed February 5, 2016).

[41]   A. Baddeley, R. Turner, spatstat: An R Package for Analyzing Spatial Point Patterns, J. Stat. Softw. 12(6). (2005). http://www.jstatsoft.org/v12/i06/ (accessed October 23, 2014).

[42]   A. Guisan, C. Rahbek, SESAM - a new framework integrating macroecological and species distribution models for predicting spatio-temporal patterns of species assemblages, J. Biogeogr. 38 (2011) 1433–1444. https://doi.org/10.1111/j.1365-2699.2011.02550.x.

[43]   M. D'Amen, A. Dubuis, R.F. Fernandes, J. Pottier, L. Pellissier, A. Guisan, Using species richness and functional traits predictions to constrain assemblage predictions from stacked species distribution models, J. Biogeogr. 42 (2015) 1255–1266. https://doi.org/10.1111/jbi.12485.

[44]   V. Di Cola, O. Broennimann, B. Petitpierre, F.T. Breiner, M. D'Amen, C. Randin, R. Engler, J. Pottier, D. Pio, A. Dubuis, L. Pellissier, R.G. Mateo, W. Hordijk, N. Salamin, A. Guisan, ecospat: an R package to support spatial analyses and modeling of species niches and distributions, Ecography (Cop.). 40 (2017) 774–787. https://doi.org/10.1111/ecog.02671.

[45]   M. D'Amen, J.-N. Pradervand, A. Guisan, Predicting richness and composition in mountain insect communities at high resolution: a new test of the SESAM framework, Glob. Ecol. Biogeogr. 24 (2015) 1443–1453. https://doi.org/10.1111/geb.12357.

[46]   O. ALLOUCHE, A. TSOAR, R. KADMON, Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS), J. Appl. Ecol. 43 (2006) 1223–1232. https://doi.org/10.1111/j.1365-2664.2006.01214.x.

[47]   H. Rue, S. Martino, N. Chopin, Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximations, J. R. Stat. Soc. B. 71 (2009) 319–392. https://folk.ntnu.no/hrue/r-inla.org/papers/inla-rss.pdf (accessed November 8, 2017).

[48]   M. Begon, Effects of Host Diversity on Disease Dynamics, in: R.S. Ostfeld, F. Keesing, V.T. Eviner (Eds.), Infect. Dis. Ecol. Eff. Ecosyst. Dis. Dis. Ecosyst., First, Princeton University Press, Princeton, 2008: pp. 12–29.

[49]   D.A. Wilkinson, J.C. Marshall, N.P. French, D.T.S. Hayman, Habitat fragmentation, biodiversity loss and the risk of novel infectious disease emergence, J. R. Soc. Interface. 15 (2018) 20180403. https://doi.org/10.1098/rsif.2018.0403.

[50]   C.K. Johnson, P.L. Hitchens, P.S. Pandit, J. Rushmore, T.S. Evans, C.C.W. Young, M.M. Doyle, Global shifts in mammalian population trends reveal key predictors of virus spillover risk, Proceedings. Biol. Sci. 287 (2020) 20192736. https://doi.org/10.1098/rspb.2019.2736.

[51]   R. Gibb, D.W. Redding, K.Q. Chin, C.A. Donnelly, T.M. Blackburn, T. Newbold, K.E. Jones, Zoonotic host diversity increases in human-dominated ecosystems, Nature. 584 (2020) 398–402. https://doi.org/10.1038/s41586-020-2562-8.

[52]   M.K. Goverdhan, P.K. Rajagopalan, D.P. Narasimha Murthy, S. Upadhyaya, J. Boshell-M, H. Trapido, T. Ramachandra Rao, Epizootiology of Kyasanur Forest Disease in wild monkeys of Shimoga district, Mysore State (1957-1964)., Indian J. Med. Res. 62 (1974) 497–510. http://www.ncbi.nlm.nih.gov/pubmed/4215749 (accessed March 6, 2019).

[53]   M.A. Sreenivasan, H.R. Bhat, P.K. Rajagopalan, The epizootics of Kyasanur Forest disease in wild monkeys during 1964 to 1973., Trans. R. Soc. Trop. Med. Hyg. 80 (1986) 810–4. http://www.ncbi.nlm.nih.gov/pubmed/3603621 (accessed October 30, 2014).

[54]   R.H. Kenyon, M.K. Rippy, K.T. McKee, P.M. Zack, C.J. Peters, Infection of Macaca radiata with viruses of the tick-borne encephalitis group., Microb. Pathog. 13 (1992) 399–409. http://www.ncbi.nlm.nih.gov/pubmed/1297916 (accessed March 6, 2019).

[55]   P. Rajagopalan, A. Patil, J. Boshell, Ixodid ticks on their mammalian hosts in the Kyasanur Forest disease area of Mysore State, India, 1961-64 , Indian J. Med. Res. 56 (1968) 510–525. https://pubmed.ncbi.nlm.nih.gov/4971268/ (accessed March 23, 2021).

[56]   K. Muraleedharan, Wildlife Arthropods of Karnataka with Special Reference to KFD Endemic Area of Shivamogga District⬚: Those Parasitic on Small and Large Mammals, Vet. Res. Int. 4 (2016) 114–123.

Table 1. Unadjusted (A) and adjusted (B and C) regression coefficients and 95% confidence intervals for the associations between Kyasanur Forest disease virus outbreaks and landscape features as derived from inhomogeneous Poisson models of the outbreak point process. Coefficients in A represent crude associations from bivariate PPMs, whereas those in B and C represent adjusted associations wherein each landscape feature is adjusted for all other features included in the multiple PPMs. The Akaike information criterion (AIC) and area under the receiver operating characteristic curve (AUC) show model fit and performance, respectively.

Landscape features	Coefficient	95% confidence interval	AIC	AUC
A. Bivariate PPM				
Mammal richness	0.08	0.02 – 0.14	34.81	0.62
Forest loss (deciles)	1.27	0.90 – 1.64	26.37	0.76
Annual precipitation (100 mm)	0.06	0.04 - 0.08	30.23	0.63
Annual temperature (Celsius)	-0.19	-0.31 – -0.07	57.13	0.52
B. Multiple PPM – without climate				
Mammal richness	0.41	0.21 – 0.61	-9.71	0.78
Forest loss (deciles)	3.62	2.68 – 4.56		
Mammal richness:forest loss interaction	-0.26	-0.35 – -0.16		
C. Multiple PPM – with climate				
Mammal richness	0.27	0.03 – 0.51	-30.19	0.74
Forest loss (deciles)	4.02	2.94 – 5.09		
Mammal richness:forest loss interaction	-0.28	-0.40 – -0.17		
Annual precipitation (100 mm)	0.06	0.03 – 0.09		
Annual temperature (Celsius)	-0.03	-0.04 – -0.02		

Table 2. Taluk-level integrated nested Laplace approximation model (binomial family) of Kyasanur Forest Disease virus outbreaks. Mammal species richness was adjusted for the biotic constraints of sympatric species using the SESAM framework. The Watanabe-Akaike information criterion (WAIC) was used to assess model fit.

Host and landscape features	Coefficient	95% credible interval	WAIC
Mammal richness$_{SESAM}$	0.86	0.39 – 1.35	150.03
Forest loss (deciles)	5.62	2.67 – 8.79	
Mammal richness:forest loss interaction	-0.54	-0.85 – -0.26	
Annual precipitation (100 mm)	0.00	0.000 – 0.001	
Annual temperature (Celsius)	-0.20	-0.34 – -0.04	

Figure legends

Figure 1. The distribution of taluks (beige) across the general region of the Western Ghats comprises five states (Maharashtra, Goa, Karnataka, Kerala, and Tamil Nadu) in Peninsular India (left panel). Kyasanur Forest Disease virus outbreaks captured via ProMED-mail (red) and independently sourced (black) are superimposed over taluks. The distribution of mammalian species richness as estimated from a sample of Western Ghats species from 2010 to 2020 is presented in the right panel.

Figure 2. The distribution of Kyasanur Forest Disease virus outbreak risk derived from the inhomogeneous Poisson model is presented in the centre panel with the lower and upper 95% confidence limits to the left and right, respectively.

Figure 3. The adjusted distribution of mammalian species richness as estimated from a sample of Western Ghats species. Species richness was adjusted using the SESAM framework to introduce biotic constraints to adjust the presence of each species for the presence of other species within each taluk.

Figure 4. The distribution of Kyasanur Forest Disease virus outbreak risk derived from the integrated nested Laplace approximation model (binomial family) is presented in the centre panel with the lower and upper 95% credible limits to the left and right, respectively.

Figure 5. Taluks identified as priority targets (red) for implementing animal-human surveillance in Karnataka state. Having located priority taluks, a sample of these can be taken either by random draw or proportional to the distribution of key landscape features, such as species richness and/or forest loss. Once the taluks are selected, One Health surveillance infrastructure can be developed by 1) working in concert with the forest department to select sampling grids from within the sampled taluks for the monitoring and sampling of wildlife, 2) working in concert with animal husbandry and veterinary services to generate random samples of farms/livestock holders from animal censuses to monitor and sample cattle, and 3) working in concert with primary healthcare centres to improve human case detection and care delivery, conduct serosurveys, and implement prevention outreach.

Lower 95% C.L
□ under 0.011
□ 0.011 to 0.026
■ 0.026 to 0.052
■ 0.052 to 0.094
■ over 0.094

Fitted outbreaks
□ under 0.011
□ 0.011 to 0.026
■ 0.026 to 0.052
■ 0.052 to 0.094
■ over 0.094

Upper 95% C.L.
□ under 0.011
□ 0.011 to 0.026
■ 0.026 to 0.052
■ 0.052 to 0.094
■ over 0.094