Personalized immunopeptidomics using differential ion mobility technology enables direct profiling of neoantigens on colorectal cancer tissues

Yuriko Minegishi[1], Kazuma Kiyotani[2], Kensaku Nemoto[2], Yoshikage Inoue[3], Yoshimi Haga[1], Risa Fujii[1], Naomi Saichi[1], Satoshi Nagayama[4], Koji Ueda[1*]

[1] Cancer Proteomics Group, Cancer Precision Medicine Center, Japanese Foundation for Cancer Research, Tokyo, Japan.

[2] Project for Immunogenomics, Cancer Precision Medicine Center, Japanese Foundation for Cancer Research, Tokyo, Japan.

[3] Department of Surgery, Kyoto University, Kyoto, Japan.

[4] Development of gastroenterological surgery, Cancer Institute Hospital of Japanese Foundation for Cancer Research, Tokyo, Japan.

*Corresponding author: Koji Ueda

3-8-31, Ariake, Koto-ku, Tokyo, 135-8550, Japan

Tel.: +81 3 3570 0658, Fax.: +81 3 3570 0644, e-mail: koji.ueda@jfcr.or.jp

Keywords: Immunopeptidomics, HLA peptides, Mass spectrometry, FAIMS, differential ion mobility (DIM), neoantigen, cancer immune therapy

Main Text:

Knowing the nature of human leukocyte antigen (HLA)-presented peptides (HLAp) is indispensable to realize the precision medicine [1] [2] while direct identification of HLAp from scarce tissue specimens still confronts technical challenges [3]. Here we optimized the isolation procedures for HLAp and employed the technology of multiple gas phase fractionation differential ion mobility mass spectrometry (DIM-MS) to overcome that limitation. From $1e^8$ cells of HCT116, our method detected 9,249 unique HLAp including 11 peptides with missense mutations (neoantigens). Next, from individual differential immunopeptidome analysis using 40 mg each of colon cancer and adjacent normal tissues (n = 17), 44,785 HLAp including 2 neoantigens, $KRAS_{7-16}$(G12V) and $CPPED1_{226-234}$(R228Q), were identified. Furthermore, colon cancer-specific amino acid usage at C-terminus of HLAp was also found. Thus, this technology drastically improves the depth of immunopeptidomic analysis and allows direct determination of clinically important cancer antigens.

Neoantigen, a cancer specific somatic mutation-carrying HLAp, is presented on cellular surface as non-self antigen and activates the associated repertoire of T cells that will evoke anti-cancer immunity [1]. The immune checkpoint inhibitors (ICIs) have been established to invigorate this immune system as cancer immunotherapies [4]. Although immunotherapies are efficacious in patients with certain cancer types [4], there are still obstacles of less efficacy in cancers with less mutational burden such as microsatellite stable (MSS) colorectal cancer (CRC) compare to the microsatellite instable (MSI) CRC with DNA mismatch-repair deficiency [4, 5]. Validating and predicting the efficacy and choosing the right strategy for ICIs based on the background of individual patient are urgent tasks to grant the benefits by cancer precision immunotherapy. For that purpose, the direct scrutiny of HLAp on clinical tissue specimens could provide fundamental

information to facilitate them.

The immunopeptidomics by mass spectrometry (MS) is the only technology to directly

interrogate the HLAp. According to the guidelines for immunopeptidomics[3, 6], one of the

disadvantages of MS-based immunopeptidomics is incapability of robust analysis from a

scarce sample, such as tiny tissues of endoscopic biopsy, compared to the genomic

prediction of HLAp by *in-silico* [7]. To obtain the thousands scale of convincing

immunopeptidome with current technology, it generally requires at least $1e^8$ of cultured

cells or nearly 1000 μg of tissue sample [8-10] with inevitable chemical prefractionation and

this becomes hindrance for the analysis from scarce sample. Actually, direct detection of

neoantigens from tissue samples, such as colorectal, liver, or ovarian cancer, have been

previously unsuccessful [11, 12] [6]. Since the variety of HLAp is dynamic under cancer-

associating microenvironments including interferon gamma (IFN-γ) stimulation,

comprehending the HLAp under the native pathological environment is crucial for further

progress in cancer immunotherapies. Therefore, it is essential to establish more efficient

technology that enables the decent immunopeptidomics analysis from scarce tissues.

For that purpose, we adopted the differential ion mobility mass spectrometry (DIM-MS)

equipped with the interface of high field symmetric wave form ion mobility mass

spectrometry (FAIMS). The seamless gas phase fractionation by FAIMS secured the

sufficient depth of immunopeptidome from the scarce samples. First, we fully optimized

the parameters for DIM-MS to maximize the identification number of HLAp. Especially, a

set of 9 compensation voltages (CVs), from -80 to -35 V (see **Methods**), were selected

for the best performance of gas phase fractionation of HLAp. Other conditions such as

collision mode, detector type, maximum injection time, and scan speed were also fitted

accordingly (**Figure 1a, Methods**). By the FAIMS-assisted immunopeptidome analysis

using $5e^6$ cells of HCT116, 3,215 HLAp on average were identified, resulting in 1.3-fold

increase from the without-FAIMS condition (2,482 HLAp on average, **Fig. 1b**). Accordingly,

the number of identified source proteins was also greater in the FAIMS-assisted condition

(2,381 proteins on average) than the without-FAIMS condition (2,077 proteins on average,

Fig. 1b). This result was consistent throughout the experiments and became statistically

significant when the sample scale was increased to $1e^7$ cells (3,893 HLAp identification

on average) or $1.5e^7$ cells (4,160 HLAp identification on average, **Fig. 1b**). The frequency

of neoantigen identification was also enhanced accordingly to the sample scale by FAIMS

up to 2.4-fold (**Fig. 1c**). Thus, the DIM-applied gas phase fractionation provided

strengthened identification of HLAp as well as neoantigens. These results indicated the

potential of DIM-MS in immunopeptidomics from samples that even with an order of

magnitude smaller than the most previous reports [8] [11] [9].

From the biologically triplicated analyses using $1e^8$ cells of HCT116, 6,915 HLAp derived

from 4,109 source proteins were identified on average for a single analysis, which yielded

9,249 unique HLAp from 5,103 unique source proteins in total (**Fig. 1d, Supplementary**

Table 1). Importantly, 7.3 neoantigens on average were identified per independent

experiments, which yielded 11 neoantigens in total (**Fig. 1e**). Among these, 7 were not

found in the immune epitope database (IEDB) (http://www.iedb.org/home_v3.php, as of

November 27, 2020), hence regarded as newly identified ones (**Table 1**). The distribution

of length and the assignment of HLAp against HLA allotypes exhibited typical and

reasonable manner in the context of HLA genotype (**Supplementary Figure 1**). These

results indicated that the FAIMS-applied DIM-MS system could provide an ideal

technological platform for in-depth profiling of HLAp, as well as direct sequencing of

neoantigens.

We next applied this technology to personalized HLAp profiling analysis of CRC. In this

analysis, tumor or adjacent normal regions of CRC tissues (n = 17) were analyzed

(**Supplementary Table 2**). Based on the whole exome sequencing (WES) analysis, the

mutation-included proteome database was individually constructed and used for database

search analysis of the DIM-MS datasets. In total, 44,785 unique HLAp were identified

(6,578 peptides per patient on average) (**Fig. 2a**, **Supplementary Table 3**). Among these,

5,603 HLAp (12.5%) were normal-exclusively identified and 14,052 HLAp (31.4%) were

tumor-exclusive, while 25,130 HLAp (56.1%) were commonly detected from both normal

and tumor tissues. Thus we succeeded to improve the peptide identification number and

quantitative depth of immunopeptidome analysis drastically, compared to the previous

reports analyzing solid tumor tissues such as colon[6], liver[12], and ovarian[11] cancers. For

the 11,117 source proteins (4,149 proteins per patient on average), 817 (9.7 %) or 1839

proteins (16.5 %) were identified normal-exclusively or tumor-exclusively, respectively

(**Fig. 2b**). The individual datasets of immunopeptidomic profiling can be found in

Supplementary figure 2. The identified HLAp exhibited features of HLA-allotype

corresponding features of unsupervised clustering and assignment against the binding

motifs (**Supplementary figure 3**). The length of HLAp were variable from patient to patient

according to their HLA-allotypes, but the typical dominance of 9-mer in length of HLA class

I was common among patients and no statistically significant difference was found

between normal and tumor tissue HLAp (**Supplementary figure 4**). Notably, 2

neoantigens were identified from 2 distinct patients' tumor tissues (**Fig. 2c**). The first

neoantigen identified from the primary tumor tissue of ID 172 carried a well-known CRC

driver mutation, $KRAS_{7-16}$(G12V). The other one carrying a R228Q mutation of

$CPPED1_{226-234}$, a serine/ threonine-protein phosphatase, was identified from ID 261's

hepatic metastasized tumor tissue. According to the motif information at IEDB and

SYFPEITYHI (http://www.syfpeithi.de/0-Home.htm), the neoantigen carrying KRAS-G12V

was predicted to have higher affinity against A*11:01 than wild type sequence due to the

preferred amino acid substitution of valine at position 6 within the epitope (**Fig. 2c**,

depicted as "Preferred: V" by arrow). On the other hand, the mutation CPPED1-R228Q

lost its deleterious amino acid of arginine at position 3 by substitution to glutamine (**Fig.**

2c, depicted as "Deleterious: P/R" by arrow). Therefore the identified neoantigens

exhibited two opposite patterns of gaining affinity against A*11:01 by acquiring a preferred

residue or losing a deleterious residue within epitope, compared to the corresponding wild

type (WT) sequences. Indeed, the predicted IC_{50} value or $KRAS_{7-16}$(WT), $KRAS_{7-}$

$_{16}$(G12V), $CPPED1_{226-234}$(WT), or $CPPED1_{226-234}$(R228Q) was 299.7 nM, 137.3 nM, 729.9

nM, or 52.2 nM, respectively. It has been a long-standing challenge to identify neoantigens

directly from small clinical tissues, but our personalized immunopeptidomic analysis

overcame that task.

We next compared HLAp of three groups; HLAp only found in normal tissue (normal-

exclusive), HLAp found in both normal and tumor tissues (shared), and HLAp only found

in tumor tissue (tumor-exclusive) per patient. If there is a bias in the mechanisms of

antigen processing/ presentation between normal and tumor tissues, HLAp should

manifest with the characteristics by groups. When we checked the overall usage of amino

acid for all 44,785 HLAp, 12 amino acids covered 99.2% of C-terminus usage, in which

the lowest frequency was cysteine (0.5%) (**Fig. 2d**). In contrast, tryptic (R, K) and

chymotryptic (L, I, V, F, Y, W, A, M) amino acids were more frequently observed for HLAp

trimming at the end of C-terminus (pΩ). The usage of amino acid frequency at pΩ was

calculated as ratio (%) against subtotal of each groups (normal-exclusive, shared, and

tumor-exclusive peptide, respectively) and compared the difference against based on the

frequency of the shared-peptide to clarify whether there would be a processing bias shown

or not between normal-exclusive and tumor-exclusive groups. Among the 12 pΩ used

amino acids, 9 patients carried at least one R-restraint HLA allotypes (A*11:01 and

A*33:03) at pΩ, and the usage of R at pΩ exhibited the tendency of reduction in tumor-

exclusive HLAp (normal- exclusive vs tumor- exclusive: -0.06 % vs -5.10 % compared to

the shared-peptide frequency, p = 0.0975, **Fig. 2e**, [R]). The usage of K exhibited no

statistical difference compared to the frequency of shared-peptide (data not shown). The

R and the K are the same tryptic amino acids while exhibited distinct dynamics in normal

and tumor tissue. For chymotryptic peptides, only W exhibited statistically significant

increase in tumor-exclusive group compared to the shared-peptide (**Fig. 2e**, [W]). The

trimming at pΩ of HLAp by L, I, and V which were generally grouped as similar

hydrophobic amino acids exhibited inconsistent and non-significant results between

normal and tumor-exclusive HLAp (data not shown). Since the usages of these three

hydrophobic residues are naturally abundant among the HLAp, it is expected that

statistical significance was diminished and was not shown as major differences when

compared at the admixture condition of tissue. Thirteen out of 17 patients carry at least

one pΩ W-restraint HLA allotypes (A*24:02, B*4402 and B*44:03) and the usage of W at

pΩ was -0.45 % in normal-exclusive HLAp and +0.8 % in tumor-exclusive HLAp compare

to the shared-peptide group (p = 0.0185) (**Fig. 2e**, [W]). A negative correlation in W usage

at pΩ between normal-exclusive and tumor-exclusive HLAp in identical patients indicated

the shift of HLAp processing at pΩ in tumor tissue (r = -0.788, p = 0.0014) (**Fig. 2f**). In

addition to these two amino acids, the cysteine usage at C-terminus was found to be

increased in tumor-exclusive HLAp (+ 0.25 %) than that of normal-exclusive HLAp

(0.00 %) compared to the shared-peptide group (p = 0.016) (**Fig. 2e**, [C]). Since there was

no clear information available for the binding motif that requires cysteine at pΩ, all 17

samples are included for this calculation. It is of note that this distinct cysteine usage

between normal and tumor-exclusive peptide may imply the unknown mechanisms of

HLAp processing under carcinogenesis and of value for further investigation.

Through optimization of the protocols for HLAp isolation and the parameters for FAIMS-

applied DIM-MS measurement, we here established an ideal technology for

immunopeptidome profiling, allowing comprehensive HLAp profiling with clinically-

meaningful depth. Indeed, cancer specific signatures of HLAp, such as neoantigens and

altered amino acid motifs, were identified from a small amount of CRC tissues. Previously,

massive amount of sample was necessary for immunopeptidomic analysis especially from

clinical tissue samples [9, 11, 12]. Alternatively, patient-derived cancer cells have to be

prepared as primary culture to secure the enough sample amount [13, 14]. The purified

culture of patient-derived cancer cells is convenient for in-depth analyses, however, this

approach conceived possibility that prolonged cultivation affects the antigen presentation

due to the lack of cancer microenvironment. Although a patient-derived xenograft in

mouse is also feasible to secure the cancer cell mass, the response against ICIs may

differ in host organisms, as shown in the discrepancy result of clinical trial of MEK inhibitor

[15] with PD-L1 treatment [16]. Our advanced immunopeptidomics by FAIMS-applied DIM-MS

paved the way to identify approximately 5,000 of HLAp from 40 mg of tissue without bias-

prone chemical prefractionation procedures [17]. Indeed, since the peptides are fractionated

by CVs which uniquely reflect the structural bulkiness of the peptides [18], the majority of

HLAp were identified from -60 V to -40 V of CV regardless of HLA allotypes. In addition,

rapid transition time for CV stepping (25 ms/ transition) [19] enabled simultaneous

measurements with 3 CVs in a single LC/MS run. Eventually, the ratio of the number of

peptide spectrum match (PSM) to acquired MS/MS spectra was effectively high, 23.1 ±

1.1% on average. Thus, the DIM-MS is considered to be ideal platform for rapid and

comprehensive analysis of HLAp.

Biosynthesis of HLAp is affected by various external conditions such as stimulation by

IFN-γ that increases the number of HLAp by inducing immunoproteasome and modifies

the variation of HLAp [6, 31]. However, there was no report showing direct comparison of

HLAp profile in cancerous region with that in normal region using individual fresh tumor

tissues. In this study, we for the first time demonstrated the detailed difference between

HLAp presented on cancer cells and that on normal cells in individual CRC tissues. The

total number of identified HLAp was relatively more abundant in tumor tissues than normal

tissues although the difference was not significant when normalized by the protein amount

(**Supplementary Figure 5**). In the previous *in vitro* experiments under stimulatory

condition using IFN-γ, chymotryptic trimming of HLAp was reported to occur frequently.

However, among chymotrypsin-targeted amino acids, only tryptophan trimming was

statistically increased in the tumor regions of CRC tissues. Regarding trypsin-targeted

amino acids, arginine exhibited the tendency of decrease in the tumor regions of CRC

tissues, whereas the frequency of lysine at pΩ did not show significant difference. These

facts indicated that the cancer-specific biosynthesis of HLAp should not be regulated only

by expression level or activity of the trimming enzymes. To fully understand the specific

mechanism of HLAp presentation in cancer cells, further large-scaled and HLA allotype-

specific HLAp analysis is needed, in combination with comprehensive expression profile

of source proteins.

It has been suggested that the frequency of neoantigen presentation on solid tumors

might be low [6, 12, 13], especially in CRC [4, 5], because of eradication of neoantigen-presenting

cancer cells by anti-tumor immunity [20, 21]. However, we determined 2 neoantigens directly

from stage IV CRC tissues in this study. In that respect, the identification of $KRAS_{7-16}$(G12V) HLAp in this study is noteworthy because this may overturn the prevailing view

that cancer driver mutation-containing neoantigens would be depleted in the later clinical

stages of cancer. Since the KRAS-G12D-reactive autologous T cells in lung-metastasized

CRC was reported in the previous clinical trial of ICI [22], further identification and utilization

of neoantigens containing driver mutations in clinical settings would be anticipated.

Another source protein of neoantigen identified in this study, CPPED1, is known as a

negative regulator of Akt signaling, which directly dephosphorylates Akt at Ser473 [23-25].

Since the amino acid substitution R228Q was found within the phosphatase domain of

CPPED1, this may attenuate the phosphatase activity and induce aberrant activation of

Akt signaling in CRC cells. Therefore, therapeutic elimination of $CPPED1_{226-234}$(R228Q)-

presenting cancer cells could be also considered as an effective strategy to halt

progression of CRC.

In principle, more mutation burden is considered to be associated with better outcome

by ICI treatment [26]. However, in our immunopeptidome dataset, the frequency of

neoantigen presentation did not correlated with genomic mutation burden. Indeed, no

neoantigen was detected from the tissue specimen derived from the patient ID 119 who

carried the highest mutation burden (1,752), even though 6,983 HLAp were identified in

total. This fact stresses the importance of taking actually-presented HLAp profiles into

account in order to precisely estimate the effect of cancer immunotherapy. These in-depth

immunopeptidomic datasets could also lead to interpretation of the complicated biological

mechanisms for biogenesis and presentation of antigens.

Thus, our FAIMS-applied immunopeptidomic analysis drastically expanded the size of

immnopeptidome dataset. Particularly, for the first time, identification of neoantigens from

clinical tissue samples was successful. Moreover, individual immunopeptidome profiling

of normal and tumor tissues revealed the distinct features of C-terminal trimming of HLAp.

Such precision immunopeptidomics holds promise for opening a next phase of cancer

immunology with new outlook from both basic and clinical aspects.

Online Methods

Clinical samples and ethical approval

  Metastatic colon cancer tissues at pathological stage IV and its adjacent normal tissues were obtained from patients who all had provided written informed consent for this study. This study (ethical committee number 2010-1058) was approved by the ethical committee of the Japanese Foundation for Cancer Research (JFCR). The clinical information of them are listed in **Supplementary Table 2**. All tissue samples were frozen and maintained at least once before the MS sample preparation. Approximately 40 mg in net weight of tissue was dissected from the mass of tumor/ normal tissues respectively and further processed for immunopeptidomics.

Cell culture

  The colon cancer cell line HCT-116 was maintained under the general condition. Briefly, the cells were cultured in RPMI1640 media (WAKO) with 10% heat-inactivated fetal bovine serum (FBS), penicillin G and streptomycin. Cells were washed once with PBS before harvest. Desired number of cell pellets were stored at -80 °C until use.

In-house purification of antibody and the IP-beads preparation

  A hybridoma clone of anti-panHLA alpha chain antibody (HB95) was obtained from ATCC. Hybridoma cells were first expanded in 10% FBS containing DMEM and then cultured in CELL Line flask (WHEATON) with COS medium (COSMO BIO) for antibody production. After 1-week of cultivation, the culture media was collected and affinity purified by Protein A Fast flow (GE). Small amount of aliquots of eluates were electrophoresed and then stained by GelCode Blue Stain Reagent (Thermo Scientific) to check IgG fractions. The IgG enriched fractions were pooled and then dialyzed to replace the buffer into PBS. The antibody concentration was examined by BCA assay. To confirm the sufficient affinity of

our in-house purified W6/32 antibody against HLA molecules, we tested all batches of

W6/32 by general IP and Western blotting by antibody anti-pan HLA class I α-chain

molecules (clone EMR8-5, MBL) and β-2-microglobulin (β2M) antibody (PROTEINTECH)

in advance. High affinity confirmed W6/32 antibody (800 ug) was then cross linked onto

200 ul of slurry of Protein G Sepharose (GE) by DMP (20 mM) in HEPES (pH 8.4) for 1

hour. After quenching by TBS-T for 1 hour, beads were used for IP or stored at 4 °C before

use.

Immunoprecipitation of HLA-Class I Complex and HLAp purification for

immunopeptidomics

Desired cell pellet was lysed by 1 ml of lysis buffer containing [20 mM] HEPES, [150 mM]

NaCl, 1% NP-40, 0.1% SDS and 10% Glycerol on ice. For tissue samples, disposable

homogenizer Bio Masher System (nippi) was used firstly with 500 µl of lysis buffer for

approximately 40 mg of tissue on ice and then add another 500 µl of lysis buffer to make

sample volume as 1 ml. Both cells and tissues samples were cleared by centrifugation at

4°C, 15,000 rpm for 15min, the supernatant was collected and conducted

immunoprecipitation for 4°C overnight on rotating rotor. For thorough washing, Bio Spin

Column (BioRad) was used afterwards. The washing steps were as follows: 1 ml of lysis

buffer for 5 times, 1 ml of PBS-T for 5 times, 1 ml of PBS for 5 times, 1 ml of high-salt PBS

for twice, 1 ml of PBS for 5 times and then 1 ml of mass spectrometry-grade water for

twice. The trimer of HLA complex was dissociated and eluted by 500 µl of 1% TFA from

PI-beads. Obtained HLA enriched eluate was further processed by tC18 SepPak. For

class I HLA peptide, 500 µl of 20% acetonitrile (ACN) was firstly used for elution. HLA

peptide including eluates were then dried up by evaporator. To elute the other components

of HLA complex to confirm the IP-efficiency, 500 µl of 80% ACN was used for second

elution and the obtained eluates were dried up by evaporator as well. Sample purity was

assessed by silver staining (Silver Quest).

Culture Cells and the construction of WES-based customized database

A colon cancer cell line, HCT116, was maintained in 10% fetal bovine serum including

RPMI1640 in humidified atmosphere at 37°C with 5% CO_2. Customized database for

HCT116 was stablished from the WES data deposited in COSMIC Cancer Cell Line

Project. Listed somatic mutations were processed by in-house built pipeline previously

reported [27, 28]. Obtained mutations were then combined with Swiss-Prot human proteome

database (20,181 entries) from UniProt website and used for neoantigen search.

HLA typing, whole exome sequencing and pipe lines for tailored protein database

For clinical tissue samples, genomic DNA was extracted by QIAmp DNA mini Kit

(QIAGEN) from normal and tumor tissues. To establish the personalized database for

neoantigen identification by immunopeptodomics, whole exome sequencing (WES) was

conducted to all samples. In-house pipe line to extract the somatic mutations has been

previously reported [28, 29]. Briefly, the genome libraries for WES were established from the

DNA samples using xGen Exome Research Panel (IDT, Coralville, IA), according to the

manufacturer's instructions. The exons were sequenced as 150 base pairs paired-end

reads by a NovaSeq6000 system (Illumina, San Diego, CA). The sequence data obtained

were then analyzed to select possible germ line variations and somatic mutations. In short,

sequences of the whole exome were compared with a reference human genome (hs37d5),

and the possible germ line variants and the somatic mutations were identified. The

variants in public databases (> 1%) were also excluded. Then cancer somatic mutations

were extracted by subtraction of germ line variation identified in normal tissue from tumor

tissue respectively. Among those mutations, only the missense mutations were selected

and translated into full length of protein and added to the same Swissprot/ Uniprot human

database described above as personalized database. The established personalized data

base was used for immunopeptidomics both normal and tumor tissues corresponding

patient's samples. The OptiType was employed for 4-digit HLA typing from deep

sequencing data [30]. The genetic details including mutation burden, oncogenic KRAS

mutations and the HLA genotypes of clinical samples can be found in **Supplementary**

Table 2.

Differential ion mobility (DIM) Mass Spectrometry (MS) by High-Field Asymmetric-

waveform Ion Mobility Spectrometry (FAIMS) for Advanced Immunopeptidomics

The sample includes HLAp was first trapped by precolumn (C18 Accleim PepMap 100

C18 Trap Cartridge, Thermo Scientific) and then separated by analytical column (Aurora

UHPLC Column (C18, 0.075 × 250 mm, 1.6 um FSC, ESI, ion opticks) coupled with a

nanospray Flex ion source for electrospray ionization (Thermo Scientific), were used

through an Ultimate 3000 RSLC nano HPLC system at a flow rate of 200 nl/ min using a

linear gradient starting from 2% solvent B (0.1% Formic acid in acetonitrile) to 28 over 55

min, a 1 min hold at 95% Solvent B a prior to a 1 min analytical column equilibration with

2% B. In order to validate the benefit of FAIMS applied DIM-MS in immunopeptidomics,

liquid chromatography tandem mass spectrometry (LC-MS/MS) and LC-FAIMS-MS/MS

conditions were firstly compared. For LC-FAIMS-MS/MS, the FAIMS-Pro interface

(Thermo Fisher Scientific) was installed onto an Orbitrap Fusion Lumos Tribrid mass

spectrometer and operated with default parameters except for the compensation voltage

(CV) settings for gas-phase fractionation. For LC-MS/MS, we uninstalled FAIMS device

from mass spectrometer and performed analyses. Under LC-FAIMS-MS/MS condition,

sample was seamlessly fractioned by 3 CVs per single analysis. Total 3 CV sets were

used per sample and each CV set includes as follows; set1 (CV = -40, -60, -80 V), set 2

(CV = -35, -50, -70 V) and set 3 (CV = -45, -55, -65 V). Therefore, total 3 analyses (raw

data), were acquired per sample. The parameters for mass spectrometry were optimized

for immunopeptidomics with small amount of sample. Full MS (rage from 320 to 850 m/z)

in the Orbitrap were acquired at resolution of 60 k followed by a MS^2 acquisition at

resolution of 15k. Maximum injection time for full MS scan was 50 ms with an auto gain

control (AGC) of $4e^5$. Maximum injection time for MS^2 scan was 100 ms with an AGC $1e^4$

followed by top speed MS^2 acquisition by the ion trap. Charge state 2 and 3 were selected

for fragmentation by collision-induced dissociation (CID) at rapid scan rate at 30% collision

energy. The easy-IC system was used for the internal calibration lock mass during the

data acquisition. The HCT116 samples were prepared from $1e^8$ cells and proportionally

injected into MS to match the indicated number of cells. For the analyses of $1e^8$ of HCT116

cells and clinical tissue samples, firstly 1/ 20 amount of sample volume was used for

analyses of 3 CV sets respectively to check sample conditions, and then from the

remaining, 5/ 20 amount of sample volume was used for 3 CV sets respectively. Therefore,

total 6 analyses were performed per clinical tissue sample.

The LC/MS raw data and summarized result files are deposited at a public proteomic

database, Japan Proteome Standard Repository/Database (jPOST) as follows; HCT116

HLAp without FAIMS in JPST001072, HCT116 HLAp with FAIMS in JPST001066, Global

identification of HCT116 HLAp in JPST001068, HLAp from normal regions of CRC tissues

in JPST001070, and HLAp from tumor regions of CRC tissues in JPST001069.

Data analysis by tailored database search for immunopeptidome

Obtained raw files were processed by Proteome Discoverer version 2.4 using Sequest

HT engine against corresponding tailored database. The search was set to no-enzyme

(unspecific) for the enzymatic designation. The precursor mass tolerance was set to 10

ppm and the fragment mass tolerance to 0.6 Da. The maximum number of missed

cleavages was set to zero. Methionine oxidation and cysteine carbamidomethylation were

included as dynamic modifications. A false discovery rate (FDR) of 0.01 (Strict) was used

for percolator algorithm to identify highly confident peptides. The output "peptide groups"

was exported as a list. Then the peptide sequence was deduplicated for unique peptide

sequences only and filtered by the amino acid length from 8 to 15 as a candidate of HLAp.

To narrow candidate sequences down to the reliable ones, NetMHCpan 4.0[31] was

employed to exclude the no-binding peptides from the list. Simultaneously, unsupervised

clustering for HLA sequence motifs, Gibbs Cluster 2.0 [32] and Seq2 logo [33] were employed

to confirm whether the identified motifs match HLA allotypes accordingly.

Profiling of HLAp in Normal and Tumor Tissues

In order to interrogate the possible distinctions of normal and tumor-derived HLAp, the

intersection of HLAp between normal and tumor tissues was calculated and draw Venn

diagrams　　　　by　　　　publicly　　　　available　　　　website

(http://bioinformatics.psb.ugent.be/webtools/Venn/). Then classified the HLAp into 3

groups; (1) normal-exclusive, (2) shared in both normal and tumor tissues (as "shared-

peptide") and (3) tumor-exclusive ones per patient. The frequency of amino acid usage at

position Ω (pΩ), the end position at C-terminus of HLAp, was calculated as percentage

against the total number of HLAp in each group. Based on the frequency of shared-

peptide's amino acid usage at pΩ, the distinction of frequency was calculated by

comparing the value between normal-exclusive and shared-peptide or tumor-exclusive

and shared-peptide.

Statistical Analysis.

All data were expressed as mean ± standard error (SE). Student's t-test was used for

statistical analyses. Pearson correlation-coefficient was used to assess the correlation.

Statistical significance thresholds were determined at $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$

(***) values. Pearson correlation-coefficient was used to assess the correlation.

Acknowledgements

This work was supported by "the Development of Technology for Patient Stratification

Biomarker Discovery" of the Japan Agency for Medical Research and Development

(20ae0101074s0302) and "Grant-in-Aid for Scientific Research (C)" of Japan Society for

the Promotion of Science (20K05759).

Author Contributions

Y.M. and K.U. designed the experiments. Y.M., K.K., K.N., Y.H., R.F., and N.S. performed

the experiments. Y.I. and S.N. provided the CRC tissue specimens. Y.M., K.K., K.N., Y.H.,

R.F., and N.S. analyzed and interpreted the data. Y.M. and K.U. wrote the article.

Competing Interests statement

The authors declare no competing interests.

References:

1. Jhunjhunwala, S., Hammer, C. & Delamarre, L. Antigen presentation in cancer: insights into
tumour immunogenicity and immune evasion. *Nat Rev Cancer* (2021).
2. Cancer Genome Atlas, N. Comprehensive molecular characterization of human colon and
rectal cancer. *Nature* **487**, 330-337 (2012).
3. Caron, E., Aebersold, R., Banaei-Esfahani, A., Chong, C. & Bassani-Sternberg, M. A Case for
a Human Immuno-Peptidome Project Consortium. *Immunity* **47**, 203-208 (2017).
4. Le, D.T. et al. PD-1 Blockade in Tumors with Mismatch-Repair Deficiency. *N Engl J Med*
372, 2509-2520 (2015).
5. de Weger, V.A. et al. Clinical effects of adjuvant active specific immunotherapy differ between
patients with microsatellite-stable and microsatellite-instable colon cancer. *Clin Cancer Res*
18, 882-889 (2012).
6. Loffler, M.W. et al. Mapping the HLA Ligandome of Colorectal Cancer Reveals an Imprint of
Malignant Cell Transformation. *Cancer Res* **78**, 4627-4641 (2018).
7. Shukla, S.A. et al. Comprehensive analysis of cancer-associated somatic mutations in class I
HLA genes. *Nat Biotechnol* **33**, 1152-1158 (2015).
8. Bassani-Sternberg, M. et al. Direct identification of clinically relevant neoepitopes presented
on native human melanoma tissue by mass spectrometry. *Nat Commun* **7**, 13404 (2016).
9. Chong, C. et al. High-throughput and Sensitive Immunopeptidomics Platform Reveals
Profound Interferongamma-Mediated Remodeling of the Human Leukocyte Antigen (HLA)
Ligandome. *Mol Cell Proteomics* **17**, 533-548 (2018).
10.   Liepe, J. et al. A large fraction of HLA class I ligands are proteasome-generated spliced
peptides. *Science* **354**, 354-358 (2016).
11.   Schuster, H. et al. The immunopeptidomic landscape of ovarian carcinomas. *Proc Natl*
Acad Sci U S A **114**, E9942-E9951 (2017).
12.   Loffler, M.W. et al. Multi-omics discovery of exome-derived neoantigens in
hepatocellular carcinoma. *Genome Med* **11**, 28 (2019).
13.   Newey, A. et al. Immunopeptidomics of colorectal cancer organoids reveals a sparse
HLA class I neoantigen landscape and no increase in neoantigens with interferon or MEK-
inhibitor treatment. *J Immunother Cancer* **7**, 309 (2019).
14.   Demmers, L.C. et al. Single-cell derived tumor organoids display diversity in HLA
class I peptide presentation. *Nat Commun* **11**, 5338 (2020).
15.   Eng, C. et al. Atezolizumab with or without cobimetinib versus regorafenib in
previously treated metastatic colorectal cancer (IMblaze370): a multicentre, open-label, phase
3, randomised, controlled trial. *Lancet Oncol* **20**, 849-861 (2019).
16.   Ebert, P.J.R. et al. MAP Kinase Inhibition Promotes T Cell and Anti-tumor Activity in
Combination with PD-L1 Checkpoint Blockade. *Immunity* **44**, 609-621 (2016).

17.   Demmers, L.C., Heck, A.J.R. & Wu, W. Pre-fractionation Extends but also Creates a
Bias in the Detectable HLA Class Iota Ligandome. *J Proteome Res* **18**, 1634-1643 (2019).

18.   Purves, R.W. et al. Using gas modifiers to significantly improve sensitivity and
selectivity in a cylindrical FAIMS device. *J Am Soc Mass Spectrom* **25**, 1274-1284 (2014).

19.   Hebert, A.S. et al. Comprehensive Single-Shot Proteomics with FAIMS on a Hybrid
Orbitrap Mass Spectrometer. *Anal Chem* **90**, 9529-9537 (2018).

20.   Rooney, M.S., Shukla, S.A., Wu, C.J., Getz, G. & Hacohen, N. Molecular and genetic
properties of tumors associated with local immune cytolytic activity. *Cell* **160**, 48-61 (2015).

21.   Davoli, T., Uno, H., Wooten, E.C. & Elledge, S.J. Tumor aneuploidy correlates with
markers of immune evasion and with reduced response to immunotherapy. *Science* **355**
(2017).

22.   Tran, E. et al. T-Cell Transfer Therapy Targeting Mutant KRAS in Cancer. *N Engl J*
Med **375**, 2255-2262 (2016).

23.   Zhuo, D.X. et al. CSTP1, a novel protein phosphatase, blocks cell cycle, promotes cell
apoptosis, and suppresses tumor growth of bladder cancer by directly dephosphorylating Akt
at Ser473 site. *PLoS One* **8**, e65679 (2013).

24.   Zhuo, D., Wu, Y., Luo, J., Deng, L. & Niu, X. CSTP1 inhibits IL-6 expression through
targeting Akt/FoxO3a signaling pathway in bladder cancer cells. *Exp Cell Res* **380**, 80-89
(2019).

25.   Yang, R. et al. Combined Transcriptome and Proteome Analysis of Immortalized
Human Keratinocytes Expressing Human Papillomavirus 16 (HPV16) Oncogenes Reveals
Novel Key Factors and Networks in HPV-Induced Carcinogenesis. *mSphere* **4** (2019).

26.   Hugo, W. et al. Genomic and Transcriptomic Features of Response to Anti-PD-1
Therapy in Metastatic Melanoma. *Cell* **165**, 35-44 (2016).

27.   Kato, T. et al. Effective screening of T cells recognizing neoantigens and construction
of T-cell receptor-engineered T cells. *Oncotarget* **9**, 11009-11019 (2018).

28.   Kiyotani, K., Chan, H.T. & Nakamura, Y. Immunopharmacogenomics towards
personalized cancer immunotherapy targeting neoantigens. *Cancer Sci* **109**, 542-549 (2018).

29.   Choudhury, N.J. et al. Low T-cell Receptor Diversity, High Somatic Mutation Burden,
and High Neoantigen Load as Predictors of Clinical Outcome in Muscle-invasive Bladder
Cancer. *Eur Urol Focus* **2**, 445-452 (2016).

30.   Szolek, A. et al. OptiType: precision HLA typing from next-generation sequencing
data. *Bioinformatics* **30**, 3310-3316 (2014).

31.   Jurtz, V. et al. NetMHCpan-4.0: Improved Peptide-MHC Class I Interaction Predictions
Integrating Eluted Ligand and Peptide Binding Affinity Data. *J Immunol* **199**, 3360-3368
(2017).

32.   Andreatta, M., Alvarez, B. & Nielsen, M. GibbsCluster: unsupervised clustering and
alignment of peptide sequences. *Nucleic Acids Res* **45**, W458-W463 (2017).

33.   Thomsen, M.C. & Nielsen, M. Seq2Logo: a method for construction and visualization

of amino acid binding motifs and sequence profiles including sequence weighting, pseudo

counts and two-sided representation of amino acid enrichment and depletion. *Nucleic Acids*

Res **40**, W281-287 (2012).

Figure Legends

Figure 1. Schematic workflow and data validation of personalized immunopeptidome analysis assisted by differential ion mobility mass spectrometry (DIM-MS). a. A proteogenomic workflow for personalized immunopeptidome analysis is shown. After constructing the personalized proteome database in which cancer-specific mutated amino acid sequences were involved, the DIM-MS analysis was performed for isolated HLAp. The FAIMS-Pro interface was coupled to mass spectrometer for seamless gas-phase fractionation to enable the DIM-MS. CV; compensation voltage. **b.** The number of identified HLAp from HCT116 cells was displayed with bars and the left y-axis (white bars; without FAIMS, black bars; with FAIMS). The number of identified source proteins was shown with the dashed line and the right y-axis. * $p < 0.05$, # $p < 0.1$. **c.** The average number of identified neoantigens from the same experiments of b was shown (white bars; without FAIMS, black bars; with FAIMS). * $p < 0.05$, ** $p < 0.01$. **d.** The number of identified HLAp from three independent analyses using $1e^8$ of HCT116 cells was shown with bars and the left y-axis. White bars; the number of HLAp from respective analyses, gray bar; an average number of HLAp from 3 independent analyses, and dark gray bar; total number of non-redundant identified HLAp. The number of source proteins were depicted in dashed line and the right y-axis. **e.** The number of neoantigens identified from the same experiments as d was shown.

Figure 2. Personalized differential immunopeptidome analysis using cancer tissues and adjacent normal tissues of colorectal cancer (n = 17). a. An average of total number of HLAp identified per patient (HLAp/ Patient) and the total number of unique HLAp from all patients (Total Unique HLAp) were displayed in bar graphs with respective numerals. **b.** An average of total number of source proteins identified per patient (Protein/ Patient) and the total number of unique HLAp from all patients (Total Unique Protein) were

displayed in bar graphs with respective numerals. Full information for 17 patients can be

found in **Supplementary Table 3**. **c.** The MS/MS spectra for $KRAS_{7-16}$(G12V) and

$CPPED1_{226-234}$(R228Q) neoantigens were shown. Possible causes of enhanced affinity

by mutations were also depicted with motif sequence of A*11:01. **d.** A pie chart depicts the

frequency of amino acid usage at the C-terminus (pΩ) of HLAp. **e.** The difference of amino

acid usage (%) at pΩ in normal-exclusive and tumor-exclusive HLAp compared to the

shared-HLAp. **f.** The dot plot demonstrates negative correlation of tryptophan (W) usage

at pΩ in normal-exclusive and tumor-exclusive HLAp.

Table 1. Neoantigens identified from HCT116 by immunopeptidomic analysis

Mutation	Neoantigen Sequence	Assigned HLA	Affinity [nM]	Detection	First Report (*)
					No WT in IEDB (#)
AGO2-H336Y	QEQKYTYLP	B4501	980.7	1/3	*
CHMP7-A324T	QTDQMVFNTY	A0101	33.48	3/3	
FNBP4-K318E	EEEEKGVAA	B4501	83.24	3/3	*
NAPA-A181V	KVIDIYEQV	A0201	50.07	3/3	*
NR1D1-G39D	YSDNSNDSF	A0101/ C0701	28.34/ 5.89	3/3	
GAPDH-I69T	AENGKLVTN	B4501	963.6	3/3	*, #
IQGAP-S1070T	VLEDKVLTV	A0201	35.1	1/3	*
RBBP7-N17D	EERVIDEEY	B1801	194.48	1/3	
NOTCH2-C41S	NEGMSVTY	B1801	6.2	2/3	*, #
PDP1-N379D	DEYTKFIPP	B1801	158.49	1/3	*
UQCRB-N88K	EEEKFYLEP	B4501	728.67	1/3	

The position of mutation is highlighted by underline. The affinity prediction by NetMHC4.0 was shown. The number of times identified out of 3 independent MS analyses was shown.

* Newly identified neoantigens.

There was no epitope information of the corresponding wild type sequence at IEDB.

Figure 1

Figure 2

Supplementary Figure 1

Supplementary Figure 1. Characteristics of immunopeptidomics and neoantigen identification from HCT116 cells

a. The length of HLAp identified from HCT116 in Figure 2d was counted. **b.** Representative image of unsupervised HLA motif clustering and the population of HLA assignment against genotype-matched HLA allotypes.

Supplementary Figure 2

Supplementary Figure 2. Number of HLAp per patients and Venn diagrams

The details of identified number of HLAp in each normal and tumor tissue samples and the overlap of HLAp in identical patients were shown. Venn diagram does not reflect the sample size in this figure.

Supplementary Figure 3

Supplementary Figure 3. Individual dataset for HLA allotypes and identified HLAp

HLA allotype and its motif sequence, unsupervised clustering of actually identified HLAp, HLAp assignment to corresponding HLA allotypes, and the length distribution of HLAp were shown individually. The details of identified HLAp in each patient was validated by using Gibbsclustering2.0, NetMHCpan. The length of HLAp were compared between normal and tumor tissues in identical patients.

Supplementary Figure 4

Supplementary Figure 4. Comparison of personalized immunopeptidome analysis for CRC tissue samples

Length distribution of HLAp shown in Figure 2a.

Supplementary Figure 5

Supplementary Figure 5. Sample conditions and their correlation with identified HLAp

Box plots depict **a.** net weight of tissues, **b.** total number of identified HLAp, **c.** protein amount included in lysate and **d.** the number of identified HLAp normalized by protein amount. There was no statistically significant increase of HLAp presentation in tumor tissues compare to the normal tissue when the identified number of HLAp was normalized by protein amount in the starting sample of lysate. **e.** A composite graph of depicts the identified number of HLAp in each samples (white bars indicates normal tissues, gray bars indicates tumor tissues) and the protein amount used. The identified number indicates the variation of HLAp and is vary from sample to samples. Asterisk indicates the sample in which neoantigen was identified respectively.