Shorter effective lifespan in laboratory populations of *D. melanogaster* can reduce sexual

selection

Avani Mital[1#], Manaswini Sarangi[2], Bodhisatta Nandy[3], Neha Pandey[4], Amitabh Joshi[5]

Evolutionary Biology Laboratory, Jawaharlal Nehru Centre for Advanced Scientific Research, Bengaluru 560064

India.

1: Avani Mital (# Corresponding author)

Present address: Biology Division, Department of Physics, Chemistry and Biology (IFM), Linköping University

Campus Valla, 58183, Sweden.

avani.mital@liu.se

2: Manaswini Sarangi

Present address: 4202 Biological Sciences Building, University of Michigan, MI 48109, USA

msarangi@umich.edu

3. Bodhisatta Nandy

Present address: Department of Biological Sciences, Indian Institute of Science Education and Research,

Berhampur 760010 India

nandy@iiserbpr.ac.in

4.nehapande2@jncasr.ac.in

5. ajoshi@jncasr.ac.in

Acknowledgment: We are grateful to Sajith V.S, Rajanna N., and Muniraju P. for help with the experiments. This work was funded by Jawaharlal Nehru Centre for Advanced Scientific Research (in-house funding to AJ), the Science and Engineering Research Board (SERB) JC Bose fellowship (recipient: AJ), and in part by personal funds of AJ. JNCASR provided doctoral fellowships to AM, MS, and NP, and a postdoctoral fellowship to BN.

Conflict of interest: The authors declare no conflict of interest.

Author contributions: AJ, AM and BN developed the overall theme of the study; AM and BN designed the specific experiments; AM, MS, BN and NP carried out the experiments; AM analysed the data and wrote the first draft of the paper; all authors contributed to the final draft.

Data Availability statement: The raw data based on which we report results in this article is available as supplementary material (SM_2).

Abstract

The role of sexual selection in mediating levels of sexual conflict has been demonstrated in many experimental evolution studies on *Drosophila* sp. where competition among males for mating was the target of selection. Sexual selection has also been shown to affect the evolution of life-histories. However, the influence of divergent life-histories on reproductive strategies and, therefore, sexual selection and sexual conflict, have been less well studied. We investigated *D. melanogaster* populations selected for a short development time and early age at reproduction for changes in reproductive behaviour and levels of sexual selection. We report a large reduction in reproductive competition experienced by the males of these populations, compared to their ancestral controls, leading to reduced levels of sexual selection. We show that rapidly developing and early reproducing populations have very low levels of mating in their lifetime (are more or less monogamous), low courtship levels, shorter copulation duration and longer maturation time, compared to their ancestral controls. These results are discussed in the context of the previously demonstrated reduction of sexual conflict in these populations. We show that life-history strategies can have a large and significant impact on sexual selection, with each influencing the other and contributing to the complexities of adaptation.

Key Words: Sexual conflict, life-history evolution, male-male competition, courtship, reproductive behaviour, experimental evolution.

Significance statement

Sexual conflict, an arms-race between males and females trying to enhance their own reproductive success at some cost to the other, is of great evolutionary interest because it can also promote speciation. The intensity of sexual selection, or variation in mating success, is well known to affect levels of sexual conflict. However, it is not so clear whether, and how, the regular evolution of life-histories affects levels of sexual conflict. Here, we show that life-history evolution in fruit fly populations selected for traits not directly related to sexual conflict can, nevertheless, mediate the evolution of altered sexual conflict levels through effects on the intensity of sexual selection. Populations that evolved to develop to adulthood fast, and reproduce relatively early in life, are shown to undergo less sexual selection, which can explain the low sexual conflict levels earlier observed in them.

Introduction

Due in part to Darwin's independent treatment of natural selection (Darwin 1859, 1868) and sexual selection (Darwin 1871), the two sub-fields have largely developed separately, and the causes and consequences of viability/fecundity selection and sexual selection have often been studied by evolutionary biologists with somewhat differing backgrounds and interests. An ironic consequence of this outcome has been that studies of life-history evolution and sexual selection did not meaningfully intersect for many decades, even though they both focus on opportunities for, and timing and distribution of, reproductive output, albeit from somewhat different points of view. Parental investment per offspring differs between males and females of most sexually reproducing species (Bateman 1948; Trivers 1972), resulting in different reproductive strategies for male or female fitness maximization, often leading to differences in optimal mating rates for males and females (Bateman 1948; Andersson 1994). Such differences can give rise to inter-locus sexual conflict (Parker 1979; Chapman et al. 2003; Anrqvist and Rowe 2013), resulting in arms-race like dynamics with males evolving to manipulate female reproductive choices, and females, in turn, evolving to circumvent such manipulation (Chapman et al. 2003; Anrqvist and Rowe 2013). Sexual selection and sexual conflict are clearly intertwined and, indeed, have been studied together in some detail over the past few decades (e.g., Wigby and Chapman 2004; Linklater et al. 2007; Edwards et al. 2010; Nandy et al. 2013a, b). However, even though viability and fecundity selection can shape life-histories in ways that can either heighten or reduce sexual selection and, thereby, sexual conflict, life-history evolution has not yet been integrated with studies of sexual selection and sexual conflict in similar detail. In this paper, we report a reduced opportunity for sexual selection in laboratory populations of *Drosophila melanogaster* (first described in Prasad et al. 2000) that have earlier been shown to have evolved reduced levels of inter-locus sexual conflict as a correlated response to selection for rapid egg-to-adult development and early reproduction (Ghosh and Joshi 2012; Mital et al. 2021). This is, to our knowledge, one of very few studies to directly connect patterns of life-history evolution to altered levels of sexual selection and sexual conflict.

Several studies on insects have investigated how reproductive strategies may evolve upon changing their breeding ecology. These studies establish a strong relationship between levels of sexual selection, achieved by manipulating the degree of competition among males for mating, and sexual conflict related traits. For example, populations of *Drosophila* sp. were subjected to either monogamy (Holland and Rice 1999; Pitnick et al. 2001; Crudgington et al. 2005; Hollis et al. 2014; Wensing et al. 2017), or different operational sex ratios (Wigby and Chapman 2004; Linklater et al. 2007; Edwards et al. 2010; Nandy et al. 2013a, b) in order to experimentally alter the degree of

polygamy and, therefore, sexual selection experienced by the flies. Traits that promote male-specific fitness

reduced in the monogamy adapted populations (Holland and Rice 1999; Pitnick et al. 2001; Crudgington et al.

2005; Wensing et al. 2017) and females from these populations experienced greater fitness loss when paired with

males adapted to polygamy than to monogamy (Holland and Rice 1999; Crudgington et al. 2005). Similarly,

females from populations with male-biased sex ratio had higher resistance to mate harm compared to females

from female biased sex ratio populations (Wigby and Chapman 2004; Nandy et al. 2014).

There is also some evidence for sexual selection affecting life histories. For example, Hollis et al (2017) show the

evolution of faster development and maturation in monogamous compared to polygamous *D. melanogaster*

populations. In males of the decorated cricket, *Gryllodes pirillas*, increasing reproductive effort with age was

shown to correlate with slower ageing and longer lifespan (Archer et al. 2012). Similarly, Zajitschek et al. (2009)

demonstrated sexual dimorphism in age dependent patterns of survival and reproduction in the field cricket

Teleogryllus commodus, likely due to differences is age-dependent reproductive strategies between males and

females. Recently, evidence for the evolution of longer development times and decreased desiccation and

starvation resistance in polyandrous *D. pseudoobscura* lines compared to monogamous lines has also been

reported (Garlovsky et al. 2021).

The direct influence of life-history changes on levels of sexual selection and sexual conflict has been investigated

relatively rarely, even though these influences can potentially have large effects on both sexual selection and

sexual conflict. For example, a life-history providing a relatively short duration of time for reproduction can

reduce sexual selection by constraining opportunities to re-mate, driving lower levels of competition among males.

The latter, in turn, could drive an evolutionary reduction in sexual conflict levels. Thus, effective adult life span,

in particular, can alter the level of sexual selection, all else being equal. For example, selection for early

reproduction in seed beetles *Callosobruchus maculatus* resulted in the evolution of more frequent early-life mating

compared to those selected for late reproduction (Maklakov et al. 2010). Similarly, populations of *D.*

melanogaster that experienced longer durations of effective adult life evolved increased male offense and defense

ability at late ages and an improved ability to induce typical female post mating responses, compared to their

controls (Service 1993; Service and Fales 1993; Service and Vossbrink 1996).

One of the clearest demonstrations of life-history evolution affecting sexual conflict levels has come from a study

showing considerably reduced sexual conflict in populations of *D. melanogaster* subjected to long term selection

for rapid development and early reproduction in the laboratory relative to that in ancestral control populations

(Ghosh and Joshi 2012; Mital et al. 2021). This was shown to be largely driven by the much smaller body size of

flies from the selected populations (Mital et al. 2021), a consistent correlated response to strong selection for rapid

development in *D. melanogaster* across multiple studies (Zwaan et al. 1995; Nunney 1996; Chippindale et al.

1997a; Prasad et al. 2000). The observation that size reduction in these selected populations likely drove the

evolution of reduced levels of sexual conflict appears to be consistent with studies showing smaller males to be

less harmful to females than larger males (Partridge et al. 1987a, b; Pitnick 1991; Pitnick and Garcia-Gonzales

2002).

In addition to being selected for rapid development, which results in reduced body size, the selected populations

used by Ghosh and Joshi (2012) were also selected for relatively early reproduction (around day 3-4 of adult life)

compared to controls (around day 10-12 of adult life). The selected populations, thus, have an effective lifespan

that is about one-third that of the controls. In a 3-4 day adult lifespan in this system, there is limited scope of

multiple matings by females, especially if smaller, resource deprived females fail to remate readily. This can result

in a significant reduction in the intensity of sexual selection and sexual conflict in these populations (Ghosh and

Joshi 2012; Mital et al. 2021). Therefore, in these rapidly developing populations, selection for early reproduction

might also have led to further reduction in sexual conflict, beyond that due to the smaller body size of selected

population males (Mital et al. 2021). Here, we examine some aspects of the breeding ecology of these populations

to establish if there are differences in levels of sexual selection between selected and control populations.

We looked at lifetime mating frequency, courtship frequency and copulation duration in our selected and control

populations. Mating frequency has been used to estimate extent of sexual selection previously (Kuijper and

Morrow 2009). Courtship frequency is indicative of male mating effort since *D. melanogaster* males perform

extensive and energetically expensive courtship for which they likely bear a fitness cost (Cordts and Partridge

1996; Anholt et al. 2020). Moreover, courtship by males potentially provides an opportunity for females to

exercise mate choice (Gavrilets et al. 2001; see Anholt et al. 2020 for review). Copulation duration is another

estimate of reproductive effort/ investment by males as copulation in *D. melanogaster* lasts substantially longer

than required for sperm transfer alone, with the additional time spent in the transfer of accessory gland proteins

(Acps) (Gilchrist and Partridge 2000). Many of these Acps are known to mediate sexually antagonistic effects in

females (Chapman et al. 1995; Wolfner et al. 1999; Chapman 2001). We also estimated the time from eclosion to

first mating by females (maturation time), which was shown earlier to have lengthened in our selected populations,

relative to controls (Prasad 2004; Ghosh-Modak 2009).

Methods

<u>Study Populations</u>

We used eight large, outbred *D. melanogaster* populations that have a common ancestry. Four of these

populations were selected for rapid pre-adult development and early reproduction and are referred to as the **FEJ**s

(**F**aster developing, **E**arly reproducing, **J**B derived); the other four were ancestral controls, called the **JB**s (**J**oshi

Baseline). Details of the ancestry of these populations and their maintenance have been reported previously (JB:

Sheeba et al. 1998; FEJ: Prasad et al. 2000).

In summary, JBs are maintained on a 21day discrete generation cycle with eggs collected into 40 replicate vials

(9.5 cm ht × 2.4 cm dia) per population (60-80 eggs/ 6mL of banana-jaggery medium). All adults typically emerge

by the 12[th] day from egg collection and are transferred to fresh food vials on days 12, 14 and 16. On day 18, all

flies are collected into to a Plexiglas cage ($25 \times 20 \times 15$ cm^3) provided with food supplemented with additional

yeast for three days before collecting eggs to start a new generation. The FEJ maintenance is similar except that

only the first 20-25% of eclosing flies from each vial are collected directly into a Plexiglas cage with food

supplemented with additional yeast. After three days, eggs are collected to initiate the next generation. All

populations are maintained at a breeding adult number of 1500-1800 flies, under constant light, 25ºC ± 1ºC and

about 90% relative humidity. Thus, only the fastest developing flies in the FEJs make it to the breeding pool and

reproduce relatively early, i.e., on day three after eclosion. Consequently, by the time of this study (~600

generations of FEJ selection), FEJs were being maintained on a 10-day discrete generation cycle. Since each of

the four FEJ populations has been derived from one JB population, we could account for ancestry by treating FEJs

and JBs with matched subscripts as random blocks in the statistical analyses. Standardization (common control-

type rearing conditions for a full generation in order to equalize non-genetic parental effects) was done only prior

to assaying reproductive maturity and copulation duration in the FEJ and JB populations. It was not possible to

record behaviour data blind as the focal flies from the JB and FEJ populations are easily distinguishable based on

their body size.

<u>Mating and Courtship Frequencies</u>

For this assay, we used flies derived from eggs laid in the running cultures, without standardization, as

we wanted to assess breeding ecology differences that FEJ and JB flies experience during their maintenance,

including those potentially due to non-genetic parental effects. For JBs, flies from these assay populations were

collected on day 12 from egg-lay, pairing five males and five females per vial with fresh banana-jaggery food

medium (12 vials per population) to facilitate courtship observations. Observation vials were again set up on days

14 and 16 as described for day 12. For observing flies in the last two days in cages prior to egg lay, we set up 100

pairs of flies per Plexiglas cage ($22 \times 18 \times 18$ cm^3) with a plate of fresh food medium covered with a paste of live

yeast, as in their regular maintenance. We set up three replicate observation cages per population; these were

slightly smaller than the maintenance cages, to facilitate behavioural observations. Flies were first collected into

a regular cage (from holding vials in case of JBs and culture vials in case of FEJs) and then 100 males and 100

females were lightly anesthetized (using carbon dioxide) and transferred into each replicate observation cage. In

case of FEJs, as in regular stock maintenance, only the first ~25% of the eclosing flies were selected to become

part of the observation set.

We took three 'instantaneous' observations every four hours (corresponding to one time point) starting from day

one of observation (i.e., day 12 from JB culture initiation (egg-lay), ~day 6.5 from FEJ culture initiation) noting

the number of males observed to be performing a courtship behaviour (Spieth 1974) towards a female, and the

number of pairs in copula. Observations were conducted over 144 hours (six days) in the vials for only JBs, as the

FEJs do not have a corresponding vial stage as adults in their regular maintenance, and for 72 hours (three days)

in Plexiglas cages for both JBs and FEJs, for each replicate population. We calculated the daily average of

courtship and mating frequency (fraction of males that were courting or mating) for each day. Daily average

frequencies were then summed over all days of the effective lifetime, i.e., nine days for JB populations and three

days for FEJ populations, each for mating or courtship.

<u>Copulation Duration and Female Maturation Time</u>

To obtain adults of the same age at the beginning of the assay, FEJ cultures were started about 60 hours

after the JB cultures, to account for the difference in their egg to adult development times. We allowed an egg

laying time of only 30 minutes, in order to have a narrow eclosion distribution and collected females that emerged

within a one-hour time window. Females were paired with one day old virgin males (single pair per food vial),

with 20-25 pairs per population, under each of two conditions (presence or absence of live-yeast supplement on

food) and kept undisturbed for continuous observation. In their regular maintenance, the first matings occur in the

presence of food supplemented with additional yeast for FEJs, but without it for JBs. We recorded time till first

mating (maturation time) and copulation duration under both yeasted and un-yeasted conditions. The duration of

observation was for 56 hours, or until 80% of pairs per population, per treatment, had mated, whichever occurred

first.

Statistical Analyses

We used mixed model analyses of variance (ANOVA) for data from all the assays. Average time till

maturation for a population, lifetime courtship and mating frequency per population, and proportion of wild-type

offspring averaged across females per population were used as the response variables. Tukey's HSD test was used

for comparing individual means and calculating all confidence intervals for post-hoc analyses. Since some data

were fractional, we also repeated each analysis for those response variables with arcsine square root transformed

values to check if there were any differences obtained in the significance of the fixed factors. Selection regime

and yeasted/non-yeasted conditions for time till maturation and copulation duration, and selection regime for

courtship and mating rate assays were treated as fixed factors in the analysis. All experiments were performed on

four replicate populations of a maintenance regime. Replicate populations with matching numerical subscripts

(FEJ$_i$ and JB$_i$; i = 1-4) were assayed together, and also share the same ancestry. Those pairs were therefore treated

as random blocks in the analyses. All analyses were carried out using STATISTICA™ using Windows Release

5.0B (StatSoft Inc. 1995).

Results

Mating and Courtship Frequencies

The differences between FEJ and JB lifetime mating and courtship frequencies were statistically

significant, with JBs showing many fold higher frequencies of both courtship and mating (Fig. 1a, b; Table 1).

When we compared the mating and courtship frequencies only within the cage stage (the last three days of their

effective lifespan), courtship frequencies were still significantly greater for JB than for FEJ, but their mating

frequencies did not differ significantly (Fig. 1c, d; Table 1). The sheer magnitude of difference in the mating and

courtship frequencies experienced by the FEJ and JB males over their effective lifetime revealed the potential for

large differences in levels of sexual selection operating in these populations, with the FEJs being more or less

monogamous, in stark contrast to the JBs. Moreover, although overall courtship frequencies were very low for

FEJ, the results from the analysis of data from only within the cages indicated a very low courtship requirement

for arousal of FEJ females for copulation, given that mating frequencies in the cage were very similar for JB and

FEJ, despite courtship frequencies being much lower in FEJ (Fig. 1c, d). It is also worth noting that both courtship

and mating frequencies dropped by almost an order of magnitude for JB when only the cage stage was considered,

indicating that most mating in the regular cultures of JBs probably occurs during the vial transfer stages, before

the flies from all vials representing one replicate population are collected into a cage.

<u>Copulation Duration and Female Maturation Time</u>

Maturation time of the JB and FEJ differed significantly (Table 2), as did copulation duration (Table 3),

with JB females maturing faster, consistent with previous reports (Prasad 2004; Ghosh-Modak 2009) and mating

for longer than FEJ females (Fig. 2a, b). The drastic reduction in FEJ development time, especially pupal duration,

may have resulted in the postponement of many aspects of reproductive development to adulthood (Prasad 2004),

causing FEJ females to take ~10 hours longer than JB females to mature (Fig. 2a). Although there is considerable

evidence for change in mating behaviour with adult diet and nutrition in flies (Fricke et al. 2008; Schultzhaus et

al. 2018; Duxbury and Chapman 2020), our results did not demonstrate any effect of additional yeast on female

maturation time or copulation duration; there was no significant main effect of environment or of the selection-

by-environment interaction (Table 2, 3). Female condition (adults being fed a protein rich diet in this case) is also

known to increase male courtship vigour through an increased attractiveness of the female (Long et al. 2009), but

such effects on the FEJ or JB males, or differences in these effects, did not manifest in our assays.

Discussion

Sexual conflict is thought to be an incidental by-product of sexual selection (Trivers 1972, Parker 1979)

and several investigations have shown that the level of sexual conflict in a population can be modulated by the

intensity and scope of sexual selection (see Introduction). Most of these experimental evolution studies have found

rapid evolution of reproductive traits, including those directly or indirectly related to sexual conflict. Alterations

in life history are also predicted to have considerable consequences on the ecology of sexual selection, and as a

result, on the evolution of sexually antagonistic traits (Bonduriansky et al. 2008; Adler and Bonduriansky 2014),

and at least a subset of these effects is expected to be through alterations in the breeding ecology of a population.

Breeding ecology, which includes not only the timing of reproductive activity, but also various other aspects of

reproduction and mating behaviour such as, male-female encounter rate, promiscuity, sexual dimorphism etc., is

expected to be defined by the life history strategy adopted by the population. For example, if males in a population

have a "live-fast, die-young" life history strategy, male-male competition is expected to be intense, all else being

equal (Bonduriansky et al. 2008, and references therein). A few investigations have shown the evolution of

sexually selected traits as a consequence of selection for life history traits such as lifespan (Service 1993; Service

and Fales 1993; Service and Vossbrink 1996; Makklakov et al. 2010). These reports generally support the idea

that evolution of life history traits can result in changes in male reproductive investment, which may further result

in reduction in sexual conflict in the population. Yun et al. (2017) showed that the sexual conflict in a population

living in a physically complex habitat can be less than that in a simple habitat. If this effect of habitat type is due

to females' access to a spatial refuge, then male-female encounter rate is directly implicated as an important

determinant of the level of conflict in that population. Therefore, factors such as, locomotor activity, opportunity

of interaction etc. are expected to be important modulators of the conflict.

In the study reported here, our observations on the breeding ecology of *D. melanogaster* populations selected for

rapid development and early reproduction suggest there is reduced scope for sexual selection in them compared

to their ancestral controls. A reduction in the time available for mating in the FEJ populations appears to

considerably reduce reproductive competition by constraining re-mating opportunities. Thus, the lower levels of

sexual conflict in these populations, compared to their ancestral controls (Ghosh and Joshi 2012), earlier shown

to be partly due to their smaller body size (Mital et al. 2021), are also likely to be in part a consequence of reduced

sexual selection and male-male competition for mates. Supporting this theory further, we also find males from

our selected populations to have substantially lower courtship frequency and shorter copulation duration,

indicating reduced investment in pre- and post-copulatory competition.

To better interpret the large difference in life-time mating rates between our selected populations and their

controls, we scaled the mating frequency results by the maximum possible number of matings achievable in that

time for that population. We multiplied the FEJ mating rate by 24 (maximal possible mating events being once

every hour) to obtain an expected number of life-time mating events, which got rounded off to one mating during

their effective adult life. In comparison, the projected life-time mating events for a JB fly is six matings during

their effective adult life (scaling factor of 12, mating once every two hours). These scaling factors were chosen

based on the mating duration and likely recovery required by FEJ and JB males before attempting re-mating (A

Mital, personal observation) and we believe these are overestimates of the maximum number of matings possible

within a day. Therefore, one and six are conservative estimates of the expected lifetime mating events per fly for

FEJs and JBs, respectively. We conclude that males from the faster developing and early reproducing population

likely experience very low levels of competition for mates as they are effectively monogamous.

Another important result indicating considerable change in reproductive behaviour of our selected population is

the large difference in their lifetime courtship frequency from that of the ancestral controls (Fig. 1b), which

persisted even in the cage stage (last three days of effective adult life, Fig. 1d) while similar differences in the

mating frequencies were not seen in the cage stage (Fig. 1c). We speculate multiple plausible causes for this result

and discuss them as follows.

The period between eclosion and reproduction is typically the only opportunity for females to feed and thereby

accumulate additional resources for egg production (Luckinbill et al 1985; Chippindale et al. 1997b). Such

compensatory feeding would be especially important for FEJ females given their small size and low lipid content

at eclosion (Prasad 2004). The already resource limited FEJ females are expected to then reduce all energetically

expensive activities, which may include mating and resistance to mate harm, making increased receptivity

selectively advantageous. That FEJ males are significantly less harmful and females less resistant has already been

reported earlier (Ghosh and Joshi 2012, Mital et al. 2021). Moreover, frugal courtship by males and a low arousal

threshold in females is likely to be selectively advantageous also because mating at least once is necessary to have

any fitness, and time available for reproduction in the FEJ breeding ecology is short (further reduced by a long

maturation time, Fig. 2a). In addition, since FEJs have also evolved a significantly different developmental

schedule and maturation time (Prasad 2004; Ghosh-Modak 2009; Fig. 2a), developmental changes may further

contribute to poor courtship by males, especially since males complete reproductive activity at a very young age

of three days. For instance, in some studies on *D. melanogaster*, males have been found to keep maturing

reproductively, especially in terms of accessory gland growth, for up to six days after eclosion, with younger

males tending to have poor sperm competitive ability and courtship effort as compared to more mature males

(Ruhmann et al. 2016). Furthermore, since courtship significantly contributes to the cost of mating in females

(Partridge and Fowler 1990), reduced courtship intensity may be selectively favoured in the FEJ regime where

male and female fitness are expected to be highly correlated. Finally, males from the ancestral controls (i.e., JBs),

may be selected for high courtship effort during the final three days of their adult life, as they have about 80%

paternity share of the offspring that form the next generation if they mate with females during the 'cage' stage

(Supplementary Material, Fig. S1) (Fig. 2c, d). JB males would gain a high fitness reward for mating effort in this

time despite low chance of mating success. This may also indicate high variation in male mating success in the

control population, a long-standing argument for stronger selection in males for exaggerated reproductive traits

(Bateman 1948; Andersson 1994).

More generally, however, low courtship frequencies despite similar mating frequencies among males that we see

in the cage stage (Fig. 1c, d) can also be explained in the context of the chase-away selection model for the

evolution of exaggerated sexual traits in *D. melanogaster* (Holland and Rice 1998; Gavrilets et al. 2001). The

model suggests that if persistent male mating effort is harmful to females (sexual conflict) then high female

resistance towards such sensory manipulation will evolve. This may be interpreted as increased female

bias/preference, in turn selecting for higher courtship effort from males (Holland and Rice 1998, 1999). Since

persistent male courtship is known to reduce female survival in *D. melanogaster* (Partridge and Fowler 1990),

reduced sexual conflict driven by low male-male competition in FEJs may release the sexes from this co-

evolutionary cycle, and appear as lack of female choice and male courtship effort. Therefore, low courtship

frequency in FEJs (for similar mating frequency as in JBs) in the last three days of their adult life may be driven

by female ecological constraints, changes in development, reduced male-male competition, or a combination of

these. Put together, our results on the lifetime mating and courtship frequencies in these populations demonstrate

the evolution of reduced sexual selection and conflict as a consequence of strong selection for rapid development

and an extremely short adult life.

The interplay between life-history and sexual selection is expected to ultimately determine the optimum

investment strategy for a population. In our study populations, the changes that have likely resulted in reduced

sexual selection are also expected to release the flies from certain reproductive investment demands. We observed

that FEJ flies mate for around 15 minutes on average, compared to about 25 minutes for JB flies (Fig. 2b).

Production of Acps is known to be costly to males (Chapman and Edwards 2011) and it has been suggested that

in *D. melanogaster*, most sperm transfer occurs during the first half of mating (Gilchrist and Partridge 2000), the

remaining time being spent in the transfer of Acps responsible for the female post-mating response. We speculate

that, reduced energy investments in courtship and in the non-sperm components of the ejaculate in the FEJs would

not only explain their shorter copulation duration (Fig. 2b) but potentially also permit the evolution of even smaller

flies, further pushing the boundaries of rapid development for which they experience strong selection. Changes in

resource use in FEJs have been demonstrated previously; FEJs have evolved larger egg size (B M Prakash and A

Joshi, unpublished data) and more eggs produced per unit dry weight by females (Prasad 2004; Ghosh-Modak

2009) than JBs. It is therefore probable that the optimum investment strategy for males has also diverged from

that of the ancestral controls. This is further supported by the observation that, greater starvation resistance per

unit lipid has evolved in the FEJs than JBs (Prasad 2004) and that Acp genes in young FEJ males are

downregulated (Satish 2010; K M Satish, P Dey and A Joshi, unpublished data).

Conclusion

Our results demonstrate that the breeding ecology of our fast developing and early reproducing

populations has likely resulted in reduced levels of sexual selection, and consequently, facilitated the evolution of

reduced sexual conflict experienced by them, as compared to their ancestral controls. The shorter effective adult

life of FEJs, together with strong directional selection for rapid development, has resulted in small flies exhibiting

significantly lower competition among males for mating. These observations complement other studies that have

reported reduced sexual conflict upon directly selecting for different levels of male-male competition

(Introduction) and exemplify the potential impact of selection on life-history related traits (development time and

age at reproduction) on levels of sexual selection and conflict via major changes in reproductive behaviour. This

is especially relevant as differences in sexual selection and sexual conflict are often implicated in the emergence

of speciation phenotypes (Parker and Partridge 1998). We note that the FEJs and JBs have diverged in their

reproductive behaviour to a degree that incipient reproductive isolation has occurred (Ghosh and Joshi 2012). Our

work, therefore, also highlights that selection on traits not directly associated with sexual conflict can, nonetheless,

drive the evolution of reproductively isolating mechanisms, either directly through rapid development (Mital et

al. 2021) or indirectly by affecting overall levels of sexual selection, or both. Dissecting out such nuanced

interactions of life-history and reproductive strategies in affecting sexual selection and sexual conflict is important

for understanding the myriad evolutionary consequences that may accompany adaptation to specific ecological

challenges.

  Tables

  Table 1 Results summary of ANOVA for effect of selection on mating and courtship frequencies[1]

Response Variable	df	MS	*F*	*P*
Lifetime Mating Fr	1	0.5502	1020.928	< 0.0001
Lifetime Courtship Fr	1	120.2025	1372.439	< 0.0001
Mating Fr in cage	1	0.0001	4.163	0.1339
Courtship Fr in cage	1	0.0010	910.352	<0.001

Table 2 Results summary of ANOVA for maturation time[2]

Effect	df	MS	*F*	*P*
Selection	1	535.6813	149.885	0.0011
Environment	1	1.6626	0.203	0.6823
Selection × Environment	1	1.0137	0.003	0.9542

Table 3 Results summary of ANOVA for copulation duration[3]

Effect	df	MS	*F*	*P*
Selection	1	143.8742	15.260	0.0297
Environment	1	0.6389	0.6460	0.4802
Selection × Environment	1	0.0867	0.0326	0.8680

[1] Summary of results of a one-way ANOVA each for lifetime and cage mating and courtship frequencies. Main effects of selection regime for the two traits are shown. In this design, random factors and interaction effects cannot be tested for significance and have been left out for brevity.

[2] Selection regime and environment (yeasted or non-yeasted) are fixed factors in the ANOVA. Random factors and interactions cannot be tested for significance in this design, and have been left out for brevity.

[3] Selection regime and environment (yeasted or non-yeasted) are fixed factors in the ANOVA. Random factors and interactions cannot be tested for significance in this design, and have been left out for brevity.

Figures

Fig. 1 a) Mean lifetime mating frequency averaged across the four replicate populations of JB and FEJ. b) Mean lifetime

courtship frequency averaged across four replicate populations of JB and FEJ. c) Mean mating frequency in the cage, during

the final three days of effective adult life, averaged across four replicate populations of JB and FEJ. d) Mean courtship

frequency in the cage, averaged across the four replicate populations of JB and FEJ. Error bars are 95% confidence intervals

around the means.

Fig. 2 a) Mean maturation time, or mating latency of virgins from eclosion, averaged across four replicate populations of JB

and FEJ, under yeasted (Y) and non-yeasted (NY) conditions. b) Mean copulation duration of the first mating averaged across

four replicate populations of JB and FEJ, under yeasted (Y) and non-yeasted (NY) conditions. Error bars are 95% confidence

intervals around the means.

Fig. 3 Mean proportion of wild-type flies among the progeny of the SE flies resulting from eggs laid on day 21, from each of

the four phases in the breeding success assay, averaged across the four replicate JB populations. Error bars are 95% confidence

intervals around the means. The results show that the JB males have the greatest fertilization success when they mate during

Phase 4, immediately before egg-collection on day 21.

References

Adler MI, Bonduriansky R (2014) Sexual conflict, life span, and aging. Cold Spring Harb Perspect Biol. 6:a017566. https://doi: 10.1101/cshperspect.a017566

Andersson M (1994) Sexual Selection. Princeton University Press, Princeton, USA

Anholt RHR, O'Grady P, Wolfner MF, Harbison ST (2020) Evolution of reproductive behaviour. Genetics 214:49–73 https://doi.org/10.1534/genetics.119.302263

Archer CR, Zajitschek F, Sakaluk SK, Royle NJ, Hunt J (2012) Sexual selection affects the evolution of lifespan and ageing in the decorated cricket *Gryllodes sigillatus*. Evolution 66:3088–100 https://doi.org/10.1111/j.1558-5646.2012.01673.x

Arnqvist G, Rowe L (2013) Sexual Conflict. Princeton University Press, Princeton, USA

Bateman AJ (1948) Intra-sexual selection in *Drosophila*. Heredity 2:349–368 https://doi.org/10.1038/hdy.1948.21

Bonduriansky R, Makklakov A, Zajitschek F, Brooks R (2008) Sexual selection, sexual conflict and the evolution of ageing and lifespan. Func Ecol 22:443–453 https://doi.org/10.1111/j.1365-2435.2008.01417.x

Chapman T, Liddle LF, Kalb JM, Wolfner MF, Partridge L (1995) Cost of mating in *Drosophila melanogaster* females is mediated by male accessory gland products. Nature 373:241–244 https://doi.org/10.1038/373241a0

Chapman T (2001) Seminal fluid-mediated fitness traits in *Drosophila*. Heredity 87:511 https://doi.org/10.1046/j.1365-2540.2001.00961.x

Chapman T, Arnqvist G, Bangham J, Rowe L (2003) Sexual conflict. Trends Ecol Evol 18:41–47

Chapman T and Edward D (2011) Mechanisms underlying reproductive trade-offs: cost of reproduction. Pp. 137– 152 *in* T. Flatt and A. Heyland, eds. Life-history evolution: understanding the proximate mechanism. Oxford Univ. Press, Oxford, U.K. https://doi.org/10.1093/acprof:oso/9780199568765.003.0011

Chippindale AK, Alipaz AJ, Chen HW, Rose MR (1997)*a* Experimental evolution of accelerated development in *Drosophila*. 1. Developmental speed and larval survival. Evolution 51:1536–1551 https://doi.org/10.1111/j.1558-5646.1997.tb01477.x

Chippindale AK, Leroi AM, Saing H, Borash DJ, Rose MR (1997)*b*. Phenotypic plasticity and selection in *Drosophila* life history evolution. 2. Diet, mates and the cost of reproduction. J Evol Biol 10:269–293

Cordts R, Partridge L (1996) Courtship reduces longevity of male *Drosophila melanogaster*. Anim Behav 52:269–278

Crudgington HS, Beckerman AP, Brüstle L, Green K, Snook RR (2005) Experimental removal and elevation of

sexual selection: does sexual selection generate manipulative males and resistant females? Am Nat 165:72–87

https://doi.org/10.1086/429353

Darwin C (1859) On the origin of species by means of natural selection, or the preservation of favoured races in

the struggle for life. Murray, London, UK

Darwin C (1868) The variation of animals and plants under domestication (2 Vols). Murray, London, UK

Darwin C (1871) The descent of man and selection in relation to sex (2 Vols). Murray, London, UK (Facsimilie:

Princeton University Press, Princeton, NJ, USA, 1981)

Duxbury EML, Chapman T (2020) Sex-specific responses of life span and fitness variation in developmental vs

adult diets in *Drosophila melanogaster.* J Gerontol Biol Sci 75:1431–1438 https://doi.org/10.1093/gerona/glz175

Edward DA, Fricke C, Chapman T (2010) Adaptations to sexual selection and sexual conflict: insights from

experimental evolution and artificial selection. Philos Trans R Soc Lond., B, Biol Sci 365:2541–2548

https://dx.doi.org/10.1098%2Frstb.2010.0027

Fricke C, Bretman A, Chapman T (2008) Adult male nutrition and reproductive success in *Drosophila*

melanogaster. Evolution 62:3170–3177 https://doi.org/10.1111/j.1558-5646.2008.00515.x

Garlovsky MD, Holman L, Brooks AL, Snook RR (2021) Sexual selection affects the evolution of physiological

and life-history traits. BioRxiv https://doi.org/10.1101/2021.03.23.436586

Gavrilets S, Arnqvist G, Friberg U (2001) The evolution of female mate choice by sexual conflict. Proc Royal

Soc B: Biol Sci 268:531–539 https://dx.doi.org/10.1098%2Frspb.2000.1382

Ghosh-Modak, SG (2009) Evolution of life-history traits, canalization and reproductive isolation in laboratory

populations of *Drosophila melanogaster* selected for faster pre-adult development and early reproduction

(Doctoral dissertation), Jawaharlal Nehru Centre for Advanced Scientific Research, Bengaluru, India

Ghosh SM, Joshi A (2012) Evolution of reproductive isolation as a by-product of divergent life-history evolution

in laboratory populations of *Drosophila melanogaster*. Ecol Evol 2:3214–3226

https://dx.doi.org/10.1002%2Fece3.413

Gilchrist AS, Partridge L (2000) Why it is difficult to model sperm displacement in *Drosophila melanogaster*: the

relation between sperm transfer and copulation duration. Evolution 54:534–542 https://doi.org/10.1111/j.0014-

3820.2000.tb00056.x

Holland B, Rice WR (1998) Perspective: chase-away sexual selection: antagonistic seduction versus resistance.

Evolution 52:1–7 https://doi.org/10.1111/j.1558-5646.1998.tb05132.x

Holland B, Rice WR (1999) Experimental removal of sexual selection reverses intersexual antagonistic co-

evolution and removes a reproductive load. PNAS 96:5083–5088 https://doi.org/10.1073/pnas.96.9.5083

Hollis B, Houle D, Yan Z, Kawecki TJ, Keller L (2014) Evolution under monogamy feminizes gene expression

in *Drosophila melanogaster*. Nat Commun 5:3482 https://doi.org/10.1038/ncomms4482

Hollis B, Keller L, Kawecki TJ (2017) Sexual selection shapes development and maturation rates in *Drosophila*.

Evolution 71:304–314 https://doi.org/10.1111/evo.13115

Kuijper B and Morrow EH (2009) Direct observation of female mating frequency using time lapse photography.

FLY 3:118–120 https://doi.org/10.4161/fly.8053

Linklater JR, Wertheim B, Wigby S, Chapman T (2007) Ejaculate depletion patterns evolve in response to

experimental manipulation of sex ratio in *Drosophila melanogaster*. Evolution 61:2027–2034

https://doi.org/10.1111/j.1558-5646.2007.00157.x

Long TAF, Pischedda A, Stewart AD, Rice WR (2009) A cost of sexual attractiveness to high-fitness females.

PLoS Biol 7:e1000254 https://doi.org/10.1371/journal.pbio.1000254

Luckinbill LS, Clare MJ (1985) Selection for life span in *Drosophila melanogaster*. Heredity 55:9–18

https://doi.org/10.1038/hdy.1985.66

Makklakov AA, Cayeteno L, Brooks RC, Bonduriansky R (2010) The roles of life-history selection and sexual

selection in the adaptive evolution of mating behaviour in a seed beetle. Evolution 64:1273–1282

https://doi.org/10.1111/j.1558-5646.2009.00904.x

Mital A, Sarangi M, Joshi A (2021) Evolution of lower levels of inter-locus sexual conflict in *D. melanogaster*

populations under strong selection for rapid development. BioRxiv https://doi.org/10.1101/2021.02.08.430125

Nandy B, Chakraborty P, Gupta V, Ali SZ, Prasad NG (2013)*a* Sperm competitive ability evolves in response to

experimental alteration of operational sex ratio. Evolution 67:2133–2141 https://doi.org/10.1111/evo.12076

Nandy B, Gupta V, Sen S, Udaykumar N, Samant MA, Ali SZ, Prasad NG (2013)*b* Evolution of mate-harm,

longevity and behaviour in male fruit flies subjected to different levels of inter-locus conflict. BMC Evol Biol

13:212 https://doi.org/10.1186/1471-2148-13-212

Nandy B, Gupta V, Udaykumar N, Samant MA, Sen S, Prasad NG (2014) Experimental evolution of female traits

under different levels of intersexual conflict in *Drosophila melanogaster*. Evolution 68:412–425

https://doi.org/10.1111/evo.12271

Nunney L (1996) The response to selection for fast larval development in *Drosophila melanogaster* and its effect

on adult weight: an example of a fitness trade-off. Evolution 50:1193–1204 https://doi.org/10.1111/j.1558-

5646.1996.tb02360.x

Parker GA (1979) Sexual selection and sexual conflict. Pages 123–166 in M. S. Blum and N. A. Blum, eds. Sexual

Selection and Reproductive Competition in Insects, Academic Press, Athens, GA, USA

Partridge L, Fowler K, Trevitt S, Sharp W (1986) An examination of the effects of males on the survival and egg-

production rates of female *Drosophila melanogaster*. J Insect Physiol 32:925–929 https://doi.org/10.1016/0022-

1910(86)90140-X

Partridge L, Hoffmann A, Jones JS (1987)*a* Male size and mating success in *D. melanogaster* and *D.*

pseudoobscura under field conditions. Anim Behav 35:468–476 https://doi.org/10.1016/S0003-3472(87)80272-5

Partridge L, Ewing A, Chandler A (1987)*b* Male size and mating success in *Drosophila melanogaster*: the roles

of male and female behaviour. Anim Behav 35:555–562 https://doi.org/10.1016/S0003-3472(87)80281-6

Partridge L, Fowler K (1990) Non-mating costs of exposure to males in female *Drosophila melanogaster*. J Insect

Physiol 36:419–425 https://doi.org/10.1016/0022-1910(90)90059-O

Pitnick S (1991) Male size influences mate fecundity and remating interval in *Drosophila melanogaster*. Anim

Behav 41:735–745 https://psycnet.apa.org/doi/10.1016/S0003-3472(05)80340-9

Pitnick S, Miller GT, Reagan J, Holland B (2001) Males' evolutionary responses to experimental removal of

sexual selection. Proc Royal Soc B: Biol Sci 268:1071–1080 https://doi.org/10.1098/rspb.2001.1621

Pitnick S, García–González F (2002) Harm to females increases with male body size in *Drosophila melanogaster*.

Proc Royal Soc B: Biol Sci 269:1821–1828 https://doi.org/10.1098/rspb.2002.2090

Prasad NG, Shakarad M, Gohil VM, Sheeba V, Rajamani M, Joshi A (2000) Evolution of reduced pre-adult

viability and larval growth rate in laboratory populations of *Drosophila melanogaster* selected for shorter

development time. Genet Res 76:249–259 https://doi.org/10.1017/s0016672300004754

Prasad NG (2004) *Life-history evolution in laboratory populations of Drosophila melanogaster subjected to*

selection for faster development and early reproduction (Doctoral dissertation). Jawaharlal Nehru Centre for

Advanced Scientific Research, Bengaluru, India

Parker GA, Partridge L (1998) Sexual conflict and speciation. Philos Trans R Soc Lond B, Biol Sci 353:261–74

https://doi: 10.1098/rstb.1998.0208

Ruhmann H, Wensing KU, Neuhalfen N, Specker JH, Fricke C (2016) Early reproductive success in *Drosophila*

males is dependent on maturity of the accessory gland. Behav Ecol 27:1859–1868

https://doi.org/10.1093/beheco/arw123

Satish KM (2010) Reverse evolution and gene expression studies on populations of *Drosophila melanogaster*

selected for rapid pre-adult development and early reproduction (Doctoral dissertation). Jawaharlal Nehru Centre

for Advanced Scientific Research, Bengaluru, India

Schultzhaus JN, Bennet CJ, Iftikar H, Yew JY, Mallet J, Carney GE (2018) High-fat diet alters Drosophila

melanogaster sexual behaviour and traits: decreased attractiveness and changes in pheromone profiles. Sci Rep

8:5387 https://doi.org/10.1038/s41598-018-23662-2

Service PM (1993) Laboratory evolution of longevity and reproductive fitness components in male fruit flies:

mating ability. Evolution 47:387–399 https://doi.org/10.2307/2410059

Service PM, Fales AJ (1993) Evolution of delayed reproductive senescence in male fruit flies: sperm competition.

Genetica 91:111–125 https://doi.org/10.1007/BF01435992

Service PM, Vossbrink RE (1996) Genetic variation in "first" male effects on egg laying and re-mating by female

Drosophila melanogaster. Behav Genet 26:39–48 https://doi.org/10.1007/BF02361157

Sheeba V, Madhyastha NA, Joshi A (1998) Oviposition preference for novel versus normal food resources in

laboratory populations of *Drosophila melanogaster*. J Biosci 23:93–100 https://doi.org/10.1007/BF02703000

Spieth HT (1974) Courtship behaviour in *Drosophila*. Annu Rev Entomol 19:385–405

https://doi.org/10.1146/annurev.en.19.010174.002125

StatSoft Inc. (1995) STATISTICA^{TM} for Windows. Tulsa, OK, USA

Trivers R (1972) Parental Investment and Sexual Selection. Pages 136–179 *in* B. Cambell, ed. Sexual Selection

& the Descent of Man, Aldine de Gruyter, New York, USA

Wensing KU, Koppik M, Fricke C (2017) Precopulatory but not postcopulatory male reproductive traits diverge

in response to mating system manipulation in *Drosophila melanogaster*. Ecol and Evol 7:10361–10378

https://doi.org/10.1002/ece3.3542

Wigby S, Chapman T (2004) Female resistance to male harm evolves in response to manipulation of sexual

conflict. Evolution 58:1028–1037 https://doi.org/10.1111/j.0014-3820.2004.tb00436.x

Wolfner MF (1997) Tokens of love: functions and regulation of *Drosophila* male accessory gland products. Insect

Biochem Mol Biol 27:179–192 https://doi.org/10.1016/s0965-1748(96)00084-7

Yun L, Chen PJ, Singh A, Agrawal AF, Rundle HD (2017) The physical environment mediates male harm and

its effects on selection in females. Proc Royal Soc B: Biol Sci 284:20170424

https://doi.org/10.1098/rspb.2017.0424

Zajitschek F, Bonduriansky R, Zajitschek SR, Brooks RC (2009) Sexual dimorphism in life history: age,

survival, and reproduction in male and female field crickets *Teleogryllus commodus* under seminatural

conditions. Am Nat 173:792–802 https://doi.org/10.1086/598486

Zwaan B, Bijlsma R, Hoekstra RF (1995) Artificial selection for developmental time in *Drosophila melanogaster*

in relation to the evolution of aging: direct and correlated responses. Evolution 49:635–648

https://doi.org/10.1111/j.1558-5646.1995.tb02300.x