Intranasal gene therapy to prevent infection by SARS-CoV-2 variants

Short title: Gene Therapy to Prevent COVID-19

Joshua J. Sims, Jenny A. Greig, Kristofer T. Michalson, Sharon Lian, R. Alexander Martino,

Rosemary Meggersee, Kevin B. Turner, Kalyani Nambiar, Cecilia Dyer, Christian Hinderer,

Makoto Horiuchi, Hanying Yan, Xin Huang, Shu-Jen Chen, James M. Wilson*

Gene Therapy Program, Department of Medicine, Perelman School of Medicine, University of

Pennsylvania, Philadelphia, PA 19104, USA

*Corresponding author:

James M. Wilson, MD, PhD

Gene Therapy Program

Perelman School of Medicine

University of Pennsylvania

125 South 31st Street, Suite 1200

Philadelphia, PA 19104, USA

Phone: 215-573-9020; Fax: 215-494-5444

E-mail: wilsonjm@upenn.edu

Abstract

SARS-CoV-2 variants have emerged with enhanced pathogenicity and transmissibility, and

escape from pre-existing immunity, suggesting first-generation vaccines and monoclonal

antibodies may now be less effective. This manuscript demonstrates an approach for preventing

clinical sequelae and the spread of SARS-CoV-2 variants. First, we affinity-matured an

angiotensin-converting enzyme 2 (ACE2) decoy protein, achieving 1000-fold binding

improvements that extend across a wide range of SARS-CoV-2 variants and distantly related,

ACE2-dependent coronaviruses. Next, we demonstrated the expression of this decoy in proximal

airway when delivered via intranasal administration of an AAV vector. This intervention

significantly diminished clinical and pathologic consequences of SARS-CoV-2 challenge in a

mouse model and achieved therapeutic levels of decoy expression at the surface of proximal

airways when delivered intranasally to nonhuman primates. Importantly, this long-lasting,

passive protection approach is applicable in vulnerable populations such as the elderly and

immune-compromised that do not respond well to traditional vaccination. This approach could

be useful in combating COVID-19 surges caused by SARS-CoV-2 variants and should be

considered as a countermeasure to future pandemics caused by pre-emergent members, ACE2-

dependent CoVs that are poised for zoonosis.

Author summary

SARS-CoV-2 variants have emerged with enhanced pathogenicity and transmissibility, and escape from pre-existing immunity, suggesting first-generation vaccines and monoclonal antibodies may now be less effective. This manuscript demonstrates an approach for preventing clinical sequelae and the spread of SARS-CoV-2 variants. First, we affinity-matured an angiotensin-converting enzyme 2 (ACE2) decoy protein, achieving 1000-fold binding improvements that extend across a wide range of SARS-CoV-2 variants and distantly related, ACE2-dependent coronaviruses. Next, we demonstrated the expression of this decoy in proximal airway when delivered via intranasal administration of an AAV vector. This intervention significantly diminished clinical and pathologic consequences of SARS-CoV-2 challenge in a mouse model and achieved therapeutic levels of decoy expression at the surface of proximal airways when delivered intranasally to nonhuman primates. Importantly, this long-lasting, passive protection approach is applicable in vulnerable populations such as the elderly and immune-compromised that do not respond well to traditional vaccination. This approach could be useful in combating COVID-19 surges caused by SARS-CoV-2 variants and should be considered as a countermeasure to future pandemics caused by pre-emergent members, ACE2-dependent CoVs that are poised for zoonosis.

Introduction

Developing a soluble form of angiotensin-converting enzyme 2 (ACE2)—referred to as a

decoy—is considered a protein therapeutic in the treatment of COVID-19 patients (1, 2). We

isolated an ACE2 decoy that broadly neutralizes SARS-CoV-2 variants and demonstrate its

potential for preventing COVID-19 when expressed from an adeno-associated virus (AAV)

following intranasal (IN) delivery. We have reported previously on the effectiveness of IN AAV

to express antibodies that broadly neutralize pandemic strains of influenza (3-6).

Results

ACE2 decoy affinity maturation enhances neutralization of SARS-CoV-2 100-fold

We initially constructed a decoy receptor by fusing a human ACE2 fragment to the human IgG4

Fc domain. We cloned this first-generation decoy into AAV and delivered it as a nasal spray into

nonhuman primates (NHPs). Although we detected decoy expression in nasal lavage fluid

(NLF), the decoy was not produced at levels sufficient to overcome the low neutralizing potency

of this protein (Figure S1). We therefore set out to affinity-mature the ACE2 protein sequence.

We generated diverse ($>10^8$ transformants) ACE2 variant libraries in a yeast-display format(7)

using error-prone polymerase chain reaction (PCR; Figure 1A and Figure S2). We screened the

primary libraries in rounds of fluorescence-activated cell sorting (FACS; Figure 1B). We

selected populations with better binding to SARS-CoV-2 receptor binding domain (RBD) and

tracked library convergence with deep sequencing (Figure 1C and 1D). Frequently observed

mutations from our primary library sorts overlap partially with mutations reported by others(2,

8), including substitutions at T27 and N90 glycan disruption (Figure S2). Validated clones from

the sorted primary libraries (Figure S3) seeded a secondary library formed by mutagenic

recombination(9, 10), which we screened using stringent off-rate sorting(11) (Figure 1C).

We expressed ACE2 variants from several stages of the yeast-display screening as soluble IgG4

Fc fusions, evaluated expression titers, and predicted IC_{50} for SARS-CoV-2 neutralization using

reporter virus (Figure S3). The most potent neutralizing variants converged upon similar

substitutions at five positions: 31, 35, 79, 330, and N90 glycan disruption (Figure 1E). After

further characterization, we selected CDY14-Fc4 as the most improved ACE2 decoy variant. To

avoid off-target effects *in vivo*, we ablated ACE2 enzyme activity by introducing H345L(12) at

no cost to potency. By surface plasmon resonance (SPR) the active site-null CDY14HL-Fc4

bound SARS-CoV-2 RBD with 1,000-fold improved affinity (29 nM for wtACE2 vs. 31 pM for

CDY14HL-Fc4; see Figure 1, panels F and G). CDY14HL-Fc4 neutralized Wuhan-Hu-1 SARS-

CoV-2 reporter nearly 100-fold better than the un-engineered ACE2 decoy (IC_{50} 127 ng/ml for

CDY14HL-Fc4 vs. 11 µg/ml for ACE2-wt-Fc4; see Figure 1H).

ACE2 decoy is effective against SARS-CoV-2 variants and SARS-CoV-1

Escape mutations at the immunodominant ACE2 binding site of the RBD is of major concern for

emerging SARS-CoV-2 variants(13). RBDs from more distant ACE2-dependent CoVs also

differ substantially from the original SARS-CoV-2 at the ACE2 interface (Figure 2A). Unlike

antibodies, decoy inhibitors may achieve broad neutralization and escape mutant resistance;

changes that reduce decoy binding would also decrease ACE2 receptor binding, thus reducing

viral fitness. To assess this potential, we measured ACE2-Fc4 (a surrogate for the native

receptor) and CDY14HL-Fc4 (the therapeutic decoy) affinities across a diverse panel of CoV

RBDs using SPR. We chose RBDs from strains under positive selection in the course of the 2020

pandemic (e.g., 439K (14), B.1.1.7 and B1.351, first isolated in the EU, UK, and the Republic of

South Africa, respectively(15)), and mink-adapted isolates(16). Several emerging SARS-CoV-2

variants with improved affinity for ACE2-Fc4 (B1.1.7, 453F, and 501T) also bind CDY14HL-

Fc4 more tightly (Figure 2B and Figure S4), while B.1.351, 439K, and 439K/417V only

modestly alter binding to ACE2-Fc4 or CDY14HL-Fc4. While 486L reduces affinity for

CDY14HL-Fc4, it does so for ACE2-Fc4 proportionally. Remarkably, decoy and receptor RBD

affinities are tightly coupled even for the distantly related SARS-pandemic CoV-1 and the pre-

emergent bat WIV1-CoV(17).

Next, we compared decoy neutralization across diverse SARS-CoVs using pseudotyped

lentivirus reporters. The SARS-CoV-2 variant reporter viruses in the 614G (18, 19) background

(439K, B1.1.7, B.1.351) are neutralized near or below the IC_{50} of the 614G reporter (IC_{50} values:

77 ng/ml for 614G, 42 ng/ml for 439K, 93 ng/ml for B.1.1.7, and 45 for B.1.351; Figure 2C).

Remarkably, selecting for increased binding to the SARS-CoV-2 RBD resulted in very potent

neutralization of the phylogenetically distinct SARS-CoV-1 reporter virus ($IC_{50} = 53$ ng/ml).

Taken together with the binding survey, these data indicate that structural features of the ACE2

interface have been retained through the stages of directed evolution. Moreover, the data predict

that CDY14HL-Fc4 could protect against current, emerging, and future pandemic ACE2-

dependent CoVs.

ACE2 decoy diminishes SARS-CoV-2 sequelae in transgenic mice

We considered SARS-CoV-2 challenge studies in hamsters, macaques, and the hACE2

transgenic (TG) mice to evaluate the *in vivo* efficacy of an AAV vector expressing CDY14HL-

Fc4. In all models, achieving evidence of viral replication *in vivo* requires virus doses that far

exceed those required for human transmission. Furthermore, the clinical and pathologic sequalae

of SARS-CoV-2 exposure is attenuated in these species compared to severely affected humans.

The most significant limitation, however, is that all the challenge models require direct pathogen

delivery to the lung in order to demonstrate pathology, which does not simulate the mechanism

of the AAV decoy product, which focuses on localized expression in the proximal airway

following intranasal delivery to reduce SARS-CoV-2 infection and its consequences. We

therefore selected the hACE2 TG mouse model for three reasons: 1) we can characterize disease

by measuring viral loads, clinical sequalae, and histopathology; 2) we can use an IN route of

administration as we would in humans, realizing this deposits vector in the proximal and distal

airways of the mouse, while IN delivery in humans is restricted to the proximal airway; and 3)

we can leverage the extensive experience of murine models in de-risking human studies of AAV

gene transfer.

We conducted pilot studies in wild-type mice to determine which decoy protein (CDY14-Fc4 vs.

CDY14HL-Fc4) and capsid (clade F AAVhu68 vs. clade A AAVrh91) maximized expression

following *in vivo* gene delivery. We administered 10^{11} GC of vector and recovered broncho-

alveolar lavage samples (BAL) 7 days later to evaluate ACE2 decoy protein expression and

activity (Figure 3A-C). Based on mass spectrometry (MS) protein measurements, the AAVhu68

capsid was more efficient than the AAVrh91 capsid in transducing mouse lung. The HL

mutation modestly reduced expression (p<0.007). Importantly, we found a direct correlation

between decoy expression levels and the ability to bind to SARS-CoV-2 spike protein and

neutralize a SARS-CoV-2 pseudotype, demonstrating function of the decoy expressed from

airway tissues (Figure 3A-C). We selected CDY14HL-Fc4 as the clinical candidate transgene

and the AAVhu68 capsid for the mouse challenge studies (Figure 3D).

We IN delivered AAVhu68-CDY14HL-Fc4 or vehicle to hACE2-TG mice. Seven days later,

animals were challenged with SARS-CoV-2 (280 pfu), followed clinically (observation and daily

weights), and necropsied on days 4 and 7 after challenge for tissue and BAL analysis (Figure

3D). Expression of CDY14HL-Fc4 in BAL normalized for dilution was in the range of the IC_{50}

measured *in vitro* and in the pilot studies (Figure 3F). Sham-treated SARS-CoV-2 challenged

animals demonstrated statistically significant weight loss as has been described by others(20-22).

We observed significantly less weight loss amongst vector-treated animals (which we followed

for 7 days) compared to untreated animals (observed on days 4 and 7; p<0.05, linear mixed effect

modeling). The vector-treated animals also significantly differed from the untreated,

unchallenged animals (Figure 3E). Interestingly, the clinical outcome of the treatment was better

among females than males, although we noted significant variations within the treated group

(Figure S5).

Histopathology of the lungs from vehicle treated animals challenged with SARS-CoV-2 revealed

findings similar to that previously described in this model(22). As expected, tissues from animals

not challenged with SARS-CoV-2 demonstrated no histopathology. Samples from days 4 and 7

showed reduced lung pathology in AAVhu68.CDY14HL-Fc4 treated animals vs. the vehicle-

treated animals; the day-4 samples achieved statistical significance ($p < 0.05$; Wilcoxon Rank

Sum Test). (Figure 3G). Compared to vehicle-treated animals, viral RNA in BAL and lung

homogenate was diminished at day 4 and 7 in AAVhu68.CDY14HL-Fc4 treated animals (Figure

3H and 3I). The greatest reductions were at day 7 for both BAL (26-fold) and lung tissue (35-

fold). Impact of the AAVhu68.CDY14HL-Fc4 on SARS-CoV-2 replication, as determined by

median sgRNA levels, was greatest at day 7 (27-fold reduction, Figure 3J). Although there was

substantial inter-animal variation, 2/5 animals in the treated groups showed nearly complete

abrogation of viral replication and little weight loss by day 7.

AAV delivery yields therapeutic ACE2 decoy levels in nonhuman primates

Next, we determined which AAV capsid is most efficient at transducing cells of the nonhuman

primate (NHP) proximal airways—the desired cellular targets for COVID-19 prophylaxis

following nasal delivery of vector. We administered vector using a previously approved

intranasal mucosal atomization device (MAD Nasal™), which comprises an atomizing tip with a

soft conical nostril seal fit on a standard syringe (Figure 4A). A mixture of 9 AAV serotypes

with uniquely barcoded transgenes were administered via the MAD Nasal™ to an NHP. Tissues

were harvested 14 days later for evaluation of relative transgene expression using the mRNA

bar-coding technique (Figure 4B)(23). The novel Clade A capsid (AAVrh91) we isolated from

macaque liver performed best in the nasopharynx and septum (Figure 4C and 4D) with low but

detectable expression levels in large airways and distal lung (Figure S6A-G). Clade E and F

capsids performed better than AAVrh91 in some non-target tissues such as distal lung (Figure

S6A-G). The profile of expression from AAVrh91 illustrates relative distribution of transgene

expression with proximal airway structures>intra-pulmonary conducting airway>distal lung

(Figure 4E).

To determine the candidate for clinical evaluation, we conducted a final NHP study where

groups of 2 animals were administered $5x10^{12}$ GC of vectors that differed with respect to capsid

(AAVhu68 vs. AAVrh91) and transgene cassettes (CDY14-Fc4 vs CDY14HL-Fc4). NLFs were

harvested on days 7, 14, and 28, and animals were necropsied on day 28 for biodistribution.

Analysis of pulmonary tissues from day 28 revealed broad distribution throughout the proximal

and distal airway, with AAVrh91 demonstrating superior gene transfer to proximal airway

structures, as suggested by the barcode study (Figure S6H-I). We estimated decoy protein

concentrations in the air-surface liquid (ASF) based on dilution-adjusted MS measurements of

NLF (Figure 4F). The effective concentrations at the ASF were in the range that demonstrated

neutralization in the *in vitro* assay. Expression was slightly higher with AAVrh91 vs. AAVhu68,

and CDY14HL-Fc4 vs. CDY14-Fc4. A subset of samples evaluated for binding to the spike

protein of SARS-CoV-2 showed a good correlation with decoy protein as measured by MS. This

indicates that the decoy protein produced *in vivo* in proximal airways is indeed functional (Figure

4G).

Based on these data, we selected a candidate for subsequent clinical evaluation called GTP404.

This candidate utilizes AAVrh91 as the capsid because of its transduction profile and

CDY14HL-Fc4 as the transgene because it retained broad and potent neutralizing activity in the

setting of an ACE2-disabling mutation. In preparation for IND-enabling studies, we administered

GTP404 at a 10-fold lower dose to two additional NHPs (Figure 4F). Impressive levels of decoy

protein were present in nasal ASF with concentrations only slightly reduced in comparison to

those achieved with the higher dose.

Discussion

The rapid emergence of more dangerous and transmissible variants of SARS-CoV-2 in this

pandemic is troubling, but not unexpected. The immunological pressures on the virus during

natural infections, following antibody therapies, and active vaccines have promoted the

emergence of variants(24). It appears that SARS-CoV-2 improved fitness through mutations that

both increased affinity for ACE2 and decreased neutralization by antibodies elicited to precursor

strains of the virus(15, 25, 26). The density of ACE2 in the nose and airway has been linked to

pathogenicity and transmissibility of SARS-CoV-2(27). The lower levels of ACE2 in the

proximal airways of children may be responsible for the lower infection rates and milder

symptoms in this group(28). It has been proposed that SARS-CoV-2 variants achieve greater

infection and transmission through increased affinity(25, 29); here we confirm increased affinity

for ACE2 in SARS-CoV-2 strains under positive selection during 2020.

Our original goal in engineering the ACE2 decoy was to improve its potency against SARS-

CoV-2, which we accomplished through affinity maturation against the Wuhan-Hu-1spike

protein in a yeast display system. Our selection strategy yielded a decoy with high binding and

neutralizing activity against a full range of SARS-CoV-2 variants, including B1.1.7 and B1.351,

which emerged from the UK and Republic of South Africa, respectively. However, we were

surprised to see equally potent binding and neutralization against other betacoronaviruses,

including SARS-CoV-1, which was responsible for the 2003 SARS pandemic. The presence of

several second-shell mutations in the affinity-matured decoy may contribute to this breadth since

the majority of the ACE2 contact surface was preserved (Figure S3). The engineered decoy may

be the Achilles' heel of any ACE2-dependent CoV whose primary driver of fitness – higher

binding to its receptor – should further enhance the potency of the ACE2 decoy.

We focused on IN delivery of AAV to express CDY14HL-Fc4 to prevent COVID-19. We used

the previously described hACE2-TG mouse challenge model to demonstrate efficacy of the

decoy *in vivo*. Treated animals lost less weight, showed reduced lung pathology, and showed less

replication of the challenge virus. Our results are consistent with the use of this model to

evaluate convalescent plasma(21), protease inhibitors(30), and monoclonal antibodies(31), where

weight loss, pulmonary pathology, and viral load were decreased, but not completely

abrogated.(21) We believe that the mouse challenge model underestimates the potential efficacy

of IN AAV-CDY14HL-Fc4. The dose of SARS-CoV-2 that results in human infection is likely

much lower, and therefore, easier to neutralize than the inoculating dose used in the mouse

challenge model (2.5×10^6 particles or 280 PFU). We used a novel AAV Clade A capsid called

AAVrh91 to maximize transduction in the proximal airways of NHPs. At a relatively low dose

(5×10^{11} GC), we achieved levels of CDY14HL-Fc4 in the ASF that should be sufficient to

neutralize SARS-CoV-2 variants. Based on our previous studies—using AAV to deliver broadly

neutralizing antibodies against influenza—we believe expression should be durable for at least

six months and can be effectively readministered(3-6),(32).

The emergence of three lethal and highly contagious CoV outbreaks in two decades – SARS in

2003, MERS in 2012, and COVID-19 in 2019 – suggests that CoVs will remain a threat to global

health. Surveillance of potential zoonotic sources of these CoVs, such as bats, revealed reservoirs

of related viruses capable of evolution and cross-species transmission(33). One possible

therapeutic application of CDY14HL-Fc4 is in the prevention and treatment of future outbreaks

caused by new CoVs that utilize ACE2 as a receptor. GTP404 could be rapidly deployed from

stockpiles to contain the initial outbreak and the CDY14HL-Fc4 protein can be leveraged to

improve outcomes in those who are infected. The CDY14HL-Fc4 products may be useful in the

current COVID-19 pandemic if SARS-CoV-2 variants confound current treatment and

prevention strategies. An immediate application could be in immune-suppressed individuals who

do not respond to traditional vaccines, develop chronic infection with SARS-CoV-2, and may be

reservoirs for new variants(24). The advantage of vector-expressed decoy in preventing COVID-

19 infections in immune-suppressed individuals is that this therapy does not rely on the

recipient's adaptive immune system to be effective.

Materials and Methods

Yeast display

We generated mutagenized ACE2 gene fragments by error prone PCR using the Diversify PCR
Random Mutagenesis Kit (TakaraBio) at multiple mutation levels, mixing the PCR products.
We used Gap-repair cloning and high-efficiency LiAc transformation(34) to assemble the ACE2
gene fragments into a centromeric plasmid. The plasmid contained an upstream Aga2 gene
fragment, a downstream HA epitope tag with flexible GSG linkers, and was driven by an
inducible GAL1 promoter, and contained a low-copy centromeric origin, similar to
pTCON2(35). Following transformation or sorting rounds, we passaged the libraries at 10X
diversity 3 times in SD-trp before inducing in log phase for 24 hrs at 30°C in SG-CAA(35). For
FACS, we stained the yeast with recombinant CoV2 RBD-His6 (Genscript) at diminishing
concentrations through the rounds (5nM, 1nM, 0.1nM +/- extended washes of up to 17 hrs for
off-rate sorting). We followed up with Mouse anti His6 (Genscript), rabbit anti-HA-PE (Cell
Signaling Technology), and goat anti mouse-488 secondary antibody (ThermoFisher) all in
phosphate buffered saline (PBS) with 0.1% BSA. Libraries were sorted and analyzed for RBD
binding and ACE2 expression at the Penn Flow Core on BD Influx and BD FACSAria II
instruments. We extracted plasmid from clones or pools of clones using the Yeast Plasmid
MiniPrep Kit (Zymo) and transformed this into bacteria for amplification.

Next-Generation Sequencing and Analysis

We performed 2x250 paired-end Illumina sequencing on randomly sheared and size-selected
ACE2 amplicons from the yeast display rounds. After removing adapters and low-quality reads,

we mapped clean reads no shorter than 200bp to the WT ACE2 nucleotide sequences using

NovoAlign (v.4.03.01). We translated in-frame sequences with mapping quality scores no lower

than 30 and without indels into amino acid sequences and compared to the WT ACE2 protein

sequence. We tallied non-synonymous changes for each codon across the ACE2 sequence (18-

615). We calculated mutation rates at each codon as follows: (sum of non-synonymous

AAs)/(sum of all AAs).

Expression of ACE2 variant IgG Fc4 fusions and RBDs

We sub-cloned candidate ACE2 synthetic DNA or decoy sequences from the yeast display

format into pCDNA3.1, using the endogenous ACE2 signal peptide and appending a human

IgG4 Fc domain (residues 99 to 327 from Uniprot reference sequence P01861) and a C-terminal

His6 tag. To generate protein for screening we transiently transfected HEK293 cells with

plasmid DNA in six-well plates using PEI and collected and clarified supernatant 72 hours later.

We quantified expression using the IgG4 Human ELISA kit (Invitrogen BMS2095) with IgG4

standards provided in the kit. For CDY14-Fc4 and CDY14HL-Fc4, we produced the protein in a

similar manner but purified it on protein A sepharose followed by dialysis and SDS-Page

analysis. We determined the concentration using the predicted extinction coefficient at 280nm.

We cloned the synthetic sequences (IDT gBlocks) of RBD [Spike amino acids 330-530 (CoV2),

317-516 (CoV1), and 318-517 (WIV1-CoV)] into pCDNA3.1 between an IL2 signal peptide plus

Gly-Ser and a C-terminal His6 tag. We transfected RBD plasmids into HEK293 cells using PEI

and collected supernatants 72 hrs later for clarification, concentration, and purification on Ni-

NTA resin, followed by dialysis into PBS. We confirmed purity using Coomassie-stained SDS-

PAGE analysis. We determined concentrations of the RBD using predicted extinction

coefficient at 280nm.

RBD Binding with SPR

We performed SPR binding analysis using a Biacore T200 instrument (GE Healthcare) at room

temperature in HBS-EP(+) buffer (10 mM HEPES pH 7.4, 150 mM NaCl, 3 mM EDTA, and

0.05% P20 surfactant, Cat# BR100669, Cytiva) using a protein A/G derivatized sensor chip

(Cat# SCBS PAGHC30M, XanTec Bioanalytics). We injected WT ACE2-hFc4 or CDY14-hFc4

diluted to 60nM in HBS-EP(+) at a flow rate of 10 µL/min for 3 min to capture ~1,000 response

units (RU) on the sensor surface in each cycle. We measured binding of various SARS-CoV

RBD proteins to this surface at concentrations ranging from 200 nM to 0.195 nM. RBD binding

was measured at a flow rate of 30 µL/min, with a 3 min association time and a 15 min

dissociation time. We performed regeneration between binding cycles using 10 mM glycine pH

1.5 injected at a flow rate of 60 µL/min for 1 min. KD values were determined for each

interaction using kinetics parameter fitting in the Biacore T200 Evaluation software. We used a

global 1:1 binding model and did not adjust for refractive index shift. Data presented are the

averate of two or more replicates were measured for each RBD domain tested.

CoV Pseudotyped Lentiviral Neutralization Assay

We obtained non-replicating lentivirus pseudotyped with CoV spike proteins from Integral

Molecular. The reporter virus particles encoded a renilla luciferase reporter gene. We set up

neutralization reactions with 100 ul of inhibitor diluted in full serum media and 10 ul of reporter

virus. After 1 hour at 37°C, we added 20,000 cells/well in 50 ul of a HEK 293T cell line

overexpressing ACE2 (Integral Molecular) and incubated the cells for 48 hours. We measured

reporter virus transduction activity on a luminometer (BioTek) using the Renilla Glo Kit

(Promega) following manufacturer's instructions. For higher throughput screens of neutralizing

potency, we used crude expression supernatant (described above) in the neutralization assay at 1

or 2 dilutions (typically 10- or 100-fold). We transformed the luciferase reading to an estimated

potency (EP) using the following formula: EP = (L*[decoy])/(1-L), where L is the fractional

luciferase level as compared to a mock sample (no inhibitor), and [decoy] is the concentration of

the decoy in the neutralization well. This was sufficient to rank clones without performing a full

titration.

AAV Vector Production

The University of Pennsylvania Vector Core produced recombinant AAV vectors as previously

described (36, 37)**.**

Decoy Quantification by Mass Spectrometry

Standards

Soluble hACE2Fc (produced in-house) was spiked at different levels (0.5-500 ng/mL) into PBS

or NLF acquired from a naïve rhesus macaque. Samples were denatured and reduced at 90°C for

10 minutes in the presence of 10mM dithiothreitol (DTT) and 2M Guanadinium-HCl (Gnd-HCl).

We cooled the samples to room temperature, then alkylated samples with 30mM iodoacetamide

(IAM) at room temperature for 30 minutes in the dark. The alkylation reaction was quenched by

adding 1µL DTT. We added 20mM ammonium bicarbonate to the denatured protein solution,

pH 7.5-8 at a volume to dilute the final Gnd-HCl concentration to 200mM. Trypsin solution was

added at ~4ng of trypsin per sample ratio and incubated at 37°C overnight. After digestion,

formic acid was added to a final of 0.5% to quench digestion reaction.

LC–MS/MS

We performed online chromatography with an Acclaim PepMap column (15 cm long, 300-µm

inner diameter) and a Thermo UltiMate 3000 RSLC system (Thermo Fisher Scientific) coupled

to a Q Exactive HF with a NanoFlex source (Thermo Fisher Scientific). During online analysis,

the column temperature was regulated to a temperature of 35°C. Peptides were separated with a

gradient of mobile phase A (MilliQ water with 0.1% formic acid) and mobile phase B

(acetonitrile with 0.1% formic acid). We ran the gradient from 4% B to 6% B over 15 min, then

to 10% B for 25 min (40 minutes total), then to 30% B for 46 min (86 minutes total). Samples

were loaded directly to the column. The column size was 75 cm x 15 um I.D. and was packed

with 2 micron C18 media (Acclaim PepMap). Due to the loading, lead-in, and washing steps, the

total time for an LC-MS/MS run was about 2 hours.

We acquired MS data using a data-dependent top-20 method for the Q Exactive HF; we

dynamically chose the most abundant not-yet-sequenced precursor ions from the survey scans

(200–2000 m/z). Sequencing was performed via higher energy collisional dissociation

fragmentation with a target value of 1e5 ions, determined with predictive automatic gain control.

We performed an isolation of precursors with a window of 4 m/z. Survey scans were acquired at

a resolution of 120,000 at m/z 200. Resolution for HCD spectra was set to 30,000 at m/z200 with

a maximum ion injection time of 50 ms and a normalized collision energy of 30. We set the S-

lens RF level at 50, which gave optimal transmission of the m/z region occupied by the peptides

from our digest. We excluded precursor ions with single, unassigned, or six and higher charge

states from fragmentation selection.

Data processing

We used BioPharma Finder 1.0 software (Thermo Fischer Scientific) to analyze all data. For

peptide mapping, we used a single-entry protein FASTA database to perform searches. The mass

area of the target peptide was plotted against the spike concentration to complete a standard

curve.

Selection of target peptide

Based on initial *in silico* studies, we selected four peptides as possible sequence-specific matches

for targeted quantification. We evaluated sensitivity performance for quantification of the four

peptide targets in the NLF background matrix. Following blank injections to establish system

cleanliness, replicate injections (n = 3) were made at all levels, from 0.5 ng/mL to 500 ng/mL.

Three of the peptides were detected with ANHYEDYGDYWR providing the greatest response

across the whole range. We determined retention time (RT) reproducibility across all samples (n

= 24) and determined peak area reproducibility and quantification accuracy for each level.

Excellent linearity was observed for the levels tested with typical $R^2 > 0.94$ for

ANHYEDYGDYWR. For ANHYEDYGDYWR, we observed excellent precision and accuracy

at all levels, with all replicates within 10% CV. For test articles, 1x or 10x NLF and/or

bronchoalveolar lavage fluid (BAL) is treated as previously described without any dilution or

protein precipitation. The mass area of target peptide in test articles was compared to the linear

calibration generated for the spiked material to determine the level of decoy present in the test

article.

ASF dilutions from serum and lavage urea

We used the urea concentrations in BAL or NLF and in serum collected at the same time to

determine the dilution that the lavage introduced to the ASF(38). We quantified urea in mouse

BAL and serum, and in NHP NLF using the Urea Assay Kit (Abcam). We obtained serum urea

concentrations from NHP from the blood urea nitrogen as part of standard bloodwork lab panels

(Antech).

Spike Binding ELISA

SARS-CoV-2 Spike Protein RBD (Sinobio #40592-V08H) was immobilized on a 96-well plate

(0.25ug/mL in PBS, 100ul/well) at 4^{o}C overnight. Plates were then washed 5x with PBS/0.05%

Tween and blocked with PBS/1.0% BSA for 1 hour with shaking. Samples (2x dilution in

PBS/2.0% BSA) and standards (soluble hACE2-Fc at starting concentration 100ng/ml, 12-point,

1:2 serial dilution, plus a 0.0ng/ml blank, in PBS/1.0% BSA) were added at 100ul/well in

duplicate and incubated for 2 hours at room temperature with shaking. Wells were washed as

described and biotin conjugated goat anti-human IgG (Jackson AffiniPure #109-065-098;

1:30,000 or Southern Biotech 2049-08; 1:1,000) in PBS/1.0% BSA detection antibody was

added to the wells at 100ul/well and incubated for 2 hours at room temperature with shaking.

Wells were washed as described, followed by the addition of 100ul/well Streptavidin-HRP

(Abcam #ab7403; 1:30,000) in PBS/1.0% BSA for 30 minutes with shaking. Wells were washed

as described and incubated in 100ul/well TMB substrate (Seracare #5120-0076) in the dark at

room temperature with shaking until reaction was stopped with 100ul/well TMB Stop Solution

(Seracare #5150-0021). Absorbances were read at 450 nm using a Spectramax M3 plate reader.

We exported and analyzed the data in GraphPad Prism Version 9.0.2. All raw data was blank

subtracted. We plotted a standard curve of soluble hACE2-Fc, and the X-axis (concentration)

was log_{10} transformed. We performed a 4-parameter nonlinear regression upon the transformed

standard curve, and interpolated sample concentrations.

Determination of Matrix Interference in BAL and NLF Samples

Soluble hACE2Fc was spiked into NLF (0.0, 0.5, 2.0, and 10.0 ng/ml) acquired from a naïve

rhesus macaque on the same plate with a standard curve (soluble hACE2Fc starting

concentration 100ng/ml, 12-point, 1:2 serial dilution, plus a 0.0ng/ml blank) in PBS/1.0% BSA.

We performed the spike binding assay and data analysis as described above.

Expression study in mice

All animal procedures were performed in accordance with protocols approved by the

Institutional Animal Care and Use Committee of the University of Pennsylvania. C57BL/6J mice

were purchased from The Jackson Laboratory. Anesthetized mice received an IN administration

of 10^{11} GC of AAVhu68.CDY14-Fc4, AAVrh91.CDY14-Fc4, AAVhu68.CDY14HL-Fc4, or

AAVrh91.CDY14HL-Fc in a volume of 50 µL or the same volume of vehicle control (PBS) on

day 0. On day 7, mice were euthanized and BAL was collected (1 ml of PBS administered

intratracheally).

hACE2 TG Mouse Study

To evaluate the prophylactic efficacy potential of AAV expressing hACE2 receptor decoys

against SARS-CoV-2, BIOQUAL, Inc. (Rockville, MD) conducted a challenge study using

hACE2 TG mice (Stock No: 034860, The Jackson Laboratory). Mice were administered with

either vehicle or 10^{11} GC of AAVhu68.CDY14HL-Fc4 IN on day -7 as described above. On day

0, mice were administered with mock or the SARS-CoV-2 challenge (50 µl of $2.8x10^2$ pfu of

SARS-CoV-2, USA_WA1/2020 isolate [NR-52281, BEI Resources]). Mice were euthanized on

either day 4 or 7 via cervical dislocation. BAL was collected as described above and aliquoted

for viral load assays into Trizol LS (Thermo Fisher Scientific, Waltham, MA) or heat inactivated

(60°C for 30 minutes) for decoy protein expression. The lung was collected and split for

histopathology into 10% neutral buffered formalin or snap frozen for viral load analysis. RNA

extraction for RT-qPCR, the quantitative RT-PCR assay for SARS-CoV-2 RNA, and

subgenomic RNA were performed as described(39).

Histopathology of Collected Organs

The organs collected at necropsy were trimmed and routinely processed for hematoxylin and

eosin (H&E) staining. Slides were blindly evaluated by a blinded pathologist using a severity

score of 0 (no lesions observed), 1 (minimal), 2 (mild), 3 (moderate), 4 (marked) and 5 (severe)

for each finding.

Intranasal capsid comparison by AAV barcoding

We generated a set of custom barcoded plasmids using degenerate nucleotides that anneal

immediately downstream of the stop codon in a GFP reporter construct that contains the AAV2

ITRs. We produced barcoded AAV vectors for each serotype in the study separately by

transfecting HEK293 cells as described(36), replacing the typical single ITR-containing plasmid

in the transfection mix with an equimolar mixture of 4 uniquely barcoded reporter constructs.

We pooled the individual vector preps on an equimolar basis using their digital droplet PCR

titers. We determined the absolute barcode distribution in the AAV pool by deep sequencing; we

extracted AAV genomes from the pool and performed linear-range PCR using primers that flank

the barcode region to generate an amplicon for paired-end Illumina sequencing.

NHP studies

Rhesus and cynomolgus macaques were obtained from Primgen (PreLabs). NHP studies were

conducted at the University of Pennsylvania or Children's Hospital of Philadelphia within

facilities that are United States Department of Agriculture-registered, Association for

Assessment and Accreditation of Laboratory Animal Care-accredited, and Public Health Service-

assured. For the barcode study, 4×10^{12} GC of the pool AAV preps was delivered IN in a total

volume of 0.28 ml to an adult male rhesus macaque using the MAD Nasal™ device. After 14

days, we collected airway tissues at necropsy, and extracted total RNA using Trizol Reagent

(Thermo Fisher). We generated cDNAs using Superscript III reverse transcriptase

(ThermoFisher) and an oligo dT primer. We used the cDNAs to prepare barcode amplicons for

Illumina sequencing as described above. We extracted the relative barcode abundances in input

(AAV mixture) and output (tissue cDNAs) from Illumina data. The ratio of output to input

relative abundances for each barcode in each tissue is proportional to the relative efficiency of

the capsid linked to that barcode in that tissue. Agreement among the 4 barcodes assigned to

each capsid allows us to assess assay noise, and detect rare, tissue-specific effects of the barcode

itself on transcript stability (none detected). For each tissue, we quantified the total capsid-

derived transcript per ug of total RNA using qPCR with a primer/probe set common to all the

barcoded reporters.

For the decoy expression in NHPs, cynomolgus macaques (n=2/vector) were administered IN

with 5×10^{12} GC of AAVhu68.CDY14, AAVrh91.CDY14, AAVhu68.CDY14HL, or

AAVrh91.CDY14HL as described above. An additional two NHPs were administered with

5×10^{11} GC of AAVrh91.CDY14HL. All NHPs were negative for pre-existing neutralizing

antibody titres to the administered AAV capsid prior to study initiation (Immunology Core at the

Gene Therapy Program). Animals were monitored throughout the in-life phase for complete

blood counts, clinical chemistries, and coagulation panels by Antech Diagnostics (Lake Success,

NY). On days 7, 14, and 28 NLF was collected (animals placed in ventral recumbency with head

tilted to the right, up to 5 mL of PBS delivered in 1mL aliquots, and fluid collected via gravity).

Animals were necropsied on day 28 and a full histopathological evaluation was performed.

Ethics Statement for Study Conducted at BIOQUAL (hACE2 TG Mouse Challenge Study)

This research was conducted under BIOQUAL Institute Institutional Animal Care and Use

Committee (IACUC) approved protocol number 21-005, in compliance with the Animal Welfare

Act and other federal statutes, and regulations relating to animals and experiments involving

animals. BIOQUAL is accredited by the Association for Assessment and Accreditation of

Laboratory Animal Care International and adheres to principles stated in the Guide for the Care

and Use of Laboratory Animals, National Research Council. Animals were monitored twice

daily for clinical signs (specifically ruffled fur, heavy breathing, lethargy) and weighed daily.

Statistical analysis

Statistical analyses performed using *R* (version 4.0.0). Statistical tests described in figure

legends.

Acknowledgments

We thank Hailey Shankle, Henry Hoff, Shiva Shrestha for notable contributions to the decoy development and characterization. We thank Nathan Denton for assistance with manuscript preparation and graphics. We thank Kirsten Copren and Maggie Shaw for sequencing and analysis support. We thank Victoria Kehm for histology processing and James Tarrant for histopathology analysis. We thank the Immunology Core and the Program for Comparative Medicine of the Gene Therapy Program at the University of Pennsylvania for study support. We thank the Flow Cytometry and Cell Sorting Resource Laboratory at the University of Pennsylvania for cell sorting. All vectors were produced by the Penn Vector Core.

Funding

This work was funded by the Harrington Discovery Institute, the NHLBI Gene Therapy Resource Program (75N92019D00016), and internal Gene Therapy Program resources.

Author Contributions

J.J.S. – conceptualization, investigation, methodology, project administration, writing-original draft, writing-review and editing; J.A.G. – formal analysis, methodology, project administration, writing-review and editing; K.T.M. – methodology, resources, writing-original draft, writing-review and editing; S.L. – investigation; R.A.M. – investigation; R.M. – investigation; K.B.T. – investigation, methodology; K.N. – resources; C.D. – methodology, resources; C.H. – conceptualization, methodology; M.H. investigation; H.Y. – formal analysis; X.H. – formal

565 analysis, software; S.J.C. – investigation; J.M.W. – conceptualization, funding acquisition,

566 methodology, supervision, writing-original draft, writing-review and editing.

568 **Conflict of Interest Statement**

576 **Data Availability Statement**

577 All datasets presented in this study are included in the article/Supplementary Material.

579 **List of Supplementary Materials**

580 Figures S1-S6

References

1. Chan KK, Tan TJC, Narayanan KK, Procko E. An engineered decoy receptor for SARS-CoV-2 broadly binds protein S sequence variants. Sci Adv. 2021;7(8).

2. Glasgow A, Glasgow J, Limonta D, Solomon P, Lui I, Zhang Y, et al. Engineered ACE2 receptor traps potently neutralize SARS-CoV-2. Proc Natl Acad Sci U S A. 2020;117(45):28046-55.

3. Adam VS, Crosariol M, Kumar S, Ge MQ, Czack SE, Roy S, et al. Adeno-associated virus 9-mediated airway expression of antibody protects old and immunodeficient mice against influenza virus. Clin Vaccine Immunol. 2014;21(11):1528-33.

4. Laursen NS, Friesen RHE, Zhu X, Jongeneelen M, Blokland S, Vermond J, et al. Universal protection against influenza infection by a multidomain antibody to influenza hemagglutinin. Science. 2018;362(6414):598-602.

5. Limberis MP, Adam VS, Wong G, Gren J, Kobasa D, Ross TM, et al. Intranasal antibody gene transfer in mice and ferrets elicits broad protection against pandemic influenza. Sci Transl Med. 2013;5(187):187ra72.

6. Limberis MP, Racine T, Kobasa D, Li Y, Gao GF, Kobinger G, et al. Vectored expression of the broadly neutralizing antibody FI6 in mouse airway provides partial protection against a new avian influenza A virus, H7N9. Clin Vaccine Immunol. 2013;20(12):1836-7.

7. Angelini A, Chen TF, de Picciotto S, Yang NJ, Tzeng A, Santos MS, et al. Protein Engineering and Selection Using Yeast Surface Display. Methods Mol Biol. 2015;1319:3-36.

8. Chan KK, Dorosky D, Sharma P, Abbasi SA, Dye JM, Kranz DM, et al. Engineering human ACE2 to optimize binding to the spike protein of SARS coronavirus 2. Science. 2020;369(6508):1261-5.

9. Aguinaldo AM, Arnold FH. Staggered extension process (StEP) in vitro recombination. Methods Mol Biol. 2003;231:105-10.

10. Eckert-Boulet N, Pedersen ML, Krogh BO, Lisby M. Optimization of ordered plasmid assembly by gap repair in Saccharomyces cerevisiae. Yeast. 2012;29(8):323-34.

11. Boder ET, Midelfort KS, Wittrup KD. Directed evolution of antibody fragments with monovalent femtomolar antigen-binding affinity. Proc Natl Acad Sci U S A. 2000;97(20):10701-5.

12. Guy JL, Jackson RM, Jensen HA, Hooper NM, Turner AJ. Identification of critical active-site residues in angiotensin-converting enzyme-2 (ACE2) by site-directed mutagenesis. FEBS J. 2005;272(14):3512-20.

13. Starr TN, Greaney AJ, Addetia A, Hannon WW, Choudhary MC, Dingens AS, et al. Prospective mapping of viral mutations that escape antibodies used to treat COVID-19. Science. 2021;371(6531):850-4.

14. Thomson EC, Rosen LE, Shepherd JG, Spreafico R, da Silva Filipe A, Wojcechowskyj JA, et al. Circulating SARS-CoV-2 spike N439K variants maintain fitness while evading antibody-mediated immunity. Cell. 2021;184(5):1171-87 e20.

15. Wang P, Nair MS, Liu L, Iketani S, Luo Y, Guo Y, et al. Antibody Resistance of SARS-CoV-2 Variants B.1.351 and B.1.1.7. Nature. 2021.

16. Oude Munnink BB, Sikkema RS, Nieuwenhuijse DF, Molenaar RJ, Munger E, Molenkamp R, et al. Transmission of SARS-CoV-2 on mink farms between humans and mink and back to humans. Science. 2021;371(6525):172-7.

17. Menachery VD, Yount BL, Jr., Sims AC, Debbink K, Agnihothram SS, Gralinski LE, et al. SARS-like WIV1-CoV poised for human emergence. Proc Natl Acad Sci U S A. 2016;113(11):3048-53.

18. Weissman D, Alameh MG, de Silva T, Collini P, Hornsby H, Brown R, et al. D614G Spike Mutation Increases SARS CoV-2 Susceptibility to Neutralization. Cell Host Microbe. 2021;29(1):23-31 e4.

19.   Yurkovetskiy L, Wang X, Pascal KE, Tomkins-Tinch C, Nyalile TP, Wang Y, et al. Structural and
Functional Analysis of the D614G SARS-CoV-2 Spike Protein Variant. Cell. 2020;183(3):739-51 e8.
20.   Winkler ES, Bailey AL, Kafai NM, Nair S, McCune BT, Yu J, et al. SARS-CoV-2 infection of human
ACE2-transgenic mice causes severe lung inflammation and impaired function. Nature Immunology.
2020;21(11):1327-35.
21.   Zheng J, Wong L-YR, Li K, Verma AK, Ortiz ME, Wohlford-Lenane C, et al. COVID-19 treatments
and pathogenesis including anosmia in K18-hACE2 mice. Nature. 2021;589(7843):603-7.
22.   Yinda CK, Port JR, Bushmaker T, Offei Owusu I, Purushotham JN, Avanzato VA, et al. K18-hACE2
mice develop respiratory disease resembling severe COVID-19. PLOS Pathogens. 2021;17(1):e1009195.
23.   Adachi K, Enoki T, Kawano Y, Veraz M, Nakai H. Drawing a high-resolution functional map of
adeno-associated virus capsid by massively parallel sequencing. Nat Commun. 2014;5:3075.
24.   Kemp SA, Collier DA, Datir RP, Ferreira I, Gayed S, Jahun A, et al. SARS-CoV-2 evolution during
treatment of chronic infection. Nature. 2021.
25.   Ozono S, Zhang Y, Ode H, Sano K, Tan TS, Imai K, et al. SARS-CoV-2 D614G spike mutation
increases entry efficiency with enhanced ACE2-binding affinity. Nat Commun. 2021;12(1):848.
26.   Zhou D, Dejnirattisai W, Supasa P, Liu C, Mentzer AJ, Ginn HM, et al. Evidence of escape of SARS-
CoV-2 variant B.1.351 from natural and vaccine-induced sera. Cell. 2021.
27.   Zhang H, Rostami MR, Leopold PL, Mezey JG, O'Beirne SL, Strulovici-Barel Y, et al. Expression of
the SARS-CoV-2 ACE2 Receptor in the Human Airway Epithelium. Am J Respir Crit Care Med.
2020;202(2):219-29.
28.   Bunyavanich S, Do A, Vicencio A. Nasal Gene Expression of Angiotensin-Converting Enzyme 2 in
Children and Adults. JAMA. 2020;323(23):2427-9.
29.   Ramanathan M, Ferguson ID, Miao W, Khavari PA. SARS-CoV-2 B.1.1.7 and B.1.351 Spike
variants bind human ACE2 with increased affinity. bioRxiv. 2021.
30.   Joaquín Cáceres C, Cardenas-Garcia S, Carnaccini S, Seibert B, Rajao DS, Wang J, et al. Efficacy of
GC-376 against SARS-CoV-2 virus infection in the K18 hACE2 transgenic mouse model. bioRxiv.
2021:2021.01.27.428428.
31.   Rosenfeld R, Noy-Porat T, Mechaly A, Makdasi E, Levy Y, Alcalay R, et al. Post-exposure
protection of SARS-CoV-2 lethal infected K18-hACE2 transgenic mice by neutralizing human monoclonal
antibody. Nature Communications. 2021;12(1):944.
32.   Limberis MP, Wilson JM. Adeno-associated virus serotype 9 vectors transduce murine alveolar
and nasal epithelia and can be readministered. Proc Natl Acad Sci U S A. 2006;103(35):12993-8.
33.   Latinne A, Hu B, Olival KJ, Zhu G, Zhang L, Li H, et al. Origin and cross-species transmission of bat
coronaviruses in China. Nat Commun. 2020;11(1):4235.
34.   Gietz RD, Schiestl RH. Large-scale high-efficiency yeast transformation using the LiAc/SS carrier
DNA/PEG method. Nat Protoc. 2007;2(1):38-41.
35.   Chao G, Lau WL, Hackel BJ, Sazinsky SL, Lippow SM, Wittrup KD. Isolating and engineering
human antibodies using yeast surface display. Nat Protoc. 2006;1(2):755-68.
36.   Lock M, Alvira M, Vandenberghe LH, Samanta A, Toelen J, Debyser Z, et al. Rapid, simple, and
versatile manufacturing of recombinant adeno-associated viral vectors at scale. Hum Gene Ther.
2010;21(10):1259-71.
37.   Lock M, Alvira MR, Chen SJ, Wilson JM. Absolute determination of single-stranded and self-
complementary adeno-associated viral vector genome titers by droplet digital PCR. Hum Gene Ther
Methods. 2014;25(2):115-25.
38.   Kaulbach HC, White MV, Igarashi Y, Hahn BK, Kaliner MA. Estimation of nasal epithelial lining
fluid using urea as a marker. J Allergy Clin Immunol. 1993;92(3):457-65.
39.   Baum A, Ajithdoss D, Copin R, Zhou A, Lanza K, Negron N, et al. REGN-COV2 antibodies prevent
and treat SARS-CoV-2 infection in rhesus macaques and hamsters. Science. 2020;370(6520):1110-5.

40.   Xu C, Wang Y, Liu C, Zhang C, Han W, Hong X, et al. Conformational dynamics of SARS-CoV-2
trimeric spike glycoprotein in complex with receptor ACE2 revealed by cryo-EM. Sci Adv. 2021;7(1).

 Figure Legends

Fig. 1. ACE2 Decoy Receptor Engineering. A. Yeast display (YD) system **B**. Decoy affinity

maturation and candidate selection process. **C**. Flow cytometry analysis of the naïve (purple)

and sorted populations (gray) from the secondary YD library. **D**. NGS analysis of plasmid

populations recovered from rounds of YD. **E**. Mutations accumulated in the top five decoy-Fc4

candidates. **F and G**. SPR binding analysis for CoV-2 RBD injected over surface-immobilized

ACE2-wt (F) or CDY14HL (G). **H**. Wuhan CoV-2-Pseudotyped lentiviral reporter neutralization

assay of ACE2-wt-Fc4, CDY14-Fc4, and CDY14HL-Fc4. Data for at least three independent

measurements are presented as average ± standard deviation.

Fig. 2 ACE2 Decoy Binding and Neutralization Across Diverse CoVs. A. Structural models

(7DF4.pdb (40)) of Wuhan CoV-2 RBD (gray) bound to human ACE2 (blue) RBD residues that

distinguish each variant (listed below each panel) from the Wuhan CoV-2 RBD are shown in red

spheres. **B**. SPR measurements: ACE2-wt-Fc4 or CDY14HL-Fc4 binding to various purified

recombinant RBD proteins. **C**. We used reporter lentiviruses pseudotyped with CoV spike

proteins from five isolates to determine the neutralizing potencies of CDY14HL-Fc4. Reporter

virus activity data are presented as mean ± standard deviation for at least three replicate

titrations**.**

718
Figure 3. Protection in the human ACE2 transgenic mouse model. BAL from vector-treated

animals analyzed for: **A.** decoy protein by MS; **B.** SARS-CoV-2 spike ELISA; **C.** neutralization

of SARS-CoV-2 pseudotyped lentivirus. **D.** Challenge study design. **E.** Weight loss in the

animals that were sustained for 7 days; one animal in the vehicle and vector treated groups

required euthanasia. **F**. MS assay of expression in ASF (corrected for BAL dilution). **G.**

Pulmonary inflammation histopathology scores of tissues harvested at days 4 and 7. **H.** Viral

RNA in BAL. **I.** Viral RNA in lung. **J.** Sub-genomic RNA in lung. Outliers are indicated with X.

738
Figure 4. **AAV-Delivered Decoy Expression in the Airway of NHP. A.** NHP vector delivery via

MAD Nasal™. **B**. Pooled capsid comparison using mRNA barcoding. mRNA barcode

enrichment scores (average across 4 barcodes $\pm$ SD) in (**C**) nasopharynx and (**D**) septum. **E.**

AAVrh91 transduction profile in airway barcode study. **F**. MS assay of expression in ASF

(corrected for BAL dilution) for NHPs (2/group) dosed with ACE2 decoy vector. **G.** Correlation

between decoy protein by MS and spike binding by ELISA. Data includes d7 and d14 samples

from (F) plus NLF of 3 naïve macaques. We excluded one naïve animal from ELISA analysis

because of background binding presumably due to a prior coronavirus infection.