State-dependent network interactions differentially gate sensory input at the motor and command neuron level in *Caenorhabditis elegans*

Zachary T. Cecere[1,2,3,6], Kathleen T. Quach[1,2,6], Eviatar Yemini[4,5], Javier J. How[1,2], Tatyana O. Sharpee[1,3], and Sreekanth H. Chalasani[1,2]*

[1]Neurosciences Graduate Program, University of California, San Diego, La Jolla, CA 92093

[2]Molecular Neurobiology Laboratory, The Salk Institute for Biological Studies, La Jolla, CA 92037

[3]Computational Neurobiology Laboratory, The Salk Institute for Biological Studies, La Jolla, CA 92037

[4]Department of Biological Sciences, Howard Hughes Medical Institute, Columbia University, New York, NY 10027

[5]University of Massachusetts Chan Medical School, Worcester, MA 01655

[6]These authors contributed equally.

*Corresponding author

Email: schalasani@salk.edu

Author Contributions: Z.T.C. designed experiments. Z.T.C. and J.J.H. performed experiments. E.Y. contributed provided strains. Z.T.C. and K.T.Q. analyzed data. K.T.Q., Z.T.C, and S.H.C. wrote the paper. T.O.S. and S.H.C. supervised research.

Competing Interest Statement: The authors declare no competing interests.

Classification: Biological Sciences, Neuroscience

Keywords: *C. elegans,* whole-brain imaging, network states, sensory gating, soft decision trees

This PDF file includes:

      Main Text
      Figures 1 to 4

Abstract

Neural responses are influenced by both external stimuli and internal network states. While
network states have been linked to behavioral and stimulus states, little is known about how
sensory inputs are filtered by whole-brain activity to affect motor and command neurons. Here,
we recorded whole-brain activity of *Caenorhabditis elegans* experiencing bacterial food stimuli,
and modeled how sensory inputs affect motor and command neurons in a network state-
dependent manner. First, we classified active neurons into six functional clusters: two sensory
neuron clusters (ON, OFF), and four motor/command neuron clusters (AVA, RME, SMDD,
SMDV). Using encoding models, we found that ON and OFF sensory neurons that respond to
onset and removal of bacteria, respectively, employ different adaptation strategies. Next, we used
decoding models to show that bacterial onset and removal differentially drive AVA and RME
cluster activity. To explore state-dependent effects on AVA and RME clusters, we developed a
model that identified network states and fitted submodels for each state to predict how each of
the six functional clusters drive AVA and RME cluster activity. We also identified network states in
which AVA and RME clusters were either largely unperturbed by or receptive to bacterial sensory
input. Furthermore, this model allowed us to disentangle the state-dependent contributions of
stimulus timescales and bacterial content to neural activity. Collectively, we present an
interpretable approach for modeling network dynamics that goes beyond implication of neurons in
particular states, and moves toward explicitly dissecting how neural populations work together to
produce state dependence.

Significance Statement

A major function of the brain is to transform sensory information into behavior. As the first
receiver of sensory input, sensory neuron activity is often most correlated with stimulus features.
However, this high-fidelity representation of sensory input becomes diluted as it travels to
downstream neurons, where sensory information is integrated with network activity. By the time
sensory information reaches motor neurons, it is often difficult to dissociate the influence of
sensory input from the influence of network activity. Here, we describe a method that is fully
interpretable such that we can show how neural populations on a whole-brain scale interact to
produce network states. From there, we can attribute motor neuron activity to network history and
sensory input.

Main Text

Introduction

Constant stimuli can have variable influences on neural responses and behavior. On the neural
population level, this variability becomes more pronounced as sensory information is transformed
by downstream neurons. As the initial receiver of sensory information, sensory neurons exhibit
activity that is the most correlated with stimulus features, but are susceptible to sensor errors that
can propagate to behavior (1, 2). At the perception level, sensory input can be modulated by
attention (3, 4), cognitive load (5), perceptual learning (6, 7), internal noise (8, 9), and internally
generated coordinated activity (10). At the motor coordination level, gating of sensory input has
been observed occurring in phase with behaviors such as locomotion (11, 12) and active
whisking (13). However, little is known about how neural populations interact on a global scale to
produce network states that modulate how sensory input is gated at the motor neuron level.
        To disentangle the influences of stimuli and internal network state on downstream neural
activity, precise stimulus control and whole-brain imaging are both needed to accurately account
for experienced stimuli and global network states. Despite advances in modern imaging
technology, most studies are limited to imaging small subsets of a brain's total activity (14–16).
This problem is alleviated by studying simpler animals like zebrafish, fly larvae, and adult *C.
elegans* where neural activity at single-cell resolution can be monitored across the entire brain

(17–20). With only 302 neurons, 189 of which are located in the head, the nematode *C. elegans*
is ideally suited for whole-brain functional imaging studies. *C. elegans* whole-brain activity can be
monitored both in restrained and freely moving animals (20–22). Moreover, whole-brain imaging
of restrained animals in a microfluidic chip (20) allows for precise, fast, and complex stimuli
presentations, thereby enabling investigations of stimulus-evoked whole-brain global dynamics
underlying sensory coding (23, 24), motor states (25), and physiological states (26, 27). While
these studies demonstrate the utility of whole-brain imaging and provide insights into the
nematode nervous system, modeling of global network activity has largely focused on identifying
which neurons and activity trends underlie particular behavioral or sensory states, but often fall
short of explaining how populations work together to influence neural activity.
86   Here, we imaged the calcium activity of the entire *C. elegans* head while we presented it
with rapidly fluctuating sequences of bacterial food stimuli and control buffer. We chose bacterial
food instead of single-compound odorants in order to study a complex stimulus that is
immediately relevant to *C. elegans*. We correlated activity of individual neurons to show that
active neurons can be divided into six functional clusters groups. We then used encoding and
decoding models to characterize basic properties of how stimuli drive sensory neurons and
motor/command neurons. Finally, we built a hybrid model to identify network states and build
submodels to explicitly show how sensory and motor populations drive motor/command neuron
activity in each state. This model allowed us to identify network states in which motor/command
neurons were either unresponsive or responsive to sensory input. Furthermore, we revealed how
stimulus features and sensory context were differentially gated in a state-dependent fashion.
97
98   **Results**
99
Food-stimulated whole-brain activity reveals six functional cell clusters
We used an automated microfluidic system (28) to simultaneously image calcium activity in *C.*
elegans head neurons and present the animal's nose with pulse-based stimulus sequences that
rapidly fluctuated between liquid flows of bacterial food stimulus (from channel 1) and control
buffer (from channel 2) (Fig. 1A, see Materials and Methods: Stimulus delivery). We refer to this
as the bacteria↔buffer stimulus sequence. To control for artifacts intrinsic to the microfluidic
setup, we also imaged activity while *C. elegans* was presented with a control buffer↔buffer
stimulus sequence that fluctuated between two chemically identical buffer flows. This microfluidic
system was previously used to show that individual chemosensory neurons detect and respond to
bacteria (29, 30). To monitor whole-brain activity (see Materials and Methods: Whole-brain
imaging), we used a strain that expressed a genetically-encoded nuclear-localized calcium
indicator (GCaMP5K) (25,31). This strain was previously used to obtain whole-brain activity from
restrained *C. elegans* during controlled delivery of stimuli (20, 25). In addition to this primary
strain, we also recorded calcium activity (GCaMP6s) (32) from a strain that expressed NeuroPAL
(24), which labels all *C. elegans* neurons with an invariant multicolor fluorescence map and
allows for unambiguous identification of neurons (Fig. 1B-D). We used this as a supplementary
strain to confirm cell identity associated with activity patterns found in the primary strain.
117   We found that active neurons could be divided into six functional clusters based on
correlated changes in GCaMP fluorescence: two sensory neuron clusters (ON and OFF) and four
motor/command neuron clusters (AVA, RME, SMDD, SMDV) (Fig. 1E–H). We first identified
primary sensory neurons by looking for neurons that were either positively correlated (ON cells)
or negatively correlated (OFF cells) with bacteria pulse onset during bacteria↔buffer stimulus
sequences. Specifically, for each neuron, we first subtracted the changes in fluorescence induced
by bacteria removals from the changes in fluorescence induced by bacteria onsets, and then
ranked each neuron according to this score (see Materials and Methods: ON and OFF sensory
neuron classification). The neurons with the highest and lowest scores were then checked for
additional criteria before being classified as ON cells and OFF cells, respectively. ON cells were
categorized as those neurons that obviously and immediately increased activity upon all bacteria
onsets, and immediately decreased upon bacteria removals (Fig. 1I). Conversely, OFF cells were
classified as neurons that decreased activity upon bacteria onsets and increased upon bacteria

removals (Fig. 1I). OFF cells were additionally required to rapidly decrease activity in response to
initial bacterial onsets, dropping well below their pre-stimulus baseline, indicating that these
sensory neurons are inhibited by bacteria rather than activated by the control buffer flow (Fig. 1I).
This OFF cell behavior of being inhibited by a stimulus was absent in the control buffer↔buffer
stimulus sequences. Instead, we found only ON cells that were activated by either the onset of
the channel 1 buffer (ON-1 cells) or the onset of channel 2 buffer (ON-2 cells) (Fig. S1).
Additionally, we identified fewer sensory neurons in animals presented with buffer↔buffer
stimulus sequences (Fig. S1A) than with bacteria↔buffer stimulus sequences (Fig. 1F,G). This
suggests that additional sensory neurons respond during bacteria ↔ buffer stimulus sequences,
compared to buffer↔buffer stimulus sequences. Therefore, bacteria↔buffer stimulus sequences
activate ON and OFF sensory neurons that respond to bacteria onset and removal, respectively,
rather than to bacteria onset and buffer onset.
142   The vast majority of neurons did not respond immediately to stimulus changes (Fig. 1F–
H). Across these neurons, we observed two pairs of stereotypical activity patterns: a pair of
clusters that were anti-correlated and bistable (Fig. 1H, red and blue), and a pair of clusters that
were anti-correlated and moderately fast (Fig. 1H, purple and yellow). Using the NeuroPAL–
GCaMP6s strain to identify neurons, we found that RME motor neurons and AVE command
neurons were anti-correlated and bistable (Fig. 1J), consistent with previous findings (25). We
also found that SMDD and SMDV neurons were anti-correlated and exhibited moderately fast
dynamics (Fig. 1K). These 4 representative motor and command neurons are associated with
forward locomotion (RME), reverse locomotion (AVA), dorsal turning (SMDD), and ventral turning
(SMDV) in *C. elegans* (25, 34–37). Using AVA, RME, SMDD, and SMDV as representative
neurons, we sorted non-sensory neurons into clusters based on how their activity correlated with
these four representative neurons (Fig. 1I–K). Many of the low noise neuronal traces strongly
correlated (>85%) with either the identified RME motor neurons or AVA command neurons (Fig.
1F,G). A number of other calcium traces appeared as distorted, noisy versions of AVA or RME
(Fig. S2), and the counts shown in the Fig. 1F,G are likely an underestimate of the true number of
cells that are highly correlated with AVA and RME. In contrast, SMDV and SMDD were often the
sole members of their eponymous clusters, usually with no other neurons that strongly correlated
with these neurons (Fig. 1F,G). AVA, and RME were previously shown to have strong positional
stereotypy, and thus, after identifying these neurons with NeuroPAL, their activity signature and
stereotyped location made them easy to identify in the absence of NeuroPAL. SMDV and SMDD
neurons were also readily identified by their previously reported distinctive activity signature (37).
Therefore, we were able to subsequently identify all four motor/command neuron clusters in non-
NeuroPAL-GCaMP5K animals without a coinciding NeuroPAL map (Fig. 1F,G).
165   These six cell clusters appeared in every animal, and we could not identify any other cell
cluster that appeared consistently across all animals exposed to bacteria↔buffer stimulus
sequences. We also observed similar clustering of active neurons in animals exposed to
buffer↔buffer stimulus sequences (Fig. S1A). Putative cells not appearing in one of the
aforementioned six cell clusters typically resembled noise or noisier versions of the activity
patterns exhibited by the six cell clusters. To reduce dimensionality of the dataset with little loss of
information (due to high correlation), we averaged across the activity traces of all neurons within
a cluster (1I–K). Hence, all subsequent modeling used cluster-averaged representations to
simplify model structure.
174
Bacteria addition and removal differentially drive activity of sensory, AVA, and RME
clusters
To analyze how ON and OFF sensory neurons differ in their responses beyond the single-pulse
timescale, we built encoding models to predict how sensory neuron activity adapts to repeated
stimuli presentations (Fig. 2A, see Materials and Methods: Encoding model). Both ON and OFF
cells are driven away from baseline activity (in opposite directions) upon bacteria onset and return
to baseline upon bacteria removal. ON and OFF cells habituate to repeated bacteria
presentations such that the change (increase for ON, decreases for OFF) in fluorescence from
baseline is smaller in subsequent bacteria pulses (Fig. 1I). Here, we explored which timescale of

adaptation best described ON and OFF cells: 1) perfect adaptation, in which neurons cease
responding despite persistent stimulation, 2) imperfectly adapting adaptation, in which neurons
attenuate but do not fully terminate their responses, and 3) non-adaptation, in which neural
responses are unaffected by recent stimulus history.
To examine the timescales of adaptation in ON and OFF cells, we used a cascade model
that was previously used to describe adaptation to odors in *C. elegans* sensory neurons (38).
This model is built on a cascade of simple ordinary differential equation (ODE) models of the
form:

$$dx_i/dt = \tau * ([input] - x_i(t))$$
$$X(t) = x_1(t) - x_2(t)$$

where τ is the time constant that controls how fast a linear temporal filter responds to stimulus
. The temporal filter describes how the recent history of the stimulus contributes to the
current value of inferred calcium level of the cell,   . With τ constrained to be positive,   is
guaranteed to exhibit perfect adaptation. That is, when encountering a step-change stimulus,
will briefly change activity before terminating the response and returning to its baseline level.
This model performed well for uncorrelated stimulus patterns but struggled considerably on
correlated patterns (38). Kato and colleagues supposed these issues could be mediated by
including more than two of the simple ODE filters. In other words, the authors hypothesized that
C. elegans sensory cells adapt at more than one timescale. To test this hypothesis, as well as to
ascertain whether the perfect adaptation assumption is justified, we simplified and generalized
this model:

$$\frac{dx_i}{dt} = \tau \cdot ([input] - x_i(t))$$
$$x_i(t) = \sum a_i x_i(t)$$
$$g(t) = (x^P \circledast f_g)(t)$$

where   is the inferred calcium level of the cell and is calculated as the sum of the temporal
filters (ODE model solutions),   is the GCaMP filter, and   is a positive value required for the
GCaMP transformation. This model learns the coefficients τ and   for an arbitrary number of
ODE basis function, and thus can learn adaptation on multiple timescales. Moreover, this model
can test the effects of perfect and imperfect adaptation on model fit by toggling the coefficient
constraint, such that perfect adaptation entails the following constraint:

$$\sum a_i = 0$$

This model formulation produced consistent and robust fits to *C. elegans* sensory
neurons and allowed us to test the effects of different model complexities. In this approach, each
sensory neuron's fluorescence trace was divided into three contiguous blocks. We used a 3-fold
cross validation approach (i.e., two of the blocks are used to fit the model, while the third is used
for testing) to assess model performance as a function of the number of basis filters and the type
of adaptation. For ON cells, all adapting models significantly outperformed the non-adapting
model according to a hierarchical bootstrap ($p < 0.05$ with Bonferroni correction, Fig. 2B, S3A). A
perfectly adapting model with one primary filter and one adapting filter performed best, matching
previous findings (38). Similarly, OFF cell adapting models outperformed the non-adapting model
(Fig. 2C, S3B). However, unlike ON cells, OFF cells were best described by an imperfectly
adapting model with one primary filter and one adapting filter, while the perfectly adapting model
was the worst-performing model (not significantly better than non-adapting model) (Fig. 2C, S3B).
It is possible that OFF sensory cells perfectly adapt over a longer timescale, but the fast

adaptation relevant to this study is imperfect in OFF cells. Using more than two basis functions
(one primary filter and one adapting filter) hurt cross-validation performance (data not shown).
While previous studies hypothesized that more than two cascade equations were required to
model adaptation on multiple timescales (38), here we found that the type of adaptation and not
the number of cascade equations to be more important for modeling OFF sensory neurons.
We also compared the timescales of ON and OFF temporal filters. For both ON cells and
OFF cells, primary filters are fast and follow the fluctuation of stimuli, while the adapting filters
reflect the slower timescale of adaptation. While ON and OFF cells displayed similar timescales in
their primary filters, OFF cells have faster adapting filters (higher τ and lower time-to-half-peak)
than ON cells (Fig. 2D,E). Thus, the OFF cell rebound was fast relative to ON cell habituation.
Moreover, the OFF cell rebound was weak compared to ON cell habituation. In the best-
performing OFF cell model, the adapting filter coefficient was typically smaller in magnitude than
the primary filter coefficient (Fig. 2E). Thus, the OFF cell adapting filter does little to temper
stimulus inhibition of OFF cells. Instead, it seems to be designed to produce fast rebounds to
bacteria removal.
Adaptation appears to have different goals in ON and OFF cells. ON cells obey a fairly
straightforward perfect adaptation law that can be explained by calcium depletion. OFF cells, on
the other hand, exhibit an imperfect, fast rebound strategy. This latter strategy will not efficiently
encode stimulus across large concentration scales. It will, however, maintain a higher dynamic
range for constrained concentrations scales. We surmise that ON sensory cells may be designed
to work across larger concentration scales, engaging specific OFF cells for specific concentration
ranges. Overall, we show that ON and OFF sensory neurons have different adaptation kinetics to
rapidly fluctuating bacterial stimulus sequences.
To determine how different phases of stimulus presentation affects motor neurons, we
next built decoding models to predict stimulus state from motor/command cluster activity (see
Materials and Methods: Decoding model). In bacteria↔buffer stimulus sequences, bacteria
pulses are effectuated by directing bacteria flow from channel 1 over the *C. elegans* nose (Fig.
2F), while buffer pulses result when buffer from channel 2 flows over the nose and displaces the
bacterial flow (Fig. 2G). Since *C. elegans* is positioned asymmetrically in the microfluidic chip
relative to the two channels (channel 1 is slightly closer to the nose), we also analyzed animals
presented with buffer↔buffer stimulus sequences to control for mechanosensory responses to
differences in flow properties between the two channels (Fig. 2H).
Unlike sensory neuron clusters, shifts in motor/command neuron cluster activity do not
reliably coincide with stimulus transitions (Fig. 1H–K). Therefore, we used multinomial logistic
regression (MLR) to predict the probabilities of a particular stimulus state given motor neuron
cluster activity inputs. MLR (39) is a robust classification model that, when combined with class
balancing, has a very natural null model: prediction from worm identity only. Here, linear
predictions about stimulus state are generated by linearly combining a set of weights with
explanatory variables of a given observation:
264

265
$$\mathrm{P}(\mathrm{S}_i | X(t) = softmax(\sum_{n=1}^{k} \beta_{i,k} X_k(t))$$

266
where　　is the stimulus class　at time window　. The　-dimensional　captures motor
neuron cluster activity at time window　along with worm identity information.　　is made up of
the coefficients for stimulus state　, which are regularized using an L1 norm and learned using
gradient ascents. 16-second time windows of motor/command neuron cluster activity data were
divided into thirds (5.33 seconds each). To predict the stimulus state in the middle subwindow (*t-*
2.67s to *t+2.67s*, with *t* as the halfway point of the prediction window), motor/command neuron
cluster activity data from the first subwindow (*t-8s* to *t-2.67s*) and the last subwindow (*t+2.67s* to
t+8s) were used for the decoding task (see Materials and Methods: Decoding model).
We used this model to decode neural activity inputs from AVA, RME, SMDD, and SMDV
clusters (Fig. 2I). Worm identity was also considered to capture variability across animals. The
decoding model predicted four stimulus states for bacteria↔buffer stimulus sequences:

prolonged bacteria, prolonged buffer, bacteria-to-buffer transition, and buffer-to-bacteria transition
(Fig. 2I). Importantly, initial onsets and removals of bacteria are included in transition states, but
not in prolonged states. For buffer↔buffer stimulus sequences, corresponding states based on
channel activation were predicted. In predicting stimulus states associated with bacteria↔buffer
stimulus sequences, decoding from the activity of the RME and AVA cluster pair improved
performance over decoding from identity alone in both non-NeuroPAL–GCaMP5K (99.3% of
bootstraps, Fig. 2J) and NeuroPAL–GCaMP6s (99.4% of bootstraps, Fig. S4A) strains. In
contrast, decoding from the activity of the SMDD and SMDV cluster pair did not perform better
than the null model (Fig. S4B). Moreover, RME and AVA cluster activity could also be used to
predict stimulus state in buffer↔buffer stimulus sequences, outperforming prediction from worm
identity alone (Fig. S4C).
The decoding model produced linear temporal filters that described how stimulus states
contributed to motor/command neuron cluster activity. Temporal filters for buffer↔buffer stimulus
sequences revealed that flow from both buffer channels had similar effects on motor/command
cluster activity. Prolonged buffer from either channel was associated with similar transient
increases in both AVA and RME cluster activity (Fig. 2K,L). Both types of buffer-to-buffer
transitions (channel 1 → channel 2, channel 2 → channel 1) reduced AVA cluster activity while
RME cluster activity remained near baseline (Fig. 2M,N). In contrast, temporal filters for
bacteria↔buffer stimulus sequences indicated that bacteria and buffer differentially drove AVA
and RME cluster activity. Prolonged bacteria was associated with sustained increase in RME
cluster activity and strong decrease in AVA cluster activity (Fig. 2O), while prolonged buffer was
associated with sustained inhibition of RME cluster activity and low AVA cluster activity (Fig. 2P).
The bacteria-to-buffer transition induced a slow increase in RME cluster activity, while AVA
cluster activity remained near baseline (Fig. 2Q). The temporal filter predicting bacteria-to-buffer
transition from RME cluster activity (Fig. 2Q) resembled a diminished version of the temporal filter
predicting prolonged bacteria (Fig. 2O), suggesting that bacteria removal did not immediately
alter RME activity. The converse buffer-to-bacteria transition is associated with rapid peak in AVA
cluster activity and near-baseline RME cluster activity (Fig. 2R). The temporal filter for predicting
buffer-to-bacteria transition from AVA cluster activity (Fig. 2R) is higher in magnitude than the
temporal filter predicting prolonged buffer from AVA cluster activity (Fig. 2P), suggesting that
bacteria onset has an immediate effect on the AVA cluster. Bacteria↔buffer decoding models
were also remarkably similar across non-NeuroPAL–GCaMP5K and NeuroPAL–GCaMP6s
strains (Fig. S4D–G). Altogether, these results suggest that bacteria presentation biases the
AVA–RME cluster pair towards RME cluster activation (associated with forward locomotion),
while bacteria removal biases the cluster pair towards AVA cluster activation (associated with
reverse locomotion). Furthermore, motor/command responses to buffer differ depending on
whether the overall sensory context also includes bacterial stimuli.
315
Identification of interpretable network states that vary in sensory gating properties
Based on our previous decoding results that AVA and RME motor/command clusters are
influenced by sensory input, we next investigated how global network activity and sensory input
drive AVA and RME cluster activity under different network states. We created a hierarchical
model that used a soft decision tree (SDT) gating model (40) to identify relevant network states,
and then we fitted MLR forecasting submodels for each network state to predict AVA and RME
cluster activity (see Materials and Methods: SDT–MLR model). This combination of models,
which we refer to as the SDT–MLR model, overcomes the limitation of using a single linear model
to describe motor neuron cluster activity. For instance, members of the AVA cluster are bistable:
they have upper and lower stable states (boundedness). Also, activity in these cells appears to
have momentum: when one of these cells begins a transition between its stable states, it will tend
to complete that transition. A single, linear model is unable to describe both momentum and
boundedness. An appropriate model should learn the positive correlation between future AVA
cluster rise and past AVA cluster increases to capture momentum. At the same time, this positive
correlation should weaken and become negative as AVA nears its upper bound as these cells do
not rise above that limit. Here, we employed multiple MLR models to capture nonlinear dynamics,

including features like momentum and boundedness. We then used an SDT gating model to route
in recent network activity (AVA, RME, SMDD, SMDV clusters) to different network states. Each of
these network states is associated with a different MLR model. In so doing, the SDT parceled the
space of network trajectories into subspaces in which network evolution can be approximated by
linear, probabilistic models. This parcellation and linearization strategy is similar to what has been
previously used (41). In our SDT–MLR modeling, there were $M \times N$ MLR submodels, where M is
the number of models being compared and N is the number of states parceled by the SDT. If the
stimulus effect depends on network history, then we expect that the stimulus filters will differ
across MLR submodels. As an end-to-end interpretable distillation of a neural network, the SDT is
capable of learning complex, non-linear features. Moreover, since both the SDT and MLR
submodels are differentiable, they can be simultaneously fit using gradient-based optimization
methods.
We started by comparing forecasting models that predict AVA and RME cluster activity
from recent network and sensory neuron activity, with models that predict from network history
alone. This allowed us to assess whether sensory input from bacteria↔buffer stimuli were more
relevant in some network states compared to others. Using recent network history inputs from
AVA, RME, SMDD, and SMDV clusters, the full SDT-MLR model was fitted to a hyperparameter
set for predicting RME and AVA calcium change (rise and fall). This model generated a tree with
three levels, consisting of top level filters, along with right and left subtree filters (Fig. S5A), and
four network states that produced the best results (Fig. 3A, S5B,C, see Materials and Methods:
SDT–MLR model). State 1 and 4 were transient states, with state 1 associated with AVA cluster
peaks, and state 4 associated with RME cluster peaks (Fig. 3A). In contrast, state 2 and 3 were
persistent states (Fig. 3A). State 2 was characterized by high AVA, low RME, low SMDV, and
high SMDD cluster activities, while state 3 exhibited the opposite (Fig. 3A). After network states
were identified, the SDT was then frozen, and the MLR submodels were fitted to the rest of the
data in an out-of-bootstrap cross-validation strategy to assess feature variability (42).
In the out-of-bootstrap cross-validation, inclusion of ON and OFF sensory neuron activity
improved overall model performance (summed across states) in >95% of bootstraps. However,
when considering model performance for individual network states, inclusion of sensory activity
was only useful in states 3 and 4, but provided little predictive value in states 1 and 2 (Fig. 3B),
despite the variance in stimulus sequence being similar across all states. Both sensory-
responsive states (states 3 and 4) are characterized by low AVA and high RME cluster activities,
both of which are associated with forward locomotion (Fig. 3B). Conversely, the sensory-
unresponsive states (states 1 and 2) display the opposite activity trends with high AVA and low
RME cluster activities, which are associated with reverse locomotion (Fig. 3B). This suggests that
sensory input is gated more heavily during reverse locomotion than during forward locomotion,
and that forward locomotion is a sensory-responsive behavior.
For each state, the SDT–MLR model generated linear temporal filters that predicted how
recent network history from AVA, RME, SMDV, and SMDD clusters affected the probability of
AVA and RME activity trends. In general, temporal filters were similar across predicted neural
activity associated with forward locomotion (AVA fall, RME rise; Fig. 3C, middle two rows), as well
as across activity associated with reverse locomotion (AVA rise, RME fall; Fig. 3C, top and
bottom rows). Additionally, temporal filters were similar across the sensory-unresponsive states
(state 1 and 2), except that recent SMDV history had a diminished effect on forward-associated
activity in state 2. Linear filters across the sensory-responsive states (state 3 and 4) resembled
each other, except for how network history drove AVA fall in state 4. Notably, RME cluster history
drove AVA and RME cluster activity in general in sensory-unresponsive states, but had a neutral
or suppressive effect in sensory-responsive states. Some state-invariant trends included the
suppressive effect of SMDD history on AVA rise, as well as the suppressive effect of AVA on
RME rise (Fig. 3C). Here, we describe some of the general trends of how network history
influenced AVA and RME cluster activity in a state-dependent fashion, but Fig. 3C can be readily
interpreted to understand in detail how each motor/command cluster affected AVA and RME
cluster activity in each state.

Next, we looked at how sensory input influences particular AVA and RME cluster activity
trends under different network states. As expected from Figure 3B, temporal filters predicting AVA
and RME cluster activity from ON and OFF sensory activity were generally flat in states 1 and 2,
indicating that sensory information was broadly suppressed from motor/command activity in these
states (Fig. 3D). In contrast, states 3 and 4 exhibited more temporal filters in which sensory
activity either increased or decreased the probability of AVA and RME activity trends (Fig. 3D).
The largest effects of sensory input are on reverse-associated motor/command activity (AVA rise,
RME fall), which is suppressed by ON activity and elevated by OFF activity (Fig. 3D). Within
reverse-associated activity, OFF activity has a greater influence on AVA rise, while ON activity
has a greater influence on RME fall (Fig. 3D). Additionally, ON sensory input drives RME rise in
state 3 (Fig. 3D), suggesting that forward locomotion that characterizes state 3 can be maintained
with ON activation of RME and OFF suppression of AVA. Overall, we show that SDT–MLR
models can be used to identify relevant network states, characterize how those states are
generated by network history, and delineate the state-dependent effects of stimuli on
motor/command neuron activity.
400
Effect of stimulus timescales and sensory context on AVA and RME clusters
To further deconstruct sensory influences on AVA and RME cluster activity, we divided
bacteria↔buffer stimulus sequences into stimulus patterns with either low- or high-variance pulse
protocols, and fitted separate SDT–MLR models using corresponding neural activity (Fig. S6, S7).
The main difference between low- and high-variance protocols is that alternating stimulus blocks
consist of a single long pulse in the former, and composed of multiple short pulses in the latter
(see Materials and Methods: Division of stimulus sequences). In general, high-variance pulses
generally had more effect on AVA and RME cluster activity than low-variance pulses (Fig. 4A). To
compare the overall effect of each network state on sensory gating, we summed the absolute
values of magnitudes from all temporal filters within a state (Fig. 4A, last row). While high- and
low-variance stimulus pulses had similar overall effects in state 4, high-variance pulses had more
influence than low-variance pulses in state 3 (Fig. 4A, last row). Particularly in state 4, low-
variance pulses can result in uneven influence from ON and OFF neurons, such as greater ON
influence on RME fall and greater OFF influence on AVA rise (Fig. 4A). While state 1 and 2 were
initially deemed to be broadly unresponsive to sensory input (Fig. 3B), enriching the model with a
subset of pulse lengths revealed some sensory influence in state that was previously masked
when all timescales were considered. For example, in state 1, high-variance-pulse ON activity
promoted AVA fall, while high-variance-pulse OFF activity suppressed AVA fall (Fig. 4A). These
opposite effects likely canceled each other out when all stimulus pulse lengths were considered,
thereby resulting in a sensory-unresponsive model prediction (Fig. 3D). Furthermore, segregation
of stimuli by low- or high-variance pulses revealed more sensory influence on forward locomotion
in states 3 and 4 (Fig. 4A) that was previously undetected in the all-pulse SDT–MLR model (Fig.
3D). Thus, the SDT–MLR model can be used to also detect feature-specific effects of stimuli on
motor/command neuron activity.
We next explored the ability of SDT–MLR models to differentiate sensory contexts by
comparing buffer↔buffer and bacteria↔buffer stimulus sequences. Since high-variance pulses
were shown to have greater overall effect (Fig. 4A), we compared only high-variance pulses from
buffer↔buffer and bacteria↔buffer stimulus sequences. In the SDT–MLR model for
buffer↔buffer stimulus sequences (Fig. S8), inclusion of buffer↔buffer sensory activity improved
overall performance only for state 4 (Fig. S8A). Compared to bacteria↔buffer sensory input,
temporal filters predicting AVA and RME cluster activity from buffer↔buffer sensory input typically
displayed slower timescales (Fig. S8C). Based on the sum of absolute magnitudes of all temporal
filters within a state, buffer↔buffer stimuli generally had more influence than bacteria↔buffer
stimuli in states 1 and 2, and less influence in state 3 (Fig. 4B). While the state-agnostic decoding
model found no difference between channel 1 and channel 2 buffer flows in how they drove AVA
and RME cluster activity (Fig. 2K–L), the SDT–MLR model found that the channel 1 buffer had a
greater effect on AVA and RME cluster activity than the channel 2 buffer in states 2, 3, and 4 (Fig.
4B, gray). This bias may be due to channel 1 being closer to the *C. elegans* nose, as previously

surmised. However, this bias disappeared when bacteria was included in the sensory context
(Fig. 4B, cyan). Therefore, while channel 2 is chemically the same for both buffer↔buffer and
bacteria↔buffer stimulus sequences, the effect that the channel 2 buffer had on AVA and RME
cluster activity changed depending on whether the greater sensory context involves switching
between chemically identical buffers, or switching between bacterial stimuli and buffer.
444
445
446 **Discussion**
447
Here, we demonstrate that the SDT–MLR model can identify interpretable global network states
that underlie differential gating of sensory input at the motor and command neuron level. First, we
used state-agnostic encoding and decoding models to confirm that bacteria onset and removal
differ in how they drive activity of sensory neurons, AVA, and RME clusters. Using the SDT–MLR
model, we identified two sensory-unresponsive network states (states 1 and 2) and two sensory-
responsive network states (states 3 and 4) in the bacteria↔buffer sensory context. For each
network state, we explicitly characterized how the history of each of the four functionally defined
motor/command neural populations (AVA, RME, SMDD, SMDV) drive AVA and RME cluster
activity to produce each network state. Finally, we used the SDT–MLR model to identify how
sensory input in general, as well as stimulus features and sensory context in particular, influence
AVA and RME cluster activity.
The encoding model showed that sensory neural responses to the addition of bacterial
stimuli (ON) and its removal (OFF) can be modeled by two linear ordinary differential equations
using one fast and one slow filter. Linearity in sensory neurons is observed in both vertebrates
and invertebrate photoreceptors (43), rat trigeminal neurons (44), and primate vestibular neurons
(45), implying that this might be a common phenomenon. We also show that ON sensory neurons
perfectly adapt to the bacterial stimulus, consistent with previous studies (38), while OFF neurons
adapt imperfectly. Given that our analysis included all sensory neurons responding to bacterial
stimuli, we suggest that this might be a general principle of *C. elegans* sensory neurons. Similar
differences in adaptation in ON and OFF neurons are also observed in single-cell
electrophysiological recordings from vertebrate photoreceptors and olfactory sensory neurons
(46). Moreover, these results also provide some hints about the encoding strategies of these two
sensory-neuron classes. We suggest that ON neurons encode stimulus over a larger dynamic
range compared to OFF neurons. Consistently, we have previously shown that AWA sensory
neurons (ON) have a larger dynamic range compared to AWC sensory neurons (OFF) in
detecting benzaldehyde (47). Moreover, studies in the vertebrate retina have shown that the
dynamic range of the ON pathway is much greater than that of the OFF pathway, likely due to a
selective effect of pre-synaptic inhibition on the ON, but not OFF, bipolar cells (48), confirming the
validity of our hypothesis.
The decoding model informed which motor/command neural populations were targets of
the sensory input, and also served as a state-agnostic model to compare with the state-
dependent SDT–MLR model. Both models were used to assess how sensory input affects AVA
and RME cluster activity. While the decoding model found no difference in effect between either
of the buffer channels in buffer↔buffer stimulus sequences, the SDT–MLR model revealed that
buffer↔buffer sensory input does indeed have a channel bias that is more pronounced in some
states than others. This suggests that the SDT–MLR model is more sensitive to sensory effect
compared to the decoding model. This sensitivity is further amplified by subdividing sensory
effects by state, channel, sensory neuron classification (ON/OFF), prediction class (AVA rise,
AVA fall, RME rise, RME fall), and stimulus features (high-variance pulses, low-variance pulses).
This granular approach allowed us to identify specific sensory effects that were obscured in more
general analyses. We found that high-variance and low-variance pulses had similar influences in
one state, but sensory input from low-variance pulses were gated more heavily in another state.
Additionally, we revealed that state-dependent gating of buffer stimulus is dependent on whether
the greater sensory context involves switching between bacteria and buffer, or between buffer
and buffer.

The SDT–MLR model differs in both goal and interpretability from recent studies that
described network states in *C. elegans* whole-brain activity (25, 27, 37, 49). In these studies,
whole-brain activity is analyzed with the purpose of understanding the temporal dynamics of
neuronal populations in terms of how the network state evolves over time. Often, probabilities of
network state transitions are related to corresponding stimulus or behavioral transitions. A
dimensionality reduction technique called principal component analysis (PCA) (50) is used to
quantify brain-wide correlations that reflect signals shared by clusters of neurons. These shared
signals are referred to as temporal principal components (PCs). A key operation of PCA-based
analyses is to transform whole-brain activity to new axes defined by the top PCs that explain the
most variance in data. However, the meaningful linear axes that originally described the data are
lost in the process. While PCA-based analyses are useful for distinguishing between network
states and their transition between each other, these networks states and transitions are
described in terms of PCs, which can be difficult to interpret. Nonetheless, PC weights can still be
used to identify relevant neurons and activity trends associated with each state (25, 27, 37, 49).
In our study, we are less concerned with the probabilities of transitioning between states, and
more interested in the within-state conditional probabilities of individual neuronal populations
interacting with each other. Instead of using PCA, our model preserves interpretability by using
soft decision trees, such that network states and network interactions are always described by the
identities of neuronal populations and their corresponding activity patterns. Rather than
supplanting PCA-based analyses, our SDT–MLR model serves as a complementary method for
focusing on the network interactions within individual network states rather than the temporal
dynamics that connect those network states.
Interestingly, the sensory-responsive states identified by the SDT–MLR are characterized
by neural activity that has been previously shown to be associated with forward locomotion, while
the sensory-unresponsive states are characterized by neural activity associated with reverse
locomotion. This is consistent with previous reports showing that inhibition of sensory input
occurs at particular phases of the locomotory cycle (11, 12). Inhibition of sensory input during
movement may serve to distinguish between external stimuli and self-generated stimuli (51),
which is accomplished by integrating sensory inputs with motor inputs (corollary discharge) (52).
Since typical *C. elegans* locomotion consists primarily of forward locomotion punctuated by
transient reversals (53), one possibility is that sensation of stimuli is suppressed during reversals
to temporarily pause processing of stimulus flows until a stable locomotion state is restored. It is
important to note that our study was conducted with *C. elegans* trapped in an immobilized
position in a microfluidic chip, and therefore behavioral associations were inferred purely from
motor and command neuron activity. Moreover, a recent study showed that the set of neurons
correlated with AVA differs depending on whether *C. elegans* is immobilized or freely moving
(49). Thus, our association of AVA and RME clusters activity with forward and reverse locomotion
is tentative and should be confirmed in freely moving animals. However, while imaging freely
moving *C. elegans* would provide rich behavioral information that can be added to the SDT–MLR
model, complex and precise stimuli presentation can be difficult to achieve when the stimulus
target is mobile. There are some efforts to study sensation in freely moving animals (22, 54), and
a reasonable balance of behavioral and sensory information richness may be achieved with a
microfluidic chip that allows semi-restricted locomotion and somewhat fast waves of liquid stimuli
(55).
Overall, we present an approach for understanding how sensory information filters
through whole-brain network interactions to affect downstream motor and command neurons in a
state-dependent manner. Currently, there is an epistemological bias towards identifying network
states that correspond with a particular stimulus or motor state. In contrast, there has been less
focus on network states that are defined by altered network interactions. Our computational
approaches provide a method for investigating network mechanisms at the level of pairwise
interactions between neuronal populations. While our study only looked at the network
mechanisms underlying sensorimotor integration, this model can be leveraged to also understand
how network inputs are integrated at any network level. More broadly, we suggest that our
approach of combining soft decision trees with multinomial logistic regression can be used to

identify relationships, not only in neural networks, but also in cellular signaling pathways,
transcription factor networks, and between other complex biological or physical entities.
549
550   **Materials and Methods**
551
Whole-brain imaging
We used two transgenic strains that expressed GCaMP. The primary strain (ZIM294) expressed
GCaMP5K in the nuclei of all neurons (mzmEx199 [*Punc-31::NLSGCaMP5K; Punc-122::GFP*]).
To identify neurons associated with activity patterns observed in ZIM294, we used a strain
(OH15500) that expressed GCaMP6s and NeuroPAL (*otIs669[NeuroPAL];otIs672[Panneuronal*
GCaMP6s]). Cells were identified according to the map described by Yemini and colleagues (24).
We monitored changes in GCaMP fluorescence using a Zeiss LSM 880 with Airyscan. Acquisition
was done in 2 micron z-steps. In 'Fast' mode, the Airyscan images the entire head of the adult
worm at about 1.5 volumes per second. Worms were typically imaged for approximately ten
minutes. We then used piecewise rigid registration to remove motion artefacts (56) and non-
negative matrix factorization to isolate individual neurons and extract their fluorescence values
(33). Out of a total 189 neurons in the head, our approach identified 50-100 neurons per animal.
564
Stimulus delivery
Day 1 adult animals were washed in M9 and loading into in a microfluidic device that trapped the
worm body while exposing only the nose to stimulus flows (28). Animals were also treated with
1.5 mMol of the paralytic tetramisole hydrochloride to suppress most perceivable worm
movement. The movement of untreated worms proved too difficult to motion correct. We
delivered precise patterns of fluctuating bacteria and M9 buffer liquid flows using a custom
designed Arduino device to send pulses to a valve controller. The bacteria solution was prepared
as a 1:1 resuspension of a bacterial culture (OD_{600} = 0.4) in M9 buffer as previously described
(29). The controller determines whether bacteria or buffer is routed to the nose of the trapped
worm or away from the worm. Worms were exposed to binary patterns of bacteria and buffer. A
number of different stimulus protocols are used in this study. In the base protocol, the trial is
divided up into pulse blocks of ~15 seconds. The pattern is constructed using transition
probabilities: p(switch on | off) = 0.2 and p(switch off | on) = 0.4. In the faster protocols, the same
switch probabilities are used but the pulse blocks have length ~1.5 seconds. The patterned
protocols are effectively the same as the base protocol. The only difference is that their 'stimulus
on blocks' are composed of multiple pulses.
581
GCaMP filters
The GCaMP filter *g(x)* is modeled as a difference of exponentials with parameters matching those
of Chen and colleagues (32). This procedure is complicated by the volumetric nature of the
imaging data. Consider two sensory neurons with identical calcium dynamics; they respond to
stimulus with the same timescale. Neuron A is in imaging slice 0, while neuron B is in slice 7.
These two neurons will have the same calcium timings relative to the stimulus. However, Neuron
B will appear to have faster response kinetics since it is acquired over half a second later (relative
to stimulus onset/removal) compared to neuron A. Thus, the slice in which the neuron appears
needs to be considered in the creation of its GCaMP filter:

$$g_f(t) = \int_{t_1}^{t_2} g(x)dx$$

$$t_1 = t + \frac{z}{n_z}$$

$$t_2 = t + \frac{z+1}{n_z}$$

591
592   where *g(x)* refers to the difference of exponentials and n_z refers to the number of z slices. The
593   normalized *g(t)* filter is applied via a linear convolution to transition between calcium and GCaMP
594   dynamics.

595
ON and OFF sensory neuron classification
596
597 Sensory neurons were classified as either ON or OFF for neuronal activity collected during
598 bacteria↔buffer stimulus sequences. The change in normalized fluorescence over a series of ≥
599 10 stimulus pulses (all trials have at least a few of these). Both the first 10 volumes into a bacteria
600 pulse and the first 10 volumes into a buffer pulse (following bacterial removal) were considered.
601 The following metric was then calculated:

$$rank = \left(\sum_i^P (f(t_i + 10) - f(t_i)) \right) - \left(\sum_j^Q (f(t_j + 10) - f(t_j)) \right)$$

602
where f is normalized fluorescence, t is pulse onset, P is the number of bacteria pulses, and Q is
the number of buffer pulses. The cells with the highest ranks considered as potential ON cells,
and the cells with the lowest ranks were considered as potential OFF cells. Rank cutoffs were
selected manually for each trial. ON cells were categorized as those neurons that obviously and
immediately increased activity upon bacteria onsets, and immediately decreased upon bacteria
removals (Fig. 1I). In contrast, OFF cells were classified as neurons that decreased activity upon
bacteria onsets and increased upon bacteria removals. For both ON and OFF cells, baseline (low
variance) activity occurred when the stimulus of interest was absent (during buffer pulses). This
distinguishes OFF-bacteria sensory neurons from hypothetical ON-buffer sensory neurons, for
which baseline activity would occur during bacteria pulses.
613
614 **Encoding model**
615 The encoding model predicted ON and OFF sensory neuron activity from stimulus features. The
616 core primary sensory model consists of three parts: (1) a set of cascade basis functions, (2)
617 coefficients for the basis functions, and (3) a GCaMP transformation.
618

$$\frac{dx_i}{dt} = \tau \cdot ([input] - x_i(t))$$
$$x_i(t) = \sum a_i x_i(t)$$
$$g(t) = (x^p \circledast f_g)(t)$$

619
620
where   is the inferred calcium level of the cell and is calculated as the sum of the temporal
filters (ODE model solutions),   is the GCaMP filter, and   is a positive value required for the
GCaMP transformation. The fitting procedure learns values for the time constant τ and the basis
coefficients   . This model uses the solutions to these ordinary differential equations   as
basis functions. For a given model instantiation, each equation is assigned a value for τ. After the
ordinary differential equations are solved analytically, the model can be linearized, allowing for
robust estimates of the a_i coefficients (see Methods). We then used a random search strategy to
obtain estimates of τ. Moreover, by toggling the coefficient constraint, we can test whether perfect
adaptation is necessary to predict sensory neuron activity. This constraint yields perfect
adaptation as all the ordinary differential equation basis functions saturate at the same value
(   ).
We used a random search strategy to find the correct set of basis functions. The strategy
chooses N basis time constants for an N cascade model and an initial estimate of p power. It
solves the basis equations analytically for a given stimulus pattern and produces an initial
estimate for the coefficients by fitting to a linearized approximation of the neuron's calcium trace.
In this approximation, the raw sensory neuron fluorescence trace is deconvolved using the
Richardson-Lucy method (57) and taken to the *(1/p)* power. The initial estimate for the
coefficients is the solution to the resulting linear regression equation. Finally, this system fits the
full model (free variables consist of basis coefficients and *p* power) is fit to the raw fluorescence
trace using gradient descent. We then repeat this process for a large number of random searches
to define the basis functions.

Decoding model

The decoding model predicted stimulus states activity from stimulus features. For both the decoding and prediction analyses, we split the data into contiguous blocks of ~10 second (16 volume) duration. Within each block, subwindows were created in a rolling fashion. For instance, for an N volume prediction window, there are 16-N legal, overlapping subwindows within each block. It should be noted that there is no overlap between prediction windows of adjacent blocks. Test/train sampling is done at the block level, guaranteeing no train/test prediction overlap. This system captures sharp transitions while measuring targets over multiple time bins, thereby limiting noise in the targets. We chose a length of 16 volumes because this captures the entirety of motor neuron event initiation (for all motor neuron classes). This causes bootstrap sampling to be performed at the event level, stopping a small, handful of events from dominating model outcomes. We did not test other blockstrap sizes.

We used Gaussian basis functions for both decoding and prediction tasks. This involves filtering network history and stimulus data through these basis functions before being fed into decoding and prediction models. We chose specific gaussians (parametrized by mean and variance) by hand-tuning model performance on the hyperparameter set.

Given the choice of 16 volume blocks and 8 volume time windows, there are 16-8=8 prediction windows within each time block. Unlike the prediction task, the decoding task centers the prediction windows relative to the input data. For instance, in one decoding task, RME/AVA cell cluster data from t+4-8 to t+4+8 volumes is used to predict whether the stimulus is on/off/altered from t to t+8 volumes. The unit of 8 volumes was chosen because many of the pulse protocols use 10 volume pulses as a base unit. Thus, using 8 volumes guarantees a fair number of samples of the on stimulus class. We did not test other volume lengths.

In the decoding analyses, we predict five classes of stimulus patterns from network activity (58). Prediction from worm identity alone serves as the null model for all decoding analyses. In order to ensure best performance for the null model, we balanced the classes within each worm. For each worm, if there are N occurrences of class A in the training set, there are N occurrences of class A in the test set. We ensured this by randomly removing prediction windows.

SDT–MLR model

We used different L1 norms for each of the multinomial logistic regression input classes: AVA/RME/SMDV/SMDD terms, ON/OFF cell terms, and worm identity terms. The worm identity terms essentially make the forecasting models into a random-intercept model (59). Initial experiments found no benefit in random-intercept style models. The number of network states and number/shape of gaussian filters were additional hyperparameters. A combination of hand-turning and grid search were used on the hyperparameter set to find good regularizers. This hyperparameter set was also used for the soft decision tree, which could also be thought of as an additional hyperparameter in this study. These hyperparameters were frozen on the out-of-bootstrap analysis (42).

In the forecasting task, we predicted changes in GCaMP fluorescence from time t to t+T1 using network history from t-T2 to t and sensory neuron activity from t-T2+T1 to t+T1. We used a length of twenty-four volumes for T2 for all prediction analyses. We treated the length of T1 as an additional hyperparameter. As in the decoding analysis, these prediction windows are contained inside larger data blocks. We performed train/test sampling at the block level. We chose sixteen seconds (24 volumes) for T1 in all models, as 16 seconds is sufficient to capture the majority of command neuron events.

RME and AVA activation and inactivation is clearly probabilistic. We used a multinomial logistic regression (MLR) as the base model for motor-neuron activity prediction. In order to use MLR, we discretized GCaMP fluorescence activity in every prediction window. This is done by subtracting the average GCaMP fluorescence over the prediction window by the GCaMP fluorescence level at the beginning of that window.

RME and AVA neurons have non-linear calcium dynamics. Two of these features are boundedness and momentum. Members of the AVA cluster exhibit boundedness: their activity is

limited to a range between their upper and lower stable states. Also, activity in these neurons
appears to have momentum: when one of these neurons begins a transition between the stable
states, it will tend to complete that transition. A single, linear model is unable to describe both
momentum and boundedness. It must learn the positive correlation between a future AVA rise
and past AVA activity to capture momentum. However, this positive correlation should weaken
and become negative as AVA nears its upper bound, since AVA activity does not rise above this
bounded limit. Thus, a gating model is required to capture the change in this positive correlation.
Here, we employ multinomial logistic regression (MLR) models to capture nonlinear
dynamics, allowing us to capture features like momentum and boundedness. A gating model is
used to divide the space of network histories into subspaces. Each of these different subspaces
is associated with a different MLR model. This approach with two submodels can simultaneously
model both momentum and boundedness in AVA. In one theoretical solution, one MLR submodel
is only active when the AVA cell cluster is near its lower bound, while another is active when AVA
near its upper bound. The first submodel learns a positive correlation between a past AVA
increase in activity and future AVA rise (momentum), while the second learns a weak positive or
negative correlation (boundedness).
We use a hybrid model combing soft decision trees and multinomial logistic regression
(SDT–MLR) method to divide the network trajectory space into different subspaces. We assumed
that GCaMP fluorescence changes in each of these subspaces can be well-described by an
MLR. A soft decision tree is a form of oblique decision tree that is end-to-end differentiable (40).
Each branch of the soft decision tree is a different logistic regression model on the same input
vector, which outputs a left vs right probability. These left and right probabilities are multiplied by
the predicted class probabilities of the corresponding left and right subtrees. In this manner, soft
decision trees are essentially hierarchical filters that can be learned through gradient descent.
The outputs of the soft decision tree weigh the different MLR models. For instance, a soft
decision tree with depth 2 and width 2 will have 4 output states. Each of these output states are
associated with a different MLR. All of these MLRs are trained against the entire dataset.
However, the data points are weighted by the particular soft decision tree output leaf. Hence,
different MLR models will focus on different subsets of the data.
We found that averaging predictions across SDT–MLR models improves cross-validation
performance. This averaging is done at two levels. First, within a SDT–MLR model, the
predictions of each MLR are weighted by the soft decision tree and averaged. Second, these
averaged predictions are further averaged across several SDT–MLR models. Here, twenty-five
SDT–MLR models were fit separately to the hyperparameter set. The best hybrid model was
chosen for analysis on the cross-validation set. For training on the cross-validation set, the Soft
Trees were frozen; only the MLRs were trained. Hence, training is convex (weighted multinomial
logistic regression) on the cross-validation set. Freezing the soft decision during cross-validation
allows us to easily align model data across bootstraps. This, in turn, gives us information on the
variance of different features of the MLR models.
In our exploration of the hyperparameter set, we found that SDT–MLR models tend to
converge on bad solutions if not regularized. These bad solutions are characterized by poor
training and test set performance as well as state imbalance. The Soft Tree assigns most data
points to one of its submodels, while its other MLR submodels are trained on very small subsets
of the data resulting in state imbalance. We solved this issue by maximizing entropy regularizer
H(X):

$$V_k(X) = \frac{1}{N} \sum_{i}^{N} S_k(X_i)$$

$$H(X) = \sum_{k} V_k(X) log \left(\frac{1}{V_k(X)} \right)$$

741
742 where X_i is the i^{th} input data point in the minibatch. S_k is the probability assigned to the kth
743 state/submodel by the soft decision tree. Thus, V_k is the average probability of state k across the

744 minibatch. Therefore, *H(X)* is high when all states are equally represented in the minibatch. It
745 should be noted that this regularizer does not directly penalize high state probabilities.

Division of stimulus sequences

To compare the effect of short versus long stimulus pulses in the SDT–MLR model, complete
stimulus sequences were divided into low-variance (long pulses) and high-variance (short pulses)
stimulus patterns. Low-variance stimulus patterns were constructed by dividing the trial into pulse
blocks of ~15 seconds, with the transition probabilities p(switch on | off) = 0.2 and p(switch off |
on) = 0.4. High-variance stimulus patterns were also constructed by dividing the trial into bacteria
or buffer blocks with the same transition probabilities as for low-variance stimulus patterns. The
difference between low- and high- variance stimulus patterns is that the bacteria block consists of
a series of sub-pulses rather than a single constant pulse. These sub-pulses are ~3 seconds
bacteria and ~3 seconds buffer.

756
Acknowledgments

758
We thank Manuel Zimmer, Oliver Hobert and CGC (Caenorhabditis Genetics Center) for worm
strains and Uri Manor, Tong Zhang and the Waitt Advanced Biophotonics Center for advice and
assistance with our imaging experiments. We also thank Saket Navlakha, Ryan Rowekamp,
Javier How, Molly Matty, Jessica Haley, Michael Rieger, and Eric Edsinger for discussions and
comments on the manuscript. This work was supposed by grants from the National Institutes of
Health NIH 5R01MH096881 (S.H.C); 5T32DK007328-35, 5T32DK007328-37, 5T32MH015174-
37 and 5T32MH015174-38 (E.Y). U19NS112959, P30AG068635, AHA-Allen Initiative in Brain
Health and Cognitive Impairment award 19PABH134610000 (T.O.S.). Z.T.C. was supported by a
fellowship from the Dan and Marina Lewis Foundation. K.T.Q. is supported by a Postdoctoral
fellowship from the Paul F. Glenn Foundation. CGC is funded by the NIH Office of Research
Infrastructure Programs (P40 OD010440).

770
References

1. J. M. Beck, W. J. Ma, X. Pitkow, P. E. Latham, A. Pouget, Not noisy, just wrong: the role
of suboptimal inference in behavioral variability. *Neuron* **74**, 30-39 (2012).
2. L. C. Osborne, S. G. Lisberger, W. Bialek, A sensory source for moto variation. *Nature*
437, 412-416 (2005).
3. J. H. Reynolds, D. J. Heeger, The normalization model of attention. *Neuron* **61**, 168-185
(2009).
4. J. H. Reynolds, T. Pasternak, R. Desimone, Attention increases sensitivity of V4 neurons.
Neuron **26**, 703-714 (2000).
5. P. Sörqvist, S. Stenfelt, J. Rönnberg, Working memory capacity and visual–verbal
cognitive load modulate auditory–sensory gating in the brainstem: Toward a unified view
of attention. *J. Cogn. Neurosci.* **24**, 2147-2154 (2012).
6. A. Schoups, R. Vogels, N. Qian, G. Orban, Practising orientation identification improves
orientation coding in V1 neurons. *Nature* **412**, 549-553 (2001).
7. A. F. Teich, N. Qian, Learning and adaptation in a recurrent model of V1 orientation
selectivity. *J. Neurophysiol.* **89**, 2086-2100 (2003).
8. A. A. Faisal, L. P. Selen, D. M. Wolpert, Noise in the nervous system. *Nat. Rev. Neurosci.*
9, 292-303 (2008).
9. M. N. Shadlen, K. H. Britten, W. T. Newsome, J. A. Movshon, A computational analysis of
the relationship between neuronal and behavioral responses to visual motion. *J*
Neurosci. **16**, 1486-1510 (1996).
10. Z. W. Davis, L. Muller, J. Martinez-Trujillo, T. Sejnowski, J. H. Reynolds, Spontaneous
travelling cortical waves gate perception in behaving primates. *Nature* **587**, 432-436
(2020).

11. K. T. Sillar, A. Roberts, A neuronal mechanism for sensory gating during locomotion in a
vertebrate. *Nature* **331**, 262-265 (1988).
12. M. H. Ouellette, M. J. Desrochers, I. Gheta, R. Ramos, M. Hendricks, A Gate-and-Switch
Model for Head Orientation Behaviors in Caenorhabditis elegans. *eNeuro* **5** (2018).
13. E. Eggermann, Y. Kremer, S. Crochet, C. C. Petersen, Cholinergic signals in mouse
barrel cortex during active whisker sensing. *Cell reports* **9**, 1654-1660 (2014).
14. J. M.Blackwell, M. N. Geffen, Progress and challenges for understanding the function of
cortical microcircuits in auditory processing. *Nat. Commun.* **8**, 1-9 (2017).
15. M. M. Churchland, J. P. Cunningham, M. T. Kaufman, J. D. Foster, P. Nuyujukian, *et al.*,
Neural population dynamics during reaching. *Nature*, **487**, 51-56 (2012).
16. R. C. Williamson, B. Doiron, M. A. Smith, M. Y. Byron, Bridging large-scale neuronal
recordings and large-scale network models using dimensionality reduction. *Curr. Opin.*
Neurobiol. **55**, 40-47 (2019).
17. M. B. Ahrens, J. M. Li, M. B. Orger, D. N. Robson, A. F. Schier, *et al.*, Brain-wide
neuronal dynamics during motor adaptation in zebrafish. *Nature* **485**, 471-477 (2012).
18. M. B. Ahrens, M. B. Orger, D. N. Robson, J. M. Li, P. J. Keller, Whole-brain functional
imaging at cellular resolution using light-sheet microscopy. *Nat Methods* **10**, 413-420
(2013).
19. W. C. Lemon, S. R. Pulver, B. Höckendorf, K. McDole, K. Branson, *et al.*, Whole-central
nervous system functional imaging in larval Drosophila. *Nat. Commun.* **6**, 1-16 (2015).
20. T. Schrödel, R. Prevedel, K. Aumayr, M. Zimmer, A. Vaziri, Brain-wide 3D imaging of
neuronal activity in Caenorhabditis elegans with sculpted light. *Nat. Methods* **10**, 1013-
1020 (2013).
21. J. P. Nguyen, F. B. Shipley, A. N. Linder, G. S. Plummer, M. Liu, *et al.* (2016). Whole-
brain calcium imaging with cellular resolution in freely behaving Caenorhabditis
elegans. *Proceedings of the National Academy of Sciences*, *113*(8), E1074-E1081.
22. V. Venkatachalam, N. Ji, X. Wang, C. Clark, J. K. Mitchell, *et al.,* Pan-neuronal imaging in
roaming Caenorhabditis elegans. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E1082-E1088
(2016).
23. J. J. How, S. Navlakha, S. H. Chalasani, Neural network features distinguish
chemosensory stimuli in Caenorhabditis elegans. *bioRxiv* [Preprint] (2020).
https://www.biorxiv.org/content/10.1101/2020.02.18.955245v1 (accessed 31 October
2021).
24. E. Yemini, A. Lin, A. Nejatbakhsh, E. Varol, R. Sun, *et al.*, NeuroPAL: a multicolor atlas
for Whole-Brain neuronal identification in C. elegans. *Cell* **184**, 272-288 (2021).
25. S. Kato, H. S. Kaplan, T. Schrödel, S. Skora, T. H. Lindsay, *et al.*, Global brain dynamics
embed the motor command sequence of Caenorhabditis elegans. *Cell*, **163**, 656-669
(2015).
26. A. L. Nichols, T. Eichler, R. Latham, M. Zimmer, A global brain state underlies C. elegans
sleep behavior. *Science* **356** (2017).
27. S. Skora, F. Mende, M. Zimmer, Energy scarcity promotes a brain-wide sleep state
modulated by insulin signaling in C. elegans. *Cell Rep.* **22**, 953-966 (2018).
28. N. Chronis, M. Zimmer, C. I. Bargmann, Microfluidics for in vivo imaging of neuronal and
behavioral activity in Caenorhabditis elegans. *Nat. Methods* **4**, 727-731 (2007).
29. A. J. Calhoun, A. Tong, N. Pokala, J. A. Fitzpatrick, T. O. Sharpee, *et al.* Neural
mechanisms for evaluating environmental variability in Caenorhabditis elegans. *Neuron*
86, 428-441 (2015).
30. S. H. Chalasani, N. Chronis, M. Tsunozaki, J. M. Gray, D. Ramot, *et al.*, Dissecting a
circuit for olfactory behaviour in Caenorhabditis elegans. *Nature* **450**, 63-70 (2007).
31. J. Akerboom, N. Carreras Calderón, L. Tian, S. Wabnig, M. Prigge, *et al.* Genetically
encoded calcium indicators for multi-color neural activity imaging and combination with
optogenetics. *Front Mol Neurosci.* **6** (2013).
32. T. W. Chen, T. J. Wardill, Y. Sun, S. R. Pulver, S. L. Renninger, *et al.*, Ultrasensitive
fluorescent proteins for imaging neuronal activity. *Nature* **499**, 295-300 (2013).

33. E. A. Pnevmatikakis, D. Soudry, Y. Gao, T. A. Machado, J. Merel, *et al.* Simultaneous
denoising, deconvolution, and demixing of calcium imaging data. *Neuron*, *89*(2), 285-299.
34. Hendricks, M., Ha, H., Maffey, N., Zhang, Y. (2012). Compartmentalized calcium
dynamics in a C. elegans interneuron encode head movement. *Nature* **487**, 99-103
(2016).
35. Y. Shen, Q. Wen, H. Liu, C. Zhong, Y. Qin, *et al.*, An extrasynaptic GABAergic signal
modulates a pattern of forward movement in Caenorhabditis elegans. *Elife* **5**, e14197
(2016)..
36. A. Kocabas, C. H. Shen, Z. V. Guo, S. Ramanathan, Controlling interneuron activity in
Caenorhabditis elegans to evoke chemotactic behaviour. *Nature* **490**, 273-277 (2012).
37. H. S. Kaplan, O. S. Thula, N. Khoss, M. Zimmer, Nested neuronal dynamics orchestrate
a behavioral hierarchy across timescales. *Neuron* **105**, 562-576 (2020).
38. S. Kato, Y. Xu, C. E. Cho, L. F. Abbott, C. I. Bargmann, Temporal responses of C.
elegans chemosensory neurons are preserved in behavioral dynamics. *Neuron* **81**, 616-
628 (2014).
39. D. W. Hosmer, S. Lemeshow, R. X. Sturdivant. *Applied logistic regression* (Wiley, 2000)
40. N. Frosst, G. Hinton, Distilling a neural network into a soft decision tree. *arXiv* [Preprint]
(2017). https://arxiv.org/abs/1711.09784 (accessed 31 October 2021).
41. S. Linderman, A. Nichols, D. Blei, M. Zimmer, L. Paninski, Hierarchical recurrent state
space models reveal discrete and continuous dynamics of neural activity in C.
elegans. *BioRxiv* [Preprint] (2019). https://www.biorxiv.org/content/10.1101/621540v1
(accessed 31 October 2021).
42. I. Tsamardinos, E. Greasidou, G. Borboudakis, Bootstrapping the out-of-sample
predictions for efficient and accurate cross-validation. *Machine Learning* **107**, 1895-1922
(2018).
43. J. Pahlberg, A. P. Sampath, Visual threshold is set by linear and nonlinear mechanisms
in the retina that mitigate noise: How neural circuits in the retina improve the
signal-to-noise ratio of the single-photon response. *Bioessays* **33**, 438-447 (2011).
44. L. M. Jones, D. A. Depireux, D. J. Simons, A. Keller, Robust temporal coding in the
trigeminal system. *Science* **304**, 1986-1989 (2004).
45. C. Massot, A. D. Schneider, M. J. Chacron, K. E. Cullen, The vestibular system
implements a linear–nonlinear transformation in order to encode self-motion. *PLoS*
Biol. **10**, e1001365 (2012).
46. G. De Palo, G. Facchetti, M. Mazzolini, A. Menini, V. Torre, C. Altafini, Common
dynamical features of sensory adaptation in photoreceptors and olfactory sensory
neurons. *Sci. Rep.*, **3**, 1-8 (2013).
47. S. G. Leinwand, C. J. Yang, D. Bazopoulou, N. Chronis, J. Srinivasan, *et al.*, Circuit
mechanisms encoding odors and driving aging-associated behavioral declines in
Caenorhabditis elegans. *Elife* **4**, e10181 (2015).
48. B. T. Sagdullaev, M. A. McCall, P. D. Lukasiewicz, Presynaptic inhibition modulates
spillover, creating distinct dynamic response ranges of sensory output. *Neuron* **50**, 923-
935 (2006).
49. K. M. Hallinen, R. Dempsey, M. Scholz, X. Yu , A. Linder, *et al.*, Decoding locomotion
from population neural activity in moving C. elegans. *Elife* **10**, e66135 (2021).
50. I.T. Jolliffe, Principal Component Analysis, Second Edition (Springer, 2002).
51. H. Straka, J. Simmers, B. P. Chagnaud, A new perspective on predictive motor
signaling. *Curr. Biol.* **28**, R232-R243 (2018).
52. T. B. Crapse, M. A. Sommer, Corollary discharge across the animal kingdom. *Nat. Rev.*
Neurosci. **9**, 587-600 (2008).
53. N. A. Croll, Components and patterns in the behaviour of the nematode Caenorhabditis
elegans. *J. Zoology* **176**, 159-176 (1975).
54. V. Susoy, W. Hung, D. Witvliet, J. E. Whitener, M. Wu, *et al.*, Natural sensory context
drives diverse brain-wide activity during C. elegans mating. *Cell* **184**, 5122-5137 (2021).

55. D. R. Albrecht, C. I. Bargmann, High-content behavioral analysis of Caenorhabditis
elegans in precise spatiotemporal chemical environments. *Nat. Methods* **8**, 599-605
(2011).
56. E. A. Pnevmatikakis, A. Giovannucci, NoRMCorre: An online algorithm for piecewise rigid
motion correction of calcium imaging data. *J Neurosci Methods.* **291**, 83-94 (2017).
57. D. A. Fish, A. M. Brinicombe, E. R. Pike, J. G. Walker, Blind deconvolution by means of
the Richardson–Lucy algorithm. *JOSA A* **12**, 58-65 (1995).
58. L. Paninski, J. Pillow, J. Lewi, Statistical models for neural encoding, decoding, and
optimal stimulus design. *Prog Brain Res.* **165**, 493-507 (2007).
59. D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting linear mixed-effects models using
lme4. *arXiv* [Preprint] (2014). https://arxiv.org/abs/1406.5823 (accessed 31 October
2021).

Figures and Tables

Figure 1. Food-stimulated whole-brain activity reveals six functional clusters. (A) Stimulus
presentation and imaging setup. A computer precisely controls delivery of a stimulus sequence
that alternates between variable-length pulses of two liquid flows: bacterial food stimulus (gold)
and control buffer (light blue). This stimulus sequence is presented to the nose of a *C. elegans*
animal that is restrained in a microfluidic chip. Volumes of the *C. elegans* head are acquired and
subsequently processed to acquire calcium traces. (B–D) Identification of neurons in the
NeuroPAL–GCaMP6s strain. Some photobleaching occurred due to acquisition after calcium
imaging (see Methods: Cell Identification). (B) RME motor neuron and AVA command neuron. (C)
SMDD motor neuron. (D) SMDV motor neuron. (E) Neurons and their associated direction of
locomotion. (F–G) Number of neurons in each functional cluster for all animals, based on > 85%
correlation with representative neurons. (F) non-NeuroPAL–GCaMP5K animals. (G) NeuroPAL–
GCaMP6s animals. (H) Calcium activity of low-noise active neurons for a single worm. Gray
shading represents bacterial stimulus pulse duration. (I–K) Calcium activity averaged across all
traces within a functional cluster. Colors of average calcium traces corresponds with colors of
individual traces in (H). (I) OFF and ON clusters. (J) AVA and RME clusters. (K) SMDD and
SMDV clusters.
935

Figure 2. Bacteria onset and removal differentially drive activity of sensory, AVA, and RME
clusters. (A) Stimulus sequence features are fed into an encoding model (see Materials and
Methods: Encoding model) to predict sensory cluster activity. (B–C) Representative examples of
sensory neurons raw traces compared with model predictions. (B) ON cell. (C) OFF cell. (D–E)
Time-to-half-peak distributions for the best performing (D) ON and (E) OFF cell adapting models.
(F–H) Stimulus states (left, highlighted in gray) and their corresponding flow configurations
relative to the *C. elegans* nose. (F) Bacteria from channel 1 flows over nose. (G) Buffer from
channel 2 flows over nose. (H) As a control, buffer emanates from both channel 1 and channel 2.
(I) AVA and RME cluster activity, as well as worm identity, is fed into a decoding model (see
Materials and Methods: Decoding model) to predict different stimulus states (highlighted in gray).
In the null model, only worm identity is used. (J) Out-of-bootstrap cross-validation performance of
the full decoding model that includes AVA and RME cluster activity. (K–R) Temporal filters
predicting stimulus states from AVA and RME cluster activity for buffer↔buffer stimulus
sequences (K-N) and bacteria↔buffer sequences (O–R). Gray shading represents the prediction
time window (with 0 s as the halfway point of the prediction window), such that preceding time
represents baseline activity and subsequent time represents delayed effects. Median bootstrap
temporal filters are plotted, with graded shading indicating 50%, 75%, and 90% of bootstraps.
956
957

Figure 3. Identification of interpretable network states that vary in sensory gating properties. (A) The top four network states identified by the soft decision tree gating model portion of the SDT-MLR forecasting model (see Fig. S5, Materials and Methods: SDT-MLR model). Cyan shading represents the time windows for which a particular state's probability exceeds 0.75. (B–C) Temporal filters predicting rise and fall of AVA and RME cluster activity from (B) recent history of AVA, RME, SMDV, and SMDD cluster motor/command activity; and (C) ON and OFF cell cluster sensory activity. (B–C) Median bootstrap temporal filters are plotted, with graded shading indicating 50%, 75%, and 90% of bootstraps. (D) Difference in out-of-bootstrap cross-validation performance between models that included both network history and sensory activity and models that that only included network history.

Figure 4. Effect of stimulus timescales and sensory context on AVA and RME clusters. (A–B) Maximum magnitudes of temporal filters predicting rise and fall of AVA and RME activity from sensory neuron clusters. Maximum magnitudes are calculated from t-12s to t-0s relative to the start of the prediction window (t). Sums of absolute values of magnitudes measure the overall sensory influence within a state. (A) Comparison of all stimulus pulse lengths, only high-variance pulses, and only low-variance pulses. (B) Comparison of buffer↔buffer and bacteria↔buffer stimulus sequences. Only high-variance pulses were compared. Channel 1 and 2 are the same as ON and OFF, respectively, for bacteria↔buffer stimulus sequences, as seen in (A).