2

Title: **Multi-scale phenological niches of seed fall in diverse Amazonian plant communities**

Damie Pak[1]*, Varun Swamy[2], Patricia Alvarez-Loayza[3], Fernando Cornejo[4], Simon A.
Queenborough[5], Margaret R. Metz[6], John Terborgh[7], Renato Valencia[8], S. Joseph Wright[9],
Nancy C. Garwood[10], and Jesse R. Lasky[1]*

[1]Department of Biology
Pennsylvania State University
University Park, PA
USA

[2]San Diego Zoo Institute for Conservation
Research
Escondido, CA
USA

[3]Duke University Hospitals
Durham, NC
USA

[4]Proyecto Castaña
Puerto Maldonado, Madre de Dios
Perú

[5]Yale School of the Environment
Yale University
New Haven, CT
USA

[6]Department of Biology
Lewis & Clark College
Portland, OR
USA

[7]Center for Tropical Conservation
Nicholas School of the Environment
Duke University
Durham, NC
USA

[8]Pontificia Universidad Católica del
Ecuador
Quito
Ecuador

[9]Smithsonian Tropical Research Institute
Apartado 0843–03092
Balboa
Republic of Panama

[10]School of Biological Sciences
Southern Illinois University Carbondale
Carbondale, IL 62901
USA

*Corresponding authors:
damiepak@umich.edu, lasky@psu.edu

Keywords: community assembly,
rainforest, spectral analysis, reproduction,
frugivory, dispersal syndrome, functional
traits

Abstract

Phenology has long been hypothesized as an avenue for niche partitioning or interspecific facilitation, both promoting species coexistence. Tropical plant communities exhibit striking diversity in reproductive phenology, but many are also noted for large synchronous reproductive events. Here we study whether the phenology of seed fall in such communities is non-random, what are the temporal scales of phenological patterns, and ecological factors that drive reproductive phenology. We applied multivariate wavelet analyses to test for phenological synchrony versus compensatory dynamics (*i.e.* anti-synchronous patterns where one species' decline is compensated by the rise of another) among species and across temporal scales. We used data from long-term seed rain monitoring of hyperdiverse plant communities in the western Amazon. We found significant synchronous whole-community phenology at multiple time scales, consistent with shared environmental responses or positive interactions among species. We also observed both compensatory and synchronous phenology within groups of species likely to share traits (confamilials) and seed dispersal mechanisms. Wind-dispersed species exhibited significant synchrony at ~6 mo scales, suggesting these species share phenological niches to match seasonality of wind. Our results suggest that community phenology is shaped by shared environmental responses but that the diversity of tropical plant phenology may partly result from temporal niche partitioning. The scale-specificity and time-localized nature of community phenology patterns highlights the importance of multiple and shifting drivers of phenology.

INTRODUCTION

Species within ecological communities often exhibit interspecific diversity in the phenology of key life events. This diversity may represent an axis of niche partitioning that reflects community assembly and evolutionary history (Ashton et al. 1988, Gonzalez and Loreau 2009, Wolkovich and Cleland 2011, Bernard-Verdier et al. 2012, Godoy and Levine 2013). Species differences in phenology may limit interspecific competition and promote species coexistence by causing niche complementarity through time in resource use or interactions with mutualists like pollinators, or in apparent competition mediated by natural enemies (Robertson 1895, Rathcke and Lacey 1985). Alternatively, phenological synchrony may be promoted by pulses in resource supply or periodically harsh environmental conditions that limit the possible phenological options (Gentry 1974, Rathcke and Lacey 1985, Vasseur et al. 2014, Usinowicz et al. 2017, Detto et al. 2018). Facilitation due to enhanced attraction of mutualist animals or predator satiation may also promote synchronous reproduction (Janzen 1974). However, phenology remains a relatively poorly characterized dimension of functional diversity in many communities, owing to a lack of long-term monitoring and the multi-scale complexity of phenology (Wolkovich et al. 2014).

Co-occurring plants with similar reproductive phenology might be more likely to compete for frugivores (Saracco et al. 2005) or other resources, given the requirements of reproduction (Karlsson and Méndez 2005). As a result, those species capable of coexisting might partition phenological space. Researchers have studied phenological niches in tropical forests (e.g. Gentry 1974, Stiles 1977, Wheelwright 1985, Ashton et al. 1988, Poulin et al. 1999, Jones and Comita 2010) and elsewhere (Elzinga et al. 2007, Botes et al. 2008, Albrecht et al. 2015). Within a community, phenological niche partitioning might be strongest among species with shared mutualists or shared resource requirements (Encinas-Viso et al. 2012), similarities that

often characterize phylogenetically related species (Robertson 1895, Prinzing et al. 2001,

Donoghue 2008, Davies et al. 2013). However, there are simultaneous and opposing processes

interacting with phenology, such as abiotic seasonality versus resource competition. As a result,

phenological patterns indicative of shared temporal niches (interspecific synchrony) versus

temporal niche partitioning (interspecific compensation, or anti-synchrony) may only emerge at

certain time scales or over certain periods of time (Baird 1980, Vasseur et al. 2005, Keitt 2008,

Lasky et al. 2016).

Tropical plant communities have highly varied phenology and there are often multiple

species reproducing at any given time of the year (Frankie et al. 1974, Gentry 1974, van Schaik

et al. 1993), including for the stage that we study here: seed fall (Smythe 1970, Chang-Yang et

al. 2016, Detto et al. 2018). The phenological diversity of tropical plants may be made possible

by favorable temperature and (in rainforests) moisture for much of the year (Gentry 1974,

Usinowicz et al. 2017). Despite this potential, tropical plant communities often do exhibit

synchrony among at least a subset of the community, perhaps due to shared responses to abiotic

seasonality (van Schaik et al. 1993, Detto et al. 2018) or seasonality in frugivory and seed

dispersal (Poulin et al. 1999). It seems likely that different species may be limited by different

conditions fluctuating across the year (e.g. light, moisture, heat), thus diversity in the phenology

of tropical seed fall may be a consequence of distinct strategies or sensitivities to seasonality in

resources (Lasky et al. 2016). Additionally, positive density dependent interactions among

species may promote synchrony, for example masting that overwhelms seed predators (Ashton et

al. 1988, Jones and Comita 2010) or facilitation by frugivore attraction (Carlo 2005). Abiotic

conditions may control species interactions, such that synchronous or compensatory reproduction

occur during periods with specific abiotic conditions (Vasseur et al. 2005).

138   We used wavelet analyses to determine whether species exhibited synchronous or

139 compensatory (anti-synchronous) seed rain (illustrated in Figure 1) (Lasky et al. 2016). Wavelets

140 are basis functions, combinations of which can be used to characterize signals in data (here, time

141 series of seed rain). Wavelet transformations decompose signals into patterns at different scales,

142 like other spectral analyses, but with the added advantage that wavelets can characterize time-

143 localized and nonstationary patterns, i.e. patterns that are inconsistent over a time series

144 (Terrence and Compo 1998, Keitt 2008). In the wavelet transformation, the base wavelet is

145 translated across the time series at varying scales/frequencies of the wavelet to identify the

146 important time scales that contribute to the variability in the signal (Cazelles et al. 2008). By

147 resolving non-stationary and scale-specific patterns, we may improve our ability to detect

148 multiple opposing processes affecting seed rain dynamics at different temporal scales (scale-

149 specific) or points in time (non-stationary). For example, while species may all increase

150 reproduction during once-a-year seasons of high resource supply (annual-scale synchrony),

151 species may peak in reproduction at different points within a favorable season (Figure 1, within-

152 season-scale compensatory dynamics, Lasky et al. 2016).

153   We addressed the following questions:

154  1. Do communities exhibit compensatory patterns or synchronous patterns of reproduction

155   (seed fall) through time?

156  2. Is evidence for compensatory and synchronous dynamics scale-specific or non-stationary

157   (i.e. changing through time)?

158  3. Is evidence for niche partitioning of seed fall phenology strongest among functionally

159   similar species, potentially those with the greatest likelihood of interspecific competition?

Do related species exhibit stronger compensatory dynamics? Or alternatively do

morphologically similar or related species exhibit synchrony?

4. Does community phenology differ between sites and time periods of contrasting climate,

suggesting phenological niches may be mediated by fluctuations in environment?

METHODS

Study sites - We studied two forest plots in the western Amazon basin, in Cocha Cashu, Peru

and Yasuní, Ecuador (Figure S1). These plots were monitored across different intervals, from

February 2000–February 2017 in Yasuní and September 2002–January 2011 in Cocha Cashu.

The study plot in Ecuador was located in Yasuní National Park at the Estación Científica Yasuní

(0° 41' S, 76° 24' W), a research station maintained by Pontificia Universidad Católica del

Ecuador. The Yasuní lowland rain forest is in the wettest and least seasonal region of the

Amazon (Xiao et al. 2006, Silman 2007). Mean annual rainfall is 2826 mm, with no months

having <100 mm rainfall on average (Valencia et al. 2004b, 2004a). Seed traps were placed

within the 50-ha Yasuní Forest Dynamics Plot (YFDP, established in 1995), where elevation

ranged from 216 to 248 m. In 25 ha of this forest plot, 1104 tree species were recorded (Valencia

et al. 2004b, 2004a).

The study plot in Peru is located at Cocha Cashu Biological Station (11°54'S, 71°22'W),

which is situated at 360 m mean elevation within the core area of Manu National Park, at the

western margin of the Madre de Dios river basin. The study plot is located in mature floodplain

forest habitat, which comprises over 700 tree species (Pitman et al. 2002). Annual precipitation

ranges between 2000–2500 mm, with a pronounced dry season from June to October with

typically less than 100 mm monthly rainfall (Gentry 1993). In the period from September and

April there is an excess of fruit available for frugivorous vertebrates (Terborgh 1986b), which

may indicate that plants compete to attract frugivores during this period.

Seed rain data - In each plot, an array of seed traps was established. At Yasuní we followed the

methods of Wright and Calderon (1995). In February 2000, 200 seed traps were placed in the 50-

ha YFDP along trails but >50 m from the plot border. Every 13.5 m along the trails, a trap was

placed a random distance between 4 and 10 m perpendicular from the trail, alternating left and

right. Traps were constructed of PVC tubes and 1-mm fiberglass mesh, positioned 0.75 m above

ground, with an area of 0.57 m^2. Twice a month (median interval = 20 days, SD = 7.0 days),

from February 2000 to February 2017 all reproductive parts in each trap were counted and

identified to species or morphospecies using a reference collection of seeds and fruits maintained

on site.

At Cocha Cashu, year-round fruit and seed fall were counted between 2002 and 2011

within a 4-ha plot. A 17 × 17 array of 289 evenly spaced seed-fall traps was installed within the

central 1.44 ha (120 × 120 m) of the plot at the beginning of the study. Seed traps consisted of

0.49 m^2 (70 × 70 cm) open bags made of 1-mm nylon mesh sewn to wire frames with 0.5-mm

monofilament line. Corners of the traps were attached to nearby trees with 1-mm monofilament

line so that the traps were suspended approximately 1 m above the ground. The contents of the

traps were collected every ~2 weeks (median interval = 15 days, SD = 3.3 days), and all seeds,

fruit and fruit parts (capsules, valves, pods, etc.) were identified to species and recorded.

For fruit counts at both sites we estimated number of seeds collected by multiplying by

the average number of seeds per fruit. We divided counts by the number of days in intervals to

control for slight variation in intervals. Further detail is available in the Supplemental Material.

Seed dispersal mechanisms - We grouped species into different dispersal syndromes to test for non-random WMR within each syndrome, using slightly different approaches for each site depending on available data. At each site, we conducted two separate classification efforts, one for all species, and another more granular classification focused on tree species (excluding lianas, herbaceous, and woody shrubs).

At both sites for all plants we first focused our analysis on classifications as animal (Yasuní, N = 741, Cocha Cashu N = 174) or wind (Yasuní N = 139, Cocha Cashu N = 48) dispersed. Species with fruit that contain pulp or aril were considered animal-dispersed, while those with fruits or seeds adapted for flight were considered wind-dispersed. For the details of the dispersal syndrome classification of only tree species, see the Supplemental Material.

For all analyses on the taxonomic and dispersal groups, we only included groups that had at least 5 species. We did not use a lower threshold on number of records for inclusion of a species, as species contributions to group-wide phenological dynamics are weighted by number of seeds in the analyses below.

Weather data - We estimated monthly precipitation and minimum temperature at the plot level for each study site. Because local weather station data contained many missing observations, we used remotely sensed data. We used a ten-day precipitation time series estimated on a 0.05º grid by (Funk et al. 2014) using both remote and locally-sensed data. We used ECMWF/ERA-Interim reanalysis 4-hr temperature data at 2 m height estimated on a N128 Gaussian (~2º) grid (European Centre for Medium-Range Weather Forecasts 2009) and calculated daily minimum temperatures and then monthly values.

To estimate the rough pattern of wind seasonality in the study region, we used weather station data. For Cocha Cashu, we calculated average monthly wind speed from a station 150 km

away, within 100 m elevation of Cocha Cashu, for the years 2004–2009

(http://atrium.andesamazon.org/meteo_station_display_info.php?id=12). For Yasuní, we used a

weather station (http://www.serviciometeorologico.gob.ec/biblioteca/) 115 km away within 70 m

elevation of the Yasuní plot for the years 2005-2012.

Statistical analysis

Wavelet transformation of seed rain data - For each species, we summed seed rain for each

time point across all traps in the plot, resulting in a single time series for each species. Species

seed abundance distributions were extremely right-skewed, with a few small-seeded species

making up most seed counts. For example, the top five species make up 78% of seeds at Yasuní

(2 Melastome species, 2 *Cecropia* species, and *Peperomia macrostachya*, all small-seeded), and

90% of seeds at Cocha Cashu (all five were small-seeded *Ficus*). To capture broader community

patterns, rather than allowing patterns to be dominated by the few most abundant species, we

transformed the counts to reduce skew (Vasseur et al. 2005, Lasky et al. 2016), specifically a

natural log transformation of the counts + 1. To explore the sensitivity of our analyses to

transformation, we also conducted all analyses in parallel without any transformation, however

we focus our discussion on results from ln(counts + 1). In a subset of our analyses, we also

explored how removing rarer species influenced our results (see Supplemental Material, also for

a justification of using 1 as opposed to another constant). Note we did not address within-plot

intraspecific variation in phenology as we were primarily interested in population- and

community-wide phenology and its potential relationship to seed dispersers (that move across the

plots) and temporal climate fluctuations (that are limited within the plot).

We then applied a continuous Morlet wavelet transformation to each species' time series

(either log transformed, or untransformed, greater detail in the Online Supplement). To

characterize individual species' phenology in relation to the community or group of species, we

calculated the wavelet modulus ratio (WMR). WMR quantifies the relationship between the

variation in the aggregate community-wide reproduction (numerator of Eqn. S3) relative to

variation in species-level reproduction (Eqn. S3, Figure 1). When species seed rain dynamics

through time cancel each other out, aggregate variation is zero (declining seed rain is balanced

by increasing seed rain). Thus at zero, the WMR indicates compensation: all species-level

dynamics are compensated so that community level reproduction is constant (Figure 1). At unity,

the WMR signifies complete phenological synchrony among the species, as species-level

phenological dynamics are completely reflected at the community level.

To answer Questions 1 and 2 above, we calculated the whole-community WMR for all

species in Yasuní (0.10 to 8.5 yr periods) and Cocha Cashu (0.08 to 4.2 yr periods). The periods

differed between sites because of the frequency of trap collection and duration. Trap observation

dates were regularized to a constant interval following (Keitt 2014), and the minimum periods

were twice the median sampling interval, limiting artifacts associated with a few days difference

in intervals. The maximum scales were calculated as half the total duration of the study period at

each site. We tested statistical significance of whole-community WMR using bootstrapping

where each species had the phase of their wavelet at a given scale randomly shifted (Keitt 2008).

This shift in phase has the effect of shifting where a wave is located in time, resulting in random

patterns of among-species phenology. All analyses were run in R (v 3.3.2). WMR was calculated

using the package 'mvcwt' (Keitt 2014, https://github.com/thk686/mvcwt).

Taxonomic and seed dispersal groups - To investigate if species that are closely related share

similar phenological niches or partition phenological niches (Question 3 above), we focused on

taxonomic groups. Our analyses were done at the family-level to ensure sufficient sample size.

Confamilials often share characteristics making them likely to exhibit evolutionary niche

conservation or character displacement. Similarly, groups with shared dispersal mechanisms

might be more likely to exhibit non-random phenology (Question 3) so we separately grouped

species based on their dispersal syndromes.

For these grouped analyses (family or dispersal syndrome), we first calculated WMR (as

with the whole-community analyses above) for each taxonomic or dispersal group averaged

across the time series. We excluded WMR values within one scale unit of the beginning and end

of time series to reduce boundary artifacts (i.e. the cone of influence). To test the hypothesis that

species within a group exhibited more synchronous seed fall or compensatory seed fall than the

greater community, we generated a null distribution of each group's WMR using permutations.

We permuted species labels over the entire community while maintaining the number of species

in each group, calculated WMR for the members of the permuted group, and then repeated this

permutation 1000 times. If the observed WMR of a group averaged across time points was above

the 97.5th percentile of the permutation-based null distribution, we considered it as nominally

significant synchrony, if the observed WMR of a group averaged across time points was below

the 2.5th percentile of the null distribution, we considered it as nominally significant

compensation. We calculated two-tailed p-values from permutations and implemented false

discovery rate (FDR) control across the multiple hypothesis tests using the method of (Benjamini

and Yekutieli 2001).

Climatic association with synchrony vs compensatory dynamics - To determine whether

climatic fluctuations might influence phenology among community members (Question 4

above), we investigated the association of local temperature and precipitation on whole

community WMR calculated above. We calculated monthly average climate data and we

aggregated seed rain data to monthly average seed counts. We then calculated WMR for Yasuní

(2–70 mo scales) and Cocha Cashu (2–50 mo scales). We used wavelet transformation of the

climate variables at the specific scales so that we could calculate their relationship with

community WMR. Specifically, at each scale, we calculated the Pearson correlation coefficient

between the community WMR and the wavelet-transformed minimum temperature or

precipitation. Null distributions for correlations were generated by permuting the starting point

of the wavelet transformed climate time series while maintaining periodic boundaries (i.e.,

adding climate values from before the randomly chosen staring points to the end of the permuted

climate series) and calculating the Pearson correlation between the randomized climate wavelet

and the WMR (n=1000, but with only 70 or 50 unique possible values for Yasuní and Cocha

Cashu, respectively, and ignoring observations within one period of the boundary). The wavelet

transform of climate was done in the R package "WaveletComp" (Rösch and Schmidbauer

2016).

RESULTS

Community-wide phenology – At the ever-wet site Yasuní we found a general trend of strong

whole-community synchrony in transformed seed counts at scales of less than ~50 days, while

larger sub-annual periods were weaker (significant at FDR = 0.05, Figure 2A). At the annual

scale we also found significant synchrony for most of the study, and we found significant

synchrony at scales greater than ~2 yrs, strongest at ~3.85 and 7 yrs (FDR = 0.05). These

patterns were largely consistent (stationary) across the time period of the study, especially the

pattern of strong synchrony at ~50 days. However, there was a weakening of annual-scale

synchrony from 2012 to 2015. For untransformed seed counts, we found significant synchrony

only at ~50 days and ~4 yrs (FDR = 0.05, Figure S2).

In contrast to Yasuní, at the seasonally dry Cocha Cashu site we found weaker whole-

community synchrony in transformed counts at the sub-annual time scales before 2006, but for

2006 into early 2008 we found significant community-wide synchrony across a wide range of

temporal scales (FDR = 0.05, Figure 2B). Additionally, there was consistent significant

synchrony at the ~1 year and ~2 year, and >3 year scales across the duration of the study,

indicating some shared annual, biannual, and multi-year dynamics among species (FDR = 0.05).

However, for untransformed seed counts we found almost no evidence of non-random WMR

patterns at Cocha Cashu (Figure S2).

Phenology among confamilials - At Yasuní among the 28 families analyzed, we found that

species of some families exhibited significant compensatory dynamics of transformed seed rain

at sub-annual timescales (Figure 3A). Note that the null model for confamilials and dispersal

groups was the community background, so significance indicates a stronger pattern than the rest

of the community. In particular, species in the Annonaceae, Euphorbiaceae, Malpighiaceae,

Myristicaceae, and Urticaceae families exhibited significant compensatory dynamics at different

timescales up to 8 months (FDR = 0.05). That is, species that declined in reproduction over a few

months tended to be replaced by other species in the same family increasing in reproduction over

that timescale. Additionally, we found significant compensatory dynamics at the longer time

scales (e.g. ~2.9 yrs) for Annonaceae. By contrast, at Cocha Cashu among the 27 studied

families, some families exhibited synchrony and other families compensation, compared to the

rest of the community. That is, confamilial species often showed significant synchrony or they

showed significant compensation, especially at sub-annual time scales (Figure 3B). For

untransformed data, we found only significant compensatory patterns within families, some of

which (Annonaceae, Menispermaceae, Malpighiaceae) also were significantly compensatory in

the analyses of transformed counts (FDR = 0.05, Figure S3).

We found some consistency of family patterns across sites. Annonaceae at Cocha Cashu

exhibited significant compensatory dynamics at sub-annual timescales (1–4 mos) similar to at

Yasuní (transformed counts). Additionally, Myristicaceae at both sites showed nominally

significant compensatory dynamics at 1–3 mo scales (though these were not significant after

FDR control at Cocha Cashu). By contrast, Fabaceae species showed nominally significant

synchrony for sub-annual to annual time scales at both sites (though these were not significant

after FDR control at Yasuní, transformed counts).

To help illustrate the patterns identified, we present one family with synchrony among

species and one family with compensatory dynamics among species (again compared to the other

species in the community). At Cocha Cashu, Bignoniaceae shows strong synchrony at ~0.5 and 1

year scales, which corresponds to a strong twice yearly peak in seed fall of multiple species and

many times of year with no seeds at all (Figure 4A, B). At Yasuní, Urticaceae showed significant

compensatory dynamics at scales of ~0.34–0.52 years, which corresponds to a family-wide

pattern where multiple species are typically releasing seed at any given point in time, and while

some annual synchrony was evident, different species were often reproductive at different times

of year so that seed counts never reached zero (Figure 4C, D, also see WMR figures for all

families in Supplemental Material).

Phenology among species sharing dispersal modes - Among the species with shared dispersal

syndrome, we found significant synchrony in log transformed counts at multiple scales at the

ever-wet Yasuní (Figure 5C & 5D). When considering all growth forms, we found that animal-

dispersed species exhibited significant synchrony at ~4 month scales but no significant

compensatory dynamics (FDR = 0.05). For wind-dispersed species, we found nominally

significant (p < 0.05) synchrony at ~6 month scales that did not pass FDR control. For only trees,

we did not find significantly non-random phenology for groups of species with similar size fruits

or similar abiotic dispersal mechanisms (though some were nominally significant p < 0.05,

Figure S5). Untransformed seed counts generally did not show significantly different WMR

within dispersal groups at Yasuní (Figures S4 and S6).

At seasonally dry Cocha Cashu, we found significant synchrony among log transformed

seed counts for dispersal groups of all growth forms (Figure 5). Wind-dispersed species

exhibited strong and significant synchrony at ~1, ~6, and ~12 month timescales (FDR = 0.05,

Figure S9). Wind variation also showed a peak in variability at ~12 month scales (and ~6 month,

depending on the metric, Figure S7). Animal-dispersed species showed significant synchrony at

~1–3 month scales (FDR = 0.05). For only tree species, we did not find non-random phenology

for species with similar dispersal syndromes, except for ~1–2 month synchrony for large

vertebrate-dispersed species (FDR = 0.05, Figure S8). Untransformed seed counts within

dispersal groups were not significantly different from the null, with the exception of animal-

dispersed species of all growth forms at ~1.7 yr scales (Figures S10–11).

Temperature, precipitation, and community-wide phenology - There was a significant

relationship between temperature (but not precipitation) and community-wide WMR for log

transformed seed counts (FDR = 0.05, Figure S13–14). Yasuní had a positive temperature–WMR

relationship most strongly at ~1.5 and 2 yr scales, indicating periodic increases in among-species

seed rain synchrony with warming temperatures. At the seasonally dry Cocha Cashu at ~1.3 yr

scales, there was a significant positive correlation between the WMR and minimum temperature,

similar to Yasuní. Unexpectedly, there was no increase in synchrony at Cocha Cashu during wet

or dry periods. There were some similar patterns observed in the untransformed counts' WMR–

climate correlations (Figure S15-16).

DISCUSSION

Ecological communities harbor extensive phenological diversity among member species,

especially in tropical wet forests (Frankie et al. 1974). This phenological diversity might help

explain species coexistence within communities if phenology is a key axis of niche partitioning

among species. Despite this great phenological diversity, there are well-known examples of

strong synchrony in tropical wet forests that lack strong seasonality (Ashton et al. 1988).

However, temporal traits (e.g. aspects of phenology) remain lesser-known dimensions of

diversity in ecological communities.

We found evidence that Amazonian plant communities frequently exhibit significant

reproductive synchrony at certain time scales at the whole-community level, suggesting that

shared responses to environment or positive interactions among species structure community

phenology. However, groups of species sharing dispersal mechanisms or related groups of

species sometimes exhibited compensatory (anti-synchronous) and sometimes synchronous

dynamics.

Do communities exhibit synchronous or compensatory reproduction? - Overall, we

found many cases of strong synchronous dynamics in seed rain at the whole-community level,

where across the whole community species tended to rise and fall in seed rain in concert. We

found almost no significant whole-community compensatory dynamics. The synchrony we

observed suggests whole-community dynamics are driven by shared responses to fluctuations in

environmental (abiotic or biotic) conditions such as rainfall or frugivore abundance, or by

positive heterospecific interactions such as enhanced frugivore attraction or natural enemy

satiation (van Schaik et al. 1993). Note that in our study we were only able to observe patterns

deviating from our permutation-based null. Processes that influence the shape (but not the phase,

which we randomized to generate our null, see Figure 2 A, B insets) of the phenology of the

species pool are outside our scope of detection (Case et al. 1983).

The ecological forces shaping phenology of fruiting and seed release are also tied to other

developmental stages. Flowering, leaf emergence and senescence, and seed germination all can

interact with abiotic and biotic conditions. Future research will benefit from an integrative

understanding of how full life-cycle phenology interacts with environment (Borchert 1996).

Are compensatory and synchronous dynamics scale-specific or non-stationary? - At both

sites, synchronous seed fall was strongest and most consistent at scales of approximately 1–2

mos, 1 yr, and >3 yrs. The fastest scale of synchrony at ever-wet Yasuní (~1.5-2 mos) may

represent shared and rapid community responses to relatively brief cloudless periods of high

radiation alternating with cloudy periods, similar to what was observed in response to rainfall in

a seasonally dry forest (Lasky et al. 2016). However, we did not find a correlation between

temperature or rainfall and WMR at the ~1 mo scale, and inconsistent correlations at larger time

scales. At both sites, the yearly and super-annual patterns of synchrony were fairly consistent

throughout the study. Clear annual peaks were visible even in the raw species or seed counts

over time, highlighting the strength of synchrony at this scale, despite the presence of many

reproductive species year-round (Figure 2 C, D, E, F).

At seasonally dry Cocha Cashu, we observed strong non-stationarity, in particular a shift

from approximately random to synchronous phenology, across all time scales <1 yr, with a peak

in WMR in 2008 (Figure 2B). The ecological explanation for this shift is unclear, but in 2008

there was the lowest nadir in species and total seed counts followed shortly by the highest peak

in species number (Figure 2 D, F). The decrease followed by a high peak in reproduction might

indicate multiple species were accumulating resources synchronously (reducing reproduction) in

order to subsequently invest a large amount in reproduction, akin to bursts of reproduction in

masting species (Janzen 1974, Ashton et al. 1988).

Is evidence for phenological partitioning strongest among ecologically similar species? - In

contrast to our findings at the whole-community level, we often found significant compensatory

dynamics within focal groups of species (confamilials and species with similar animal seed

dispersers), regardless of whether or not we transformed seed counts. Whole-community patterns

may obscure phenological niche partitioning that occurs within groups of closely interacting

species. This hypothesis is consistent with findings of ecological character displacement among

closely related species in a wide variety of taxa (Robinson and Wilson 1994, Dayan and

Simberloff 2005), including in time series of phytoplankton populations (Barraquand et al.

2018). Previous studies that have shown evidence for such partitioning have been largely focused

on groups of closely related species hypothesized to be closely interacting due to shared

pollinators or mutualists (Gentry 1974, Ashton et al. 1988, Botes et al. 2008). Future efforts

might also identify these species based on scale-specific, non-random phenological patterns

instead of relying on prior ecological or phenotypic knowledge. Annonaceae species showed

similar patterns of compensatory seed fall for within-year timescales at both sites, perhaps

indicating consistency in phenological niche partitioning among species in this family. Many

Annonaceae species have a similar fruit morphology of fleshy stipitate monocarps and animal

dispersal, such that phenological niche partitioning may promote coexistence.

It is challenging with our approach to determine the specific mechanisms behind any

phenological niche partitioning. An example misleading scenario is where species share dispersal

mechanisms but also other traits, of which the latter lead to resource competition that is

ameliorated by phenological niche partitioning. To demonstrate niche partitioning with respect to

animal dispersers at our sites requires additional evidence. For example, (Botes et al. 2008)

showed how species that deposit pollen on the same location on pollinators' bodies (suggesting

competition or interference) exhibited compensatory flowering, but species depositing pollen on

different body locations did not.

Synchrony among species sharing dispersal mode or related species - We found that among

all growth forms, wind-dispersed species exhibit strong synchrony of seed rain compared to the

whole community, particularly at time scales indicative of shared abiotic niches. At both sites we

found some degree of ~6 mo synchrony among wind-dispersed species (as well as the wind-

dispersed family Bignonicaceae at Cocha Cashu), consistent with the twice-yearly peaks in wind

speed observed at nearby weather stations (Figures S7, S9, S12). Near Cocha Cashu there is a

strong peak in wind speed in September (Figure S7), also consistent with the synchrony at this

site observed among wind-dispersed species at ~1 yr periods (Figures 5 & S9). Near Yasuní the

wind had a broad peak in average speed from September to November with a small peak in April

(Figure S12). The tendency for wind-dispersed tropical species to synchronously release seed

during windy seasons has been reported in the literature (Frankie et al. 1974, Janzen 1974, Detto

et al. 2018) and can be considered a positive control for our approach.

We found significantly greater synchrony for several speciose families at ~1 yr scales or

sub-annual scales for Arecaceae, Euphorbiaceae, Fabaceae, and Solanaceae (in addition to

aforementioned Bignoniaceae at Cocha Cashu). Given the dominance of climate variation at 1

year scales (Figures S14–15) the ~1 year synchrony in these families may be tied to shared

responses to abiotic fluctuations, although there are no clear ecological similarities among these

ecologically distinct families. Our finding may partly indicate greater power to detect patterns in

more speciose groups.

Community phenology and abiotic fluctuations - Gentry (1974) hypothesized that

phenological diversity of communities was promoted by permissive abiotic conditions and

longer growing seasons compared to seasonally harsh environments, where the range of potential

phenologies is narrower. We saw mixed evidence for this hypothesis based on fluctuations in

climate at our sites. At both sites, sub-annual WMR was only weakly (if significantly) associated

with temperature. Super-annual WMR showed some strong associations with temperature, in

particular at Yasuní were we found at >4.5 yr scales higher synchrony in warmer periods.

Similarly, at Barro Colorado Island in Panama, community-wide peaks in seed rain occur during

ENSO events (Detto et al. 2018), presumably as trees had shared (synchronous) responses to

increased light during these dry and warm periods (Wright and Calderón 2006). However, we

also found lower WMR in warmer periods at ~3.3 yr scales at Yasuní, leaving our conclusions

unclear.

We did not observe differences between ever-wet Yasuní and seasonally dry Cocha

Cashu that can easily be explained by the sites' precipitation seasonality. Even though Yasuní

may represent the most climatically favorable site, the whole community still showed strong

synchrony. Furthermore, there was no link between precipitation and the strength of synchrony

at seasonally dry Cocha Cashu (Figure 6). While we found temporal associations with climate,

we had too few sites to ask whether spatial climatic differences are associated with whole-

community differences in synchrony and compensation. Also note that the larger area covered by

traps at Yasuní and the longer time series may have increased the precision of our estimates of

population phenology and our power to detect non-random patterns, compared to Cocha Cashu.

Conclusion - Here we showed how whole-community phenology in diverse plant communities

is largely characterized by synchrony. However, we also found that groups of related or

ecologically similar species often show compensatory patterns of seed rain, indicating potential

phenological axes of niche partitioning that might promote species coexistence. Our results

highlight the scale-specific and sometimes non-stationary characteristics of community

phenology that may arise from the multiple opposing drivers of phenology. These changing

patterns across temporal scales are similar to those observed in trait based tests of community

assembly across spatial scales, where authors have found trait diversity patterns can reverse

across scales indicating potentially opposing community assembly forces acting at different

scales (Swenson and Enquist 2009). Flexible multi-scale analyses may reveal evidence of scale-

specific niche partitioning and environmental filtering and suggest the mechanisms that structure

communities.

Acknowledgements - This manuscript benefited from comments of Tomás Carlo, Kathryn

Cottingham, and two anonymous reviewers. Work at Yasuní was supported by funding to NCG

and collaborators from the Andrew W. Mellon Foundation, Natural Environment Research

Council (GR9/04037), British Airways, Department of Botany, Natural History Museum, and the

National Science Foundation (DEB-0614525, DEB-1122634, DEB-1754632, DEB-1754668).

We thank the Ecuadorian Ministerio del Ambiente for permission to work in Yasuní National

Park (under Nº 014-2019-IC-PNY-DPAO/AVS, Nº 012-2018-IC-PNY-DPAO/AVS, Nº 008-

2017-IC-PNY-DPAO/AVS, Nº. 012–2016-IC-FAU-FLO-DPAO-PNY, Nº. 014-2015-FLO-

MAE-DPAO-PNY, and earlier permits). We very gratefully thank Milton Zambrano for

collecting most of the Yasuní trap data from 2002–2017. We also thank Viveca Persson for help

initiating the censuses in 2000–2002, with assistance from Zornitza Aguilar, Paola Barriga and

Matt Priest, and Gorky Villa, Alvaro Perez and Pablo Alvia for help identifying species. Data collection at Cocha Cashu was supported by funding from the Andrew W. Mellon Foundation and the National Science Foundation (DEB-0742830). We thank the Peruvian authorities INRENA and SERNANP for permission to work in Manu National Park (under N° 020-CC-2008-INRENA-IANP, 05-CC-2009-SERNANP-PNM, 010-2010-SERNANP-JPNM, and earlier permits). More than 25 Peruvian undergraduate students assisted with data collection from 2002-11. Vishnu Viswanathan provided assistance digitizing weather records.

References

Albrecht, J., V. Bohle, D. G. Berens, B. Jaroszewicz, N. Selva, and N. Farwig. 2015. Variation in neighbourhood context shapes frugivore-mediated facilitation and competition among co-dispersed plant species. Journal of Ecology 103:526–536.

Ashton, P. S., T. J. Givnish, and S. Appanah. 1988. Staggered Flowering in the Dipterocarpaceae: New Insights Into Floral Induction and the Evolution of Mast Fruiting in the Aseasonal Tropics. The American Naturalist 132:44–66.

Bagchi, R., V. Swamy, J.-P. L. Farfan, J. Terborgh, C. I. A. Vela, N. C. A. Pitman, and W. G. Sanchez. 2018. Defaunation increases the spatial clustering of lowland Western Amazonian tree communities. Journal of Ecology 106:1470–1482.

Baird, J. W. 1980. The Selection and Use of Fruit by Birds in an Eastern Forest. The Wilson Bulletin 92:63–73.

Barraquand, F., C. Picoche, D. Maurer, L. Carassou, and I. Auby. 2018. Coastal phytoplankton community dynamics and coexistence driven by intragroup density-dependence, light and hydrodynamics. Oikos 127:1834–1852.

548 Benjamini, Y., and D. Yekutieli. 2001. The control of the false discovery rate in multiple testing

549   under dependency. Annals of Statistics 29:1165–1188.

550 Bernard-Verdier, M., M.-L. Navas, M. Vellend, C. Violle, A. Fayolle, and E. Garnier. 2012.

551   Community assembly along a soil depth gradient: contrasting patterns of plant trait

552   convergence and divergence in a Mediterranean rangeland. Journal of Ecology

553   100:1422–1433.

554 Borchert, R. 1996. Phenology and flowering periodicity of Neotropical dry forest species:

555   evidence from herbarium collections. Journal of Tropical Ecology 12:65–80.

556 Botes, C., S. D. Johnson, and R. M. Cowling. 2008. Coexistence of succulent tree aloes:

557   partitioning of bird pollinators by floral traits and flowering phenology. Oikos 117:875–

558   882.

559 Carlo, T. A. 2005. Interspecific Neighbors Change Seed Dispersal Pattern of an Avian-Dispersed

560   Plant. Ecology 86:2440–2449.

561 Case, T. J., J. Faaborg, and R. Sidell. 1983. THE ROLE OF BODY SIZE IN THE ASSEMBLY

562   OF WEST INDIAN BIRD COMMUNITIES. Evolution; International Journal of Organic

563   Evolution 37:1062–1074.

564 Chang-Yang, C.-H., I.-F. Sun, C.-H. Tsai, C.-L. Lu, and C.-F. Hsieh. 2016. ENSO and frost

565   codetermine decade-long temporal variation in flower and seed production in a

566   subtropical rain forest. Journal of Ecology 104:44–54.

567 Davies, T. J., E. M. Wolkovich, N. J. B. Kraft, N. Salamin, J. M. Allen, T. R. Ault, J. L.

568   Betancourt, K. Bolmgren, E. E. Cleland, B. I. Cook, T. M. Crimmins, S. J. Mazer, G. J.

569   McCabe, S. Pau, J. Regetz, M. D. Schwartz, and S. E. Travers. 2013. Phylogenetic

570   conservatism in plant phenology. Journal of Ecology 101:1520–1530.

Dayan, T., and D. Simberloff. 2005. Ecological and community-wide character displacement: the

next generation. Ecology Letters 8:875–894.

Detto, M., S. J. Wright, O. Calderón, and H. C. Muller-Landau. 2018. Resource acquisition and

reproductive strategies of tropical forest in response to the El Niño–Southern Oscillation.

Nature Communications 9:1–8.

Donoghue, M. J. 2008. A phylogenetic perspective on the distribution of plant diversity.

Proceedings of the National Academy of Sciences of the United States of America

105:11549–11555.

Elzinga, J. A., A. Atlan, A. Biere, L. Gigord, A. E. Weis, and G. Bernasconi. 2007. Time after

time: flowering phenology and biotic interactions. Trends in Ecology & Evolution

22:432–439.

Encinas-Viso, F., T. A. Revilla, and R. S. Etienne. 2012. Phenology drives mutualistic network

structure and diversity. Ecology Letters 15:198–208.

European Centre for Medium-Range Weather Forecasts. 2009. ERA-Interim Project. Research

Data Archive at the National Center for Atmospheric Research, Computational and

Information Systems Laboratory.

Frankie, G. W., H. G. Baker, and P. A. Opler. 1974. Comparative phenological studies of trees in

tropical wet and dry forests in the lowlands of Costa Rica. The Journal of Ecology:881–

919.

Funk, C. C., P. J. Peterson, M. F. Landsfeld, D. H. Pedreros, J. P. Verdin, J. D. Rowland, B. E.

Romero, G. J. Husak, J. C. Michaelsen, and A. P. Verdin. 2014. A Quasi-Global

Precipitation Time Series for Drought Monitoring.

Gentry, A. H. 1974. Flowering Phenology and Diversity in Tropical Bignoniaceae. Biotropica

6:64–68.

Gentry, A. H. 1993. Four neotropical rainforests. Yale University Press.

Godoy, O., and J. M. Levine. 2013. Phenology effects on invasion success: insights from

coupling field experiments to coexistence theory. Ecology 95:726–736.

Gonzalez, A., and M. Loreau. 2009. The Causes and Consequences of Compensatory Dynamics

in Ecological Communities. Annual Review of Ecology, Evolution, and Systematics

40:393–414.

Harrison, R. D., S. Tan, J. B. Plotkin, F. Slik, M. Detto, T. Brenes, A. Itoh, and S. J. Davies.

2013. Consequences of defaunation for a tropical tree community. Ecology Letters

16:687–694.

Janzen, D. H. 1974. Tropical Blackwater Rivers, Animals, and Mast Fruiting by the

Dipterocarpaceae. Biotropica 6:69–103.

Jones, F. A., and L. S. Comita. 2010. Density-dependent pre-dispersal seed predation and fruit

set in a tropical tree. Oikos 119:1841–1847.

Keitt, T. H. 2008. Coherent ecological dynamics induced by large-scale disturbance. Nature

454:331–334.

Keitt, T. H. 2014. mvcwt: Wavelet analysis of multiple time series.

Lasky, J. R., M. Uriarte, and R. Muscarella. 2016. Synchrony, compensatory dynamics, and the

functional trait basis of phenological diversity in a tropical dry forest tree community:

effects of rainfall seasonality. Environmental Research Letters 11:115003.

Pitman, N. C. A., J. W. Terborgh, M. R. Silman, P. Núñez V., D. A. Neill, C. E. Cerón, W. A.

Palacios, and M. Aulestia. 2002. A comparison of tree species diversity in two upper

amazonian forests. Ecology 83:3210–3224.

Poulin, B., S. J. Wright, G. Lefebvre, and O. Calderón. 1999. Interspecific synchrony and

asynchrony in the fruiting phenologies of congeneric bird-dispersed plants in Panama.

Journal of Tropical Ecology 15:213–227.

Prinzing, A., W. Durka, S. Klotz, and R. Brandl. 2001. The niche of higher plants: evidence for

phylogenetic conservatism. Proceedings of the Royal Society of London. Series B:

Biological Sciences 268:2383–2389.

Rathcke, B., and E. P. Lacey. 1985. Phenological patterns of terrestrial plants. Annual Review of

Ecology and Systematics:179–214.

Robertson, C. 1895. The Philosophy of Flower Seasons, and the Phaenological Relations of the

Entomophilous Flora and the Anthophilous Insect Fauna. The American Naturalist

29:97–117.

Robinson, B. W., and D. S. Wilson. 1994. Character Release and Displacement in Fishes: A

Neglected Literature. The American Naturalist 144:596–627.

Rösch, A., and H. Schmidbauer. 2016. WaveletComp 1.1: A guided tour through the R package.

URL: http://www. hsstat. com/projects/WaveletComp/WaveletComp_guided_tour. pdf.

Saracco, J. F., J. A. Collazo, M. J. Groom, and T. A. Carlo. 2005. Crop Size and Fruit

Neighborhood Effects on Bird Visitation to Fruiting Schefflera morototoni Trees in

Puerto Rico. Biotropica 37:81–87.

van Schaik, C. P., J. W. Terborgh, and S. J. Wright. 1993. The Phenology of Tropical Forests:

Adaptive Significance and Consequences for Primary Consumers. Annual Review of

Ecology and Systematics 24:353–377.

Silman, M. R. 2007. Plant species diversity in Amazonian forests. Pages 269–294 *in* M. B. Bush

and J. R. Flenley, editors. Tropical Rainforest Responses to Climatic Change. Springer

Berlin Heidelberg, Berlin, Heidelberg.

Smythe, N. 1970. Relationships between Fruiting Seasons and Seed Dispersal Methods in a

Neotropical Forest. The American Naturalist 104:25–35.

Stiles, F. G. 1977. Coadapted Competitors: The Flowering Seasons of Hummingbird-Pollinated

Plants in a Tropical Forest. Science 198:1177–1178.

Swenson, N. G., and B. J. Enquist. 2009. Opposing assembly mechanisms in a Neotropical dry

forest: implications for phylogenetic and functional community ecology. Ecology

90:2161–2170.

Usinowicz, J., C.-H. Chang-Yang, Y.-Y. Chen, J. S. Clark, C. Fletcher, N. C. Garwood, Z. Hao,

J. Johnstone, Y. Lin, M. R. Metz, T. Masaki, T. Nakashizuka, I.-F. Sun, R. Valencia, Y.

Wang, J. K. Zimmerman, A. R. Ives, and S. J. Wright. 2017. Temporal coexistence

mechanisms contribute to the latitudinal gradient in forest diversity. Nature 550:105–108.

Valencia, R., R. Condit, R. B. Foster, K. Romoleroux, G. Villa Munoz, J.-C. Svenning, E.

Magård, M. S. Bass, E. C. Losos, and H. Balslev. 2004a. Yasuni forest dynamics plot,

Ecuador. Tropical forest diversity and dynamism: findings from a large-scale plot

network:609–620.

Valencia, R., R. B. Foster, G. Villa, R. Condit, J.-C. Svenning, C. Hernández, K. Romoleroux, E.

Losos, E. Magård, and H. Balslev. 2004b. Tree species distributions and local habitat

658   variation in the Amazon: large forest plot in eastern Ecuador. Journal of Ecology 92:214–

659    229.

660 Vasseur, D. A., J. W. Fox, A. Gonzalez, R. Adrian, B. E. Beisner, M. R. Helmus, C. Johnson, P.

661    Kratina, C. Kremer, C. de Mazancourt, E. Miller, W. A. Nelson, M. Paterson, J. A.

662    Rusak, J. B. Shurin, and C. F. Steiner. 2014. Synchronous dynamics of zooplankton

663    competitors prevail in temperate lake ecosystems. Proceedings of the Royal Society B:

664    Biological Sciences 281.

665 Vasseur, D. A., U. Gaedke, and K. S. McCann. 2005. A seasonal alternation of coherent and

666    compensatory dynamics occurs in phytoplankton. Oikos 110:507–514.

667 Wheelwright, N. T. 1985. Competition for dispersers, and the timing of flowering and fruiting in

668    a guild of tropical trees. Oikos:465–477.

669 Wolkovich, E. M., and E. E. Cleland. 2011. The phenology of plant invasions: a community

670    ecology perspective. Frontiers in Ecology and the Environment 9:287–294.

671 Wolkovich, E. M., B. I. Cook, and T. J. Davies. 2014. Progress towards an interdisciplinary

672    science of plant phenology: building predictions across space, time and species diversity.

673    New Phytologist 201:1156–1162.

674 Wright, S. J., and O. Calderon. 1995. Phylogenetic Patterns among Tropical Flowering

675    Phenologies. Journal of Ecology 83:937–948.

676 Wright, S. J., and O. Calderón. 2006. Seasonal, El Niño and longer term changes in flower and

677    seed production in a moist tropical forest. Ecology Letters 9:35–44.

678 Xiao, X., S. Hagen, Q. Zhang, M. Keller, and B. M. Iii. 2006. Detecting leaf phenology of

679    seasonally moist tropical forests in South America with multi-temporal MODIS images.

680    Remote Sensing of Environment 103 :465–473.

Figure 1. Illustration of wavelet modulus ratio (WMR) as a measure of simulated multi-species patterns in reproduction. (A-C) Shows two-year time series of reproduction (excerpted from a 20-year series) for three different scenarios for a 5-species community (each species a different colored line). (D-F) Demonstrates WMR for these scenarios across a whole 20-year time series, where WMR is determined by the variation in total aggregate community reproduction relative to the variation in individual species reproduction. Thick black lines indicate regions of significantly non-random WMR (FDR = 0.05). When there is synchrony (A), the aggregate community variation is similar to the individual species variation, and WMR is high at the relevant scale (1 yr). When there are compensatory dynamics (B), the aggregate community variation is low but the individual species variation is high, and WMR is low at the relevant scale (1 yr). Both synchrony and compensatory dynamics can occur at different scales (C), in this case long-scale synchrony and short scale compensation. A small amount of noise was added to each scenario. The cone of influence (white shading in D-F) marks the regions where the wavelet transforms are affected by the boundaries of the sampling period.

Figure 2: The whole-community wavelet modulus ratio (WMR) of log-transformed seed counts (A,B) at Yasuní (1059 species) and Cocha Cashu (654 species), and time series of the total species in traps (C,D) and total estimated seeds in traps (E,F, natural log) in each sampling period. In (A,B) red indicates high WMR while blue indicates low WMR. When WMR is greater than the null expectation this is significant synchrony, and when WMR is lower than expected we term this significant compensation. Insets (in A and B) show WMR averaged across the time series (gray ribbon is null distribution, the line indicates observed average WMR with yellow being scales where WMR is significantly different than the null) The thin dashed contour lines

bound the points in time and scale (years) when the WMR was nominally significant (p<0.05)

based on bootstrapping, while thick black lines bound regions significant with a false discovery

rate (FDR) = 0.05. Nearly all significant regions in the plot are high WMR (yellow to red). The

cone of influence (white shading, A,D) marks the regions where the wavelet transforms are

affected by the boundaries of the sampling period. Note the null distribution of WMR values

changes with scales. Log transformed counts are analyzed here. See Figure S2 for

untransformed.

Figure 3: The averaged wavelet modulus ratio of families at Yasuní (A) and Cocha Cashu (B)

at the sub-annual (left) and interannual (right) scales compared to a community-wide null. The

number in parenthesis represents the number of species analyzed within the family. Colored

points represent nominally significant (p<0.05) synchronous (red) or compensatory (blue)

dynamics at the time scale. Thick borders around the points indicate significant points at false

discovery rate (FDR) = 0.05. Log transformed counts are presented here. See Figure S3 for

untransformed.

Figure 4. Two families at each site with distinct patterns of phenology, with each species' time

series of estimated seed counts in traps shown as a different colored line. Bignoniaceae (A-B) at

Cocha Cashu (60 species) shows synchrony especially at ~1 year scales, with multiple species

rising and falling in reproduction together in concert at these scales, and many times where no

species are releasing seed. By contrast, Urticaeae (C-D) at Yasuní (13 species) shows significant

compensatory dynamics compared the community-wide null at ~0.34–0.52 year scales. Although

there is some synchrony, the most abundant Urticaeae species peak at different parts of the year,

and even as some species decline in reproduction over these scales, others replace them, so that

there is always substantial reproduction by some members of the family (D).

Figure 5. The averaged wavelet modulus ratio for plant species of all growth forms based on

dispersal syndrome (thick lines). The number in parentheses represents the number of species

within the animal and wind-dispersed groups. The light blue ribbon represents the 2.5–97.5th

percentiles of the null-distribution generated through bootstrapping. Any points that lie above the

ribbon were considered nominally significant and synchronous while any points below the

ribbon were considered nominally significant and compensatory dynamics (p<0.05). Enlarged

points indicate significant at FDR = 0.05.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.