Plasmid fitness costs are caused by specific genetic conflicts

Authors

James P. J. Hall[1,2*], Rosanna C. T. Wright[2,3], Ellie Harrison[3], Katie J. Muddiman[2], A. Jamie Wood[4,5], Steve Paterson[1], Michael A. Brockhurst[3]

Affiliations

[1]Department of Evolution, Ecology and Behaviour, Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Crown Street, Liverpool, L69 7ZB

[2]Department of Animal and Plant Sciences, University of Sheffield, Western Bank, Sheffield, S10 2TN

[3]Division of Evolution and Genomic Sciences, University of Manchester, Oxford Road, Manchester, M13 9PL

[4]Department of Biology, University of York, Wentworth Way, York, YO10 5DD

[5]Department of Mathematics, University of York, Wentworth Way, York, YO10 5DD

*Correspondence to j.p.j.hall@liverpool.ac.uk

Abstract

Plasmids play an important role in bacterial genome evolution by transferring genes between lineages. Fitness costs associated with plasmid acquisition are expected to be a barrier to gene exchange, but the causes of plasmid fitness costs are poorly understood. Single compensatory mutations are often sufficient to completely ameliorate plasmid fitness costs, suggesting that such costs are caused by specific genetic conflicts rather than generic properties of plasmids, such as their size, metabolic burden, or expression level. Here we show — using a combination of experimental evolution, reverse genetics, and transcriptomics — that fitness costs of two divergent large plasmids in *Pseudomonas fluorescens* are caused by inducing maladaptive expression of a chromosomal tailocin toxin operon. Mutations in single genes unrelated to the toxin operon, and located on either the chromosome or the plasmid, ameliorated the disruption associated with plasmid acquisition. We identify one of these compensatory loci, the chromosomal gene *PFLU4242*, as the key mediator of the fitness costs of both plasmids, with the other compensatory loci either reducing expression of this gene or mitigating its deleterious effects by upregulating a putative plasmid-borne ParAB operon. The chromosomal mobile genetic element Tn6291, which uses plasmids for transmission, remained upregulated even in compensated strains, suggesting that mobile genetic elements communicate through pathways independent of general physiological disruption. Plasmid fitness costs caused by specific genetic conflicts are unlikely to act as a long-term barrier to horizontal gene transfer due to their propensity for amelioration by single compensatory mutations, explaining why plasmids are so common in bacterial genomes.

Introduction

Plasmid-mediated horizontal gene transfer (HGT) is a powerful force in bacterial evolution. Many of the traits necessary for the ecological success of bacteria in new environments are carried between bacterial strains and species by plasmids. In hospitals, plasmids mobilise antimicrobial resistance genes between pathogens, commensals, and environmental bacteria of diverse species [1–3]. In soil, plasmids are responsible for bioremediation of contaminated sites, and provide new ecological opportunities both to free-living and to plant-associated bacteria [4,5]. In infections, plasmids provide key virulence determinants necessary for colonisation and establishment, and factors that assist in immune evasion and pathogen transmission [6,7]. Thus in diverse habitats, plasmids are integral constituents of microbiomes, acting as potent drivers of bacterial evolutionary ecology, in large part through their role as vehicles of HGT [8–11].

Nevertheless, there exist various features of their bacterial hosts that restrict plasmid transmission, and thus the flow of genetic information [12]. Some of these features impede acquisition of plasmids, such as entry exclusion systems, CRISPR, and restriction systems [13,14]. Other barriers emerge from the inability of plasmids to replicate or to segregate effectively to daughter cells. However, even if a plasmid can enter, replicate, and be passed to daughter cells, its longer term persistence is contingent upon the fitness costs that emerge from plasmid carriage [15,16].

Plasmid-imposed fitness costs are the negative impacts on growth, replication, and/or survival that emerge as a consequence of plasmid carriage [15]. These costs are thought to act as major barriers to plasmid maintenance — and plasmid-mediated HGT — because cells burdened by the costs of plasmid carriage are outcompeted by others that lack the plasmid and its associated costs [17]. Importantly, selection for plasmid-borne genes (for example, by antibiotics) is insufficient to explain plasmid persistence in the long term, because those genes can become relocated to the chromosome through recombination, resulting in net fitness costs

of plasmid carriage, favouring plasmid loss [17–19]. Therefore, to understand plasmid

maintenance — and by extension the ability of plasmids to act as vehicles of gene exchange in

bacterial communities — it is crucial to understand the mechanisms that underlie plasmid fitness

costs, and how these costs may be resolved.

A variety of causes of plasmid fitness costs have been proposed [15], but in most cases the

underlying molecular mechanisms remain unknown. General properties of plasmids may explain

costs because cellular resources — not least nucleotides, amino acids, and the machineries of

replication, transcription, and translation — must be redirected to plasmid activity, imposing a

metabolic burden [16,20]. These costs would be expected to scale with plasmid size, copy

number, and gene expression, being higher for plasmids that are large, multi-copy, and/or have

highly active promoters. Plasmid DNA composition, particularly GC content and bias in codon

use, is another general property that can be a potential source of plasmid costs, exacerbating

the inefficiencies in bacterial physiology caused by plasmid maintenance [21,22]. Alternatively,

plasmid fitness costs can derive from the effects of specific plasmid genes. Some plasmid-borne

genes are costly in and of themselves, for example some efflux pumps [23] or the conjugative

pilus which can disrupt the cell membrane and exposes bacteria to phage predation [24],

whereas other plasmid genes elicit deleterious interactions with chromosomal genes, for

example by triggering stress responses [25] or interfering with metabolism [26,27].

Commonly, plasmids or the host bacterium undergo compensatory evolution, reducing the costs

of plasmid carriage and hence the strength of purifying selection against plasmids, enabling

maintenance [15,23,28–33]. Intuitively, the nature of compensatory mutations can provide an

insight into the causes of plasmid fitness costs. If costs emerge from general properties of

plasmids, we might expect large scale changes, such as deletions that reduce plasmid size, or

many individual changes to plasmid codon usage. Whereas, if costs derive from specific

plasmid genes or specific plasmid-chromosomal gene interactions, we might instead expect

targeted deletions, mutations, or regulatory changes affecting only those genes. Whilst some studies of compensatory evolution do report large-scale plasmid deletions [28,34], the majority implicate one or few mutations to specific genes as responsible for ameliorating the cost of plasmid carriage [23,25,29–33], suggesting that specific genetic interactions may commonly underlie plasmid fitness costs. Identifying these interactions, and how they are resolved by compensatory evolution, is crucial for understanding the evolution and ecology of horizontal gene transfer. In particular, the location of compensatory evolution is predicted to have important consequences: if the compensatory mutation occurs to the plasmid then it is propagated with onward horizontal transmission [30,35], whereas mutations occurring on the chromosome can potentially facilitate accumulation of many different plasmids [32,36]. However, chromosomal and plasmid compensations may not both be available for some plasmid-host combinations owing to the processes by which plasmid costs emerge and are resolved. Although many compensatory mutations have been identified for different plasmid-bacterial pairings, the mechanistic bases of compensation, and thus the nature of the resolved fitness costs, often remain unclear.

Here, we take advantage of a plasmid-host system where multiple genetic pathways of compensatory evolution exist, to investigate the mechanistic bases of plasmid fitness costs and their amelioration. We have previously described two pathways of chromosomal compensatory mutation in *Pseudomonas fluorescens* SBW25, targeting the *gacA/S* global regulatory system or the hypothetical protein *PFLU4242*, which individually ameliorate the fitness costs of megaplasmids pQBR103 (425 kb), pQBR57 (307 kb), or pQBR55 (157 kb) [29,37]. In this study, we describe an additional plasmid-borne compensatory locus, the lambda-repressor-like protein PQBR57_0059, that ameliorates the cost of pQBR57. Using a combination of experimental evolution, transcriptomics and reverse genetics we then show that specific genetic conflicts between chromosomal and plasmid genes, not plasmid size or gene expression *per se*, are the

117 principal cause of plasmid fitness costs in this system, and identify a single chromosomal locus

118 that acts as the key mediator of plasmid fitness costs.

Materials and methods

Bacterial strains

Pseudomonas fluorescens SBW25 was cultured in 6 ml King's B (KB) media in 30 ml glass

universals ('microcosms') at 28°C shaking at 180 rpm. Strains were labelled with either

gentamicin resistance (GmR) or streptomycin resistance (SmR) by insertion of mini-Tn7

resistance cassettes into *attTn7* sites according to the protocol of Choi & Schweizer [38]. Allelic

replacement of *gacA*, *gacS* and *PFLU4242* genes was achieved using the pUIC3 suicide vector

with a two-step protocol as described previously [33]. The 'evolved' pQBR57 plasmids came

from strains A.01.65.G.002 (ancestral type) and B.09.65.G.029 (V100A variant; Samples 1 and

49 respectively from [39]). These plasmids were allowed to transfer into an ancestral SmR

background which were used as donor strains for conjugation into the experimental lines.

Plasmid pQBR103 was from [40].

Transconjugants for use in experiments were generated by mixing equal volumes of overnight

cultures of donors and recipients and co-culturing for 24 hours before spreading on KB plates

containing mercuric chloride (20 μM) and either gentamicin (30 μg/ml) or streptomycin (250

μg/ml). Candidate transconjugant colonies were re-streaked on selective media and tested by

PCR for the presence of the plasmid. Independent transconjugants were used for each

replicate.

To express PQBR57_0059 independently of pQBR57, amplicons from the 'anc' and 'V100A'

strains were cloned EcoRI/KpnI into pUCP18 [41]. Inserts were sequenced, and 500 ng plasmid

was transformed into *P. fluorescens* SBW25 GmR using the sucrose method [38]. Three

independent transformants were generated (one for each replicate). Putative pQBR57 *par*

genes were cloned into pUCP18 and transformed into *P. fluorescens* SBW25 in a similar

manner. pUCP18-containing lines were selected and maintained on kanamycin 50 μg/ml, and

144 | pQBR57 and pQBR103 variants were transferred to these lineages as described above. To test

145 | effects of expressing plasmid-regulated genes, candidates were cloned EcoRI/KpnI into

146 | pME6032 [42], sequenced, and transformed into *P. fluorescens* as above. pME6032-containing

147 | lines were selected and maintained on 10 µg/ml (*E. coli*) or 100 µg/ml (*P. fluorescens*)

148 | tetracycline, and expression was induced with 100 µM IPTG.

Competitions

| For all competitions, GmR strains were competed against SmR-*lacZ* [43] strains. Overnight

| cultures of competing strains were washed and resuspended in KB, mixed in approximately

| equal ratios, and used to inoculate a KB microcosm at a 1:100 dilution. Samples of the

| competitors were spread on KB agar to estimate colony forming units at the beginning of the

| experiment and after 48 hours, and relative fitness was calculated as the ratio of Malthusian

| parameters $W = \ln(\text{test}_{end}/\text{test}_{start})/\ln(\text{reference}_{end}/\text{reference}_{start})$ [44]. These assays are designed

| to assess the fate of lineages that initially start with the plasmid, rather than plasmids *per se*,

| since competitors may gain or lose the plasmid during the assay. Nevertheless, replica plating

| showed that segregation had a negligible effect on the overall proportions of plasmid-bearers

| over the 48 hours of the competition (<0.01% of initially resistant colonies became susceptible,

| see also Hall et al. [45]). For the highest conjugation rate conditions (into *P. fluorescens*, where

| the donor has both compensatory mutations), transconjugants comprised ~10% of the final

| plasmid-bearing population. Measuring the fitness of the plasmid (rather than fitness of plasmid-

| bearers) increased calculated fitness measurements by <7% on average.

| 'Medium-term' competitions in soil were performed as described in [39]. Briefly, equal ratios of

| plasmid bearers and plasmid-free strains were mixed and added to soil microcosms, and

| transferred into fresh media every four days for 5 transfers in total. Populations were

| enumerated by plating or replica-plating onto selective media.

Growth curves

Growth curves of pME6032-carrying lines were established from a 1:1000 dilution of overnight culture into 150 µl KB + tetracycline ± 100 µM IPTG. Plates with growth curve cultures were incubated under humid conditions at 28°C shaking at 180 rpm (3 mm oribital radius), and OD600 was measured every 15 minutes on a Tecan Spark 10M plate reader. Four independent replicate cultures were established for each of three independent transformants.

RNA extraction and sequencing

RNA was extracted using TRI Reagent (T9424, Sigma Aldrich). Overnight cultures of bacteria were subcultured 1:50 into 6 ml prewarmed KB microcosms and grown at 28°C shaking at 180 rpm. When cultures reached intermediate exponential phase (OD600 ~ 0.6, as assessed on a Tecan Spark plate reader) 1 ml samples were added to 0.4 volumes of ice-cold 95% ethanol 5% phenol 'stop solution' [46], mixed gently, and incubated on ice for at least 30 minutes. Each of the three replicates was processed separately as a block. Samples were harvested at 4.5 G for 10 minutes, resuspended in 1 ml TRI reagent, and incubated at room temperate for 5 minutes. Amylene-stabilised chloroform (400 µl, Sigma-Aldrich 34854-M) was added to each tube, and incubated for 2-15 minutes at room temperature. Samples were then transferred to 5Prime Phase Lock Gel 'Heavy' tubes (VWR 733-2478) and centrifuged at 17 G for 15 minutes at room temperature. The aqueous phase was collected into a fresh microfuge tube and 450 µl isopropanol was added to precipitate RNA. Samples were incubated at room temperature for 30 minutes before pelleting at 17 G for 30 minutes. The pellet was rinsed by gently adding 1 ml of 70% ethanol and pelleting at 7.5 G for 5 minutes. The supernatant was carefully removed and the pellet allowed to air dry before resuspension in RNase-free water at 65°C. To remove any residual contaminating DNA, RNA was diluted to ≤200 ng/µl and treated with TURBO DNA-free RNase-free DNase (Invitrogen AM1907) according to the manufacturer's instructions. Reaction

products were purified with Agencourt RNAClean XP beads. Ribosomal RNA was removed

using the Bacterial RiboZero rRNA depletion kit (Illumina), and stranded libraries generated

using NEBNext Ultra-Directional RNA library preparation kit (Illumina). Libraries were

sequenced using paired-end 2x150 bp sequencing on two lanes of a HiSeq 4000. Short read

sequences are available on NCBI-GEO short-read archive, project accession GSE151570.

RNAseq analysis

Illumina adapter sequences were trimmed from the raw FASTQ files using Cutadapt version

1.2.1, using the option -O 3. The reads were further trimmed using Sickle version 1.200

(https://github.com/najoshi/sickle), with a minimum window quality score of 20, and reads

shorter than 20 bp after trimming were removed. FastQC

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and MultiQC [47] were used to

analyse the reads. Warnings arising from "Sequence content," "Sequence length" and

"Sequence duplication" modules were followed up and found to be largely due to an

overrepresentation of functional RNAs, namely tmRNA and RnaseP. Reads from each sample

were aligned strand-specifically using HISAT2 version 2.1.0 [48] to the corresponding reference

sequence(s): either the *P. fluorescens* SBW25 chromosome (AM181176) alone, the

chromosome with pQBR57 (LN713926), or the chromosome with pQBR103 (AM235768). The

options `phred33`, `no-spliced-alignment`, `new-summary`, and `rna-strandness RF` were used,

and the output was filtered using samtools to remove mappings with PHRED-scaled quality <

10. The function featureCounts from the Rsubread package version 1.30.9 [49] was used to

identify the reads mapping to each feature, and the table of counts was analysed in edgeR

version 3.22.5 [50]. Here, we restricted differential expression analysis to putative protein-

coding genes. Expression of chromosomal genes was analysed in a set of negative binomial

GLMs and all treatments were compared with the plasmid-free ancestor with a series of quasi-

likelihood F-tests. P-values were corrected using the Benjamini-Hochberg false-discovery rate

of q = 0.05.

Statistical analyses

Relative fitness of knockout strains was analysed in a linear mixed effects model using the R

package nlme. The data were Box-Cox transformed to meet model assumptions. A model was

fitted to all interactions of plasmid and host genotype. Transconjugant replicate was included as

a random effect. Model reduction was performed using stepwise AIC comparisons and

likelihood ratio tests. Effects of mutations and of plasmid carriage were estimated using post-

hoc tests with p value adjustment using a multivariate *t* distribution using the `lsmeans()`

function in the R package emmeans.

Relative fitness of plasmid mutants was analysed in a linear model with Box-Cox transformation.

Though the effect of the *lacZ* marker was found to be non-significant (one-sample t-test, t_5 =

0.32, p = 0.76), fitness values were initially corrected for marker effects by dividing each

measurement by the mean fitness of plasmid-free strains (1.01). A linear model was performed

with plasmid and marker and their interaction as fixed effects. Model reduction was performed

using stepwise AIC comparisons and F tests. We detected no interaction effect of the marker

($F_{1,25}$ = 0.03, p = 0.86). Post-hoc pairwise contrasts were performed as above.

Relative fitness of plasmid mutants was analysed in a linear model, with plasmid and knockout

and their interactions as fixed effects, with post-hoc contrasts performed as above.

Relative fitness of strains expressing PQBR57_0059 *in trans*, or *par* in trans, were each

analysed with linear mixed-effects models, with transformant as a random effect, and pUCP

variant and pQBR variant and their interactions as fixed effects. Model reduction was performed

using stepwise AIC comparisons and likelihood ratio tests, and post-hoc tests performed as

above. The experiments with pQBR103 and *par* were analysed used linear models, as

transconjugant replicates were not repeatedly-measured.

The dynamics of different PQBR57_0059 variants over time were analysed using a linear

mixed-effects model. We used Box-Cox-transformed cumulative plasmid density as the

response variable, and PQBR57_0059 status as a fixed effect. PQBR57_0059 variant was

modelled using a random effect.

Growth curves were analysed using the ratios of cumulative densities of IPTG-induced and non-

induced conditions as the response variable, and pME6032 insert as a fixed effect.

Transformant replicate was modelled as a random effect. Post-hoc contrasts were performed for

each strain against the no-insert control using the `trt.vs.ctrl` option in `lsmeans()`, with p

value adjustment using the multivariate *t* distribution.

Data and example analysis scripts are available at https://github.com/jpjh/COMPMUT/ and at

the University of Liverpool Datacat, doi: **10.17638/datacat.liverpool.ac.uk/1275**.

____

253   Results

254   Mutations to PFLU4242 and pQBR57_0059 emerged in parallel in evolution
255   experiments

Previous evolution experiments with *P. fluorescens* SBW25 and megaplasmids pQBR103,

pQBR57, or pQBR55, have identified four key genes implicated in plasmid compensatory

evolution in this strain [29,33,37,39,51] (**Fig. 1**, full details and discussion in **Supplementary**

Text): the *gacA/S* two-component signalling pathway, which has repeatedly been associated

with pQBR103 and pQBR55 carriage; *PFLU4242*, a chromosomal gene that has been mutated

in evolution experiments with all three pQBR plasmids tested; and *PQBR57_0059*, the only

plasmid-borne gene that was a target of parallel evolution in pQBR57 [52]. Parallel mutations

are a strong signal of selection [53,54], and the appearance of similar mutations in experiments

with different plasmids led us to hypothesise that mutations to *PFLU4242* and *gacS* were

general mechanisms plasmid compensation in *P. fluorescens* SBW25. Given the difficulties of

isolating wild-type pQBR55 transconjugants (i.e. without *de novo* compensation [29]), we

focused on identifying the basis of fitness cost and amelioration in the the two larger

megaplasmids, pQBR57 and pQBR103.

Figure 1. *Targets of compensatory mutation. Diagrams show the four genes associated with*

parallel compensatory mutations from Harrison et al. [33]; Hall et al. [29,39,52] and Carrilero et

al. [37]. Symbols indicate the location of the mutation in the nucleic acid sequence. The colour

of the symbol indicates the plasmid with which that mutation was associated, and the shape of

the symbol indicates the type of mutation. Amino acid-identical mutations (or identical insertion

sites) that emerged in parallel in >1 population for that plasmid are shown as a larger symbol.

The full list of mutations across these studies, as well as a discussion of how these genes were

*identified, is provided in **Supplementary Information.***

Mutations to PFLU4242, to gacS, or to pQBR57_0059 ameliorate the costs of plasmid carriage

First, to test whether *PFLU4242* and *gacS* mutations indeed had a general effect on plasmid cost, we generated pQBR57 and pQBR103 transconjugants in either wild-type bacteria or in mutant backgrounds where we had deleted *PFLU4242* or *gacS*, and measured fitness by direct competition against a plasmid-free isogenic strain (**Fig. 2A**). Knockouts were used since the diversity of naturally-arising mutations to each target (including deletions) was indicative of loss-of-function (**Fig. 1**). Consistent with our hypothesis, we detected a significant effect of mutation on the fitness costs of plasmid carriage (linear mixed-effects model (LMM), plasmid:host interaction likelihood ratio test (LRT) χ^2 = 37.3, p < 1e-6). Each plasmid levied a significant cost in the wild-type background, corresponding with previous studies [37,45]: for pQBR57, 17.8% (*t* = 4.8, p = 0.003); for pQBR103, 52% (*t* = 10.2, p < 0.001). Mutation of *PFLU4242* resulted in substantial amelioration such that we did not detect a significant fitness cost of either plasmid (post-hoc contrasts with multivariate *t* ('mvt') adjustment pQBR57 *t* = 0.87, p = 0.99; pQBR103 *t* = 1.19, p = 0.95). Mutation to *gacS* was more equivocal, with more effective compensation of pQBR103 than of pQBR57 [37]. Neither the Δ*gacS* nor the Δ*PFLU4242* mutants had a detectable fitness difference from wild-type (*gacS t* = -0.24, p = 1; *PFLU4242 t* = 0.27, p = 1), indicating that the effects of mutation were due to interaction with the plasmid rather than providing a direct fitness benefit.

Figure 2. *Relative fitness and conjugation rates of plasmid bearing strains and putative*

compensatory mutations. (A) Fitness of wild-type, ∆gacS, and ∆PFLU4242 strains of P.

fluorescens SBW25 carrying either no plasmid (left panel), pQBR57 (middle panel) or pQBR103

(right panel). All strains were competed against an isogenic, differentially-marked plasmid-free

wild-type P. fluorescens SBW25. Experiments were performed with four independent

transconjugant strains as replicates, and were performed twice (light and dark grey). Each

measurement is plotted, and the mean across all measurements for each condition indicated

with an empty black circle, with error bars corresponding to standard error. (B) Fitness of strains

with ancestral and V100A variant pQBR57 plasmids. Strains were competed against isogenic,

differently-marked plasmid-free strains. Experiments were performed with a total of twelve

independent transconjugants into either gentamicin-resistant (dark grey, n=6) or streptomycin-

resistant (light grey, n=6) recipients. Relative fitness was corrected for marker effects as

competitions were performed with both streptomycin- and gentamicin-resistant test strain

backgrounds. For panels A and B, relative fitness of 1 (horizontal line) indicate no fitness

difference between strains. (C) Conjugation rates of ancestral and V100A variant pQBR57

plasmids from wild-type, ∆gacS, or ∆PFLU4242 strains of P. fluorescens SBW25 into either

isogenic P. fluorescens, or into P. putida KT2440. Experiments were performed with four

independent transconjugant strains as replicates. Conjugation was measured according to

Simonsen et al. [55]. For all panels, each measurement is plotted, and the mean across all

measurements for each condition indicated with an empty circle, with error bars corresponding

to standard error.

To investigate the role of PQBR57_0059, we focused our attention on two sequenced, evolved

pQBR57 variants clones from an evolution experiment [39]. One of these, A.01.65.G.002 ('anc'),

was identical to the ancestor, while the other B.09.65.G.029 ('V100A') had acquired only a

single basepair mutation across its 307,330 bp length, resulting in a Val100Ala mutation in the

predicted C-terminal domain of PQBR57_0059. By conjugating the A.01.65.G.002 ('anc') and

the B.09.65.G.029 ('V100A') evolved pQBR57 variants into the ancestral strain of *P. fluorescens*

SBW25, and each of the chromosomal mutants, we could specifically measure the effects of

this mutation (**Fig. 1B**). Competitions showed that the V100A mutation largely ameliorated

plasmid cost, (ANOVA, effect of plasmid $F_{2,27}$ = 49.8, p = 8.65e-10; post-hoc contrasts with `mvt`

adjustment anc vs. V100A, p < 1e-8; plasmid-free vs. V100A, p = 0.037), but that possessing

this plasmid mutation provided no additional benefit in either the *gacS* or the *PFLU4242* mutant

backgrounds (posthoc contrasts, effect of plasmid mutation given chromosomal mutation ($|t|$ <

1.7, p > 0.74) (**Fig. S1**). We also tested 10 other naturally-emerging PQBR57_0059 variants

and likewise found that, in general, plasmids with a disrupted PQBR57_0059 allele were

significantly better at persisting (LMM, effect of PQBR57_0059 disruption χ^2 = 5.6, p = 0.02,

Fig. S2, Supplementary Text), though variance was high between evolved plasmids, probably

due to second-site mutations in pQBR57.

Supplementary Figure S1. *Relative fitness as determined by direct competition for ancestral*

and V100A variants of pQBR57, in wild-type, ∆gacS, or ∆PFLU4242 strains of P. fluorescens

SBW25, against plasmid-free wild-type P. fluorescens SBW25. Competitions were performed

with four independent transconjugants. Each measurement is plotted and the mean across all

measurements for each condition indicated with an empty black circle, with bars indicating

standard error.

Supplementary Figure S2. *Serial passage of plasmid-bearing strains in competition in soil microcosms. Panels show the relative proportions of different subpopulations over five transfers for three independent replicates (rows a, b, c). Columns correspond to different pQBR57 variants which were conjugated into an ancestral strain for this experiment. Columns are arranged according to whether that variant had a disruption in PQBR57_0059 (right-hand columns) or not (left-hand columns). All detected mutations in that variant are described at the top of the column, with numbers corresponding to the affected locus_tag (with leading zeroes removed for reasons of space). 'Tn' indicates a transposon insertion. Mutations in PQBR57_0059 are indicated in* **bold***, mutations affecting the region immediately upstream of PQBR57_0059 are in italics. Columns 1 and 2 contain data reproduced from Hall et al. [39].*

Together, these data confirm that mutation to the chromosomal gene *PFLU4242* effectively

ameliorates the cost of both pQBR57 and pQBR103 (as we previously showed both for these

plasmids [37] and for pQBR55 [29]) and similarly, that mutation to the plasmid gene

PQBR57_0059 ameliorates the cost of pQBR57. As megaplasmid cost can be ameliorated by

disruption to just one gene (*PFLU4242*, *gacS*, or *PQBR57_0059*) our results suggest that the

major burden of these megaplasmids comes not from the maintenance of plasmid DNA but

rather from an interaction that depends on the functioning of specific genes.

Plasmid compensatory evolution need not negatively affect plasmid

transmission

Next, we tested whether the different modes of plasmid compensatory evolution affected

conjugation rates. For this, we focussed on pQBR57, since both chromosomal and plasmid-

borne genes are implicated. We measured both intraspecific (between isogenic *P. fluorescens*

SBW25 strains) and interspecific (to *P. putida* KT2440) conjugation (**Fig. 1C)**. Consistent with

previous measurements made in soil [19], conjugation into *P. putida* occurred much less readily

than between *P. fluorescens* (linear model, effect of recipient $F_{1,43}$ = 925, p < 1e-8. The Δ*gacS*

mutation had a significant effect, reducing both intraspecific and interspecific conjugation

efficiency by approximately 10-fold (effect of chromosomal mutation $F_{2,43}$ = 110, p < 1e-8).

However, neither the Δ*PFLU4242* nor the PQBR57_0059_V100A mutations significantly

reduced conjugation, within or between species (p > 0.14 for both modes of compensation).

This suggests that there is no necessary trade-off between vertical and horizontal modes of

plasmid replication, and that compensatory mutations can potentially enhance both

maintenance and spread of conjugative plasmids.

Interestingly, our previous study [33] that identifed GacA/S mutation as a route to pQBR103

amelioration did not find a significant reduction in conjugation rates in the evolved lineages.

However, the lack of significance in that study was driven principally by replicates that had

mutations in PFLU4242 rather than GacA/S, consistent with our findings with the clean knockout

strains reported here.

Distinct pQBR plasmids have common and divergent effects on

chromosomal gene expression

The observation that mutations to *PFLU4242* or *gacA/S* ameliorate the costs of different pQBR

plasmids suggests that there are convergent physiological responses to the acquisition and

compensation of these different plasmids. To understand these responses and their resolution,

we performed RNAseq on plasmid-free bacteria, and from bacteria carrying either pQBR57 or

pQBR103, either without amelioration mutations, or with the Δ*PFLU4242* or Δ*gacS* knockouts,

or the PQBR57_0059 V100A mutation.

First, we analysed the effect of plasmid acqusition without amelioration. The different plasmids

had some common and some divergent effects on chromosomal gene expression (**Fig. 3, Fig.**

S3). pQBR57 affected the expression of 398 genes (FDR<0.05), with 47 upregulated >2x and

60 downregulated >2x. pQBR103 affected expression of 254 genes, with 77 upregulated >2x

and only 11 downregulated >2x. GO terms enriched amongst the upregulated genes for both

plasmids included those associated with the SOS response and signal transduction. However

there were divergent effects of each plasmid, particularly for 33 genes which were upregulated

by pQBR103 but downregulated by pQBR57. Carbohydrate metabolism was overrepresented in

this set (7/33 GO:0005975 "carbohydrate metabolic process," p_{adj} = 0.008) suggesting that while

there was a common response to megaplasmid acquisition, each plasmid also had specific

effects.

Figure 3. *Common and divergent effects on chromosomal gene expression following plasmid*

acquisition and amelioration. Each column in the heatmap refers to a differentially-expressed

(FDR<0.05, absolute log₂ fold-change > 1) gene following acquisition of either plasmid. Note:

there were no additional genes meeting these criteria in the ∆PFLU4242 or

PQBR57_0059_V100A ameliorated strains, however, there were 401 additional differentially-

expressed genes as a result of gacS deletion, which have not been shown for clarity. Columns

have been grouped by pattern of response to pQBR57 and pQBR103. A coloured symbol below

each gene indicates whether it is associated with a chromosomal MGE and/or the SOS

response. Red indicates increase and blue indicates decreases in expression; cells with a solid

border had FDR<0.05 for that specific condition. Genes selected for cloning are labelled.

Supplemental Figure S3. *Volcano plots showing changes to gene expression following*

plasmid acquisition. Points are coloured according to Figure 4. PFLU1169 is highlighted with a

black border.

A closer examination of the differentially-expressed genes shows a set of 50 genes that were at

least two-fold upregulated by both plasmids. Many of these were co-localised in the genome,

indicating co-regulation, and the possible activation of particular operons by pQBR103 and

pQBR57. Of these 50 genes, 31 were in putative prophages, while 3 were in the Tn6291

transposon [39]. Since prophage induction is a signature of the SOS response, and the *P.*

fluorescens SBW25 *lexA* homologues *PFLU1560* and *PFLU3605* were also upregulated by both

pQBR57 and pQBR103, we investigated how many of the 16 remaining upregulated genes

were located downstream of a predicted LexA binding site. By scanning the genome for the

conserved pattern `CTGKMTNNWHDHHCAG` [56] we identified 12 genes <151 bp dowstream of a

predicted LexA site, of which 5 were upregulated by both plasmids. The common transcriptional

response to pQBR57 and pQBR103 acquisition by *P. fluorescens* SBW25 can thus be

characterised as activation of the SOS response and chromosomal mobile genetic elements,

which constitute 42/50 (84%) of the >2-fold upregulated genes. This is not the same as the

transient SOS response caused by plasmid acquisition reported by Baharoglu et al. [57], as the

strains used in these experiments are many generations removed from the original

transconjugant due to periods of growth during colony isolation and culture.

Plasmid genes were transcriptionally active. pQBR103 carries 478 predicted ORFs, while

pQBR57 carries 426; for both plasmids the majority of ORFs have unknown function. In the

ancestral host, expression of pQBR103 ORFs ranged from 0.9–2323 transcripts per million

(mean 87, median 39), while expression from pQBR57 ranged from 1.8–2532 (mean 132,

median 43). In general, plasmid-borne genes were expressed at levels approximately 2x higher

than those of the chromosome, after accounting for gene length (**Fig. 4**). In both cases, the

most highly-expressed genes from the plasmids were uncharacterised proteins (**Fig. S4**).

447
Figure 4. *Megaplasmid genes are expressed, and overall levels of expression remain the same*

regardless of amelioration. Violin plots show log-transformed transcripts-per-million (TPM) of

chromosomal (green) and plasmid (orange) genes, averaged across experimental replicates.

Solid dots indicate the grand mean for that replicon and condition, calculated from log-

transformed values.

Supplementary Figure S4. *Gene expression varies across the plasmids, with the most highly-*

expressed genes outwith regions with predicted functions. Heatmaps show level of expression

in transcripts-per-million (TPM) across each plasmid under different amelioration conditions.

Coloured bars across the top indicate regions annotated in Hall et al. [45]: grey = rep; green =

che; orange = par; purple = pil; yellow = SAM; red = Tn5042 mercury resistance transposon;

pink = uvr; blue = tra. Loci are numbered according to the `locus_tag` *feature.*

Plasmid amelioration has broadly similar effects on transcription regardless of mechanism

We then investigated how compensatory mutations affected the changes in transcription caused by plasmid acquisition.

Effects of PFLU4242 on plasmid cost and amelioration

Without pQBR57 or pQBR103, disruption of *PFLU4242* had no detectable effect on chromosomal gene expression compared with wild-type bacteria ($p_{adj} > 0.05$ for all genes). However this same mutation in plasmid-bearing bacteria had a striking effect on plasmid-induced changes in gene expression: the reversion to ancestral levels of almost all (pQBR57: 376/398, 94%; pQBR103: 251/254, 99%) of the chromosomal genes differentially regulated following plasmid acquisition (**Fig. 3**), including all of the SOS-associated genes upregulated by both plasmids, and the majority of plasmid-specific differentially-expressed chromosomal genes.

Plasmid genes were also differentially expressed following amelioration by PFLU4242 mutation, but only a subset of them (pQBR57: 57/426; pQBR103: 85/478). The responses of pQBR57 and pQBR103 were largely divergent, with PFLU4242 disruption generally resulting in downregulation of pQBR57 genes, particularly those encoding putative conjugation apparatus (*tra*) and a type IV pilus (*pil*) (**Fig. 5**), while pQBR103 genes were upregulated by PFLU4242 disruption (including *tra* and *pil*) (**Fig. S6**). Though modulation of plasmid gene expression is likely to have some effect on cell physiology, these divergent responses lead us to conclude that the effects of plasmid transcriptional burden or plasmid gene expression *per se* are not primarily responsible for the costs of these plasmids, since the phenotypic costs of carriage could be abolished by *PFLU4242* disruption without major convergent effects on plasmid gene expression.

Figure 4. *Common and divergent effects on pQBR57 gene expression following different modes of compensatory evolution. (A) Heatmap of differentially-expressed (FDR<0.05) genes on pQBR57 following either chromosomally-located amelioration via ΔPFLU4242, or plasmid-located amelioration via the V100A mutation to PQBR57_0059. A coloured symbol below each gene indicates the predicted plasmid region affected. Red indicates increase and blue indicates decreases in expression; cells with a solid border had FDR<0.05 for that specific condition. (B) The region upstream of the predicted pQBR57 par genes. Predicted transcription and translation factor binding sites are highlighted, as are identified repeats, and mutations identified by Hall et al. [39]. TSS = Transcription Start Site; RBS = Ribosome Binding Site.*

Notably, expression of the three genes from Tn6291 was unaffected by PFLU4242 mutation

amelioration for either plasmid. These genes (*PFLU1886-1888*) are hypothesised to encode the

transposase subunits for Tn6291. The fact that these genes remain upregulated in the

compensated strains suggests that these are not causally responsible for the main negative

effects of plasmid acquisition, and also suggests that they are activated through a different route

to the other plasmid-upregulated genes.

PFLU4242 mutation therefore had the general effect of bringing chromosomal gene expression

back to levels similar to plasmid-free bacteria, suggesting wild-type *PFLU4242* is instrumental in

interacting with pQBR plasmids to generate a fitness cost.

Effects of *gacS* on plasmid cost and amelioration

Compared with *PFLU4242*, unpicking the specific transcriptional effects of *gacS* deletion on

plasmid carriage was more difficult, as *gacS* deletion by itself caused differential expression of

733 chromosomal genes affecting numerous phenotypes (see also [58]). One hypothesis was

that disruption of GacA/S signalling ameliorates the cost of plasmid carriage by independently

downregulating genes that were upregulated by the plasmids, thus 'balancing out' the effects of

plasmid acquisition. However, while we found a number of genes that were differentially

expressed in the plasmid bearers and the Δ*gacS* mutant, the direction of exprssion varied

between pQBR57 and pQBR103, and of the 50 genes upregulated by both plasmids (logFC>0,

FDR<0.05) only 6 were independently downregulated by the Δ*gacS* mutation, providing little

support for this hypothesis.

We therefore considered an alternative hypothesis: that there exists an interaction between

GacA/S signalling and the pQBR plasmids that is responsible for plasmid cost, and disruption of

the GacA/S system prevents this interaction, resulting in amelioration. One pathway for an

interaction between GacA/S and the plasmids could be through PFLU4242. The GacA/S system

operates by interfering with the activity of inhibitory mRNA-binding Rsm proteins, causing

translation. Other studies have identified putative RsmA binding sites in *PFLU4242*, suggesting

that *PFLU4242* may form part of the GacA/S regulon [[59]; Jenna Gallie, pers. comm.]. Indeed,

we found that in the ∆*gacS* mutant, *PFLU4242* transcription was reduced by approximately

50%, but *gacS* expression was not affected in the ∆*PFLU4242* mutant (**Fig. S5**). These data

provide an explanation for some of the convergence between these two mutational routes to

compensation, namely, that PFLU4242 is (more) proximally responsible for plasmid cost, and

that disruption to GacA/S could reduce plasmid cost by reducing expression of PFLU4242.

Consistent with this hypothesis, and with the phenotypic fitness data, GacA/S disruption

affected plasmid-induced changes in gene expression by reverting some of the differentially-

expressed genes back to ancestral levels, but this effect was incomplete, particularly with

pQBR57 (**Fig. 3**). The fact that PFLU4242 is expressed under wild-type plasmid-free conditions

also suggests that PFLU4242 by itself is not toxic, and it is interactions with the megaplasmids

that underlies the fitness consequences of this gene.

Supplementary Figure S5. *Expression of PFLU3777 (gacS) and PFLU4242 in the different*

treatments. Note different y-axis scaling. Points indicate different replicates, with values for each

replicate scaled by effective library size.

Supplementary Figure S6. *Differentially-expressed plasmid genes across all amelioration conditions. Image as **Fig. 4A**, but showing also the effect of ∆gacS and differentially-expressed genes on pQBR103.*

Disruption of GacA/S signalling affected gene expression from both plasmids in a manner

similar to PFLU4242 disruption, but there was an additional effect on pQBR57 which showed 26

genes, mainly of unknown function, specifically downregulated in the Δ*gacS* mutant relative to

the wild-type (**Figs. S4, S6**). There was no overall pattern in the predicted functions of these

genes, other than their location near the putative origin of replication, and start of the sequence,

of pQBR57.

The overlap between plasmid and gac-regulated genes likely represents a complex interaction

between the plasmids and the gac system. However, the direct effect that GacA/S has on

PFLU4242 transcription can effectively provide a unified explanation for the two chromosomal

modes of amelioration, converging on PFLU4242.

Effects of PQBR57_0059 on plasmid cost and amelioration

Mutations to PQBR57_0059 had a broadly similar effect on chromosomal gene expression to

PFLU4242 (**Fig. 3**). Each amelioration had similar effects on gene expression from pQBR57

(**Fig. 5**), though the effects of the chromosomal mutation were clearer, with many genes

trending in a similar direction with the plasmid-borne mutation but not exceeding the FDR<0.05

significance threshold. One clear exception were the *par* genes, which were upregulated by the

PQBR57_0059_V100A mutation but not by the ΔPFLU4242 mutation (see below).

A mechanism for plasmid cost and its amelioration

Together, our transcriptomic data indicates a set of genes that are upregulated by both

plasmids, and then downregulated by the different pathways of compensatory evolution.

Expression of these genes therefore correlates with the phenotypic fitness costs of plasmid

carriage. These genes may be responsible for physiological disruption and fitness costs, but it is

also possible that their expression is a downstream effect of other disruptive interactions

emerging as a consequence of plasmid acquisition. To test whether expression of these genes

causally affects bacterial fitness, we selected 12 candidates for further investigation. Eight

candidates were selected as genes which were >2x upregulated by both plasmids in the wild-

type but not the Δ*PFLU4242* background, and four additional genes that did not meet these

criteria were also selected as controls (**Fig. 3**). The ORF of each gene was cloned into the

inducible *Pseudomonas* expression vector pME6032 [60], and the effects of expression on

growth were tested by comparing growth curves between induced and uninduced cultures (**Fig.**

5).

Figure 5. *Expression of genes upregulated by megaplasmid plasmid acquisition imposes a*

fitness cost. Different panels indicate cloned genes. Each vector was transformed into P.

fluorescens three times independently, and the growth of each transformant was tested with and

without IPTG induction four times, with the mean (solid line) and standard error (shaded area)

calculated and plotted for each transformant separately, under 100 µg/ml IPTG (red) or

uninduced (blue) conditions. One of the three pME6032::PFLU1169 transformants did not show

a large growth defect, but resequencing this plasmid from P. fluorescens revealed a 16-bp

deletion in the insert in this replicate. No other mutations were identified in the other replicates,

nor in any of the other inserts.

We identified one gene, *PFLU1169*, which, when expressed, resulted in a considerable

reduction in growth representative of a large fitness cost (LMM effect of insert χ^2 = 49.2, p <

0.0001; Dunnet's post-hoc test *PFLU1169* vs. empty vector t_{26} = 5.36, p < 0.0001). Indeed, this

gene was difficult to clone and express as growth in culture imposed strong selection for

mutation of the insert or loss of the expression vector. *PFLU1169* is part of the *P. fluorescens*

SBW25 'prophage 1' locus, however, further investigation revealed that this prophage is in fact

a tailocin: a protein complex related to a phage tail which acts as a bacteriocin against

competitors [61]. *PFLU1169* encodes the holin, expression of which, in concert with other

prophage genes, causes cell lysis and tailocin release [62]. The prophage 1 locus is itself

regulated through the SOS response, as a LexA binding site was found upstream of this operon.

PFLU1170, another 'prophage 1' gene, had a more marginal effect (*PFLU1170* vs. empty vector

t_{26} = 3.2, p < 0.03). Overexpression of any of the remaining 10 genes had no detectable effect

(p > 0.99 for all comparisons) under the tested conditions.

We have therefore identified one causal link between plasmid-induced gene expression

changes and fitness costs. Plasmid acquisition activates the SOS response, which activates the

'prophage 1' tailocin, causing increased cell permeability and lysis. Lysis is likely to be

stochastic, the result of which is an apparent reduction in fitness at the population level.

pQBR57_0059 interacts with other plasmid genes to generate a fitness cost

PQBR57_0059 is predicted to encode a DNA-binding protein of the lambda-repressor family

(HHPRED top hit 3BDN 'lambda repressor,' p = 4E-23, [63]). We therefore considered it unlikely

that PQBR57_0059 was a toxin that directly interfered with cellular physiology. Instead, we

predicted that it affected the expression of other genes, either on pQBR57, on the chromosome,

or both, resulting in a negative effect on competitive fitness. To distinguish between these

alternatives, we cloned either the ancestral-type ('anc') or mutant ('V100A') variants of the

PQBR57_0059 open reading frame (ORF) under the control of the P_{lac} promoter, which is

615	constitutively active in *Pseudomonas*, on the expression vector pUCP18 [41] and introduced
616	these variants to *P. fluorescens* SBW25. We then introduced either the evolved variant of
617	pQBR57 with the V100A mutation, or the unmutated variant. If the costs associated with wild-
618	type PQBR57_0059 were due to direct interaction with the cell, or solely through effects on
619	chromosomal gene expression, fitness costs would be apparent by expressing the ancestral
620	type variant from pUCP18. However if other megaplasmid genes were involved, then pQBR57
621	would be required to recapitulate the effect, and the wild-type *PQBR57_0059* would
622	complement the otherwise costless V100A pQBR57 variant.

623	Expressing just the wild-type PQBR57_0059 from pUCP18 was insufficient to generate a fitness
624	cost (**Fig. 6**; LMM, post-hoc tests for pUCP18 variants without pQBR57 p > 0.91 for all
625	comparisons), and other genes on pQBR57 were required for PQBR57_0059 to be costly
626	(interaction between pUCP18 and pQBR57 variant χ^2 = 29.4, p = 1e-4). In fact, expression of
627	PQBR57_0059 from pUCP18 appeared to decrease the fitness of pQBR57-bearers even further
628	(effect of anc vs. no insert p = 0.0078), probably due to the increased effect of higher
629	PQBR57_0059 expression from the multi-copy expression vector. Therefore, the fitness effects
630	of PQBR57_0059 are likely to emerge from its effects on other plasmid genes, hinting that a
631	more complex interaction between the plasmid and chromosome is responsible for the cost of
632	pQBR57.

Figure 6. *Relative fitness as determined by direct competition for P. fluorescens SBW25*

overexpressing either the ancestral or the V100A variant of PQBR57_0059 (or no-insert

expression vector, or no-vector controls), and carrying either no pQBR57 or the ancestral or

V100A variants of pQBR57. Competitions were performed with three independent

transformants/transconjugants indicated by shades of grey, and the fitness of each

transconjugant/transformant was tested four times. For the 'none/no-pQBR57' control treatment,

all measurements were taken on the same strain.

We therefore investigated plasmid-borne genes that were specifically differentially-expressed by the PQBR57_0059 amelioration. Amongst the plasmid genes differentially expressed by amelioration, only three were specifically affected by pQBR57_0059 (**Fig. 4A**). One, PQBR57_0215, was down regulated in a similar direction by the $\Delta PFLU4242$ mutation ($0.05 <$ $p_{adj} < 0.1$) and thus is likely to be a general response to amelioration. However the other two, which form an operon PQBR57_0054-0055, were exclusively upregulated by pQBR57_0059 disruption. These genes are predicted to encode a ParAB system responsible for efficient segregation of plasmids to daughter cells (HHPRED top hit for PQBR57_0055 6SDK 'Spo0J/ParB' p = 2E-24; PQBR57_0054 matches 2OZE_A 'ParA/Walker-type ATPase' p = 4.2E-25). Examination of the region upstream of this operon reveals three ATTGC repeats and an inverted repeat (CTTGA(N)$_9$TCAAG) that might interact with pQBR57_0059, which is predicted to bind DNA (**Fig. 4B**). Interestingly this region was also the target of parallel mutations in a previous evolution experiment [39]: in one population, evolved pQBR57 carried by *P. putida* had acquired an insertion of Tn6291 transposon 78 bases upstream from the predicted transcriptional start site, while in another replicate, evolved plasmids carried both by *P. fluorescens* and *P. putida* gained a single T>C transition 38 bases upstream of the transcriptional start site. Together this suggests that pQBR57_0059 acts as a repressor of the *par* locus, and disruption of this repressor, or disruption of its binding site, results in increased transcription. The fact that there is no additive benefit of mutations in PFLU4242 and pQBR57_0059 suggests that increased par gene expression somehow prevents a deleterious interaction with PFLU4242, directly or indirectly, to ameliorate plasmid cost.

Increased expression of the *par* genes from pQBR57 is sufficient to ameliorate the costs of pQBR57, but not pQBR103

To test the hypothesis that increased *par* gene expression as a consequence of the disruption of the repressor PQBR57_0059 was responsible for ameliorating the costs of pQBR57, we

cloned the putative *par* genes PQBR57_0054-0055 into pUCP18 and introduced this vector, or

an empty pUCP18 control, into *P. fluorescens* SBW25. We then introduced pQBR57, and

measured the effects on relative fitness. Consistent with our hypothesis, overexpression of the

par genes ameliorated the costs of pQBR57 (**Fig. 7A**; LMM, interaction of pUCP18 insert and

pQBR57 carriage χ^2 = 22.2, p < 1e-4; pairwise contrasts no insert pQBR57 vs. *par* pQBR57 t_{10}

= 5.15, p = 0.004), but had no significant fitness effects by themselves (no insert no pQBR57

vs. *par* no pQBR57 t_{10} = 1.82, p = 0.5).

Figure 7. *Effects of pQBR57 par gene expession on megaplasmid cost. **(A)** Competitions were performed with three independent transformants/transconjugants indicated by shades of grey, and the fitness of each transconjugant/transformant was tested four times. Empty circles indicate the grand mean for that set of conditions. **(B)** Expression of pQBR57's par genes in trans exacerbates the fitness costs of the distantly-related megaplasmid pQBR103. Competitions were performed with five independent transconjugants, with each one tested once.*

We then investigated whether overexpression of pQBR57's *par* genes had a general effect on

plasmid fitness costs. We introduced pQBR103 to the strains that were overexpressing

PQBR57_0054-0055 and again measured relative fitness. In contrast to the amelioration of

pQBR57, expression of PQBR57_0054-0055 had no beneficial effect on pQBR103 (LM,

pUCP18:pQBR plasmid interaction $F_{2,24}$ = 1.79, p = 0.18; pairwise contrast pQBR103 no insert

vs. pQBR103 *par* t_{26} = 2.41, p = 0.12), and we had weak evidence that costs were in fact

exacerbated (effect of pUCP18 insert $F_{2,26}$ = 4.4, p = 0.02). Increased expression of pQBR57

par therefore reduced the fitness costs of pQBR57, but did not have any ameliorative effect on

an heterologous plasmid.

Discussion

Plasmids are known to impose fitness costs, but the molecular causes of such costs are far

from clear. Here we show that, rather than emerging from the general bioenergetic demands of

replicating DNA, transcribing into mRNA, or translating into protein [20], the majority of the

fitness costs of plasmid acquisition come from specific deleterious interactions between plasmid

and chromosomal genes. As a consequence, small mutations — in some cases just single base

changes — can be sufficient to ameliorate even a very large plasmid with hundreds of genes.

Previous studies have tended to focus on smaller, non-conjugative replicons [25,30,64], or have

identified plasmid compensatory mutations with substantial pleiotropic effects: disruption or

deletion of conjugative machinery from the plasmid (rendering it immobile) [28,65]; deletion of

large portions of the plasmid [34,66]; or disruption of extensive and multi-functional regulatory

circuits [31,33,67]. In contrast, our data shows that single mutations can enable large, natural,

conjugative, plasmids to become rapidly accommodated with few obvious tradeoffs, and

furthermore describes the molecular negotiation by which this occurs (**Fig. 8**).

Figure 8. Interactions leading to pQBR plasmid fitness costs. Dotted lines indicate speculative

interactions, solid lines indicate experimentally- or bioinformatically-determined interactions

(from this study or others). Targets of compensatory mutation are shown in blue. Genes are

solid boxes whereas pathways/groups of genes are outlines. (A) Both plasmids interact with

PFLU4242. (B) PFLU4242 activates the SOS response, probably through causing DNA breaks.

(C) One outcome of the SOS response is expression of PFLU1169 which directly causes cell

damage. (D) The effects of plasmid interactions with PFLU4242 are exacerbated by GacA/S,

which increases expression of PFLU4242. pQBR103 may also interact directly with the GacA/S

system. (E) Mutation of PQBR57_0059 increases expression of par, which reduces the

disruption of pQBR57 but not pQBR103. (F). Both plasmids activate the Tn6291 transpose

through a different route to PFLU4242 or GacA/S, which is likely to increase transposition

activity and mobilisation of Tn6291. Overall, specific gene conflict costs greatly exceed the

metabolic costs of pQBR57 and pQBR103.

──

Specifically, we have linked the fitness costs of both pQBR57 and pQBR103 to the SOS

response in *P. fluorescens* SBW25. SOS response-associated genes comprised 42/50 of the

genes upregulated in common, and we found that increasing expression of just one gene in the

SOS regulon — the 'prophage 1' holin *PFLU1169* — had a severe effect on fitness, implicating

the SOS response as a direct proximal cause of plasmid-induced fitness costs. The SOS

response has wide-ranging impact on bacterial physiology [68], meaning that *PFLU1169* is

unlikely to be the *sole* mechanism of plasmid-induced fitness costs. How is the SOS response

triggered by the pQBR megaplasmids? Single-stranded DNA, such as is acquired in the early

stages of conjugative transfer, can directly activate the SOS response [57], but as the strains in

this study were many generations removed from the original conjugation event due to picking

and restreaking of transconjugant colonies, the SOS response in the pQBR/SBW25 system is

clearly being perennially activated through a different route. DNA damage is the most likely

candidate, implicating the SOS response as essentially an amplifier of disruptions caused by

plasmid acquisition. Like an amplification circuit, the SOS response can itself cause DNA

fragmentation, forming a positive feedback loop, and further disruption [69].

How does plasmid carriage cause DNA breaks? Clashes between the machineries of

transcription and replication put stress on DNA molecules, and such clashes have been shown

to drive the fitness costs of the minimal (2.1 kb) plasmid pIND in *E. coli* [64]. In *Shewanella*

oneidensis, the replication machinery of the IncP1-beta mini-replicon pMS0506 (13.1 kb)

probably triggers the SOS response by binding to the host DNA helicase DnaB and blocking the

| replication fork, exposing vulnerable ssDNA intermediates to breaks [70]. In *P. aeruginosa*

| PAO1, the small (5.1 kb) plasmid pNUK73 triggered the SOS response due to increased

| expression of the plasmid replication machinery, which may cause breaks due to increased

| stress on DNA molecules within the cell [25]. However, other costly plasmids did not have the

| same effect on *P. aeruginosa* PAO1, suggesting that SOS response activation was not a

| general source of plasmid fitness costs in that strain [26]. The fact that different species of

| *Pseudomonas* had differing SOS responses to the same pCAR1 metabolic plasmid [71] likewise

| suggests that activation of the SOS response usually occurs in a plasmid-by-host specific

| manner, again consistent with our hypothesis that plasmid fitness costs emerge principally

| through specific genetic conflicts.

| It is unclear whether the conflicts between transcription and replication seen with small plasmids

| is responsible for DNA damage for much larger plasmids like pQBR103 and pQBR57. We saw

| no significant change in megaplasmid copy number [39] or *rep* gene expression when

| comparing ancestral with ameliorated strains, hinting that other processes may be at work.

| What is clear is that mutations to a limited number of targets can prevent activation of the SOS

| response, presumably by preventing DNA breaks from occurring. Principal amongst these

| mutations, in terms of degree and scope of amelioration, are loss-of-function mutations to

| PFLU4242. While the molecular and evolutionary functions of PFLU4242 remain unknown, its

| DUF262 domain pattern resembles that of known endonucleases that target modified phage

| DNA [72], and related genes are located in known genome defence islands [73]. The hypothesis

| that PFLU4242 has an nuclease function in phage defence would explain how PFLU4242

| disruption might prevent DNA breaks, and also why this gene has been maintained despite

| having no detectable effect on fitness or gene expression in plasmid-free bacteria in an axenic

| laboratory setting. Work is ongoing to understand the molecular and evolutionary functions of

| this gene. The concept of a trade-off between openness to HGT and resistance to phage

infection has been explored extensively in the context of CRISPR-Cas and restriction-

modification modules [14,74,75]. Our data hints at another mechanism by which genome

defence might have side effects: an inappropriately excessive response driving fitness costs,

analogous to immune hypersensitivity.

We show that PFLU4242 expression is induced by a global regulator of secondary metabolism,

the GacA/S two-component system, and so our data is consistent with the hypothesis that

disruption to GacA/S could exert its effect through repression of PFLU4242. It is possible that

global regulators have been implicated in plasmid persistance due to their regulatory effects on

more proximal causes of plasmid conflict [31,67]. The combined size of the *gacA* and *gacS*

genes (~3.4 kb) is over twice that of *PFLU4242* (~1.6 kb) (**Fig. 1**), providing a larger mutational

target, which, alongside the mutability of the GacA/S system [76], may explain why GacA/S

rather than PFLU4242 was the primary target of mutation in one of our previous studies [33].

However, we have hints that PFLU4242 is not the only mechanism by which pQBR plasmids

and the GacA/S system interact. Acqusition of pQBR103 has a similar effect on PFLU4242

expression as deletion of *gacS*, while 26 pQBR57 genes appear to be differentially regulated in

the Δ*gacS* background but not in the Δ*PFLU4242* background. Moreover, pQBR103 appears to

carry a homologue of the Gac system signal transducer RsmA, in common with pQBR55 but not

pQBR57 [45]. Understanding how plasmids may manipulate core pathways in bacterial

signalling is an important subject for future investigation [77].

Compensatory mutations occuring on a plasmid are thought to have increased potential to

enhance plasmid persistence and accelerate horizontal gene transfer relative to mutations of

similar effect on the chromosome, because plasmid-borne compensation is inherited both by

daughter cells and by transconjugants [35]. Mutation of the putative lambda-family repressor

PQBR57_0059 caused increased expression of the putative partitioning machinery of pQBR57,

reducing the cost of that plasmid. The molecular mechanism by which *par* expression reduced

pQBR57's cost remains unclear. Chromosomal ParB is known to bind DNA both specifically (at

parS sites) and non-specifically, impeding access by DNA-damaging agents like nucleases,

forming bridges and loops between distant parts of a DNA sequence, and affecting gene

expression [78–83]. The putative ParB of pQBR57 may also act to prevent DNA damage,

perhaps by restricting nuclease access to otherwise vulnerable DNA. What, then, is the

evolutionary function of PQBR57_0059, in repressing the protective *par* module? Tuning *par*

expression for plasmid cost reduction may be a delicate matter, as our data suggests that high

levels of pQBR57 *par* expression alone has a cost (as has also been found for chromosomal

ParB expression [79,82]), and may exacerbate the costs of heterologous plasmids as we see for

pQBR103. A highly conjugative plasmid such as pQBR57 is expected to frequently find itself in

co-existence with other plasmids [84], and PQBR57_0059 may be one mechanism by which

pQBR57 can modulate its own activity in multi-replicon cells. Other plasmids are known to carry

genes that prevent cellular disruption, for example the anti-SOS genes PsiB [85], or the

nucleoid-like proteins H-NS that repress plasmid gene expression [86], reducing costs. In this

context, it is interesting that we have witnessed the evolution of this 'stealth' function for the

pQBR57 *par* genes during the course of our evolution experiment, showing that plasmids can

contain the genetic resources necessary to evolve their own amelioration.

The principal group of genes upregulated by both plasmids were located in chromosomal

regions annotated as mobile genetic elements. Foremost amongst these were prophages 1 and

3. However, the prophages of SBW25 are not predicted to be intact integrated phage, and

rather seem to be relics which have lost the genes involved in DNA replication and packaging

[62,87,88]. These prophages are therefore candidate tailocins — phage tails that have been

evolutionarily co-opted to act as toxins used in inter-strain competition [61]. To exert their

activity, tailocins translated in the cytoplasm must be released by cell lysis [61]. Tailocin

production is therefore an example of bacterial spite, whereby an individual suffers a fitness cost

to harm a competitor [89]. Spite, like altruism, is evolutionarily stable only within a narrow range

of conditions — specifically, where those benefitting from the action are likely to be related, and

those targeted by the action are unlikely to be related [90]. Receiving a new plasmid is a clear

signal of the presence of an unrelated competitor, and thus an opportune time for spiteful

tailocin activation. The SOS response can be considered a strategy against aggressions, rather

than a solely defensive reaction [68]. It is therefore tempting to speculate, that, rather than being

solely a side effect of DNA damage, expression of tailocins following plasmid acquisition could

be an adaptive trait, that both inhibits competitors [90] and potentially also prevents invasion of

parasitic plasmids through cell suicide in a manner similar to abortive infection [91].

The other chromosomal mobile genetic element upregulated by both pQBR57 and pQBR103

was the transposon Tn6291. The upregulated genes are thought to encode the transposase,

which provide the machinery enabling Tn6291 to relocate both to other chromosomal locations

and, importantly, to conjugative plasmids. Tn6291 does not appear to encode conjugative

machinery of its own, and therefore depends on other apparatus to move horizontally between

bacterial lineages. Indeed, we previously showed using experimental evolution that Tn6291

exploits pQBR57 to move between *Pseudomonas* species in soil [39]. This dependence of

Tn6291 on other mobile genetic elements for transmission provides an adaptive explanation for

why the transposase genes are sensitive to conjugative plasmid acquisition: the presence of a

conjugative plasmid signals the presence of a vehicle for Tn6291 to use. Communication

between the pQBR plasmids and Tn6291 appears to be independent of the SOS response, as

the Tn6291 transposase remains upregulated in the otherwise ameliorated plasmid-bearing

strains. There are numerous adaptive reasons why mobile genetic elements might be expected

to communicate with one another, since conflict and collaboration between mobile genetic

elements is expected to be a key factor influencing horizontal gene transfer and hence bacterial

evolution [e.g. 2,92]. Interpreting this language could prove key to understanding the factors

driving the spread of ecologically- and clinically-important traits.

Our work has focused on megaplasmids: large plasmids, generally exceeding 150 kb in size.

Megaplasmids are likely to be especially effective vectors of horizontal gene transfer as they

can mobilise many traits at once — a particular concern for the spread of antimicrobial

resistance because selection with one drug may drive the spread of multi-drug resistance [93].

Similarly, compensation of megaplasmids potentially facilitates the maintenance of many

different traits in a lineage, provided that (consistent with our findings here) costs do not emerge

from plasmid size *per se*. Advances in long-read sequencing technologies are beginning to

expose the diversity, ubiquity, and complexity of megaplasmids in diverse bacterial genera, and

therefore our findings are likely to apply beyond this specific case of mercury resistance in a soil

Pseudomonad [e.g. 94,95]. For example, megaplasmids pQBR103 and pQBR57 are distant

relatives of a large family of clinically important plasmids that appear to be disseminating multi-

drug resistance in clinical isolates of *Pseudomonas aeruginosa* from opportunistic infections

[96,97]. In those strains, megaplasmids appear to be stable even without the selective pressure

of antimicrobials, suggesting that those strains are either pre-adapted for megaplasmid carriage,

or have undergone compensatory evolution.

The fact that plasmid costs emerge principally from specific gene conflicts is not to say that

replication and expression of megaplasmid genes *per se* levies no burden, nor that such costs

are invisible to selection. Lynch and Marinov [20] calculated the costs of bacterial gene

replication, transcription, and translation, and found that the costs of even replicating short

pieces of non-beneficial DNA are sufficient for purifying selection in moderately-sized bacterial

populations, with the costs of transcribing and translating those genes several orders of

magnitude greater. The assays we used to measure fitness have limits, and it is likely that the

residual bioenergetic burden of plasmid carriage would be detected by more sensitive methods

[98]. Nevertheless, it is clear that such bioenergetic costs are many fold smaller than those that

emerge from conflicts between specific genes or their products, and may well be overwhelmed

by other selective pressures in natural environments (such as phage [99]) or rendered negligible

in the context of horizontal gene transfer [100].

Our findings have important implications for understanding the evolution of bacterial genomes.

Fitness costs of plasmid acquisition are, in theory, a barrier to HGT, however, our data suggest

that when such costs are caused by specific genetic conflicts they will be readily and rapidly

ameliorated by single compensatory mutations thus enabling the long-term maintenance of

plasmids in bacterial genomes. This helps to explain why plasmids are so common in bacterial

genomes and, moreover, suggests that such compensated plasmid-carrying lineages will

become important hubs of HGT within bacterial communities [19,101]. The fact that plasmid

costs depend on specific conflicts is relevant to understanding the dissemination of ecologically

and clinically important bacterial traits, such as antimicrobial resistance, because the immediate

fitness costs of the encoding plasmids are unlikely to predict their long-term dynamics: purifying

selection against plasmids will diminish over time due to rapid compensatory evolution. More

generally, the dynamics of mobile elements in bacterial genomes are unlikely to be predictable

from their generic properties, and thus more sophisticated analyses of genetic associations in

pangenomes [e.g. 102] will be required to predict the patterns and outcomes of gene exchange

and how these will shape pangenome dynamics.

____

Author contributions

James P. J. Hall: Funding acquisition, Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Visualization, Writing — Original Draft Preparation, Writing — Review & Editing

Rosanna C. T. Wright: Resources, Investigation, Writing — Review & Editing

Ellie Harrison: Funding acquisition, Conceptualization, Writing — Review & Editing

Katie Muddiman: Resources

A. Jamie Wood: Funding acquisition, Conceptualization, Writing — Review & Editing

Steve Paterson: Funding acquisition, Supervision, Conceptualization, Writing — Review & Editing

Michael A. Brockhurst: Funding acquisition, Conceptualization, Project Administration, Writing — Review & Editing

Acknowledgements

We would like to thank staff at the NERC Biomolecular Analysis Facility, University of Liverpool, for assistance with the RNA-seq, and Jenna Gallie (Max Planck Institute for Evolutionary Biology) for a list of putative RsmA targets in *P. fluorescens* SBW25.

Funding

This work was supported by funding from NERC [to M.A.B., S.P., A.J.W., J.P.J.H. NE/R008825/1; to M.A.B., A.J.W. NE/K011774/1], an ERC Consolidator Grant [to M.A.B.; 11490-COEVOCON], an Academy of Medical Sciences Springboard Award [to J.P.J.H.;

908   SBF005\1062], and funding from the Institutional Strategic Support Fund (ISSF) awarded by

909   Wellcome Trust via the University of Liverpool [to J.P.J.H.; 204822/Z/16/Z].

References

1. Weingarten RA, Johnson RC, Conlan S, Ramsburg AM, Dekker JP, Lau AF, et al. Genomic analysis of hospital plumbing reveals diverse reservoir of bacterial plasmids conferring carbapenem resistance. MBio. 2018;9. doi:10.1128/mBio.02011-17

2. Sheppard AE, Stoesser N, Wilson DJ, Sebra R, Kasarskis A, Anson LW, et al. Nested russian Doll-Like genetic mobility drives rapid dissemination of the carbapenem resistance gene blaKPC. Antimicrob Agents Chemother. 2016;60: 3767–3778. doi:10.1128/AAC.00464-16

3. Peter S, Bosio M, Gross C, Bezdan D, Gutierrez J, Oberhettinger P, et al. Tracking of antibiotic resistance transfer and rapid plasmid evolution in a hospital setting by nanopore sequencing. mSphere. 2020;5. doi:10.1128/mSphere.00525-20

4. Garbisu C, Garaiyurrebaso O, Epelde L, Grohmann E, Alkorta I. Plasmid-Mediated bioaugmentation for the bioremediation of contaminated soils. Front Microbiol. 2017;8: 1966. doi:10.3389/fmicb.2017.01966

5. Savory EA, Fuller SL, Weisberg AJ, Thomas WJ, Gordon MI, Stevens DM, et al. Evolutionary transitions between beneficial and phytopathogenic rhodococcus challenge disease management. Elife. 2017;6. doi:10.7554/eLife.30925

6. Bruto M, James A, Petton B, Labreuche Y, Chenivesse S, Alunno-Bruscia M, et al. Vibrio crassostreae, a benign oyster colonizer turned into a pathogen after plasmid acquisition. ISME J. 2017;11: 1043–1052. doi:10.1038/ismej.2016.162

7. Pilla G, Tang CM. Going around in circles: Virulence plasmids in enteric pathogens. Nat Rev Microbiol. 2018;16: 484–495. doi:10.1038/s41579-018-0031-2

933   8.   Sentchilo V, Mayer AP, Guy L, Miyazaki R, Green Tringe S, Barry K, et al. Community-

934   wide plasmid gene mobilization and selection. ISME J. 2013;7: 1173–1186.

935   doi:10.1038/ismej.2013.13

936   9.   Xue H, Cordero OX, Camas FM, Trimble W, Meyer F, Guglielmini J, et al. Eco-

937   Evolutionary dynamics of episomes among ecologically cohesive bacterial populations.

938   MBio. 2015;6: e00552–15. doi:10.1128/mBio.00552-15

939   10.   Brown Kav A, Rozov R, Bogumil D, Sørensen SJ, Hansen LH, Benhar I, et al.

940   Unravelling plasmidome distribution and interaction with its hosting microbiome. Environ

941   Microbiol. 2020;22: 32–44. doi:10.1111/1462-2920.14813

942   11.   Stalder T, Press MO, Sullivan S, Liachko I, Top EM. Linking the resistome and

943   plasmidome to the microbiome. ISME J. 2019;13: 2437–2446. doi:10.1038/s41396-019-0446-

944   4

945   12.   Popa O, Dagan T. Trends and barriers to lateral gene transfer in prokaryotes. Curr Opin

946   Microbiol. 2011;14: 615–623. doi:10.1016/j.mib.2011.07.027

947   13.   Benz F, Huisman JS, Bakkeren E, Herter JA, Stadler T, Ackermann M, et al. Plasmid-

948   and strain-specific factors drive variation in ESBL-plasmid spread in vitro and in vivo.

949   ISME J. 2021;15: 862–878. doi:10.1038/s41396-020-00819-4

950   14.   Price VJ, Huo W, Sharifi A, Palmer KL. CRISPR-Cas and Restriction-Modification act

951   additively against conjugative antibiotic resistance plasmid transfer in enterococcus

952   faecalis. mSphere. 2016;1. doi:10.1128/mSphere.00064-16

953   15.   San Millan A, MacLean RC. Fitness costs of plasmids: A limit to plasmid transmission.

954   Microbiol Spectr. 2017;5. doi:10.1128/microbiolspec.MTBP-0016-2017

955   16.   Baltrus DA. Exploring the costs of horizontal gene transfer. Trends Ecol Evol. 2013;28:

956   489–495. doi:**10.1016/j.tree.2013.04.002**

957   17.   Bergstrom CT, Lipsitch M, Levin BR. Natural selection, infectious transfer and the

958   existence conditions for bacterial plasmids. Genetics. 2000;155: 1505–1519. Available:

959   **https://www.ncbi.nlm.nih.gov/pubmed/10924453**

960   18.   Kottara A, Hall JPJ, Harrison E, Brockhurst MA. Variable plasmid fitness effects and

961   mobile genetic element dynamics across pseudomonas species. FEMS Microbiol Ecol.

962   2018;94. doi:**10.1093/femsec/fix172**

963   19.   Hall JPJ, Wood AJ, Harrison E, Brockhurst MA. Source-sink plasmid transfer dynamics

964   maintain gene mobility in soil bacterial communities. Proc Natl Acad Sci U S A.

965   2016;113: 8260–8265. doi:**10.1073/pnas.1600974113**

966   20.   Lynch M, Marinov GK. The bioenergetic costs of a gene. Proc Natl Acad Sci U S A.

967   2015;112: 15690–15695. doi:**10.1073/pnas.1514974112**

968   21.   Agashe D, Shankar N. The evolution of bacterial DNA base composition. J Exp Zool B

969   Mol Dev Evol. 2014;322: 517–528. doi:**10.1002/jez.b.22565**

970   22.   Dietel A-K, Merker H, Kaltenpoth M, Kost C. Selective advantages favour high genomic

971   AT-contents in intracellular elements. PLoS Genet. 2019;15: e1007778.

972   doi:**10.1371/journal.pgen.1007778**

973   23.   Bottery MJ, Wood AJ, Brockhurst MA. Adaptive modulation of antibiotic resistance

974   through intragenomic coevolution. Nat Ecol Evol. 2017;1: 1364–1369. doi:**10.1038/s41559-**

975   **017-0242-3**

24. Jalasvuori M, Friman V-P, Nieminen A, Bamford JKH, Buckling A. Bacteriophage

selection against a plasmid-encoded sex apparatus leads to the loss of antibiotic-

resistance plasmids. Biol Lett. 2011;7: 902–905. doi:10.1098/rsbl.2011.0384

25. San Millan A, Toll-Riera M, Qi Q, MacLean RC. Interactions between horizontally

acquired genes create a fitness cost in pseudomonas aeruginosa. Nat Commun. 2015;6:

6845. doi:10.1038/ncomms7845

26. San Millan A, Toll-Riera M, Qi Q, Betts A, Hopkinson RJ, McCullagh J, et al. Integrative

analysis of fitness and metabolic effects of plasmids in pseudomonas aeruginosa PAO1.

ISME J. 2018. doi:10.1038/s41396-018-0224-8

27. Takahashi Y, Shintani M, Takase N, Kazo Y, Kawamura F, Hara H, et al. Modulation of

primary cell function of host pseudomonas bacteria by the conjugative plasmid pCAR1.

Environ Microbiol. 2015;17: 134–155. doi:10.1111/1462-2920.12515

28. Porse A, Schønning K, Munck C, Sommer MOA. Survival and evolution of a large

multidrug resistance plasmid in new clinical bacterial hosts. Mol Biol Evol. 2016;33:

2860–2873. doi:10.1093/molbev/msw163

29. Hall JPJ, Wright RCT, Guymer D, Harrison E, Brockhurst MA. Extremely fast

amelioration of plasmid fitness costs by multiple functionally diverse pathways.

Microbiology. 2019. doi:10.1099/mic.0.000862

30. Sota M, Yano H, Hughes JM, Daughdrill GW, Abdo Z, Forney LJ, et al. Shifts in the host

range of a promiscuous plasmid through parallel evolution of its replication initiation

protein. ISME J. 2010;4: 1568–1580. doi:10.1038/ismej.2010.72

31.   Stalder T, Rogers LM, Renfrow C, Yano H, Smith Z, Top EM. Emerging patterns of

998   plasmid-host coevolution that stabilize antibiotic resistance. Sci Rep. 2017;7: 4853.

999   doi:10.1038/s41598-017-04662-0

32.   Loftie-Eaton W, Bashford K, Quinn H, Dong K, Millstein J, Hunter S, et al. Compensatory

mutations improve general permissiveness to antibiotic resistance plasmids. Nat Ecol

Evol. 2017;1: 1354–1363. doi:10.1038/s41559-017-0243-2

33.   Harrison E, Guymer D, Spiers AJ, Paterson S, Brockhurst MA. Parallel compensatory

evolution stabilizes plasmids across the parasitism-mutualism continuum. Curr Biol.

2015;25: 2034–2039. doi:10.1016/j.cub.2015.06.024

34.   Lee M-C, Marx CJ. Repeated, selection-driven genome reduction of accessory genes in

experimental populations. PLoS Genet. 2012;8: e1002651.

doi:10.1371/journal.pgen.1002651

35.   Zwanzig M, Harrison E, Brockhurst MA, Hall JPJ, Berendonk TU, Berger U. Mobile

compensatory mutations promote plasmid survival. mSystems. 2019;4: e00186–18.

doi:10.1128/mSystems.00186-18

36.   Jordt H, Stalder T, Kosterlitz O, Ponciano JM, Top EM, Kerr B. Coevolution of host-

plasmid pairs facilitates the emergence of novel multidrug resistance. Nat Ecol Evol.

2020. doi:10.1038/s41559-020-1170-1

37.   Carrilero L, Kottara A, Guymer D, Harrison E, Hall JPJ, Brockhurst MA. Positive

selection inhibits plasmid coexistence in bacterial genomes. Cold Spring Harbor

Laboratory. 2020. p. 2020.09.29.318741. doi:10.1101/2020.09.29.318741

38.   Choi K-H, Schweizer HP. Mini-Tn7 insertion in bacteria with single attTn7 sites: Example

pseudomonas aeruginosa. Nat Protoc. 2006;1: 153–161. doi:10.1038/nprot.2006.24

39.   Hall JPJ, Williams D, Paterson S, Harrison E, Brockhurst MA. Positive selection inhibits

gene mobilisation and transfer in soil bacterial communities. Nat Ecol Evol. 2017;1:

1348–1353. doi:10.1038/s41559-017-0250-3

40.   Lilley AK, Bailey MJ. The acquisition of indigenous plasmids by a genetically marked

pseudomonad population colonizing the sugar beet phytosphere is related to local

environmental conditions. Appl Environ Microbiol. 1997;63: 1577–1583. Available:

https://www.ncbi.nlm.nih.gov/pubmed/16535580

41.   Schweizer HP. Escherichia-Pseudomonas shuttle vectors derived from pUC18/19.

Gene. 1991;97: 109–121. Available: https://www.ncbi.nlm.nih.gov/pubmed/1899844

42.   Heeb S, Itoh Y, Nishijyo T, Schnider U, Keel C, Wade J, et al. Small, stable shuttle

vectors based on the minimal pVS1 replicon for use in gram-negative, plant-associated

bacteria. Mol Plant Microbe Interact. 2000;13: 232–237. doi:10.1094/MPMI.2000.13.2.232

43.   Zhang X-X, Rainey PB. Construction and validation of a neutrally-marked strain of

pseudomonas fluorescens SBW25. J Microbiol Methods. 2007;71: 78–81.

doi:10.1016/j.mimet.2007.07.001

44.   Lenski RE, Rose MR, Simpson SC, Tadler SC. Long-Term experimental evolution in

escherichia coli. I. Adaptation and divergence during 2,000 generations. Am Nat.

1991;138: 1315–1341. doi:10.1086/285289

45.   Hall JPJ, Harrison E, Lilley AK, Paterson S, Spiers AJ, Brockhurst MA. Environmentally

co-occurring mercury resistance plasmids are genetically and phenotypically diverse and

confer variable context-dependent fitness effects. Environ Microbiol. 2015;17: 5008–5022. doi:10.1111/1462-2920.12901

46. Colgan AM, Kröger C, Diard M, Hardt W-D, Puente JL, Sivasankaran SK, et al. The impact of 18 ancestral and Horizontally-Acquired regulatory proteins upon the transcriptome and sRNA landscape of salmonella enterica serovar typhimurium. PLoS Genet. 2016;12: e1006258. doi:10.1371/journal.pgen.1006258

47. Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: Summarize analysis results for multiple tools and samples in a single report. Bioinformatics. 2016;32: 3047–3048. doi:10.1093/bioinformatics/btw354

48. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol. 2019;37: 907–915. doi:10.1038/s41587-019-0201-4

49. Liao Y, Smyth GK, Shi W. The R package rsubread is easier, faster, cheaper and better for alignment and quantification of RNA sequencing reads. Nucleic Acids Res. 2019;47: e47. doi:10.1093/nar/gkz114

50. Robinson MD, McCarthy DJ, Smyth GK. edgeR: A bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics. 2010;26: 139–140. doi:10.1093/bioinformatics/btp616

51. Harrison E, Hall JPJ, Paterson S, Spiers AJ, Brockhurst MA. Conflicting selection alters the trajectory of molecular evolution in a tripartite bacteria-plasmid-phage interaction. Mol Ecol. 2017;26: 2757–2764. doi:10.1111/mec.14080

1061   52.   Hall JPJ, Harrison E, Brockhurst MA. Competitive species interactions constrain abiotic

1062   adaptation in a bacterial soil community. Evol Lett. 2018;2: 580–589. doi:10.1002/evl3.83

1063   53.   Bailey SF, Blanquart F, Bataillon T, Kassen R. What drives parallel evolution?: How

1064   population size and mutational variation contribute to repeated evolution. Bioessays.

1065   2017;39: e201600176. doi:10.1002/bies.201600176

1066   54.   Orr HA. The probability of parallel evolution. Evolution. 2005;59: 216–220. Available:

1067   https://www.ncbi.nlm.nih.gov/pubmed/15792240

1068   55.   Simonsen L, Gordon DM, Stewart FM, Levin BR. Estimating the rate of plasmid transfer:

1069   An end-point method. J Gen Microbiol. 1990;136: 2319–2325. doi:10.1099/00221287-136-11-

1070   2319

1071   56.   Cirz RT, O'Neill BM, Hammond JA, Head SR, Romesberg FE. Defining the

1072   pseudomonas aeruginosa SOS response and its role in the global response to the

1073   antibiotic ciprofloxacin. J Bacteriol. 2006;188: 7101–7110. doi:10.1128/JB.00807-06

1074   57.   Baharoglu Z, Bikard D, Mazel D. Conjugative DNA transfer induces the bacterial SOS

1075   response and promotes antibiotic resistance development through integron activation.

1076   PLoS Genet. 2010;6: e1001165. doi:10.1371/journal.pgen.1001165

1077   58.   Cheng X, Bruijn I de, Voort M van der, Loper JE, Raaijmakers JM. The gac regulon of

1078   pseudomonas fluorescens SBW25. Environ Microbiol Rep. 2013;5: 608–619.

1079   doi:10.1111/1758-2229.12061

1080   59.   Kulkarni PR, Jia T, Kuehne SA, Kerkering TM, Morris ER, Searle MS, et al. A sequence-

1081   based approach for prediction of CsrA/RsmA targets in bacteria with experimental

1082   validation in pseudomonas aeruginosa. Nucleic Acids Res. 2014;42: 6811–6825.

1083   doi:10.1093/nar/gku309

60.   Heeb S, Blumer C, Haas D. Regulatory RNA as mediator in GacA/RsmA-dependent

global control of exoproduct formation in pseudomonas fluorescens CHA0. J Bacteriol.

2002;184: 1046–1056. Available: https://www.ncbi.nlm.nih.gov/pubmed/11807065

61.   Ghequire MGK, De Mot R. The tailocin tale: Peeling off phage tails. Trends Microbiol.

2015;23: 587–590. doi:10.1016/j.tim.2015.07.011

62.   Mavrodi DV, Loper JE, Paulsen IT, Thomashow LS. Mobile genetic elements in the

genome of the beneficial rhizobacterium pseudomonas fluorescens pf-5. BMC Microbiol.

2009;9: 8. doi:10.1186/1471-2180-9-8

63.   Zimmermann L, Stephens A, Nam S-Z, Rau D, Kübler J, Lozajic M, et al. A completely

reimplemented MPI bioinformatics toolkit with a new HHpred server at its core. J Mol

Biol. 2018;430: 2237–2243. doi:10.1016/j.jmb.2017.12.007

64.   Wein T, Hülter NF, Mizrahi I, Dagan T. Emergence of plasmid stability under non-

selective conditions maintains antibiotic resistance. Nat Commun. 2019;10: 2595.

doi:10.1038/s41467-019-10600-7

65.   Turner PE, Williams ESCP, Okeke C, Cooper VS, Duffy S, Wertz JE. Antibiotic

resistance correlates with transmission in plasmid evolution. Evolution. 2014;68: 3368–

3380. doi:10.1111/evo.12537

66.   Morton ER, Merritt PM, Bever JD, Fuqua C. Large deletions in the pAtC58 megaplasmid

of agrobacterium tumefaciens can confer reduced carriage cost and increased

expression of virulence genes. Genome Biol Evol. 2013;5: 1353–1364.

doi:10.1093/gbe/evt095

67.   Kloos J, Gama JA, Hegstad J, Samuelsen Ø, Johnsen PJ. Piggybacking on niche-
adaptation improves the maintenance of multidrug resistance plasmids. Mol Biol Evol.
2021. doi:10.1093/molbev/msab091

68.   Baharoglu Z, Mazel D. SOS, the formidable strategy of bacteria against aggressions.
FEMS Microbiol Rev. 2014;38: 1126–1145. doi:10.1111/1574-6976.12077

69.   Dwyer DJ, Camacho DM, Kohanski MA, Callura JM, Collins JJ. Antibiotic-induced
bacterial cell death exhibits physiological and biochemical hallmarks of apoptosis. Mol
Cell. 2012;46: 561–572. doi:10.1016/j.molcel.2012.04.027

70.   Yano H, Wegrzyn K, Loftie-Eaton W, Johnson J, Deckert GE, Rogers LM, et al. Evolved
plasmid-host interactions reduce plasmid interference cost. Mol Microbiol. 2016;101:
743–756. doi:10.1111/mmi.13407

71.   Shintani M, Takahashi Y, Tokumaru H, Kadota K, Hara H, Miyakoshi M, et al. Response
of the pseudomonas host chromosomal transcriptome to carriage of the IncP-7 plasmid
pCAR1. Environ Microbiol. 2010;12: 1413–1426. doi:10.1111/j.1462-2920.2009.02110.x

72.   Machnicka MA, Kaminska KH, Dunin-Horkawicz S, Bujnicki JM. Phylogenomics and
sequence-structure-function relationships in the GmrSD family of type IV restriction
enzymes. BMC Bioinformatics. 2015;16: 336. doi:10.1186/s12859-015-0773-z

73.   Doron S, Melamed S, Ofir G, Leavitt A, Lopatina A, Keren M, et al. Systematic discovery
of antiphage defense systems in the microbial pangenome. Science. 2018; eaar4120.
doi:10.1126/science.aar4120

74. Oliveira PH, Touchon M, Rocha EPC. Regulation of genetic flux between bacteria by

restriction-modification systems. Proc Natl Acad Sci U S A. 2016;113: 5658–5663.

doi:10.1073/pnas.1603257113

75. Jiang W, Maniv I, Arain F, Wang Y, Levin BR, Marraffini LA. Dealing with the

evolutionary downside of CRISPR immunity: Bacteria and beneficial plasmids. PLoS

Genet. 2013;9: e1003844. doi:10.1371/journal.pgen.1003844

76. Broek D van den, Chin-A-Woeng TFC, Bloemberg GV, Lugtenberg BJJ. Molecular

nature of spontaneous modifications in gacS which cause colony phase variation in

pseudomonas sp. Strain PCL1171. J Bacteriol. 2005;187: 593–600.

doi:10.1128/JB.187.2.593-600.2005

77. Madsen JS, Hylling O, Jacquiod S, Pécastaings S, Hansen LH, Riber L, et al. An

intriguing relationship between the cyclic diguanylate signaling system and horizontal

gene transfer. ISME J. 2018. doi:10.1038/s41396-018-0183-0

78. Song D, Loparo JJ. Building bridges within the bacterial chromosome. Trends Genet.

2015;31: 164–173. doi:10.1016/j.tig.2015.01.003

79. Kawalek A, Glabski K, Bartosik AA, Fogtman A, Jagura-Burdzy G. Increased ParB level

affects expression of stress response, adaptation and virulence operons and potentiates

repression of promoters adjacent to the high affinity binding sites parS3 and parS4 in

pseudomonas aeruginosa. PLoS One. 2017;12: e0181726.

doi:10.1371/journal.pone.0181726

80. Kusiak M, Gapczynska A, Plochocka D, Thomas CM, Jagura-Burdzy G. Binding and

spreading of ParB on DNA determine its biological function in pseudomonas aeruginosa.

J Bacteriol. 2011;193: 3342–3355. doi:10.1128/JB.00328-11

81.   Murray H, Ferreira H, Errington J. The bacterial chromosome segregation protein Spo0J

1149   spreads along DNA from parS nucleation sites. Mol Microbiol. 2006;61: 1352–1361.

1150   doi:10.1111/j.1365-2958.2006.05316.x

1151   82.   Bartosik AA, Lasocki K, Mierzejewska J, Thomas CM, Jagura-Burdzy G. ParB of

1152   pseudomonas aeruginosa: Interactions with its partner ParA and its target parS and

1153   specific effects on bacterial growth. J Bacteriol. 2004;186: 6983–6998.

1154   doi:10.1128/JB.186.20.6983-6998.2004

1155   83.   Baek JH, Rajagopala SV, Chattoraj DK. Chromosome segregation proteins of vibrio

1156   cholerae as transcription regulators. MBio. 2014;5: e01061–14. doi:10.1128/mBio.01061-14

1157   84.   San Millan A, Heilbron K, MacLean RC. Positive epistasis between co-infecting plasmids

1158   promotes plasmid survival in bacterial populations. ISME J. 2014;8: 601–612.

1159   doi:10.1038/ismej.2013.182

1160   85.   Petrova V, Chitteni-Pattu S, Drees JC, Inman RB, Cox MM. An SOS inhibitor that binds

1161   to free RecA protein: The PsiB protein. Mol Cell. 2009;36: 121–130.

1162   doi:10.1016/j.molcel.2009.07.026

1163   86.   Doyle M, Fookes M, Ivens A, Mangan MW, Wain J, Dorman CJ. An H-NS-like stealth

1164   protein aids horizontal DNA transmission in bacteria. Science. 2007;315: 251–252.

1165   doi:10.1126/science.1137550

1166   87.   Silby MW, Cerdeño-Tárraga AM, Vernikos GS, Giddens SR, Jackson RW, Preston GM,

1167   et al. Genomic and genetic analyses of diversity and plant interactions of pseudomonas

1168   fluorescens. Genome Biol. 2009;10: R51. doi:10.1186/gb-2009-10-5-r51

88. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, et al. PHASTER: A better, faster

version of the PHAST phage search tool. Nucleic Acids Res. 2016;44: W16–21.

doi:10.1093/nar/gkw387

89. West SA, Diggle SP, Buckling A, Gardner A, Griffin AS. The social lives of microbes.

Annu Rev Ecol Evol Syst. 2007;38: 53–77. doi:10.1146/annurev.ecolsys.38.091206.095740

90. Gardner A, West SA. Spite and the scale of competition. J Evol Biol. 2004;17: 1195–

1203. doi:10.1111/j.1420-9101.2004.00775.x

91. Lopatina A, Tal N, Sorek R. Abortive infection: Bacterial suicide as an antiviral immune

strategy. Annu Rev Virol. 2020;7: 371–384. doi:10.1146/annurev-virology-011620-040628

92. Carraro N, Rivard N, Ceccarelli D, Colwell RR, Burrus V. IncA/C conjugative plasmids

mobilize a new family of multidrug resistance islands in clinical vibrio cholerae Non-

O1/Non-O139 isolates from haiti. MBio. 2016;7. doi:10.1128/mBio.00509-16

93. Pal C, Bengtsson-Palme J, Kristiansson E, Larsson DGJ. Co-occurrence of resistance

genes to antibiotics, biocides and metals reveals novel insights into their co-selection

potential. BMC Genomics. 2015;16: 964. doi:10.1186/s12864-015-2153-5

94. Aviv G, Tsyba K, Steck N, Salmon-Divon M, Cornelius A, Rahav G, et al. A unique

megaplasmid contributes to stress tolerance and pathogenicity of an emergent

salmonella enterica serovar infantis strain. Environ Microbiol. 2014;16: 977–994.

doi:10.1111/1462-2920.12351

95. Cusick KD, Polson SW, Duran G, Hill RT. Multiple mega-plasmids confer extremely high

metal tolerance in alteromonas strains. Appl Environ Microbiol. 2019.

doi:10.1128/AEM.01831-19

96. Cazares A, Moore MP, Hall JPJ, Wright LL, Grimes M, Emond-Rhéault J-G, et al. A

megaplasmid family driving dissemination of multidrug resistance in pseudomonas. Nat

Commun. 2020;11: 1370. doi:10.1038/s41467-020-15081-7

97. Urbanowicz P, Bitar I, Izdebski R, Baraniak A, Literacka E, Hrabák J, et al. Epidemic

territorial spread of IncP-2-Type VIM-2 Carbapenemase-Encoding megaplasmids in

nosocomial pseudomonas aeruginosa populations. Antimicrob Agents Chemother.

2021;65. doi:10.1128/AAC.02122-20

98. Gallet R, Cooper TF, Elena SF, Lenormand T. Measuring selection coefficients below

10(-3): Method, questions, and prospects. Genetics. 2012;190: 175–186.

doi:10.1534/genetics.111.133454

99. Harrison E, Truman J, Wright R, Spiers AJ, Paterson S, Brockhurst MA. Plasmid

carriage can limit bacteria-phage coevolution. Biol Lett. 2015;11. doi:10.1098/rsbl.2015.0361

100. Lopatkin AJ, Meredith HR, Srimani JK, Pfeiffer C, Durrett R, You L. Persistence and

reversal of plasmid-mediated antibiotic resistance. Nat Commun. 2017;8: 1689.

doi:10.1038/s41467-017-01532-1

101. Valle AA, Leon-Sampedro R, Rodriguez-Beltran J, DelaFuente J, Hernandez-Garcia M,

Ruiz-Garbajosa P, et al. The distribution of plasmid fitness effects explains plasmid

persistence in bacterial communities. 2020. p. 2020.08.01.230672.

doi:10.1101/2020.08.01.230672

102. Whelan FJ, Rusilowicz M, McInerney JO. Coinfinder: Detecting significant associations

and dissociations in pangenomes. Microb Genom. 2020;6. doi:10.1099/mgen.0.000338