Respiratory dysfunctions in two rodent models of chronic epilepsy and acute

seizures and their link with brainstem serotonin system

Hayet Kouchi [1,2], Michaël Ogier [3], Gabriel Dieuset [4], Anne Morales [1,2],

Béatrice Georges [1,2], Jean-Louis Rouanet [5], Benoît Martin [4], Philippe Ryvlin [6],

Sylvain Rheims [1,2,7*], Laurent Bezin [1,2*]

1.   Lyon Neuroscience Research Center, Inserm U1028, CNRS UMR5292, Lyon 1 University,
Lyon, France ;
2.   Epilepsy institute, Bron, France;
3.   IRBA, Institut de Recherche Biomédicale des Armées, Brétigny-sur-Orge, France;
4.   Univ Rennes, CHU Rennes, Inserm, LTSI (Laboratoire de Traitement du Signal et de
l'Image), UMR-1099, Rennes, France;
5.   Laboratoire d'Ecologie des Hydrosystèmes Naturels et Anthropisés, U5023, Lyon 1
University, Lyon, France;
6.   Department of Clinical Neurosciences, Centre Hospitalier Universitaire Vaudois,
Université de Lausanne, Lausanne, Switzerland.
7.   Department of Functional Neurology and Epileptology, Hospices Civils de Lyon, Lyon,
France.
19
20   * co-last authors
21

Corresponding author:

Laurent Bezin, CRNL UMR5292 U1028, Epilepsy institute IDEE, 59 boulevard Pinel, F-69500

Bron, France; laurent.bezin@univ-lyon1.fr; tel: +33 472 138 970.

ORCID ID:
Laurent Bezin: 0000-0001-6411-4267
Gabriel Dieuset: 0000-0002-6300-7352
Hayet Kouchi: 0000-0003-1443-0645
Benoît Martin: 0000-0002-8412-1488
Sylvain Rheims: 0000-0002-4663-8515
32

33   Number of figures: 5
34   Number of tables: 1
35   Supplementary figures: 2
36   Supplementary table: 1
37

Abbreviations: 5-HT, serotonin; AGS, audiogenic seizures; EPI; CIH, chronic intermittent

hypoxia; EEG, electro-encephalography; epileptic; EPI/SDOC-, epileptic rat without sharp

decrease in oxygen consumption; EPI/SDOC+, epileptic rat with sharp decrease in oxygen

consumption; f_R, respiratory frequency; FRA, fatal respiratory arrest; HCVR, hypercapnia

ventilatory response; HIS, handling-induced seizures; HVR, hypoxia ventilatory response;

NFRA, non-fatal respiratory arrest; GTCS, generalized tonic-clonic seizures; NTS, nucleus

tractus solitarius; OSA, obstructive sleep apnea syndrome; Pilo-SE, pilocarpine-induced

status epilepticus; SE, *status epilepticus*; SERT, serotonin transporter; SSRI, selective

serotonin reuptake inhibitor; SUDEP, sudden unexpected death in epilepsy; T_E, expiration

time; T_I, inspiration time; Tph2, tryptophan hydroxylase 2; T_{tot}, total respiratory cycle

duration; $\dot{V}_E$, minute volume; VLM, ventrolateral medulla; $\dot{V}O_2$, oxygen consumption; V_T, tidal

volume

Key words: *status epilepticus*, Sprague-Dawley rat, DBA/2 mice, Metabolic rate of oxygen,

SUDEP.

Abstract

Patients with drug-resistant epilepsy can experience respiratory alterations notably during seizures. The mechanisms underlying this long-term alteration of respiratory function remain unclear. This study aimed at determining in rats whether epilepsy is associated with alterations of both the respiratory function and brainstem serotonin (5-HT) system. Epilepsy was triggered by pilocarpine-induced *status epilepticus* in rats. 30-50% of epileptic (EPI) rats exhibited sharp decrease of oxygen consumption (SDOC), low metabolic rate of oxygen and slow regular ventilation; these rats were called EPI/SDOC+ rats. These alterations were only detected in rats with chronic epilepsy, independent of behavioral seizures, persisted over the time, and were not associated with death. In these rats, 5-HT fiber density in the nucleus *tractus solitarius* was below that of control and EPI/SDOC- rats. Both EPI/SDOC+ rats and DBA/2 mice presenting with fatal respiratory arrest following an audiogenic-induced seizure, a model of sudden and expected death in epilepsy, had increased transcript levels of tryptophan hydroxylase 2 (p<0.001 for both strains) and 5-HT presynaptic transporter (rats: p=0.003; mice: p=0.001). Thus, our data support that 5-HT alterations are associated with chronic and acute epilepsy-related respiratory dysfunctions.

Introduction

Seizure-related respiratory dysfunctions in patients with epilepsy are well characterized and

can be observed either during focal seizures or during generalized tonic-clonic seizures

(GTCS) [1, 2]. In patients with drug-resistant epilepsy, oxygen desaturation <90%, as

measured by pulse oximetry, has thus been reported in 33% of focal seizures [3] and in up to

86% of GTCS [2]. In both seizure types, these transient hypoxemias result from ictal and/or

post-ictal central apnea [1, 4]. In addition, respiratory dysfunctions are not only limited to

seizure periods in some patients with epilepsy. Obstructive sleep apnea syndrome (OSA) is

thus prevalent in people with drug-resistant epilepsy ranging between 30-49% and markedly

exceeding the estimated rate in the general population [5, 6].

Importantly, these respiratory dysfunctions might represent a risk factor for sudden

unexpected death in epilepsy (SUDEP), which primarily results from post-ictal central apnea

[3], urging to fill the gap of knowledge regarding epilepsy-related respiratory alteration and

its pathological mechanisms. Post-mortem examination of the brainstem of patients who

died from SUDEP showed changes in the ventrolateral medulla (VLM) [7], a region containing

different populations of respiratory neurons including those of the pre-Bötzinger Complex, a

kernel generator of the inspiratory rhythm [8]. Specifically, SUDEP victims presented with an

alteration of brainstem serotonin (5-HT) system with notable decrease in both tryptophan

hydroxylase (TPH2), the rate-limiting enzyme in 5-HT synthesis, and 5-HT presynaptic

transporter (SERT) within medullary raphe nuclei and/or the VLM [7].

Similarly, brainstem 5-HT abnormalities have been reported in DBA/1 and DBA/2 mouse

strains, which are frequently used as a SUDEP model in preclinical studies because they can

exhibit a fatal respiratory arrest (FRA) after induction of an audiogenic seizure (AGS) [9-11].

Interestingly, FRAs are prevented when DBA/2 and DBA/1 mice are treated with fluoxetine, a

selective serotonin reuptake inhibitor (SSRI) before AGS induction, which is consistent with

the hypothesis that 5-HT system may have a strong implication in FRA following AGS [12, 13].

Nevertheless, animals prone to AGS-induced FRA present some limitations to completely

characterize respiratory dysfunction in chronic epilepsy and its potential associated 5-HT

abnormalities, mostly because they represent models of induced seizure(s) and do not

develop spontaneous recurrent seizures.

In this study, we used one of the most studied rat model of chronic epilepsy, following

pilocarpine-induced *status epilepticus* (Pilo-SE). As for a large number of patients with

temporal lobe epilepsy, Pilo-SE rats are notably characterized by chronic, spontaneous and

drug-resistant seizures [14]. In this model, while mortality is present during SE, no mortality

has been reported once chronic phase of epilepsy has developed. Thus, to further explore

the hypothesis that epilepsy-related respiratory alterations are associated with altered

brainstem 5-HT system, we first explored whether respiratory alterations occurred with

development of epilepsy, and then investigated whether they were associated with

brainstem 5-HT system alterations; i.e. number of 5-HT-positive cell bodies, density of 5-HT-

positive terminals, TPH2 and SERT mRNA levels. To help better conclude on the potential

involvement of 5-HT system alterations with respect to the respiratory disorders observed

during the chronic phase of epilepsy and independently of any risk of SUDEP-like mortality,

we finally compared brainstem TPH2 and SERT mRNA levels between EPI rats and DBA/2

mice exhibiting FRA after AGS.

Results

Development of behavioral spontaneous recurrent seizures (SRSs) in rats after Pilo-SE

The numbers of rats that did not develop Pilo-SE or died during SE are reported in the Supplementary information section. SRSs were identified as detailed in the Methods section. Briefly, out of the 82 rats subjected to Pilo-SE in experiments 1-3, 59 developed at least two consecutive SRSs by the end of the 2nd week post-SE, and were then stated as EPI rats.

According to the evaluation of two experimenters, the number of recorded seizures (stages 3-5 of severity according to Racine's scale) per 12 hours during the daylight period was about 1.1 ± 0.1 in EPI rats from weeks 3 to 8 after SE. In experiment 1, none of the Pilo-SE rats had developed 2 consecutive Handling-Induced Seizures (HIS) at one-week post-SE, in agreement with previous studies [15].

A subset of EPI rats presented with abnormal pattern of oxygen consumption

In experiment 1, metabolic rate of oxygen ($\dot{V}O_2$) of 20 EPI rats and 8 controls was recorded for 24h each week from weeks 1 to 8 after SE. A thorough screening of $\dot{V}O_2$ was then made for each rat.

Presence of sharp decreases in oxygen consumption (SDOCs). Control rats (n=8) presented with slight up and down fluctuations of $\dot{V}O_2$ during both light and dark periods, as illustrated in one rat (Figure 1A). For a better characterization of $\dot{V}O_2$ fluctuations within the daily pattern, $\dot{V}O_2$ was averaged for each rat during the light and dark periods, separately. Afterwards, the lowest value below the averaged $\dot{V}O_2$ was determined for each rat; it ranged between 38% and 70% of their average $\dot{V}O_2$, independently of the circadian rhythm. This pattern remained unchanged during the next 7 weeks of recordings. In rats subjected to Pilo-SE, the pattern of daily $\dot{V}O_2$ was similar to that of controls during the first two weeks

post-SE, as illustrated in one Pilo-SE rat (Figure 1B). However, by the end of the 2^{nd} week

post-SE, two groups of EPI rats could be identified according to $\dot{V}O_2$ fluctuations. A subset of

EPI rats presented with a large amplitude of $\dot{V}O_2$ decrease ("Sharp Decrease(s) in Oxygen

Consumption"; SDOCs), defined as a decrease greater than -62%, a percentage that

corresponds to the maximal decrease of $\dot{V}O_2$ in control rats. Such EPI rats were thus

identified as EPI/SDOC+ rats and their lowest $\dot{V}O_2$ values ranged between 0% and 32% of the

average $\dot{V}O_2$, as illustrated at 3 weeks post-SE in one rat (Figure 1C). By contrast, the

remaining EPI rats (EPI/SDOC- rats) were similar to controls with regard to the lowest $\dot{V}O_2$

values, which ranged between 41% and 67% of the average $\dot{V}O_2$. Video monitoring

confirmed that SDOCs were not associated with behavioral seizures, as illustrating while rat

adopted a sleep-like position (Figure 2A). By contrast, behavioral seizures were associated

with a long-lasting increase of the $\dot{V}O_2$, as illustrated during a stage-5 seizure (Figure 2B).

Regarding the highest $\dot{V}O_2$ values in control rats, they were ranged between 164% and 229%

of the average $\dot{V}O_2$ (median=183%). The highest fluctuations within the daily $\dot{V}O_2$ pattern of

EPI/SDOC+ and EPI/SDOC- rats were not statistically different from controls, and ranged

between 144% and 227% of the average $\dot{V}O_2$ (median=180%). However, EPI/SDOC+ rats had

greater increases of $\dot{V}O_2$ than those observed in EPI/SDOC- rats (+93 ± 5% *vs* +75 ± 4%;

P=0.01).

As a good indicator of the stability of $\dot{V}O_2$ pattern, the intra-individual variation (coefficient

of variation, CV) of the daily oxygen consumption pattern (CV-$\dot{V}O_2$) was calculated

independently of light and dark periods. The mean CV-$\dot{V}O_2$ of the control group was

estimated to be 20.8 ± 0.5% and 22 ± 0.7% for the light and the dark periods, respectively.

The mean CV-$\dot{V}O_2$ of EPI/SDOC- rats was almost similar to that of controls with 19.1 ± 0.8%

and 22.8 ± 0.7% for the light and dark periods, respectively. However, in EPI/SDOC+ rats, it

was significantly higher than that measured in both control (P= 0.002 and p= 0.004 for light

and dark periods, respectively) and EPI/SDOC- groups (P< 0.0001 and p= 0.017 for light and

dark periods, respectively), reaching 27.4 ± 2.9% and 26.5 ± 1.3% for light and dark periods,

respectively.

Low metabolic rate of oxygen in EPI/SDOC+ rats. Since $\dot{V}O_2$ was measured weekly for all

rats for 8 weeks, it was possible to retrospectively assign each EPI rat into either the

EPI/SDOC+ or the EPI/SDOC- group from the first week after SE.

We investigated the evolution of the daily $\dot{V}O_2$ in each group. We first verified that SDOC

events in EPI/SDOC+ rats did not affect the average value of the daily $\dot{V}O_2$, both during the

light (P=0.86) and the dark (P=0.93) periods. Indeed, in order to compare between the three

groups of rats the daily $\dot{V}O_2$ during the 8 weeks of measurement, we preferred to use the

data without SDOCs in EPI/SDOC+ rats. We show in control and EPI/SDOC- rat groups that

the daily $\dot{V}O_2$ decreased progressively by 21.3 ± 2.4% (P=0.05) and 22.0 ± 2.2% (P=0.02),

respectively, between weeks 1-2 and 8 post-SE. By contrast, the decrease observed in

EPI/SDOC+ rats was much larger during the same period (-69.5 ± 1.1%; P=0.002), and already

reached -53 ± 6.0% (P=0.02) by the fifth week post-SE (Figure 3A). We also found that the

decrease of daily $\dot{V}O_2$ coincided with the onset of SDOCs (Figure 3B).

In each group, we verified that the daily $\dot{V}O_2$ did not change significantly between 5 and 8

weeks post-SE. Then, the daily $\dot{V}O_2$ has been averaged during that period for each group to

ease the comparison. We found that the daily $\dot{V}O_2$ was greatly reduced in EPI/SDOC+ rats at

5-8 weeks post-SE compared to both EPI/SDOC- rats and controls (-39 ± 7 % and -34 ± 7 %,

respectively). This decrease was observed during both the light period (EPI/SDOC+ *vs.*

EPI/SDOC- rats: p< 0.001; EPI/SDOC+ *vs.* control rats: P=0.002) and the dark period

(EPI/SDOC+ *vs.* EPI/SDOC- rats: p< 0.001; EPI/SDOC+ *vs.* control rats: P=0.002)

(Supplementary Figure S1A). At weeks 1-2 post-SE, no difference was observed between

groups during both light and dark periods (Supplementary Figure S1A).

In the 3 groups of rats (controls, EPI/SDOC+ and EPI/SDOC-), we evaluated the effect of

three factors that have been referenced as affecting $\dot{V}O_2$ [16]: time (t_0 corresponds to the

time of induction of SE in EPI rats), body weight and circadian rhythm. Full description of

these effects is detailed in the Supplementary information section.

Characterization of SDOCs. The number of SDOCs during the 24-hour measurement ranged

between 12 and 414 events (median = 44), with no difference of SDOC numbers between

dark and light periods. The median duration of SDOCs was 5 min (range: 20 sec to 13 min).

The longest SDOC event is illustrated in figure 2A; it lasted 13 min and occurred, according to

the video, during the daylight period while the animal was sleeping without any evidence of

behavioral seizure. Full description of $\dot{V}O_2$ pattern during the manifestation of seizures is

detailed in the Supplementary information section.

Body position during SDOCs was far different from freezing posture frequently observed

before rats had a behavioral seizure that was easily discernible by visual observation alone

(stages 3 to 5 on Racine's scale). It is also noteworthy that rats always exhibited a quiet state

during SDOC events. The proportion of EPI/SDOC+ rats increased between weeks 3 and 5

post-SE and then stabilized (from weeks 5 to 8 post-SE). By the end of the 5[th] week post-SE,

30% (6/20) of EPI rats exhibited SDOCs (Figure 3C).

Ventilatory and cardiac functions in epileptic rats

Experiment 2 aimed to further investigate respiratory function by evaluating ventilatory and

cardiac functions of EPI rats during chronic phase of epilepsy. It should be noted that we

could not record respiratory variables using whole-body plethysmography before the 10th

week post-SE because rats were not able to stay calm in the apparatus chamber, likely due

to enhanced reactivity to stressful situations.

EPI rats presented with a low steady respiratory pattern and enhanced ventilatory

response to hypoxia and hypercapnia. Under normoxia, EPI rats (n=16) presented a slow

respiratory pattern with a significant decrease of minute ventilation ($\dot{V}_E$) compared to

controls (n=5) (-12 ± 3%, P=0.03) (Supplementary Figure S2A). The reduction of $\dot{V}_E$ in EPI rats

resulted from a decrease of respiratory frequency (f_R) (-12 ± 2% *vs* controls, P=0.001) while

tidal volume (V_T) was similar to controls (P=0.92). The decrease of f_R of EPI rats was related

to a simultaneous increase of inspiratory time (T_I) and expiratory time (T_E), that reached +20

± 3% (P=0.01) and +10 ± 2 % (P=0.001) compared to the values measured in controls,

respectively. Consequently, the total respiratory cycle duration (T_{tot}) of EPI rats was

significantly higher than controls (+14 ± 2 % ; P=0.002) (Supplementary Figure S2A).

Regarding the stability of breathing pattern, the coefficient of variation of breathing, apnea

frequency and duration were similar between EPI and control rats (p=0.16, p=0.61 and

p=0.09, respectively) (Supplementary Figure S2B).

Compared to control rats (n=13), EPI rats (n=18) showed a greater hypoxia ventilatory

response (HVR) with higher $\dot{V}_E$ (+23 ± 6 % *vs* controls, P=0.007), higher f_R (+16 ± 2 % *vs*

controls, P<0.001) and similar V_T (P=0.06). Similarly, the hypercapnia ventilatory response

(HCVR) of EPI rats (n=18) was higher than controls (n=12) with greater $\dot{V}_E$ (+14 ± 4 % *vs*

controls, P=0.01) and greater f_R (+10 ± 3 % *vs* controls, p=0.02), while V_T remained similar to

that of controls (P=0.09).

Abnormal $\dot{V}O_2$ pattern is associated with pronounced decrease of ventilatory rhythm at

rest. Since oxygen is strictly provided during ventilatory process, we aimed to compare

ventilation between EPI rats with or without abnormal $\dot{V}O2$ pattern. In experiment 2, we

were able to discriminate 8 EPI/SDOC+ rats, which presented with a daily $\dot{V}O_2$ lower than

that of both EPI/SDOC- rats (n=15, -51 ± 9%; P< 0.001) and control rats (n=16, -41 ± 11%;

P=0.003).

Under normoxia, the ventilation of EPI/SDOC- rats was significantly altered compared to

control rats (Table 1A), but to a lesser extent than that of EPI/SDOC+ rats. Indeed, compared

to control rats, EPI/SDOC- rats presented with a slight but not significant 6 ± 3% reduction of

$\dot{V}_E$, associated with an 8 ± 2% reduction (P=0.015) of f_R, a 13 ± 1% increase (P<0.001) of T_I

and a 9 ± 2% increase (P=0.025) of T_{tot} (Table 1A). Furthermore, compared to EPI/SDOC- rats,

EPI/SDOC+ rats presented with an additional 13 ± 3% reduction (P=0.013) of $\dot{V}_E$, that resulted

from an 8 ± 2% reduction (P=0.036) of f_R, a 13 ± 3% increase (P<0.001) of T_I and an 8 ± 3%

increase (P=0.013) of T_{tot} (Table 1A). Tidal volume (V_T), the coefficient of variation of the

breathing frequency, apnea frequency and duration were similar in all groups (Table 1A).

Under hypoxic and hypercapnic conditions, $\dot{V}_E$, V_T, and f_R was similar between EPI/SDOC+

(n=6) and EPI/SDOC- rats (n=12) (Table 1B).

No alteration of cardiac frequency and body-temperature in EPI rats. Respiratory system,

cardiac function and body-temperature have a tight relationship. In this experiment, we

showed that cardiac frequency and body-temperature of EPI/SDOC+ rats (n=2), EPI/SDOC-

rats (n=3) and control rats (n=3) were similar during both light and dark periods (Table 1C,

1D).

Brainstem 5-HT system of EPI rats

In experiment 3, 20 rats developed epilepsy. Among those rats, we identified 8 EPI/SDOC+

and 8 EPI/SDOC- rats. Eight control rats were used. Brainstem was processed for either

histological (n=3/8 rats in each group) or gene expression (n=5/8 rats in each group) analyses.

Alteration of the density of brainstem 5-HT immunopositive processes in EPI rats. We

determined whether metabolic and ventilatory alterations developed by EPI/SDOC+ rats

were associated with dysfunction of the 5-HT system in the brainstem. Therefore, we

focused our investigation within caudal raphe nuclei, which are known to innervate neuronal

groups closely implicated in the regulation of respiratory rhythm generation, such as the NTS

and the VLM. 5-HT-immunolabeling was detected in both processes and cell bodies. Within

caudal raphe nuclei (pallidus, magnus and obscurus), although we observed a downward

trend in epileptic rats, there was no statistical difference in the number of 5-HT-positive

neuronal cell bodies between the three groups of rats (control rats: 22 ± 6, EPI/SDOC-: 18 ± 4,

EPI/SDOC+: 13 ± 5, P=0.339). In addition, there was no difference in surface area of 5-HT

positive cell bodies (control rats: 383 ± 21 pixel^2, EPI/SDOC- rats: 445 ± 37 pixel^2, EPI/SDOC+

rats: 397 ± 54 pixel^2, P=0.383) nor in the index of 5-HT neuronal concentration (control rats:

109 ± 19 A.U., EPI/SDOC- rats: 125 ± 23 A.U., EPI/SDOC+ rats: 119 ± 18 A.U., P=0.864)

between the three rat groups. The density of 5-HT immunopositive processes within both

the VLM (Figure 4A) and the NTS (Figure 4B) was significantly higher in EPI/SDOC- rats

compared to EPI/SDOC+ rats and control rats (Figures 4A4 and 4B4). Furthermore, the

density of 5-HT-immunolabeled processes within the NTS was significantly lower in

EPI/SDOC+ rats compared to controls (Figure 4B4).

Alteration in brainstem transcript level of TPH2 and SERT in EPI/SDOC+ rats. TPH2

transcript level was significantly increased in EPI/SDOC+ compared to both EPI/SDOC- rats

and control rats. There was no statistical difference in transcript level of TPH2 between

EPI/SDOC- and control rats (Figure 5A). SERT transcript level was significantly increased in

EPI/SDOC+ compared to EPI/SDOC- rats, but not compared to control rats. In addition, SERT

transcript level did not differ statistically between control and EPI/SDOC- rats (Figure 5B).

Brainstem 5-HT system of DBA/2 mice prone to AGS

Previous studies demonstrated that almost 75% of DBA/2 mice developing AGS died from

fatal respiratory arrest (FRA) following a tonic hindlimb extension [17]. Among the remaining

mice, 15% developed non-fatal respiratory arrest (NFRA), meaning that they spontaneously

recovered from post-AGS respiratory arrest. Brainstem 5-HT system of DBA/2 mice

exhibiting FRA in response to AGS was shown to be altered in comparison with C57Black/6

mice that are resistant to AGS [10]. Thus, in experiment 4, we examined whether the

alterations of TPH2 and SERT gene expression that we have observed between EPI/SDOC+

rats and EPI/SDOC- rats could be found between FRA and NFRA DBA/2 mice. Among the

eleven male DBA/2 mice that underwent AGS with respiratory arrest, 64% of DBA/2 mice

presented with FRA (n=7) and 36% with NFRA (n=4). We found that the levels of TPH2 and

SERT transcripts were significantly elevated in FRA DBA/2 mice compared to NFRA DBA/2

mice (Figure 5A-B). It was striking that the level of TPH2 and SERT transcripts of FRA DBA/2

mice followed a trend that was similar to that of EPI/SDOC+ rats when both were compared

to their respective controls (NFRA DBA/2 mice and EPI/SDOC- rats).

Discussion

This study reports chronic alterations of respiratory system in epileptic (EPI) rats associated with 5-HT dysfunctions within the brainstem.

Exploration of respiratory function in epilepsy models.

To date, studies in animal models of seizures, aimed at understanding respiratory dysfunction in the context of epilepsy, were mostly performed during the time period surrounding seizures, i.e. during ictal and post-ictal periods [17-27]. Respiratory alterations have been detected in these abovementioned studies from short recording periods (2-30 min). In view of the respiratory disorders that can affect patients suffering from temporal lobe epilepsy (ref), this work carried out on a widely used preclinical model of this type of epilepsy (following status epilepticus induced by intraperitoneal administration of pilocarpine, Pilo-SE) had two major objectives.

The first was to study respiratory function in rats with a long history of epilepsy and at a distance from the period of brain remodeling that follows Pilo-SE-induced severe brain damages [28]. After a 10-week period of active epilepsy (12 weeks post-SE), we showed a decrease of breathing rhythm in epileptic (EPI) rats at rest. This result is different from that reported in a recent study showing no alteration in resting ventilation in rats 2-4 weeks after SE developed following intrahippocampal administration of pilocarpine [29]. However, beyond the fact that we did not use the same strain of rats (Wistar vs. Sprague-Dawley in our study) nor the same route of pilocarpine administration (intrahippocampally vs. intraperitoneally in our study), we agree with the authors of the above-mentioned study that the mechanisms leading to altered breathing may depend on the stage of epilepsy. In

their study, the assessment was rather at an early, whereas it was clearly at a later stage of

epilepsy in our study.

The second objective was to perform a long duration (24h) recording of the respiratory

function in order to extend investigation beyond the period surrounding seizure. Using

indirect respiratory thermochemistry, we achieved a 24-hour monitoring to determine the

amount of oxygen consumed by the whole-body. Subsequently, we provide evidence of the

presence of abnormal $\dot{V}O_2$ pattern in 30-50% of epileptic rats, which started with the

installation of epilepsy (2-3 weeks post-SE). These EPI rats (EPI/SDOC+ group) presented with

chronic $\dot{V}O_2$ abnormalities characterized by low daily pattern among which intermittent and

repetitive SDOC events were observed.

Until now, both preclinical and clinical studies investigated changes in the metabolic rate of

O_2 in epilepsy mostly at the brain level, reporting a decline in the activity of mitochondrial

respiratory complexes in rats subjected to Pilo-SE and in patients with TLE [30, 31]. In our

study, we measured the whole-body O_2 intake which is mostly related to gas exchange

during ventilatory process [32]. However, whole-body O_2 intake can be also affected directly

or indirectly by several factors including age, circadian rhythm, body-weight, body-

temperature, cardiac function and finally activity [16, 32-37]. None of these factors was

found to be associated with abnormal pattern of $\dot{V}O_2$ in our study. We then focused our

attention on the ventilatory variables that may explain abnormal $\dot{V}O_2$ pattern in a subset of

EPI rats.

Respiratory system at rest in a subset EPI rats.

The identification of several groups within a single disease is a common feature in human

epilepsy, but rarely described in preclinical studies. One study has dichotomized two groups

among rats subjected to kainic acid-induced SE according to the extent of brain damage and

metabolic alterations [38]. In our study, EPI/SDOC+ subset has presented with a decrease of

daily $\dot{V}O2$ and a severe decrease of $\dot{V}E$ compared to EPI/SDOC- group.

During a 20-min monitoring period at rest, decreases of both $\dot{V}E$ and $\dot{V}O_2$ have already been

described as an inherent dysfunction in a genetic model of audiogenic seizure in the WAR rat

strain [26]. Because of technical limitations (24-hour $\dot{V}O_2$ monitoring), we could not jointly

assess $\dot{V}E$ and $\dot{V}O_2$ in our study, which prevented us from addressing the direct relationship

that may exist between $\dot{V}E$ and $\dot{V}O_2$. Nevertheless, we estimated for each rat the $\dot{V}E/\dot{V}O_2$

ratio as an indicator of ventilatory efficiency; this ratio was significantly higher in EPI/SDOC+

rats compared with EPI/SDOC- (+52%, P=0.007) and healthy control rats (+35%, P=0.04), and

not statistically different between these two latter rat groups.

It is difficult at this stage to interpret these results, especially in the context of epilepsy.

However, high $\dot{V}E/\dot{V}O_2$ has already been described as an index of poor gas exchange during

physical exercise of individual with heart failure disease [39]. Further exploration using blood

oxygen measurement will be required to determine the direct relationship between $\dot{V}E$ and

$\dot{V}O_2$ in epileptic rats following Pilo-SE. Another limitation of this study was the short duration

of the ventilatory recording compared with the 24 hours of $\dot{V}O_2$ assessments. Significant

fluctuations in $\dot{V}E$ could have been observed, similar to those seen for $\dot{V}O_2$, if the $\dot{V}E$

recordings had been longer. Further studies with devices allowing to concomitantly measure

in free-moving rats both $\dot{V}E$ and $\dot{V}O_2$ over sufficiently long periods of time will be needed to

establish a direct link between $\dot{V}O_2$ and $\dot{V}E$ profiles and specifically during SDOC events.

Indeed, these events could represent a disruption of O_2 uptake balance. But, to our

knowledge, events similar to SDOCs have not yet been described and we can only interpret

them in the light of what we know about a dyspneic situation. A first explanation for SDOCs

could be the presence of ventilatory pauses; however, they would have to be extremely

numerous and of long duration to induce such a collapse of $\dot{V}O_2$. A second explanation could

be a combination of several factors; for example, the absence of gasping, especially in

hypoxic conditions, could lead to poor tissue oxygenation [40]. In EPI/SDOC+ rats, the $\dot{V}E$

signal recording was too short in duration to determine whether the absence of gasping

could be part of the ventilatory dysregulation.

Ventilatory regulation in EPI rats.

We observed an increased hypoxia ventilatory response (HVR) in EPI rats, in contrast to

some studies that report no alteration in HVR in Pilo-SE rats [29, 41]. In addition, EPI/SDOC+

and EPI/SDOC- rats exhibited a sustained hypercapnia ventilatory response (HCVR) whereas

previous studies have reported a diminished response in Pilo-SE-induced epilepsy, rapid

amygdala kindling, and WAR rat models [26, 29, 42]. It is very difficult to compare our data

with those in the literature, due to the fact that there are differences in models, timing of

respiratory recordings, and severity of exposure to hypoxia and hypercapnia between our

study and all others. Given that low HVR and HCVR are generally indicative of alterations in

peripheral or central respiratory components, their maintenance in our study may indicate

that if alterations occurred in the early phases, then they were compensated for or repaired

in the late stages of epilepsy at which we performed our investigations.

Brainstem 5-HT alterations.

5-HT is mostly described as an excitatory drive toward respiratory network and subsequently

the motor output [43]. Dysregulation of brainstem neuronal groups involved in the control

of breathing, particularly those of the 5-HT system, has been found in patients with drug-

resistant epilepsy and has been associated with an increased risk of SUDEP [7]. We found

that brainstem 5-HT innervation was impaired in EPI rats compared to control rats, with

notable differences between EPI/SDOC+ and EPI/SDOC- rats. Indeed, EPI rats presented with

an opposite 5-HT innervation pattern within the NTS. Since the impairment of respiratory

function in the EPI/SDOC- rats seem to be moderate by opposition to EPI/SDOC+ rats, it is

conceivable that, in these rats, the increase of 5-HT innervation in both the NTS and VLM

may be part of protecting mechanisms of respiratory function. Such profile of 5-HT

innervation was lacking in EPI/SDOC+ rats, but these rats presented with an increase of

brainstem levels of both TPH2 and SERT transcripts. Such mechanisms in EPI/SDOC+ rats

might support an increased concentration of 5-HT in neuron terminals by increasing 5-HT

synthesis and re-uptake of intact 5-HT molecules, respectively, and thus may represent

adaptive mechanisms to compensate, although not sufficiently, for the decreased brainstem

5-HT innervation. This conclusion is supported by results reported in conditional knock-out

Lmx1b mice and WAR rats, which, as for EPI/SDOC+ rats, present similar respiratory

alterations at rest and concomitant decrease of 5-HT drive [26, 43]. We also believe

according to the difference of 5-HT innervation between VLM and NTS of the same

EPI/SDOC+ rats that the decreasing of 5-HT could be structure specific and could be present

in other primary regions of respiratory regulations.

We also compared brainstem 5-HT system of DBA/2 mice with FRA to their closest controls,

DBA/2 mice with NFRA, contrasting with previous studies that compared DBA/2 mice with

FRA to C57BL/6 mice resistant to AGS [10]. We found an increased level of mRNA coding for

TPH2 and SERT in DBA/2 mice with FRA. This 5-HT profile in DBA/2 mice with FRA is

intriguingly similar to that of EPI/SDOC+ rats. Altered 5-HT system is thus likely involved in

respiratory dysfunctions in both acute and chronic seizure models.

In conclusion, Pilo-SE rat model, which shows both chronic respiratory alterations and

spontaneous seizures, could be a relevant model to further explore neurobiological

mechanisms underlying chronic alteration of respiratory regulation in epilepsy. Identifying

the mechanisms that protect EPI/SDOC+ rats from mortality might also be an interesting

approach to pave the way for new therapeutic strategies to prevent SUDEP.

Methods

418   All animal procedures were in compliance with the guidelines of the European Union

419   regulating animal experimentation (directive 2010-63), taken in the French law (decree

420   2013-118), and have been approved by the ethical committee of the Claude Bernard Lyon 1

421   University (protocol # BH-2008-11).

Animals

Five week-old male Sprague–Dawley rats (Envigo, The Netherlands) were delivered to the

animal facility of the Lyon 1 University (Villeurbanne, France) and housed in groups of 5 in

transparent polycarbonate cages. Male DBA/2 mice of breeding colony of the LTSI laboratory

were housed in groups of five. Rats and mice were maintained under a controlled 12-h/12-h

light–dark cycle, in a temperature-controlled room (24-26 °C and 21-23 °C for rats and mice,

respectively), with free access to food and water.

Experimental design

Full description of methods have been described in details elsewhere [11, 28, 44-46]. Rats

subjected to Pilo-SE are daily observed to detect spontaneous seizures until animals had

developed epilepsy (EPI rats). Respiratory thermochemistry set-up was used to measure

oxygen consumption ($\dot{V}O_2$) during 24 consecutive hours (daily $\dot{V}O_2$) for each rat. Animals of

this study were subjected to one of following experiments:

Experiment 1. To establish the time course of respiratory alterations occurring among Pilo-

SE rats, 4 series of 10 rats were used in this experiment. In each series, Pilo-SE was induced

in 8 rats and 2 rats were used as controls. Daily $\dot{V}O_2$ of each rat was measured once a week

from the 1[st] to the 8[th] week post-SE.

Experiment 2. To further investigate cardio-respiratory functions in EPI rats during the

chronic phase of epilepsy, Pilo-SE was triggered in 30 rats and 16 rats were used as controls.

At the chronic period of epilepsy (12 weeks post-SE), the resting ventilation (normoxia, 21%

O_2) was measured using whole-body plethysmography. Between 13-14 weeks post-SE, a

subset of EPI and control rats were used to evaluate ventilatory responses to hypoxia (HVR,

FiO_2= 9%) and to hypercapnia (HCVR, $FiCO_2$= 5%), while others underwent surgery

implantation of telemetric device to evaluate cardiac frequency and body temperature at 15

weeks post-SE. As $\dot{V}O_2$ and ventilation are intermingled processes, to visualize global state of

respiratory system in EPI rats, we had in-depth our data exploitation by characterizing

ventilatory pattern according to $\dot{V}O_2$ pattern. To note, daily $\dot{V}O_2$ of EPI and control rats was

measured short before plethysmography at 7-9 weeks post-SE.

Experiment 3. To determine whether integrity of the 5-HT system is altered in the brainstem

of EPI rats presenting with respiratory alterations, Pilo-SE was induced in 20 rats and 8 rats

were used as controls. Daily $\dot{V}O_2$ was evaluated between the 6[th] and the 8[th] week post-SE.

Subsequently, animals were transcardially perfused and their brains were collected and

processed either for histological procedures (5-HT immunodetection; n=3 in each group) or

for gene expression analyses (SERT and TPH2 mRNA; n=5 in each group).

Experiment 4. To determine whether integrity of the 5-HT system is compromised in the

brainstem of DBA/2 mice presenting with FRA, 11 DBA/2 mice underwent AGS followed by

respiratory arrest. DBA/2 mouse brains were removed right after AGS for both FRA and non-

fatal respiratory arrest (NFRA) DBA/2 mice. Mice with FRA were resuscitated and as for

NFRA mice were injected with a lethal dose of pentobarbital and perfused with NaCl.

Brainstem SERT and TPH2 gene expressions were measured and then compared between

mice with either FRA or NFRA.

***In vivo* procedures**

Pilocarpine-induced SE. Epilepsy was triggered following Pilo-SE in 180-200 g male rats at

the age of 7 weeks, as previously described [28, 45]. Scopolamine methylnitrate (1 mg×kg^{-1},

s.c.; S-2250, Sigma-Aldrich) was administered 30 min prior to pilocarpine hydrochloride (350

mg×kg^{-1}, i.p.; P-6503, Sigma-Aldrich). After an initial period of immobility, the onset of SE

was characterized by repetitive tonic-clonic activity of the trunk and the limbs, occurring

after repeated rearing with forelimb clonus and falling. Because SE event is a life-threating

condition in rats, there is a need to stop it after 2 hours by a single injection of diazepam

(Valium®, 10 mg×kg^{-1}, i.p.; Roche). Rats were then hydrated with 2 mL of saline solution

(0.9% NaCl; s.c.) and transferred to individual cages. After a one-week recovery period, all

rats (control and treated rats) were housed in groups of 5 to avoid any stress that may result

from social isolation.

Animal care after Pilo-SE. Control and treated rats were weighted daily during the first two

weeks following Pilo-SE, and then every week until termination of the experiment. Daily

abdominal massage was performed twice a day during the first week to activate intestinal

motility, which is disrupted following Pilo-SE.

Detection of spontaneous recurrent seizures after Pilo-SE. To maintain as low as possible

the level of anxiety in rats subjected to Pilo-SE, electro-encephalographic recordings (EEG)

were not performed to detect the onset of epilepsy in our experiments. Indeed,

conventional EEG set-up requires that rats are housed individually in a restrictive

environment, which is known to enhance stress and anxiety and worsens epilepsy

phenotype in Pilo-SE models [47]. In addition, as mentioned below in the "whole-body

plethysmography" section, it is difficult to perform ventilatory recordings in animals with

high anxiety level.

It is now clearly established that rats of the Sprague-Dawley strain from the Janvier

Laboratories develop spontaneous recurrent seizures between 10 and 36 days after

induction of Pilo-SE at 42 days of age [48, 49]. Because we were using Sprague-Dawley rats

from Envigo Laboratories, we had to monitor the time of onset of spontaneous recurrent

seizures (SRSs) in our experiments. To determine whether rats were in the chronic phase of

epilepsy, they were manipulated daily between 09:00 and 11:00 a.m. to define the onset of

handling-induced seizures (HIS), which was induced by restraining rats at the level of the

chest with gentle pressure for 10 seconds [48]. HIS were used in our study to help

experimenters to define the time to start detecting the first behavioural spontaneous

recurrent seizures (SRSs). Thus, upon detection of 2 HIS, rats were observed by

experimenters for 5 consecutive hours (between 01:00 and 06:00 p.m.) over the following

days to detect the presence or absence of behavioural SRSs. Following Pilo-SE, rats are

considered epileptic once they developed at least two SRSs during the periods of the daily

observations. The severity of these seizures was scored from 3 to 5 on the Racine's scale [50].

Seizure severity of less than 3 was difficult to detect visually in this study, which was

conducted without daily video monitoring. Stage 3 corresponds to forelimb clonus, stage 4

to bilateral forelimb clonus with rearing, and stage 5 to rearing and falling or to tonic-clonic

seizures.

During the 24 hour-$\dot{V}O2$ recordings, rats were video-monitored for 12h during the light

period, making it possible to evaluate the number of seizures and their severity per session.

Evaluation of respiratory and cardiac functions

Open respiratory thermochemistry. To detect respiratory alterations in epileptic rats, we

used respiratory thermochemistry as an alternative approach to plethysmography, allowing

respiratory function to be recorded for a long period of time on fed, unrestrained and

unanesthetized animal. Respiratory thermochemistry mostly reflects gas exchanges and

allows more specifically the measurement of the oxygen consumption ($\dot{V}O_2$) with no bias

effect from animal's body movements on the accuracy of respiratory acquisition.

Oxygen consumption ($\dot{V}O_2$) was measured with sample rate of 3 acquisitions per minute,

under standard conditions (variation in temperature = 0°C; Ambient pressure = 760 mm Hg;

dry air) using an open respiratory thermochemistry system, as previously described [51]. $\dot{V}O_2$

was continuously recorded during 24 consecutive hours on the same thermochemistry

system; therefore, only one rat could be recorded each day in our experimental setting.

Briefly, rats were placed individually in a thermostatic recording chamber (12 L) with steady

airflow (4 L × min^{-1}), under a controlled 12-h/12-h light–dark cycle (light from 06:00 a.m. to

06:00 p.m.), and were allowed free access to food and water. Ambient temperature inside

the thermostatic chamber was monitored with copper-constantan thermocouples and was

maintained at 24-26 °C using a thermostatic container. Airflow rate was measured using a

Platon volumeter and converted to standard STPD (standard temperature and pressure, dry)

values. Fractional O_2 concentration was measured using a Servomex 1100 paramagnetic gas

analyzer (Taylor Instrument Analytics ltd, Sussex, UK) and $\dot{V}O_2$ was calculated. The O_2

analyzer was calibrated daily with pure nitrogen gas (0% O_2) and room air (20.93% O_2). Rats

were weighted before and after recording then $\dot{V}O_2$ values were normalized to the average

body weight (calculated from the weight measured before and after each recording session).

Rats were video-monitored during the daylight period.

Whole-body plethysmography. Plethysmographic chamber allows collecting ventilatory

variables from acquired breathing pattern signal. Ventilatory function was evaluated

between 09:00 a.m. and 04:00 p.m. in unrestrained, unanesthetized rats under BTPS (body

temperature and pressure, saturated) conditions using whole-body plethysmography

(EMKA; technology). Calibration of the system was performed before each recording session

by injecting a 1 mL air pulse inside the recording chamber. Briefly, each rat was individually

placed in the recording chamber (600 mL) that is connected to a constant bias flow supply

providing continuous inflow of fresh air (2.5 L × min^{-1}). Minute volume ($\dot{V}_E$; mL × min^{-1}) was

calculated as the product of tidal volume (V_T; mL) and respiratory frequency (f_R; breath x

min^{-1}). Total respiratory cycle duration (T_{tot}) was calculated as the sum of the inspiratory

time (T_I; msec) and expiratory time (T_E; msec). We also calculated for each animal the

coefficient of variation of the respiratory frequency, which is an index of the variability of

the respiratory pattern [52] and the frequency of spontaneous apnea, which is defined as a

cessation of breathing lasting more than two respiratory cycles (>2 x T_{tot}). Apnea number

and duration were calculated. Apnea frequency was normalized to recording duration and

expressed per min of breathing. Recordings were started after a 40-minute habituation

period. Then, resting ventilation was monitored during 20 minutes under normoxic condition

(21% O_2). After recordings, rats were brought back to their home cage.

Ventilatory responses to hypoxia (FiO_2= 9%) and to hypercapnia ($FiCO_2$= 5%) were evaluated

during two distinct recording sessions, which were performed a week apart from each other.

After a 40-minute habituation period, plethysmographic recordings under basal condition

(normoxia) were started 10 minutes before gas variations. Ventilatory responses to a 10-

minute exposure to hypoxia or hypercapnia were subsequently evaluated. All

plethysmography recordings were made between 09:00 a.m. and 04:00 p.m.; because,

beyond 04:00 p.m., EPI rats were more stressed in the plethysmography chamber, requiring

longer time to reach a quiet state in this novel and restricted environment. At the end of

each recording session, rats were brought back to their home cage. Repeated measurements

are randomly performed for each rat.

Abdominal implantation of telemetry transmitters to evaluate cardiac function and body-

temperature. Rats were anesthetized with a mix of ketamine and xylazine (75 mg × kg^{-1} and

25 mg × kg^{-1}, respectively; i.p.). A midline abdominal incision was made and a sterile

telemetry transmitter (Data Sciences International, France) was placed inside the abdominal

cavity. A pair of biocompatible wire electrodes originating from the transmitter were

tunnelled subcutaneously and implanted to the left intercostal muscles and between the

right shoulder blades. The transmitter was attached to the abdominal wall during wound

closure with absorbable sutures. After surgery, animals received an intramuscular injection

of penicillin (15,000 units) once a day for 3 consecutive days and were then returned to their

home cage. Rats were monitored daily to control general health state. Two weeks after

surgery, telemetry transmitters were magnetically activated and freely moving animals were

placed in individual cages placed under platform-receiver units. Variation in cardiac

frequency and body-temperature were recorded during 24 consecutive hours.

Induction of audiogenic seizures in DBA/2 mice

AGS were induced as previously described [11, 17], with slight modifications. Briefly, at age

of 23 days, mice were individually placed in a plastic cylinder (40 cm diameter, 20 cm height)

and exposed to a broadband acoustic stimulus (intensity of 110 dB SPL with a peak

frequency at 12 kHz) generated by a mechanical sonicator (Deltasonic model Delta 920). The

stimulus was given for a maximum duration of 60 seconds or until the mouse exhibited AGS,

characterized by an initial wild running and ending by a tonic hindlimb extension. All mice

exhibited a tonic hindlimb extension leading to a respiratory arrest identified by pinnae

relaxation and a complete cessation of chest movements. Respiratory arrest was followed by

spontaneous recovery of the respiratory function within 5 seconds in mice identified as

NFRA, or was followed by death for the other mice (FRA). Mouse brain was systematically

removed within 2-3 min after acoustic stimulus was triggered.

Ex vivo **procedures**

DBA/2 mice and rats were injected with a lethal dose of pentobarbital (250 mg × kg^{-1}; i.p.)

before being sacrificed. For gene expression analyses, animals were transcardially perfused

(30 mL x min^{-1}) for 2 min with ice-cold saline and their brain was rapidly removed. Then, the

brainstem region was isolated, frozen in liquid nitrogen and stored at -80°C. For

immunohistochemistry procedure, animals were transcardially perfused (30 mL x min^{-1}) for 2

min with ice-cold saline and then for 10 min with ice-cold 4% paraformaldehyde in 0.2 M

phosphate buffer (30 mL x min^{-1}). After cryoprotection in a 25% sucrose solution, brains

were frozen at -40°C in isopentane and stored at -80°C.

5-HT immunohistochemistry. Coronal sections (40 µm thick) were cryostat-cut throughout

the medulla oblongata and were processed free-floating for 5-HT immunostaining. Using the

rat brain atlas, sections from each experimental group were selected at interaural

coordinates -2.60 mm, and were all processed together for 5-HT immunofluorescent labeling.

Selected sections were rinsed in 0.1M Phosphate Buffered Saline (PBS) and incubated in PBS

containing 0.3% Triton X-100 and 3% normal donkey serum. Then, sections were incubated

with a rabbit polyclonal antibody against 5-HT (Sigma; S5545; final dilution = 1:3,000). After

few washes, sections were incubated with an AlexaFluor-488 conjugated donkey anti-rabbit

antibody (Molecular Probes; A-21206; final dilution = 1:1,000). At the end of the

immunostaining protocol, sections were mounted onto SuperFrost®Plus slides and cover

slipped using Vectashield mounting medium (H-1500, Vector).

Semi-quantification of brainstem 5-HT immunostaining. Images (1152 x 1024 pixels) were

captured with a HLX PL APO x63/0.70 objective and with the same conditions of

photomultiplier gain, offset and pinhole aperture using a TCS SP2 confocal microscopy

system (Leica), allowing the comparison of fluorescence intensity between sections. Image

analysis was conducted on an average of 8 z-compressed images stacks (each image = 4 μm

depth), which include the whole region of interest. Using the image J software (NIH), we first

measured the average grey value within each 5-HT-positive neuronal cell body of caudal

raphe nuclei (pallidus, magnus and obscurus), as an index of 5-HT neuronal concentration

(A.U.). Using the same software, we also measured the density of 5-HT-immunopositive

processes within the surface area of the NTS and the VLM, expressed as 5-HT-positive pixel/

pixel^2.

Quantification of transcript level variation by RT-qPCR. Total RNA was extracted from

brainstem samples using Tri-reagent LS (Molecular Research Center, Inc.). Contaminant

genomic DNA was subsequently removed from the samples by treatment with Turbo DNA-

free™ kit (Ambion). The messenger RNAs (mRNAs) contained in 1 µg of purified RNA extracts

were reverse-transcribed using the reverse transcriptase RNase H minus (Promega) and

oligo-d(T)15. To normalize the RT step, a synthetic external and non-homologous poly(A)

standard RNA (SmRNA; Morales and Bezin, patent WO2004.092414) was added to the RT

reaction mix (150,000 copies in each experimental sample), as previously described [28, 44-

46]. PCR amplification of selected cDNAs, i.e. those complementary to the mRNA encoding

rate-limiting enzyme of brainstem 5-HT synthesis (tryptophan hydroxylase: TPH2) and the 5-

HT transporter (SERT), were performed using the Rotor-Gene Q system (Qiagen) and the

QuantiTect SYBR Green PCR Kit (Qiagen). Sequences of the different primer pairs used for

PCR amplification of mouse and rat TPH2 and SERT cDNAs are listed in Supplementary Table

S1.

Statistical analysis

The data for each experimental group were pooled and expressed as the mean ± SEM in the

text and figures, unless otherwise specified. All statistics were performed using Sigma plot

software 3.5 (Systat Software Inc., Point Richmond, California, USA). A Kolmogorov-Smirnov

test was used to test if the data follows a normal distribution, then paired or unpaired t-test

was performed for the two-group comparison. For multiple group comparisons, one-way

ANOVA followed by post-hoc Fisher's LSD test was used. Statistical significance was

predefined as $P < 0.05$.

Authorship

L.B and S.R. contributed equally to the article. L.B., S.R and P.R conceived and designed the experiments. A.M., J.R. and H.K performed respiratory thermochemistry experiments. H.K. performed plethysmography and telemetry experiments. B.M., G.D., L.B. and H.K., performed AGS in DBA/2 mice. H.K. and B.G. performed histochemistry and RT-qPCR experiments. L.B., S.R., M.O. and H.K. analyzed data. L.B., S.R. and H.K. wrote the paper.

Acknowledgements

We thank for their assistance: Frédérique Cohen-Adad (scientific documentation; TIGER team, CRNL), and Denis Ressnikoff (confocal microscopy; Centre Commun de Quantimétrie of the Claude Bernard Lyon 1 University). Benoît Martin has received subventions from Eisai, UCB-Pharma and Zogenix.

Conflicts of interest

None of the authors has any conflict of interest to disclose. We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this report is consistent with those guidelines.

References

1. Lacuey, N., et al., The incidence and significance of periictal apnea in epileptic seizures. Epilepsia, 2018. **59**(3): p. 573-582.
2. Bourdillon, P., et al., Letter to the Editor. Temporal lobe epilepsy: open or stereotactic surgery? J Neurosurg, 2019. **131**(3): p. 989.
3. Ryvlin, P., et al., Incidence and mechanisms of cardiorespiratory arrests in epilepsy monitoring units (MORTEMUS): a retrospective study. Lancet Neurol, 2013. **12**(10): p. 966-77.

4. Vilella, L., et al., Incidence, Recurrence, and Risk Factors for Peri-ictal Central Apnea and
Sudden Unexpected Death in Epilepsy. Front Neurol, 2019. **10**: p. 166.

5. Sivathamboo, S., et al., Sleep-disordered breathing in epilepsy: epidemiology,
mechanisms, and treatment. Sleep, 2018. **41**(4).

6. Somboon, T., M.M. Grigg-Damberger, and N. Foldvary-Schaefer, Epilepsy and Sleep-
Related Breathing Disturbances. Chest, 2019. **156**(1): p. 172-181.

7. Patodia, S., et al., The ventrolateral medulla and medullary raphe in sudden unexpected
death in epilepsy. Brain, 2018. **141**(6): p. 1719-1733.

8. Garcia, A.J., 3rd, et al., Chapter 3--networks within networks: the neuronal control of
breathing. Prog Brain Res, 2011. **188**: p. 31-50.

9. Faingold, C.L., et al., Differences in serotonin receptor expression in the brainstem may
explain the differential ability of a serotonin agonist to block seizure-induced sudden
death in DBA/2 vs. DBA/1 mice. Brain Res, 2011. **1418**: p. 104-10.

10. Uteshev, V.V., et al., Abnormal serotonin receptor expression in DBA/2 mice associated
with susceptibility to sudden death due to respiratory arrest. Epilepsy Res, 2010. **88**(2-
3): p. 183-8.

11. Martin, B., et al., Audiogenic seizure as a model of sudden death in epilepsy: A
comparative study between four inbred mouse strains from early life to adulthood.
Epilepsia, 2020.

12. Faingold, C.L., S. Tupal, and M. Randall, Prevention of seizure-induced sudden death in a
chronic SUDEP model by semichronic administration of a selective serotonin reuptake
inhibitor. Epilepsy Behav, 2011. **22**(2): p. 186-90.

13. Tupal, S. and C.L. Faingold, Evidence supporting a role of serotonin in modulation of
sudden death induced by seizures in DBA/2 mice. Epilepsia, 2006. **47**(1): p. 21-6.

14. Curia, G., et al., The pilocarpine model of temporal lobe epilepsy. J Neurosci Methods,
2008. **172**(2): p. 143-57.

15. Biagini, G., et al., Proepileptic influence of a focal vascular lesion affecting entorhinal
cortex-CA3 connections after status epilepticus. J Neuropathol Exp Neurol, 2008. **67**(7):
p. 687-701.

16. Feher, J.J., Quantitative human physiology : an introduction. Second edition. ed.
Academic Press series in biomedical engineering. 2017, Amsterdam ; Boston:
Elsevier/AP, Academic Press is an imprint of Elsevier. xiii, 968 pages.

17.   Venit, E.L., B.D. Shepard, and T.N. Seyfried, Oxygenation prevents sudden death in
seizure-prone mice. Epilepsia, 2004. **45**(8): p. 993-6.

18.   Aiba, I. and J.L. Noebels, Spreading depolarization in the brainstem mediates sudden
cardiorespiratory arrest in mouse SUDEP models. Sci Transl Med, 2015. **7**(282): p.
282ra46.

19.   Jefferys, J.G.R., et al., Brainstem activity, apnea, and death during seizures induced by
intrahippocampal kainic acid in anaesthetized rats. Epilepsia, 2019. **60**(12): p. 2346-2358.

20.   Faingold, C.L., M. Randall, and S. Tupal, DBA/1 mice exhibit chronic susceptibility to
audiogenic seizures followed by sudden death associated with respiratory arrest.
Epilepsy Behav, 2010. **17**(4): p. 436-40.

21.   Johnston, S.C., et al., Central apnea and acute cardiac ischemia in a sheep model of
epileptic sudden death. Ann Neurol, 1997. **42**(4): p. 588-94.

22.   Kim, Y., et al., Severe peri-ictal respiratory dysfunction is common in Dravet syndrome. J
Clin Invest, 2018. **128**(3): p. 1141-1153.

23.   Lertwittayanon, W., O. Devinsky, and P.L. Carlen, Cardiorespiratory depression from
brainstem seizure activity in freely moving rats. Neurobiol Dis, 2019. **134**: p. 104628.

24.   Nakase, K., et al., Laryngospasm, central and obstructive apnea during seizures: Defining
pathophysiology for sudden death in a rat model. Epilepsy Res, 2016. **128**: p. 126-139.

25.   St-John, W.M., et al., Changes in respiratory-modulated neural activities, consistent with
obstructive and central apnea, during fictive seizures in an in situ anaesthetized rat
preparation. Epilepsy Res, 2006. **70**(2-3): p. 218-28.

26.   Totola, L.T., et al., Impaired central respiratory chemoreflex in an experimental genetic
model of epilepsy. J Physiol, 2017. **595**(3): p. 983-999.

27.   Villiere, S.M., et al., Seizure-associated central apnea in a rat model: Evidence for
resetting the respiratory rhythm and activation of the diving reflex. Neurobiol Dis, 2017.
101: p. 8-15.

28.   Sanchez, P.E., et al., Optimal neuroprotection by erythropoietin requires elevated
expression of its receptor in neurons. Proc Natl Acad Sci U S A, 2009. **106**(24): p. 9848-53.

29.   Maia, O.A.C., et al., Pilocarpine-induced status epilepticus reduces chemosensory
control of breathing. Brain Res Bull, 2020. **161**: p. 98-105.

30.   Kudin, A.P., et al., Seizure-dependent modulation of mitochondrial oxidative
phosphorylation in rat hippocampus. Eur J Neurosci, 2002. **15**(7): p. 1105-14.

31. Kunz, W.S., et al., Mitochondrial complex I deficiency in the epileptic focus of patients
with temporal lobe epilepsy. Ann Neurol, 2000. **48**(5): p. 766-73.

32. Gastinger, S., et al., A comparison between ventilation and heart rate as indicator of
oxygen uptake during different intensities of exercise. J Sports Sci Med, 2010. **9**(1): p.
110-8.

33. Gautier, H., Interactions among metabolic rate, hypoxia, and control of breathing. J Appl
Physiol (1985), 1996. **81**(2): p. 521-7.

34. Refinetti, R., Metabolic heat production, heat loss and the circadian rhythm of body
temperature in the rat. Exp Physiol, 2003. **88**(3): p. 423-9.

35. Refinetti, R., The circadian rhythm of body temperature. Front Biosci (Landmark Ed),
2010. **15**: p. 564-94.

36. Seifert, E.L. and J.P. Mortola, Circadian pattern of ventilation during acute and chronic
hypercapnia in conscious adult rats. Am J Physiol Regul Integr Comp Physiol, 2002.
282(1): p. R244-51.

37. Stephenson, R., et al., Circadian rhythms and sleep have additive effects on respiration
in the rat. J Physiol, 2001. **536**(Pt 1): p. 225-35.

38. Pearce, P.S., et al., Metabolic injury in a variable rat model of post-status epilepticus.
Epilepsia, 2016. **57**(12): p. 1978-1986.

39. Luks, A., R.W. Glenny, and H.T. Robertson, Introduction to cardiopulmonary exercise
testing. 2013, New York: Springer. vii, 147 pages.

40. Poets, C.F., et al., Gasping and other cardiorespiratory patterns during sudden infant
deaths. Pediatr Res, 1999. **45**(3): p. 350-4.

41. Campos, R.R., F.R. Tolentino-Silva, and L.E. Mello, Respiratory pattern in a rat model of
epilepsy. Epilepsia, 2003. **44**(5): p. 712-7.

42. Totola, L.T., et al., Amygdala rapid kindling impairs breathing in response to chemoreflex
activation. Brain Res, 2019. **1718**: p. 159-168.

43. Hodges, M.R. and G.B. Richerson, Interaction between defects in ventilatory and
thermoregulatory control in mice lacking 5-HT neurons. Respir Physiol Neurobiol, 2008.
164(3): p. 350-7.

44. Morales, A., et al., Unexpected expression of orexin-B in basal conditions and increased
levels in the adult rat hippocampus during pilocarpine-induced epileptogenesis. Brain
Res, 2006. **1109**(1): p. 164-75.

45. Nadam, J., et al., Neuroprotective effects of erythropoietin in the rat hippocampus after
pilocarpine-induced status epilepticus. Neurobiol Dis, 2007. **25**(2): p. 412-26.

46. Navarro, F.P., et al., Brain heparanase expression is up-regulated during postnatal
development and hypoxia-induced neovascularization in adult rats. J Neurochem, 2008.
105(1): p. 34-45.

47. Manouze, H., et al., Effects of Single Cage Housing on Stress, Cognitive, and Seizure
Parameters in the Rat and Mouse Pilocarpine Models of Epilepsy. eNeuro, 2019. **6**(4).

48. Dube, C., et al., Relationship between neuronal loss and interictal glucose metabolism
during the chronic phase of the lithium-pilocarpine model of epilepsy in the immature
and adult rat. Exp Neurol, 2001. **167**(2): p. 227-41.

49. Leroy, C., et al., Temporal changes in mRNA expression of the brain nutrient
transporters in the lithium-pilocarpine model of epilepsy in the immature and adult rat.
Neurobiol Dis, 2011. **43**(3): p. 588-97.

50. Racine, R.J., Modification of seizure activity by electrical stimulation. II. Motor seizure.
Electroencephalogr Clin Neurophysiol, 1972. **32**(3): p. 281-94.

51. Belouze, M., et al., Leanness of Lou/C rats does not require higher thermogenic capacity
of brown adipose tissue. Physiol Behav, 2011. **104**(5): p. 893-9.

52. Ogier, M., et al., Brain-derived neurotrophic factor expression and respiratory function
improve after ampakine treatment in a mouse model of Rett syndrome. J Neurosci,
2007. **27**(40): p. 10912-7.

Figure legends

Figure 1: Detection of SDOC events during a 24-hour monitoring of oxygen consumption in a subgroup of epileptic rats. Oxygen consumption ($\dot{V}O_2$) (light period from 06:00 am to 6:00 pm) was measured using respiratory thermochemistry. Grey bar above each x-axis indicates the dark period. A single profile was observed in control rats (A), while two distinct profiles were distinguished in epileptic rats (B, C). The first profile was similar to that observed in control rats (B), the second one was characterized by several episodes of sharp decrease in oxygen consumption (SDOC) (C), sometimes reaching the zero value. These SDOC events were only seen in epileptic rats, thereby discriminating two groups of epileptic rats upon their presence (EPI/SDOC+) or their absence (EPI/SDOC-).

Figure 2: SDOC can occur independently of behavioral spontaneous seizures. (A) Example of a SDOC event recorded in an EPI rat at 5 weeks post-SE, when it was adopting a sleep posture. This SDOC event occurred during the light period and was identified with the decline in $\dot{V}O_2$ values. No behavioral seizure was observed before, during and after the SDOC occurrence. By contrast, (B) a sharp increase in $\dot{V}O_2$ values was observed after a 43-second stage-5 seizure according to Racine's scale in an EPI/SDOC+ rat whose values of $\dot{V}O_2$ were above pre-ictal level for about 30 minutes.

Figure 3: Time course of daily $\dot{V}O_2$ pattern. Respiratory thermochemistry was recorded once a week for each rat, from the 1st to the 8th week post-SE. (A) Daily $\dot{V}O_2$ is calculated for each group (control rats (n=8), EPI/SDOC- rats (n=14), EPI/SDOC+ rats (n=6)) from the 1 to 8 weeks post-SE. (B) The daily $\dot{V}O_2$ of animals is calculated at 3-4 weeks post-SE in animals identified or not yet as EPI/SDOC+ rats. Paired comparison of daily $\dot{V}O_2$ over weeks showed statistical significance with only 1-2 weeks post-SE. Results are expressed in mean ± SEM of

̇VO2, *, p<0.05; **, P=0.01 in comparison of daily $\dot{V}O_2$ at 1-2 weeks post-SE. (C) the first

SDOC was detected in one rat (1/20) during the 3rd post-SE week. The maximum number of

EPI rats exhibiting SDOCs (EPI/SDOC+, 30%) was reached by the end of the 5th week post-SE.

The proportion of EPI/SDOC+ rats was expressed in percent of the total number of rats

developing epilepsy (yellow curve) or in percent of the total number of rats presenting with

SDOCs by the end of the 8th post-SE (red curve).

Figure 4: Semi-quantification of 5-HT immunopositive terminals of caudal raphe nuclei

within the medulla oblongata of control and EPI rats. Fluorescent 5-HT labeling was

performed on sections selected at interaural -2.60 mm within two brainstem regions

involved in the regulation of respiratory rhythm generation: the ventrolateral medulla (VLM)

and nucleus tractus solitarius (NTS). Images of 5-HT positive fibers were captured within the

VLM (A1, A2, A3) and the NTS (B1, B2, B3) using confocal microscopy (x63 objective). Density

of 5-HT positive fibers was calculated in the VLM (A4) and the NTS (B4). Data are expressed

as the mean ± SEM (n=3 per group). *, P<0.05; ***, P<0.001, compared to controls and †††,

P<0.001, compared to EPI/SDOC-.

Figure 5: TPH2 and SERT mRNA transcript levels were measured in the brainstem of EPI

rats and DBA/2 mice. Changes in brainstem levels of TPH2 (A) and SERT (B) transcripts were

measured in EPI rats and DBA/2 mice at 8 weeks post-SE and 2-3 minutes after an

audiogenic seizure (AGS), respectively. In contrast to EPI rats, DBA/2 mice had no history of

seizure before AGS. Some mice developed a fatal respiratory arrest (DBA/2 with FRA) or a

non-fatal respiratory arrest (DBA/2 with NFRA) after an AGS. TPH2 and SERT transcript levels

in EPI/SDOC+ rats and in DBA/2 with FRA mice are expressed in percentage (mean ± SEM) of

EPI/SDOC- rats and DBA/2 mice with NFRA, respectively (n=5 in each group). †: P<0.05, ††:

P<0.01, †††: P<0.001, in comparison to EPI/SDOC- group for rat and DBA/2 with FRA group

for mice. *, P<0.05; **, P<0.01, in comparison to control rats.

Table legend

Table 1: Monitoring of cardio-respiratory functions in control and EPI rats between 12 and

15 weeks post-SE. Using plethysmography, the ventilatory variables were measured under

normoxic condition (21% O_2) at 12 weeks post-SE (A) and in response to hypoxia (9% O_2) or

hypercapnia (5% CO_2) exposures between 13-14 weeks post-SE (B). Cardiac frequency (C)

and body-temperature (D) were concomitantly evaluated at 15 weeks post-SE.

Abbreviations: Fc, cardiac frequency; f_R, respiratory frequency; T_B, body-temperature; T_E,

expiration time; T_I, inspiration time; T_{tot}, total respiratory cycle duration; $\dot{V}_E$, minute volume;

V_T, tidal volume. Statistics: NT, not statistically significant.

Supplementary information

Figure S1: Metabolic rate of oxygen at 1-2 weeks and 5-8 weeks post-SE. The average

oxygen consumption ($\dot{V}O2$) was calculated for the light and dark periods at 5-8 weeks post-

SE (A) and then at 1-2 weeks post-SE (B) using the same experimental groups,

retrospectively. Dichotomization of SDOC+ and SDOC- EPI rats was based on data collected

at 5-8 weeks post-SE, which showed a significant decline in the average $\dot{V}O2$ of EPI/SDOC+

rats during both the light and dark periods compared to EPI/SDOC- and control rats (A).

There was no difference the three rat groups at 1-2 weeks post-SE, during which time no

SDOCs were detected (B). Results are expressed in percent of control rats *, P<0.05; ***,

p<0.001, compared to control rats. †††, P<0.001, compared to EPI/SDOC- rats.

Figure S2: Ventilatory function in epileptic and control rats under normoxia. (A) ventilation

(V_E), respiratory frequency (F_R) and tidal volume (V_T) of epileptic (n=16) and control (n=5)

rats at 12 weeks post-*SE.* (B) Index of pattern stability includes coefficient of variation of

ventilatory pattern and apnea number/duration. The results are expressed as the mean ±

SEM. *, P<0.05; **, P<0.01, in comparison to control rats.

Supplementary Table S1: Primer pairs used for qPCR amplification of TPH2 and SERT

cDNAs of rat and mouse.

Key Points

• There are mild and severe patterns of respiratory dysfunctions among rats with epilepsy

developed after pilocarpine induced *status epilepticus*.

• Severe respiratory dysfunctions were evidenced in 30-50% of rats with epilepsy.

• These dysfunctions consisted with intermittent episodes of sharp decrease in oxygen

consumption (SDOC), slow breathing pattern and low metabolic rate of oxygen at rest.

• SDOCs were associated with decreased density of 5-HT-containing processes in the

nucleus tractus solitarius.

• Rats with SDOCs had SERT and TPH2 transcript level alterations similar to DBA/2 mice

subjected to AGS-induced fatal respiratory arrest.

Figure 1: Detection of SDOC events during a 24-hour monitoring of oxygen consumption in a subgroup of epileptic rats. Oxygen consumption ($\dot{V}O_2$) (light period from 06:00 am to 6:00 pm) was measured using respiratory thermochemistry. Grey bar above each x-axis indicates the dark period. A single profile was observed in control rats (A), while two distinct profiles were distinguished in epileptic rats (B, C). The first profile was similar to that observed in control rats (B), the second one was characterized by several episodes of sharp decrease in oxygen consumption (SDOC) (C), sometimes reaching the zero value. These SDOC events were only seen in epileptic rats, thereby discriminating two groups of epileptic rats upon their presence (EPI/SDOC+) or their absence (EPI/SDOC-).

A. **Epileptic rat during sleep**

B. **Rat undergoing stage-5 seizure**

Figure 2: SDOC can occur independently of behavioral spontaneous seizures. (A) Example of a SDOC event recorded in an EPI rat at 5 weeks post-SE, when it was adopting a sleep posture. This SDOC event occurred during the light period and was identified with the decline in $\dot{V}O_2$ values. No behavioral seizure was observed before, during and after the SDOC occurrence. By contrast, (B) a sharp increase in $\dot{V}O_2$ values was observed after a 43-second stage-5 seizure according to Racine's scale in an EPI/SDOC+ rat whose values of $\dot{V}O_2$ were above pre-ictal level for about 30 minutes.

A. Time course of the daily V̇O2 in experimental groups

Weeks post-SE	Daily V̇O2 (mL x min^{-1} x kg^{-1})		
	Control rats (n=8)	EPI/SDOC- rats (n=14)	EPI/SDOC+ rats (n=6)
8	19.2 ± 0.6 *	18.0 ± 0.5 *	8.1 ± 0.3 **
7	18.1 ± 0.4 *	19.8 ± 0.5	10.1 ± 1.6 *
6	20.0 ± 0.4	19.9 ± 0.8	11.0 ± 2.5 $^{p=0.09}$
5	19.8 ± 0.1	22.3 ± 1.2	12.5 ± 1.6 *
4	20.8 ± 0.5	20.1 ± 0.6	
3	21.2 ± 0.9	21.0 ± 0.8	
1-2	24.4 ± 1.1	23.1 ± 1.7	26.6 ± 1.0

B. Time course of the daily V̇O2 in EPI/SDOC+ rats

Weeks post-SE	Rats identified as EPI/SDOC+		Rats not yet identified as EPI/SDOC+	
	n	Daily V̇O2 (mL x min^{-1} x kg^{-1})	n	Daily V̇O2 (mL x min^{-1} x kg^{-1})
4	3/6	13.5 ± 3.5	3/6	20.1 ± 1.8
3	1/6	17.2	5/6	21.1 ± 1.9

C. Time course of SDOC appearance by EPI rats

Figure 3: Time course of daily V̇O$_2$ pattern. Respiratory thermochemistry was recorded once a week for each rat, from the 1st to the 8th week post-SE. (A) Daily V̇O$_2$ is calculated for each group (control rats (n=8), EPI/SDOC- rats (n=14), EPI/SDOC+ rats (n=6)) from the 1 to 8 weeks post-SE. (B) The daily V̇O$_2$ of animals is calculated at 3-4 weeks post-SE in animals identified or not yet as EPI/SDOC+ rats. Paired comparison of daily V̇O$_2$ over weeks showed statistical significance with only 1-2 weeks post-SE. Results are expressed in mean ± SEM of V̇O2, *, p<0.05; **, p=0.01 in comparison of daily V̇O$_2$ at 1-2 weeks post-SE. (C) the first SDOC was detected in one rat (1/20) during the 3rd post-SE week. The maximum number of EPI rats exhibiting SDOCs (EPI/SDOC+, 30%) was reached by the end of the 5th week post-SE. The proportion of EPI/SDOC+ rats was expressed in percent of the total number of rats developing epilepsy (yellow curve) or in percent of the total number of rats presenting with SDOCs by the end of the 8th post-SE (red curve).

Figure 4: Semi-quantification of 5-HT immunopositive terminals of caudal raphe nuclei within the medulla oblongata of control and EPI rats. Fluorescent 5-HT labeling was performed on sections selected at interaural -2.60 mm within two brainstem regions involved in the regulation of respiratory rhythm generation: the ventrolateral medulla (VLM) and nucleus tractus solitarius (NTS). Images of 5-HT positive fibers were captured within the VLM (A1, A2, A3) and the NTS (B1, B2, B3) using confocal microscopy (x63 objective). Density of 5-HT positive fibers was calculated in the VLM (A4) and the NTS (B4). Data are expressed as the mean ± SEM (n=3 per group). *, p<0.05; ***, p<0.001, compared to controls and †††, p<0.001, compared to EPI/SDOC-.

A. Level expression of mRNA encoding TPH2

B. Level expression of mRNA encoding SERT

Figure 5: TPH2 and SERT mRNA transcript levels were measured in the brainstem of EPI rats and DBA/2 mice. Changes in brainstem levels of TPH2 (A) and SERT (B) transcripts were measured in EPI rats and DBA/2 mice at 8 weeks post-SE and 2-3 minutes after an audiogenic seizure (AGS), respectively. In contrast to EPI rats, DBA/2 mice had no history of seizure before AGS. Some mice developed a fatal respiratory arrest (DBA/2 with FRA) or a non-fatal respiratory arrest (DBA/2 with NFRA) after an AGS. TPH2 and SERT transcript levels in EPI/SDOC+ rats and in DBA/2 with FRA mice are expressed in percentage (mean ± SEM) of EPI/SDOC- rats and DBA/2 mice with NFRA, respectively (n=5 in each group). †: $p<0.05$, ††: $p<0.01$, †††: $p<0.001$, in comparison to EPI/SDOC- group for rat and DBA/2 with FRA group for mice. *, $p<0.05$; **, $p<0.01$, in comparison to control rats.

A- Basal respiratory variables under normoxia at 12 weeks post-*SE*

	Control (n=5)	EPI/SDOC- (n=8)	EPI/SDOC+ (n=8)	Ctrl vs SDOC- (P value)	Ctrl vs SDOC+ (P value)	SDOC+ vs SDOC- (P value)
$\dot{V}_E$ (mL×min^{-1}×100g^{-1})	32 ± 1	30 ± 1	26 ± 1	*0.274*	0.002	0.013
f_R (breath×min^{-1})	101 ± 3	93 ± 2	87 ± 2	0.015	<0.001	0.036
V_T (mL×100g^{-1})	0.32 ± 0.01	0.33 ± 0.01	0.31 ± 0.01	*NT*	*NT*	*NT*
T_I (msec)	207 ± 7	233 ± 3	263 ± 8	0.015	<0.001	<0.001
T_E (msec)	385 ± 12	413 ± 9	438 ± 10	*0.088*	0.003	*0.083*
T_{tot} (msec)	593 ± 14	646 ± 11	700 ± 17	0.025	<0.001	0.013
Apnea frequency x min^{-1}	0.32 ± 0.17	0.27 ± 0.13	0.10 ± 0.07	*NT*	*NT*	*NT*
Apnea duration (msec)	1339 ± 67	1435 ± 65	1477 ± 75	*NT*	*NT*	*NT*
Coefficient of variation	0.49 ± 0.01	0.53 ± 0.01	0.52 ± 0.02	*NT*	*NT*	*NT*

B- Respiratory response to hypoxia and hypercapnia at 13-14 weeks post-*SE*

% of baseline		Control (n=13)	EPI/SDOC- (n=12)	EPI/SDOC+ (n=6)	Ctrl vs SDOC- (P value)	Ctrl vs SDOC+ (P value)	SDOC+ *vs* SDOC- (P value)
Hypoxia 9% of O$_2$	$\dot{V}_E$ (mL×min^{-1}×100g^{-1})	225 ± 9	273 ± 18	278 ± 19	0.019	0.046	*0.847*
	f_R (breath×min^{-1})	146 ± 5	171 ± 4	165 ± 3	<0.001	0.024	*0.463*
	V_T (mL×100g^{-1})	153 ± 5	169 ± 8	170 ± 7	*NT*	*NT*	*NT*
Hypercapnia 5% of CO$_2$	$\dot{V}_E$ (mL×min^{-1}×100g^{-1})	187 ± 6	211 ± 8	219 ± 14	0.028	0.023	*0.562*
	f_R (breath×min^{-1})	128 ± 4	141 ± 4	141 ± 8	*NT*	*NT*	*NT*
	V_T (mL×100g^{-1})	141 ± 3	148 ± 5	153 ± 6	*NT*	*NT*	*NT*

C- Cardiac frequency at 15 weeks post-*SE*

		Control (n=3)	EPI/SDOC- (n=3)	EPI/SDOC+ (n=2)	Ctrl vs SDOC- (P value)	Ctrl vs SDOC+ (P value)	SDOC+ vs SDOC- (P value)
Day	F_C (beats×min^{-1})	315 ± 17	338 ± 11	333 ± 8	*NT*	*NT*	*NT*
Night	F_C (beats×min^{-1})	332 ± 19	355 ± 9	351 ± 12	*NT*	*NT*	*NT*

D- Body-temperature at 15 weeks post-*SE*

		Control (n=3)	EPI/SDOC- (n=3)	EPI/SDOC+ (n=2)	Ctrl vs SDOC- (P value)	Ctrl vs SDOC+ (P value)	SDOC+ vs SDOC- (P value)
Day	T_B (degree C°)	37.5 ± 0.1	37.6 ± 0.1	37.7 ± 0.2	*NT*	*NT*	*NT*
Night	T_B (degree C°)	38 ± 0.2	37.9 ± 0.1	38.1 ± 0.1	*NT*	*NT*	*NT*

Table 1: Monitoring of cardio-respiratory functions in control and EPI rats between 12 and 15 weeks post-SE. Using plethysmography, the ventilatory variables were measured under normoxic condition (21% O$_2$) at 12 weeks post-SE (A) and in response to hypoxia (9% O$_2$) or hypercapnia (5% CO$_2$) exposures between 13-14 weeks post-SE (B). Cardiac frequency (C) and body-temperature (D) were concomitantly evaluated at 15 weeks post-SE. Abbreviations: Fc, cardiac frequency; f_R, respiratory frequency; T_B, body-temperature; T_E, expiration time; T_I, inspiration time; T_{tot}, total respiratory cycle duration; $\dot{V}_E$, minute volume; V_T, tidal volume. Statistics: NT, not statistically significant.