Asialoglycoprotein receptor 1 is a novel PCSK9-independent ligand of liver LDLR that is shed by Furin

Delia Susan-Resiga[1,†], Emmanuelle Girard[1,†], Rachid Essalmani[1], Anna Roubtsova[1], Jadwiga Marcinkiewicz[1], Mouna Rabeb Derbali[1], Alexandra Evagelidis[1], Paul F. Lebeau[2], Richard C. Austin[2], Nabil G. Seidah[1,]*

[1]Montreal Clinical Research Institute (IRCM), affiliated to the University of Montreal, Laboratory of Biochemical Neuroendocrinology, Montreal, Quebec H2W 1R7, Canada;
[2]Department of Medicine, Division of Nephrology, McMaster University, St. Joseph's Healthcare Hamilton, Ontario L8N 4A6, Canada;

[†] Authors contributed equally to this work.

* **To whom correspondence should be addressed:** Dr. Nabil G. Seidah, Laboratory of Biochemical Neuroendocrinology, Clinical Research of Montreal, 110 Pine Ave. West, Montréal, QC H2W 1R7, Canada. Tel: 514- 987-5609; E-mail address: seidahn@ircm.qc.ca (N.G. Seidah).

Running title: ASGR1 regulates hepatic LDLR expression

Key words: ASGR1, Cardiovascular disease, LDL cholesterol, Lipid metabolism, Liver

This manuscript contains a supporting information as a supplementary Table S1.

ABSTRACT
The hepatic carbohydrate-recognizing
asialoglycoprotein receptor (ASGR1) mediates
the endocytosis/lysosomal degradation of
desialylated glycoproteins following binding
to terminal galactose/N-acetylgalactosamine.
Human heterozygote-carriers of *ASGR1*-
deletions exhibited ~34% lower risk of
coronary artery disease and ~10-14% non-
HDL-cholesterol reduction. Since PCSK9 is a
major degrader of LDLR, we examined the
regulation of LDLR and/or PCSK9 by ASGR1.
We investigated the role of
endogenous/overexpressed ASGR1 on LDLR
degradation and functionality in naïve HepG2
and HepG2-PCSK9-knockout cells by
Western-blot and immunofluorescence.
ASGR1, like PCSK9, targets LDLR and both
interact with/enhance the degradation of the
receptor independently. The lack of
cooperativity between PCSK9 and ASGR1 on
LDLR expression was confirmed in livers of
wild-type (WT) *versus* *Pcsk9*$^{-/-}$ mice. ASGR1-
knockdown in naïve HepG2 cells significantly
increased total (~1.2-fold) and cell-surface
(~4-fold) LDLR protein. In HepG2-PCSK9-
knockout cells ASGR1-silencing led to ~2-fold
higher levels of LDLR protein and DiI-LDL
uptake associated with ~4-fold increased cell-
surface LDLR. Overexpression of WT-ASGR1
reduced primarily the immature non-O-
glycosylated LDLR (~110 kDa), whereas the
triple Gln^{240}/Trp^{244}/Glu^{253} Ala-mutant (loss of
carbohydrate-binding) reduced the mature
form of the LDLR (~150 kDa), suggesting that
ASGR1 binds the LDLR in sugar-dependent
and -independent fashion. Furin sheds ASGR1
at **R**KM**K**^{103}↓ into a secreted form, likely
resulting in a loss-of-function on LDLR.
LDLR is the first example of a liver-receptor
ligand of ASGR1. Additionally, we
demonstrate that lack of ASGR1 enhances
LDLR levels and DiI-LDL incorporation,
independently of PCSK9. Overall, silencing of
ASGR1 and PCSK9 may lead to higher LDL-
uptake by hepatocytes, thereby providing a
novel approach to further reduce LDL-
cholesterol.

INTRODUCTION
Elevated levels of plasma low-density
lipoprotein cholesterol (LDLc) and
inflammation are the leading modifiable risk
factors that drive the development and
progression of atherosclerosis, the underlying
cause of cardiovascular disease (CVD) (1). The
removal of plasma LDLc is mediated by the
LDL receptor (LDLR) located on the surface of
hepatocytes. The proprotein convertase
subtilisin/kexin type 9 (PCSK9) is highly
expressed in the liver (2) and upon its secretion
into plasma, binds to and enhances the
degradation of the LDLR in a non-enzymatic
fashion (3-5), thereby increasing circulating
LDLc. Unlike gain-of-function (GOF) PCSK9
variants (6), loss-of-function (LOF) variants
enhance LDLR levels and promote LDLc
clearance (4,7). The development of inhibitory
human monoclonal antibodies against PCSK9
represents a unique treatment strategy for the
management of CVD, which has been shown
to reduce the risk of cardiovascular events in
clinical populations (8). Despite these hallmark
studies and their utility for lipid lowering to
prevent CVD, other surface proteins on
hepatocytes that bind to and modulate their
interaction with PCSK9 and/or LDLR as well
as LDLc uptake have not been thoroughly
characterized (5).
The dimeric asialoglycoprotein receptor
(ASGR) (9) is a hepatocyte type-II
transmembrane heterodimeric glycoprotein
highly conserved among mammals. It consists
of two subunits ASGR1 (major; Fig. 1A) and
ASGR2 (minor) and plays a critical role in
serum glycoprotein homeostasis by mediating
the endocytosis and lysosomal degradation of
desialylated glycoproteins with exposed
terminal galactose or N-acetylgalactosamine
residues (10). In mouse liver ASGRs are highly
expressed during development and in the adult
(Fig. 1B). Recent studies showed that
heterozygous carriers of the early termination
LOF del12 mutation in the ASGR1 gene (1 in
120 persons in the Icelandic study) and another
early termination heterozygote LOF ASGR1
mutant (W158X) had lower plasma levels of
non-HDL cholesterol than non-carriers, and
that the del12 mutation was associated with a

significant 34% lower risk of CVD (11).
Comparison of plasma LDLc decrease in LOF
PCSK9 R46L (-17 mg/dL; ~30% reduction in
CVD risk) with that in LOF ASGR1 del12
mutation (-11 mg/dL) illustrates that reduction
in CVD risk observed in this ASGR1 mutant
(~34%) was far greater than would have been
predicted (i.e., ~18%) from the associated
modest reduction of LDLc (11). This suggests
that the atheroprotective effects of ASGR1
del12 go beyond the lowering of serum LDLc.
Strategies aimed at inhibiting ASGR1
expression/activity are being explored as
another approach to lowering plasma LDLc
and CVD risk (10). Despite the importance of
PCSK9 and ASGR1 in lipid lowering and
reduced CVD risk, there are currently no
studies to suggest that ASGR1 and PSCK9 act
in concert to impact lipid metabolism and/or
atherosclerosis by modulating the
binding/uptake of LDLc *via* the LDLR. The
purpose of the present study was to investigate
the possible regulation of the LDLR by
ASGR1 and the role of PCSK9 in this process.
124
RESULTS
Expression of ASGR1 and ASGR2 in mice
during development and in the adult, and in
human hepatocyte cell lines
Human ASGR is composed of a major subunit
(ASGR1) and a minor subunit (ASGR2) (12).
Both subunits are type II, single-pass
transmembrane proteins. In ASGR1, residues
Gln_{240}, Trp_{244} and Glu_{253} in the carbohydrate
recognition domain (CRD) are critical for
binding exposed terminal galactose or N-
acetylgalactosamine residues
(https://www.uniprot.org/uniprot/P07306)
(Fig. 1A). In mouse, whole body *in situ*
hybridization histochemistry of ASGR1 and
ASGR2 mRNA expression during
development and in the adult revealed that both
transcripts are mainly expressed in liver
starting at embryonic days 17 and 15,
respectively, and showed that their expression
increases until adulthood (Fig. 1B). We next
estimated the relative mRNA expression of
ASGR1 and ASGR2 by quantitative RT-PCR
in adult mouse liver and in human hepatocyte
HepG2, HepG2-PCSK9-KO cells (lacking

endogenous PCSK9) (13), and the
immortalized human primary hepatocytes IHH
cell line (Fig. 1C) (13). In all cases the mRNA
levels of ASGR1 were higher than those of
ASGR2, especially in mouse liver.
155
PCSK9 reduces LDLR but not ASGR1 levels
Since circulating PCSK9 is responsible for the
majority of its activity to reduce the levels of
cell-surface LDLR in hepatocytes (5), in all
our experiments cells were incubated with
purified recombinant PCSK9 (14). We thus
compared the known ability of *exogenous*
PCSK9 to enhance the degradation of LDLR to
its effect on ASGR1 in HepG2 cells by
Western blot (WB; Fig. 2A), and
immunofluorescence (IF; Fig. 2B). As
expected, PCSK9 reduced the endogenous
levels of total LDLR (-40% by WB) and cell-
surface LDLR (-90% by IF). In contrast,
PCSK9 did not alter the levels of endogenous
ASGR1 in both assays.
The effect of the lack of *endogenous*
PCSK9 on LDLR and ASGR1 in HepG2 cells
was next investigated following its mRNA
knockdown (KD) using a smart pool of 4
siRNAs. This resulted in ~70% reduction in
PCSK9 mRNA and protein levels (Figs. 2C,
D). As expected (15,16), this treatment did not
affect the mRNA levels of LDLR. In addition,
no effect was observed on the mRNA levels of
ASGR1, ASGR2 and SREBP2 (Fig. 2C). At
the protein level, PCSK9 KD resulted in ~1.5-
fold increase in LDLR, with no change in
ASGR1 (Fig. 2D). We conclude that in HepG2
cells *endogenous* or *exogenous* PCSK9 does
not affect ASGR1 mRNA and/or protein
levels.
To address the effect of PCSK9 on ASGR1
in vivo, we analyzed the levels of ASGR1 in
Pcsk9-knockout (KO) mice (15), by liver
immunohistochemistry (IHC) under non-
permeable conditions (17) (Fig. 3A) and WB
(Fig. 3B). The data show that while LDLR
protein levels at the cell-surface of hepatocytes
(IHC) and in whole tissue (WB) are increased
in *Pcsk9*-KO livers, in accordance with (17),
those of ASGR1 are not affected. Overall, we
conclude that, like in HepG2 cells, *in vivo*
endogenous PCSK9 does not affect the levels
of ASGR1 in mouse hepatocytes.

To analyze the impact of the lack of LDLR on ASGR1 protein, we now show that in the liver of *Ldlr*-KO mice both ASGR1 cell-surface immunoreactivity (Fig. 3A) and total ASGR1 protein by WB (Fig. 3B) are significantly increased by ~1.6 and ~1.3 fold, respectively. This suggests that ASGR1 levels could also be regulated by those of the LDLR, as are circulating PCSK9 levels (5).

ASGR1 regulates LDLR levels and functionality independently of PCSK9

In order to examine the impact of ASGR1 on the LDLR, KD of ASGR1 (-75%) was performed in naïve HepG2 and HepG2-PCSK9-KO cells (Figs. 4, 5, 6). We first demonstrated that KD of ASGR1 in naïve HepG2 cells does not affect the levels of intracellular and secreted PCSK9 (Fig. 4). In contrast, ASGR1 KD resulted in a significant increase in total (WB) and cell-surface (IF) LDLR in both cell lines. Specifically, we estimated a total LDLR increase of ~1.3-fold (Fig. 5A) and ~2-fold (Fig. 5B) in each cell line, respectively. The corresponding cell-surface increase in LDLR was ~4-fold in both cell lines (Figs. 5C, D top left panels). Transcript analysis by quantitative RT-PCR revealed that KD of ASGR1 did not modify the mRNA levels of LDLR and SREBP2 (Fig. 5E). This suggests that endogenous ASGR1 reduces the total levels of LDLR protein, especially those at the cell-surface, leading to the hypothesis that LDLR is a target of ASGR1, independent of PCSK9.

We next investigated whether ASGR1 affects the ability of PCSK9 to enhance the degradation of the LDLR. In both cell lines KD of ASGR1 had no significant impact on the activity of exogenous PCSK9 on the LDLR (Figs. 5A, B, C). Thus, we can conclude that independently PCSK9 and ASGR1 can reduce the levels of LDLR protein.

To probe the functionality of the LDLR upon KD of ASGR1, we incubated naïve HepG2 (Fig. 6) and HepG2-PCSK9-KO cells (Fig. 5D) with fluorescent DiI-LDL and estimated its uptake by IF. The data show that the KD of ASGR1 resulted in at least ~2-fold increase in DiI-LDL uptake mirroring the increase in LDLR protein levels (Figs. 5D, 6).

Thus, endogenous ASGR1 regulates protein levels of LDLR and its functionality independently of PCSK9.

Since ASGR1 can act as a homodimer or heterodimer with ASGR2, we next examined the effect of overexpression of ASGR1/2 in HepG2-PCSK9-KO cells. Co-expression of ASGR1 and ASGR2 resulted in a significant decrease in the total levels of both endogenous (-20%) and overexpressed (-30%) LDLR (WB; Fig. 7A), of the cell-surface endogenous LDLR (-80%) and DiI-LDL uptake (-70%) (IF; Fig. 7B). Altogether, KD and overexpression analyses demonstrated that, independent of PCSK9, ASGR1 negatively regulates LDLR levels and its functionality.

ASGR1 can bind LDLR and enhance its degradation by carbohydrate-dependent and independent mechanisms

Co-immunoprecipitation analyses were used to probe the possible interaction between endogenous ASGR1 and LDLR in HepG2-PCSK9-KO cells (Fig. 8A). Thus, immunoprecipitation of cell lysates under non-denaturing conditions with an ASGR1 antibody, followed by separation of the immune complex by SDS-PAGE and WB with an LDLR antibody revealed that human ASGR1 (~48 kDa) co-immunoprecipitated primarily with mature mLDLR (N- and O-glycosylated; ~150 kDa) and not immature iLDLR (non-O-glycosylated; ~110 kDa) (18). This suggests that endogenous ASGR1 and LDLR form a complex independent of PCSK9. This observation was confirmed also in *vivo*, in mouse liver from wild type (WT) and *Pcsk9*-KO mice (Fig. 8B).

Since ASGR1 is a lectin-binding protein and Gln_{240}, Trp_{244} and Glu_{253} in its CRD are critical for carbohydrate binding, we next probed by immunoprecipitation the WT ASGR1 and its triple Ala-mutant (QA/WA/EA), following their co-expression with LDLR in HEK293 cells, which do not endogenously express ASGR1/2. Immunoprecipitation with Flag antibody (ASGR1-Flag) and WB with an ASGR1 antibody and LDLR antibody, respectively, revealed that WT ASGR1 binds to both immature iLDLR (~110 kDa) and mature

mLDLR (~150 kDa) (Fig. 8C, left panel). In contrast the triple Ala-mutant, which lost its carbohydrate-binding capacity, primarily binds the non-O-glycosylated iLDLR (Fig. 8C, left panel). Binding of the LDLR to ASGR1 was further confirmed in a reciprocal binding assay where the LDLR-V5 is immunoprecipitated first with V5 antibody and the immune complex is then separated by SDS-PAGE and revealed by WB to ASGR1 (Fig. 8C, right panel).

We next investigated whether the enhanced degradation of functional LDLR requires the intact carbohydrate-binding motif of ASGR1. Thus, the levels of cell-surface LDLR and DiI-LDL uptake were estimated in HEK293 cells expressing either LDLR alone or LDLR and ASGR2 together with either WT ASGR1 or its triple Ala-mutant (QA/WA/EA). In non-permeabilized cells, IF showed that both cell-surface LDLR and DiI-LDL uptake are significantly reduced in the presence of WT ASGR1, but not of its QA/WA/EA triple Ala-mutant (Fig. 9A). Thus, functional cell-surface LDLR levels are reduced only by WT ASGR1 and require its critical carbohydrate-binding residues. We extended these data by biotinylating cell-surface proteins of the above HEK293 cells and analyzed the biotinylated proteins by WB. The data showed that the ratio of cell-surface [iLDLR] / [total LDLR] is significantly reduced (~40%) by WT ASGR1 but not by the triple mutant (Fig. 9B, lower left panel). Furthermore, we noticed that the cell-surface levels of the triple-Ala mutant of ASGR1 are ~60% (~1.6 fold) higher than the WT (Fig. 9B, top lelft panel), mirroring its ~70% (~1.7 fold) increased total cellular levels (Fig. 9B, right panel) and suggesting that even at higher levels, this mutant that binds iLDLR (Fig. 8C; left panel) does not efficiently reduce its levels (Fig. 9A). Collectively, the data suggest that the integrity of ASGR1 CRD regulates the receptor's ability to reduce the levels of cell-surface iLDLR, and that in its absence iLDLR is less degraded and more apparent at the cell-surface.

In contrast, WB analyses of the total levels of LDLR in these cells showed that as for WT, the triple QA/WA/EA and double QA/WA mutants of ASGR1 can also significantly reduce total LDLR levels (~ 40%) relative to vector control lacking ASGR1/2 (Fig 9C, lower right panel). As observed at the plasma membrane (Fig. 9B, left panel), in total cell lysate (intracellular and plasma membrane) WT ASGR1 also reduced the levels of iLDLR, which is not seen with the CRD mutants (Fig 9C; lower left panel). This is evident from the relative ratio of [iLDLR] / [total LDLR], which is ~40% lower in WT ASGR1 *versus* double or triple mutants and vector control (Fig. 9C; lower left panel). Since some of the LDLR is shed into the medium by undefined matrix metalloproteases (19), we also analyzed the fate of media LDLR. The data also showed that ASGR1 reduces the levels of the shed forms of LDLR by carbohydrate-dependent (WT) and carbohydrate-independent (Ala-mutants) activities (Fig. 9C; top right panel). Conversely, silencing ASGR1 in HepG2-PCSK9-KO cells significantly increased total cellular LDLR levels, associated with enhanced shedding of the LDLR (~10% of total LDLR), especially iLDLR (Fig. 9D).

ASGR1 specificity for LDLR
The present data are the first to identify the secretory membrane-bound LDLR as a physiological target of ASGR1, as opposed to soluble desialylated glycoproteins, e.g., von Willebrand factor (20), normally thought to be sensitive to hepatocyte-derived ASGR1 regulation (10). However, since we do not know the repertoire of liver glycoproteins that are targeted by ASGR1, we next investigated the PCSK9-independent ability of ASGR1 to enhance the degradation of other membrane-bound liver receptors. Accordingly, we analyzed by WB the endogenous levels of CD36 and insulin receptor following KD of ASGR1 in HepG2-PCSK9-KO cells. The data show that although KD of ASGR1 increased the levels of LDLR by ~2-fold, those of CD36 and insulin receptor were not affected (Fig. 10). Thus, even though like the LDLR, the highly N-glycosylated CD36 is a target of PCSK9 (21), it is not sensitive to ASGR1 in the absence of PCSK9. We conclude that the ability of ASGR1 to enhance the degradation of the LDLR may not extend to all liver-derived secretory membrane-bound receptors.

Furin sheds ASGR1

We previously showed that Furin, the third member of the proprotein convertase (PC) family (22), can cleave and inactivate PCSK9 in liver hepatocytes at the general PC-like motif (**R/K**)xx(**R/K**)↓ (23,24). Scanning the primary sequence of the luminal domain of human ASGR1 revealed the presence of two potential Furin-like cleavage sites, namely **R**GL**R**$_{74}$↓ET and **R**KM**K**$_{103}$↓SL. Thus, co-expression in HEK293 cells of human ASGR1 with each of the liver-expressed PCs (22) revealed that only Furin can cleave the protein into a soluble ~28 kDa secreted form (Fig. 11A). Whereas Ala-mutation of the **R**GL**R**$_{74}$↓ET site had no effect (Fig. 11B), only Ala-mutation of the underlined basic aa in the **R**KM**K**$_{103}$↓SL site abrogated Furin-cleavage and hence sheding of ASGR1 (Fig. 11C). Indeed, insertion of an optimal Furin-like site **RRRR**$_{103}$↓**E**L resulted in complete cleavage of mature ~48 kDa ASGR1 by endogenous Furin into a soluble ~28 kDa form (Fig. 11D). A similar observation was made for PCSK9 optimally cleaved by Furin (23). A comparable processing pattern was observed in hepatic HepG2-PCSK9-KO cells (Fig. 11E). In conclusion, Furin sheds ASGR1, likely resulting in a loss-of-function towards LDLR.

DISCUSSION

The hepatic ASGR is pivotal for the efficient clearance of desialylated glycoproteins from circulation by receptor-mediated endocytosis and lysosomal degradation, whereupon the receptor is recycled back to the cell-surface (12,25). Glycoprotein ASGR ligands contain terminal galactose or N-acetylgalactosamine, with the latter being the preferred residues (25). In humans, the functional receptor is composed of two subunits, ASGR1 (major, 46 kDa) and ASGR2 (minor, 50 kDa) that exist as homo-oligomers or hetero-oligomers in a molar ratio of ~3:1, respectively (12,25). Both subunits are type-II transmembrane proteins comprising a cytosolic, transmembrane domain, a luminal stalk region and a calcium-dependent CRD (Fig. 1A). Oligomerization of the ASGR subunits *via* their respective stalk regions occurs in the endoplasmic reticulum (ER) (25). ASGRs are internalized from the cell-surface *via* clathrin-coated vesicles into endosomes and are rapidly recycled constitutively even in the absence of a ligand (10). Thus, at steady-state only ~40-60% of total cellular ASGRs are on the cell-surface (25). So far, the known ligands of ASGRs are soluble desialylated proteins, including von Willebrand factor (10) and exogenous proteins (e.g. asialofuetin, asialo-orosomucoid) (26).

Recently, it was observed that conjugation of antisense oligonucleotides (ASO) to N-acetylgalactosamine mediated their efficient uptake into liver hepatocytes *via* ASGR1, providing a powerful targeted delivery of these drugs to the liver (27). Indeed, this ASGR1-targeted antisense technology is now successfully applied to silence liver PCSK9 in hypercholesterolemic patients worldwide (28).

Circulating PCSK9 originating from hepatocytes (2) binds to cell-surface LDLR and targets it to degradation in endosomes/lysosomes (5). This mechanism led to a powerful new treatment to reduce LDLc *via* inhibitory monoclonal antibodies or antisense mRNA silencing (29). In 2016, Nioi *et al.* reported that two human *ASGR1*-deletion variants found in Iceland, resulting in truncated, likely inactive, proteins, are associated with a modest ~10-14% reduction in LDLc, yet with a significant ~34% reduced risk of coronary artery disease (11). This suggested that the atheroprotective effects of *ASGR1* LOF extended beyond reduction of serum LDLc and could implicate, as of yet unknown functions of ASGR1.

Since PCSK9 is a major player in the regulation of LDLc (6) and its receptor LDLR (3,5), we investigated the possible involvement of ASGR1 in the regulation of LDLR (10) and/or PCSK9 functions. The data presented in this work clearly eliminated a reciprocal direct effect between PCSK9 and ASGR1 proteins (Figs. 2, 3, 4, 5), and demonstrated that loss of endogenous ASGR1 in hepatic cells lines enhanced the levels of the LDLR and its ability to uptake DiI-LDL (Fig. 5), whereas the reverse is true upon overexpression of ASGR1 (Fig. 7), and both phenotypes are independent of PCSK9. This was confirmed *in vivo* in mice

lacking PCSK9, which as expected exhibit higher levels of LDLR (15), but no significant change in ASGR1 protein levels (Fig. 3). Finally, lack of endogenous ASGR1 in HepG2 cells does not affect PCSK9 mRNA or its protein levels in cells and media (Fig. 4).

In the absence of PCSK9, co-immunoprecipitation experiments in HepG2-PCSK9-KO cells and in mouse *Pcsk9*-KO livers revealed that endogenous ASGR1 and LDLR form a complex (Figs. 8A, B). Since ASGR1 binds N-acetyl-galactosamine/galactose, and its Gln_{240}, Trp_{244} and Glu_{253} are implicated in carbohydrate recognition (26), we tested the importance of these ASGR1 residues in the regulation of the LDLR. The triple Ala-mutant of these residues largely failed to co-immunoprecipitate with mLDLR (~150 kDa) in HEK293 cells, but still bound non-O-glycosylated iLDLR (~110 kDa) (Fig. 8C; left panel). This unexpected result revealed that ASGR1 could bind mostly the mLDLR (N- and O-glycosylated) in a sugar-dependent fashion, and iLDLR (N-glycosylated) in an O-glycosylation-independent manner. These data suggest that the galactose/N-acetyl-galactosamine residues recognized by the CRD of ASGR1 likely reside in part on O-glycosylated chains of the mLDLR. The latter are predominantly present close to the C-terminal transmembrane domain, but are also found in the linker regions separating the N-terminal repeat domains of the LDLR (18). Finally, shedding of the LDLR by metalloproteases is thought to mostly target iLDLR (19). Overexpression of WT ASGR1 decreased LDLR shedding, likely due to degradation of mainly iLDLR (Fig. 9C), and silencing endogenous ASGR1 resulted in elevated shedding of primarily iLDLR, in line with increased cellular LDLR and higher exposure to endogenous sheddase(s) (Fig. 9D). It was of interest to note that not all cell surface glycoproteins are targeted by ASGR1, since the insulin receptor and CD36 were insensitive to it (Fig. 10). It is possible that the O-glycosylation-independent binding may be implicated in the selectivity of ASGR1 towards the LDLR.

The 9-membered family of proprotein convertases (PCs) is composed of secretory serine proteases implicated in a wide variety of functions both in health and disease (22,30). The first seven members of the family cleave substrates exhibiting a canonical R/K-(2X)n-R/K↓ sequence, where X is any residue and n=0-3 spacer aa (22). The presence of two such motifs in the primary sequence of ASGR1 suggested that it may be cleaved by one or more member(s) of the PC-family. Indeed, our data showed that only Furin can cleave ASGR1 at **R**KM**K**$_{103}$↓SL and shed it into the media (Figs. 11A, C, E). We therefore expect that some of the hepatic ASGR1 may be circulating as a soluble form in plasma and conceivably bind to non-hepatic targets and regulate their function. The generation of an optimally shed form of ASGR1 with the above sequence replaced by **RRRR**$_{103}$↓**E**L (Fig. 11D) may lead in the future to the identification of some of the possible extrahepatic roles of the resulting circulating ASGR1. One additional suggested consequence of the lack of ASGR1 in human was on receptors implicated in modulating inflammation (11).

Finally, it is interesting to note that liver ASGR1 has recently been shown to bind the N-terminal domain and ACE2 receptor binding domain of the Spike-glycoprotein of SARS-CoV-2 (31), the etiological agent of COVID-19 (32). Since liver does not significantly express ACE2, the main SARS-CoV-2 receptor (33), it is possible that ASGR1 can act as an alternative receptor in hepatocytes, thereby expanding the tropism of this deadly virus (34). It would be thus be informative to define the mode of interaction of ASGR1 with the Spike-glycoprotein of SARS-CoV-2 and whether the LDLR may participate in this process.

We conclude that the LDLR is a ligand of hepatic ASGR1, representing the first case of a membrane-bound protein targeted by ASGR1 for degradation. This is the first instance in which surface ASGR1 interacts with another surface protein to induce its degradation, since ASGRs are only known to interact with circulating factors (10,25). Our data provide a mechanism for the modest reduction of LDLc in humans lacking functional ASGR1 (11). The other functions of ASGR1 that are not related to LDLc regulation but affect inflammation,

608 atherosclerosis and cardiovascular health (11)
609 are yet to be identified.
610
611

EXPERIMENTAL PROCEDURES
Plasmids and site directed mutagenesis
The cDNAs encoding for human ASGR1 and
human ASGR2 were obtained from GenScript
(NJ). Human ASGR1 and its mutants
(Q240A/W244A and Q240A/W244A/E253A)
and human ASGR2 were sub-cloned into
pcDNA 3.1 vector (Invitrogen) containing a C-
terminal Flag tag (ASGR1) or HA tag
(ASGR2). Point mutations of ASGR1 were
created by site-directed mutagenesis, and the
identity of each mutant was confirmed by DNA
sequencing. The constructions containing the
human LDLR, human PACE4, human PC7,
human Furin or mouse PC5A, cloned in
pIRES2-EGFP (Clontech), were previously
described (35,36).
629

Cell culture, transfections and treatments
HepG2 (human hepatocellular carcinoma),
HepG2-PCSK9-KO (CRISPR cas9 for
PCSK9), IHH (immortalized human primary
hepatocytes) and HEK293 (human embryonic
kidney-derived epithelial) cells were cultured
at 37°C under 5% CO_2 in complete medium
(Eagle's Minimal Essential Medium for
HepG2 and IHH cells; Dulbecco's Modified
Eagle's Medium for HEK293 cells) (Gibco,
Thermofisher Scientific) containing 10% (v/v)
fetal bovine serum (Invitrogen).
Protein overexpression was achieved using
JetPEI (PolyPlus) transfection reagent in
HepG2 and HepG2-PCSK9-KO cells and
JetPRIME (PolyPlus) in HEK293 cells,
following the manufacturer's
recommendations. 24 hours post-transfection,
the culturing medium was changed from
complete to serum-free to achieve maximal
expression of the LDLR, and the cells were
treated according to each experiment.
Incubations with *exogenous* purified PCSK9
(ACRO Biosytems) was carried out for an
additional 20 hours. LDLR functionality was
assessed by incubation with Dil-LDL (Alfa
Aesar, Kalen Biochemicals). Small interfering
RNA (siRNA) targeted against ASGR1 or
PCSK9 were purchased from siGenome,

Horizon Discoveries and transfections were
carried out using INTERFERin (PolyPlus) as
per manufacturer's instructions.
662

Western blotting
Cells were washed twice with ice-cold PBS
and lysed 60 min on ice with ice-cold non-
denaturing cell lysis buffer (20 mM Tris-HCl,
pH 8, 137 mM NaCl, 2 mM Na_2EDTA, 1%
Nonidet P-40, 10% glycerol, supplemented
with protease inhibitor cocktail without
EDTA). Mouse livers were homogenized and
protein extraction was performed in
radioimmune precipitation assay buffer (50
mM Tris-HCl pH 7.8, 150 mM NaCl, 1%
Nonidet P-40, 0.5% sodium deoxycholate,
0.1% SDS, supplemented with complete
protease inhibitor cocktail) for 40 min on ice.
Cell lysates and conditioned media were
electrophoretically resolved on 8 or 10% Tris-
glycine SDS-polyacrylamide gels,
respectively, and transferred to PVDF
membranes using a Trans-Blot Turbo transfer
System (Bio-Rad). Mouse liver lysates were
subjected to 8% Tris-tricine SDS-PAGE before
transfer to PVDF membrane. Membranes were
blocked with 10% skim milk in tris-buffered
saline containing Tween-20 for 1 hour and
subsequently incubated with primary antibody
according to the manufacturer's
recommendations. Antigen-primary antibodies
complexes were detected with secondary
antibodies conjugated to horse radish
peroxidase and developed using a
chemiluminescent reagent (Clarity ECL, Bio-
Rad). Images of the western blots were
acquired using a ChemiDoc MP System (Bio-
Rad) and analyzed with Image Lab (version
6.0) software (Bio-Rad). Immunoblots from
cell lysates and liver lysates were quantified
and normalized to membranes probed for α-
Tubulin or β-Actin, respectively. See
Supplementary Table S1 for a list of the
antibodies used for Western blot analysis.
703

Immunofluorescence
For immunofluorescence experiments,
HEK293, HepG2 or HepG2-PCSK9-KO cells
($0.5x10^5$ cells/well) were plated on poly-L-
lysine-coated round microscope coverslips that
were placed in a 24-well cell culture plate.

Cells were then treated as required for each
experiment (PCSK9 incubation, siRNA
transfection, protein overexpression). To
analyze plasma membrane LDLR and ASGR1
expression, the cells were washed twice with
PBS and fixed with a solution of 4%
paraformaldehyde in PBS (10 min). After
blocking with PBS + 2% BSA (1 hour),
samples were incubated at 4 °C overnight with
the respective primary antibodies (see Table
S1), washed with PBS, and incubated with the
appropriate fluorescent secondary antibody
(see Table S1) for 1 hour at room temperature.
To analyze LDLR and ASGR1 total
expression, the same protocol was followed,
except after the blocking step and prior to
incubation with primary antibody, cells were
permeabilized with PBS containing 0.2%
Triton X-100 for 5 min. When LDLR
functionality was tested, prior to fixation cells
were incubated for 2 hours at 37°C with 5µg/ml
DiI-LDL in SFM media. Coverslips were
mounted on a glass slide with ProLong Gold
antifade reagent with DAPI. Samples were
visualized using a Plan-Apochromat 63x 1.4
oil objective of an LSM-710 confocal laser-
scanning microscope (Carl Zeiss) with
sequential excitation and capture image
acquisition with a digital camera. Images were
processed with ZEN software. Image analysis
to quantify the fluorescence intensities was
accomplished using Volocity®6.0.
742
Animal experiments
In situ hybridization studies of ASGR1 and
ASGR2 mRNA expression were performed on
whole body tissue cryostat sections (8- to 10-
µm) from WT (C57BL/6J) mice, embryonic
day 10 to adult, as described previously
(2,13,37). Mouse antisense and sense (negative
control) cRNA (complementary RNA)
riboprobes coding for ASGR1 and ASGR2
were labeled with ^{35}S-UTP and ^{35}S-CTP (1,250
Ci/mmol; Amersham Pharmacia), to obtain
high specific activities of ≈1,000 Ci/mmol (2).
755
Immunohistochemistry and Western Blot
LDLR and ASGR1 protein expression in
mouse liver was assessed by IHC on
cryosections (9-10 mice) and WB of tissue
extracts (3-4 mice) from 12-16-week-old male

Pcsk9$^{-/-}$ and *Ldlr$^{-/-}$* mice on a C57BL/6J
background and age-matched C57BL/6J
controls. For IHC, mice liver cryosections (8
µm thick) were fixed in 4% paraformaldehyde
in PBS for 1 hour, rinsed in PBS with 0.1%
glycine, washed in PBS, and blocked in 1%
BSA (Sigma-Aldrich, Oakville, ON) in PBS
for 1 hour at room temperature. Sections were
then incubated overnight at 4°C with the LDLR
and ASGR1 primary antibodies (see Table S1)
and washed three times for a total of 15 min in
PBS. Labeling was visualized by incubation
with Alexa Fluor 488-labeled secondary
antibodies (see Table S1) for 1 hour at room
temperature in PBS. After three washes with
PBS, nuclei were counterstained with Hoechst
dye (Sigma). Images were acquired as
described previously using a LSM 700
confocal microscope equipped with ZEN 2011
software (38). Data quantification was
achieved using Matlab software. For all images
the minimum threshold was set to a value of 20,
negative control values were subtracted and
final values were analyzed using Microsoft
Excel. For each genotype, the most
representative image was chosen.
787
Quantitative RT-PCR
Quantitative RT-PCR was performed as
previously described (15). Briefly, total RNA
from liver or cells was extracted with TRIzol
(Invitrogen). cDNA was generated from 1µg of
total RNA using a SuperScript IV cDNA
reverse transcriptase (Invitrogen). Quantitative
PCR was performed using the SYBR Select
Master Mix (Applied Biosystems, Carlsbad,
CA) and the ΔΔct method. Expression of each
human gene was normalized to that of Tata-
binding protein (TBP), while expression of the
mouse genes was normalized to that of
hypoxanthine phosphoribosyl transferase
(HPRT). The sets of primers (ThermoFisher
Scientific) were as follows: human ASGR1, 5´-
GGGAAGAAAGATGAAGTCGCTAGA
versus 5´-GCAGGCTGGAGTGATCTTCAC;
human ASGR2, 5´-
GACGGAGGTCCAGGCAATC
versus 5´-
TGGCTCCTAGGGATGTGATCTT; human
LDLR, 5´- AGGAGACGTGCTTGTCTGTC

versus 5′- CTGAGCCGTTGTCGCAGT;
human PCSK9, 5′-
TGGAGCTGGCCTTGAAGTTG
versus 5′-
GATGCTCTGGGCAAAGACAGA; human
SREPB2, 5′-
AGAATGTCCTTCTGATGTCC
versus 5′- GGAGAGTCTGGCTCATCTT;
human TBP, 5′-
CGAATATAATCCCAAGCGGTTT
versus 5′-
GTGGTTCGTGGCTCTCTTATCC; mouse
ASGR1, 5′- TCTGACGTGCGAAGCTTGAG
versus 5′-GGTCCTTTCAGAGCCATTGC;
mouse ASGR, 5′-
CGATGATGAACATGGCTCTCA
versus 5′-AGGCTGCCCTTTCCAGTGT;
mouse HPRT, 5′-
CCGAGGATTTGGAAAAAGTGTT
versus 5′-
CCTTCATGACATCTCGAGCAAGT.
832
Co-immunoprecipitation
For co-immunoprecipitation of LDLR-ASGR1
complexes from HepG2-PCSK9-KO cells
(endogenous LDLR-ASGR1 complex) and
transfected HEK293 cells, cells were lysed in
Pierce non-denaturing IP buffer supplemented
with complete protease inhibitor cocktail.
Lysates containing 0.5 mg total protein
(HepG2-PCSK9-KO cells) or 0.25 mg total
protein (HEK293 cells) were exposed
overnight, at 4 °C on a rocker, to 2 µg of IP
antibody (rabbit anti-ASGR1 antibody for
HepG2-PCSK9-KO cell lysates; mouse
(IgG1)MAB Flag M2 or mouse (IgG2a) anti-
V5 MAB antibodies for HEK293 cell lysates).
The following day, 70 µl True Blot anti-rabbit
IgG beads (HepG2-PCSK9-KO cell lysates) or
40 µl of protein A/G PLUS-Agarose (HEK293
cell lysates) were added for an additional
overnight or 1 hour incubation, respectively.
Following three washes with lysis
buffer/protease inhibitors and two washes with
PBS and elution in 70 µl 2x laemmli sample
buffer, the pull-downs were separated by 8%
Tris-glycine SDS-PAGE and analyzed by WB
for LDLR and ASGR1 using their respective
primary antibodies. Rabbit IgG TrueBlot was
used for the detection of immunoblotted target
proteins from HepG2-PCSK9-KO cell lysates

(endogenous LDLR-ASGR1 complex), which
eliminated the hindrance by interfering
immunoprecipitating immunoglobulin heavy
and light chains. TrueBlot preferentially
detects the non-reduced form of rabbit IgG
over the reduced, SDS-denatured form of IgG.
For co-immunoprecipitation of LDLR-ASGR1
complexes from mice livers, fresh livers from
Pcsk9-KO mice litter-mates were lysed in in
Pierce non-denaturing IP buffer supplemented
with complete protease inhibitor mixture.
Liver lysates containing 1 mg of total protein
were exposed to 2 µg of rabbit anti-ASGR1
antibody overnight at 4 °C on a rocker. The
following morning, 100 µl of Surebeads
Protein G magnetic beads (BioRad) was added
to each sample tube, including "beads only"
negative control. Isolation and purification was
carried out using magnets as per
manufacturer's instructions. A "control" was
included, which consisted of a liver lysate from
a wild type (C57BL/6J) mouse to which
ASGR1 antibody was added to help identify
the IgGs. See Table S1 for antibodies used.
886
Cell surface biotinylation
Biochemical detection of cell surface LDLR
and ASGR1 by Western blot analysis was
performed following the protocol described in
(39). Namely, HEK293 cells seeded in 6-well
plates and after reaching 80% confluency were
transiently co-transfected with 1.5 µg DNA
consisting of V5-tagged LDLR and empty
vector or a combination of LDLR-V5, HA-
tagged ASGR2 and Flag-tagged ASGR1, WT
or carbohydrate-binding mutants
Q240A/W244A/E253A or Q240A/W244A.
Following 48 hours post-transfection, cells
were washed twice with ice-cold PBS and
biotinylated with 2 ml of EZ-link sulfo-NHS-
LC-Biotin (0.5 mg/ml in PBS) for 20 min at
4°C, on ice. After 10 min incubation on ice
with ice-cold BSA (0.1 mg/ml) in PBS to
quench the reaction and two washes with cold
PBS, cells were lysed for 10 min on ice in
biotinylation lysis buffer (20 mM Tris pH 7.5,
5 mM EDTA, 5 mM EGTA, 0.5% N-dodecyl-
N-maltoside, supplemented with protease
inhibitor cocktail without EDTA). Cell lysates
were collected after centrifugation at 3,000
rpm for 10 min at 4 °C, and quantified for

protein concentration. Biotinylated cell surface
proteins (240 µg total protein) were
immunoprecipitated overnight with 30 µl
Streptavidin agarose. Following centrifugation
at 7,000 rpm for 2 min at 4°C, the pellet
containing the plasma membrane pool was
washed three times, eluted in 60 µl 2x laemmli
sample buffer and separated by 8% Tris-
glycine SDS-PAGE. A negative control (-
Biotin) was included, to which PBS was added
instead of Biotin at the biotinylation step. Total
proteins (plasma membrane + intracellular)
were detected from a fraction of the lysates
before Streptavidin immunoprecipitation,
which was referred to as "input".
928
929 *Statistical analysis*
930 Quantifications are defined as averages ± S.D.
931 The statistical significance was evaluated by
932 Student's *t* test, and probability values (*p*)
933 <0.05 were considered significant.
934
AUTHORS CONTRIBUTIONS
NGS and RCA conceived the studies and wrote
the manuscript. DSR, EG, RE, AR, JM, MD,
AE and PFL performed all experiments.

SOURCES OF FUNDING
This work was supported in part by research
grants to Nabil G. Seidah from the CIHR
Foundation Scheme grant (148363), a Canada
Research Chair (950-231335) and a Leducq
Foundation grant (13 CVD 03), as well as to
Richard C. Austin from the Heart and Stroke
Foundation of Canada (G-13-0003064, G-15-
0009389) and the Canadian Institutes of Health
Research (74477, 173520). Financial support
from the Research Institute of St. Joe's
Hamilton is acknowledged. Richard C. Austin
is a Career Investigator of the Heart and Stroke
Foundation of Ontario and holds the Amgen
Canada Research Chair in the Division of
Nephrology at St. Joseph's Healthcare and
McMaster University.
956
957 **CONFLICT OF INTEREST**
958 The authors declare that they do not have
959 anything to disclose regarding funding or
960 conflict of interest regarding this manuscript.
961

962 References
963
1. Wadhera, R. K., Steen, D. L., Khan, I.,
Giugliano, R. P., and Foody, J. M.
(2016) A review of low-density
lipoprotein cholesterol, treatment
strategies, and its impact on
cardiovascular disease morbidity and
mortality. *J. Clin. Lipidol.* **10**, 472-489
2. Seidah, N. G., Benjannet, S., Wickham,
L., Marcinkiewicz, J., Jasmin, S. B.,
Stifani, S., Basak, A., Prat, A., and
Chretien, M. (2003) The secretory
proprotein convertase neural
apoptosis-regulated convertase 1
(NARC-1): liver regeneration and
neuronal differentiation. *Proc. Natl.*
Acad. Sci. U. S. A. **100**, 928-933
3. Maxwell, K. N., and Breslow, J. L.
(2004) Adenoviral-mediated
expression of Pcsk9 in mice results in
a low-density lipoprotein receptor
knockout phenotype. *Proc. Natl. Acad.*
Sci. U. S. A. **101**, 7100-7105
4. Cohen, J., Pertsemlidis, A., Kotowski, I.
K., Graham, R., Garcia, C. K., and
Hobbs, H. H. (2005) Low LDL
cholesterol in individuals of African
descent resulting from frequent
nonsense mutations in PCSK9. *Nat.*
Genet. **37**, 161
5. Seidah, N. G., Abifadel, M., Prost, S.,
Boileau, C., and Prat, A. (2017) The
Proprotein Convertases in
Hypercholesterolemia and
Cardiovascular Diseases: Emphasis on
Proprotein Convertase
Subtilisin/Kexin 9. *Pharmacol. Rev.* **69**,
33-52
6. Abifadel, M., Varret, M., Rabes, J. P.,
Allard, D., Ouguerram, K., Devillers,
M., Cruaud, C., Benjannet, S.,
Wickham, L., Erlich, D., Derre, A.,
Villeger, L., Farnier, M., Beucler, I.,
Bruckert, E., Chambaz, J., Chanu, B.,
Lecerf, J. M., Luc, G., Moulin, P.,
Weissenbach, J., Prat, A., Krempf, M.,
Junien, C., Seidah, N. G., and Boileau,

C. (2003) Mutations in PCSK9 cause autosomal dominant hypercholesterolemia. *Nat. Genet.* **34**, 154-156

7. Cohen, J. C., Boerwinkle, E., Mosley, T. H., Jr., and Hobbs, H. H. (2006) Sequence variations in PCSK9, low LDL, and protection against coronary heart disease. *New England Journal of Medicine* **354**, 1264-1272

8. Sabatine, M. S., Leiter, L. A., Wiviott, S. D., Giugliano, R. P., Deedwania, P., De Ferrari, G. M., Murphy, S. A., Kuder, J. F., Gouni-Berthold, I., Lewis, B. S., Handelsman, Y., Pineda, A. L., Honarpour, N., Keech, A. C., Sever, P. S., and Pedersen, T. R. (2017) Cardiovascular safety and efficacy of the PCSK9 inhibitor evolocumab in patients with and without diabetes and the effect of evolocumab on glycaemia and risk of new-onset diabetes: a prespecified analysis of the FOURIER randomised controlled trial. *Lancet Diabetes Endocrinol* **5**, 941-950

9. Ashwell, G., and Harford, J. (1982) Carbohydrate-specific receptors of the liver. *Annu. Rev. Biochem.* **51**, 531-554

10. Igdoura, S. A. (2017) Asialoglycoprotein receptors as important mediators of plasma lipids and atherosclerosis. *Curr. Opin. Lipidol.* **28**, 209-212

11. Nioi, P., Sigurdsson, A., Thorleifsson, G., Helgason, H., Agustsdottir, A. B., Norddahl, G. L., Helgadottir, A., Magnusdottir, A., Jonasdottir, A., Gretarsdottir, S., Jonsdottir, I., Steinthorsdottir, V., Rafnar, T., Swinkels, D. W., Galesloot, T. E., Grarup, N., Jorgensen, T., Vestergaard, H., Hansen, T., Lauritzen, T., Linneberg, A., Friedrich, N., Krarup, N. T., Fenger, M., Abildgaard, U., Hansen, P. R., Galloe, A. M., Braund, P. S., Nelson, C. P., Hall, A. S., Williams, M. J., van Rij, A. M., Jones, G. T., Patel, R. S., Levey, A. I., Hayek, S., Shah, S. H., Reilly, M., Eyjolfsson, G. I., Sigurdardottir, O., Olafsson, I., Kiemeney, L. A., Quyyumi, A. A., Rader, D. J., Kraus, W. E., Samani, N. J., Pedersen, O., Thorgeirsson, G., Masson, G., Holm, H., Gudbjartsson, D., Sulem, P., Thorsteinsdottir, U., and Stefansson, K. (2016) Variant ASGR1 Associated with a Reduced Risk of Coronary Artery Disease. *N. Engl. J. Med.* **374**, 2131-2141

12. Bischoff, J., and Lodish, H. F. (1987) Two asialoglycoprotein receptor polypeptides in human hepatoma cells. *J. Biol. Chem.* **262**, 11825-11832

13. Ben Djoudi Ouadda, A., Gauthier, M. S., Susan-Resiga, D., Girard, E., Essalmani, R., Black, M., Marcinkiewicz, J., Forget, D., Hamelin, J., Evagelidis, A., Ly, K., Day, R., Galarneau, L., Corbin, F., Coulombe, B., Caku, A., Tagliabracci, V. S., and Seidah, N. G. (2019) Ser-Phosphorylation of PCSK9 (Proprotein Convertase Subtilisin-Kexin 9) by Fam20C (Family With Sequence Similarity 20, Member C) Kinase Enhances Its Ability to Degrade the LDLR (Low-Density Lipoprotein Receptor). *Arterioscler. Thromb. Vasc. Biol.* **39**, 1996-2013

14. Susan-Resiga, D., Girard, E., Kiss, R. S., Essalmani, R., Hamelin, J., Asselin, M. C., Awan, Z., Butkinaree, C., Fleury, A., Soldera, A., Dory, Y. L., Baass, A., and Seidah, N. G. (2017) The Proprotein Convertase Subtilisin/Kexin Type 9-resistant R410S Low Density Lipoprotein Receptor Mutation: A novel mechanism causing familial hypercholesterolemia. *J. Biol. Chem.* **292**, 1573-1590

15. Zaid, A., Roubtsova, A., Essalmani, R., Marcinkiewicz, J., Chamberland, A., Hamelin, J., Tremblay, M., Jacques, H., Jin, W., Davignon, J., Seidah, N. G., and

Prat, A. (2008) Proprotein convertase subtilisin/kexin type 9 (PCSK9): Hepatocyte-specific low-density lipoprotein receptor degradation and critical role in mouse liver regeneration. *Hepatology* **48**, 646-654

16. Gupta, N., Fisker, N., Asselin, M. C., Lindholm, M., Rosenbohm, C., Orum, H., Elmen, J., Seidah, N. G., and Straarup, E. M. (2010) A locked nucleic acid antisense oligonucleotide (LNA) silences PCSK9 and enhances LDLR expression in vitro and in vivo. *PLoS. ONE.* **5**, e10682

17. Roubtsova, A., Chamberland, A., Marcinkiewicz, J., Essalmani, R., Fazel, A., Bergeron, J. J., Seidah, N. G., and Prat, A. (2015) PCSK9 deficiency unmasks a sex/tissue-specific subcellular distribution of the LDL and VLDL receptors in mice. *J. Lipid Res.* **56**, 2133-2142

18. Pedersen, N. B., Wang, S., Narimatsu, Y., Yang, Z., Halim, A., Schjoldager, K. T., Madsen, T. D., Seidah, N. G., Bennett, E. P., Levery, S. B., and Clausen, H. (2014) Low density lipoprotein receptor class A repeats are O-glycosylated in linker regions. *J. Biol. Chem.* **289**, 17312-17324

19. Nassoury, N., Blasiole, D. A., Tebon, O. A., Benjannet, S., Hamelin, J., Poupon, V., McPherson, P. S., Attie, A. D., Prat, A., and Seidah, N. G. (2007) The Cellular Trafficking of the Secretory Proprotein Convertase PCSK9 and Its Dependence on the LDLR. *Traffic.* **8**, 718-732

20. Ward, S. E., O'Sullivan, J. M., Drakeford, C., Aguila, S., Jondle, C. N., Sharma, J., Fallon, P. G., Brophy, T. M., Preston, R. J. S., Smyth, P., Sheils, O., Chion, A., and O'Donnell, J. S. (2018) A novel role for the macrophage galactose-type lectin receptor in mediating von Willebrand factor clearance. *Blood* **131**, 911-916

21. Demers, A., Samani, S., Lauzier, B., Des Rosiers, C., Sock, E. T., Ong, H., and Mayer, G. (2015) PCSK9 Induces CD36 Degradation and Affects Long-Chain Fatty Acid Uptake and Triglyceride Metabolism in Adipocytes and in Mouse Liver. *Arterioscler Thromb Vasc Biol.* **35**, 2517-2525

22. Seidah, N. G., and Prat, A. (2012) The biology and therapeutic targeting of the proprotein convertases. *Nat Rev Drug Discov.* **11**, 367-383

23. Benjannet, S., Rhainds, D., Hamelin, J., Nassoury, N., and Seidah, N. G. (2006) The proprotein convertase PCSK9 is inactivated by furin and/or PC5/6A: Functional consequences of natural mutations and post-translational modifications. *J. Biol. Chem.* **281**, 30561-30572

24. Essalmani, R., Susan-Resiga, D., Chamberland, A., Abifadel, M., Creemers, J. W., Boileau, C., Seidah, N. G., and Prat, A. (2011) In vivo evidence that furin from hepatocytes inactivates PCSK9. *J Biol. Chem.* **286**, 4257-4263

25. Weigel, P. H., and Yik, J. H. (2002) Glycans as endocytosis signals: the cases of the asialoglycoprotein and hyaluronan/chondroitin sulfate receptors. *Biochim Biophys Acta* **1572**, 341-363

26. Meier, M., Bider, M. D., Malashkevich, V. N., Spiess, M., and Burkhard, P. (2000) Crystal structure of the carbohydrate recognition domain of the H1 subunit of the asialoglycoprotein receptor. *J. Mol. Biol.* **300**, 857-865

27. Tanowitz, M., Hettrick, L., Revenko, A., Kinberger, G. A., Prakash, T. P., and Seth, P. P. (2017) Asialoglycoprotein receptor 1 mediates productive uptake of N-acetylgalactosamine-conjugated and unconjugated phosphorothioate antisense oligonucleotides into liver

hepatocytes. *Nucleic Acids Res.* **45**, 12388-12400

28. Stoekenbroek, R. M., Kallend, D., Wijngaard, P. L., and Kastelein, J. J. (2018) Inclisiran for the treatment of cardiovascular disease: the ORION clinical development program. *Future Cardiol.* **14**, 433-442

29. Seidah, N. G., Prat, A., Pirillo, A., Catapano, A. L., and Norata, G. D. (2019) Novel strategies to target proprotein convertase subtilisin kexin 9: beyond monoclonal antibodies. *Cardiovasc. Res.* **115**, 510-518

30. Thomas, G. (2002) Furin at the cutting edge: from protein traffic to embryogenesis and disease. *Nature Reviews Molecular Cell Biology* **3**, 753-766

31. Gu, Y., Cao, J., Zhang, X., Gao, H., Wang, Y., Wang, J., Zhang, J., Shen, G., Jiang, X., Yang, J., Zheng, X., Xu, J., Zhang, C. C., Lan, F., Qu, D., Zhao, Y., Xu, G., Xie, Y., Luo, M., and Lu, Z. (2020) Interaction network of SARS-CoV-2 with host receptome through spike protein. *bioRxiv*, 2020.2009.2009.287508

32. Morens, D. M., and Fauci, A. S. (2020) Emerging Pandemic Diseases: How We Got to COVID-19. *Cell* **182**, 1077-1092

33. Zamorano Cuervo, N., and Grandvaux, N. (2020) ACE2: Evidence of role as entry receptor for SARS-CoV-2 and implications in comorbidities. *Elife* **9**

34. Mokhtari, T., Hassani, F., Ghaffari, N., Ebrahimi, B., Yarahmadi, A., and Hassanzadeh, G. (2020) COVID-19 and multiorgan failure: A narrative review on potential mechanisms. *J. Mol. Histol.* **51**, 613-628

35. Munzer, J. S., Basak, A., Zhong, M., Mamarbachi, A., Hamelin, J., Savaria, D., Lazure, C., Hendy, G. N., Benjannet, S., Chretien, M., and Seidah, N. G. (1997) In vitro characterization of the novel proprotein convertase PC7. *J. Biol. Chem.* **272**, 19672-19681

36. Ozden, S., Lucas-Hourani, M., Ceccaldi, P. E., Basak, A., Valentine, M., Benjannet, S., Hamelin, J., Jacob, Y., Mamchaoui, K., Mouly, V., Despres, P., Gessain, A., Butler-Browne, G., Chretien, M., Tangy, F., Vidalain, P. O., and Seidah, N. G. (2008) Inhibition of Chikungunya Virus Infection in Cultured Human Muscle Cells by Furin Inhibitors: Impairment of the maturation of the E2 surface glycoprotein. *J. Biol. Chem.* **283**, 21899-21908

37. Seidah, N. G., Mowla, S. J., Hamelin, J., Mamarbachi, A. M., Benjannet, S., Toure, B. B., Basak, A., Munzer, J. S., Marcinkiewicz, J., Zhong, M., Barale, J. C., Lazure, C., Murphy, R. A., Chretien, M., and Marcinkiewicz, M. (1999) Mammalian subtilisin/kexin isozyme SKI-1: A widely expressed proprotein convertase with a unique cleavage specificity and cellular localization. *Proc. Natl. Acad. Sci. U. S. A.* **96**, 1321-1326

38. Roubtsova, A., Munkonda, M. N., Awan, Z., Marcinkiewicz, J., Chamberland, A., Lazure, C., Cianflone, K., Seidah, N. G., and Prat, A. (2011) Circulating Proprotein Convertase Subtilisin/Kexin 9 (PCSK9) Regulates VLDLR Protein and Triglyceride Accumulation in Visceral Adipose Tissue. *Arteriosclerosis, Thrombosis, and Vascular Biology* **31**, 785-791

39. Hassan, H. H., Bailey, D., Lee, D. Y., Iatan, I., Hafiane, A., Ruel, I., Krimbou, L., and Genest, J. (2008) Quantitative analysis of ABCA1-dependent compartmentalization and trafficking of apolipoprotein A-I: implications for determining cellular kinetics of nascent high density lipoprotein biogenesis. *J. Biol. Chem.* **283**, 11164-11175

FIGURES AND FIGURE LEGENDS

Figure 1: Expression of ASGR1 and ASGR2 in mice and cell lines

(A) Schematic representation of ASGR1, depicting its cytoplasmic N terminus (40 amino acids), the single transmembrane domain (TM) and its exoplasmic C terminus (~230 aa) that is composed of a stalk region and a C-terminal carbohydrate recognition domain (CRD) with three reported residues important in carbohydrate binding. (B) *In situ* hybridization of ASGR1 and ASGR2 in mice before and after birth. (C) ASGR1 and ASGR2 quantification by quantitative RT-PCR in mouse (m) liver, HepG2 naïve, HepG2-PCSK9-KO, and IHH cells.

Figure 2: Comparative effects of PSCK9 on cellular LDLR and ASGR1 levels

HepG2 naïve cells were incubated for 20 hours with media only (- PCSK9) or media containing purified PCSK9 (3 μg/ml) (+ PCSK9). (A) WB analysis of cellular LDLR and ASGR1 expression and of PCSK9 in the media. (B) Immunofluorescence microscopy of plasma membrane LDLR and ASGR1 (green signal, non-permeabilized conditions). Scale bar, 15 μm. HepG2 naïve cells were transfected with non-targeting siRNA (siCnt) or siRNA PCSK9 (siPCSK9) at final concentrations of 10 nM and analyzed 48 hours post-transfection. (C) Quantitative RT-PCR analysis. (D) WB analysis of cellular LDLR and ASGR1 and of secreted PCSK9. Data are representative of at least three (A-B, D) and at least two independent experiments (C), respectively. Quantifications are averages ± SD. * p<0.05; *** p<0.001 (t-test).

Figure 3: LDLR and ASGR1 expression levels in livers from wild-type (WT) mice and mice lacking mPCSK9 (*PCSK9*-KO) or mLDLR (*Ldlr*-KO), respectively

(A) LDLR and ASGR1 staining by immunohistochemistry (green signal) in mouse liver sections. (B) WB analysis of total ASGR1 and LDLR proteins in mouse livers. Immunoblots were cropped for clarity. Data are representative of 9-10 mice (A) and 3-4 mice (B) and quantifications are averages ± SD. * p<0.05; ** p<0.01; *** p<0.001 (t-test).

Figure 4: ASGR1 does not affect PCSK9 expression and secretion

HepG2 naïve cells were transfected for 48 hours with non-targeting siRNA (siCnt) or siRNA ASGR1 (siASGR1) at final concentrations of 20 nM and analyzed by quantitative RT-PCR (A), or WB for cellular LDLR, PCSK9, ASGR1 and for secreted PCSK9 (B). Data are representative of three independent experiments. Quantifications are averages ± SD. *** p<0.001 (t-test).

Figure 5: ASGR1 regulates LDLR levels and functionality independently of PCSK9

HepG2 naïve cells (A, C) or HepG2-PCSK9-KO cells (B, D) were transfected for 48 hours with non-targeting siRNA (si-Cnt) or siRNA ASGR1 (si-ASGR1) at final concentrations of 20 nM. Cells were then incubated for the last 20 hours with media only (- PCSK9) or media containing purified PCSK9 (3 μg/ml) (+ PCSK9) (A-C). (A-B) Total protein levels of LDLR and ASGR1 were assessed by WB. (C) Immunofluorescence microscopy of LDLR and ASGR1 at the plasma membrane (green signal, non-permeabilized condition). (D) Immunofluorescence microscopy of LDLR and ASGR1 (green signal) at the plasma membrane (PM, non-permeabilzed condition) (top) and total levels (Total, permeabilized condition) (bottom). LDLR functionality is assessed by DiI-LDL (5 μg/ml) internalization for 2 hours at 37°C before fixation (red signal in upper left panel). Scale bar, 15 μm. (E) Quantitative RT-PCR analysis was performed on HepG2 naïve cells transfected for 48 hours with non-targeting siRNA (si-Cnt) or siRNA ASGR1 (si-ASGR1) at final concentrations of 20 nM. Data are representative of at least three independent experiments. Quantifications are averages ± SD. * p<0.05; *** p<0.001 (t-test).

Figure 6: Knockdown of either PCSK9 or ASGR1 enhances LDLR levels

HepG2 naïve cells were transfected for 48 hours with 10 nM non-targeting siRNA (siCnt), siRNA PCSK9 (siPCSK9), or siRNA ASGR1 (siASGR1). Immunofluorescence microscopy of plasma membrane LDLR (green signal, non-permeabilized condition) and DiI-LDL (5 µg/ml) internalization for 2 hours at 37°C before fixation (red signal) and quantifications are shown. Data are representative of two independent experiments. Quantifications are averages ± SD. *** $p < 0.001$ (t-test).

Figure 7: Overexpressed ASGR1 and ASGR2 reduce LDLR levels independent of PCSK9

HepG2-PCSK9-KO cells were transfected with Flag-tagged ASGR1 (A1) and HA-tagged ASGR2 (A2) (A1+A2) or control empty vector (v), (LDLR endogenous) (A-B), or in combination with LDLR (LDLR overexpressed) (A). (A) WB analyses and quantification of total cellular LDLR, ASGR1 (Flag-HRP) and ASGR2 (HA-HRP). (B) Immunofluorescence microscopy of endogenous cell-surface LDLR (green signal, non-permeabilized condition) and DiI-LDL uptake (red signal) for 2 hours at 37°C before fixation. Quantification was performed only in cells transfected with ASGR1 and ASGR2 (white signal). Scale bar, 15 μm. Immunoblots were cropped for clarity. Data are representative of at least four independent experiments. Quantifications are averages ± SD. *** p<0.001 (t-test).

Figure 8: LDLR and ASGR1 form a complex that is not limited to lectin binding

(A) Co-immunoprecipitation of endogenous LDLR and ASGR1 in HepG2-PCSK9-KO cells. Pull-downs with ASGR1 antibody were analyzed by TrueBlot for LDLR and ASGR1. To "Beads only" negative control addition of ASGR1 antibody was omitted. "Input" represents 10% of the original material subjected to co-immunoprecipitation. (B) Co-immunoprecipitation from liver extracts of WT and *PCSK9*-KO mice. Pull-downs with ASGR1 antibody were analyzed by WB for LDLR and ASGR1. To "Beads only" negative control addition of ASGR1 antibody was omitted. "Control" (IgG migration control) consists of WT mouse liver lysate (30 µg) plus ASGR1 antibody. (C) Co-immunoprecipitation from HEK293 cells transfected with either vector (v), V5-tagged LDLR or Flag-tagged ASGR1, WT or mutant Q240A/W244A/E253A (QA/WA/EA), or a combination of ASGR1 and LDLR. Pull-downs with Flag M2 antibody (left panel) or V5 antibody (right panel) were analyzed by WB for LDLR and ASGR1. "Input" represents 10% of the original material subjected to co-immunoprecipitation.

Immunoblots were cropped for clarity. Data are representative of two independent experiments.

Figure 9: The lectin binding domain in ASGR1 is important for the ASGR1-mediated degradation of the functional, mainly non-O-glycosylated, LDLR at the plasma membrane

HEK293 cells were co-transfected with LDLR and empty vector (v) or LDLR, HA-tagged ASGR2 and Flag-tagged ASGR1, WT or carbohydrate-binding mutants Q240A/W244A/E253A (QA/WA/EA) (A-C) or Q240A/W244A (QA/WA) (C). (A) Immunofluorescence microscopy of plasma membrane (non-permeabilized condition) overexpressed LDLR (PM LDLR) (green signal) and DiI-LDL uptake (red signal) for 2 hours at 37°C before fixation. Non-transfected cells (v) expressed ~100-fold less LDLR than transfected ones. Scale bar, 15 μm. (B) Surface protein biotinylation: cells incubated with Biotin (+ Biotin) or PBS (-Biotin) were lysed and plasma membrane (PM) proteins were pulled-down with streptavidin and analyzed by WB for LDLR (V5-HRP) and ASGR1 (Flag-HRP) (left panel). For reference, one eigth (30 μg) of the (+Biotin) cell lysate before streptavidin pull-down (input) was analyzed by WB (right panel). (C) WB analyses of total cellular LDLR (mature mLDLR, and immature iLDLR), ASGR1 (Flag-HRP) and ASGR2 (HA-HRP), and of shed LDLR in the media. (D) HepG2-PCSK9-KO cells were transfected for 48 hours with 20 nM non-targeting (Cnt) siRNA or siRNA ASGR1. WB analyses of total cellular LDLR (mLDLR and iLDLR), ASGR1 and of shed LDLR in the media are shown. Note that 13% of total media and 8% of total lysates are loaded on the gel and that the exposure times are 15 min and 3 min, respectively. Immunoblots were cropped for clarity. Data are representative of at least three independent experiments. Quantifications are averages ± SD. * p<0.05; ** p<0.01; *** p<0.001 (t-test).

Figure 10: ASGR1 specificity for LDLR

HepG2-PCSK9-KO cells were transfected with 20 nM non-targeting (Cnt) siRNA or siRNA ASGR1 for 48 hours. WB analyses and quantification of total cellular LDLR, ASGR1, CD36 and insulin receptor (IR β) from three independent experiments are shown. Immunoblots were cropped for clarity. Quantifications are averages ± SD. *** p<0.001 (t-test).

Figure 11: ASGR1 is shed in the media by Furin

(A) HEK293 cells were co-transfected with Flag-tagged ASGR1 and either one of the liver expressed PCs, PACE4, PC5A, PC7, Furin, or empty vector (v), as negative control. WB of cellular ASGR1 (Flag-HRP) and of shed ASGR1 (Flag-HRP) in the media is shown. HEK293 cells (B-D) or HepG2-PCSK9-KO cells (E) were co-transfected with Flag-tagged ASGR1, WT or Furin-like cleavage site mutants $R_{71}A$-$R_{74}A$ (B), $R_{100}A$-$K_{103}A$ (C, E), $RRRREL_{105}$ (D, E), and V5-tagged Furin (+) or empty vector (-). WB analyses of cellular Furin (V5-HRP) and ASGR1 (Flag-HRP) and of shed ASGR1 (Flag-HRP) in the media are depicted. The slower SDS-PAGE migration of shed ASGR1 $RRRREL_{105}$ mutant in (D) and (E) is due to an increased negative charge (glutamic acid, $\underline{E}$) following the Furin cleavage site in mutant protein $RRRR_{103}\downarrow ELE$ compared to WT $RKMK_{103}\downarrow SLE$. Immunoblots were cropped for clarity. Data are representative of at least three independent experiments.