To buffer or to be labile? A framework to disentangle demographic

patterns and evolutionary processes

Gabriel Silva Santos [1,*], Roberto Salguero-Gómez [2], André Tavares Corrêa Dias[3],

Maja Kajin [1,4]

1 Department of Ecology, Graduate Program in Ecology and Evolution, Rio de Janeiro

State University, 524 São Francisco Xavier street, 20550-900, Maracanã, Rio de

Janeiro, Brazil

2 Department of Zoology, University of Oxford, 11a Mansfield Road, OX1 3SZ,

Oxford, UK

3 Departamento de Ecologia, Instituto de Biologia, Universidade Federal do Rio de

Janeiro (UFRJ), Rio de Janeiro, RJ, Brazil.

4 Department of Biology, Biotechnical Faculty, University of Ljubljana, Ljubljana,

Slovenia

[*] Corresponding author: ssantos.gabriel@gmail.com

Abstract

Until recently, natural selection was assumed to reduce temporal fluctuation in vital rates due to its negative effects on population dynamics – the so-called Demographic Buffering Hypothesis (DBH). After several failures to support the DBH in the two decades since it was first posited, an alternative hypothesis was suggested; the Demographic Lability Hypothesis (DLH), where population vital rates should track rather than buffer the environmental conditions. Despite the huge contribution of both hypotheses to comprehend the demographic strategies to cope the environmental stochasticity, it remains unclear if they represent two competing patterns or the extreme ends of a continuum encompassing all demographic strategies. To solve this historical debate, we unify several methods with an integrative theoretical approach where: i) using the sum of stochastic elasticity with respect to mean and variance – a first-order derivative approach – we rank species on a Buffering-Lability (DB-DL) continuum and ii) using the second-order derivative, we examine how vital rates are shaped by natural selection. Our framework, applied to 40 populations of 34 mammals, successfully placed the species on the DB-DL continuum. We could also link the species' position on the DB-DL continuum to their generation time and time to recovery. Moreover, the second-order derivative unveiled that vital rates with lower temporal variation are not necessarily under a strong pressure of stabilizing selection, as predicted by DBH and DLH. Our framework provides an important step towards unifying the different perspectives of DBH and DLH with key evolutionary concepts.

Keywords: COMADRE Animal Matrix Database, elasticity, life-history evolution, second-order derivative, sensitivity, vital rate variance.

Introduction

Modern demographic theory offers a wide array of hypotheses to explain the evolutionary processes that shape the diversity of demographic responses to environmental stochasticity (Charlesworth 1994, Pfister 1998, Healy *et al.* 2019, Hilde *et al*. 2020). Of these, two stand out as the most promising in linking evolutionary processes to demographic patterns: the Demographic Buffering Hypothesis (DBH) and its alternative hypothesis, the Demographic Lability Hypothesis (DLH) (Koons *et al.* 2009, Jongejans *et al*. 2010, McDonald *et al*. 2017, Hilde *et al*. 2020). Together, the DBH and DLH (hereafter *DB-DL perspective*), represent two contrasting ways by which natural populations cope with environmental stochasticity by either buffering (DBH) or tracking (DLH) changes in environmental conditions (Morris *et al*. 2008, McDonald *et al*. 2017, Hilde *et al*. 2020).

Coined over two decades ago, the DBH postulates that vital rates (*e.g.* survival, fecundity) that have the greatest influence on population growth should buffer the negative effects of the environmental stochasticity by presenting lower temporal variation than less influential vital rates (Pfister 1998). Because local extinction risk typically increases with increasing fluctuations in population size (Ovaskainen and Meerson 2010), demographic buffering is hypothesized to be a result of natural selection. This evolutionary process reducing variation in vital rates, called environmental canalization (Debat and David 2001, Gaillard and Yoccoz 2003), is expected to be more intense in vital rates that contribute the most to population growth rate (Pfister 1998).

Demographic buffering is expected to be particularly important for species that are less capable to recover after perturbations, which have also a higher risk of extinction due to demographic stochasticity (*e.g.* long-lived trees and primates, Morris

et al. 2008, 2011, Hilde *et al.* 2020). Contrastingly, the DLH postulates that some

species might take advantage if their vital rates were labile enough to permit population

size track rather than buffering the environmental stochasticity (Koons *et al.* 2009,

2014, Jongejans *et al.* 2010, Hilde *et al.* 2020).

The DB-DL perspective simultaneously reflects *demographic patterns, i.e.* how

populations respond to the environmental stochasticity (Morris *et al.* 2008, Jongejans *et*

al. 2010, McDonald *et al.* 2017), and *evolutionary processes, i.e.* how the life-histories

of the individuals that make up those populations were shaped by natural selection

(Pfister 1998, Gaillard and Yoccoz 2003, Koons *et al.* 2009). Both demographic and

evolutionary facets are important to predict the eco-evolutionary dynamic in response to

climate changes (Metcalf and Pavard 2007, Burns *et al.* 2010, Salguero-Gómez *et al.*

2018). Despite the importance of understanding demographic responses to

environmental changes, existing approaches to assess the DBH and DLH present

several constraints and can therefore hinder our understanding of the eco-evolutionary

processes behind the DB-DL perspective (Box 1, 2).

After two decades of consolidating the methodological bases of the DB-DL

perspective (Hilde *et al.* 2020), a lack of consensus on how to identify evidence in favor

of the DBH and DLH still persists (Box 1). Moreover, the bases of the DBH and DLH

have been mainly developed in terms of mathematical models rather than trait-based

mechanisms (Hilde *et al.* 2020). Indeed, only few authors have suggested trait-based

mechanisms to support the DB-DL perspective (Morris *et al.* 2008, McDonald *et al.*

2017). Of those, even the prevalence of DBH in long-lived species is not well supported

(Morris *et al.* 2011, Reed and Slade 2012, McDonald *et al.* 2017). Additionally, other

attempts have found no correlation between the DBH and DLH and other life-history

traits (*e.g.* generation time), environmental variables, nor phylogenetic relationships

(McDonald *et al.* 2017).

In this study, we show that the lack of patterns in the assessment of DBH and

DLH arises from different interpretations of what the two main approaches to assess

them consider as evidence for DBH and DLH. Here we will refer to these approaches as

multi-species comparative approaches (*e.g.* Morris *et al.* 2008, McDonald *et al.* 2017)

and (very often single population) hypothesis testing approaches (*e.g.* Reed and Slade

2012, Jäkäläniemi *et al.* 2013). We argue that these approaches represent two

perspectives with very distinctive eco-evolutionary implications that should be

disentangled to better understand the performance and the evolutionary processes of the

buffering and lability demographic strategies. To better assess the multiple facets of the

DB-DL perspective, we reconcile different methods developed over the last two decades

in a unified framework. Finally, we demonstrate the insights provided by our

framework to better understand how species and populations both respond to and are

shaped by environmental stochasticity.

Box 1. Standard protocol to assess the DBH and DLH

The usage of simple correlation analyses (particularly the Spearman's correlation) has been the most frequent approach to assess the DBH and DLH ever since they were posited (Pfister 1998, Hilde *et al.* 2020). Yet, two competing interpretations exist regarding the outcome of such correlation analyses (below). The lack of agreement between the two interpretations highlights the lack of consensus on how the demographic buffering vs. lability strategies manifest themselves in life histories.

The two competing interpretations of the correlation analyses to assess the DBH and DLH are based on different perspectives. Some researchers are interested in unveiling the patterns and distribution of the two demographic strategies - DBH and DLH across the tree of life in a multi-species comparative approaches (Jongejans *et al*. 2010, McDonald *et al*. 2017). Contrastingly, other researchers who assess DBH and DLH in terms of hypothesis testing are interested in finding support for one of these hypotheses by examining how natural selection have shaped the variance in vital rates (Reed and Slade 2012, Rotella *et al*. 2012). While the hypothesis testing approach looks for evidence to corroborate the patterns expected by either DBH or DLH (*e.g.* Jäkäläniemi *et al*. 2013), the multi-species comparative approaches assumes that all demographic strategies should lie somewhere between completely buffered (DBH) and completely labile (DLH)(*e.g.* McDonald *et al*. 2017). However, despite the well documented evidence in favor of the DBH, only scarce evidence exists to support the DLH (Hilde *et al*. 2020). Furthermore, the quantitative assessment of DBH and DLH reflects these different views of the comparative and hypothesis testing approaches as we show below.

Consider the example in Fig 1, where a negative Spearman's correlation coefficient of -0.45 with $P > 0.05$ is shown. Despite the non-significant results, patterns

like this have been used as evidence of demographic buffering in the multi-species

comparative approaches (*e.g.* Doak *et al.* 2005, McDonald *et al.* 2017). Thus, the

correlation coefficient is used as an index to place populations into the continuum,

ranging from completely demographically buffered (Spearman's correlation ρ = -1) to

completely demographically labile (ρ = 1), regardless of the P-value (McDonald *et al.*

2017) - hereafter, *Demographic buffering-lability continuum* (or *DB-DL continuum*).

However, other authors interpret the absence of statistical support for the DBH as

evidence in favor of the DLH (*e.g.* Jäkäläniemi *et al.* 2013) or the absence of any

particular strategy, neither DBH nor DLH (*e.g.* Reed and Slade 2012).

Figure 1. A hypothetical Matrix Population Model comparing the temporal variation in

each matrix element. Letters represent (st-)age transitions (growth, stasis, and shrinkage

– t) and recruitment (r), and the subscripts in the matrix element a_{ij} correspond to

transitions or recruitment events from stage *j* (column in matrix ***A***) in time *t,* to state *i*

(row in ***A***) in t+1. The solid line represents the linear trend calculated by the least

squares regression. The Spearman's correlation coefficient (ρ) between the two

variables and its P-value are informed in the figure.

Disentangling demographic patterns and evolutionary processes in the DB-DL perspective: buffering mechanisms vs. canalization of variance

All populations are buffered in someway (Hunter 2009). Thus, the major question is not *if* but *to what extent* the populations are dependent on environmental stochasticity. Some traits, like endothermy or migration, can help individuals to buffer (or avoid) the negative effects of environmental conditions (Fuller *et al*. 2010, Edic *et al*. 2020). On the another hand, traits like fixed litter size and long parental care may be buffered, once they remain constant independently of environmental conditions (Meiri *et al*. 2020, Santos *et al*. *in prep*). All these traits which increase the capacity of an individual to buffer the effects of environmental stochasticity may be considered as buffering mechanisms (Santos *et al. in prep*). Ultimately, the capacity of the species to buffer the environmental stochasticity should be proportional to the set of buffering mechanisms of these species - including those traits which limit the population to increase in the good years (*e.g.* bonanzas).

Some patterns are expected to arise from these multiple traits. For example, bigger, long-lived species should be more buffered than smaller, shorter-lived species (Morris *et al*. 2008). Yet, such patterns do not tell us how these buffering traits have been shaped by natural selection (*i.e.* what processes led to these patterns). To assess the processes which underly these patterns, we need to analyze the *force and direction of natural selection acting on them*. Thus, if the temporal variation in an important buffering trait (*e.g.* longevity, age of first reproduction) exerts a negative impact on individual performance, its temporal variation might be canalized - *i.e. stabilizing selection* (Debat and David 2001) - increasing the population's capacity to buffer the environmental stochasticity. Otherwise, natural selection might promote variation in the

171 expression of these traits if they improve the individual performance in a variable

172 environment, *i.e. disruptive selection* (Pasztor *et al*. 2000, Koons *et al*. 2009, 2014).

A unified framework to assess the multiple facets of the DBH and DLH

Here, we propose a framework that overcomes the limitations of previous

approaches (Box 1, 2) by explicitly disentangling demographic patterns from the

evolutionary processes shaping the different vital rates. The application of this

framework requires three steps which include two additional steps to the most largely

used approaches to quantify DBH *vs*. DLH: the correlative analyses (Box 1). The **step 1**

in the framework consists of comparing to what extent the life histories are variable and

rank the populations of multiple species according to whether they are buffered or

environmental dependent (labile) (Fig. 2A). **Step 2** and **step 3** are focused on

populations rather than species. The **step 2** is analogous to the correlative analyses.

However, we limited this step to visual inspection of biplots representing the relation

between the temporal variance of vital rates and their contribution to population growth

rate for each population. The graphical inspection of biplots helps to make inferences

about the forces of natural selection acting on each vital rate before the empirical

assessment by step 3. Finally, **step 3** enables to assess how temporal variance in

survival and reproduction are shaped by natural selection (Figs. 2 B, C).

Our unified framework provides a link between the DBH and DLH to

evolutionary mechanisms, *i.e.* disruptive and stabilizing selection, where natural

selection acts to reduce or increase the temporal variance in a specific vital rate (Fig.

2C). The full approach of our framework is explained in detail below, after a concise

description of structured population models (in terms of both matrix population models

[MPM hereafter; Caswell 2001] and integral projection models [IPM hereafter; Ellner *et*

al. 2016]) - the common tools to assess the relative contribution of vital rates to

population growth rate (de Kroon *et al*. 1986, Pfister 1998, Hilde *et al*. 2020).

The mathematical bases of the framework

The DBH and DLH have been mainly assessed by MPM (*e.g.* Pfister 1998,

Rotella *et al*. 2012) and IPM (*e.g.* Rodríguez-Caro *et al*. 2020), which represent the

analytical foundation for our framework. Both MPMs and IPMs are stage-structured,

discrete-time demographic models (Caswell 2001, Ellner et al. 2016). In both,

individuals in a population are categorized according to discrete (*e.g.* age, stage, as in

MPMs; Caswell 2001) or continuous traits (*e.g.* size, height, as in IPMs; Easterling *et*

al. 2000). For simplicity, for the rest of the framework description we focus on the

usage of MPMs, but note that the same approaches are applicable to IPMs (*e.g.* Griffith

2017).

MPMs are matrix representations of life cycles (Ebert 1999). In an MPM **A**,

each element a_{ij} represents a demographic process, for example, a transition from a (st-

)age to another or a contribution of a given (st-)age to the recruitment of new

individuals. It is well known that each demographic process, a_{ij}, contributes differently

to the population growth rate (λ) depending on the life cycle and life history of the

species (Silvertown *et al*. 1993). This contribution of a_{ij} to λ, named *sensitivity*, can be

assessed by simulating small changes on $a_{i,j}$ or by the partial first-order derivative of

each element in the matrix **A** with respect to λ (Caswell 2001, Tuljapurkar *et al*. 2003).

As the sensitivities are dependent on the initial values of a_{ij}, they are not comparable as

some represent survival (bounded from 0-1) while others represent recruitment ($\geq$0).

Thus, to compare sensitivities for survival and recruitment, they must be rescaled to

proportional contributions to λ, the so-called *elasticities* e_{ij} (Caswell 2001).

As survival and recruitment are always, in some way, dependent on the current

environmental conditions (Hunter 2009), we must expect (1) different values of a_{ij} for

each time t sampled and (2) a different population growth rate for each time t, λ_t

(Tuljapurkar *et al*. 2003). Moreover, in a stochastic environment, stochastic population

growth rate (λ_s) is a multiplicative process of bad and good years for the population

(Morris and Doak 2002). Due to the multiplicative process that characterizes the λ_s, the

arithmetic mean of multiples λ_t ($\bar{\lambda}$) is not equivalent to λ_s because the temporal

dependence of λ_s was not accounted for in $\bar{\lambda}$ (Tuljapurkar *et al*. 2003). Indeed, very

often $\lambda_s < \bar{\lambda}_t$ (Morris and Doak 2002). Additionally, elasticity in a stochastic

perspective, namely stochastic elasticity, e_{ij}^s, is composed of its mean and its variance,

which both contribute to the stochastic population growth rate λ_s (Tuljapurkar *et al*.

2003). Integrating these stochastic elasticities allow us to assess the effect of variation

in each vital rate weighted by its relative contribution to λ_s as explained below.

Comparing different life history responses to the environmental conditions

The sum of all stochastic elasticities ($\Sigma E_{a_{ij}}^S$) of an MPM **A** must be equal to 1

because it represents the contribution of all elements a_{ij} to the stochastic population

growth rate (λ_s) (Tuljapurkar *et al.* 2003, Haridas and Tuljapurkar 2005). Moreover the

sum of all stochastic elasticities ($\Sigma E_{a_{ij}}^S$) can be decomposed to assess how temporal

variance and mean values of each vital rates contribute to λ_s – they are respectively the

sum of stochastic elasticity with respect of the variance $\Sigma E_{a_{ij}}^{S^\sigma}$, and the *sum of stochastic*

elasticity with respect of the mean $\Sigma E_{a_{ij}}^{S^\mu}$. Once again, $\Sigma E_{a_{ij}}^{S^\sigma} + \Sigma E_{a_{ij}}^{S^\mu}$ should be equal to

1 because $\Sigma E_{a_{ij}}^S = \Sigma E_{a_{ij}}^{S^\sigma} + \Sigma E_{a_{ij}}^{S^\mu}$.

The $\Sigma E_{a_{ij}}^{S^\sigma}$, if applied to MPMs, might range from 0 to -1, producing a continuum

of species' responses that is similar to the predictions provided by the DBH and DLH

(Box 1) – the *DB-DL* continuum (Haridas and Tuljapurkar 2005, Morris *et al.* 2008). A

population with $\Sigma E_{a_{ij}}^{S^\sigma}$ close to 0 - the *Demographic Buffered-end* (or *DB-end*) - has only

a minor temporal variation in its vital rates, particularly in those vital rates that have a

major contribution to λ_s. The higher the variance in the most important vital rates, the

further the population will be from $\Sigma E_{a_{ij}}^{S^\sigma} = 0$ (Fig. 2A). In other words, the further a

population is from $\Sigma E_{a_{ij}}^{S^\sigma} = 0$, the closer it is to the *DL-end* of the DB-DL continuum.

As the $\Sigma E_{a_{ij}}^{S} = \Sigma E_{a_{ij}}^{S^\sigma} + \Sigma E_{a_{ij}}^{S^\mu}$, any change in $\Sigma E_{a_{ij}}^{S^\sigma}$ will reflect in $\Sigma E_{a_{ij}}^{S^\mu}$, thus, both

values will provide the same information.

The values of $\Sigma E_{a_{ij}}^{S^\sigma}$ and $\Sigma E_{a_{ij}}^{S^\mu}$ represent how species responded to the

environmental stochasticity during the time sampled (Tuljapurkar *et al.* 2003). These

values should vary from population to population according to the environmental

conditions which populations were exposed to at the time, thus, the $\Sigma E_{a_{ij}}^{S^\sigma}$ and $\Sigma E_{a_{ij}}^{S^\mu}$

have also been referred as *the relative effect of environmental variation to population*

growth rate (Morris *et al.* 2008). Yet, the $\Sigma E_{a_{ij}}^{S^\sigma}$ and $\Sigma E_{a_{ij}}^{S^\mu}$ should vary according to

species-specific range, because there are limits to how the most important vital rates can

vary without the population becoming extinct, particularly regarding survival (*e.g.*

Rodríguez-Caro *et al.* 2020). We also expect limits on reproductive output imposed by

physiology and morphology of some species (*e.g.* Meiri *et al.* 2020). Ultimately, both

$\Sigma E_{a_{ij}}^{S^\sigma}$ and $\Sigma E_{a_{ij}}^{S^\mu}$ and the coefficient of correlation, can be used to place populations in a

DB-DL continuum (Box 1). However, unlike previous methods, the $\Sigma E_{a_{ij}}^{S^\sigma}$ and $\Sigma E_{a_{ij}}^{S^\mu}$

overcomes the key limitations characteristic of the correlative analyses and allows

assessing DBH and DLH evidence more clearly (Box 2).

Evolutionary processes have been immediately evoked to explain the patterns

arising from the comparative approach when assessing the position of species along the

DB-DL continuum (Morris *et al.* 2011, McDonald *et al.* 2017). Thus, species placed at

the DB-end of the continuum are commonly expected to be under strong selection

pressures to reduce temporal variation in their most important vital rates (Gaillard and

Yoccoz 2003). Contrastingly, natural selection is claimed to promote temporal variation

in the most important vital rates in those species placed at DL-end of the continuum

(Koons *et al.* 2009, 2014). Such evolutionary claims behind the DB-DL continuum are

key to bridge the gap between demographic and evolutionary processes but remain

unsupported by previous studies which failed to find consistent patterns (McDonald *et*

al. 2017). To test these predictions, we use the evolutionary assumptions of stage-

structured population models and their mathematical proprieties, as detailed below.

Again, we focus on MPMs, but the same properties also emerge from IPMs (Ellner *et*

al. 2016).

Matrix Population Models have been used numerous times to identify the

evolutionary processes at the population level (*e.g.* Lande 1982, Coulson *et al.* 2006, de

Vries *et al.* 2020). Because the mean values of the matrix elements a_{ij} represent the

mean performance of individuals at each (st-)age in surviving and transmitting their

genes to the next generation, the population growth rate assessed by the MPM (λ) is

assumed to be a proxy for the mean fitness of a population (Lande 1982, Brodie et al.

1995, Coulson et al. 2006, Metcalf and Pavard 2007, Engen et al. 2011). If λ can be

used as a proxy for the mean fitness, each a_{ij} is one of the fitness components and its

contribution to the mean fitness can be quantified with its respective *sensitivity*.

Nevertheless, the sensitivity obtained as the first-order derivative of a_{ij}, assumes a

292 linear relationship between a demographic process a_{ij} and λ, even though this

293 assumption is often violated in natural systems (Caswell 1996). The non-linearity

294 between the contribution of each a_{ij} and λ is adequately verified (and quantified) by

295 using the second-order derivatives of pairs of a_{ij} (Caswell 1996).

296   The assumption that λ_s is approximately the long-term mean fitness in a

297 population in a stochastic environment (*e.g.* Lande 1982, Coulson *et al.* 2006, de Vries

298 *et al*. 2020), allows us to interpret the results from second-order derivative in terms of

299 disruptive and stabilizing selection (Caswell 1996, McCarthy et al. 2008, Shyu and

300 Caswell 2014). Therefore, if the second-order derivative of demographic process a_{ij} is

301 negative – representing a concave function (∩-shaped) - natural selection tends to

302 reduce temporal variation in a_{ij} (Brodie et al. 1995). We can expect a negative second-

303 order derivative for demographic processes a_{ij} with high contribution to fitness and low

304 temporal variance, evidencing buffering as well as an environmental canalization

305 process – a prediction from the DBH (Pfister 1998, Gaillard and Yoccoz 2003). On the

306 other hand, if the second-order derivative is positive (U-shaped), the temporal variance

307 of demographic process should increase, thus evidencing a disruptive selection, as

308 predicted by the DLH (Pasztor *et al*. 2000, Koons *et al*. 2009, 2014).

Figure 2. A unified framework placing life histories along a continuum of demographic buffering *vs.* demographic lability and linking the demographic processes to the underlying key evolutionary processes. Arrows show the flow between each step. Step 1 allocates (rank) each life history into the DB-DL continuum using the $\Sigma E_{a_{ij}}^{S^\mu}$ and $\Sigma E_{a_{ij}}^{S^\sigma}$. This step produces a diagonal line where species are placed according to their responses to environmental stochasticity, from the more buffered on the top left corner, to more labile on the bottom right corner. A population will move along the diagonal continuum, from the top left towards the bottom right corner, if its demographic processes a_{ij} vary more across time – particularly the most contributive a_{ij}. Dots represent 50 hypothetical life histories and colors represent the relative place along the continuum from more buffered (blue) to more labile (red). To perform steps 2 and 3, we must look at each population (each dot) in step 1, separately. Here we represent a hypothetical population from a hypothetical wolf species to assess whether the evolutionary process acting on its vital rates or demographic processes (or a_{ij} for MPMs) correspond to expectations posit by the DBH and DLH. In step 2, we plot the elasticity of each vital rate or demographic process, a_{ij}, *vs.* its coefficient of temporal variation providing a visual guide to evaluate

the trend between temporal variance and its contribution to λ. Note that this step is

common with most approaches that assess DBH and DLH, but, in our framework, it

only serves to visually inspect the relationship. Finally, step 3 quantifies the type of

natural selection driving the temporal variation in each demographic process, by

obtaining the second-order derivatives, represented by different colors – from

stabilizing selection (second-order derivatives < 0) in blue to disruptive selection

(second-order derivatives > 0) in red.

Box 2. Advantages of the unified analytical framework

Numerous limitations in using correlative analyses to assess the DBH and DLH

have been pointed out since it was first proposed by Pfister (1998) (Morris and Doak

2004, Doak et al. 2005, McDonald et al. 2017). While some of these limitations have

been overcome with corrective methods, others remain neglected. Below, we point out

these limitations and how the framework proposed to handle them in contrast to the

correlative analyses – the dominant approach to assess DBH and DLH (Table 1).

Mean-variance scale: Different demographic processes cannot be compared

because they have different magnitudes of mean and variance (Morris and Doak 2004).

For instance, consider two survival rates s_1 ($\bar{x} = 0.6 \pm 0.1$) and s_2 ($\bar{x} = 0.9 \pm 0.1$). Both

present similar variances, yet s_1 varies proportionally more than s_2. Additionally,

variation in s_2 is constrained by the maximum possible survival, which means that in

some time intervals, no individuals died (*survival* = 1.0). A possible solution for such

issues is rescaling variance by the maximum possible variance (Morris and Doak 2004,

Morris *et al.* 2011). However, there is still no consensus on whether this is the best

practice as different rescaling methods provide different results (Bjørkvoll *et al.* 2016,

McDonald *et al.* 2017). Nevertheless, this is not a problem for our framework once vital

rate variance is weighted by its proportional contribution to λ – the elasticity, a

comparable metric between any kinds of demographic processes (Haridas and

Tuljapurkar 2005).

Independent and identically distributed (iid) variables: A common

assumption in correlative analyses is the iid, which assumes that all demographic

processes are independent from each other (Zar 2010). However, this might often not

hold, since many vital rates can be linked by trade-offs (Doak et al. 2005). Thus, the

selection pressure on a given demographic process (*e.g.* promoting variation) might

affect other demographic processes in the opposite direction (Doak et al. 2005, Doak

and Morris 2010). Since the data are not iid, the analytical framework must not be

constrained by this assumption. In turn, our framework computes $\Sigma E_{a_{ij}}^{S^{\mu}}$ and $\Sigma E_{a_{ij}}^{S^{\sigma}}$,

which inherently capture environmental and trade-off related variance. Additionally, the

second-order derivatives (step 3), may also assess the force and direction of natural

selection in trade-offs which is a potential source of correlation between different

demographic processes (cross-correlative second-order derivative, not described here

but see Caswell 1996, 2001, Shyu and Caswell 2014).

Life histories $\times$ demographic processes or patterns $\times$ causality: Correlative

analyses fail to provide a formal link between *processes* (or *causal mechanisms*) and

patterns (Shipley 1999)*,* especially because different demographic processes, which are

under different selective pressures, are analyzed as part of the same pattern (Doak *et al*.

2005). For instance, Doak *et al*. (2005) demonstrate that even if high temporal variation

in most of the demographic processes reduces λ, the variation of one or a few vital rates

might have an opposite (and positive) contribution to λ. Mixing demographic processes

under different evolutionary pressures in simple correlative analyses limits our

understanding of the behavior of each demographic process and might lead to finding

no pattern at all. Indeed, this might be a potential explanation of why most of the

coefficients of correlation are close to zero in one of the most comprehensive uses of

correlation analyses assessing DBH and DLH in plants by McDonald *et al.* (2017).

Equal weight: Correlative analyses (Spearman's correlation) are equally

sensitive to demographic processes with high and low contributions to population

growth rate. However, demographic processes with low contributions to population

growth rate are free to vary under the predictions from the DBH and DLH (Doak *et al*.

2005). In this sense, it is preferable to think the predictions by DBH and DLH as

boundaries, confining the maximum possible temporal variance of the demographic

processes, rather than a trend which they should follow (Fig. 3). By using the step 1 of

our framework, demographic processes with low contributions to λ, even if varying

freely, should have minor contribution to place populations into the DB-DL continuum

(Haridas and Tuljapurkar 2005). This characteristic of $\Sigma E_{a_{ij}}^{S^{\mu}}$ and $\Sigma E_{a_{ij}}^{S^{\sigma}}$ fits conveniently

with the predictions posted by the DBH and DLH.

Table 1. Performance comparison between the framework proposed here and the largely

used approaches to test DBH and DLH (correlative analyses).

	Largely used protocol	New framework proposal	
Limitation \ Approach	correlative analyses	Sum of stochastic elasticity (step 1)	Second-order derivative (steps 2 and 3)
Mean-variance scale	Solved[1,2,3,*]	Accounted for	Not relevant
iid	Neglected[4]	Accounted for	Accounted for
Life history x vital rates	Accounted for but without consensus**	Accounted for, but each step of the present framework has a clear distinction between patterns and processes	
Equal weight	Neglected	Accounted for	Not relevant

*There are multiple ways to solve the scale problem but there is no consensus of

which one is the best ([1]Morris *et al*. 2011, [2]Bjørkvoll *et al*. 2016, [3]McDonald *et*

al. 2017)

[4] Doak *et al.* 2005.

** Two different approaches exist and are assessed using the same methods *i.e.*

multi-species comparative approach and hypothesis testing approach (Box 1).

Figure 3. Theoretical relationship between stochastic elasticity and the coefficient of

temporal variation under the predictions of DBH and DLH. The predictions from DBH

and DLH are depicted in two different ways. Continuous and dashed lines represent a

perfect support for the DBH and DLH, respectively, if we assume demographic

processes should follow specific trends. Blue and red points represent the positions,

where hypothetical vital rates should be placed, considering that natural selection acts

by inducing boundaries to temporal variation under DBH and DLH predictions. Areas

with co-occurrence of blue and red points together, cannot provide insights about the

evolutionary process acting on demographic processes, thus, they represent

uninformative areas. Assuming the DBH and DLH as boundaries to temporal variation,

the right quadrant is expected to be empty.

Demographic buffering in mammals: a case study using the proposed framework

Study system and predictions

We demonstrate the performance of our framework to assess the DBH *vs.* DLH using MPMs from mammal species. Mammals are of special interest here for two reasons: first, mammalian life histories have been well studied (Gillespie 1977, Stearns 1983, Bielby *et al*. 2007, Jones 2011); and second, some of their populations have already been assessed in terms of DBH and DLH, particularly for primates (Morris *et al.* 2008, 2011, Reed and Slade 2012, Rotella *et al*. 2012, Campos *et al*. 2017). Together, the well-studied life histories and previous information about the DBH and DLH in mammals provide the necessary information to make accurate predictions to validate the performance of the proposed framework.

First step predictions – demographic responses in mammal life history: Long-lived species should be characterized by vital rates that are less dependent on environmental conditions (Gaillard and Yoccoz 2003, Morris e*t al*. 2008, 2011). This might be particularly evident for primates, which have intense parental care, high longevities and flexible diets (Jones 2011). These aforementioned traits are expected to increase species' capacity to buffer the negative effects of the environmental conditions. In addition, limited litter size imposed by physiological restrictions limits variation in recruitment even in the bonanza years (Campos *et al.* 2017). Therefore, we predict minimum variation on demographic processes for primates. The capacity to buffer the negative effects of the environmental stochasticity and physiological limitations should vary across the different lineages of mammals. Therefore, at the DL-end, we should expect short-lived species with a highly variable brood size, as is shown in small rodents (de Andreazzi *et al.* 2011) (Fig. 4).

Ultimately, we tested whether the demographic responses of mammals are

related to their longevities and recovery capacities. The relationship between

demographic responses has been suggested by Morris *et al.* (2008) but was not

supported by McDonnald *et al.* (2017). On the other hand, a negative relationship

between buffering and recovery rates has been suggested as an ecological reason why

demographic buffer should evolve, yet this claim remains untested (Ovaskainen and

Meerson 2010, McDonald *et al.* 2017).

Second and third step predictions – patterns of variance and natural selection in

mammal's life history: **Step 2:** As we posited in the predictions for step 1, it is unlikely

that primates should be placed at the DL-end in the DB-DL continuum. Yet, previous

studies have found no support for the idea that the most important vital rates vary less in

primates – a pattern expected to support the DBH (Morris *et al.* 2011). Thus, we do not

expect clear patterns from the relationship between temporal variation of the

demographic processes and their contribution to λ (step 2), particularly for primates

(Fig. 5). **Step 3**: This step is more critical in our framework because only humans

(Caswell 1996) and a species of perennial herb (Shyu and Caswell 2014) had their

demographic processes evaluated in terms of second-order derivatives so far. Yet, as the

DBH and DLH predict, we should expect the most important demographic processes to

be under stabilizing selection to reduce their temporal variation, supporting the DBH, or

to be under disruptive section to increase temporal variation, supporting the DLH (Fig.

5).

Data and analyses

We used Matrix Population Models from 40 out of 139 studies with mammals

available in the COMADRE database v.3.0.0 (Salguero-Gómez *et al.* 2016). These 40

populations encompass 34 species from eight taxonomic orders, and have been included

in our analyses because they provide values of demographic processes (a_{ij}) for three or

more different moments, which allowed us to obtain the stochastic elasticity of each a_{ij}.

Although we are aware that not all possible temporal variation in vital rates may have

been expressed within this time period, we assumed this was enough to provide

sufficient variation for population comparison and to test and showcase our framework.

Fortunately, several long-lived species, characterized by low variation in their vital

rates, were studied for a long time (*e.g.* some primates have been studied for more than

20 years – Morris *et al.* 2011). We also removed the populations where either only

survival or only reproduction rates were reported, because we could not calculate the

stochastic growth rate for these populations. A detailed description of the analyzed data

and their original sources is available in supplementary material (Table S1).

We included *Homo sapiens* in our analyses because it is the only mammal

species, where second-order derivatives have been applied (Caswell 1996), thus making

them ideal for species comparisons. The data for *Homo sapiens* were pooled from 26

modern populations from different cities to construct a variance. Note that in this case

we are not working with true temporal variance but a spatiotemporal variance instead.

A subset of 16 populations, which had their MPMs organized by age (plus *Homo*

sapiens), were used to run steps 2 and 3 of our framework. We selected these

populations because the complexity of life histories is summarized only to two

demographic processes: survival (t) and contribution to recruitment of new individuals

(r), which is equivalent to or approximately equal to the mean reproductive output for

each age, depending on the organization of the matrices (Ebert 1999). The advantage of

using such matrices is that there are only two kinds of demographic processes included,

survival and recruitment, instead of multiple transitions from one stage to another. We

provide the results for all 40 populations in the supplementary material (Fig. S2). As the

non-linear aspect of the contribution of vital rates to individual fitness is often

neglected, we decided to provide an example to explore in detail how this nonlinear

relationship appears in life histories in the DB-DL context (Fig. S3).

To perform the step 1 of our framework, the $\Sigma E_{a_{ij}}^{S^{\sigma}}$ and $\Sigma E_{a_{ij}}^{S^{\mu}}$, we followed

Tuljapurkar *et al.* (2003). We checked if the position of species within the DB-DL

continuum is correlated to longevity (here represented by generation time) and recovery

rate (represented by damping ratio) using simple Spearman's correlations. Generation

time and damping ratio were extracted using the *Popbio* package (Stubben 2007). To

perform step 2 of our framework, we calculated the coefficient of temporal variation of

each demographic process corrected by its maximum variance, as suggested by Morris

et al. (2011) (Box 2), and plotted the variance against the respective deterministic

elasticity, also extracted using the *Popbio* package. All analyses were performed using

R version 3.5.1 (R Core team, 2018).

Finally, to perform the step 3 of our framework, the second-order derivatives

were adapted from *demogR* (Jones 2007) following Caswell (1996) and applied for the

mean MPM. Most contributive demographic processes usually have higher absolute

values of second-order derivative (Carslake *et al.* 2008) as they are expected to be under

stronger pressure of natural selection (Brodie *et al.* 1995, Pfister 1998, Benton and

Grant 1999). On another hand, second-order derivative rarely equals zero for the

demographic processes, even for those less contributive demographic processes

(Carslake *et al.* 2008, McCarthy *et al.* 2008, Shyu and Caswell 2014). However, less

contributive demographic processes are either less important or more prone to have

neglectable curvature (particularly because they are expected to be under less pressure

of the natural selection [Pfister 1998]). As there is no approach to assess whether a

second-order derivative is significantly different from zero or not (see McCarthy *et al.*

2008), we arbitrary rounded the values of second-order derivative for the first decimal

place and considered zero as non-informative.

Results

We ranked 40 populations from 34 species according to the cumulative impact of

variation in demographic processes on λ_s using the first step of our framework ($\Sigma E_{a_{ij}}^{S^{\sigma}}$

and $\Sigma E_{a_{ij}}^{S^{\mu}}$). We show both values in Figure 4, but as the terms $\Sigma E_{a_{ij}}^{S^{\sigma}}$ and $\Sigma E_{a_{ij}}^{S^{\mu}}$ are

complementary and add up to 1, we only discuss the values from of $\Sigma E_{a_{ij}}^{S^{\sigma}}$ for simplicity.

Additional information is provided in the supplementary material (Table S1). The

smallest contribution of variation in vital rates (*i.e.* maximum value of $\Sigma E_{a_{ij}}^{S^{\sigma}}$, note that

$\Sigma E_{a_{ij}}^{S^{\sigma}}$ ranges from 0 to -1), meaning a less environmentally-dependent life history, was

assigned to the Northern muriqui (*Brachyteles hyphoxantus*, Primates) ($\Sigma E_{a_{ij}}^{S^{\sigma}}$ = -0.09 $\times$

$10^{-4} \pm 0.12 \times 10^{-4}$) (mean $\pm$ standard deviation) and the most environmentally-dependent

population was the stoat (*Mustela erminea*, Carnivora) ($\Sigma E_{a_{ij}}^{S^{\sigma}}$ = -0.35 $\pm$ 0.02).

All 14 primate populations were posited at the DB-end of demographic

responses to environmental stochasticity, occupying the upper-left corner of the DB-DL

continuum (Fig. 4). Other species in groups such as Carnivora (N = 10) and

Artiodactyla (N = 5) are spread across the DB-DL continuum (Fig. 4). Our findings

support that life histories with longer generation time are less affected by the

environmental stochasticity, $|\Sigma E_{a_{ij}}^{S^{\sigma}}|$ (Spearman's correlation, ρ = -0.83, P < 0.01), which

means, more buffered. On another hand, life histories more dependent of the

environmental stochasticity have also higher recovery capacities, $|\Sigma E_{a_{ij}}^{S^{\sigma}}|$ (Spearman's

538 correlation, ρ = 0.45, P = 0.01). These results are in concordance with the predictions

539 from step 1 of our framework.

540  As we predicted for the step 2 (graphical inspection), we could not observe a

541 clear negative correlation between the contribution to λ from separate demographic

542 processes and their temporal variation. Particularly for primates, with the lack or minor

543 temporal variation in demographic processes, an almost flat trend was created for most

544 of the analyzed populations, *e.g.* Northen muriqui, and Montain gorilla (*Gorilla*

545 *beringei*) (Fig. 5, S2). Yet, these abovementioned populations were all placed into the

546 extreme DB-end using the first step of our framework ($\Sigma E_{a_{ij}}^{S^{\sigma}}$ and $\Sigma E_{a_{ij}}^{S^{\mu}}$). Such result

547 highlights the importance of a clear definition of the expectations and the importance to

548 choose the right method to assess the aspects of the demographic strategy in the wildlife

549 populations. A full discussion is provided below.

550  By using the second-order derivative (step 3), our framework explicitly assessed

551 the evolutionary forces of natural selection acting on each demographic process.

552 However, no clear pattern arose from this step (Fig. 5, S2). Instead, we could identify

553 some evidence of stabilizing or disruptive selection in almost one half of demographic

554 processes (47% and 23 % respectively), mainly mixed within each life history (Fig S4).

555 For instance, evidence of disruptive and stabilizing selection is present in the most

556 important demographic parameters of White face capuchin monkey (*Cebus capucinus*),

557 the rhesus monkey (*Macaca mulata*), and the tamar wallaby (*Macropus eugenii*)

558 populations (Fig. 5, S4). We also use these populations to exemplify how variations in

559 some of the most important demographic processes determine the long-term mean

560 fitness (λ_s) (Fig. S3). Only a few populations, like humans and Columbian ground

561 squirrel (*Urocitellium colombianus*), presented negatives values of second-order

562 derivatives for all most important demographic processes – a pattern expected by DBH

563 (Fig. 3, 5). Similarly, only a few populations (*e.g.* killer whale [*Orcinus orca*])

564 presented all most important demographic processes under disruptive selection –

565 evidencing DLH (Fig. 5, S4). A general profile on the frequency of what kind of natural

566 selection the demographic processes are under is provided in the supplementary

567 material (S4).

Figure 4. Results of the sum of stochastic elasticities with respect to their variance $\Sigma E_{a_{ij}}^{S^{\sigma}}$ and mean $\Sigma E_{a_{ij}}^{S^{\mu}}$, as suggested by our framework (step 1), ranking the 40 populations from 34 species of mammals from the COMADRE database onto a DB-DL continuum from more buffered (top left corner) to more labile (bottom right corner). Colors represent different taxonomic orders with Primates occupying the top left corner. Primate populations show the highest capacity to buffer environmental stochasticity, as expected.

Figure 5. Results from steps 2 and 3 of the proposed framework (see Fig. 2B,C). The 16 populations presented here are those MPMs built by ages available in the COMADRE database. Different shapes represent different demographic processes (survival and recruitment). As in Figure 2B (step 2), the coefficient of temporal variation of each demographic process is plotted against its deterministic elasticity (see Box 1,2). We complete the graphical information with the value of the second-order derivative (denoted by size) and the direction of natural selection (denoted by color) acting on each demographic process represented by a specific matrix element. Full results are presented in supplementary material (Fig. S2).

Discussion

Our framework allows researchers to detect, quantify, and compare the extent to which

natural populations are demographically buffered or labile in the environment, as well

as the evolutionary processes shaping this capacity to buffer or track the environmental

stochasticity (Fig. 2). In addition, our framework enables the development of an explicit

link between the demographic buffering hypothesis (DBH; Pfister 1998) and the

demographic labile hypothesis (DLH; Koons et al. 2009), and key evolutionary

concepts such as disruptive and stabilizing selection (Fig. 2C). Below, we discuss the

performance and contribution provided by each step of our framework to unveil the

patterns and processes behind the DB-DL perspective.

Framework performance

By using the sum of stochastic elasticity of the stochastic population growth rate

with respect of the mean ($\Sigma E_{a_{ij}}^{S^{\mu}}$) and variance ($\Sigma E_{a_{ij}}^{S^{\sigma}}$, step 1), our framework supports

the existing link between species' buffering capacity, their capacity to recover after

disturbances, and their longevity (Boyce *et al.* 2006, Morris *et al.* 2008, McDonald *et*

al. 2017). Particularly for the recovery rate, our results represent the first empirical

evidence between species' recovery capacity and the DB-DL perspective. This

particular result signifies an important gain, as both, the DBH and DLH, are mainly

considered in terms of species' capacity to track environmental conditions - particularly

for the DL-end's species (*e.g.* McDonald *et al.* 2017, Hilde *et al.* 2020). In terms of the

relationship between longevity and the DB-DL continuum, our findings support recent

work. Morris *et al.* (2008), also using $\Sigma E_{a_{ij}}^{S^{\sigma}}$ and $\Sigma E_{a_{ij}}^{S^{\mu}}$, found that longevity determines a

species' position along the DB-DL continuum for 36 species of plants, mammals, and

other vertebrates. Together, the emerging evidence is starting to indicate that life-history

traits should drive the species' position in the DB-DL continuum regardless of the

taxonomic group. It is worth pointing out that other attempts, specifically those using

correlative analyses (*e.g.* Morris *et al*. 2011, Reed and Slade 2012, McDonald *et al*.

2017)), did not reach the same conclusions.

The contrasting results of the correlative analyses versus the $\Sigma E_{a_{ij}}^{S^\sigma}$ and $\Sigma E_{a_{ij}}^{S^\mu}$,

highlight the importance of carefully defining what should be expected for buffered or

labile life histories. By clearly stating what we should expect based on the DBH and

DLH, our framework using the $\Sigma E_{a_{ij}}^{S^\sigma}$ and $\Sigma E_{a_{ij}}^{S^\mu}$ (step 1 of our framework; Fig. 2), ranks

species on a DB-DL continuum even when all demographic processes present low (or

minimum) temporal variance. Temporal variance in vital rates is a necessary component

to assess the DBH and DLH predictions and to rank species by using correlative

analyses. Yet, several populations – particularly from highly buffered species – should

present low (or minimal) temporal variance (*e.g.* Morris *et al.* 2011). A flat trend

between temporal variation of vital rates and their contributions to λ might arise if all

vital rates present similar and low temporal variation, which cannot be interpreted in

terms of DBH and DLH using correlative analyses. Examples of flat or quasi-flat trends

appear several times when visually representing the temporal variation of demographic

processes and their contributions to λ. Such a pattern is not surprising for primates,

where prolonged parental care and limited brood-size keep juvenile survival and adult

reproductive rates constant in good or bad years (Morris *et al*. 2011; Campos *et al*.

2017, Salguero-gómez 2021).

Finally, the second-order derivative, step 3 in our framework (Fig. 2), reveals

that the role of natural selection shaping temporal variation in vital rates is much more

complex than expected by DBH and DLH. Indeed, we found that demographic

processes within a population are often under a mix of stabilizing and disruptive

selection - a pattern already suggested by Doak *et al.* (2005). Two findings here deserve

further attention: firstly, stabilizing selection is not universal but is still dominant in the

most important demographic processes. In fact, evidence for demographic processes

under the influence of stabilizing selection might reach up to 46%, if we consider only

those 25% of most contributing demographic processes in the life-histories. Such a

result might help explain why evidence for the DBH has been dominant in the literature

(see McDonald *et al.* 2017; Hilde *et al.* 2020). Secondly, we report evidence of

beneficial variation - positive values of second-order derivative (U-shaped), even for

most important vital rates with low temporal variation. Beneficial temporal variation in

most important vital rates was suggested by Koons *et al.* (2009), particularly for short-

lived organisms. Here, we show that natural selection might also increase the variance

in demographic parameters in long-lived organisms. We found such pattern particularly

for primates where most important demographic processes are the survival of oldest

individuals (*e.g.* White face capuchin monkey [*C. capucinus*]). This result is probably a

consequence of natural selection promoting longer lifespans in some populations, like

those of primates, since we found positive second-order derivatives for adult survival of

primates (Caswell 1996, Jones 2011, Jones and Tuljapurkar 2015, Colchero *et al.* 2016).

General insights: challenges and future directions

Investigating what mechanisms drive species' positioning along the DB-DL

continuum is a promising avenue of further investigation. Our results from step 1, $\Sigma E_{a_{ij}}^{S^\sigma}$

and $\Sigma E_{a_{ij}}^{S^\mu}$, show that demographic buffering is a stronger pattern in Primates than

Carnivora and Artiodactyla, for example. This pattern in some but not other taxa

challenges the view that demographic strategies may not be predicted via phylogenetic

distance as suggested by Morris *et al*. (2008). While we did not test this hypothesis

explicitly here, however, our trait-based prediction to species' buffering capacity

supports the idea that more phylogenetically related species (or species with more

similar traits) should present more similar buffering capacities. Indeed, the clustering of

Primates but not Carnivora – a widespread taxa in terms of longevity and reproduction

rate (van de Kerk *et al*. 2013) – supports the idea that similar species should be placed

closer on the DB-DL continuum. We suggest further research should focus on exploring

how the DB-DL continuum manifests across the Tree of Life and the role that morpho-

physiological and life-history traits play in determining the demographic strategies of

these species.

The insights provided by the second-order derivatives used in our framework

(step 3) represent an important step to integrate ecological and evolutionary

implications of selection acting on vital rate temporal variance. The contribution

provided by the second-order derivatives was critical to link the evolutionary

predictions posited by the DBH and DLH (Pfister 1998, Gaillard and Yoccoz 2003,

Hilde *et al*. 2020). Moreover, the second-derivatives provide the necessary information

when simple linear relationship fails to predict the effect of temporal variation

(perturbation) of a given demographic processes to the mean fitness (Carslake *et al*.

2008, Shyu and Caswell 2014). We suggest the second-order derivatives to become

more integrated into the ecological and evolutionary aspects of the demographic

analyses with the attempt to link trait-based demographic models. For example, Integral

Projection Models (IPMs; Easterling *et al.* 2000) usually use continuous traits (*e.g.* size)

as a proxy to predict survival and fecundity (Ellner *et al.* 2016). As such, the same

approach presented here might be adaptable to assess the evolutionary forces shaping

the temporal variation of continuous traits used to build the IPMs.

Regardless to the potential of second-order derivatives in stochastic

demography, their application is still rather limited (but see Caswell 1996, Carslake *et*

al. 2008, Shyu and Caswell 2014, 2016), and has for the most part focused on

methodological aspects (*e.g.* Caswell 1996, Shyu and Caswell 2014). However, despite

the progress in the field since the implementation of the second-order derivatives for

structured population models led by Caswell (1996), this method had not yet been

applied interspecifically for comparative purposes. Thus, in order to make such

comparison, we designed a specific profile of each life history based on the

characteristics of the second-order derivative values to make them comparable. Despite

the success of our approach to perform descriptive analyses, this approach offer limited

statistical inference and remains a challenge for the future.

Conclusions

Both, the Demographic Buffering Hypothesis (DBH) and the Demographic Lability

Hypothesis (DLH) have become important pieces of an ecological puzzle to predict

population responses to environmental stochasticity, climate change, and direct

anthropogenic disturbances (Pfister 1998, Boyce *et al*. 2006, McDonald *et al*. 2017,

Vázquez *et al*. 2017). Yet, the ecological and evolutionary facets of the DBH and DLH

remain tangled in different perspectives that could not have been properly assessed by

correlative analyses, as done to date. As a consequence, the literature exploring these

two hypotheses is full of contradiction (as reviewed in Hilde et al. 2020). Even

comparing small groups of closely related species (*e.g.* primates [Morris *et al*. 2011,

Campos *et al*. 2017], rodents [Reed and Slade 2012], or herbs [Jäkäläniemi *et al*. 2013])

the results can be contradictory. The framework we propose here disentangles the

ecological and evolutionary facets of population responses to stochastic environments,

and provides the necessary tools to quantify and link them through two steps (sum of

stochastic elasticity of the stochastic population growth rate with respect of the mean

($\Sigma E^{S^{\mu}}_{a_{ij}}$) and variance ($\Sigma E^{S^{\sigma}}_{a_{ij}}$) and the second-order derivative), in addition to the usual

approach to assess the DBH and DLH (Pfister 1998, Morris and Doak 2004, McDonald

et al. 2017).

Applied for 40 populations of different species, our framework helps to solidify

classic and more recent avenues of research. Particularly, we highlight the capacity to

link the DB-DL perspective with life-history traits (*e.g.* generation time), recovery rates

and the possibility to link such results to the recent emphasis in linking functional traits

and demographic performance (*e.g.* Adler *et al.* 2014, Salguero-Gómez *et al.* 2018).

Thus, future investigations should benefit from this framework to explore, for example,

how body mass and behavioral traits correlate with different demographic strategies in

environmentally stochastic conditions. Furthermore, the usage of second-order

derivatives in the DB-DL perspective represents a promising research avenue to explore

the role natural selection to shape the temporal variation of vital rates in different taxa.

Acknowledgements

This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de

Nível Superior - Brasil (CAPES) - Finance Code 001. We are grateful for Samuel

Gascoigne for providing valuable comments to improve the science and English of the

manuscript. RS-G was supported by a NERC Independent Research Fellowship

(NE/M018458/1). MK thanks Prociência-UERJ (#363861) and Ad Futura - Public

Scholarship, Development, Disability and Maintenance Fund of the Republic of
Slovenia (#11013-16/2018) for their support.
737

References

Adler, P. B. et al. 2014. Functional traits explain variation in plant life history strategies.
- Proc. Natl. Acad. Sci. 111: 740–745.

Benton, T. G. and Grant, A. 1999. Elasticity analysis as an important tool in
evolutionary and population ecology. - Trends Ecol. Evol. 14: 467–471.

Bielby, J. et al. 2007. The Fast☐Slow Continuum in Mammalian Life History: An
Empirical Reevaluation. - Am. Nat. 169: 748–757.

Bjørkvoll, E. et al. 2016. Demographic buffering of life histories? Implications of the
choice of measurement scale. - Ecology 97: 40–47.

Boyce, M. S. et al. 2006. Demography in an increasingly variable world. - Trends Ecol.
Evol. 21: 141–148.

Brodie, E. D. et al. 1995. Visualizing and quantifying natural selection. - Trends Ecol.
Evol. 10: 313–318.

Burns, J. H. et al. 2010. Empirical tests of life-history evolution theory using
phylogenetic analysis of plant demography. - J. Ecol. 98: 334–344.

Campos, F. A. et al. 2017. Does climate variability influence the demography of wild
primates? Evidence from long-term life-history data in seven species. - Glob.
Chang. Biol. 23: 4907–4921.

Carslake, D. et al. 2008. Nonlinearity in eigenvalue-perturbation curves of simulated
population projection matrices. - Theor. Popul. Biol. 73: 498–505.

Caswell, H. 1996. Second derivatives of population growth rate: Calculation and
applications. - Ecology 77: 870–879.

Caswell, H. 2001. Matrix population models: Construction, analysis, and interpretation.

Charlesworth, B. 1994. Evolution in age-structured populations. - Cambridge University

Press.

Colchero, F. et al. 2016. The emergence of longevous populations. - Proc. Natl. Acad.

Sci. U. S. A. 113: E7681–E7690.

Coulson, T. et al. 2006. Putting evolutionary biology back in the ecological theatre: A

demographic framework mapping genes to communities. - Evol. Ecol. Res. 8:

1155–1171.

de Andreazzi, C. S. et al. 2011. Population ecology of small rodents and marsupials in a

semi-deciduous tropical forest of the southeast Pantanal, Brazil. - Zoologia 28:

762–770.

de Kroon, H. et al. 1986. Elasticity: The Relative Contribution of Demographic

Parameters to Population Growth Rate. - Ecology 67: 1427–1431.

de Vries, C. et al. 2020. A matrix model for density-dependent selection in stage-

classified populations, with application to pesticide resistance in Tribolium. - Ecol.

Modell. 416: 108875.

Debat, V. and David, P. 2001. Mapping phenotype: canalization, plasticity and

developmental stability. - Trends Ecol. Evol. 16: 555–561.

Doak, D. F. and Morris, W. F. 2010. Demographic compensation and tipping points in

climate-induced range shifts. - Nature 467: 959–962.

Doak, D. F. et al. 2005. Correctly estimating how environmental stochasticity

influences fitness and population growth. - Am. Nat. 166: E14–E21.

Easterling, M. R. et al. 2000. Size-Specific Sensitivity: Applying a New Structured

Population Model. - Ecology 81: 694–708.

Ebert, T. A. 1999. Populations Methods in Demography. - Methods &Demography in

press.

Edic, M. N. et al. 2020. Heritable variation in the timing of emergence from

hibernation. - Evol. Ecol. in press.

Ellner, S. P. et al. 2016. Data-driven Modelling of Structured Populations. in press.

Engen, S. et al. 2011. Evolution of a plastic quantitative trait in an age-structured

population in a fluctuating environment. - Evolution (N. Y). 65: 2893–2906.

Fuller, A. et al. 2010. Physiological mechanisms in coping with climate change. -

Physiol. Biochem. Zool. 83: 713–720.

Gaillard, J. M. and Yoccoz, N. G. 2003. Temporal variation in survival of mammals: A

case of environmental canalization? - Ecology 84: 3294–3306.

Gillespie, J. H. 1977. Natural Selection for Variances in Offspring Numbers: A New

Evolutionary Principle. - Am. Nat. 111: 1010–1014.

Griffith, A. B. 2017. Perturbation approaches for integral projection models. - Oikos

126: 1675–1686.

Haridas, C. V. and Tuljapurkar, S. 2005. Elasticities in Variable Environments:

Properties and Implications. - Am. Nat. 166: 481–495.

Healy, K. et al. 2019. Animal life history is shaped by the pace of life and the

distribution of age-specific mortality and reproduction. - Nat. Ecol. Evol. 3: 1217–

1224.

Hilde, C. H. et al. 2020. The Demographic Buffering Hypothesis: Evidence and

Challenges. - Trends Ecol. Evol. 35: 523–538.

Hunter, P. 2009. Robust yet flexible. In biological systems, resistance to change and

innovation in the light of it go hand in hand. - EMBO Rep. 10: 949–952.

Jäkäläniemi, A. et al. 2013. Variability of important vital rates challenges the

demographic buffering hypothesis. - Evol. Ecol. 27: 533–545.

Jones, J. H. 2011. Primates and the evolution of long, slow life histories. - Curr. Biol.
21: R708–R717.

Jones, J. H. and Tuljapurkar, S. 2015. Measuring selective constraint on fertility in
human life histories. - Proc. Natl. Acad. Sci. U. S. A. 112: 8982–8986.

Jongejans, E. et al. 2010. Plant populations track rather than buffer climate fluctuations.
- Ecol. Lett. 13: 736–743.

Koons, D. N. et al. 2009. Is life-history buffering or lability adaptive in stochastic
environments? - Oikos 118: 972–980.

Koons, D. N. et al. 2014. Drivers of waterfowl population dynamics: From teal to
swans. - Wildfowl: 169–191.

Lande, R. 1982. A Quantitative Genetic Theory of Life History Evolution. - Ecology
63: 607–615.

McCarthy, D. et al. 2008. On second order sensitivity for stage-based population
projection matrix models. - Theor. Popul. Biol. 74: 68–73.

McDonald, J. L. et al. 2017. Divergent demographic strategies of plants in variable
environments. - Nat. Ecol. Evol. in press.

Meiri, S. et al. 2020. The global diversity and distribution of lizard clutch sizes. - Glob.
Ecol. Biogeogr.: 1515–1530.

Metcalf, C. J. E. and Pavard, S. 2007. Why evolutionary biologists should be
demographers. - Trends Ecol. Evol. 22: 205–212.

Morris, W. F. and Doak, D. F. 2002. Quantitative conservation biology: theory and
practice of population viability analysis. - In: Quantitative Conservation Biology:
Theory and Practice of Population Viability Analysis. in press.

Morris, W. F. and Doak, D. F. 2004. Buffering of Life Histories against Environmental
Stochasticity: Accounting for a Spurious Correlation between the Variabilities of

Vital Rates and Their Contributions to Fitness. - Am. Nat. 163: 579–590.

Morris, W. F. et al. 2008. Longevity can buffer plant and animal populations against

changing climatic variability. - Ecology 89: 19–25.

Morris, W. F. et al. 2011. Low Demographic Variability in Wild Primate Populations:

Fitness Impacts of Variation, Covariation, and Serial Correlation in Vital Rates. -

Am. Nat. 177: E14–E28.

Ovaskainen, O. and Meerson, B. 2010. Stochastic models of population extinction. -

Trends Ecol. Evol. 25: 643–652.

Pasztor, L. et al. 2000. Jensen's inequality and optimal life history strategies in

stochastic environments [3]. - Trends Ecol. Evol. 15: 117–118.

Pfister, C. A. 1998. Patterns of variance in stage-structured populations: Evolutionary

predictions and ecological implications. - Proc. Natl. Acad. Sci. U. S. A. 95: 213–

218.

Reed, A. W. and Slade, N. A. 2012. Buffering and plasticity in vital rates of oldfield

rodents. - J. Anim. Ecol. 81: 953–959.

Rodríguez-Caro, R. C. et al. 2020. The demographic buffering strategy has a threshold

of effectiveness to increases in environmental stochasticity. - bioRxiv: 1–41.

Rotella, J. J. et al. 2012. Evaluating the demographic buffering hypothesis with vital

rates estimated for Weddell seals from 30years of mark-recapture data. - J. Anim.

Ecol. 81: 162–173.

Salguero-gómez, R. 2021. The fast-slow continuum is not the end-game of life history

evolution, human or otherwise. - bioRxiv: 2021.03.09.434595.

Salguero-Gómez, R. et al. 2016. COMADRE: A global data base of animal

demography. - J. Anim. Ecol. 85: 371–384.

Salguero-Gómez, R. et al. 2018. Delivering the promises of trait-based approaches to

the needs of demographic approaches, and vice versa. - Funct. Ecol. 32: 1424–

1435.

Shipley, B. 1999. Testing Causal Explanations in Organismal Biology: Causation,

Correlation and Structural Equation Modelling. - Oikos 86: 374.

Shyu, E. and Caswell, H. 2014. Calculating second derivatives of population growth

rates for ecology and evolution. - Methods Ecol. Evol. 5: 473–482.

Shyu, E. and Caswell, H. 2016. A demographic model for sex ratio evolution and the

effects of sex-biased offspring costs. - Ecol. Evol. 6: 1470–1492.

Silvertown, J. et al. 1993. Comparative Plant Demography--Relative Importance of

Life-Cycle Components to the Finite Rate of Increase in Woody and Herbaceous

Perennials. - J. Ecol. 81: 465.

Stearns, S. C. 1983. The Influence of Size and Phylogeny on Patterns of Covariation

among Life-History Traits in the Mammals. - Oikos 41: 173.

Tuljapurkar, S. et al. 2003. The Many Growth Rates and Elasticities of Populations in

Random Environments. - Am. Nat. 162: 489–502.

van de Kerk, M. et al. 2013. Carnivora Population Dynamics Are as Slow and as Fast as

Those of Other Mammals: Implications for Their Conservation. - PLoS One in

press.

Vázquez, D. P. et al. 2017. Ecological and evolutionary impacts of changing climatic

variability. - Biol. Rev. 92: 22–42.

Zar, J. H. 2010. Biostatistical analysis, 5th.

883   **Appendix 1 – Data available in COMADRE Version 2.0.1 and results from step 1.**

884   Table S1. present the metadata used in step 1 of our framework and the respective results presented in the paper. The first four columns represent the information from where Matrix Populations Models (MPMs) were

885   extract exactly as presented in COMADRE 2.0.1. Column titles differ from the database as "SpeciesAuthorComadre" is equivalent to "SpeciesAuthor" and "SpeciesName" is equivalent to "SpeciesAccepted" in

886   COMADRE 2.0.1. The remaining columns present the results of step 1, where we present the raw values of $\Sigma E^{S^\mu}_{a_{ij}}$ and $\Sigma E^{S^\sigma}_{a_{ij}}$, their respective standard deviation, population growth rate λ, number of available matrices

887   (# matrices), generation time and damping ratio for each population. For ByAge, assign "TRUE" for MPMs built by age or "FALSE" if otherwise.

SpeciesAuthorComadre	SpeciesName	CommonName	Order	$\Sigma E^{S^\mu}_{a_{ij}}$	$\Sigma E^{S^\mu}_{a_{ij}}$ (sd)	$\Sigma E^{S^\sigma}_{a_{ij}}$	$\Sigma E^{S^\sigma}_{a_{ij}}$ (sd)	# matrices	λ	Generation time	Damping ratio	ByAge
Homo_sapiens_subsp._sapiens	*Homo sapiens sapiens*	Human	Primates	1.003	0.003	1.003	0.004	13.000	1.064	5.499	NA	TRUE
Alces_alces	*Alces alces*	Moose	Artiodactyla	1.001	0.001	1.001	0.001	13.000	1.205	10.628	2.168	TRUE
Antechinus_agilis	*Antechinus agilis*	Agile antechinus	Dasyuromorphia	1.111	0.111	1.111	0.011	2.000	0.931	2.153	NA	FALSE
Brachyteles_hypoxanthus	*Brachyteles hypoxanthus*	Northern muriqui	Primates	1.000	0.000	1.000	0.000	12.000	1.051	37.205	1.244	TRUE
Callospermophilus_lateralis	*Callospermophilus lateralis*	Golden-mantled ground squirrel	Rodentia	1.054	0.054	1.054	0.055	9.000	2.052	NA	3.955	TRUE
Cebus_capucinus	*Cebus capucinus*	White faced capuchin monkey	Primates	1.000	0.000	1.000	0.000	11.000	1.021	29.363	1.391	TRUE
Cercopithecus_mitis	*Cercopithecus mitis*	Blue monkey	Primates	1.000	0.000	1.000	0.000	14.000	1.036	21.452	1.240	TRUE
Eumetopias_jubatus	*Eumetopias jubatus*	Northern sea lion; Steller sea lion	Carnivora	1.005	0.005	1.005	0.002	2.000	0.904	9.645	NA	FALSE
Felis_catus	*Felis catus*	Feral cat	Carnivora	1.136	0.136	1.136	0.012	1.000	1.948	5.476	5.135	FALSE
Gorilla_beringei	*Gorilla beringei*	Mountain gorilla	Primates	1.000	0.000	1.000	0.000	21.000	1.027	38.817	1.239	TRUE
Hippocamelus_bisulcus	*Hippocamelus bisulcus*	Huemul deer	Artiodactyla	1.002	0.002	1.002	0.001	1.000	0.996	10.452	2.928	FALSE
Lepus_americanus	*Lepus americanus*	Snowshoe hare	Lagomorpha	1.294	0.294	1.294	0.165	2.000	0.812	2.290	NA	FALSE
Lycaon_pictus	*Lycaon pictus*	African wild dog	Carnivora	1.100	0.100	1.100	0.008	1.000	1.500	4.149	2.163	FALSE
Macaca_mulatta_3	*Macaca mulatta*	Rhesus macaque	Primates	1.000	0.000	1.000	0.001	12.000	1.127	14.749	1.616	TRUE
Macropus_eugenii	*Macropus eugenii*	Tammar wallaby	Diprotodontia	1.013	0.013	1.013	0.012	7.000	0.981	4.872	10.930	TRUE
Marmota_flaviventris_2	*Marmota flaviventris*	Yellow-bellied marmot	Rodentia	1.007	0.007	1.007	0.006	4.000	0.890	4.210	1.489	FALSE
Marmota_flaviventris_3	*Marmota flaviventris*	Yellow-bellied marmot	Rodentia	1.008	0.008	1.008	0.005	4.000	0.921	5.017	3.575	FALSE
Microtus_oeconomus	*Microtus oeconomus*	Root vole	Rodentia	1.000	0.000	1.000	0.001	14.000	1.028	30.588	4.489	TRUE
Mustela_erminea	*Mustela erminea*	Stoat	Carnivora	1.334	0.334	1.334	0.117	2.000	1.258	5.803	11.201	FALSE
Orcinus_orca_2	*Orcinus orca*	Killer whale	Cetacea	1.001	0.001	1.001	0.001	24.000	0.999	28.745	NA	TRUE
Ovis_aries_2	*Ovis aries*	Soay sheep	Artiodactyla	1.033	0.033	1.033	0.020	3.000	1.099	8.036	NA	TRUE
Pan_troglodytes_subsp._schweinfurthii	*Pan troglodytes*	Eastern chimpanzee	Primates	1.000	0.000	1.000	0.001	22.000	0.982	37.652	1.132	TRUE
Papio_cynocephalus	*Papio cynocephalus*	Olive baboon	Primates	1.000	0.000	1.000	0.000	19.000	1.054	21.860	1.289	TRUE
Peromyscus_maniculatus_2	*Peromyscus maniculatus*	Deer mouse	Rodentia	1.010	0.010	1.010	0.005	2.000	1.107	5.596	1.780	FALSE
Phocarctos_hookeri	*Phocarctos hookeri*	New Zealand sea lion	Carnivora	1.005	0.005	1.005	0.003	8.000	1.023	9.681	1.256	FALSE
Propithecus_verreauxi	*Propithecus verreauxi*	Verreaux's sifaka	Primates	1.000	0.000	1.000	0.000	12.000	0.986	21.963	1.376	TRUE
Puma_concolor_8	*Puma concolor*	Cougar	Carnivora	NA	NA	NA	NA	10.000	1.115	4.806	NA	FALSE
Rattus_fuscipes	*Rattus fuscipes*	Bush rat	Rodentia	1.246	0.246	1.246	0.029	2.000	1.305	2.023	NA	FALSE
Spermophilus_armatus	*Urocitellus armatus*	Uinta ground squirrel	Rodentia	1.016	0.016	1.016	0.011	4.000	1.125	3.755	8.020	FALSE
Spermophilus_armatus_2	*Urocitellus armatus*	Uinta ground squirrel	Rodentia	1.017	0.017	1.017	0.010	3.000	1.095	4.838	2.767	FALSE
Spermophilus_columbianus	*Urocitellus columbianus*	Columbian ground squirrel	Rodentia	1.036	0.036	1.036	0.025	3.000	1.009	4.508	2.510	FALSE
Spermophilus_columbianus_3	*Urocitellus columbianus*	Columbian ground squirrel	Rodentia	1.003	0.003	1.003	0.006	3.000	1.200	4.524	NA	TRUE
Ursus_americanus_subsp._floridanus	*Ursus americanus*	Florida black bear	Carnivora	1.003	0.003	1.003	0.003	2.000	1.020	9.591	1.890	FALSE

Ursus_arctos_subsp._horribilis_5	*Ursus arctos*	Grizzly bear	Carnivora	1.001	0.001	1.001	0.001	4.000	1.026	11.558	1.158	FALSE
Ursus_maritimus_2	*Ursus maritimus*	Polar bear	Carnivora	1.019	0.019	1.019	0.007	2.000	0.941	18.491	1.552	TRUE
Brachyteles_hypoxanthus_2	*Brachyteles hypoxanthus*	Northern muriqui	Primates	1.000	0.000	1.000	0.000	12.000	1.111	21.278	3.443	TRUE
Cebus_capucinus_2	*Cebus capucinus*	WhiteNAfaced capuchin monkey	Primates	1.000	0.000	1.000	0.000	11.000	1.059	18.977	3.636	TRUE
Chlorocebus_aethiops_2	*Chlorocebus aethiops*	Vervet	Primates	1.075	0.075	1.075	0.087	5.000	1.187	5.694	3.508	FALSE
Erythrocebus_patas	*Erythrocebus patas*	Patas monkey	Primates	1.051	0.051	1.051	0.038	5.000	1.128	3.499	2.291	FALSE
Gorilla_beringei_subsp._beringei	*Gorilla beringei*	Mountain gorilla	Primates	1.000	0.000	1.000	0.000	21.000	1.053	24.536	4.161	TRUE

Appendix 2

Figure S2. Results from steps 2 and 3 for all populations analyzed (see fig. 2B,C) in this study. Differently to figure 5, which only uses

MPMs built by age with two kinds of transitions a_{ij} – survival and recruitment, figure S2 presents MPMs built with all sorts of categories

if they matched the criteria informed in *data and analyses* section. Therefore, instead of simple survival rates, we might disentangle it in

other demographic processes as self-loops (retain in same stage) and jump for next stage (growth) are depicted in "stage transition". Here

we opted to simply inform what is a transition from a stage to another and what is contribution to recruitment. Moreover, differently to

figure 5, common species names are paired with their respective identification in the "SpeciesAuthor" column in COMADRE database.

Please note that the important results present in this figure have been covered by the discussion of results for figure 5.

Appendix 3

Figure S3. Example of nonlinear contribution to long-term mean fitness (λ) by two demographic processes in three populations analyzed in figure 5. The x-axis shows multiple simulated values of recruitment and survival with their respective contribution to long-term mean fitness (λ) on the y-axis. Vertical dashed lines show the recruitment and survival values from the mean MPM, as well as the corresponding value of λ (horizontal dashed line).

Figure S4. Most of life histories present mix of stabilizing and disruptive selection in their most important demographic processes and its

proportion are presented for all life histories analyzed. From right to left, we present the proportion of disruptive and stabilizing selection

acting on all demographic processes in a life-history (all non-zero entries in the MPM), followed by the proportion of each kind of natural

selection for those 50% and 25% of most contributive demographic processes. At the end of each column the overall result including all

life-histories is presented. Each life-history analyzed is informed at the left by species common name followed by its respective

identification in the "SpeciesAuthor" column in COMADRE database.

916