1

5   *Title:* **Cortico-Fugal Regulation of Predictive Coding.**

7   *Authors:* Alexandria M.H. Lesicko[1], Christopher F. Angeloni[2], Jennifer M. Blackwell[3], Mariella De
8   Biasi[4,5,6], Maria N. Geffen[1,6,7]
9

Affiliations:

1. Department of Otorhinolaryngology, University of Pennsylvania, Philadelphia, United States
2. Department of Psychology, University of Pennsylvania, Philadelphia, United States
3. Department of Neurobiology and Behavior, Stony Brook University, Stony Brook, New York,
United States
4. Department of Psychiatry, University of Pennsylvania, Philadelphia, United States
5. Department of Systems Pharmacology and Experimental Therapeutics, University of
Pennsylvania, Philadelphia, United States
6. Department of Neuroscience, University of Pennsylvania, Philadelphia, United States
7. Department of Neurology, University of Pennsylvania, Philadelphia, United States
20

22 *Corresponding author:*   Dr. Maria N. Geffen
23   Stemmler Hall G10
24   3450 Hamilton Walk
25   Philadelphia, PA 19104
26   (215)898-0872
27   mgeffen@pennmedicine.upenn.edu
28

29 *Keywords:* prediction error, hierarchical predictive coding, inferior colliculus, auditory cortex,

30 cortico-collicular, stimulus specific adaptation, deviance detection

ABSTRACT

Sensory systems must account for both contextual factors and prior experience to adaptively engage with the dynamic external environment. In the central auditory system, neurons modulate their responses to sounds based on statistical context. These response modulations can be understood through a hierarchical predictive coding lens: responses to repeated stimuli are progressively decreased, in a process known as repetition suppression, whereas unexpected stimuli produce a prediction error signal. Prediction error incrementally increases along the auditory hierarchy from the inferior colliculus (IC) to the auditory cortex (AC), suggesting that these regions may engage in hierarchical predictive coding. A potential substrate for top-down predictive cues is the massive set of descending projections from the auditory cortex to subcortical structures, although the role of this system in predictive processing has never been directly assessed. We tested the effect of optogenetic inactivation of the auditory cortico-collicular feedback in awake mice on responses of IC neurons to stimuli designed to test prediction error and repetition suppression. Inactivation of the cortico-collicular pathway led to a decrease in prediction error in IC. Repetition suppression was unaffected by cortico-collicular inactivation, suggesting that this metric may reflect fatigue of bottom-up sensory inputs rather than predictive processing. We also discovered populations of IC neurons that exhibit repetition enhancement, a sequential increase in firing with stimulus repetition. Cortico-collicular inactivation led to a decrease in repetition enhancement in the central nucleus of IC, suggesting that it is a top-down phenomenon. Negative prediction error, a stronger response to a tone in a predictable rather than unpredictable sequence, was suppressed in shell IC units during cortico-collicular inactivation. These changes in predictive coding metrics arose from bidirectional modulations in the response to the standard and deviant contexts, such that neurons in IC responded more similarly to each context in the absence of cortical input. We also investigated how these metrics compare between the anesthetized and awake states by recording from the same neurons under both conditions. We found that metrics of predictive coding and deviance detection differ depending on the anesthetic state of the animal, with negative prediction error emerging in the central IC and repetition enhancement and prediction error being more prevalent in the absence of anesthesia. Overall, our results demonstrate that the auditory cortex provides cues about the statistical context of sound to subcortical brain regions via direct feedback, regulating processing of both prediction and repetition.

INTRODUCTION

Sensory systems differentially encode environmental stimuli depending on the context in which they are encountered (De Franceschi & Barkat, 2020; Herrmann et al., 2015; Jaramillo et al., 2014; Pakan et al., 2016; Takesian et al., 2018; Zhai et al., 2020). The same physical stimulus can elicit distinct neuronal responses depending on whether it is predictable or unexpected in a given sensory stream (Weissbart et al., 2020; Yaron et al., 2012). One framework for understanding this dynamic sensory capability is hierarchical predictive coding, which suggests that neuronal networks form predictions about incoming stimuli based on the statistics of prior experience (Friston & Kiebel, 2009). These predictions are generated at higher levels of the sensory hierarchy and broadcast to lower stations to minimize processing of redundant input and maximize coding efficiency (Friston, 2009; Friston & Kiebel, 2009). Any mismatch between predictions and representations of sensory input is coded in a neuronal response known as a "prediction error", which is further propagated up the sensory hierarchy, ultimately allowing for the formation of updated predictions (Friston & Kiebel, 2009; Shipp, 2016). Multiple sensory modalities exhibit hierarchical predictive coding, including the motor, visual, and auditory systems (Okada et al., 2018; Parras et al., 2017; Rao & Ballard, 1999; Rauss et al., 2011; Schellekens et al., 2016; Shipp et al., 2013).

Neurons in select regions of the central auditory system are sensitive to statistical context, responding more strongly to a tone when it is presented rarely (a "deviant") than when it is commonplace (a "standard") (Ulanovsky et al., 2003). This phenomenon, known as stimulus specific adaptation (SSA), is prevalent in the auditory cortex (Natan et al., 2015; Ulanovsky et al., 2003). Weaker SSA is present in regions peripheral to the AC, including the auditory midbrain, or inferior colliculus (IC), and the auditory thalamus, or medial geniculate body (MGB) (Anderson et al., 2009; Antunes et al., 2010; Duque & Malmierca, 2015; Malmierca et al., 2009; Taaseh et al., 2011; Ulanovsky et al., 2003). Subdivisions in IC and MGB that receive descending projections from AC exhibit relatively higher SSA levels than their lemniscal counterparts (Antunes et al., 2010; Duque et al., 2012), suggesting that SSA may be generated de novo in AC and subsequently broadcast to subcortical structures via cortico-fugal projections (Nelken & Ulanovsky, 2007). Silencing of AC through cooling has been shown to modulate, but not abolish, SSA in IC and MGB of anesthetized rats (Anderson & Malmierca, 2013; Antunes & Malmierca, 2011). However, it remains unknown whether these modulations in the SSA index with cortical deactivation reflect changes in predictive processing.

Recent studies have implemented additional control tone sequences to further decompose the traditional SSA index into two distinct underlying processes: repetition suppression and prediction

error (Harms et al., 2014; Parras et al., 2017; Ruhnau et al., 2012). Repetition suppression is characterized by a decrease in firing rate to each subsequent presentation of a standard tone (Auksztulewicz & Friston, 2016; Parras et al., 2017). Prediction error is thought to signal the mismatch between the predicted input, based on prior experience with repeated presentations of the standard, and the actual sensory input when a deviant tone is presented (Friston, 2009; Friston & Kiebel, 2009). Whereas repetition suppression is thought to potentially reflect synaptic depression, prediction error has been proposed to underlie true deviance detection (Parras et al., 2017; Taaseh et al., 2011). Prediction error increases along the auditory hierarchy and is more prevalent in regions of IC and MGB that receive cortical feedback (Parras et al., 2017), suggesting that these subcortical regions may engage in hierarchical predictive coding, with AC potentially providing predictive cues to IC and MGB. However, how feedback projections from AC shape predictive processing in subcortical targets has never been directly assessed. In fact, virtually all models of hierarchical predictive coding to date have focused on intra-cortical connections, with the massive system of descending cortico-fugal projections remaining unexplored (Asilador & Llano, 2020; Bastos et al., 2012).

Here, we investigated how inputs from AC to IC, the first station in the auditory system in which prediction error is found, shape metrics associated with predictive coding and deviance detection (Parras et al., 2017). To test this, we optogenetically inactivated cortico-collicular feedback while recording neuronal responses in IC and found that prediction error, negative prediction error, and repetition enhancement in IC are altered in the absence of cortical input. Our results suggest that the cortico-collicular pathway sends cues from AC to IC regarding the statistical context of auditory stimuli.

RESULTS

Experimental design

We used a Cre/FLEX viral injection strategy to selectively express the inhibitory opsin, ArchT, in cortico-collicular neurons of four mice by injecting a retroAAV-Cre-GFP construct into IC and an AAV9-FLEX-ArchT-tdTomato construct into AC (Figure 1A, left). The retroAAV-Cre-GFP construct is transported in a retrograde fashion and expressed in neurons that project to IC (Blackwell et al., 2020). The genes encoded in the AAV9-FLEX-ArchT-tdTomato construct can only be expressed in neurons containing the Cre construct, thereby limiting ArchT expression to neurons in AC that project to IC. In the presence of green light, ArchT, a light-driven outward proton pump, mediates rapid, reversible inactivation of the neurons in which it is expressed (Han et al., 2011).

We implanted cannulas over AC in mice injected with the Cre/FLEX constructs and a 532 nm laser was used to provide green light illumination to the region, allowing for inactivation of cortico-collicular neurons (Figure 1A, right). The mice were head-fixed and a 32-channel probe was lowered into IC to perform awake extracellular recordings (Figure 1A). Auditory stimuli consisted of oddball sequences of two repeated pure tones, presented at a 90:10 standard-to-deviant ratio and half-octave frequency separation (Figure 1B). On a subset of trials, presentations of either the deviant or the last standard prior to the deviant were coupled with activation of the green laser (Figure 1B, right).

Neurons that displayed a significantly higher response to the deviant than the standard were designated as "adapting" neurons, while those that exhibited a significantly higher response to the standard than the deviant were categorized as "facilitating" neurons (Figure 1D). The difference in firing rate to the standard and deviant was quantified with an index of neuronal mismatch (iMM), which is equivalent to the SSA index used in previous studies (Parras et al., 2017).

A cascade stimulus consisting of 10 evenly spaced tones, including the tone pair from the oddball sequence, was presented to further decompose the neuronal mismatch between the responses to the standard and deviant (Figure 1C, 1D). This stimulus is unique in that each tone occurs with the same likelihood as the deviant tone in the oddball stimulus (10%), but it contains no true statistical deviants: each tone has the same likelihood of presentation, and the tone sequence overall follows a regular and predictable pattern (Parras et al., 2017). Therefore, the response to a given tone when it is embedded in the cascade can be compared to the response when it is a deviant in order to isolate prediction error effects (Figure 1C; 1D, top). A neuron exhibits prediction error if it fires more strongly to a tone when it is a deviant than when it is presented in the cascade sequence (Figure 1D, top). Conversely, if a neuron responds more strongly to a tone presented in the cascade sequence than

Figure 1: Experimental design. A) Cre/FLEX dual injections for selective ArchT expression in cortico-collicular neurons. Recordings were performed in IC while inactivation was mediated by a 532 nm laser connected to cannulas implanted over AC. B) Oddball stimuli consisted of pairs of pure tones separated by 0.5 octave with a 90:10 standard-to-deviant ratio. Two sequences were constructed such that each frequency is represented as both the standard and the deviant. C) Cascade sequences consisted of 10 evenly spaced tones separated by 0.5 octaves, with both frequencies from the oddball sequence included in the sequence. Responses to tones in the cascade context were compared to responses in the standard and deviant context to analyze repetition and prediction effects, respectively. D) A positive iMM (top diagram) indicates a stronger response to the deviant than the standard (adaptation), while a negative iMM (bottom diagram) indicates a stronger response to the standard than to the deviant (facilitation). The iMM can be further decomposed into an iPE and an iRS. Positive iPE values represent prediction error and negative values convey negative prediction error. Positive iRS indices indicate repetition suppression, while repetition enhancement is represented by negative values.

when it is a deviant, the neuron encodes negative prediction error (Figure 1D, bottom). This phenomenon is quantified using an index of prediction error (iPE), with positive indices indicating prediction error and negative indices representing negative prediction error (Figure 1D).

The cascade sequence is also free from repetition effects, since adjacent tone presentations never include a tone of the same frequency (Figure 1C). Therefore, the response to a given tone embedded in the cascade sequence can be compared to the response generated when that tone is a standard. The difference in response indicates either repetition suppression (stronger response to the tone in the cascade) (Figure 1D, top) or repetition enhancement (stronger response to the tone as a standard) (Figure 1D, bottom). These contrasting processes are quantified by the index of repetition suppression (iRS), with a positive index indicating repetition suppression and a negative index representing repetition enhancement (Figure 1D).

Cre/FLEX viral injection strategy enables selective inactivation of cortico-collicular neurons

Examination of fixed tissue from injected mice revealed that expression of the retroAAV-Cre-GFP construct was restricted to IC (Figure 1 – Figure Supplement 1A, top left). Somatic expression of tdTomato (indicating the presence of ArchT) was restricted to layer 5 and deep layer 6 of AC, which contain cortico-collicular cell bodies, and was broadly distributed throughout the rostro-caudal extent of the auditory cortex (Figure 1 – Figure Supplement 1A, right) (Bajo et al., 2007; Schofield, 2009; Yudintsev et al., 2019). Axons and terminals labeled with tdTomato were distributed in IC in a manner matching the known projection pattern of this pathway, with dense, "patchy" labeling in shell regions of IC (Figure 1 – Figure Supplement 1A, bottom left) (Herbert et al., 1991; Lesicko et al., 2016; Saldaña et al., 1996; Torii et al., 2013). These data confirm that our viral injection strategy leads to selective transfection of cortico-collicular neurons.

Extracellular recordings in AC of injected mice revealed a reduction in firing rate during the duration of the laser stimulus in several neurons (Figure 1 – Figure Supplement 1B, 2C). In these putative cortico-collicular neurons, laser-induced inactivation led to a mean ~60% reduction in firing rate at baseline (Figure 1 – Figure Supplement 1C, left; 2D, top; Table 1; p=1.9e-06, Wilcoxon signed rank test) and an average ~45% reduction in firing during presentation of pure tone stimuli (Figure 1 – Figure Supplement 1C, right; 2D, bottom; Table 1; p=1.9e-06, Wilcoxon signed rank test). These results indicate that our optogenetic parameters significantly suppress cortico-collicular neurons.

Figure 1 – Figure Supplement 1: Cre/FLEX viral injection strategy enables selective inactivation of cortico-collicular neurons. A) Expression of the retroAAV-Cre-GFP construct at the injection site is restricted to IC (top left). tdTomato labeled axons are found in a pattern matching the known topographical distribution of cortico-collicular neurons in IC (left bottom). Somatic AAV9-FLEX-ArchT-tdTomato expression is present in layer 5 and 6 of AC (right, inset). B) Experimental design for recording from AC to confirm presence of inactivated neurons. C) Example of an inactivated neuron (i.e. putative cortico-collicular neuron) exhibiting a strong reduction in firing during silence and during the presentation of pure tones. D) Population data demonstrating reduced firing rates during silence and in response to pure tone stimuli in putative cortico-collicular neurons. Dots represent individual neurons. Bar plots represent means over the population n = 20 CC neurons. Error bars are standard error of the mean.

Parsing of recording sites into central and shell locations

Shell and central regions of IC differ in their tuning, degree of adaptation, and amount of input from AC, and may also play distinct roles in predictive processing (Aitkin et al., 1975; Bajo et al., 2007; Blackwell et al., 2020; Duque et al., 2012; Herbert et al., 1991; Stebbings et al., 2014; Syka et al., 2000). We quantitatively parsed our recording sites by exploiting known differences in the sharpness of tuning and direction of frequency gradients between shell and central regions: shell IC neurons tend to have broader frequency tuning (low sparseness) than central IC neurons, and the central IC is characterized by a highly stereotyped tonotopic gradient with depth (Figure 1 – Figure Supplement 2A) (Aitkin et al., 1975; Chen et al., 2012; Malmierca et al., 2008; Stiebler & Ehret, 1985; Syka et al., 2000). Similar to previously established procedures used in human and monkey IC research, we performed clustering analysis using the mean sparsity and variation in best frequency

Figure 1 – Figure Supplement 2: Parsing of recording sites into central and shell locations. A) Experimental design for awake IC recordings in the central and shell regions of IC. B) Linear fits for best frequency vs. depth in central (left) and shell (right) IC. C) Sparseness vs. R^2 value for linear fit. K-means clustering was performed using these parameters to classify recording sites as either in the shell or central nucleus of IC. D) Left: DiA labeling from an electrode penetration in a recording site classified as a central site. Atlas image overlay confirms that the dye track runs through the central IC (CIC) (Paxinos & Franklin, 2019). Right: DiD labeling from an electrode penetration in a recording site classified as a shell site. Atlas image overlay confirms that the dye track runs through the shell IC (here denoted as ECIC/DCIC). E) Example raster plots and peristimulus time histograms showing different firing types in the awake IC. F) Example tuning curves in central (left) and shell (right) IC. G) Example of a tuning curve with inhibited side-bands.

with depth from each recording site to determine whether it was from the central nucleus or shell regions of IC (Figure 1 – Figure Supplement 2B, 3C) (Bulkin & Groh, 2011; Ress & Chandrasekaran, 2013). In a subset of recordings, we also marked the recording electrode with a lipophilic dye to histologically confirm the recording location (Figure 1 – Figure Supplement 2D).

IC neurons in both regions exhibited multiple response types to pure tone stimuli (Figure 1 – Figure Supplement 2E). In addition to excitatory responses (e.g. onset and sustained responses), inhibited and offset responses were common, as has previously been characterized in IC of awake animals (Figure 1 – Figure Supplement 2E, top right, bottom middle) (Duque & Malmierca, 2015). Consistent with previous findings, tuning curves from central regions were sharp and narrow, whereas neurons in shell regions exhibited broad frequency tuning (Figure 1 – Figure Supplement 2F, left vs. right) (Aitkin et al., 1975; Syka et al., 2000). Inhibited side-bands were common in tuning curves from both regions, and some inhibited tuning curves were observed (Figure 1 – Figure Supplement 2G). These data confirm that our experimental parameters elicit sound responses and tuning properties characteristic of central and shell regions of the awake IC (Aitkin et al., 1975; Duque & Malmierca, 2015; Syka et al., 2000).

IC neurons encode different aspects of prediction and repetition in awake and anesthetized states

Much of the research regarding SSA and deviance detection in IC to date has been performed in anesthetized animals, with few studies recording from awake subjects (Duque & Malmierca, 2015; Parras et al., 2017). Given that neuronal responses to sound depend on the state of anesthesia of the subject, it is possible that there are differences in predictive coding metrics between the awake and anesthetized states (Fontanini & Katz, 2008; Gaese & Ostwald, 2001; Schumacher et al., 2011). While previous studies have characterized how anesthesia affects SSA, it remains unknown whether its component repetition and prediction metrics differ with anesthetic state (Duque & Malmierca, 2015). Therefore, we first characterized how anesthesia affects these predictive coding metrics in a subset of animals. We first performed awake recordings and then repeated our experimental procedures, leaving the animal head-fixed and the probe in place, after anesthetizing the mouse with isoflurane (Figure 2A). This protocol allowed us to compare how metrics of predictive coding differ between the awake and anesthetized preparations in the same population of neurons.

In the central IC, the mean iMM in the anesthetized condition was positive, indicative of prevalent adaptation (Figure 2B). The iMM values under anesthesia were significantly higher than those obtained while the animal was awake (Figure 2B, Table 1; p=8.8e-05, Wilcoxon rank sum test). To better understand what prediction or repetition effects underlie iMM in each condition, the iMM

for both distributions was further decomposed into an iPE and iRS. In the anesthetized condition, the
mean iPE value of 0.077 indicated the presence of modest prediction error, while an iPE value of -
0.13 indicated that negative prediction error is significantly more prevalent in the awake condition
(Figure 2C, Table 1; p=0.017, Student's T-test). Under both anesthetized and awake conditions,
prominent repetition suppression was observed in the central IC (Figure 2D).

Similar to the central IC, the mean iMM was significantly more positive in shell regions during
anesthesia (Figure 2E, Table 1; p=3.5e-08, Wilcoxon rank sum test). A greater proportion of neurons
in the awake condition had a negative iMM compared with the anesthetized distribution, indicating
that facilitation (a greater response to the standard than the deviant context) is more common in the
awake than the anesthetized condition (Figure 2E). The iPE values in shell IC suggest that prediction
error is significantly higher in the awake compared to the anesthetized condition (Figure 2F, Table 1;
p=2.6e-05, Wilcoxon rank sum test). Although the distribution for the iRS under anesthesia had a
positive mean of 0.25, indicating prevalent repetition suppression, the awake distribution exhibited a
significant leftward shift by comparison (Figure 2G). Interestingly, the mean iRS for the awake
condition was negative (mean=-0.056), indicating that repetition *enhancement*, rather than
suppression, is present in the awake shell IC (Figure 2G, Table 1; p=2.5e-16, Wilcoxon rank sum
test). These results point to differences between predictive coding metrics in the awake and
anesthetized states, with previously undescribed metrics such as repetition enhancement and negative
prediction error more prominent in awake animals.

Adapting and facilitating neurons are differentially affected by cortico-collicular inactivation

We next performed recordings in IC of awake mice to determine how neuronal mismatch and
its component repetition and prediction metrics were affected by cortico-collicular inactivation
(Figure 3A). To inactivate cortico-collicular feedback, we shined light over AC in subjects which
expressed a suppressive opsin in cortico-collicular neurons. We segregated the population of recorded
neurons according to those that exhibited a significantly stronger response to the deviant than the
standard (adapting neurons; Figure 3B, blue; 5C), those that exhibited a significantly stronger
response to the standard than the deviant (facilitating neurons; Figure 3B, red; 5F), and those that
responded equally to both stimulus contexts (non-adapting neurons; Figure 3B, green) for recordings
in both central and shell regions of IC (Figure 3B, left vs. right).

The iMM for adapting neurons in the central nucleus significantly decreased with laser
inactivation of cortico-collicular neurons (Figure 3D, top; Table 1; p=0.00034, Wilcoxon signed rank
test). The iMM at baseline for adapting neurons predominantly represents repetition suppression

Figure 2: IC neurons encode different aspects of prediction and repetition in awake and anesthetized states. A) Experimental design for recording in the awake and isoflurane anesthetized IC in the same population of neurons. B) Distribution of iMM in the awake vs. anesthetized central IC. Bar plots represent means over the population of n = 39 neurons. Error bars are standard error of the mean. C) iPE distribution in the awake vs. anesthetized central IC. D) iRS distribution in the awake vs. anesthetized central IC. E) Distribution of iMM in the awake vs. anesthetized shell IC. Bar plots represent means over the population of n = 165 neurons. Error bars are standard error of the mean. F) iPE distribution in the awake vs. anesthetized shell IC. G) iRS distribution in the awake vs. anesthetized shell IC.

(Figure 3D, bottom) and a small amount of prediction error (Figure 3D, middle). Prediction error was abolished during laser inactivation (Figure 3D, middle; Table 1; p=0.048, Wilcoxon signed rank test), while repetition suppression remained unaffected (Figure 3D, bottom). Adapting neurons in shell regions of IC exhibited a similar pattern to those in the central nucleus. At baseline, these neurons encoded both prediction error and repetition suppression (Figure 3E, middle and bottom). A significant decrease in iMM during laser inactivation (Figure 3E, top; Table 1; p=0.0023, Wilcoxon signed rank test) was driven by a decrease in prediction error (Figure 3E, middle; Table 1; p=0.034, Wilcoxon signed rank test), whereas repetition suppression remained unaffected (Figure 3E, bottom). Combined, these results suggest that removing cortical feedback reduced prediction error but not repetition suppression in adapting neurons.

Prior studies of deviance detection in IC have focused exclusively on adapting neurons. However, given the relative prevalence of facilitating neurons discovered in the awake versus anesthetized IC (Figure 2), we further investigated this population of neurons to determine whether facilitation reflects prediction or repetition effects. In the central nucleus, cortico-collicular inactivation led to a significant decrease in facilitation in facilitating neurons (Figure 3G, top; Table 1; p=0.0036, Student's t-Test). At baseline, the iMM for facilitating neurons represents a combination of negative prediction error and repetition enhancement (Figure 3G, middle and bottom). During inactivation, negative prediction error remained unaffected (Figure 3G, middle), while repetition enhancement was nearly abolished (Figure 3G, bottom; Table 1; p=0.0026, Student's t-Test). Facilitating neurons in the shell IC were also significantly affected by cortico-collicular inactivation (Figure 3H, top; Table 1; p=0.0016, Wilcoxon signed rank test). In this case, however, the change in iMM was driven by the near abolishment of negative prediction error (Figure 3H, middle; Table 1; p=0.037, Wilcoxon signed rank test), while repetition enhancement was unaffected (Figure 3H, bottom).

These data suggest that adaptation and facilitation in the awake IC are composed of distinct underlying processes: adapting populations in both central and shell regions of IC exhibit prediction error and repetition suppression, while facilitating populations are characterized by negative prediction error and repetition enhancement. In adapting neurons in both central and shell regions, cortico-collicular inactivation significantly decreases prediction error. Facilitating neurons in the central IC display decreased repetition enhancement with cortico-collicular inactivation, while those in shell regions exhibit decreased negative prediction error. To ensure that the laser-induced changes described above were opsin-mediated, we performed control experiments in two mice with identical manipulations to the experimental group, but in the absence of ArchT (Figure 3 – Figure Supplement 1A). At baseline, the control group exhibited a similar distribution of iMM values to the experimental

Figure 3: Adapting and facilitating IC neurons are differentially affected by cortico-collicular inactivation. A) Experimental design for recording in awake IC during laser inactivation of the cortico-collicular pathway. B) Categorization of neurons according to whether they displayed significant adaptation, facilitation, or neither (non-adapting). C) Average peristimulus time histogram for adapting neurons in central (top) and shell (bottom) IC. Green = during laser inactivation. D) iMM (top), iPE (middle), and iRS (bottom) for adapting neurons in the central nucleus. Dots represent individual neurons. Bar plots represent means over the population of n = 52 neurons. Error bars are standard error of the mean. E) iMM (top), iPE (middle), and iRS (bottom) for adapting neurons in shell regions of IC. Dots represent individual neurons. Bar plots represent means over the population of n = 113 neurons. Error bars are standard error of the mean. F) Average peristimulus time histogram for facilitating neurons in central (top) and shell (bottom) IC. Green = during laser inactivation. G) iMM (top), iPE (middle), and iRS (bottom) for facilitating neurons in the central nucleus. Dots represent individual neurons. Bar plots represent means over the population of n = 14 neurons. Error bars are standard error of the mean. H) iMM (top), iPE (middle), and iRS (bottom) for facilitating neurons in shell regions of IC. Dots represent individual neurons. Bar plots represent means over the population of n = 38 neurons. Error bars are standard error of the mean. This figure has Figure Supplements 1 and 2.

group in both the central and shell regions of IC (Figure 3 – Figure Supplement 1B, Table 2). We found no significant differences between baseline and laser trials for either adapting (Figure 3 – Figure Supplement 1C, D, Table 2) or facilitating (Figure 3 – Figure Supplement 1E, F) neurons in either region. This experiment confirmed that the observed effects of cortico-collicular inactivation were indeed due to opsin-mediated inactivation of the cortico-collicular projection neurons.

Figure 3 – Figure Supplement 1: Control data. A) Experimental design for control experiments. All procedures were performed identically to the experimental group, except with the omission of ArchT from the viral construct injected in AC. B) Comparison of iMM distribution for control (navy) and experimental (light blue) groups in central (left) and shell (right) IC. C) iMM (top), iPE (middle), and iRS (bottom) for control adapting neurons in the central nucleus. Dots represent individual neurons. Bar plots represent means over the population of n = 18 neurons. Error bars are standard error of the mean. D) iMM (top), iPE (middle), and iRS (bottom) for control adapting neurons in shell regions of IC. Dots represent individual neurons. Bar plots represent means over the population of n = 35 neurons. Error bars are standard error of the mean. E) iMM (top), iPE (middle), and iRS (bottom) for control facilitating neurons in the central nucleus. Dots represent individual neurons. Bar plots represent means over the population of n = 4 neurons. Error bars are standard error of the mean. F) iMM (top), iPE (middle), and iRS (bottom) for control facilitating neurons in shell regions of IC. Dots represent individual neurons. Bar plots represent means over the population of n = 21 neurons. Error bars are standard error of the mean. G) iMM (top), iPE (middle), and iRS (bottom) for control non-adapting neurons in the central nucleus. Dots represent individual neurons. Bar plots represent means over the population of n = 55 neurons. Error bars are standard error of the mean. H) iMM (top), iPE (middle), and iRS (bottom) for control non-adapting neurons in shell regions of IC. Dots represent individual neurons. Bar plots represent means over the population of n = 63 neurons. Error bars are standard error of the mean.

Adapting and facilitating neurons respond similarly to the cascade and many standards controls

        Though the cascade sequence is free of repetition effects between adjacent tone pairs, it does exhibit global repetition across the entire tone sequence. To assess whether global stimulus regularity affects the response to the cascade context, we used a shuffled version of the cascade sequence, known as the "many standards" sequence, as an additional control stimulus (Figure 3 – Figure Supplement 2A). The many standards sequence contains the same 10 tones as the cascade but presented in random order (Figure 3 – Figure Supplement 2A). This reduces the potential for adaptation across adjacent frequency channels and also eliminates the global predictability of the stimulus, both of which could lead to suppression of responses to tones in the cascade context and potentially affect the calculations of iMM, iPE, and iRS. We compared the responses of adapting and facilitating neurons in both central and shell regions of IC to tones in the cascade versus the many standards context (Figure 3 – Figure Supplement 2A). We found no significant differences in firing rates to the cascade versus the many standards contexts (Figure 3 – Figure Supplement 2B, C, Table 1), suggesting that the global structure of the cascade sequence does not significantly affect how neurons in IC respond to this stimulus, as has been shown in other structures (Casado-Román et al., 2020; Parras et al., 2021).

Figure 3 – Figure Supplement 2: Comparison of neuronal responses between the many standards and cascade sequences. A) The many standards sequence consists of the same 10 tones found in the cascade sequence, but the tone order is random. Responses to the cascade and many standards sequences were compared to assess whether cross-frequency adaptation or global stimulus regularity affect responses to the cascade condition. B) Firing rates of adapting neurons (left) and facilitating neurons (right) in the central IC to tones in the cascade and many standards contexts. Dots represent individual neurons. Bar plots represent means over the population of n = 52 adapting and n = 14 facilitating neurons. Error bars are standard error of the mean. C) Firing rates of adapting neurons (left) and facilitating neurons (right) in the shell IC to tones in the cascade and many standards contexts. Dots represent individual neurons. Bar plots represent means over the population of n = 113 adapting and n = 38 facilitating neurons. Error bars are standard error of the mean.

Non-adapting units also display top-down repetition enhancement

The majority of neurons in both central and shell IC do not exhibit either adaptation or facilitation but respond similarly to tones when they are presented as a standard or deviant (Figure 4A). However, since both negative and positive metrics are included in the calculation of iMM, it is still possible that these units exhibit predictive processing that may not be reflected in the overall iMM value. We further characterized these non-adapting neurons (Figure 4B) and tested how they are affected by cortico-collicular inactivation. Non-adapting neurons in the central nucleus exhibited a significant increase in iMM during inactivation (Figure 4C, top; Table 1; p=2.7e-06, Wilcoxon signed rank test), whereas those in the shell IC were unaffected (Figure 4D, top). The change in iMM for non-adapting neurons in the central nucleus was driven by a significant increase in iRS (Figure 4C, bottom middle; Table 1; p=0.0011, Wilcoxon signed rank test). To determine whether this reflected a change in repetition suppression or enhancement, we further segregated central non-adapting units according to whether their baseline iRS values were negative or positive (Figure 4C, bottom). Only those units with negative baseline iRS values (i.e., those units showing repetition enhancement) were significantly affected by cortico-collicular inactivation (Figure 4C, bottom; Table 1; p=0.00012, Wilcoxon signed rank test). In control experiments without ArchT, no significant changes were observed in non-adapting neurons (Figure 3 – Figure Supplement 1G, H, Table 2). These results indicate that, similar to central facilitating units, central non-adapting units display repetition enhancement, and that input from the cortex is critical for expression of this phenomenon.

Standard and deviant responses are bidirectionally modulated by cortico-collicular inactivation

The observed changes in repetition metrics with cortico-collicular inactivation could reflect an effect on either the standard or cascade context. Similarly, the shift in prediction metrics observed with inactivation could be due to altered responses to either the cascade or deviant contexts. We next determined whether the laser-induced changes in the iMM, the iPE, and the iRS for adapting neurons reflect changes in the firing rates to the standard, deviant, or cascade contexts. We found that adapting neurons in the central nucleus increased responses to the standard (Figure 5A, Table 1; p=0.0092, one sample t-test) and decreased responses to the deviant (Figure 5A, Table 1; p=0.0054, one sample t-test) during inactivation. These results explain the decrease in iMM for this population during the laser stimulus (Figure 3D, top): the firing rate to the cascade stimulus did not change during cortico-collicular inactivation, which means that the decrease in firing rate to the deviant alone underlies the decrease in prediction error observed for this population (Figure 3D, middle). Adapting neurons in the shell exhibited the same pattern of bidirectional changes to the standard (Figure 5B, Table 1;

Figure 4: Non-adapting units also display top-down repetition enhancement. A) Distribution of adapting types (adapting, facilitating,
and non-adapting) for neurons in central (left) and shell (right) regions of IC. B) Average peristimulus time histogram for non-
adapting neurons in central (top) and shell (bottom) IC. C) iMM (top), iPE (middle), and iRS (bottom) for non-adapting neurons in
central regions of IC. Dots represent individual neurons. Bar plots represent means over the population of n = 155 neurons. Error bars
are standard error of the mean. D) iMM (top), iPE (middle), and iRS (bottom) for non-adapting neurons in shell regions of IC. Dots
represent individual neurons. Bar plots represent means over the population of n = 243 neurons. Error bars are standard error of the
mean.

p=0.035, one sample Wilcoxon test) and deviant (Figure 5B, Table 1; p=0.0057, one sample
Wilcoxon test), similarly accounting for their decrease in iMM and prediction error (Figure 3E), with
no change in response to the cascade condition (Figure 5B). These data suggest that inactivation of
the cortico-collicular pathway induces bidirectional changes in firing rates to the standard and deviant
for adapting neurons in both central and shell regions of IC.

We also investigated how responses to each stimulus context changed with cortico-collicular
inactivation for facilitating neurons. For central facilitating neurons, only the firing rate to the
standard context changed during inactivation (Figure 5C, Table 1; p=0.0013, one sample t-test),
explaining the observed change in repetition enhancement for this population (Figure 3G). For shell
facilitating neurons, a decreased response to the standard (Figure 5D, Table 1; p=0.0042, one sample
t-test) and an increased response to the deviant (Figure 5D, Table 1; p=0.0013, one sample t-test)
were elicited on laser trials, accounting for changes in the iMM and the abolishment of negative
prediction error (Figure 3H). These changes are directionally opposite to the observed firing rate
changes observed for adapting neurons under inactivation, with a decrease to the standard context for
both central and shell neurons and an increase to the deviant context for shell neurons.

For non-adapting neurons, a significant decrease in response to the standard context was
observed in both central (Figure 5E, Table 1; p=1.4e-06, one sample Wilcoxon test) and shell (Figure
5F, Table 1; p=0.035, one sample Wilcoxon test) regions of IC. The decrease was only significant
enough to produce an effect on the iMM in central regions (Figure 4C, top), leading to an increase in
repetition suppression (Figure 4C, bottom).

For adapting and facilitating neurons, these data exhibit that IC responses to the standard and
deviant contexts in the absence of cortical input are bidirectionally modulated, such that neurons
respond more similarly to both contexts rather than firing differentially to each. For non-adapting
neurons, the response to the standard context alone is diminished during cortico-collicular
inactivation, causing these neurons to become more adapting. These changes suggest that under
normal conditions, AC provides information regarding sound context to neurons in IC.

Individual neurons have distinct combinations of iPE and iRS

To determine whether sensitivity to repetition and prediction is encoded in distinct neuronal
populations, or whether individual neurons exhibit particular combinations of repetition
suppression/enhancement and prediction error/negative prediction error, we plotted the iPE values
against the iRS values for each neuron in the adapting, facilitating, and non-adapting groups. Both
the adapting and non-adapting groups in the central IC contained neurons with significant values for

Figure 5: Standard and deviant responses are bidirectionally modulated by cortico-collicular inactivation. A) Responses to the
standard (left), cascade (middle left), and deviant (middle right) for adapting neurons in central regions of IC under baseline and laser

conditions. Change in firing rate between the laser and baseline condition for each stimulus (right). Dots represent individual neurons. Bar plots represent means over the population of n = 52 neurons. Error bars are standard error of the mean. B) Responses to the standard (left), cascade (middle left), and deviant (middle right) for adapting neurons in shell regions of IC under baseline and laser conditions. Change in firing rate between the laser and baseline condition for each stimulus (right). Dots represent individual neurons. Bar plots represent means over the population of n = 113 neurons. Error bars are standard error of the mean. C) Responses to the standard (left), cascade (middle left), and deviant (middle right) for facilitating neurons in central regions of IC under baseline and laser conditions. Change in firing rate between the laser and baseline condition for each stimulus (right). Dots represent individual neurons. Bar plots represent means over the population of n = 14 neurons. Error bars are standard error of the mean. D) Responses to the standard (left), cascade (middle left), and deviant (middle right) for facilitating neurons in shell regions of IC under baseline and laser conditions. Change in firing rate between the laser and baseline condition for each stimulus (right). Dots represent individual neurons. Bar plots represent means over the population of n = 38 neurons. Error bars are standard error of the mean. E) Responses to the standard (left), cascade (middle left), and deviant (middle right) for non-adapting neurons in central regions of IC under baseline and laser conditions. Change in firing rate between the laser and baseline condition for each stimulus (right). Dots represent individual neurons. Bar plots represent means over the population of n = 155 neurons. Error bars are standard error of the mean. F) Responses to the standard (left), cascade (middle left), and deviant (middle right) for non-adapting neurons in shell regions of IC under baseline and laser conditions. Change in firing rate between the laser and baseline condition for each stimulus (right). Dots represent individual neurons. Bar plots represent means over the population of n = 243 neurons. Error bars are standard error of the mean.

both iPE and iRS, most often resulting from a combination of negative prediction error and repetition suppression (Figure 6A, maroon dots). In the shell IC, a greater variety of response combinations was observed. All three groups contained neurons with both significant negative prediction error and repetition suppression, as well as a separate population exhibiting significant prediction error and repetition enhancement (Figure 6B, maroon dots). Some shell adapting neurons also exhibited a combination of both repetition suppression and prediction error (Figure 6B, left). These results suggest that individual neurons in IC exhibit distinct combinations of repetition suppression/enhancement and prediction error/negative prediction error.

Figure 6: Individual neurons exhibit distinct combinations of iPE and iRS. A) Distribution of both iRS and iPE in individual adapting (left), facilitating (middle), and non-adapting (right) neurons in central IC. B) Plots of distributions of both iRS and iPE in individual adapting (left), facilitating (middle), and non-adapting (right) neurons in shell IC. C) Response to three subsequent standards prior to or following the deviant in facilitating neurons in central IC. Comparison between the last standard before and the first standard after the deviant demonstrates significant repetition enhancement. Bar plots represent means over the population of n = 14 neurons. Error bars are standard error of the mean. D) Response to three subsequent standards prior to or following the deviant in facilitating neurons in shell IC. Comparison between the last standard before and the first standard after the deviant demonstrates significant repetition enhancement. Bar plots represent means over the population of n = 38 neurons. Error bars are standard error of the mean.

Facilitating neurons exhibit true repetition enhancement

Facilitating neurons in both central and shell regions of IC exhibited repetition enhancement

at baseline, as defined by the difference in firing rate to the last standard and the same tone embedded

in the cascade sequence (Figure 3G, 5H). We sought to further characterize the response to the

standard context to determine whether the repetition enhancement captured by the iRS indicates true

repetition enhancement (an incremental increase in firing rate on subsequent presentations of the

standard) or simply a net increase in firing rate to the standard versus cascade condition. We

calculated the mean firing rate for each of the three standards before the deviant and each of the three

standards after the deviant (Figure 6C, 8D). The progression of standards by position exhibited

subsequent enhancements in firing rate that plateaued by the second to last standard before the deviant

for both central (Figure 6C) and shell facilitating neurons (Figure 6D). The firing rate to the last

standard was significantly higher than the first in both regions (Figure 6C, Table 1; p=0.0017,

Wilcoxon signed rank test; Figure 6D, Table 1; p=9.3e-05, Wilcoxon signed rank test). These data

provide evidence that facilitating neurons in IC exhibit true repetition enhancement.

514
515   Figure 7: Cortico-fugal regulation of predictive coding. Laser inactivation led to the abolishment of repetition enhancement in central
516   facilitating units and the abolishment of negative prediction error in shell facilitating units. Prediction error decreased during
517   inactivation for adapting units in both shell and central regions of the IC. Repetition suppression remained unaffected during cortical
518   inactivation, suggestion that it may reflect fatigue of bottom-up sensory inputs.

DISCUSSION

Summary of findings

The results of the present study indicate that AC is critically involved in regulating both repetition and prediction effects in the awake IC, providing evidence for the implementation of predictive coding in cortico-subcortical networks. Adapting and facilitating neurons were bi-directionally modulated by cortico-collicular inactivation, with adapting neurons becoming less adapting and facilitating neurons becoming less facilitating on laser trials (Figure 3). The decrease in adaptation for adapting neurons was driven by a decrease in prediction error for neurons in both central and shell regions of IC (Figure 3D, 5E; Figure 7, pink arrows). For facilitating and non-adapting neurons in the central nucleus, inactivation-driven changes were caused by a decrease in repetition enhancement (Figure 3G; Figure 7, gold dashed arrows). The decrease in facilitation in the shell IC, however, was caused by the abolishment of negative prediction error (Figure 3H; Figure 7, pink dashed arrows).

In adapting neurons, these changes were modulated by an increased response to the standard and a decreased response to the deviant, while the opposite pattern was true for facilitating neurons (Figure 5). Overall, these bi-directional changes indicate that, without input from AC, IC responds more similarly to tones in the standard and deviant contexts. These findings demonstrate that AC provides critical contextual cues about the statistics of the auditory environment to targets in IC under normal conditions. We further discuss these results in the context of a hierarchical predictive coding framework below.

iMM in the awake versus anesthetized IC

Our results include the first investigation of how the repetition and prediction processes that underlie deviance detection in the awake IC compare to the anesthetized condition. Our data suggest that while iMM values are higher under anesthesia, they almost entirely reflect repetition suppression, with only a small contribution of prediction error (Figure 2). In the central IC, modest prediction error is present under anesthesia, but negative prediction error becomes dominant when the animal is awake. In the shell IC, the same neurons exhibit drastically different iPE and iRS values for the awake versus the anesthetized condition. Prediction error is substantially higher in the awake IC and repetition enhancement, rather than repetition suppression, is observed (Figure 2F, 4G). These findings suggest that the iMM values in the awake and anesthetized brain reflect different underlying processes, and that anesthesia induces bidirectional changes in metrics of repetition and prediction.

Facilitating neurons in IC

We also provide here the first analysis of facilitating neurons in IC. Previous studies that have

investigated iMM have focused selectively on the positive side of the iMM distribution, since these

neurons display adaptation. However, facilitation seems to be enriched in the awake IC (Figure 2B,

4E) and reflects other potentially interesting parameters, such as repetition enhancement (represented

as a higher response to the standard than the cascade sequence) (Figure 2G) and negative prediction

error (represented as a higher response to the cascade than the deviant) (Figure 2C).

Repetition enhancement and repetition suppression in IC

Because previous studies that have applied a predictive coding framework to decompose

neuronal mismatch have focused exclusively on adapting neurons, the repetition enhancement found

here in facilitating neurons has not been previously described (Parras et al., 2017). However, it is

well-documented in fMRI literature that repetition enhancement is a common phenomenon in

humans, existing either alongside or in place of repetition suppression (De Gardelle et al., 2013;

Müller et al., 2013; Segaert et al., 2013). Interestingly, repetition enhancement has been proposed to

reflect novel network formation and consolidation of novel sensory representations (Segaert et al.,

2013). Once new representations have been formed, repetition suppression is hypothesized to take

over, reflecting the minimization in prediction errors that occurs when new representations give rise

to accurate predictions (Auksztulewicz & Friston, 2016; De Gardelle et al., 2013; Friston & Kiebel,

2009). Though the repetition enhancement described in human studies differs drastically on spatial

and temporal scales from the phenomenon described here, we find that it similarly involves a

sequential enhancement in the response to subsequent presentations of the standard (Figure 6C, 8D).

Repetition enhancement has also been observed in the medial geniculate body in response to

temporally degraded stimuli that are hypothesized to engage top-down resources to compensate for

bottom-up acoustic information loss (Cai et al., 2016; Kommajosyula et al., 2019). Interestingly, this

enhancement is reversed when cortico-thalamic pathways are blocked, further suggesting that

repetition enhancement in the auditory system reflects a top-down phenomenon (Kommajosyula et

al., 2021).

While repetition suppression can be understood from a predictive coding framework, it can

also be viewed from the perspective of neuronal fatigue, whereby the incremental decrease in firing

rate to a repeated standard tone is simply explained by synaptic depression (Escera & Malmierca,

2014; Taaseh et al., 2011). Interestingly, we did not find any effect on repetition suppression during

cortico-collicular inactivation, suggesting that it may reflect fatigue of bottom-up sensory inputs

rather than an active predictive process (Figure 3D, 5E; Figure 7, gold arrows). While these data do
not provide definitive proof of either perspective, they do suggest that the processes that underlie
repetition suppression in IC do not involve top-down cortical signals. This notion is supported by the
fact that repetition suppression was much more prevalent when animals were under anesthesia, a state
in which the auditory responsiveness in the cortex is compromised (Figure 2G) (Brugge & Merzenich,
1973; Katsuki et al., 1959).

Prediction error in IC

In both central and shell populations that exhibited prediction error at baseline, cortico-
collicular inactivation led to a decrease, or complete abolishment, of prediction error (Figure 3D, 5E).
According to models of hierarchical predictive coding, higher-order stations generate predictions that
they broadcast to lower centers (Friston & Kiebel, 2009). These predictions are compared with
representations of the actual sensory input, and if there is a mismatch, a prediction error is generated
and forwarded up the hierarchy (Friston & Kiebel, 2009). Under this framework, the inactivation of
top-down inputs would interfere with communication of predictions, leading to dysfunction in the
prediction error response, as seen in our data. Another possibility is that prediction errors are directly
backpropagated from AC to IC. While this contradicts canonical predictive coding models, evidence
for prediction error has been found in deep layers of the cortex in which feedback neurons reside
(Asilador & Llano, 2020; Rummell et al., 2016). Though the precise mechanism underlying the
generation of prediction error in IC remains unclear, our data show that feedback from AC plays a
critical role in this process.

Negative prediction error in IC

In addition to neurons with prediction error, we found neurons in IC that responded more
strongly to the cascade than the deviant context (Figure 3G, 5H), consistent with previous reports
(Parras et al., 2017). A stronger response to a tone in the cascade sequence compared to the context
in which it is a deviant could simply reflect a relative lack of cross-frequency adaptation; the oddball
stimulus consists of repeated tone presentations of two neighboring frequencies, making it more likely
to generate cross-frequency effects than the cascade stimulus, which cycles through repetitions of 10
evenly-spaced frequencies (Parras et al., 2017; Taaseh et al., 2011). Previous studies that have
investigated the effective bandwidth for cross-frequency adaptation, however, have found that it
occurs between channels with a frequency separation of a third of an octave or less (Taaseh et al.,
2011). The stimuli used in the present study had a half-octave frequency separation, indicating that

cross-frequency effects should be minimized. Therefore, it is unlikely that the negative prediction
error responses observed in the present study simply reflect cross-frequency adaptation to the oddball
stimulus.

A stronger response to a tone when it is embedded in a completely predictable sequence, such
as the cascade sequence, than when it is a deviant could also signify that a neuron encodes predictions,
rather than prediction errors. In hierarchical predictive coding, both predictions and prediction errors
are generated at every level of the hierarchy, with prediction errors being forwarded to ascending
sensory centers and predictions being backpropagated (Friston & Kiebel, 2009). In the shell IC, the
region which receives the vast majority of descending cortical input, evidence for negative prediction
error was abolished during cortico-collicular inactivation (Figure 3H), consistent with the notion that
feedback from the cortex may carry predictions to IC (Bajo et al., 2007; Herbert et al., 1991; Saldaña
et al., 1996; Stebbings et al., 2014). Interestingly, negative prediction error in the central nucleus
remained unperturbed during inactivation of cortical feedback (Figure 3G). Given that only a small
fraction of cortico-collicular fibers terminate in the central nucleus, it is likely that it receives
predictions from another source (Bajo et al., 2007; Herbert et al., 1991; Saldaña et al., 1996; Stebbings
et al., 2014). An intriguing potential candidate for this source of predictions could be the shell IC,
given the extensive network of intracollicular connections in IC (Lesicko & Llano, 2020; Saldaña &
Merchań, 1992; Saldaña & Merchán, 2005). Future studies will be required to determine whether the
negative prediction error metric described here captures the type of top-down predictions described
in canonical predictive coding models.

Technical considerations

One limitation of the present study is that laser inactivation achieved only partial and not
complete inactivation of the cortico-collicular pathway. Given that light itself can have a modulatory
or toxic effect on neurons, these types of optogenetic experiments require a careful titration between
using enough power to substantially affect the population of interest without causing non-specific
light or heat-based perturbations (Tyssowski & Gray, 2019). Though other techniques, such as
chemogenetic approaches or cooling, provide more complete inactivation, they do not allow for rapid
and reversible inactivation (English & Roth, 2015). With our laser power parameters, we found a
mean 60% reduction in firing in putative cortico-collicular neurons at baseline and a 45% reduction
during presentation of pure tone stimuli (Figure 1 – Figure Supplement 1D). This reduction produced
clear effects on repetition and prediction processing in IC, in several cases with the severe reduction
or complete abolishment of certain metrics of deviance detection, such as prediction error and

repetition enhancement in the central nucleus and negative prediction error in the shell IC (Figure 3).

The interpretation of these results should bear in mind that they reflect only partial and not complete

inactivation.

Conclusions

Our findings indicate that deviance detection and predictive coding in IC involves additional

complexity than has been previously described. We provide here the first description of facilitating

neurons in IC, as well as evidence for the existence of repetition enhancement and negative prediction

error in these neurons. We show that AC regulates these metrics and is also involved in the

generation of prediction error in IC. Repetition suppression is unaffected by inactivation of

cortical input to IC, providing evidence that this process may reflect bottom-up fatigue rather

than top-down predictive processing. These results demonstrate the role of AC in providing

contextual cues about the auditory stream to targets in IC.

MATERIALS AND METHODS

Animals

We performed experiments in six adult Cdh23 mice (Cdh23tm2.1Kjn/J, JAX: 018399; 4 males and 2 females, age 3-8 months). This mouse line has a targeted point reversion in the Cdh23 gene that protects against the age-related hearing loss common to C57BL/6 strains (Johnson et al., 2017). Animals were housed on a reversed 12-hour light–dark cycle with water and food available ad libitum. All procedures were approved by the University of Pennsylvania IACUC and the AALAC Guide on Animal Research. We made every attempt to minimize the number of animals used and to reduce pain or discomfort.

Virus injection

Mice were continuously anesthetized with isoflurane and mounted in a stereotaxic frame. Buprenex (0.1 mg/kg), Meloxicam (5 mg/kg) and Bupivicane (2 mg/kg) were injected subcutaneously for preoperative analgesia. We performed small craniotomies bilaterally over AC (−2.6 mm caudal to Bregma, ±4.3 mm lateral, +1 mm ventral) and IC (−4.96 mm caudal to Bregma, ±0.5 mm lateral, +0.5 mm ventral and −4.96 mm caudal to Bregma, ±1.25 mm lateral, +1.0 mm ventral). A glass syringe (30-50 μm diameter) connected to a pump (Pump 11 Elite, Harvard Apparatus) was used to inject modified viral vectors (AAV9-CAG-FLEX-ArchT-tdTomato or AAV9-CAG-FLEX-tdTomato; 750 nL/site; UNC Vector Core) into AC and a retroAAV construct (retro AAV-hSyn-Cre-GFP; 250 nL/site) into IC (Figure 1A, 2A, Figure 3 – Figure Supplement 1A). Large viral injections were performed to broadly target cortico-collicular neurons throughout all regions of the auditory cortex. We implanted fiber-optic cannulas (Thorlabs, Ø200 μm Core, 0.22 NA) bilaterally over AC injection sites (0.4 mm ventral to brain surface) and secured them in place with dental cement (C and B Metabond) and acrylic (Lang Dental). IC injection sites were covered with a removable silicone plug (Kwik-Sil). A custom-built headplate was secured to the skull at the midline and a ground-pin was lowered into a small craniotomy over Bregma. We injected an antibiotic (5 mg/kg Baytril) subcutaneously for four days postoperatively. Virus injection sites were confirmed postmortem for all animals included in the study.

Extracellular recordings

We performed recordings a minimum of 21 days after virus injection surgeries to allow adequate travel time for the viral constructs (Figure 1A). Recordings were carried out inside a double-walled acoustic isolation booth (Industrial Acoustics) or a custom-built table-mounted acoustic isolation

booth. For IC recordings, mice were briefly anesthetized to remove the silicone plug over IC virus injection sites. Following recovery from anesthesia, the headplate was clamped within a custom base to provide head-fixation. We lowered a 32-channel silicon probe (Neuronexus) vertically into IC during presentation of broadband noise clicks and monitored sound responses online to confirm localization within IC (Figure 1A). In a subset of animals, the probe was first coated in a lipophilic dye (DiD or DiA; Invitrogen) to aid in posthoc reconstruction of recording sites. In each animal, two recordings were performed per IC (four total recording sessions bilaterally). Following completion of all IC recording sessions, we recorded the activity of neurons in AC using the same procedure (Figure 1 – Figure Supplement 1B). We performed a square craniotomy (2 mm x 2 mm) over AC and oriented the probe vertically to the cortical surface (35-degree angle of the stereotaxic arm). Electrophysiological data were filtered between 600 and 6000 Hz to isolate spike responses and then digitized at 32 kHz and stored for offline analysis (Neuralynx). For a subset of recordings, the experimental procedures were repeated while recording from the same units after the animal had been anesthetized with isoflurane (Figure 2A). We performed spike sorting using Kilosort2 software (https://github.com/MouseLand/Kilosort). Both single and multi-units were included for all analyses (experimental IC: 50 single units, 354 multi-units; control IC: 17 single units; 111 multi-units; anesthetized: 10 single units, 129 multi-units; AC: 95 single units, 300 multi-units; putative cortico-collicular: 9 single units; 11 multi-units).

Laser inactivation

We inactivated cortico-collicular neurons using a 532 nm DPSS laser (GL532T3-300, Slocs lasers, 3 mW power at cannula tip or OptoEngine, MGL-III-532, 15 mW power at cannula tip) connected via optical fibers to the implanted cannulas (Figure 1A, 2C, 2D). Data collected using either laser was pooled together, as no significant differences were observed in the strength of inactivation in AC during silence (p=0.054, Wilcoxon rank sum test) or the presentation of pure tone stimuli (p=0.072, Wilcoxon rank sum test) between the two lasers. Square laser pulses were timed to coincide with tone onset and lasted for 100 ms. Evidence of inactivation in putative cortico-collicular neurons (infragranular AC neurons with a minimum 30% reduction in both baseline and sound-evoked neuronal activity) was confirmed for all animals included in the study.

Stimuli

We generated an initial frequency response function from a sequence of 50 pure tones, 1-70 kHz, repeated 20 times at 70 dB SPL in pseudo-random order. This response function was generated

online to select suitable frequencies for the oddball stimuli, i.e. frequencies that would fall into the

average response area for neurons in a given recording. Each tone was 50 ms duration (1 ms cosine

squared ramps) with an inter-stimulus interval (ISI) of 200 ms and presentation rate of 4 Hz. A similar

tuning curve stimulus, with 8 amplitude levels (35-70 dB, 5 dB increments) and 5 repetitions, was

used to further characterize the tuning properties of each neuron (Figure 1 – Figure Supplement 2E,

3F).

Oddball tone pairs were chosen to fit within the average response area for neurons from a

given recording. Given the prevalence of inhibited regions in the tuning curves, and the fact that this

often led to differences in the response profile of the neuron to each frequency in the oddball tone

pair, the responses to each frequency were analyzed separately (Figure 1 – Figure Supplement 2F).

Oddball stimuli consisted of a frozen sequence of two pure tones (with the same tone parameters as

those used in the initial frequency response functions) with a 90:10 standard-to-deviant ratio and half-

octave frequency separation. The number of standards interleaved between two deviants was

counterbalanced and varied between 3 and 17 standards. The stimuli were divided into blocks (with

the end of a block defined by the presentation of a deviant), and tone type and laser pairings were

alternated on subsequent blocks. For example, on the first block the laser stimulus was paired with

the deviant, on the second block it was paired with the last standard, and the corresponding tones in

the third block served as baseline controls, with no laser stimulus. The number of preceding standards

in the blocks was balanced for all three laser conditions (deviant, last standard, and baseline). Each

block type (laser + standard, laser + deviant, no laser) was presented 45 times and the total number

of tones in each sequence was 1250. Two oddball sequences were created, both with the same frozen

pattern, but with the frequencies of the standard and the deviant switched.

Cascade sequences consisted of either an ascending or descending set of 10 evenly log-spaced

(half-octave separation) pure tones (same tone parameters as described above) (Figure 1C). The two

tones used in the oddball sequences were always included as adjacent tones in the cascade sequences,

though their position within the cascade was varied. To generate the many standards control sequence,

we shuffled the cascade sequences using an algorithm that does not allow for repetition of tones of

the same frequency on subsequent presentations.

Analysis

To distinguish between shell and central IC recording locations, we plotted the best frequency

for each neuron from a given recording against its depth and fit the data with a robust linear regression

model (Figure 1 – Figure Supplement 2B). Additionally, we computed the mean sparsity for all

neurons from a given recording site to quantify the sharpness of tuning. The R^2 metric from the linear
fit and the mean sparsity from each recording were used to perform k-means clustering with two
groups. Each recording was assigned to a location (either central or shell) according to the k-means
output, with central sites typically having high sparsity and high R^2 values and shell sites having low
sparsity and low R^2 metrics (Figure 1 – Figure Supplement 2C).

Sound response profiles were categorized quantitatively from analysis of the combined
responses to the standard and deviant tones using MATLAB's "findpeaks" function with a minimum
peak height set to the mean of the baseline period (50 ms before tone onset) +/- 3 standard deviations.
Units that did not display maxima or minima during the tone duration period (0-50 ms) or in the 50
ms after (the "offset window") were labeled as sound unresponsive and were removed from the
analysis. Units that showed only a single minimum ("inhibited" units) or only a response in the offset
window were similarly removed from the analysis. Units that showed at least one maxima during the
tone duration period were included in the analysis and further categorized as either onset (single
maxima in the first 10 ms after tone onset), sustained (single maximum after the first 10 ms after tone
onset), E-I or I-E (units that displayed both a maximum and minimum during the tone duration
period), biphasic (units that displayed two maxima during the tone duration period), or mixed (units
with greater than 2 maxima and/or minima during the tone response period). It was common for units
to display a response both during the tone duration window and the offset window, and in these cases
a combined response profile was assigned (e.g., onset/offset, sustained/inhibited-offset). Neurons
with only inhibited or offset responses were removed from the data set.

Significant adaptation or facilitation for each neuron was assessed with a Wilcoxon rank sum
test between the trial-by-trial firing rates to the standard and deviant on the 45 baseline trials. The
index of neuronal mismatch (iMM), identical to the traditional SSA index, was further deconstructed
into an index of prediction error (iPE) and an index of repetition suppression (iRS) such that iMM =
iPE + iRS. The raw firing rates to the standard, cascade, and deviant conditions were normalized by
dividing by the Euclidean norm, $N = \sqrt{FR_{Dev}^2 + FR_{Casc}^2 + FR_{Stan}^2}$. The iPE was calculated as
the difference in normalized firing rate to the deviant and cascade conditions ($iPE = \frac{FR_{Dev}}{N} - \frac{FR_{Casc}}{N}$),
while the iRS was calculated as the difference in normalized firing rate to the cascade and standard
conditions ($iRS = \frac{FR_{Casc}}{N} - \frac{FR_{Stan}}{N}$).

Statistical analysis

Shapiro-Wilk tests were used to assess normality. For normally distributed data, Student's T-tests
were performed. When the assumption of normality was violated, Wilcoxon rank sum tests were used
for nonpaired data and Wilcoxon signed rank tests were used for paired data. Cohen's d was calculated
as measure of effect size for t-tests. For Wilcoxon tests, the effect size r was calculated as the z statistic
divided by the square root of the sample size.
799

ACKNOWLEDGEMENTS

The authors thank members of the Geffen laboratory for helpful discussions. This work was

supported by funding from the National Institute of Health grants F32MH120890 to AMHL,

R01DC015527, R01DC014479 and R01NS113241 to MNG, F31DC016524 to CA, and

R01DA044205, R01DA049545 and U01AA025931 to MDB.

Table 1: Statistical comparisons for experimental data.

Comparison	Figure	Mean	Median	SD	SEM	CI (±)	Test	Test statistic	N	df	p	Effect size
Response of putative cortico-collicular neurons in silence (laser OFF vs. ON)	1S1D (top)	OFF: 11 ON: 4.1	OFF: 9.0 ON: 3.5	OFF: 8.9 ON: 3.5	OFF: 2.0 ON: 0.78	OFF: 4.2 ON: 1.6	Wilcoxon signed rank test	V = 0	20	NA	1.9e-06	0.88
Response of putative cortico-collicular neurons to pure tones (laser OFF vs. ON)	1S1D (bottom)	OFF: 18 ON: 9.6	OFF: 8.8 ON: 4.3	OFF: 24 ON: 12	OFF: 5.4 ON: 2.7	OFF: 11 ON: 5.6	Wilcoxon signed rank test	V = 0	20	NA	1.9e-06	0.88
iMM central (awake vs. anesthetized)	2B	Aw: 0.050 An: 0.25	Aw: 0.045 An: 0.28	Aw: 0.21 An: 0.49	Aw: 0.024 An: 0.074	Aw: 0.047 An: 0.15	Wilcoxon rank sum test	W = 952.5	Aw: 78 An: 43	NA	8.8e-05	0.36
iPE central (awake vs. anesthetized)	2C	Aw: -0.13 An: 0.077	Aw: -0.11 An: 0.098	Aw: 0.17 An: 0.53	Aw: 0.019 An: 0.081	Aw: 0.038 An: 0.16	Student's T-test	t = -2.5	Aw: 78 An: 43	38	0.017	0.52
iRS central (awake vs. anesthetized)	2D	Aw: 0.18 An: 0.18	Aw: 0.17 An: 0.30	Aw: 0.17 An: 0.56	Aw: 0.019 An: 0.085	Aw: 0.039 An: 0.17	Wilcoxon rank sum test	W = 1444	Aw: 78 An: 43	NA	0.21	0.12
iMM shell (awake vs. anesthetized)	2E	Aw: 0.095 An: 0.27	Aw: 0.090 An: 0.27	Aw: 0.31 An: 0.35	Aw: 0.025 An: 0.022	Aw: 0.050 An: 0.043	Wilcoxon rank sum test	W = 12502	Aw: 147 An: 254	NA	3.5e-08	0.28
iPE shell (awake vs. anesthetized)	2F	Aw: 0.15 An: 0.018	Aw: 0.15 An: -0.0075	Aw: 0.33 An: 0.39	Aw: 0.027 An: 0.025	Aw: 0.053 An: 0.049	Wilcoxon rank sum test	W = 23368	Aw: 147 An: 254	NA	2.6e-05	0.21
iRS shell (awake vs. anesthetized)	2G	Aw: -0.056 An: 0.25	Aw: -0.085 An: 0.29	Aw: 0.36 An: 0.33	Aw: 0.029 An: 0.020	Aw: 0.058 An: 0.040	Wilcoxon rank sum test	W = 9501.5	Aw: 147 An: 254	NA	2.5e-16	0.41
iMM central adapting (laser OFF vs. ON)	3D (top)	OFF: 0.26 ON: 0.21	OFF: 0.24 ON: 0.19	OFF: 0.096 ON: 0.13	OFF: 0.013 ON: 0.019	OFF: 0.027 ON: 0.037	Wilcoxon signed rank test	V = 1083	52	NA	0.00034	0.50
iPE central adapting (laser OFF vs. ON)	3D (middle)	OFF: 0.0077 ON: -0.029	OFF: 0.036 ON: -0.0041	OFF: 0.16 ON: 0.16	OFF: 0.022 ON: 0.022	OFF: 0.043 ON: 0.044	Wilcoxon signed rank test	V = 907	52	NA	0.048	0.28
iRS central adapting (laser OFF vs. ON)	3D (bottom)	OFF: 0.25 ON: 0.24	OFF: 0.24 ON: 0.24	OFF: 0.16 ON: 0.16	OFF: 0.023 ON: 0.022	OFF: 0.046 ON: 0.045	Wilcoxon signed rank test	V = 832	52	NA	0.19	0.18
iMM shell adapting (laser OFF vs. ON)	3E (top)	OFF: 0.34 ON: 0.31	OFF: 0.32 ON: 0.28	OFF: 0.19 ON: 0.20	OFF: 0.017 ON: 0.019	OFF: 0.035 ON: 0.037	Wilcoxon signed rank test	V = 4283	113	NA	0.00023	0.29
iPE shell adapting (laser OFF vs. ON)	3E (middle)	OFF: 0.15 ON: 0.14	OFF: 0.12 ON: 0.10	OFF: 0.30 ON: 0.30	OFF: 0.028 ON: 0.028	OFF: 0.056 ON: 0.056	Wilcoxon signe	V = 3963	113	NA	0.034	0.20

							d rank test					
iRS shell adapting (laser OFF vs. ON)	3E (bottom)	OFF: 0.19 ON: 0.17	OFF: 0.19 ON: 0.16	OFF: 0.24 ON: 0.24	OFF: 0.023 ON: 0.023	OFF: 0.045 ON: 0.045	Paired t-test	t = 1.6	113	112	0.11	0.15
iMM central facilitating (laser OFF vs. ON)	3G (top)	OFF: -0.32 ON: -0.13	OFF: -0.31 ON: -0.11	OFF: 0.16 ON: 0.19	OFF: 0.042 ON: 0.050	OFF: 0.090 ON: 0.11	Paired t-test	t = -3.5	14	13	0.0036	0.95
iPE central facilitating (laser OFF vs. ON)	3G (middle)	OFF: -0.20 ON: -0.17	OFF: -0.24 ON: -0.20	OFF: 0.20 ON: 0.17	OFF: 0.054 ON: 0.044	OFF: 0.12 ON: 0.095	Paired t-test	t = -1.2	14	13	0.25	0.32
iRS central facilitating (laser OFF vs. ON)	3G (bottom)	OFF: -0.12 ON: 0.036	OFF: -0.092 ON: 0.069	OFF: 0.18 ON: 0.24	OFF: 0.049 ON: 0.064	OFF: 0.11 ON: 0.14	Paired t-test	t = -3.7	14	13	0.0026	1.0
iMM shell facilitating (laser OFF vs. ON)	3H (top)	OFF: -0.29 ON: -0.19	OFF: -0.24 ON: -0.15	OFF: 0.15 ON: 0.16	OFF: 0.024 ON: 0.026	OFF: 0.048 ON: 0.052	Wilcoxon signed rank test	V = 159	38	NA	0.0016	0.50
iPE shell facilitating (laser OFF vs. ON)	3H (middle)	OFF: -0.026 ON: 0.033	OFF: 0.011 ON: 0.023	OFF: 0.26 ON: 0.29	OFF: 0.042 ON: 0.047	OFF: 0.085 ON: 0.096	Wilcoxon signed rank test	V = 227	38	NA	0.037	0.34
iRS shell facilitating (laser OFF vs. ON)	3H (bottom)	OFF: -0.26 ON: -0.23	OFF: -0.29 ON: -0.23	OFF: 0.32 ON: 0.33	OFF: 0.052 ON: 0.054	OFF: 0.11 ON: 0.11	Wilcoxon signed rank test	V = 254	38	NA	0.093	0.27
iMM central non-adapting (laser OFF vs. ON)	4C (top)	OFF: 0.022 ON: 0.072	OFF: 0.023 ON: 0.065	OFF: 0.12 ON: 0.14	OFF: 0.0094 ON: 0.011	OFF: 0.019 ON: 0.022	Wilcoxon signed rank test	V = 3419	155	NA	2.7e-06	0.38
iPE central non-adapting (laser OFF vs. ON)	4C (middle top)	OFF: -0.096 ON: -0.081	OFF: -0.098 ON: -0.093	OFF: 0.19 ON: 0.19	OFF: 0.015 ON: 0.015	OFF: 0.030 ON: 0.030	Wilcoxon signed rank test	V = 5327	155	NA	0.20	0.10
iRS central non-adapting (laser OFF vs. ON)	4C (middle bottom)	OFF: 0.12 ON: 0.15	OFF: 0.12 ON: 0.15	OFF: 0.15 ON: 0.17	OFF: 0.012 ON: 0.013	OFF: 0.024 ON: 0.027	Wilcoxon signed rank test	V = 4224	155	NA	0.0011	0.26
iRS > 0 central non-adapting (laser OFF vs. ON)	4C (bottom)	OFF: 0.17 ON: 0.19	OFF: 0.16 ON: 0.18	OFF: 0.10 ON: 0.15	OFF: 9.1e-03 ON: 0.013	OFF: 1.8e-02 ON: 0.026	Wilcoxon signed rank test	V = 3313	127	NA	0.071	0.16
iRS < 0 central non-adapting (laser OFF vs. ON)	4C (bottom)	OFF: -0.13 ON: -0.012	OFF: -0.10 ON: -0.017	OFF: 0.11 ON: 0.15	OFF: 0.021 ON: 0.029	OFF: 0.044 ON: 0.060	Wilcoxon signed rank test	V = 30	25	NA	0.00012	0.71
iMM shell non-adapting (laser OFF vs. ON)	4D (top)	OFF: 0.0053 ON: 0.023	OFF: 0.0062 ON: 0.028	OFF: 0.13 ON: 0.16	OFF: 0.0081 ON: 0.010	OFF: 0.016 ON: 0.020	Wilcoxon signed rank test	V = 12765	243	NA	0.076	0.11
iPE shell non-adapting (laser OFF vs. ON)	4D (middle)	OFF: 0.053 ON: 0.072	OFF: 0.059 ON: 0.061	OFF: 0.21 ON: 0.20	OFF: 0.013 ON: 0.013	OFF: 0.026 ON: 0.026	Wilcoxon signed rank test	V = 13474	243	NA	0.22	0.079
iRS shell non-adapting (laser OFF vs. ON)	4D (bottom)	OFF: -0.048 ON: -0.049	OFF: -0.042 ON: -0.041	OFF: 0.23 ON: 0.22	OFF: 0.015 ON: 0.014	OFF: 0.029 ON: 0.028	Wilcoxon signed rank test	V = 14344	243	NA	0.66	0.028

FR change standard central adapting	5A	2.1	2.0	5.6	0.78	1.6	One sample t-test	t = 2.7	52	51	0.0092	0.38
FR change cascade central adapting	5A	-0.38	0.67	6.9	0.95	1.9	One sample t-test	t = -0.40	52	51	0.69	0.056
FR change deviant central adapting	5A	-2.3	-2.2	5.6	0.78	1.6	One sample t-test	t = -2.9	52	51	0.0054	0.40
FR change standard shell adapting	5B	0.64	0.89	5.3	0.50	0.98	One sample Wilcoxon test	V = 3760	113	NA	0.035	0.20
FR change cascade shell adapting	5B	0.50	0.44	7.3	0.68	1.4	One sample t-test	t = 0.74	113	112	0.46	0.069
FR change deviant shell adapting	5B	-1.8	-1.3	7.4	0.69	1.4	One sample Wilcoxon test	V = 2040	113	NA	0.0057	0.26
FR change standard central facilitating	5C	-6.3	-7.3	5.8	1.6	3.4	One sample t-test	t = -4.1	14	13	0.0013	1.1
FR change cascade central facilitating	5C	-0.44	-0.89	4.1	1.1	2.4	One sample t-test	t = -0.40	14	13	0.69	0.11
FR change deviant central facilitating	5C	1.5	1.3	3.4	0.92	2.0	One sample t-test	t = 1.7	14	13	0.12	0.45
FR change standard shell facilitating	5D	-2.7	-3.1	5.4	0.87	1.8	One sample t-test	t = -3.1	38	37	0.0042	0.50
FR change cascade shell facilitating	5D	0.36	0.44	5.1	0.84	1.7	One sample t-test	t = 0.43	38	37	0.67	0.070
FR change deviant shell facilitating	5D	2.6	2.7	4.5	0.74	1.5	One sample t-test	t = 3.5	38	37	0.0013	0.57
FR change standard central non-adapting	5E	-2.5	-2.2	6.2	0.50	0.99	One sample Wilcoxon test	V = 2995	155	NA	1.4e-06	0.38
FR change cascade central non-adapting	5E	-0.68	-0.44	6.3	0.51	1.0	One sample t-test	t = -1.3	155	154	0.18	0.11
FR change deviant central non-adapting	5E	0.57	0.0	5.8	0.47	0.93	One sample t-test	t = 1.2	155	154	0.22	0.098
FR change standard shell non-adapting	5F	-0.63	-0.44	5.3	0.34	0.68	One sample Wilcoxon test	V = 11050	243	NA	0.035	0.14

Comparison	Figure	Mean	Median	SD	SEM	CI (±)	Test	Test statistic	N	df	p	Effect size
FR change cascade shell non-adapting	5F	-0.51	-0.44	5.1	0.32	0.64	One sample Wilcoxon test	V = 12157	243	NA	0.15	0.089
FR change deviant shell non-adapting	5F	-0.059	0.0	5.0	0.32	0.64	One sample t-test	t = -0.18	243	242	0.86	0.012
FR central facilitating (first vs. last standard)	6C	First: 31 Last: 36	First: 29 Last: 31	First: 15 Last: 16	First: 3.9 Last: 4.4	First: 8.5 Last: 9.5	Wilcoxon signed rank test	V = 0	14	NA	0.0017	0.87
FR shell facilitating (first vs. last standard)	6D	First: 53 Last: 57	First: 38 Last: 42	First: 38 Last: 42	First: 6.2 Last: 6.8	First: 13 Last: 14	Wilcoxon signed rank test	V = 92	38	NA	9.3e-05	0.64
FR central adapting (cascade vs. many standards)	3S2B (left)	Casc: 61 MS: 63	Casc: 50 MS: 52	Casc: 38 MS: 40	Casc: 5.2 MS: 5.6	Casc: 10 MS: 11	Wilcoxon signed rank test	V = 595	52	NA	0.39	0.12
FR central facilitating (cascade vs. many standards)	3S2B (right)	Casc: 29 MS: 31	Casc: 26 MS: 28	Casc: 14 MS: 16	Casc: 3.8 MS: 4.3	Casc: 8.2 MS: 9.3	Wilcoxon signed rank test	V = 41	14	NA	0.49	0.19
FR shell adapting (cascade vs. many standards)	3S2C (left)	Casc: 64 MS: 66	Casc: 43 MS: 41	Casc: 61 MS: 68	Casc: 5.7 MS: 6.4	Casc: 11 MS: 13	Wilcoxon signed rank test	V = 2653	113	NA	0.46	0.064
FR shell facilitating (cascade vs. many standards)	3S2C (right)	Casc: 43 MS: 45	Casc: 24 MS: 28	Casc: 41 MS: 52	Casc: 6.6 MS: 8.4	Casc: 13 MS: 17	Wilcoxon signed rank test	V = 264.5	38	NA	0.41	0.14

807   Table 2: Statistical comparisons for control data.

Comparison	Figure	Mean	Median	SD	SEM	CI (±)	Test	Test statistic	N	df	p	Effect size
iMM central (control vs. experimental)	3S1B (left)	Con: 0.092 Exp: 0.057	Con: 0.086 Exp: 0.064	Con: 0.16 Exp: 0.18	Con: 0.011 Exp: 0.012	Con: 0.022 Exp: 0.024	Wilcoxon rank sum test	W = 7919	77 (control) 221 (exp.)	NA	0.37	0.052
iMM shell (control vs. experimental)	3S1B (right)	Con: 0.083 Exp: 0.073	Con: 0.069 Exp: 0.053	Con: 0.23 Exp: 0.24	Con: 0.012 Exp: 0.012	Con: 0.023 Exp: 0.024	Wilcoxon rank sum test	W = 22364	119 (control) 394 (exp.)	NA	0.45	0.034
iMM central adapting (laser OFF vs. ON)	3S1C (top)	OFF: 0.35 ON: 0.33	OFF: 0.35 ON: 0.32	OFF: 0.11 ON: 0.15	OFF: 0.026 ON: 0.034	OFF: 0.054 ON: 0.072	Wilcoxon signed rank test	V = 124	18	NA	0.099	0.40
iPE central adapting (laser OFF vs. ON)	3S1C (middle)	OFF: 0.16 ON: 0.19	OFF: 0.10 ON: 0.081	OFF: 0.39 ON: 0.40	OFF: 0.091 ON: 0.094	OFF: 0.19 ON: 0.20	Paired t-test	t = -1.1	18	17	0.30	0.25
iRS central adapting (laser OFF vs. ON)	3S1C (bottom)	OFF: 0.19 ON: 0.14	OFF: 0.24 ON: 0.14	OFF: 0.38 ON: 0.37	OFF: 0.090 ON: 0.087	OFF: 0.19 ON: 0.18	Paired t-test	t = 1.9	18	17	0.077	0.44
iMM shell adapting (laser OFF vs. ON)	3S1D (top)	OFF: 0.38 ON: 0.38	OFF: 0.35 ON: 0.38	OFF: 0.19 ON: 0.22	OFF: 0.032 ON: 0.037	OFF: 0.065 ON: 0.075	Paired t-test	t = -0.0013	35	34	0.99	0.00022

iPE shell adapting (laser OFF vs. ON)	3S1D (middle)	OFF: 0.16 ON: 0.14	OFF: 0.12 ON: 0.15	OFF: 0.24 ON: 0.26	OFF: 0.041 ON: 0.044	OFF: 0.083 ON: 0.090	Paired t-test	t = 0.58	35	34	0.56	0.099
iRS shell adapting (laser OFF vs. ON)	3S1D (bottom)	OFF: 0.22 ON: 0.24	OFF: 0.24 ON: 0.20	OFF: 0.23 ON: 0.22	OFF: 0.040 ON: 0.038	OFF: 0.081 ON: 0.077	Paired t-test	t = -0.78	35	34	0.44	0.13
iMM central facilitating (laser OFF vs. ON)	3S1E (top)	OFF: -0.37 ON: -0.33	OFF: -0.36 ON: -0.37	OFF: 0.15 ON: 0.18	OFF: 0.077 ON: 0.090	OFF: 0.25 ON: 0.29	Paired t-test	t = -1.1	4	3	0.34	0.57
iPE central facilitating (laser OFF vs. ON)	3S1E (middle)	OFF: -0.043 ON: 0.030	OFF: -0.0047 ON: 0.077	OFF: 0.47 ON: 0.45	OFF: 0.24 ON: 0.22	OFF: 0.75 ON: 0.71	Paired t-test	t = -0.93	4	3	0.42	0.47
iRS central facilitating (laser OFF vs. ON)	3S1E (bottom)	OFF: -0.33 ON: -0.36	OFF: -0.49 ON: -0.53	OFF: 0.55 ON: 0.60	OFF: 0.27 ON: 0.30	OFF: 0.87 ON: 0.95	Paired t-test	t = 0.49	4	3	0.66	0.24
iMM shell facilitating (laser OFF vs. ON)	3S1F (top)	OFF: -0.38 ON: -0.31	OFF: -0.32 ON: -0.30	OFF: 0.22 ON: 0.20	OFF: 0.048 ON: 0.043	OFF: 0.10 ON: 0.090	Wilcoxon signed rank test	V = 63	21	NA	0.070	0.40
iPE shell facilitating (laser OFF vs. ON)	3S1F (middle)	OFF: -0.090 ON: -0.094	OFF: -0.11 ON: -0.081	OFF: 0.18 ON: 0.20	OFF: 0.040 ON: 0.044	OFF: 0.083 ON: 0.093	Wilcoxon signed rank test	V = 109	21	NA	0.84	0.050
iRS shell facilitating (laser OFF vs. ON)	3S1F (bottom)	OFF: -0.29 ON: -0.21	OFF: -0.28 ON: -0.15	OFF: 0.24 ON: 0.21	OFF: 0.053 ON: 0.047	OFF: 0.11 ON: 0.097	Paired t-test	t = -1.8	21	20	0.091	0.39
iMM central non-adapting (laser OFF vs. ON)	3S1G (top)	OFF: 0.021 ON: 0.060	OFF: 0.014 ON: 0.050	OFF: 0.24 ON: 0.23	OFF: 0.032 ON: 0.031	OFF: 0.064 ON: 0.063	Paired t-test	t = -1.8	55	54	0.075	0.24
iPE central non-adapting (laser OFF vs. ON)	3S1G (middle)	OFF: 0.12 ON: 0.14	OFF: 0.034 ON: 0.092	OFF: 0.34 ON: 0.35	OFF: 0.046 ON: 0.047	OFF: 0.092 ON: 0.095	Paired t-test	t = -1.2	55	54	0.23	0.16
iRS central non-adapting (laser OFF vs. ON)	3S1G (bottom)	OFF: -0.095 ON: -0.083	OFF: -0.064 ON: -0.072	OFF: 0.31 ON: 0.29	OFF: 0.042 ON: 0.038	OFF: 0.084 ON: 0.077	Paired t-test	t = -0.57	55	54	0.57	0.077
iMM shell non-adapting (laser OFF vs. ON)	3S1H (top)	OFF: 0.063 ON: 0.051	OFF: 0.040 ON: 0.031	OFF: 0.16 ON: 0.22	OFF: 0.021 ON: 0.027	OFF: 0.042 ON: 0.054	Wilcoxon signed rank test	V = 1133	63	NA	0.39	0.11
iPE shell non-adapting (laser OFF vs. ON)	3S1H (middle)	OFF: 0.053 ON: 0.027	OFF: 0.0 ON: 0.0	OFF: 0.25 ON: 0.26	OFF: 0.031 ON: 0.032	OFF: 0.063 ON: 0.065	Paired t-test	t = 0.88	63	62	0.38	0.11
iRS shell non-adapting (laser OFF vs. ON)	3S1H (bottom)	OFF: 0.011 ON: 0.024	OFF: 0.028 ON: 0.041	OFF: 0.27 ON: 0.28	OFF: 0.034 ON: 0.035	OFF: 0.068 ON: 0.071	Paired t-test	t = -0.43	63	62	0.67	0.054

REFERENCES

Aitkin, L. M., Webster, W. R., Veale, J. L., & Crosby, D. C. (1975). Inferior colliculus. I.

Comparison of response properties of neurons in central, pericentral, and external nuclei of

adult cat. *Journal of Neurophysiology*, *38*(5), 1196–1207.

Anderson, L A, & Malmierca, M. S. (2013). The effect of auditory cortex deactivation on stimulus-

specific adaptation in the inferior colliculus of the rat. *European Journal of Neuroscience*,

37(1), 52–62.

Anderson, Lucy A, Christianson, G. B., & Linden, J. F. (2009). Stimulus-specific adaptation occurs

in the auditory thalamus. *Journal of Neuroscience*, *29*(22), 7359–7363.

Antunes, F. M., & Malmierca, M. S. (2011). Effect of auditory cortex deactivation on stimulus-

specific adaptation in the medial geniculate body. *Journal of Neuroscience*, *31*(47), 17306–

17316.

Antunes, F. M., Nelken, I., Covey, E., & Malmierca, M. S. (2010). Stimulus-specific adaptation in

the auditory thalamus of the anesthetized rat. *PLoS One*, *5*(11), e14071.

Asilador, A., & Llano, D. A. (2020). Top-down inference in the auditory system: Potential roles for

corticofugal projections. *Frontiers in Neural Circuits*, *14*.

Auksztulewicz, R., & Friston, K. (2016). Repetition suppression and its contextual determinants in

predictive coding. *Cortex*, *80*, 125–140.

Bajo, V. M., Nodal, F. R., Bizley, J. K., Moore, D. R., & King, A. J. (2007). The ferret auditory

cortex: descending projections to the inferior colliculus. *Cerebral Cortex*, *17*(2), 475–491.

Bastos, A. M., Usrey, W. M., Adams, R. A., Mangun, G. R., Fries, P., & Friston, K. J. (2012).

Canonical microcircuits for predictive coding. *Neuron*, *76*(4), 695–711.

Blackwell, J. M., Lesicko, A., Rao, W., De Biasi, M., & Geffen, M. N. (2020). Auditory cortex

shapes sound responses in the inferior colliculus. *ELife*, *9*. https://doi.org/10.7554/eLife.51890

Brugge, J. F., & Merzenich, M. M. (1973). Responses of neurons in auditory cortex of the macaque

monkey to monaural and binaural stimulation. *Journal of Neurophysiology*, *36*(6), 1138–1158.

Bulkin, D. A., & Groh, J. M. (2011). Systematic mapping of the monkey inferior colliculus reveals

enhanced low frequency sound representation. *Journal of Neurophysiology*, *105*(4), 1785–

1797.

Cai, R., Richardson, B. D., & Caspary, D. M. (2016). Responses to predictable versus random

temporally complex stimuli from single units in auditory thalamus: impact of aging and

anesthesia. *Journal of Neuroscience*, *36*(41), 10696–10706.

Casado-Román, L., Carbajal, G. V, Pérez-González, D., & Malmierca, M. S. (2020). Prediction

error signaling explains neuronal mismatch responses in the medial prefrontal cortex. *PLoS Biology*, *18*(12), e3001019.

Chen, C., Rodriguez, F. C., Read, H., & Escabí, M. A. (2012). Spectrotemporal sound preferences of neighboring inferior colliculus neurons: implications for local circuitry and processing. *Frontiers in Neural Circuits*, *6*, 62.

De Franceschi, G., & Barkat, T. R. (2020). Task-induced modulations of neuronal activity along the auditory pathway. *BioRxiv*.

De Gardelle, V., Waszczuk, M., Egner, T., & Summerfield, C. (2013). Concurrent repetition enhancement and suppression responses in extrastriate visual cortex. *Cerebral Cortex*, *23*(9), 2235–2244.

Duque, D., & Malmierca, M. S. (2015). Stimulus-specific adaptation in the inferior colliculus of the mouse: anesthesia and spontaneous activity effects. *Brain Structure and Function*, *220*(6), 3385–3398.

Duque, D., Pérez-González, D., Ayala, Y. A., Palmer, A. R., & Malmierca, M. S. (2012). Topographic distribution, frequency, and intensity dependence of stimulus-specific adaptation in the inferior colliculus of the rat. *Journal of Neuroscience*, *32*(49), 17762–17774.

English, J. G., & Roth, B. L. (2015). Chemogenetics—a transformational and translational platform. *JAMA Neurology*, *72*(11), 1361–1366.

Escera, C., & Malmierca, M. S. (2014). The auditory novelty system: an attempt to integrate human and animal research. *Psychophysiology*, *51*(2), 111–123.

Fontanini, A., & Katz, D. B. (2008). Behavioral states, network states, and sensory response variability. *Journal of Neurophysiology*, *100*(3), 1160–1168.

Friston, K. (2009). The free-energy principle: a rough guide to the brain? *Trends in Cognitive Sciences*, *13*(7), 293–301.

Friston, K., & Kiebel, S. (2009). Predictive coding under the free-energy principle. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *364*(1521), 1211–1221.

Gaese, B. H., & Ostwald, J. (2001). Anesthesia changes frequency tuning of neurons in the rat primary auditory cortex. *Journal of Neurophysiology*, *86*(2), 1062–1066.

Han, X., Chow, B. Y., Zhou, H., Klapoetke, N. C., Chuong, A., Rajimehr, R., Yang, A., Baratta, M. V, Winkle, J., & Desimone, R. (2011). A high-light sensitivity optical neural silencer: development and application to optogenetic control of non-human primate cortex. *Frontiers in Systems Neuroscience*, *5*, 18.

Harms, L., Fulham, W. R., Todd, J., Budd, T. W., Hunter, M., Meehan, C., Penttonen, M., Schall,

U., Zavitsanou, K., & Hodgson, D. M. (2014). Mismatch negativity (MMN) in freely-moving

rats with several experimental controls. *PloS One*, *9*(10), e110892.

Herbert, H., Aschoff, A., & Ostwald, J. (1991). Topography of projections from the auditory cortex

to the inferior colliculus in the rat. *Journal of Comparative Neurology*, *304*(1), 103–122.

Herrmann, B., Henry, M. J., Fromboluti, E. K., McAuley, J. D., & Obleser, J. (2015). Statistical

context shapes stimulus-specific adaptation in human auditory cortex. *Journal of*

Neurophysiology, *113*(7), 2582–2591.

Jaramillo, S., Borges, K., & Zador, A. M. (2014). Auditory thalamus and auditory cortex are equally

modulated by context during flexible categorization of sounds. *Journal of Neuroscience*,

34(15), 5291–5301.

Johnson, K. R., Tian, C., Gagnon, L. H., Jiang, H., Ding, D., & Salvi, R. (2017). Effects of Cdh23

single nucleotide substitutions on age-related hearing loss in C57BL/6 and 129S1/Sv mice and

comparisons with congenic strains. *Scientific Reports*, *7*(1), 1–13.

KATSUKI, Y., MURATA, K., SUGA, N., & TAKENAKA, T. (1959). Electrical activity of cortical

auditory neurons of unanaesthetized and unrestrained cat. *Proceedings of the Japan Academy*,

35(9), 571–574.

Kommajosyula, S. P., Bartlett, E. L., Cai, R., Ling, L., & Caspary, D. M. (2021). Corticothalamic

Projections Deliver Enhanced-Responses to Medial Geniculate Body as a Function of the

Temporal Reliability of the Stimulus. *BioRxiv*.

Kommajosyula, S. P., Cai, R., Bartlett, E., & Caspary, D. M. (2019). Top-down or bottom up:

decreased stimulus salience increases responses to predictable stimuli of auditory thalamic

neurons. *The Journal of Physiology*, *597*(10), 2767–2784.

Lesicko, A.M.H., Hristova, T. S., Maigler, K. C., & Llano, D. A. (2016). Connectional modularity

of top-down and bottom-up multimodal inputs to the lateral cortex of the mouse inferior

colliculus. *Journal of Neuroscience*, *36*(43). https://doi.org/10.1523/JNEUROSCI.4134-

15.2016

Lesicko, Alexandria M H, & Llano, D. A. (2020). Circuit mechanisms underlying the segregation

and integration of parallel processing streams in the inferior colliculus. *Journal of*

Neuroscience, *40*(33), 6328–6344.

Malmierca, M. S., Cristaudo, S., Pérez-González, D., & Covey, E. (2009). Stimulus-specific

adaptation in the inferior colliculus of the anesthetized rat. *Journal of Neuroscience*, *29*(17),

5483–5493.

Malmierca, M. S., Izquierdo, M. A., Cristaudo, S., Hernández, O., Pérez-González, D., Covey, E.,

& Oliver, D. L. (2008). A discontinuous tonotopic organization in the inferior colliculus of the rat. *Journal of Neuroscience*, *28*(18), 4767–4776.

Müller, N. G., Strumpf, H., Scholz, M., Baier, B., & Melloni, L. (2013). Repetition suppression versus enhancement—it's quantity that matters. *Cerebral Cortex*, *23*(2), 315–322.

Natan, R. G., Briguglio, J. J., Mwilambwe-Tshilobo, L., Jones, S. I., Aizenberg, M., Goldberg, E. M., & Geffen, M. N. (2015). Complementary control of sensory adaptation by two types of cortical interneurons. *Elife*, *4*, e09868.

Nelken, I., & Ulanovsky, N. (2007). Mismatch negativity and stimulus-specific adaptation in animal models. *Journal of Psychophysiology*, *21*(3–4), 214–223.

Okada, K., Matchin, W., & Hickok, G. (2018). Neural evidence for predictive coding in auditory cortex during speech production. *Psychonomic Bulletin & Review*, *25*(1), 423–430.

Pakan, J. M. P., Lowe, S. C., Dylda, E., Keemink, S. W., Currie, S. P., Coutts, C. A., & Rochefort, N. L. (2016). Behavioral-state modulation of inhibition is context-dependent and cell type specific in mouse visual cortex. *Elife*, *5*, e14985.

Parras, G. G., Casado-Román, L., Schröger, E., & Malmierca, M. S. (2021). The posterior auditory field is the chief generator of prediction error signals in the auditory cortex. *NeuroImage*, 118446.

Parras, G. G., Nieto-Diego, J., Carbajal, G. V, Valdés-Baizabal, C., Escera, C., & Malmierca, M. S. (2017). Neurons along the auditory pathway exhibit a hierarchical organization of prediction error. *Nature Communications*, *8*(1), 1–17.

Paxinos, G., & Franklin, K. B. J. (2019). *Paxinos and Franklin's the mouse brain in stereotaxic coordinates*. Academic press.

Rao, R. P. N., & Ballard, D. H. (1999). Predictive coding in the visual cortex: a functional interpretation of some extra-classical receptive-field effects. *Nature Neuroscience*, *2*(1), 79–87.

Rauss, K., Schwartz, S., & Pourtois, G. (2011). Top-down effects on early visual processing in humans: A predictive coding framework. *Neuroscience & Biobehavioral Reviews*, *35*(5), 1237–1253.

Ress, D., & Chandrasekaran, B. (2013). Tonotopic organization in the depth of human inferior colliculus. *Frontiers in Human Neuroscience*, *7*, 586.

Ruhnau, P., Herrmann, B., & Schröger, E. (2012). Finding the right control: the mismatch negativity under investigation. *Clinical Neurophysiology*, *123*(3), 507–512.

Rummell, B. P., Klee, J. L., & Sigurdsson, T. (2016). Attenuation of responses to self-generated sounds in auditory cortical neurons. *Journal of Neuroscience*, *36*(47), 12010–12026.

Saldaña, E., Feliciano, M., & Mugnaini, E. (1996). Distribution of descending projections from
primary auditory neocortex to inferior colliculus mimics the topography of intracollicular
projections. *Journal of Comparative Neurology*, *371*(1), 15–40.

Saldaña, E., & Merchań, M. A. (1992). Intrinsic and commissural connections of the rat inferior
colliculus. *Journal of Comparative Neurology*, *319*(3), 417–437.

Saldaña, E., & Merchán, M. A. (2005). Intrinsic and commissural connections of the inferior
colliculus. In *The inferior colliculus* (pp. 155–181). Springer.

Schellekens, W., Van Wezel, R. J. A., Petridou, N., Ramsey, N. F., & Raemaekers, M. (2016).
Predictive coding for motion stimuli in human early visual cortex. *Brain Structure and*
Function, *221*(2), 879–890.

Schofield, B. R. (2009). Projections to the inferior colliculus from layer VI cells of auditory cortex.
Neuroscience, *159*(1), 246–258.

Schumacher, J. W., Schneider, D. M., & Woolley, S. M. N. (2011). Anesthetic state modulates
excitability but not spectral tuning or neural discrimination in single auditory midbrain
neurons. *Journal of Neurophysiology*, *106*(2), 500–514.

Segaert, K., Weber, K., de Lange, F. P., Petersson, K. M., & Hagoort, P. (2013). The suppression of
repetition enhancement: a review of fMRI studies. *Neuropsychologia*, *51*(1), 59–66.

Shipp, S. (2016). Neural elements for predictive coding. *Frontiers in Psychology*, *7*, 1792.

Shipp, S., Adams, R. A., & Friston, K. J. (2013). Reflections on agranular architecture: predictive
coding in the motor cortex. *Trends in Neurosciences*, *36*(12), 706–716.

Stebbings, K. A., Lesicko, A. M. H., & Llano, D. A. (2014). The auditory corticocollicular system:
Molecular and circuit-level considerations. *Hearing Research*, *314*.
https://doi.org/10.1016/j.heares.2014.05.004

Stiebler, I., & Ehret, G. (1985). Inferior colliculus of the house mouse. I. A quantitative study of
tonotopic organization, frequency representation, and tone-threshold distribution. *Journal of*
Comparative Neurology, *238*(1), 65–76.

Syka, J., Popelář, J., Kvašňák, E., & Astl, J. (2000). Response properties of neurons in the central
nucleus and external and dorsal cortices of the inferior colliculus in guinea pig. *Experimental*
Brain Research, *133*(2), 254–266.

Taaseh, N., Yaron, A., & Nelken, I. (2011). Stimulus-specific adaptation and deviance detection in
the rat auditory cortex. *PloS One*, *6*(8), e23369.

Takesian, A. E., Bogart, L. J., Lichtman, J. W., & Hensch, T. K. (2018). Inhibitory circuit gating of
auditory critical-period plasticity. *Nature Neuroscience*, *21*(2), 218–227.

Torii, M., Hackett, T. A., Rakic, P., Levitt, P., & Polley, D. B. (2013). EphA signaling impacts
975 development of topographic connectivity in auditory corticofugal systems. *Cerebral Cortex*,
976 *23*(4), 775–785.

977 Tyssowski, K. M., & Gray, J. M. (2019). Blue light increases neuronal activity-regulated gene
978 expression in the absence of optogenetic proteins. *ENeuro*, *6*(5).

979 Ulanovsky, N., Las, L., & Nelken, I. (2003). Processing of low-probability sounds by cortical
980 neurons. *Nature Neuroscience*, *6*(4), 391–398.

981 Weissbart, H., Kandylaki, K. D., & Reichenbach, T. (2020). Cortical tracking of surprisal during
982 continuous speech comprehension. *Journal of Cognitive Neuroscience*, *32*(1), 155–166.

983 Yaron, A., Hershenhoren, I., & Nelken, I. (2012). Sensitivity to complex statistical regularities in rat
984 auditory cortex. *Neuron*, *76*(3), 603–615.

985 Yudintsev, G., Asilador, A., Coppinger, M., Nair, K., Prasad, M., & Llano, D. A. (2019).
986 Connectional heterogeneity in the mouse auditory corticocollicular system. *BioRxiv*, 571711.

987 Zhai, X., Khatami, F., Sadeghi, M., He, F., Read, H. L., Stevenson, I. H., & Escabí, M. A. (2020).
988 Distinct neural ensemble response statistics are associated with recognition and discrimination
989 of natural sound textures. *Proceedings of the National Academy of Sciences*, *117*(49), 31482–
990 31493.