Title

Two tales of one neural link predict blind individual's Braille reading proficiency

Authors

Ruxue Wang[1†], Jiangtao Gong[2,3†], Chenying Zhao[1], Yingqing Xu[2,3*] and Bo Hong[1,4*]

[1]Department of Biomedical Engineering, School of Medicine, Tsinghua University,

Beijing, 100084, P.R. China;

[2]The Future Laboratory, Tsinghua University, Beijing, 100084, P.R. China;

[3]Academy of Arts & Design, Tsinghua University, Beijing, 100084, P.R. China;

[4]IDG/McGovern Institute for Brain Research, Tsinghua University, Beijing, 100084,

P.R. China

[†]R.W. and J. G. contributed equally to this work.

*Correspondence and requests for materials should be addressed to Bo Hong (email:

hongbo@tsinghua.edu.cn) or to Yingqing Xu (email: yqxu@tsinghua.edu.cn).

Number of pages: 50

Number of figures: 6

Number of tables: 2

Number of words for Abstract: 177

Number of words for Significance Statement:114

Number of words for Introduction: 583

Number of words for Discussion: 1482

Conflict of Interest: We claim no conflict of interest

Acknowledgements: This work is supported by the National Science Foundation of

China [grant number NSFC 61621136008 to B.H.] and the National Key R&D Program

of China [grant number 2017YFA0205904 B.H.] We thank Brigitte Röder at the

University of Hamburg, Ning Guo, and Fang Cai at Tsinghua University for comments

on the manuscript.

Data and code availability:

All data reported in this paper will be available by the corresponding author upon

request.

All original code has been deposited at Github and is publicly available as of the date

of publication.

Abstract

Natural Braille reading, a demanding cognitive skill, poses a huge challenge for the

brain network of the blind. Here, with behavioral measurement and functional MRI

imaging data, we pinpointed the neural pathway and investigated the neural

mechanisms of individual differences in Braille reading in late blindness. Using resting

state fMRI, we identified a distinct neural link between the higher-tier 'visual' cortex—

the lateral occipital cortex (LOC), and the inferior frontal cortex (IFC) in the late blind

brain, which is significantly stronger than sighted controls. Individual Braille reading

proficiency positively correlated with the left-lateralized LOC-IFC functional

connectivity. In a natural Braille reading task, we found an enhanced bidirectional

information flow with a stronger top-down modulation of the IFC-to-LOC effective

connectivity. Greater top-down modulation contributed to higher Braille reading

proficiency via a broader area of task-engaged LOC. Together, we established a model

to predict Braille reading proficiency, considering both functional and effective

connectivity of the LOC-IFC pathway. This 'two-tale' model suggests that developing

the underpinning neural circuit and the top-down cognitive strategy contributes

uniquely to superior Braille reading performance.

SIGNIFICANCE STATEMENT

For late blind humans, one of the most challenging cognitive skills is natural Braille reading. However, little is known about the neural mechanisms of significant differences in individual Braille reading performance. Using functional imaging data, we identified a distinct neural link between the left lateral occipital cortex (LOC) and the left inferior frontal cortex (IFC) for natural Braille reading in the late blind brain. To better predict individual Braille reading proficiency, we proposed a linear model with two variables of the LOC-IFC link: the resting-state functional connectivity and the task-engaged top-down effective connectivity. These findings suggest that developing the underpinning neural pathway and the top-down cognitive strategy contributes uniquely to superior Braille reading performance.

Introduction

For blind humans, one of the most demanding cognitive skills is natural Braille reading, which makes a big difference to their social well-being (Ryles, 1996, 2000; Schroeder, 1996; Marshall and Moys, 2020). They decode the tactile dots information into meaningful patterns with semantic properties and ultimately achieve content reading (Sadato, 2005). Learning Braille reading poses a critical challenge to the blind brain, especially to late blind people. Unlike congenital blindness, late blind individuals who lost sight several years after birth may have experienced traditional print early in life and get exposed to Braille only later. Such a huge gap between visual and tactile reading requires more effort for them to learn Braille reading (Trent and Truan, 1997). Moreover, not all blind learners excel at their Braille reading; in fact, they exhibit wide disparities in Braille reading proficiency (Coppins and Barlow-Brown, 2006). However, the neural mechanisms of individual differences in natural Braille reading performance in late blindness remain largely unknown.

Previous studies have reported the recruitment of the somatosensory cortex, the higher-order frontal language system, and the 'visual' system during Braille reading in the blind, including late blind humans (Pascual-Leone et al., 2005; Sadato, 2005). This 'visual' system stretched from the primary 'visual' cortex (V1) to the higher-tier 'visual' cortex—the lateral occipital cortex (LOC), which approximately included the consistent region as the typical visual word form area in the sighted (Reich et al., 2011; Kim et al., 2017). Moreover, the most frequently found functional reorganization of the

blind brain was the functional connectivity between the 'visual' cortex and the inferior

frontal cortex (IFC) (Liu et al., 2007; Sabbah et al., 2016; Bedny, 2017), which hinted

its role in higher-cognitive functions, including Braille reading. However, previous

studies leave open three important questions. First, for people with a late acquisition of

blindness and Braille reading, what is the specific 'visual' region playing the essential

role in bridging the tactile and language-related processing in natural Braille reading?

Furthermore, employing either only Braille letter (Cohen et al., 1997) or word

recognition (Sadato et al., 1998; Katarzyna Rączy et al., 2019), prior findings have

limited functional inference of the dynamic neural interactions under real-life, natural

Braille reading scenario. Thus, how does the information flow between the specific

'visual' region and the IFC during natural Braille reading? Finally, how can such neural

interactions determine the Braille reading proficiency of late blind individuals? In other

cognitive behaviors, individual performance has been suggested to correlate with the

specific neural connectivity (Jung et al., 2018; Ren et al., 2018). Therefore, we

hypothesized that the strength of the distinct resting-state functional connectivity and

the task-engaged effective connectivity between the 'visual' cortex and the frontal

cortex would jointly predict Braille reading proficiency of late blind individuals.

To answer these questions, we tested individual differences in Braille reading

proficiency and investigated brain signatures from both resting and task-engaged

functional magnetic resonance imaging (fMRI) acquisition in late blindness. We used

natural Braille articles in this study to resemble real-life natural Braille reading

(Figure 1). We first examined whether the V1 or the LOC in the late blind brain

developed resting-state functional connectivity with the IFC compared to sighted

controls. We then employed dynamic causal modeling (DCM) to infer their effective

connectivity in a natural Braille reading task. To test our hypothesis, we examined how

individual Braille reading proficiency was linked with the distinct 'visual'-frontal

neural connectivity. Finally, a neural predictive model was established to further reveal

the neural mechanisms of individual differences in Braille reading proficiency.

Materials and Methods

Experimental subjects

A total of 39 blind subjects (mean age $\pm$ s.d. = 21.78 $\pm$ 1.96 years; 15 females)

participated in the Braille reading performance test. They were all Braille readers. A

subgroup of 16 right-handed blind subjects with late-onset acquired blindness (Table 1,

mean age $\pm$ s.d. = 22.43 $\pm$ 1.86 years; mean age at blindness onset $\pm$ s.d. = 9.42 $\pm$ 4.32

years; 8 females) agreed to participate in the fMRI scan. Fifteen education-matched

sighted controls also participated in the fMRI scan. One sighted subject with excessive

head motion (mean framewise displacement [FD (Power et al., 2012); > 2 SD from the

group mean]) during fMRI acquisition was excluded from further analyses, leaving 14

sighted subjects (mean age $\pm$ s.d. = 21.73 $\pm$ 1.07 years; 7 females). The two groups were

matched in age (two-sample T-test, n = 16 in the late blind group and n = 14 in the

sighted group, T-value = 1.248, 95% CI = [-0.452 1.862], DF = 28, p = 0.222, Cohen's

d = 0.457, two-tailed). Sighted subjects had normal or corrected-to-normal vision and

had never learned Braille. All subjects had no history of neurological disorders. The

study was approved by the Institutional Review Board of Tsinghua University, and

informed consent was obtained from each participant.

Acquisition of biographical data on Braille reading practice

According to previous studies (Bengtsson et al., 2005; Liu et al., 2007), we obtained

the number of hours of practicing Braille reading for each blind subject during

childhood (from the start of practicing until 11 years), adolescence (12–16 years) and

adulthood (17 years until the time of the magnetic resonance scan). These values were

calculated from biographical data collected from all blind participants on the key events

from the beginning of Braille reading training until the scan and on their self-estimated

number of hours of practicing within the periods between these key time points (Table

1).

Braille reading performance test

Previous Braille reading studies have demonstrated that passage length and content

validity might influence the reliability and validity of the behavioral assessment (Mason,

2012). The content validity of Braille materials concerns the readability and difficulty

for subjects. Given that all of our blind subjects had passed the special college entrance

examination for blind individuals, we selected two natural Braille articles at the high

school level, slightly below their current levels. Each natural Braille article consisted

of approximately 500 Braille words. All reading materials were proofread by an

experienced Braille reader and were printed by a Braille printer from VIEWPLUS

(Viewplus.com) on Braille paper, measuring 27.94 cm in length and 24.89 cm in width.

The task was to read the two natural Braille articles aloud as accurately and quickly as

possible. The reading strategy was not specified. We tested individual Braille reading

proficiency by measuring the number of Braille words read correctly per minute on

average. Individual Braille reading accuracy was also assessed by calculating the

proportion of Braille words read correctly to all the words in the two Braille articles.

Blind individuals with a reading accuracy reaching 99.5% were recruited for this study

to ensure content comprehension.

Functional MRI parameters

All MR images were acquired from a Philip Achieva 3.0 Tesla TX MR scanner with a

32-channel head coil. The participants laid supine with their heads snugly fixed with

straps and foam pads to minimize head movement. Structural images were acquired

using a sagittal magnetization-prepared rapid gradient-echo T1-weighted sequence (TR

= 7.6 ms, TE = 3.7 ms, flip angle = 8°, voxel size = $1 \times 1 \times 1$ mm, FOV = 256, slices =

180). Before the task session, participants underwent two resting-state fMRI scans (8

min per scan) using an echo-planar imaging sequence (TR = 2000 ms, TE = 30 ms, flip

angle = 90°, voxel size = $3.5 \times 3.5 \times 3.5$ mm, FOV = 224, slices = 35). During the

resting-state fMRI scan, the participants were instructed to stay awake. Functional

images for the natural Braille reading task were also acquired using an echo-planar

imaging sequence (TR = 3000 ms, TE = 35 ms, flip angle = 90°, voxel size = $3 \times 3 \times 3$

mm, FOV = 220, slices = 47). All subjects were asked to keep their eyes closed during

the scans.

FMRI task paradigm

A natural Braille reading task with ten blocks was used in this study (Supplementary

Figure 4). In each block, blind subjects were instructed to silently read the page of the

Braille article and the next blank page (control: blank), mainly using the right index

finger at a constant speed, according to the audio cue of "Begin" or "Turn the page"

played in an earphone. Blind individuals were asked to thoroughly read and

comprehend the Braille article during the fMRI scan. All subjects were instructed to

carefully touch and perceive the blank page line by line as if they were reading the

Braille page. Each audio cue lasted for 1 s. We used natural Braille articles to simulate

real-life Braille reading conditions. Blank pages were printed with the same kind and

size of paper as Braille pages.

Resting-state fMRI data preprocessing and analysis

Structural data were processed and reconstructed to individual cortical surface using

FreeSurfer (http://surfer.nmr.mgh.harvard.edu). Individual resting-state data were

preprocessed using the following procedures: (i) discarding of the first four volumes;

(ii) slice-timing correction; (iii) rigid body correction for head motion with the AFNI

package (Cox, 1996); (iv) functional-anatomical image co-registration; (v) bandpass

temporal filtering (0.01-0.08 Hz); and (vi) regressing out of nuisance covariates to

control for physiological effects and head motion (6 head motion parameters and 2

regressors corresponding to the ventricular signal and white matter signal). One blind

subject (B14) with excessive head motion (mean FD (Power et al., 2012) > 2 SD from

the group mean) during resting-state fMRI acquisition was excluded, leaving 15 blind

subjects for resting-state analyses. The preprocessed fMRI data of each subject were

first aligned to the reconstructed individual cortical surface in each hemisphere. Then,

a 6-mm full-width half-maximum (FWHM) Gaussian smoothing kernel was applied to

the fMRI data in the surface space. The smoothed data were then down-sampled to the

FreeSurfer template with 2,562 vertices in each hemisphere using the mri_surf2surf

function in FreeSurfer. The fMRI data of two resting-state scans were concatenated for

further analyses to improve reliability.

The primary visual cortex (V1) was defined as the combination of the occipital pole

and calcarine sulcus in FreeSurfer's automatic anatomical parcellation (aparc2009)

(Destrieux et al., 2010), consistent with the location of V1 in previous studies (Bedny

et al., 2011; Watkins et al., 2012; Butt et al., 2013). In accord with the seed region in

prior functional connectivity analyses of blind and sighted individuals (Bedny et al.,

2011; Watkins et al., 2012; Abboud and Cohen, 2019), we used the lateral occipital

cortex as a higher-tier visual seed located in the anatomical occipital region according

to the FreeSurfer aparc2009 atlas (Destrieux et al., 2010) (encompassing the middle

occipital gyrus and sulcus, anterior occipital sulcus, inferior occipital gyrus and sulcus,

and lateral occipital-temporal sulcus). The V1 and the LOC were used as the seeds for

resting-state functional connectivity (rsFC) analyses, respectively.

To obtain each seed's whole-brain functional connectivity patterns, we computed the

correlation between the mean blood oxygen level-dependent (BOLD) signal of the seed

and the time series of every other vertex in both hemispheres for each subject. These r-

maps were next converted into z-maps using Fisher's r-to-z transformation. All

statistical group-level FC pattern analyses were carried out using the Surface-Based

Data Processing & Analysis for (Resting-State) Brain Imaging software (DPABISurf

1.1) (Yan et al., 2016). A one-sample T-test ($n = 15$, DF = 14 in the blind group and n

= 14, DF = 13 in the sighted group, one-tailed) was performed for individual FC z-maps

to generate significant and positive FC maps (vertexwise $p < 10^{-6}$, uncorrected, cluster

size > 100 mm) of the V1 or LOC seed for each group. Thus, a union mask including

identified significant vertices in either the blind or sighted group was created for further

FC pattern comparison between the two groups (Wang et al., 2015). Recent works have

consistently suggested that the influence of head motion on intrinsic functional

connectivity in the analysis of between-group functional connectivity comparison

needs to be further scrutinized regardless of the correction one employs at the individual

level (Power et al., 2012; Yan et al., 2013). We, therefore, calculated the mean FD

(Power et al., 2012), describing the mean absolute displacement of each brain volume

relative to the previous volume in translation and rotation in the x, y, and z directions

during the resting-state scan for each subject.

With mean FD as the covariate controlling for the head motion effect (Wang et al.,

2015), we conducted a two-sample T-test ($n = 15$ in the blind group and $n = 14$ in the

sighted group, DF = 27, two-tailed) based on individual FC z-maps within the union

mask to examine how the V1 or LOC FC patterns of the blind group changed compared

to the sighted group. Significant vertices were those surviving the multiple comparison

correction using non-parametric permutation testing (Winkler et al., 2016) (5,000

permutations) and threshold-free cluster enhancement (Chen et al., 2018) ($p < 0.01$,

corrected for both hemispheres, cluster size > 30 mm).

Task fMRI data preprocessing and analysis

The task fMRI data were preprocessed using the FsFast software package

(http://surfer.nmr.mgh.harvard.edu/fswiki/FsFast). The preprocessing procedures

included motion correction, slice timing correction, and boundary-based functional-

anatomical alignment. Like the resting-state fMRI, the task fMRI data of each subject

were registered to the reconstructed individual cortical surface. Next, the fMRI data

were smoothed with a Gaussian filter (FWHM = 6 mm) in the surface space. The

smoothed data were then down-sampled to the FreeSurfer template with 2,562 vertices

in each hemisphere. One blind subject (B6) out of 16 with excessive head motion (mean

FD (Power et al., 2012); > 2 SD from the group mean) during the task fMRI scan was

excluded from further general linear model (GLM) and DCM analyses.

Functional MRI analyses were carried out using Statistical Parametric Mapping toolbox

(SPM12, Welcome Department of Imaging Neuroscience, Institute of Neurology,

London). For the first-level individual analysis, functional images were modeled with

three regressors (one for the Braille condition, one for the blank condition, and another

for the turning page condition which was discarded in the analysis) using the GLM

method. Head motion regression was also performed in the analysis. We defined a

Braille > blank contrast to reveal activated brain regions (significant $p < 0.025$ for both

hemispheres, uncorrected). We did not set an extremely strict threshold here because

these GLM analyses aimed to yield brain regions showing task-relevant effects for

further DCM analyses. However, one blind subject (B16) failing to exhibit any

activated vertices in three of our ROIs above a very liberal threshold ($p < 0.1$,

uncorrected) was excluded, leaving 14 blind subjects for group-level GLM analysis and

DCM analysis. For group-level GLM analysis, data of all the blind subjects were

concatenated and then analyzed by a one-sample T-test ($n = 14$ in the blind group, DF

= 13, one-tailed). Significant brain activations in the blind group were also revealed by

the Braille > blank contrast (Supplementary Figure 4, significant $p < 0.025$ for both

hemispheres, uncorrected).

Dynamic causal modeling (DCM)

All DCM analyses were carried out using SPM12. DCM is a computational framework

for explaining the effective connectivity among brain regions that show experimental

effects (Friston et al., 2003). With the specified model and priors on the coupling and

hemodynamic parameters, posterior probability analysis in DCM determines the most

likely coupling parameters at the neuronal level. We focused on the dynamic

interactions among the IPS, LOC, V1, and IFC in the left hemisphere that were

activated when the blind subjects were reading the natural Braille article. The exact left

V1, LOC, and IFC regions in the resting-state analysis and the anatomical region of the

IPS (intraparietal sulcus, defined in the FreeSurfer aparc2009 atlas (Destrieux et al.,

2010)) were used as anatomical masks for defining ROIs later. Individual ROIs

consisted of significant vertices revealed by the Braille > blank contrast ($p < 0.025$

uncorrected, cluster size > 5 vertices) within the corresponding anatomical masks. The

time series of these ROIs were then extracted for each subject. We slightly dropped the

threshold for two blind subjects (B11: $p < 0.025$ uncorrected; B14: $p < 0.05$ uncorrected,

cluster size > 5 vertices) who lacked strong responses in one or more of the ROIs to

ensure that significant vertices could be identified (Zeidman et al., 2019).

Typically, there are three types of DCM parameters (Friston et al., 2003): 1) endogenous

parameters, reflecting the task-independent effective connectivity among brain regions;

2) driving effects, which quantify how neural responses change due to the task stimuli;

and 3) modulatory strengths, the parameters of greatest interest, which account for the

changes in specific effective connectivity due to the task-relevant conditions. We

constructed bidirectional endogenous connections among the four ROIs in DCM

models based on anatomical knowledge and our resting-state functional connectivity

results. We used the IPS to model the tactile input for the LOC, consistent with the

finding that tactile information from the somatosensory cortex was redirected to the

'visual' systems through the poly-sensory intraparietal cortex in blind people

(Wittenberg et al., 2004; Sadato, 2005; Fujii et al., 2009). We set the modulatory effect

of Braille reading on the connectivity between the left IPS and LOC in accord with a

prior study (Fujii et al., 2009). We then investigated the task modulatory effect of Braille

reading on the connection between the left LOC and the left IFC. Therefore, we

considered that the task-relevant information flow on any direction (bottom-up or top-

down) between these two brain regions might be modulated, resulting in four competing

models (Figure 4A).

We performed BMS across all blind individuals with random-effects analysis to select

the winning model after balancing the explanation of data and the model complexity.

The exceedance probability intuitively presented the relative preference across all

models (Figure 4B). BMA implements a weighted average of parameters in all

competing models where the weights were given by the evidence of each model (Penny

et al., 2010). We then employed BMA to estimate the full posterior density on model

parameters (Penny et al., 2010). Therefore, the significance of each connectivity

parameter could be assessed by the fraction of samples in the posterior density (out of

10,000 sampled data points) that were different from zero (Seghier et al., 2011).

Likewise, the significance of the difference in modulatory strengths on one connection

versus another connection was assessed by the fraction of samples that were higher for

one connection than another (Seghier et al., 2011).

Statistical analysis on the relationship between resting-state functional connectivity and Braille reading behavior data

To investigate the statistical relationship between the left LOC-IFC neural link and Braille reading behavior data of blind individuals, we calculated the Pearson correlation and Spearman correlation coefficients between their functional connectivity (normalized) and Braille reading proficiency, as well as Braille reading practice time within each age periods respectively. In addition, we also performed partial correlations between the left LOC-IFC functional connectivity and Braille reading behavior data with covariates of the onset age of blindness, the current age, and motion in the resting state scan indexed by mean FD (Jung et al., 2018).

Statistical analysis on the relationship between top-down effective connectivity and Braille reading proficiency

The Pearson correlation between top-down modulation of the left IFC-to-LOC effective connectivity in the natural Braille reading task and individual Braille reading proficiency tended to approach significance ($n = 13$, $p = 0.141$, $r = 0.432$, Supplementary Figure 5A). The top-down modulation parameter of DCM analysis provided a precise estimation at the neuronal level of how the rate of change of activity in the IFC influences the rate of change in the LOC activity (Friston et al., 2003) in the natural Braille reading task. Hence, to fully investigate the potential relationship between the top-down modulation and individual Braille reading performance, we investigated whether the area of functional LOC could act as a neural mediator of the

top-down modulation in support of the natural Braille reading task. In accord with our

DCM analysis, the neural activity of the functional LOC in the task was predominantly

influenced by the changes of activity in the IFC through the top-down effective

connectivity. Accordingly, we first identified individual peak vertex revealed by the

Braille > blank contrast as the center of individual functional LOC. Within the

anatomical LOC cortex, individual functional LOC also consisted of vertices whose

functional connectivity (normalized) with the peak vertex reached the threshold. The

area of individual functional LOC was further normalized by calculating the proportion

of these vertices in the anatomical LOC. We initially set a threshold value throughout

the functional connectivity with the peak vertex in 0.2 intervals and eventually chose

the threshold value of 0.8 since only when using this threshold did the area of functional

LOC manifest significant correlation with both top-down modulation (Figure 5A) and

Braille reading proficiency of blind individuals (Figure 5C). However, the bottom-up

modulation of the LOC-to-IFC connectivity was correlated with the area of functional

LOC using none of these threshold values.

Mediation analysis was performed using the "mediation" package (Tingley et al., 2014)

in R (https://www.r-project.org). To test the significance of the mediation effect, we

used the bootstrapping method (simulations: 5000, the bias-corrected and accelerated

confidence intervals were estimated then), which was generally considered more

advantageous.

Analysis of linear neural predictive models for individual Braille reading proficiency

Based on our hypothesis that the resting-state functional connectivity (rsFC) and task-specific top-down effective connectivity (tdEC) of the left LOC-IFC neural connection each explained unique individual variance, we first established a 'two-tale' model linearly combining these two variables to predict Braille reading proficiency of late blind individuals. Note that in the model, the area of functional LOC was used as the neural surrogate of the tdEC (Figure 5D). In this case, this model is given by:

$$y_{profi} = ax_{rsFC} + bx_{tdEC} + e \tag{1}$$

Then, derived from Eq (1), linear models with each of the two brain variables alone, either x_{rsFC} or x_{tdEC}, were fitted separately.

We assessed the linear models' performance in predicting the Braille reading proficiency of a held-out individual using a leave-one-out cross-validation procedure. On each held-out data of this cross-validation procedure, we fitted the model to the training set by minimizing the following cost function:

$$\Theta_{fit} = argmin_{\Theta} \sum_{n=1}^{N} \left(y_{train}^{(n)} - \hat{y}_{train}^{(n)} \right)^2 \tag{2}$$

Here, the $y_{train}^{(n)}$ term refers to the measured Braille reading proficiency of the *n*-th blind subject in the training set ($N = 12$), while the $\hat{y}_{train}^{(n)}$ term refers to the *n*-th individual behavior data estimated by each model during training. This minimization was performed using *fmincon* in MATLAB R2018a. We then applied these model parameters to predict the Braille reading proficiency of the held-out test individual.

393   Finally, the MSE and the Pearson correlation between predicted and corresponding

394   actual proficiency for blind individuals from the cross-validation procedure were

395   calculated for each model.

397   **Results**

399   ---------Figure 1 here---------

Individual differences in Braille reading proficiency

We assessed the Braille reading proficiency of blind individuals using a natural Braille

reading behavioral test. Thirty-nine blind participants (mean age $\pm$ s.d. = 21.78 $\pm$ 1.96

years; 15 females) read two long natural Braille articles (500 Braille words on average)

aloud as accurately and quickly as possible. Results showed that the Braille reading

proficiency of blind individuals varied considerably (Figure 1A, Braille words correctly

read per minute, mean $\pm$ s.d. = 97.80 $\pm$ 43.31). Sixteen of these participants with late-

onset acquired blindness participated in the subsequent functional imaging experiments

(Table 1, mean age $\pm$ s.d. = 22.43 $\pm$ 1.86 years; Braille reading proficiency, mean $\pm$ s.d.

= 93.69 $\pm$ 32.63; 8 females).

412   ---------Figure 2 here---------

Distinctive functional connectivity between the higher-tier 'visual' cortex and the frontal cortex in the late blind brain

Taking the sighted group as the reference, we performed a whole-brain seed-to-vertex-based functional connectivity analysis to test connectivity changes in the late blind group of two visual brain regions, V1 and LOC, for the two hemispheres separately. The whole-brain functional connectivity for V1 was not different between the blind and sighted groups. In both groups, we found significant connections of V1 with other visual areas but not with frontal language areas—the IFC (Figure 2, Supplementary Figure 1, right panel). By contrast, we observed significantly enhanced functional connectivity of the left LOC with the left IFC in the blind group relative to the sighted group (significant difference at $p < 0.01$, corrected for both hemispheres; Figure 2, left panel). Strong connections of the LOC with other visual regions and extensively with dorsal parietal regions were similarly found in both groups (Figure 2, Supplementary Figure 1, left panel). Notably, despite a significant connection between the left LOC seed and the right IFC in the blind group ($p < 10^{-6}$, uncorrected), the difference in such connectivity with the sighted group failed to reach a significant level (thresholded at $p < 0.01$, corrected for both hemispheres; Figure 2, left panel). In addition, we also found significantly stronger functional connectivity between the left LOC and the left intraparietal sulcus (IPS) in the blind group than in the sighted group (Supplementary Figure 2).

436   ---------Figure 3 here---------

Left LOC-IFC functional connectivity correlated with Braille reading proficiency

of late blind individuals

To test our hypothesis, we examined whether there was an association between the left

LOC-IFC functional connectivity and individual levels of Braille reading proficiency.

We found that the left LOC-IFC functional connectivity was significantly and positively

correlated with individual Braille reading proficiency (Figure 3A left panel, Pearson

correlation, $n = 15$, $p = 0.023$, $r = 0.580$; Spearman correlation, $n = 15$, $p = 0.005$, $r =$

0.686). This effect remained significant after controlling for the age of blindness onset,

current age, and mean framewise displacement (Power et al., 2012) (motion) in the

resting-state scan (partial Pearman correlation, $p = 0.101$, $r = 0.496$; partial Spearman

correlation, $p = 0.011$, $r = 0.704$). By contrast, the left V1-IFC functional connectivity

was not correlated with the blind individual's Braille reading proficiency (Figure 3B

left panel, Pearson correlation, $n = 15$, $p = 0.998$, $r = -0.001$; Spearman correlation, $n =$

15, $p = 0.870$, $r = 0.046$). Thus, better Braille reading proficiency was associated with

greater left LOC-IFC functional connectivity.

Next, following previous studies (Liu et al., 2007), we calculated the estimated number

of hours of practicing Braille reading for late blind participants during childhood,

adolescence, and adulthood (Table 1). We further examined whether the left LOC-IFC

functional connectivity was also associated with the individual number of hours of

practicing Braille reading. We found that the Braille reading practice time during

adolescence was significantly correlated with the left LOC-IFC functional connectivity

in blind individuals (Figure 3A right panel, Pearson correlation, $n = 15$, $p = 0.005$, $r =$

0.679; Spearman correlation, $n = 15$, $p = 0.022$, $r = 0.586$; in other age periods,

Supplementary Figure 3A). This positive relationship remained significant after

controlling for the age of blindness onset, current age, and head motion (Power et al.,

2012) (partial Pearson correlation, $p = 0.027$, $r = 0.634$; partial Spearman correlation,

$p = 0.076$, $r = 0.531$). By contrast, the left V1-IFC connectivity was not associated with

blind individual's Braille reading practice time (Figure 3B right panel, in adolescence,

Pearson correlation, $p = 0.988$, $r = 0.004$; Spearman correlation, $p = 0.970$, $r = 0.011$;

in other age periods, Supplementary Figure 3B). Thus, stronger left LOC-IFC

functional connectivity was associated with more Braille reading practice.

---------Figure 4 here---------

Bidirectional LOC-IFC information flow for Braille reading with a

preponderance in the top-down modulation

Next, we investigated the direction of the left LOC-IFC information flow and how the

information flow was modulated by the Braille reading task, employing dynamic causal

modeling (DCM). The blind subjects were asked to silently read pages containing

natural Braille articles while undergoing an fMRI scan, with no-Braille blank pages as

the control. The Braille > blank contrast elicited bilateral V1, LOC, IPS, and left IFC

activations in blind subjects (Supplementary Figure 4). In addition, we constrained our

DCM models to the left hemisphere due to the left-lateralized activation of the IFC and

the left-lateralized increased LOC-IFC and LOC-IPS functional connections in blind

individuals. Therefore, four brain regions, IPS, V1, LOC, and IFC—all frequently

implicated in Braille reading (Sadato, 2005; Katarzyna Rączy et al., 2019)—were

selected for further DCM analysis.

We specified the model space as shown in Figure 4A. The null model represents the

basic structure, including endogenous effective connectivity among regions of interest

(ROIs) and the modulation of the IPS-LOC connection confirmed in the previous study

(Fujii et al., 2009). We used the IPS to model the tactile input for the LOC because

previous studies (Wittenberg et al., 2004; Sadato, 2005; Fujii et al., 2009) agreed that

the tactile information from the somatosensory cortex was redirected to the occipital

cortex through the poly-sensory intraparietal cortex. To investigate the dynamic

interactions between the left LOC and the left IFC in late blind humans, each direction

of this effective connectivity can be either modulated or not, hence obtaining four

possible models. Random-effects Bayesian model selection (BMS) across blind

individuals showed that the bidirectional model largely outperformed other models

(Figure 4B, exceedance probability [the probability that a particular model is more

likely than any other model] of the bidirectional model = 94.56%).

Bayesian model averaging (BMA) over all the competing models showed that the

endogenous connectivity and modulatory effect parameters between the LOC and IFC

were all significant (Figure 4C, Supplementary Tables 1, 2, posterior probabilities >

0.99). As illustrated in Figure 4C, the LOC-to-IFC endogenous connectivity was

positive (0.134 s^{-1}), while the IFC-to-LOC connectivity was negative (-0.392 s^{-1}). These

results indicate that irrespective of the task-relevant modulation, the activity changes of

the LOC increased the activity of the IFC but not vice versa. The modulation effects of

Braille reading on the left LOC-IFC bidirectional connections were both positive in the

late blind group, with a modulation strength of $+$ 0.547 s^{-1} for the bottom-up connection

and $+$ 1.496 s^{-1} for the top-down connection (Figure 4C left panel). Notably, the Braille

reading task-relevant modulation of the top-down communication from the left IFC to

the LOC was significantly stronger than the bottom-up communication from the LOC

to the IFC (Figure 4C, $p < 0.001$). Such asymmetrical modulation highlighted the

remarkable functional relevance to natural Braille reading of the top-down

communication from the left IFC to the left LOC in late blind humans.

517   ---------Figure 5 here---------

519   **Greater top-down modulation of the left IFC-to-LOC connectivity contributed to**

520   **higher Braille reading proficiency via a broader area of task-engaged LOC**

521   Significant and positive top-down modulation of the IFC-to-LOC effective connectivity

522   during the natural Braille reading task suggested that at the neuronal level, the

523   information flow from the left IFC increasingly induced the task-specific activity in the

left LOC. Accordingly, to measure such neural activity, we identified the individual

peak of activation (Supplementary Figure 5C) of the left LOC during natural Braille

reading and surrounding vertices with the strongest functional connection to this LOC

peak. We referred to such vertices within the anatomical LOC region as the functional

LOC of blind individuals. Significant and positive correlations were found between

individual top-down modulatory strength and the area of functional LOC (Figure 5A,

Pearson correlation, $n = 13$, $p = 0.005$, $r = 0.730$), as well as between the area of

individual functional LOC and Braille reading proficiency (Figure 5C, Pearson

correlation, $n = 13$, $p = 0.011$, $r = 0.675$). Thus, we ran a mediation analysis to

investigate whether the top-down modulation of the left IFC-to-LOC connectivity was

associated with individual Braille reading proficiency through the area of functional

LOC. The top-down modulation during the task and individual Braille reading

proficiency acted as the independent and dependent variables respectively, whereas the

area of functional LOC acted as the neural mediator. Results showed that the area of

functional LOC significantly and completely mediated the relationship between top-

down modulation and Braille reading proficiency (Figure 5D, mediation effect = 9.282,

95% CI = [1.765, 23.870], $p = 0.043$), providing a neural surrogate of the top-down

modulation of the left IFC-to-LOC connection in support of late blind individual's

Braille reading proficiency. In contrast, the bottom-up modulation of the left LOC-to-

IFC connection was not significantly associated with either the area of functional LOC

(Figure 5B, $n = 13$, $p = 0.644$, $r = 0.142$) or individual Braille reading proficiency

(Supplementary Figure 5B).

547   ---------Figure 6 here---------

A 'two-tale' model with the LOC-IFC neural link better predicts individual Braille

reading proficiency

Finally, we established neural predictive models to further understand the neural

mechanisms of individual variance in Braille reading proficiency. We imagined a linear

'two-tale' model with two neurofunctional predictors that the resting-state functional

connectivity (rsFC) and task-specific top-down effective connectivity (tdEC) of the left

LOC-IFC neural link each explained independent variance in individual Braille reading

proficiency in late blindness. Here we used the area of functional LOC (Figure 5D) as

the neural surrogate of the top-down modulation of the left IFC-to-LOC effective

connectivity during the natural Braille reading task. We also ran linear models with each

of these two measures of the left LOC-IFC neural connection separately to predict

Braille reading proficiency. We assessed the models' performance for held-out data

using a leave-one-out cross-validation procedure. Cross-validation results showed that

the 'two-tale' model outperformed the other two models in terms of both MSE (Figure

6B, Table 2, mean $\pm$ s.e.: 'two-tale' model = 457.619 $\pm$ 158.048; Model with only rsFC

= 661.208 $\pm$ 245.886; Model with only tdEC = 651.066 $\pm$ 174.217) and R^2 (Table 2,

Model 1 = 0.490; Model 2 = 0.285; Model 3 = 0.279) between the predicted proficiency

and the held-out actual proficiency. Notably, the coefficients on these two neural

measures did not change dramatically, and the total variance explained was near

additive when considering these two variables separately and together in different

models (Table 2). These results reinforced that for the left LOC-IFC neural link of late

blind humans, the rsFC and the tdEC strength each significantly contributed to unique

individual variance in Braille reading proficiency in distinct ways.

Discussion

This study linked resting-state functional connectivity and task-engaged effective

connectivity from functional imaging data together to natural Braille reading behavior

in late blindness. The remarkable resting-state functional connectivity between the

higher-tier 'visual' cortex—the LOC, but not the V1, and the IFC in the late blind brain,

as well as its significant relationship with individual Braille reading proficiency,

concurred in suggesting the indispensable role of such functional connectivity in

bridging Braille orthographic and language processing of natural Braille reading. In a

natural Braille reading task, DCM analyses revealed increased bidirectional

information flow with a preponderance in the top-down modulation of the left IFC-to-

LOC effective connectivity. Further analyses suggested greater top-down modulation

of the left IFC-to-LOC effective connectivity contributed to higher Braille reading

proficiency through a mediation effect. Crucially, to fully reveal the neural mechanisms

of individual variance in Braille reading performance, we established a neural

predictive model with two variables of the LOC-IFC link: the functional connectivity

and the top-down effective connectivity. This 'two-tale' model suggests that developing

the underpinning neural circuit and the top-down cognitive strategy contributes

uniquely to superior Braille reading performance. This work potentially provides a

general framework for understanding crossmodal neuroplasticity during learning.

The crossmodal recruitment of the typical visual cortex for nonvisual higher cognitions

in blind individuals, including Braille reading, has been frequently observed (Pavani

and Röder, 2012; Renier et al., 2014; Bedny, 2017; Heimler and Amedi, 2020). A

particular concern regarding this plasticity is whether there is a reversed hierarchy of

functional organization across 'visual' regions or not (Amedi et al., 2003; Büchel, 2003;

Burton, 2003; Watkins et al., 2012). Studies by Amedi et al. (2003) first proposed a

reversal of the normal visual hierarchy in congenitally blind individuals by showing

that compared to higher-tier 'visual' regions, early V1 preferentially responded to more

'abstract' cognitive tasks (Amedi et al., 2003; Büchel, 2003). These findings are in

accord with studies showing functional connectivity of blind individuals between the

V1 and frontal language areas at rest (Butt et al., 2013; Striem-Amit et al., 2015). Some

studies, on the other hand, did not support such reversed hierarchy by providing

evidence of task activations (Röder et al., 2002; Burton, 2003; Noppeney, 2003) and

resting-state functional connectivity (Bedny et al., 2011; Watkins et al., 2012) of the

primary and higher-tier 'visual' regions. For the late blind who lost their sight several

years after birth, a normal hierarchy is more possible since late blind humans might

have initially developed their neural circuits as normally sighted people (Röder et al.,

2021). Here in our study, the functional connectivity between higher-tier 'visual' region

LOC and the frontal cortex (Figure 2) and its behavioral significance for natural Braille

reading (Figure 3) indicates a normal hierarchy. Notably, the anatomical LOC area in

this study, approximately encompassed the anatomical location of the visual word form

area (VWFA) as in sighted people. Studies showed the 'VWFA' activation of blind

humans not only in tactile Braille reading but also in functions such as language

processing (Reich et al., 2011; Kim et al., 2017; Katarzyna Rączy et al., 2019), which

is similar to the functional profile of the broader LOC cortex (Noppeney, 2003; Bedny

et al., 2011; Watkins et al., 2012; Bedny, 2017). In line with these findings, our data

suggest that the left LOC in late blind humans might incorporate the VWFA-like

function and extensively develop as a versatile crossmodal region, bridging Braille

orthographic and language processing in natural Braille reading via the distinctive

neural nexus of the left LOC-IFC. In line with the connectivity-based plasticity theory

(Schepers et al., 2012; Bedny, 2017), the development of such neural connectivity in

late blind humans likely depends on the prior connectivity between the VWFA and

frontal language areas as normally sighted people.

Our DCM results provide direct evidence for the direction of information flow on the

left LOC-IFC neural interaction of late blind humans during a natural Braille reading

task. In line with the Hebbian theory (Hebb, 1950), the winning model suggests that

natural Braille reading strongly modulates and strengthens the bidirectional LOC-IFC

connection (Figures 4A, 4B). This result mirrors the robust resting-state functional

connectivity in late blind humans (Figure 2A), as well as its positive relationship with

Braille reading practice time (Figure 3A), although the preferential practice time might

depend on the age of blindness onset (Liu et al., 2007). However, a closer look at the

winning model suggests that such neural connection does not simply convey balanced

reciprocal interactions but predominantly delivers the task-specific top-down

information from the higher cognitive cortex—the left IFC (Figure 4C). This top-down

influence guarantees the active prediction and perception of the tones and semantic

meanings through the use of contextual cues (Trent and Truan, 1997; Friston, 2010).

Importantly, we found that the task modulation of top-down connectivity from the left

IFC to the left LOC was strongly and positively associated with individual Braille

reading proficiency through a mediator, the area of task-engaged functional LOC

(Figure 5D). Our findings reinforced the significant contribution of the top-down

influence to natural Braille reading.

Furthermore, why the 'visual' cortex engages more actively in various crossmodal tasks

in blind humans, including late blind humans, than in sighted controls (Matuszewski et

al., 2021) is still debated (Röder et al., 2021). Using natural Braille reading as a model,

the current study provides empirical evidence for the hypothesis that 'visual' cortex

activity in late blindness predominantly arises from the top-down connectivity from the

higher-order multisensory or supramodal cortex (Collignon et al., 2013; Bedny, 2017;

Röder et al., 2021). This neural mechanism is essential for the rapid acquisition of new

skills because it might be relatively difficult for late blind humans to extract crossmodal

characteristics exclusively via bottom-up stimulus input, with their visual deprivation

occurring after the pruning of exuberant crossmodal connectivity during the typical

early development (Batardière et al., 2002; Röder et al., 2021). Such a top-down

controlled learning mechanism in late blind humans is also compatible with the typical

adult learning characteristics demonstrated by human and non-human animal studies

(Keuroghlian and Knudsen, 2007; Rohlf et al., 2017).

To our knowledge, this is the first study to date to establish the neural predictive model

for the natural Braille reading performance of late blind individuals. Our model's two

neurofunctional predictors regarding the left LOC-IFC neural interactions arguably

happen by distinct experience-dependent brain plasticity in late blindness. The

functional connectivity predictor at the resting state might stem from prolonged Braille

reading practicing of late blind humans on a much slower timescale (likely at months

to years), which is fundamental to high-level Braille reading performance. In line with

previous studies (Bengtsson et al., 2005; Scholz et al., 2009), we postulate that

increased myelination resulted from constant neural activity along the fiber tract of the

inferior fronto-occipito fasciculus (Forkel et al., 2014) during training, is a potential

mechanism underlying the facilitated communications between neural ensembles of the

LOC and the IFC. Previous work in animals (Pan et al., 2020) also showed that lifelong

learning-induced myelin plasticity affected neuronal responses and subsequent

behavior. On the other hand, the effective connectivity predictor in our model represents

the effect of selectively implementing a top-down cognitive strategy such as active

prediction in every natural Braille reading task on a faster timescale (likely at

milliseconds to seconds). Consistent with previous studies (Citri and Malenka, 2008),

we speculate that the burst of activity caused by the top-down influence of the IFC

largely increases the probability of neurotransmitter release to neurons of the LOC.

Such metabolically and behaviorally efficient synaptic plasticity (Mongillo et al., 2008)

boosts the information transmission of synapses (Citri and Malenka, 2008) and the LOC

activity and, therefore, the behavior performance. If further refined and validated in a

larger cohort, this brain-behavior model might be not only beneficial for more efficient

acquisition of Braille reading skills in blind individuals, but also instructive for more

general learning practices. Future studies are needed to provide detailed

neurobiological evidence to explicitly elucidate the mechanism of each neurofunctional

predictor.

Unlike other studies using Braille letters or words, the large-scale Braille articles used

in our study made it possible to observe task-specific top-down communication in blind

humans during natural Braille reading. The 'two-tale' model proposed in the current

study (Figure 6) suggests that for blind Braille readers to make better use of this top-

down effective connectivity during natural Braille reading, a promising strategy might

be to make the reading context more available. Compared with traditional refreshable

Braille displays offering only a single line of Braille words each time (Ren et al., 2008),

emerging technology is the large-print Braille computer (Völkel et al., 2008; Jiao et al.,

2018), which enables continuous and large-scale natural Braille context display (Figure

1C). This new type of Braille reading device arguably allows for active understanding

and prediction based on contextual information during natural Braille reading, boosting

the top-down information flow and thus promoting Braille reading proficiency of blind

individuals.

References

Abboud S, Cohen L (2019) Distinctive Interaction Between Cognitive Networks and the Visual Cortex in Early Blind Individuals. Cerebral cortex (New York, NY : 1991) 29:4725–4742.

Amedi A, Raz N, Pianka P, Malach R, Zohary E (2003) Early "visual" cortex activation correlates with superior verbal memory performance in the blind. Nature Neuroscience 6:758–766.

Batardière A, Barone P, Knoblauch K, Giroud P, Berland M, Dumas AM, Kennedy H (2002) Early specification of the hierarchical organization of visual cortical areas in the macaque monkey. Cerebral Cortex 12:453–465.

Bedny M (2017) Evidence from Blindness for a Cognitively Pluripotent Cortex. Trends in Cognitive Sciences 21:637–648.

Bedny M, Pascual-Leone A, Dodell-Feder D, Fedorenko E, Saxe R (2011) Language processing in the occipital cortex of congenitally blind adults. Proceedings of the National Academy of Sciences 108:4429–4434.

Bengtsson SL, Nagy Z, Skare S, Forsman L, Forssberg H, Ullén F (2005) Extensive piano practicing has regionally specific effects on white matter development. Nature Neuroscience 8:1148–1150.

Büchel C (2003) Cortical hierarchy turned on its head. Nature Neuroscience 6:657–658.

Burton H (2003) Visual cortex activity in early and late blind people. Journal of Neuroscience 23:4005–4011.

Butt OH, Benson NC, Datta R, Aguirre GK (2013) The fine-scale functional correlation of striate cortex in sighted and blind people. Journal of Neuroscience 33:16209–16219.

Chen X, Lu B, Yan CG (2018) Reproducibility of R-fMRI metrics on the impact of different strategies for multiple comparison correction and sample sizes. Human Brain Mapping 39:300–318.

Citri A, Malenka RC (2008) Synaptic plasticity: Multiple forms, functions, and

mechanisms. Neuropsychopharmacology 33:18–41.

Cohen LG, Celnik P, Pascual-Leone A, Corwell B, Faiz L, Dambrosia J, Honda M,

Sadatok N, Gerloff C, Catala MD, Hallett M (1997) Functional relevance of

cross-modal plasticity in blind humans. Nature 389:180–183.

Collignon O, Dormal G, Albouy G, Vandewalle G, Voss P, Phillips C, Lepore F

(2013) Impact of blindness onset on the functional organization and the

connectivity of the occipital cortex. Brain 136:2769–2783.

Coppins N, Barlow-Brown F (2006) Reading difficulties in blind, braille-reading

children. British Journal of Visual Impairment 24:37–39.

Cox RW (1996) AFNI: Software for analysis and visualization of functional magnetic

resonance neuroimages. Computers and Biomedical Research 29:162–173.

Destrieux C, Fischl B, Dale A, Halgren E (2010) Automatic parcellation of human

cortical gyri and sulci using standard anatomical nomenclature. NeuroImage

53:1–15.

Forkel SJ, Thiebaut de Schotten M, Kawadler JM, Dell'Acqua F, Danek A, Catani M

(2014) The anatomy of fronto-occipital connections from early blunt dissections

to contemporary tractography. Cortex 56:73–84.

Friston K (2010) The free-energy principle: A unified brain theory? Nature Reviews

Neuroscience 11:127–138.

Friston KJ, Harrison L, Penny W (2003) Dynamic causal modelling. NeuroImage

19:1273–1302.

Fujii T, Tanabe HC, Kochiyama T, Sadato N (2009) An investigation of cross-modal

plasticity of effective connectivity in the blind by dynamic causal modeling of

functional MRI data. Neuroscience Research 65:175–186.

Hebb DO (1950) The Organization of Behavior; A Neuropsychological Theory. The

American Journal of Psychology 63:633.

Heimler B, Amedi A (2020) Are critical periods reversible in the adult brain? Insights

on cortical specializations based on sensory deprivation studies. Neuroscience

and Biobehavioral Reviews 116:494–507.

Jiao Y, Lu X, Xu Y (2018) A graphical tactile display for the visually impaired. [C]

Proceedings of AsiaHaptics, Best Video Demo Award.

Jung WH, Lee S, Lerman C, Kable JW (2018) Amygdala Functional and Structural

Connectivity Predicts Individual Risk Tolerance. Neuron 98:394-404.e4.

Katarzyna Rączy, Urbańczyk A, Korczyk M, Szewczyk JM, Sumera E, Szwed M

(2019) Orthographic Priming in Braille Reading as Evidence for Task-specific

Reorganization in the Ventral Visual Cortex of the Congenitally Blind. Journal

of Cognitive Neuroscience 31:1065–1078.

Keuroghlian AS, Knudsen EI (2007) Adaptive auditory plasticity in developing and

adult animals. Progress in Neurobiology 82:109–121.

Kim JS, Kanjlia S, Merabet LB, Bedny M (2017) Development of the Visual Word

Form Area Requires Visual Experience: Evidence from Blind Braille Readers.

The Journal of Neuroscience 37:11495–11504.

Liu Y, Yu C, Liang M, Li J, Tian L, Zhou Y, Qin W, Li K, Jiang T (2007) Whole

brain functional connectivity in the early blind. Brain 130:2085–2096.

Marshall L, Moys J-L (2020) Readers' experiences of Braille in an evolving

technological world. Visible Language 54.

Mason LK (2012) Experimental investigation of hand and finger usage in braille

reading. Dissertations 209.

Matuszewski J, Kossowski B, Bola Ł, Banaszkiewicz A, Paplińska M, Gyger L,

Kherif F, Szwed M, Frackowiak RS, Jednoróg K, Draganski B, Marchewka A

(2021) Brain plasticity dynamics during tactile Braille learning in sighted

subjects: Multi-contrast MRI approach. NeuroImage 227:117613.

Mongillo G, Barak O, Tsodyks M (2008) SynaptiC Theory of Working Memory.

Science 319:1543–1546.

Noppeney U (2003) Effects of visual deprivation on the organization of the semantic

system. Brain 126:1620–1627.

Pan S, Mayoral SR, Choi HS, Chan JR, Kheirbek MA (2020) Preservation of a remote

fear memory requires new myelin formation. Nature Neuroscience 23.

Pascual-Leone A, Amedi A, Fregni F, Merabet LB (2005) the Plastic Human Brain

Cortex. Annual Review of Neuroscience 28:377–401.

Pavani F, Röder B (2012) Crossmodal plasticity as a consequence of sensory loss:
Insights from blindness and deafness. In: The new handbook of multisensory
processes. (Stein BE, ed), pp 737–760. Cambridge, MA: MIT Press.

Penny WD, Stephan KE, Daunizeau J, Rosa MJ, Friston KJ, Schofield TM, Leff AP
(2010) Comparing families of dynamic causal models. PLoS Computational
Biology 6.

Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE (2012) Spurious but
systematic correlations in functional connectivity MRI networks arise from
subject motion. NeuroImage 59:2142–2154.

Reich L, Szwed M, Cohen L, Amedi A (2011) A ventral visual stream reading center
independent of visual experience. Current Biology 21:363–368.

Ren K, Liu S, Lin M, Wang Y, Zhang QM (2008) A compact electroactive polymer
actuator suitable for refreshable Braille display. Sensors and Actuators, A:
Physical 143:335–342.

Ren Y, Nguyen VT, Sonkusare S, Lv J, Pang T, Guo L, Eickhoff SB, Breakspear M,
Guo CC (2018) Effective connectivity of the anterior hippocampus predicts
recollection confidence during natural memory retrieval. Nature
Communications 9.

Renier L, De Volder AG, Rauschecker JP (2014) Cortical plasticity and preserved
function in early blindness. Neuroscience and Biobehavioral Reviews 41:53–63.

Röder B, Kekunnaya R, Guerreiro MJS (2021) Neural mechanisms of visual sensitive
periods in humans. Neuroscience and Biobehavioral Reviews 120:86–99.

Röder B, Stock O, Bien S, Neville H, Rösler F (2002) Speech processing activates
visual cortex in congenitally blind humans. European Journal of Neuroscience
16:930-936.

Rohlf S, Habets B, von Frieling M, Röder B (2017) Infants are superior in implicit
crossmodal learning and use other learning mechanisms than adults. eLife 6:1–
23.

Ryles R (1996) The impact of braille reading skills on employment, income,
education, and reading habits. Journal of Visual Impairment and Blindness

90:219–226.

Ryles R (2000) Braille as a predictor of success. In: Braille into the next millennium

(Judith Dixon, ed), pp 463–491. Washington, DC: National Library Service for

the Blind and Print Disabled, and Friends of Libraries for Blind and Physically

Handicapped Individuals in North America.

Sabbah N, Authié CN, Sanda N, Mohand-Saïd S, Sahel JA, Safran AB, Habas C,

Amedi A (2016) Increased functional connectivity between language and

visually deprived areas in late and partial blindness. NeuroImage 136:162–173.

Sadato N (2005) How the blind "see" Braille: Lessons from functional magnetic

resonance imaging. Neuroscientist 11:577–582.

Sadato N, Pascual-Leone A, Grafman J, Deiber MP, Ibañez V, Hallett M (1998)

Neural networks for Braille reading by the blind. Brain 121:1213–1229.

Schepers IM, Hipp JF, Schneider TR, Röder B, Engel AK (2012) Functionally

specific oscillatory activity correlates between visual and auditory cortex in the

blind. Brain 135:922–934.

Scholz J, Klein MC, Behrens TEJ, Johansen-Berg H (2009) Training induces changes

in white-matter architecture. Nature Neuroscience 12:1370–1371.

Schroeder FK (1996) Perceptions of Braille Usage by Legally Blind Adults. Journal

of Visual Impairment & Blindness 90:210–218.

Seghier ML, Josse G, Leff AP, Price CJ (2011) Lateralization is predicted by reduced

coupling from the left to right prefrontal cortex during semantic decisions on

written words. Cerebral Cortex 21:1519–1531.

Striem-Amit E, Ovadia-Caro S, Caramazza A, Margulies DS, Villringer A, Amedi A

(2015) Functional connectivity of visual cortex in the blind follows retinotopic

organization principles. Brain 138:1679–1695.

Tingley D, Yamamoto T, Hirose K, Keele L, Imai K (2014) Mediation: R package for

causal mediation analysis. Journal of Statistical Software 59:1–38.

Trent SD, Truan MB (1997) Speed, accuracy, and comprehension of adolescent

braille readers in a specialized school. Journal of Visual Impairment and

Blindness 91:494–500.

Völkel T, Weber G, Baumann U (2008) Tactile graphics revised: The novel

BrailleDis 9000 pin-matrix device with multitouch input. Lecture Notes in

Computer Science (including subseries Lecture Notes in Artificial Intelligence

and Lecture Notes in Bioinformatics) 5105 LNCS:835–842.

Wang X, Caramazza A, Peelen M V., Han Z, Bi Y (2015) Reading without speech

sounds: VWFA and its connectivity in the congenitally deaf. Cerebral Cortex.

Watkins KE, Cowey A, Alexander I, Filippini N, Kennedy JM, Smith SM, Ragge N,

Bridge H (2012) Language networks in anophthalmia: Maintained hierarchy of

processing in "visual" cortex. Brain 135:1566–1577.

Winkler AM, Ridgway GR, Douaud G, Nichols TE, Smith SM (2016) Faster

permutation inference in brain imaging. NeuroImage 141:502–516 Available at:

http://dx.doi.org/10.1016/j.neuroimage.2016.05.068.

Wittenberg GF, Werhahn KJ, Wassermann EM, Herscovitch P, Cohen LG (2004)

Functional connectivity between somatosensory and visual cortex in early blind

humans. European Journal of Neuroscience 20:1923–1927.

Yan CG, Cheung B, Kelly C, Colcombe S, Craddock RC, Di Martino A, Li Q, Zuo

XN, Castellanos FX, Milham MP (2013) A comprehensive assessment of

regional variation in the impact of head micromovements on functional

connectomics. NeuroImage 76:183–201.

Yan CG, Wang X Di, Zuo XN, Zang YF (2016) DPABI: Data Processing & Analysis

for (Resting-State) Brain Imaging. Neuroinformatics 14:339–351.

Zeidman P, Jafarian A, Corbin N, Seghier ML, Razi A, Price CJ, Friston KJ (2019) A

guide to group effective connectivity analysis, part 1: First level analysis with

DCM for fMRI. NeuroImage 200:174–190.

Figure legends

Figure 1. Schematic overview of the study. (A) A natural Braille reading test showed a wide difference in Braille reading proficiency of blind individuals. (B) Both resting-state signals and dynamic task-related fMRI data were assessed. The whole-brain resting-state functional connectivity of two visual brain regions (V1 and LOC), respectively, especially with the frontal cortex, was compared between late blindness and sighted groups. The task fMRI paradigm was designed to simulate real-life Braille reading using natural Braille articles. (C) The current study might inspire the design change from a traditional Braille display (upper panel) with limited single-line Braille words each time to a natural Braille reading device (lower panel) with a continuous Braille article display for blind individuals.

Figure 2. Functional connectivity patterns for the primary 'visual' cortex (V1) and the lateral occipital cortex (LOC). (A) Maps show vertices with significantly positive connectivity with the left LOC seed (vertexwise $p < 10^{-6}$, uncorrected, cluster size > 100 mm) in the late blind group and sighted group (left panel). Maps show vertices with significant positive connectivity with the left V1 seed (vertexwise $p < 10^{-6}$, uncorrected, cluster size > 100 mm) in the late blind group and sighted group (right panel). (B) Differences in functional connectivity patterns for left LOC (left panel) and V1 (right panel) in the late blind group versus the sighted group, respectively. Significant vertices were corrected for multiple comparisons using permutation testing with threshold-free

cluster enhancement ($p < 0.01$, corrected for both hemispheres, cluster size > 30 mm).

Box plots of the two groups' connectivity strengths (Fisher r-to-z-transformed) between

the seed and the left IFC cluster (located from the left panel). The horizontal line

denotes the median, and the asterisk indicates significance. V1 primary visual cortex

(light green), LOC lateral occipital cortex (light yellow), IFC inferior frontal cortex. L

left hemisphere, R right hemisphere. Blind, dark color; Sighted, light color. **$p < 0.01$,

corrected for both hemispheres. n.s. non-significant.

Figure 3. Left LOC-IFC functional connectivity showed significant correlations with individual Braille reading proficiency and practice time. (A) Plots of the left LOC-IFC functional connectivity. Significant and positive correlations were found between individual strengths of the left LOC-IFC functional connectivity and Braille reading proficiency (left panel) and Braille reading practice time in adolescence (in hours, right panel). (B) The connectivity between the left V1 and the left IFC was not correlated with either Braille reading proficiency (left panel) or Braille reading practice time (right panel). L left hemisphere. All *r* and *p*-value pairs reflect Pearson correlation, consistent with the Spearman correlation results.

Figure 4. DCM analyses of modulations of the left LOC-IFC effective connectivity

for natural Braille reading. (A) Model space specified by adding task modulation to

different directions of the left LOC-IFC connection, resulting in four competing models.

(B) Exceedance probability for the DCM models. (C) Illustration of the Bayesian model

averaging (BMA) result for the late blind group (all significant at a posterior probability

threshold of 0.99). Modulators added to the LOC-IFC connection are coded in the size

of the filled circles, representing their effect size (left panel). The modulatory strength

(in Hz) of Braille reading on the IFC-to-LOC connection was significantly higher than

that on the LOC-to-IFC connection ($p < 0.001$). For each parameter (endogenous or

modulatory, in Hz) of the top-down IFC-to-LOC connection and the bottom-up LOC-

to-IFC connection, the distribution of the 10,000 samples of the posterior densities is

provided, respectively. Endogenous connectivity (gray); modulatory strength, red for

IFC-to-LOC connection, and cyan for LOC-to-IFC connection. ***$p < 0.001$.

Figure 5. Top-down modulation of the IFC-to-LOC connection facilitated individual Braille reading proficiency via a broader area of functional LOC in the task. (A) Task modulation of the top-down communication from the left IFC-to-LOC (in Hz) was positively correlated with the area of functional LOC (in %) engaging in natural Braille reading. (B) There was no significant relationship between task modulation of the bottom-up communication from the left LOC to IFC (in Hz) and the area of functional LOC (in %). (C) The area of functional LOC engaging in the natural Braille reading task was positively correlated with individual Braille reading proficiency. All r and p-value pairs reflect Pearson correlation. (D) Mediation analysis showed a significant and complete mediation effect, suggesting that the top-down modulation was significantly associated with enhanced Braille reading proficiency through the increase of the area of functional LOC. FunLOC - functional LOC.

Figure 6. Cross-validation performance of models in predicting Braille reading proficiency of late blind individuals. (A) Schematic representation of the 'two-tale' model in predicting Braille reading proficiency of late blind individuals. The resting-state functional connectivity and the task-specific top-down effective connectivity of the left LOC-IFC neural link contributed to unique variance in individual Braille reading proficiency. According to the mediation effect (Figure 5D), the area of functional LOC engaging in the natural Braille reading task was used as the neural surrogate of the top-down effective connectivity from the left IFC-to-LOC here in the model. (B) Leave-one-out cross-validation analysis showed that the 'two-tale' model outperformed two other models with a single neural predictor, either the functional or top-down effective connectivity of the left LOC-IFC neural link, in predicting a blind individual's Braille reading proficiency. (C) A significant and positive correlation was observed between the predicted Braille reading proficiency of the 'two-tale' model and the held-out test proficiency. The *r* and *p*-value pair reflect the Pearson correlation. rsFC - resting-state functional connectivity between the left LOC and the left IFC, tdEC - top-down modulation of the left IFC-to-LOC effective connectivity in the natural Braille reading task, profi - individual Braille reading proficiency, MSE - mean squared error between the predicted proficiency and the held-out test proficiency.

     Table 1 Characteristic of blind participants of the fMRI scan.

| | Sex | Age | Onset Age (years) | Braille reading practice time | | | | Braille Reading Proficiency (words/min) | Cause of Blindness |
				Start Age (years)	Childhood (h)	Adolescence (h)	Adulthood (h)		
B1	F	23.54	14.72	9.88	4255.90	8212.50	1769.21	92.41	Glaucoma
B2	M	21.39	4.41	7.68	3153.60	9125.00	2951.03	91.15	Ocular pathology
B3	F	22.54	4.12	9.84	2759.40	2737.50	1193.55	92.85	Unknown
B4	F	22.00	5.41	7.29	7736.18	6387.50	1095.00	104.19	Optic atrophy
B5	F	20.31	5.41	7.99	2194.50	6387.50	449.43	153.08	Trauma
B6	M	19.73	15.00	15.00	-	2190.00	680.50	32.10	Macular discoloration
B7	M	23.28	11.18	13.93	-	3923.50	-	62.51	Glaucoma
B8	F	23.87	7.00	7.52	1637.00	912.50	205.29	64.33	Optic atrophy
B9	M	20.35	11.00	6.99	10065.00	10037.50	339.86	147.04	Unknown
B10	M	24.44	9.00	11.08	1002.00	8212.50	220.14	107.57	Ocular pathology
B11	F	18.68	6.00	7.13	6221.43	9125.00	489.10	113.50	Trauma
B12	F	23.46	8.00	6.91	8360.33	15512.50	2917.39	135.68	Glaucoma
B13	M	24.14	10.00	7.28	4307.00	4562.50	848.89	71.07	Glaucoma
B14	M	22.28	6.45	8.62	10486.45	13687.50	19965.50	79.81	Ocular
B15	M	24.56	16.00	19.00	-	-	9937.13	66.70	Thrombocytopenia
B16	F	24.37	17.00	8.00	5475.00	10402.50	5812.63	85.04	Unknown

971 **Table 2** Summary of cross-validation results of linear neural predictive models for

972 Braille reading proficiency of late blind individuals.

	'Two-tale' Model	Model with Only rsFC	Model with Only tdEC
rsFC	21.265	29.946	-
tdEC	229.399	-	317.238
constant	38.775	60.348	54.455
MSE	457.619	661.208	651.066
R^2	0.490	0.285	0.279

Data are given as unstandardized mean coefficients of the fitted models of the leave-one-out cross-validation procedure. rsFC resting-state functional connectivity between the left LOC and the left IFC, tdEC top-down modulation of the left IFC-to-LOC effective connectivity, which contributed to individual Braille reading proficiency in the models via a neural mediator—the area of functional LOC engaging in the natural Braille reading task, MSE mean squared error between the predicted proficiency and the held-out actual proficiency, R^2 R-square between the predicted proficiency and the held-out actual proficiency.