Dynein activation in vivo is regulated by the nucleotide states of its AAA3 domain

Rongde Qiu, Jun Zhang, Jeremy D. Rotty and Xin Xiang

Department of Biochemistry and Molecular Biology, The Uniformed Services University of the Health Sciences- F. Edward Hébert School of Medicine, Bethesda, Maryland 20814, USA.

Corresponding author contact:

Xin Xiang (Ph. D.)

Department of Biochemistry and Molecular Biology
Uniformed Services University of the Health Sciences
4301 Jones Bridge Road, Bethesda, Maryland 20814
Tel: (301) 295-0000
Email: xin.xiang@usuhs.edu

Running title:

AAA3 nucleotide states regulate dynein activation

Abstract

Cytoplasmic dynein is activated by cargo adapters, dynactin, and LIS1. However, it is unclear whether there are additional regulations for coordinating cargo binding and dynein activation. Here we found that a dynein AAA4 arginine-finger mutation in *Aspergillus nidulans* bypasses the requirement of LIS1 for dynein activation driven by the early endosomal adapter HookA. As AAA4 arginine-finger is implicated in AAA3 ATP hydrolysis, we examined AAA3 mutants defective in ATP binding and hydrolysis respectively. Astonishingly, blocking AAA3 ATP hydrolysis allows dynein activation by dynactin without LIS1 or HookA. As a consequence, dynein accumulates at microtubule minus ends while early endosomes stay near the plus ends. Blocking AAA3 ATP binding abnormally stabilizes the dynein-LIS1 interaction, suggesting that ATP binding at AAA3 regulates the dynamic dynein-LIS1 interaction. Thus, the AAA3 ATPase cycle not only regulates dynein's mechanochemical cycle but also regulates the dynein-LIS1 interaction and the coordination between dynein activation and cargo binding.

Introduction

Cytoplasmic dynein-1 (called "dynein" hereafter for simplicity) is a minus-end-directed microtubule motor that transports a variety of cellular cargoes as well as viruses, and dysfunction of dynein or its regulators is linked to neurodevelopmental disorders and neurogenerative diseases (Wynshaw-Boris, 2007; Perlson et al., 2010; Schiavo et al., 2013; Reck-Peterson et al., 2018; Stavoe & Holzbaur, 2019; Scherer et al., 2020). Dynein is a multi-protein complex containing heavy chains, intermediate chains, light intermediate chains and light chains (Pfister et al., 2006). The two dynein heavy chains form a homodimer, and each heavy chain contains a motor ring with six AAA+ (ATPases Associated with diverse cellular Activities) domains, a stalk connecting the microtubule-binding domain and a buttress that interacting with the stalk (Koonce et al., 1992; Gee et al., 1997; S. M. King, 2000; Burgess et al., 2003; Carter et al., 2011; Kon et al., 2011; Kon et al., 2012; Schmidt et al., 2012; Roberts et al., 2013; Lin et al., 2014; Schmidt et al., 2015; Schmidt & Carter, 2016). At the N-terminus of AAA1 is the linker domain, whose conformational change is involved in dynein's mechanochemical cycle (Roberts et al., 2009; Kon et al., 2012; Bhabha et al., 2014; Schmidt et al., 2015; Bhabha et al., 2016), and at the N-terminus of the linker is the dynein tail that binds other dynein subunits and the multi-component dynactin complex (Karki & Holzbaur, 1995; K. T. Vaughan & Vallee, 1995; S. J. King et al., 2003; Schroer, 2004; Pfister et al., 2006; Schlager et al., 2014; Chowdhury et al., 2015; Urnavicius et al., 2015; K. Zhang et al., 2017a; Grotjahn et al., 2018; Urnavicius et al., 2018). The dynactin complex and specific cargo adapters allow dynein to associate with cargoes such as early endosomes (J. Zhang et al., 2011; Yeh et al., 2012; Bielska et al., 2014; Scherer et al., 2014; J. Zhang et al., 2014; Hoogenraad & Akhmanova, 2016; Reck-Peterson et al., 2018; Celestino et al., 2019; Olenick & Holzbaur, 2019; Gonçalves et al., 2020). Importantly, several coiled-coils-containing cargo adapters, including the BicD family and the Hook family of proteins, activate the processive movement of mammalian dynein in the presence of dynactin (McKenney et al., 2014; Schlager et al., 2014; Olenick et al., 2016; Schroeder & Vale, 2016). Dynactin and cargo adapter-mediated dynein activation is caused by a series of conformational changes (K. Zhang et al., 2017a). The dynein dimer initially forms an autoinhibited "phi" conformation (Amos, 1989; Torisawa et al., 2014; K. Zhang et al., 2017a), which can be switched to an open conformation with the motor domains being separated from each other; Binding of dynactin and a cargo adapter to the dynein tails turns the dynein dimer to a parallel configuration, allowing it to walk along a microtubule (K. Zhang et al., 2017a). Cargo adapter-mediated dynein activation is a useful mechanism that allows dynein to be activated only after it binds its cargo. Function of the cargo adapter has been implicated in enhancing the interaction between purified mammalian dynein and dynactin (Splinter et al., 2012; McKenney et al., 2014; Schlager et al., 2014; Olenick et al., 2016). However, since dynein and dynactin interact before cargo binding in some in vivo systems (Markus et al., 2020; Xiang & Qiu, 2020), additional regulatory mechanisms may exist to ensure that dynein is not activated by dynactin alone before cargo binding.

The minus-end-directed movement of dynein requires multiple elements of the dynein heavy chain, including the AAA1-6 domains, the linker, the buttress, the stalk and its distal microtubule-binding domain (Bhabha et al., 2016; Can et al., 2019; Niekamp et al., 2019; Rao et al., 2019; X. Liu et al., 2020). Although AAA1-4 domains can all bind ATP (Gibbons et al., 1991), only AAA1, 3, and 4 hydrolyze ATP, with AAA1 ATP hydrolysis being the most critical for driving motility (Kon et al., 2004; Schmidt et al., 2015; Bhabha et al., 2016; Schmidt & Carter, 2016). ATP binding at AAA1 drives the conformational change of the AAA ring and makes the ring closer (Carter et al., 2011; Bhabha et al., 2014), and it also causes the linker to change from an extended conformation with its N-terminus docked at AAA5 to a bent conformation with its N-terminus moved to AAA2/3 (Bhabha et al., 2014; Schmidt et al., 2015), and these changes are transmitted to buttress and stalk, causing the stalk to shift its coiled-coil registry, which switches the microtubule-binding domain to a low-affinity state (Schmidt et al., 2015; Bhabha et al., 2016; Schmidt & Carter, 2016;

Rao et al., 2019). ATP hydrolysis at AAA3 and ATP binding at AAA4 regulate the AAA1-dependent release of dynein from microtubule (Silvanovich et al., 2003; Kon et al., 2004; Cho et al., 2008; Bhabha et al., 2014; DeWitt et al., 2015; Nicholas et al., 2015; X. Liu et al., 2020). Most relevant to our current study, structural studies combined with biophysical analyses in vitro on purified yeast dynein motor domain have provided significant insights into the function of the AAA3 ATPase cycle (Bhabha et al., 2014; DeWitt et al., 2015; Nicholas et al., 2015). Specifically, when the AAA3 ATP hydrolysis is blocked, the linker fails to bend in response to ATP binding at AAA1 and dynein fails to release from microtubule (Bhabha et al., 2014; DeWitt et al., 2015; Nicholas et al., 2015). It has been thought that the AAA3 ATPase cycle may participate in the spatial regulation of dynein in vivo (Bhabha et al., 2014; DeWitt et al., 2015); For example, the strong microtubule binding of dynein with its ATP-bound AAA3 may help anchor dynein at the vicinity of the microtubule plus end before the initiation of minus-end-directed movement. However, this interesting idea has not yet been examined in vivo in the context of cargo adapter, dynactin and another important dynein regulator LIS1. LIS1 is a causal gene for human lissencephaly and a known dynein regulator in many organisms and cell types (Reiner et al., 1993; Wynshaw-Boris, 2007; Markus et al., 2020), and it binds to the dynein motor ring around AAA3/AAA4 and to a site on the stalk (Huang et al., 2012; Toropova et al., 2014; DeSantis et al., 2017; Htet et al., 2020). Different AAA3 nucleotide states cause the yeast dynein motor domain to respond to LIS1 differently in vitro: While LIS1 strongly inhibits the motility of apo-AAA3 dynein, it enhances the motility of ATP-AAA3 dynein (DeSantis et al., 2017). However, as the relationship between LIS1 and the AAA3 ATPase cycle has not yet been examined in the context of dynactin and cargo adapters, it would be important to do so especially given the more recent knowledge of LIS1 being positively involved in dynein activation in the context of dynactin and cargo adapters (Baumbach et al., 2017; Gutierrez et al., 2017; Jha et al., 2017; Qiu et al., 2019; Elshenawy et al., 2020; Htet et al., 2020; Marzo et al., 2020).

We have been studying the in vivo regulation of dynein using the filamentous fungus *Aspergillus nidulans* as a genetic system. Dynein in filamentous fungi is required for nuclear distribution (Plamann et al., 1994; Xiang et al., 1994; Alberti-Segui et al., 2001; Xiang, 2018) and for the transport of early endosomes and other cargoes, some of which move by hitchhiking on the motile early endosomes (Wedlich-Soldner et al., 2002; Lenz et al., 2006; Baumann et al., 2012; Etxebeste et al., 2013; Higuchi et al., 2014; Penalva et al., 2017; Salogiannis & Reck-Peterson, 2017; Steinberg et al., 2017; Hernández-González et al., 2018; Otamendi et al., 2019; Salogiannis et al., 2021). Inside the fungal hypha, dynein, dynactin and LIS1 all accumulate at the microtubule plus ends near the hyphal tip, and the plus-end accumulation of dynein and dynactin depends on kinesin-1 (Han et al., 2001; J. Zhang et al., 2003; Lenz et al., 2006; J. Zhang et al., 2010; Egan et al., 2012; Yao et al., 2012). The microtubule plus-end accumulation of dynein is important for dynein-mediated transport of early endosomes away from the hyphal tip (Lenz et al., 2006; Abenza et al., 2009; Zekert & Fischer, 2009). The physical interaction between dynein and early endosome depends on the dynactin complex as well as the FHF (Fts, Hook and Fhip) complex (note that the FHF complex was first identified in mammalian cells (Xu et al., 2008)) (J. Zhang et al., 2011; Yeh et al., 2012; Bielska et al., 2014; Yao et al., 2014; J. Zhang et al., 2014). Within the *A. nidulans* FHF complex, HookA lacking the C-terminal cargo-binding domain (ΔC-HookA) interacts with dynein-dynactin (J. Zhang et al., 2014); FhipA mediates the interaction between HookA's C-terminus and early endosome (Yao et al., 2014), most likely via its direct interaction with Rab5 on early endosome (Guo et al., 2016). Recently, we found that ΔC-HookA overexpression drives the relocation of dynein and dynactin from the microtubule plus ends to the minus ends at septa or the spindle-pole bodies (SPBs) (Qiu et al., 2019; Bieger et al., 2021), both of which contain active microtubule-organizing centers (Oakley et al., 1990; Y. Zhang et al., 2017b). This relocation is consistent with cargo-adapter-mediated dynein activation observed in vitro (McKenney et al., 2014; Schlager et al., 2014; Baumbach et al., 2017; Jha et al., 2017). By using this ΔC-HookA-driven dynein relocation as a readout for dynein activation in vivo, we found that dynein activation requires NudF/LIS1 in *A. nidulans* (Qiu et al., 2019). Importantly, by making the phi mutation

preventing the formation of the phi structure (K. Zhang et al., 2017a), the requirement for NudF/LIS1 is partially bypassed, suggesting that LIS1 promotes the open state of dynein (Qiu et al., 2019). This conclusion is largely consistent with other recent studies (Elshenawy et al., 2020; Htet et al., 2020; Marzo et al., 2020; McKenney, 2020).

A. nidulans genetics has previously revealed a NudF/LIS1-bypass suppressor mutation in the AAA4 arginine finger of the dynein heavy chain (Willins et al., 1997; Zhuang et al., 2007; Huang et al., 2012), which is implicated in AAA3 ATP hydrolysis based on the knowledge from other AAA ATPases (Tucker & Sallai, 2007; Wendler et al., 2012). Here we show that the AAA4 arginine finger mutation allows the function of NudF/LIS1 in the ∆C-HookA-driven dynein activation to be bypassed. We further created dynein AAA3 ATP-binding (Walker A or wA) and ATP-hydrolyzing (Walker B or wB) mutants and analyzed them in the presence and absence of dynactin, NudF/LIS1 or HookA. The wB-AAA3 mutation suppresses the ∆*nudF* (*lis1*) colony growth defect, and the wA-AAA3 mutation causes a relatively much weaker suppression. Surprisingly, the wB-AAA3 dynein accumulates strongly at the microtubule minus ends at septa even without the overexpression of ∆C-HookA. The wA-AAA3 dynein mainly accumulates at the hyphal tip but also shows a septal accumulation. The minus-end localization of wB-AAA3 dynein depends on the functionality of dynein as well as the dynactin complex, but it does not need NudF/LIS1 or HookA, suggesting that blocking AAA3 ATP hydrolysis would allow the requirement of LIS1 or HookA for dynein activation to be bypassed. The wA-AAA3 mutation has a relatively milder effect on bypassing the requirement of LIS1 or HookA, but it alters the dynamic nature of the dynein-LIS1 interaction, which will be discussed in more detail. Together, our data suggest that AAA3 ATPase cycle is critical for dynein regulation in vivo, and most importantly, AAA3 ATP binding modulates the LIS1-dynein interaction, and AAA3 ATP hydrolysis is needed for preventing dynein from being abnormally activated before binding its early endosome cargo.

Results:

Dynein with the AAA4 arginine-finger mutation bypasses the requirement of NudF/LIS1 for ∆C-HookA-driven activation

In *A. nidulans*, genetic studies led to the identification of a *nudF (lis1)* bypass suppressor in dynein HC, *nudA*[R3086C], which represents a mutation in the AAA4 arginine finger (Willins et al., 1997; Zhuang et al., 2007; Huang et al., 2012). Here we tested if the AAA4 arginine finger (RF-AAA4) mutation allows the requirement of NudF/LIS1 for dynein activation to be bypassed. In the strain carrying the RF-AAA4 mutation, GFP-tagged dynein heavy chain (called GFP-dynein for simplicity) localizes along microtubules with a clear plus-end enrichment (Figure 1A) (Zhuang et al., 2007). In addition, septal (minus-end) accumulation of dynein was also readily noticed in some hyphae (Figure 1A). Upon overexpression of ∆C-HookA (*gpdA*-∆C-*hookA*-S), the RF-AAA4 dynein accumulates strongly at the septal minus ends (Figure 1A), similar to that exhibited by the wild-type dynein (Qiu et al., 2019), suggesting that dynein with the RF-AAA4 mutation is activated by the cargo adapter. We then introduced the temperature-sensitive loss-of-function mutation of *nudF/lis*1, *nudF*6, to test if NudF/LIS1 is needed for this activation. Interestingly, a strong septal accumulation of the RF-AAA4 dynein was observed in the *nudF*6, *gpdA*-∆C-*hookA*-S background (Figure 1A), which is in sharp contrast to the plus-end accumulation of wild-type dynein in the *nudF*6, *gpdA*-∆C-*hookA*-S background (Figure 1A, B) (Qiu et al., 2019). Thus, the RF-AAA4 mutation allows the requirement of NudF/LIS1 for dynein activation to be bypassed.

Dynein function in vivo is affected by AAA3 ATP-binding or hydrolysis mutation; the AAA3 ATP-hydrolysis mutation suppresses the ∆*nudF/lis1* colony phenotype

The AAA4 arginine finger is implicated in ATP hydrolysis at AAA3 based on studies of other AAA+ ATPases (Wendler et al., 2012). Indeed, mutations in both the AAA4 arginine finger and the AAA3

Walker B motif affecting ATP-hydrolysis cause a significant reduction in dynein velocity in vitro (Kon et al., 2004; Cho et al., 2008; Huang et al., 2012). Thus, we sought to directly analyze the role of the AAA3 ATPase cycle in dynein activation in vivo mediated by dynactin, LIS1 and cargo adapter. Based on the similarity of dynein heavy chain sequences from mammalian dynein and *A. nidulans* dynein (Figure S1), we constructed the AAA3 ATP-binding (Walker A or wA-AAA3, *nudA*[K2599A]) and hydrolyzing (Walker B or wB-AAA3, *nudA*[E2663Q]) mutants. Both of these mutants formed small colony with a defect in asexual spore production (indicated by the lack of colony color) (Figure 2A), consistent with a defect in nuclear migration to the spores (Xiang et al., 1995a). In comparison to the RF-AAA4 mutant, which formed a colony with asexual spores (note that the green color of the colony is from the asexual spores), these AAA3 mutants are much sicker (Figure 2A). Thus, although the AAA4 arginine finger is implicated in AAA3 ATP hydrolysis, our data showed that it is not essential for this process in *A. nidulans*. Nevertheless, the colonies formed by the wA-AAA3 and wB-AAA3 mutants are obviously healthier than the Δ*nudA* (dynein heavy chain) (Xiang et al., 1995b) and the Δ*nudF* (*lis1*) (Willins et al., 1997) mutants colonies (Figure 2A). Thus, the wA-AAA3 or wB-AAA3 mutant dynein is partially functional in vivo. Consistently, both the wA-AAA3 and wB-AAA3 mutants exhibited a partial defect in nuclear distribution (Figure 2B, 2C). Importantly, the wB-AAA3 mutation clearly suppressed the colony growth defect of the Δ*nudF* mutant, suggesting that the wB-AAA3 mutation allows the function of NudF/LIS1 to be bypassed (Figure 2D). The wA-AAA3 mutation also made the Δ*nudF* colony slightly bigger at 32°C but the suppression is not as strong as that caused by the wB-AAA3 mutation (Figure 2D). Surprisingly, adding the *nudF*6 temperature-sensitive mutation to the wA-AAA3 mutant makes the wA-AAA3 mutant grew better at *nudF*6's semi-permissive temperature of 32°C (Figure 2D). Because the *nudF*6 mutation makes NudF/LIS1 protein less stable (Xiang et al., 1995a), this result suggests that a decrease in the NudF/LIS1 protein level alleviates the defect caused by the wA-AAA3 mutation.

The minus-end localization of dynein at septa is enhanced mildly by the wA-AAA3 mutation and more dramatically by the wB-AAA3 mutation.

We next used live cell imaging to observe the localization of GFP-dynein (Zhuang et al., 2007), carrying either the wA-AAA3 or wB-AAA3 mutation. We also observed early endosomes in the same hyphae labeled by mCherry-RabA (Abenza et al., 2009; Abenza et al., 2010; J. Zhang et al., 2010). In wild-type hyphae, dynein mainly accumulates at the microtubule plus ends near the hyphal tip as comet-like structures (Xiang et al., 2000; Han et al., 2001), although localization of dynein at the septa (minus ends) is also seen (B. Liu et al., 2003; Y. Zhang et al., 2017b). The wA-AAA3 mutant dynein signals are seen near the hyphal tip, although in some hyphae the signals do not look like typical plus-end comets and seem partially overlap with early endosomes abnormally accumulated at the hyphal tip (Figure 3A). In addition, the wA-AAA3 dynein signals also form a faint microtubule decoration with an enrichment at septa (Figure 3A). In comparison, the wB-AAA3 mutant dynein shows no hyphal tip-enriched signals but a more prominent accumulation at septa and microtubule decoration (Figure 3A, 3B). In some hyphae, the wB-AAA3 dynein is also accumulated as bright dots, which correspond to the SPB signals near the individual nuclei (Figure S2). In case of the wild-type dynein, such a strong septal and SPB accumulation was only observed when ΔC-HookA was overexpressed to drive dynein activation (Qiu et al., 2019). As the SPB signals are cell cycle-specific (Bieger et al., 2021), we focus only on the septal accumulation that is more consistently observed. Previously, we showed that the phi mutant dynein (open dynein) also accumulates at the septa without the overexpression of ΔC-HookA, and it is co-localized with early endosomes at the septa (Qiu et al., 2019). However, early endosomes in either the wB-AAA3 mutant or the wA-AAA3 mutant exhibited a hyphal-tip accumulation instead of a septal accumulation (Figure 3A), suggesting that dynein goes to the minus ends without its early endosome cargo.

Next, we examined the localization of dynactin in the wA-AAA3 and wB-AAA3 mutants by introducing the p150-GFP fusion into strains containing the mutations. In the wA-AAA3 mutant, the pattern of p150-GFP localization is similar to that of dynein localization, with both plus-end and septal signals (Figure S3). In the wB-AAA3 mutant, the localization pattern of p150-GFP is also similar to that of wB-AAA3 dynein with strong septal accumulation, although the microtubule decoration appears much weaker, and faint plus-end comets were also observed in some hyphal tips. These observations suggest that dynactin is associated with dynein in these mutants, although some individual dynein and dynactin complexes are also present.

Although the wB-AAA3 dynein decorates microtubules besides forming the minus-end accumulation, the microtubule-bound dynein signals are mainly non-motile. Thus, the minus-end accumulation must have been established during many hours of hyphal growth. To address whether the full-length wB-AAA3 dynein in *A. nidulans* is motile, we prepared cell extract from a strain containing GFP-labeled wB-AAA3 dynein for an in vitro motility assay. A strain containing GFP-labeled wild-type dynein was used as a positive control. In this control strain, the *gpdA*-ΔC-*hookA*-S allele was present to ensure that dynein mainly moves towards the microtubule minus end. We found that wild-type dynein moved robustly in the motility assay, and the full-length wB-AAA3 dynein also moves processively along the microtubule (when 3mM ATP instead of 1mM ATP was used), albeit with a very low speed (Figure S4A, S4B and S4C), consistent with previous in vitro data obtained from the yeast dynein motor domain (Cho et al., 2008; Bhabha et al., 2014; DeWitt et al., 2015; Nicholas et al., 2015; DeSantis et al., 2017; X. Liu et al., 2020). This very slow motility was also observed in vivo for the septum-directed movement of p150-GFP, which was detected very occasionally (Figure S4D), and it is also consistent with the notion that the wB-AAA3 dynein is partially functional in vivo (Figure 2).

Next, we performed in vivo experiments to further confirm that the septal (minus-end) accumulation of the AAA3 mutant dynein needs dynein functionality. To do that, we constructed GFP-tagged dynein heavy chain containing an AAA3 mutation and the *nudA*[F208V] dynein tail mutation that significantly impairs dynein activity but does not impair the dynein-dynactin interaction or dynein-early endosome interaction (Qiu et al., 2013, 2019). Dynein carrying the wA-AAA3 and *nudA*[F208V] double mutations formed plus-end comets near hyphal tip just like dynein with the *nudA*[F208V] single mutation (Figure 3C). Dynein carrying the wB-AAA3 and *nudA*[F208V] mutations showed a microtubule decoration with a clear plus-end enrichment (Figure 3C). Importantly, the *nudA*[F208V] mutation diminished the minus-end accumulation of the mutant dynein, as the septal accumulation disappeared in both the wA-AAA3, *nudA*[F208V] and the wB-AAA3, *nudA*[F208V] double mutants (Figure 3C). This result confirmed that the minus-end accumulation in both the wA-AAA3 or wB-AAA3 single mutants requires dynein functionality.

Activation of wA-AAA3 dynein depends largely on HookA, but activation of wB-AAA3 dynein occurs without HookA, although it still needs dynactin.

In *A. nidulans*, the early endosomal adapter HookA is a critical dynein activator, and overexpression of the cytosolic ΔC-HookA drives dynein relocation from the microtubule plus ends to the minus ends (Qiu et al., 2019). We suspected that activation of the AAA3 mutant dynein could be HookA-independent because the HookA-bound early endosomes did not co-localize with dynein at the septa. However, it would be hard to exclude the possibility that the mutant dynein is activated by the HookA-bound early endosome and then dissociates abnormally from the early endosome. To directly test whether HookA is needed for activating the wA-AAA3 and wB-AAA3 dynein, we introduced the *gpdA*-ΔC-*hookA*-S allele (causing overexpression of ΔC-HookA) (Qiu et al., 2019) or the Δ*hookA* allele (J. Zhang et al., 2014) into strain background with GFP-dynein harboring the wA-AAA3 or wB-AAA3 mutation. We found that activation of the wA-AAA3 dynein is largely but not absolutely HookA-dependent. Overexpression of ΔC-HookA (*gpdA*-ΔC-*hookA*-S)

drives the septal accumulation of the wA-AAA3 dynein, just like wild-type dynein (Figure 4A). In the Δ*hookA* mutant, wild-type dynein forms strong comets near the hyphal tip, and the wA-AAA3 dynein forms plus-end comets as well but also a microtubule decoration and a septal enrichment (Figure 4B). As a consequence, the wA-AAA3 dynein's plus-end signal intensity is lower than that of wild-type dynein in the Δ*hookA* background (Figure 4C). These results suggest that wA-AAA3 dynein needs HookA for activation just like wild-type dynein, but a small portion of the wA-AAA3 dynein molecules may be abnormally activated in a HookA-independent fashion. On the other hand, the wB-AAA3 dynein shows a strong septal accumulation not only upon overexpression of ΔC-HookA (*gpdA*-ΔC-*hookA*-S) but also in the Δ*hookA* background (Figure 4A, 4B). The septal signal intensity of the wB-AAA3 dynein in the Δ*hookA* background is significantly higher than that of the wild-type dynein or the wA-AAA3 dynein in the Δ*hookA* background (Figure 4D). Moreover, while loss of HookA clearly decreases the septal signal intensity of the wA-AAA3 dynein (Figure 4E), it does not significantly decrease the septal signal intensity of the wB-AAA3 dynein (Figure 4F). Thus, the abnormal HookA-independent dynein activation occurs at a much higher degree in the wB-AAA3 mutant than in the wA-AAA3 mutant. However, while the microtubule decoration by the wB-AAA3 dynein was obvious in the Δ*hookA* background, overexpression of ΔC-HookA in the *gpdA*-ΔC-*hookA*-S background drives almost all the wB-AAA3 mutant dynein to septa, indicating that HookA still can facilitate the activation of wB-AAA3, despite the tendency of the mutant dynein to undergo HookA-independent activation.

The dynactin complex is a critical component in dynein activation, as the Arp1 mini-filament provides the dynein tail binding sites (Schlager et al., 2014; Chowdhury et al., 2015; Urnavicius et al., 2015; K. Zhang et al., 2017a; Grotjahn et al., 2018; Urnavicius et al., 2018). However, this function of dynactin has not been directly addressed in *A. nidulans* due to its additional role in the microtubule plus-end accumulation of wild-type dynein. To determine whether dynactin is required for activating the wA-AAA3 or wB-AAA3 dynein, we introduced the Arp1 conditional-null (*alcA-nudK*[Arp1]) allele (J. Zhang et al., 2008) into the strain background with GFP-dynein harboring the wA-AAA3 or wB-AAA3 mutation. In the background of wild-type dynein, shutting down Arp1 expression using this *alcA-nudK*[Arp1] allele causes a loss of microtubule plus-end accumulation of dynein (J. Zhang et al., 2008). Since the *alcA* promoter is repressed by glucose but de-repressed by glycerol (Waring et al., 1989), we used both glycerol- and glucose-containing media to culture the cells containing wA-AAA3, *alcA-nudK*[Arp1] and wB-AAA3, *alcA-nudK*[Arp1] double mutants. In the single AAA3 mutants without the *alcA-nudK*[Arp1] allele, the same localization patterns were observed under these two culture conditions (Figure 5A). In glycerol medium that allows the expression of Arp1 from the *alcA* promoter, the double mutants containing the *alcA-nudK*[Arp1] allele showed the localization patterns similar to those in the single mutants. However, the double mutants showed drastically different localization patterns when Arp1 expression is shut off by glucose. Specifically, loss of Arp1 causes the wA-AAA3 dynein to become diffused in the cytoplasm, consistent with the previously identified role of dynactin in dynein's plus-end accumulation (Xiang et al., 2000; J. Zhang et al., 2003; J. Zhang et al., 2008; Egan et al., 2012; Yao et al., 2012). For the wB-AAA3 dynein, loss of Arp1 allows it to more obviously decorate microtubules but significantly diminishes the septal (minus-end) accumulation (Figure 5A, B), indicating that dynactin is needed for activating the wB-AAA3 dynein.

Activation of the wB-AAA3 dynein does not need NudF/LIS1, but NudF/LIS1 is useful for activating the wA-AAA3 dynein

Next, we observed GFP-dynein harboring the wA-AAA3 or wB-AAA3 mutation in the Δ*nudF* mutant background. The wA-AAA3 dynein is totally accumulated at the microtubule plus ends as represented by bright comets in the absence of NudF/LIS1, suggesting that even the residual HookA-independent septal accumulation of the wA-AAA3 dynein is still dependent on NudF/LIS1. In contrast, the wB-AAA3 dynein accumulated strongly at septa regardless of whether NudF/LIS1

is present or not (Figure 6A). This result indicates that the requirement of NudF/LIS1 can be bypassed by the wB-AAA3 mutation, consistent with the strong suppression of the Δ*nudF* colony defect by the wB-AAA3 mutation (Figure 2D).

Since the septal localization of wA-AAA3 dynein depends on NudF/LIS1, we further examined if overexpression of ΔC-HookA would help overcome this dependency. In this experiment, we used the temperature-sensitive *nudF*6 allele instead of the Δ*nudF* deletion allele to facilitate strain growth on solid medium before the live-cell imaging experiments. Previously, we found that while ΔC-HookA overexpression (*gpdA*-ΔC-*hookA*-S) causes dynein to relocate from the microtubule plus ends to the minus ends, the *nudF*6 mutation prevents this relocation and keeps dynein at the plus ends (Qiu et al., 2019). Consistent with a role of NudF/LIS1 in activation of the wA-AAA3 dynein, the *nudF*6 mutation also kept the wA-AAA3 dynein as plus-end comets upon overexpression of ΔC-HookA (*gpdA*-ΔC-*hookA*-S) (Figure 6B, 6C). However, a septal accumulation of the wA-AAA3 dynein was also obvious in the *nudF*6, *gpdA*-ΔC-*hookA*-S background, which was not observed for wild-type dynein (Figure 6B, 6D). Thus, the wA-AAA3 mutation bypasses NudF/LIS1 function to a low but detectable degree when ΔC-HookA is overexpressed. This may explain why the wA-AAA3 mutation mildly suppresses the colony growth defect of the Δ*nudF* mutant (Figure 2D).

The wA-AAA3 mutation alters the LIS1-dynein interaction

Previously, we found that during the ΔC-HookA-mediated activation of wild-type dynein, NudF/LIS1 accumulates at the microtubule plus ends instead of going to the minus ends with dynein despite its importance for dynein activation (Qiu et al., 2019). This is consistent with LIS1's dissociation from the activated dynein complex during minus-end-directed movement (Lenz et al., 2006; Egan et al., 2012; Jha et al., 2017; Elshenawy et al., 2020; Htet et al., 2020). In the wB-AAA3 dynein mutant, although dynein strongly accumulates at the septal minus ends, NudF/LIS1-GFP forms plus-end comets instead of co-localizing with dynein at the septa, which is similar to the behavior of NudF/LIS1-GFP during the ΔC-HookA-driven activation of wild-type dynein (Figure 7A). Surprisingly, in the wA-AAA3 mutant, NudF/LIS1-GFP localizes to both the hyphal tip and septa just like wA-AAA3 dynein. Next, we introduced the *gpdA*-ΔC-*hookA*-S allele into the NudF/LIS1-GFP, wA-AAA3 strain and the NudF/LIS1-GFP, wB-AAA3 strain to examine NudF/LIS1-GFP localization upon overexpression of ΔC-HookA (Figure 7B). In the wA-AAA3, *gpdA*-ΔC-*hookA*-S background, where the wA-AAA3 dynein accumulates at the septa (Figure 4A), NudF/LIS1-GFP also accumulates at the septa (Figure 7B, 7C), although plus-end comets were still observed. In contrast, NudF/LIS1-GFP does not go to the septa in the wB-AAA3, *gpdA*-ΔC-*hookA*-S strain. These results suggest that the wA-AAA3 mutation allows NudF/LIS1 to stay with dynein and co-localize with dynein at the microtubule minus ends, while the wB-AAA3 dynein tends to dissociate from NudF/LIS1 just like wild-type dynein after its cargo adapter-mediated activation.

Discussion:

Significant progress has been made in understanding dynein activation (McKenney et al., 2014; Schlager et al., 2014; K. Zhang et al., 2017a; Reck-Peterson et al., 2018; Olenick & Holzbaur, 2019; Qiu et al., 2019; Elshenawy et al., 2020; Htet et al., 2020; Marzo et al., 2020) and regulatory elements of the dynein motor domain (Bhabha et al., 2016; Carter et al., 2016). Most relevant to our current study is a detailed understanding on the role of AAA3 ATP hydrolysis in dynein's mechanochemical cycle (Bhabha et al., 2014; DeWitt et al., 2015; Nicholas et al., 2015). However, as the dynein motor mechanism is mainly studied in vitro by using the dynein motor domain itself, it has been unclear whether and how dynein's mechanochemical cycle itself is linked to dynein activation in vivo. In this current study, we first found that an *A. nidulans* AAA4 arginine-finger

mutation (implicated in AAA3 ATP hydrolysis) of dynein allows the function of NudF/LIS1 in dynein activation to be bypassed. This led us to further study the behaviors of wA-AAA3 (ATP-binding mutant) and wB-AAA3 (ATP-hydrolyzing mutant) dynein in the context of dynactin, the cargo adapter HookA and NudF/LIS1 in vivo. Our data suggest that the requirement for dynein activation is altered in both AAA3 mutants but more significantly in the wB-AAA3 mutant. In wild-type cells, dynein accumulates at the microtubule plus end and its activation needs both LIS1 and HookA, in addition to dynactin (Xiang & Qiu, 2020). In contrast, the wB-AAA3 dynein can be activated by dynactin in the absence of LIS1 or HookA, as evidenced by its dynactin-dependent microtubule minus-end accumulation in the *nudF/lis*1 or *hookA* deletion mutant. Thus, AAA3 ATP hydrolysis is needed for preventing dynein from getting fully activated before binding its early endosome cargo. Moreover, our data also suggest that the wA-AAA3 mutation stabilizes the dynein-LIS1 interaction, suggesting that the dynamic nature of the dynein-LIS1 interaction, which has been observed both in vivo (Lenz et al., 2006; Egan et al., 2012) and in vitro (Jha et al., 2017; Elshenawy et al., 2020; Htet et al., 2020), is regulated by ATP binding at AAA3.

The wB-AAA3 dynein does not accumulate at the microtubule plus end, but rather, it shows a strong accumulation at the minus end, even in the absence of NudF/LIS1 or the cargo adapter HookA. The absence of the plus-end accumulation is most likely due to an unscheduled activation of the wB-AAA3 dynein, as its plus-end accumulation can be partially restored if dynein activity is compromised by a dynein tail mutation. The unscheduled activation may also weaken the kinesin-1-dynein interaction if kinesin-1 preferentially transports dynein in the autoinhibited phi conformation (Xiang & Qiu, 2020); or, the mutant dynein may still bind kinesin-1 but overpowers it (Belyy et al., 2016). In the budding yeast, where dynein is exclusively required for spindle orientation (Winey & Bloom, 2012), plus-end dynein accumulation depends on Pac1/LIS1, Bik1/Clip170 and Kip2/kinesin-7 (Lee et al., 2003; Sheeman et al., 2003; Carvalho et al., 2004; Markus et al., 2009; Markus et al., 2011; Roberts et al., 2014). In filamentous fungi, LIS1 is not as important as dynactin for the plus-end accumulation of dynein (J. Zhang et al., 2003; Lenz et al., 2006; J. Zhang et al., 2008; Egan et al., 2012; Yao et al., 2012). This makes the activating function of LIS1 to be much more easily observed upon cargo-adapter-mediated dynein activation (compared to Num1-mediated activation in yeast (Lammers & Markus, 2015)), as the lack of NudF/LIS1 prevents dynein from leaving the plus end (Qiu et al., 2019). In addition, while the *A. nidulans* AAA4 arginine-finger mutation or the wB-AAA3 mutation suppresses the growth defect of the Δ*nudF/lis*1 mutant, the yeast AAA4 arginine-finger mutation does not suppress the Δ*pac*1/*lis1* mutant phenotype, because the additional function of Pac1/LIS1 in plus-end dynein accumulation cannot be bypassed (Huang et al., 2012). In mammalian cells or if mammalian dynein is used in vitro, LIS1 also plays an important role in dynein's plus-end accumulation (although dynactin is needed too) (Splinter et al., 2012; Baumbach et al., 2017; Jha et al., 2017). Thus, it may be easier to test if the mammalian wB-AAA3 dynein bypasses the requirement of LIS1 in an experimental setting where plus-end dynein accumulation is not a complicating factor. Despite the differences in the mechanisms of plus-end dynein localization, the dynein motor mechanism is most likely conserved in different organisms. Consistent with previous results on the yeast wB-AAA3 dynein motor domain (Cho et al., 2008; Bhabha et al., 2014; DeWitt et al., 2015; Nicholas et al., 2015; DeSantis et al., 2017; X. Liu et al., 2020), the wB-AAA3 mutation also significantly inhibits the in vitro motility of full-length dynein in *A. nidulans* extract (Figure S4).

In this study, we found that the wB-AAA3 mutation allows dynein to bypass the requirement of NudF/LIS1, consistent with its suppression on the growth defect of the Δ*nudF* (lis1) mutant colony, an effect similarly caused by the AAA4 arginine-finger mutation or the phi mutation (Willins et al., 1997; Zhuang et al., 2007; Huang et al., 2012; Qiu et al., 2019). As recent data link the function of LIS1 to promoting the open state of dynein (Qiu et al., 2019; Elshenawy et al., 2020; Htet et al., 2020; Marzo et al., 2020; McKenney, 2020), our current result suggest that the wB-AAA3 dynein is also open. In the wB-AAA3 mutant, the linker is locked at one position during the AAA1 ATPase

cycle, with its N-terminus docked at AAA5 (Bhabha et al., 2014). This is in contrast to wild-type dynein in which the linker bends upon ATP binding to AAA1, with its N-terminus close to AAA2/AAA3 and/or other positions (Bhabha et al., 2014). The phi conformation of dynein is maintained by multiple interactions, and linkers in the phi dynein (PDB code: 5NUG) are bent (K. Zhang et al., 2017a). Thus, if the linker is docked at AAA5, the phi structure may not be able to form, which implies that the wB-AAA3 dynein is unable to form the phi structure. However, the open conformation of the wB-AAA3 dynein must be different from that of the phi mutant dynein, which is open due to the weakened AAA4-linker ionic interaction between two dynein heavy chains within the dimer (K. Zhang et al., 2017a). The phi mutant dynein also strongly accumulates at the microtubule minus ends at centrosomes in mammalian cells and at the septa in *A. nidulans* (K. Zhang et al., 2017a; Qiu et al., 2019). However, the phi mutant dynein co-localizes with the early endosome cargo at the septa (Qiu et al., 2019), while the wB-AAA3 dynein does not do so. Although it is hard to completely exclude the possibility that other cargo adapters may be responsible for this minus-end localization, our data are most easily explained by the possibility that blocking AAA3 ATP hydrolysis allows the requirement of cargo adapter for dynein activation to be bypassed. The function of the cargo adapter has been implicated in enhancing the dynein-dynactin interaction (Splinter et al., 2012; McKenney et al., 2014; Schlager et al., 2014; Olenick et al., 2016). In fungi, however, the dynein-dynactin interaction clearly occurs before cargo-adapter-mediated dynein activation (Markus et al., 2020; Xiang & Qiu, 2020), especially since dynactin is absolutely required for dynein's plus-end accumulation in filamentous fungi (J. Zhang et al., 2003; Lenz et al., 2006; J. Zhang et al., 2008; Egan et al., 2012; Yao et al., 2012) and dynein is required for the plus end-accumulation of dynactin in the budding yeast (Moore et al., 2008). Thus, the function of the cargo adapter must be needed for changing the binding configuration of the two complexes so that the tails and motor domains of the dynein dimer can assume a parallel configuration (K. Zhang et al., 2017a). Given the special linker position of the wB-AAA3 dynein (Bhabha et al., 2014), could the dynactin complex be sufficient for making the wB-AAA3 dynein parallel? In this context, we would also point out that in some previous in vitro experiments, dynactin alone can indeed cause or enhance dynein processivity in a moderate way (S. J. King & Schroer, 2000; Ross et al., 2006; Kardon et al., 2009; Tripathy et al., 2014). Alternatively, could there be an in vivo regulator that reinforces the inactive state of dynein-dynactin before cargo binding and the wB-AAA3 mutation prevents dynein-dynactin from interacting with this regulator? Future work will clearly be needed to address these interesting questions.

LIS1 tends to dissociate from the dynein-dynactin-cargo adapter complex in fungal hyphae and in vitro assays (Lenz et al., 2006; Egan et al., 2012; Jha et al., 2017; Qiu et al., 2019; Elshenawy et al., 2020; Htet et al., 2020), although it co-migrates with the dynein-dynactin-cargo adapter complex in other in vitro assays (Baumbach et al., 2017; Gutierrez et al., 2017). In *A. nidulans*, the notion of LIS1-dynein dissociation after dynein activation explains the observation that LIS1 does not co-localize with activated dynein at the septal minus ends (Qiu et al., 2019). Interestingly, if AAA3 ATP binding is blocked, NudF/LIS1 strongly accumulate at the minus ends just like activated dynein, suggesting that the AAA3 ATPase cycle regulates the dynamic dynein-LIS1 interaction in vivo. It is possible that the abnormally increased dynein-LIS1 affinity contributes to the functional defect of the wA-AAA3 dynein, since reducing the NudF/LIS1 level in *A. nidulans* by using a *nudF*6 mutant grown at a semi-permissive temperature improves the growth of the wA-AAA3 mutant (Figure 2D). Based on this current study and previous studies from several labs, we propose that the AAA3 ATPase cycle is closely linked to the spatial regulation of dynein activation in vivo in the following ways: First, during kinesin-1-mediated transport to the microtubule plus end, dynein is in the autoinhibited phi conformation with an ADP-bound AAA3 (Figure 8). Second, LIS1 may preferentially bind dynein with an apo-AAA3 (note that LIS1 binding is not compatible with the phi dynein conformation (Htet et al., 2020; Marzo et al., 2020)). Third, ATP binding to AAA3 occurs after cargo binding, as otherwise dynein will be activated by dynactin alone without the cargo. Finally, after cargo-mediated dynein activation, LIS1 may dissociate at any step except when AAA3

is in the apo state. We should point out that our current study is done in the context of not only cargo adapter and dynactin, but also NudE, another dynein regulator that binds both LIS1 and dynein (Efimov & Morris, 2000; Feng et al., 2000; Niethammer et al., 2000; Sasaki et al., 2000; Efimov, 2003; Yan et al., 2003; Liang et al., 2004; Shu et al., 2004; Li et al., 2005; Stehman et al., 2007; McKenney et al., 2010; McKenney et al., 2011; Wang & Zheng, 2011; Zylkiewicz et al., 2011; Wang et al., 2013). Previous work on purified mammalian dynein showed that while LIS1 only binds dynein when ATP and vanadate are added, the inclusion of NudE allows LIS1-dynein binding in the absence of nucleotide (McKenney et al., 2010). Although the wA-AAA3 mutation clearly alters the dynamic LIS1-dynein interaction in *A. nidulans*, our hypothesis that LIS1 preferentially binds dynein with an apo-AAA3 before cargo binding will need to be further examined in the presence and absence of NudE.

In summary, our results from *A. nidulans* suggest that ATP binding and hydrolysis at AAA3 are important for different aspects of cargo-mediated dynein activation in vivo, including the dynamic dynein-LIS1 interaction, as well as the coordination between cargo binding and dynein activation. The mechanisms of dynein regulation, *e.g.*, kinesin-mediated localization of dynein-dynactin to the microtubule plus-end (J. Zhang et al., 2003; Carvalho et al., 2004; Lenz et al., 2006; Arimoto et al., 2011; Egan et al., 2012; Yao et al., 2012; Twelvetrees et al., 2016), the importance of the plus-end dynein-dynactin in cargo transport (P. S. Vaughan et al., 2002; Lenz et al., 2006; Abenza et al., 2009; Zekert & Fischer, 2009; J. Zhang et al., 2010; Egan et al., 2012; Lloyd et al., 2012; Moughamian & Holzbaur, 2012; Splinter et al., 2012; Moughamian et al., 2013), the mechanism of dynein-cargo interaction and cargo transport, are all conserved in principle from fungi to higher eukaryotic cells especially elongated neurons (J. Zhang et al., 2011; Yeh et al., 2012; Bielska et al., 2014; Yao et al., 2014; J. Zhang et al., 2014; Guo et al., 2016; Reck-Peterson et al., 2018; Olenick et al., 2019; Olenick & Holzbaur, 2019). Thus, it will be of interest to test whether the wA-AAA3 and wB-AAA3 mutations cause similar effects in higher eukaryotic cells, especially neurons. It will also be highly worthwhile to examine purified mammalian dynein containing the wA-AAA3 or the wB-AAA3 mutation to further dissect the mechanism of the AAA3-involved dynein regulation in vitro in the context of other known factors involved in dynein activation.

Materials and Methods

***A. nidulans* strains and media**

A. nidulans strains used in this study are listed in Table 1. Solid rich medium was made of either YAG (0.5% yeast extract and 2% glucose with 2% agar) or YAG+UU (YAG plus 0.12% uridine and 0.12% uracial). Genetic crosses were done by standard methods. Solid minimal medium containing 1% glucose was used for selecting progeny from a cross. For live cells imaging experiments, cells were cultured in liquid minimal medium containing 1% glycerol for overnight at 32°C. All the biochemical analyses were done using cells grown at 32°C for overnight in liquid YG rich medium (0.5% yeast extract and 2% glucose). For a few experiments using strains containing the *alcA-nudK*[Arp1] (Arp1 conditional null) allele, we harvest spores from the solid minimal medium containing 1% glycerol and cultured them in liquid minimal medium containing 1% glucose for imaging analysis.

Live cell imaging and analyses

All images were captured using an Olympus IX73 inverted fluorescence microscope linked to a PCO/Cooke Corporation Sensicam QE cooled CCD camera. An UPlanSApo 100x objective lens (oil) with a 1.40 numerical aperture was used. A filter wheel system with GFP/mCherry-ET Sputtered series with high transmission (Biovision Technologies) was used. For all images, cells were grown in the LabTek Chambered #1.0 borosilicate coverglass system (Nalge Nunc

International, Rochester, NY). The IPLab software was used for image acquisition and analysis. Image labeling was done using Microsoft PowerPoint and/or Adobe Photoshop. For quantitation of signal intensity, a region of interest (ROI) was selected and the Max/Min tool of the IPLab program was used to measure the maximal intensity within the ROI. The ROI box was then dragged to a nearby area inside the cell to take the background value, which was then subtracted from the intensity value. Hyphae were chosen randomly from images acquired under the same experimental conditions. For measuring the signal intensity of a MT plus-end comet formed by GFP-dynein or NudF/LIS1-GFP proteins, only the comet closest to hyphal tip was measured. For measuring GFP-dynein signal intensity at septa, usually only the septum most proximal to the hyphal tip was measured. Images were taken at room temperature immediately after the cells were taken out of the incubators. Cells were cultured overnight in minimal medium with 1% glycerol and supplements at 32°C or 37°C (all experiments using *nudF*6 strains and controls were done at 37°C). Note that the *nudF*6 mutant is temperature sensitive; it forms a tiny colony lacking asexual spores at a higher temperature (typical of a *nud* mutant), but some spores are produced at its semipermissive temperature of 32°C. Thus, for experiments involving *nudF*6, we harvested spores at 32°C and cultured them at 37°C for imaging analysis.

Constructing the wA-AAA3 (*nudA*^{K2599A}) and wB-AAA3 (*nudA*^{E2663Q}) mutants and other strains containing the mutant alleles

For making the AAA3 Walker A (wA-AAA3) mutant *nudA*^{K2599A}, we used fusion PCR to make a DNA fragment containing the K2599A mutation with the following primers: WAF (5'-TTCTGGTGCAACCATGACACTGTTTGCCG-3'), WAR (5'-GTGTCATGGTTGC ACCAGAACCGGGAGGAC-3'), NudA58(Zhuang et al., 2007) and NdA310 (Zhuang et al., 2007). For making the AAA3 Walker B (wB-AAA3) mutant *nudA*^{E2663Q}, we used fusion PCR to make a DNA fragment containing the E2663Q mutation with the following primers: WBF (5'-ATCTTCTGCGATCAAATCAACCTGCCGGCTC-3'), WBR (5'-CAGGTTGATTTG ATCGCAGAAGATAACCAGCCAA-3'), NudA58 (Zhuang et al., 2007) and NdA310 (Zhuang et al., 2007). Each fragment was co-transformed with a selective marker *pyrG* fragment into the RQ2 strain containing GFP-dynein HC (NudA), mCherry-RabA and the Ku70 deletion allele Δ*nkuA-argB*, which facilitates the selection of transformants in which transformed DNA fragments underwent homologous integration into the genome. Several transformants with growth defect were selected and the mutations confirmed by sequencing analysis of the genomic DNA fragments. For making the strains containing p150-GFP and the wA-AAA3 or the wB-AAA3 allele, the fusion PCR product containing the wA-AAA3 or the wB-AAA3 mutation was co-transformed with the p150-GFP-*AfpyrG* fragment (J. Zhang et al., 2011) into the XY42 strain containing mCherry-RabA and Δ*nkuA-argB.* Transformants with growth defect were selected and the mutations confirmed by sequencing analysis.

Strains containing multiple mutant alleles were constructed by genetic crosses. Genetic crosses were done by standard methods, and progeny with desired genotypes were selected based on colony phenotype, imaging analysis, Western analysis, diagnostic PCR, and/or sequencing of specific regions of the genomic DNA.

In vitro dynein motility assay using *A. nidulans* extract

About 0.25 g of hyphal mass was harvested from overnight culture for each sample, and cell extract was prepared using 150 μl of lysis buffer (30 mM HEPES pH 7.3, 50 mM potassium acetate, 2 mM magnesium acetate, 1 mM EGTA, 10% glycerol, 1 mM DTT, and 0.2 mM ATP). Cell extracts were then centrifuged at 20,000 *g* for 30min at 4°C to remove cell debris, and supernatant was kept on ice for motility assays, or snap-frozen with liquid nitrogen and stored at -80°C if motility assays cannot be performed on the same day. Immediately prior to imaging, the cell lysates were warmed to room temperature and diluted 10 times in the lysis buffer containing 1 mM ATP, 1 mM

Trolox, 20 μM Taxol, 10 mM DTT, 1 mg/mL BSA, 1 mg/mL β-casein, 15 μg/mL catalase, 40 μM glucose, and 40 μg/mL glucose oxidase.

To make X-rhodamine labeled microtubule, un-labeled tubulin (Cytoskeleton, Inc) and X-rhodamine-labeled tubulin (Cytoskeleton, Inc) are mixed with a ratio of ~12:1 in BRB80 buffer (80 mM PIPES, pH 7.0, 1 mM EGTA, and 1 mM $MgCl_2$) with 2 mM GTP. After incubating for 12 min at 35°C, the polymerized microtubule was stabilized by adding 20 μM taxol and stored at room temperature.

To prepare flow chambers, 75 mm x 25 mm glass microscope slides and 18 mm x 18 mm coverslips with #1.5 thickness were treated as described (Nicholas et al., 2014). In summary, both the microscope slides and coverslips were pre-cleaned with alkaline in bath sonicator followed by washing with ultrapure water and treating in plasma clear with RF coil set on "high" for 5 min. The pre-cleaned coverslips were further aminosilanized with 2% APTES (Sigma-Aldrich). To assemble the slide chamber, a set of parafilm strips were inserted between the aminosilane surface-coated coverslip and pre-cleaned slide to generate an approximate volume of 15-20 μL and the chamber was then formed after being heat sealed. The aminosilanized surface of the slide chamber was further activated with 8% glutaraldehyde (Sigma-Aldrich) to enhance microtubule binding prior to an experiment.

To image dynein motility on microtubule, 20 μL of ~0.1 mg/mL microtubule was flushed into the glutaraldehyde treated chamber from one side and incubate at room temperature for 90 seconds, 20 μL BRB80 with 20 μM taxol was then added from the same side and the chamber was incubated at room temperature for 20 min to allow microtubule binding. The flow chamber was then blocked with 2 mg/mL β-casein and 20 μM Taxol for 5 min, and the cell lysates were added in the presence of either 1 mM or 3 mM ATP. Time lapse movies were acquired immediately with either an Olympus IX73 inverted fluorescence microscope linked to a PCO/Cooke Corporation Sensicam QE cooled CCD camera or an Olympus IX83 microscope equipped with a one-line motorized IX3 cellTIRF illuminator and quad laser TIRF set, outfitted with an ORCA flash 4.0 v3 sCMOS camera (Hamamatsu), a 100x 1.50-NA objective, and CellSens software. A 488-nm laser and a 561-nm laser were used for acquiring images of GFP-labeled dynein and X-Rhodamine-labeled microtubules respectively. Several 1.5 to 2 min movies were acquired at room temperature per chamber with 1 second intervals of each frame. Velocities were measured by using kymographs generated by the Fiji/ImageJ software (the National institutes of Health).

Statistical analysis

All statistical analyses were done using GraphPad Prism 8 for Mac (version 8.0.0, 2018). The D'Agostino & Pearson normality test was performed on all data sets. For all data sets, non-parametric tests were used without assuming Gaussian distribution. Specifically, the Kruskal-Wallis ANOVA test (unpaired) with Dunn's multiple comparisons test was used for analyzing multiple data sets. The Mann-Whitney test (unpaired, two tailed) was used for analyzing the two data sets. Note that adjusted p values were generated from the Kruskal-Wallis ANOVA test with Dunn's multiple comparisons test. In all figures, **** indicates $p<0.0001$, *** indicates $p<0.001$, ** indicates $p<0.01$, * indicates $p<0.05$ *, and "ns" indicates $p>0.05$.

Acknowledgements

We thank Drs. Berl Oakley, Stephen Osmani and Miguel Peñalva for Aspergillus strains. This work was funded by the National Institutes of Health RO1 GM121850 (to X. Xiang) and RO1 GM134104 (to J. Rotty). J. Rotty was also supported by Uniformed Services University start-up fund. Disclaimer: The opinions and assertions expressed herein are those of the author(s) and do not necessarily reflect the official policy or position of the Uniformed Services University or the Department of Defense.

Table 1. *Aspergillus nidulans* strains used in this study

Strain	Genotype	Source
JZ383	$\Delta nudF::pyr4$; GFP-*nudA*^{HC}*; argB2::[argB*-alcAp*::mCherry-*RabA]*	(Qiu et al., 2019)
LZ36	GFP-*nudA*^{R3086C}; $\Delta nkuA::argB$; *pyroA4*	(Zhuang et al., 2007)
RQ2	GFP-*nudA*^{HC}*; argB2::[argB*-alcAp*::mCherry-*RabA]*; $\Delta nkuA::argB$; *pyrG89*; *pyroA4*; *yA2*	(Qiu et al., 2013)
RQ8	GFP-*nudA*^{F208V}*; argB2::[argB*-alcAp*::mCherry-*RabA]*; *pyrG89*; *wA2*	(Qiu et al., 2013)
RQ165	GFP-*nudF-AfpyrG*; *argB2::[argB*-alcAp*::mCherry-*RabA]*; $\Delta nkuA::argB$; *pyroA4*; *pyrG89*; *wA2*	(Qiu et al., 2019)
RQ275	*nudF6*; *gpdA-ΔC-hookA-S-AfpyrG*; GFP-*nudA*^{HC}; *pyrG89*; *yA2*; possibly $\Delta nkuA::argB$	(Qiu et al., 2019)
RQ287	GFP-*nudA*^{HC}; *gpdA-ΔC-hookA-S-AfpyrG*; *yA2*	(Qiu et al., 2019)
XX60	$\Delta nudA::pyrG$; *pyrG89*	(Xiang et al., 1995b)
XX222	GFP-*nudA*^{HC}*; argB2::[argB*-alcAp*::mCherry-*RabA]*; *yA2*; *pantoB100*	(J. Zhang et al., 2010)
XX514	GFP-*nudA*^{HC}*; ΔyA::NLS-DsRed; possibly $\Delta nkuA::argB$	(Qiu et al., 2019)
XX634	$\Delta nudF::pyr4$; GFP-*nudA*$^{R1602E,\ K1645E}$; *argB2::[argB*-alcAp*::mCherry-*RabA]*; possibly $\Delta nkuA::argB$	(Qiu et al., 2019)
XY13	p150-GFP-*AfpyrG*; $\Delta nkuA::argB$; *pyrG89*; *pyroA4*	(Yao et al., 2012)
XY42	*argB2::[argB*-alcAp*::mCherry-*RabA]*; $\Delta nkuA::argB$; *pyrG89*; *yA2*; *pantoB100*	(Qiu et al., 2018)
JZ504	GFP-*nudA*^{HC}; $\Delta hookA-AfpyrG$; *argB2::[argB*-alcAp*::mCherry-RabA]*; *pyroA4*	This work
RQ102	GFP-*nudA*^{HC}; $\Delta hookA-AfpyrG$; *argB2::[argB*-alcAp*::mCherry-RabA]*; *pantoB100*; *yA2*	This work
RQ293	*nudF6*; *gpdA-ΔC-hookA-S-AfpyrG*; GFP-*nudA*^{R3086C}	This work
RQ295	*gpdA-ΔC-hookA-S-AfpyrG*; GFP-*nudA*^{R3086C}	This work
RQ300	GFP-*nudA*^{K2599A}(wA-AAA3); *argB2::[argB*-alcAp*::mCherry-*RabA]*; $\Delta nkuA::argB$; *pyroA4*; *yA2*	This work
RQ301	GFP-*nudA*^{E2663Q}(wB-AAA3); *argB2::[argB*-alcAp*::mCherry-*RabA]*; $\Delta nkuA::argB$; *pyroA4*; *yA2*	This work
RQ306	$\Delta nudF::pyr4$; GFP-*nudA*^{K2599A}(wA-AAA3); *argB2::[argB*-alcAp*::mCherry-RabA]*	This work
RQ307	$\Delta nudF::pyr4$; GFP-*nudA*^{E2663Q}(wB-AAA3); *argB2::[argB*-alcAp*::mCherry-RabA]*	This work
RQ309	GFP-*nudA*^{K2599A}(wA-AAA3); *gpdA-ΔC-hookA-S-AfpyrG*	This work
RQ311	GFP-*nudA*^{K2599A}(wA-AAA3)*; ΔyA::NLS-DsRed; possibly $\Delta nkuA::argB$	This work
RQ312	GFP-*nudA*^{E2663Q}(wB-AAA3); *gpdA-ΔC-hookA-S-AfpyrG*	This work
RQ314	GFP-*nudA*^{E2663Q}(wB-AAA3)*; ΔyA::NLS-DsRed; possibly $\Delta nkuA::argB$	This work
RQ317	GFP-*nudA*^{K2599A}(wA-AAA3); *gpdA-ΔC-hookA-S-AfpyrG*; *argB2::[argB*-alcAp*::mCherry-RabA]*	This work
RQ318	GFP-*nudA*^{K2599A}(wA-AAA3); *gpdA-ΔC-hookA-S-AfpyrG*; *nudF6*	This work

RQ323	*nudA*[K2599A](wA-AAA3); p150-GFP-*AfpyrG*; Δ*nkuA::argB; argB2::[argB*-alcAp*::mCherry-*RabA]; pantoB100*; *yA2*	This work
RQ324	*nudA*[E2663Q](wB-AAA3); p150-GFP-*AfpyrG*; Δ*nkuA::argB; argB2::[argB*-alcAp*::mCherry-*RabA]; pantoB100*; *yA2*	This work
RQ328	GFP-*nudA*[E2663Q](wB-AAA3); *argB2::[argB*-alcAp*::mCherry-*RabA]*; Δ*hookA-AfpyrG*	This work
RQ329	GFP-*nudA*[K2599A](wA-AAA3); *argB2::[argB*-alcAp*::mCherry-*RabA]*; Δ*hookA-AfpyrG*	This work
RQ330	GFP-*nudA*[K2599A](wA-AAA3); *argB2::[argB*-alcAp*::mCherry-*RabA]*	This work
RQ331	GFP-*nudA*[E2663Q](wB-AAA3); *argB2::[argB*-alcAp*::mCherry-*RabA]*	This work
RQ332	GFP-*nudA*[K2599A](wA-AAA3); *alcAp-nudK*[Arp1]-*pyr4;* possibly *argB2::[argB*-alcAp*::mCherry-*RabA];* possibly *p25-S*	This work
RQ333	GFP-*nudA*[E2663Q](wB-AAA3); *alcAp-nudK*[Arp1]-*pyr4;* possibly *argB2::[argB*-alcAp*::mCherry-*RabA];* possibly *p25-S*	This work
RQ343	GFP-*nudA*[K2599A, F208V]*; argB2::[argB*-alcAp*::mCherry-*RabA];* Δ*nkuA::argB; pyroA4*	This work
RQ344	GFP-*nudA*[E2663Q, F208V]*; argB2::[argB*-alcAp*::mCherry-*RabA];* Δ*nkuA::argB; pyroA4*	This work
RQ354	GFP-*nudF-AfpyrG; nudA*[K2599A](wA-AAA3); *argB2::[argB*-alcAp*::mCherry-*RabA];* possibly Δ*nkuA::argB; pyrG89*	This work
RQ355	GFP-*nudF-AfpyrG; nudA*[E2663Q](wB-AAA3); *argB2::[argB*-alcAp*::mCherry-*RabA];* possibly Δ*nkuA::argB; pyrG89*	This work
XX632	GFP-*nudA*[R3086C]; Δ*nudF::pyr4; argB2::[argB*-alcAp*::mCherry-*RabA];* possibly Δ*nkuA::argB;* possibly *pyrG89*	This work
XX644	GFP-*nudA*[K2599A](wA-AAA3); *nudF6; argB2::[argB*-alcAp*::mCherry-*RabA];* possibly Δ*nkuA::argB*	This work
XX684	GFP-*nudF-AfpyrG; nudA*[K2599A](wA-AAA3); *gpdA-*ΔC-*hookA-S;* possibly Δ*nkuA::argB;* possibly *pyrG89;* possibly *argB2::[argB⁻alcAp*::mCherry-*RabA]*	This work
XX689	GFP-*nudF-AfpyrG; nudA*[E2663Q](wB-AAA3); *gpdA-*ΔC-*hookA-S;* possibly Δ*nkuA::argB;* possibly *pyrG89;* possibly *argB2::[argB⁻alcAp*::mCherry-*RabA]*	This work

Figure legends:

Figure 1. The AAA4 arginine-finer mutation (RF-AAA4 or *nudA*[R3086C]) allows NudF/LIS1 function to be bypassed during ΔC-HookA-mediated dynein activation. (A) Images of GPF-dynein are shown for a strain containing RF-AAA4, a strain containing RF-AAA4, *gpdA-ΔC-hookA*-S, a strain containing *nudF*6, RF-AAA4, *gpdA-ΔC-hookA*-S, and a strain containing *nudF*6, *gpdA-ΔC-hookA*-S. The strain containing *nudF*6, *gpdA-ΔC-hookA*-S was used as a control, as it shows plus-end comets formed by wild-type dynein. Bright-field images are shown below to indicate hyphal shape and position of septum. Hyphal tip is indicated by a yellow arrowhead and septum by a light brown arrow. Bar, 5 μm. (B) A quantitative analysis on dynein comet intensity in the *nudF*6, *gpdA-ΔC-hookA*-S strain (n=58), the RF-AAA4, *gpdA-ΔC-hookA*-S strain (n=42) and the *nudF*6, RF-AAA4, *gpdA-ΔC-hookA*-S strain (n=40). The average value for the *nudF*6, *gpdA-ΔC-hookA*-S strain is set as 1. Scatter plots with mean and S.D. values were generated by Prism 8. ****p<0.0001; ns, non-significant or p>0.05 (Kruskal-Wallis ANOVA test with Dunn's multiple comparisons test, unpaired).

Figure 2. Both the wA-AAA3 and wB-AAA3 mutations affect dynein function, and the wB-AAA3 mutation suppresses the Δ*nudF* (lis1) growth defect. (A) Colony phenotypes of the wA-AAA3 and wB-AAA3 mutants in comparison with the wild type, RF-AAA4, Δ*nudA* and Δ*nudF* strains. The plate was incubated at 32°C for 3 days. (B) Images of nuclei labeled with NLS-DsRed in wild type, the wA-AAA3 mutant and the wB-AAA3 mutant. Spore head is indicated by a brown arrow. Bar, 5 μm. (C) A quantitative analysis on the percentage of germ tubes containing 0, 1, 2, 3, 4 or 5 nuclei in the spore head (wild type: n=54; wA-AAA3: n=34; wB-AAA3: n=28). The number of nuclei in the spore head of the wA-AAA3 mutant or the wB-AAA3 mutant is higher than that in wild type (p<0.0001 in both cases, Kruskal-Wallis ANOVA test with Dunn's multiple comparisons test, unpaired). However, the number of nuclei in the spore head of the wA-AAA3 mutant is not significantly different from that in the wB-AAA3 mutant (p>0.99, Kruskal-Wallis ANOVA test with Dunn's multiple comparisons test, unpaired). (D) Colony phenotypes of various single and double mutant strains. The plate was incubated at 32°C for 3 days.

Figure 3. The wA-AAA3 or wB-AAA3 dynein partially accumulates at the microtubule minus ends at septa, and this accumulation requires dynein functionality. (A) Localization of GFP-dynein and early endosomes (mCherry-RabA) in wild type, the wA-AAA3 mutant and the wB-AAA3 mutant. Bright-field images are shown below to indicate hyphal shape and position of septum. Hyphal tip is indicated by a yellow arrowhead and septum by a light brown arrow. (B) A quantitative analysis on septal dynein intensity in wild type (n=39), wA-AAA3 (n=39), and wB-AAA3 (n=30) strains. The average value for the wB-AAA3 strain is set as 1. Scatter plots with mean and S.D. values were generated by Prism 8. ****p<0.0001, ***p<0.001, **p<0.01 (Kruskal-Wallis ANOVA test with Dunn's multiple comparisons test, unpaired). (C) The dynein tail mutation $nudA$F^{208V} diminishes the septal accumulation of the wA-AAA3 or wB-AAA3 mutant dynein. Bright-field images are shown below to indicate hyphal shape and position of septum. Hyphal tip is indicated by a yellow arrowhead and septum by a light brown arrow. Bars, 5 μm.

Figure 4. Different effects of the cargo adapter HookA on the localization of wild-type dynein, wA-AAA3 dynein and wB-AAA3 dynein. (A) Wild-type dynein, wA-AAA3 dynein and wB-AAA3 dynein in the *gpdA-ΔC-hookA*-S (overexpression of ΔC-HookA) background. Bright-field images are shown below to indicate hyphal shape and position of septum. Hyphal tip is indicated by a yellow arrowhead and septum by a light brown arrow. Bar, 5 μm. (B) Wild-type dynein, wA-AAA3 dynein and wB-AAA3 dynein in the Δ*hookA* mutant background. Bright-field images are shown below to indicate hyphal shape and position of septum. Hyphal tip is indicated by a yellow arrowhead and septum by a light brown arrow. (C) A quantitative analysis on comet intensity of wild type (n=39), wA-AAA3 (n=38), and wB-AAA3 (n=50) dynein in the Δ*hookA* background. The average value for wild-type dynein in the Δ*hookA* strain is set as 1. Scatter plots with mean and S.D. values were generated by Prism 8. ****p<0.0001 (Kruskal-Wallis ANOVA test with Dunn's multiple comparisons test, unpaired). (D) A quantitative analysis on septal intensity of wild type (n=39), wA-AAA3 (n=38), and wB-AAA3 (n=50) dynein in the Δ*hookA* background. The average value for wB-AAA3, Δ*hookA* strain is set as 1. Scatter plots with mean and S.D. values were generated by Prism 8. ****p<0.0001, ***p<0.001 (Kruskal-Wallis ANOVA test with Dunn's multiple comparisons test, unpaired). (E) A quantitative analysis on septal intensity of wA-AAA3 in the wild-type (n=39) and the Δ*hookA* (n=38) backgrounds. Scatter plots with mean and SD values were generated by Prism 8. ***p<0.001 (unpaired, Mann-Whitney test, Prism 8). (F) A quantitative analysis on septal intensity of wB-AAA3 in the wild-type (n=30) and the Δ*hookA* (n=50) backgrounds. Scatter plots with mean and SD values were generated by Prism 8. ns, non-significant or p>0.05 (unpaired, Mann-Whitney test, Prism 8).

Figure 5. Dynactin is required for the septal accumulation of the wA-AAA3 or wB-AAA3 dynein. (A) Images of GFP-labeled wA-AAA3 dynein and wB-AAA3 dynein in the wild type or the Arp1 conditional-null (*alcA-nudK*^Arp1^) background. The *alcA* promoter is active on glycerol but repressed by glucose, and thus, the expression of Arp1 is shut off in strains containing the *alcA-nudK*^Arp1^ allele grown on glucose. The strains without the *alcA-nudK*^Arp1^ allele grown on glycerol or glucose were shown as controls. Bright-field images are shown below to indicate hyphal shape and position of septum. Hyphal tip is indicated by a yellow arrowhead and septum by a light brown arrow. Bar, 5 μm. (B) A quantitative analysis on septal intensity of the wB-AAA3 strain (n=31) and the wB-AAA3, *alcA-nudK*^Arp1^ strain (n=35) grown on glucose. Scatter plots with mean and SD values were generated by Prism 8. ****p<0.0001 (unpaired, Mann-Whitney test, Prism 8).

Figure 6. NudF/LIS1 is dispensable for the septal accumulation of the wB-AAA3 dynein but not the wA-AAA3 dynein, but overexpression of ΔC-HookA allows the wA-AAA3 mutation to partially bypass the requirement of NudF/LIS1. (A) Images of GFP-labeled wild-type dynein, wA-AAA3 dynein and wB-AAA3 dynein in the Δ*nudF* background. Bright-field images are shown below to indicate hyphal shape and position of septum. Hyphal tip is indicated by a yellow arrowhead and septum by a light brown arrow. Bar, 5 μm. (B) Images of GFP-labeled wA-AAA3 dynein or wild-type dynein in the *gpdA*-ΔC-*hookA*-S or the *nudF*6, *gpdA*-ΔC-*hookA*-S background. (C) (D) Quantitative analysis on comet intensity (C) and septal intensity (D) of the wA-AAA3 dynein in the *gpdA*-ΔC-*hookA*-S (n=31) or the *nudF*6, *gpdA*-ΔC-*hookA*-S background (n=40) and that of wild-type dynein in the *nudF*6, *gpdA*-ΔC-*hookA*-S background (n=42). Scatter plots with mean and SD values were generated by Prism 8. ****p<0.0001, **p<0.01; ns, non-significant or p>0.05 (unpaired, Mann-Whitney test, Prism 8).

Figure 7. NudF/LIS1 accumulates at the septa in the wA-AAA3 mutant but not in the wB-AAA3 mutant. (A) Images of NudF/LIS1-GFP localization in wild-type, the wA-AAA3 mutant and the wB-AAA3 mutant. Bright-field images are shown below to indicate hyphal shape and position of septum. Hyphal tip is indicated by a yellow arrowhead and septum by a light brown arrow. Bar, 5 μm. (B) Images of NudF/LIS1-GFP localization in wild-type, the wA-AAA3 mutant and the wB-AAA3 mutant in the *gpdA-ΔC-hookA*-S background. (C). A quantitative analysis on septal intensity of LIS1-GFP in the backgrounds of *gpdA-ΔC-hookA*-S (n=35), wA-AAA3, *gpdA-ΔC-hookA*-S (n=25), wA-AAA3 (n=30), wB-AAA3 (n=31), and wB-AAA3, *gpdA-ΔC-hookA*-S (n=31). The average value of the LIS1-GFP septal intensity in the wA-AAA3, *gpdA-ΔC-hookA*-S strain is set as 1. Scatter plots with mean and S.D. values were generated by Prism 8. ****$p<0.0001$, ***$p<0.001$, **$p<0.01$, ns, non-significant or $p>0.05$ (Kruskal-Wallis ANOVA test with Dunn's multiple comparisons test, unpaired).

Figure 8. A hypothetic model of dynein activation involving the AAA3 ATPase cycle. Note that we only focus on a series of events at the vicinity of the microtubule plus end. During kinesin-1-mediated transport of dynein-dynactin to the microtubule plus end, dynein is most likely in the autoinhibited phi conformation. In the phi dynein, both AAA1 and AAA3 are occupied by ADP, and the linker is in a bent conformation (PDB code: 5NUG) (K. Zhang et al., 2017a). Two different states (either apo or ATP at both AAA1 and AAA3) with the linkers docked on AAA5 (Schmidt et al., 2012; Bhabha et al., 2014) are shown at the plus end. LIS1 may preferentially bind dynein with apo-AAA3, which is followed by cargo binding to cause dynein activation. ATP binding to AAA3 happens after cargo-mediated dynein activation, because otherwise dynein would be activated by dynactin alone without the cargo. During the minus-end-directed movement of the dynein-dynactin-cargo adapter complex, AAA3 is ADP-bound most of the time, and the linker is flexible and its position would depend on the AAA1 nucleotide state (Bhabha et al., 2014; DeWitt et al., 2015). Finally, LIS1 can dissociate from the activated dynein-dynactin-cargo adapter complex at any step as long as the AAA3 is not in the apo state.

Online Supplemental Materials:

Figure S1 A part of the sequence alignment of dynein heavy chains from *A. nidulans* (An), budding yeast (Sc) and human (Hs). The alignment was done using Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/). Residues that are identical (*), strongly similar (:) or weakly similar (.) are indicated. The Lysine (K) of AAA3 Walker A and the Glutamic acid (E) of AAA3 Walker B are shown using bold characters.

```
An  QVPTIDVNTHSITQTDVVIPTVDTVRHEDVLYSWLAEHKPLLLCGPPGSGKTMTLFAALR 2608
Sc  EIPSVSLEAHEVMRPDIVIPTIDTIKHEKIFYDLLNSKRGIILCGPPGSGKTMIMNNALR 2433
Hs  KVPQIEVETHKVAAPDVVVPTLDTVRHEALLYTWLAEHKPLVLCGPPGSGKTMTLFSALR 2610
    ::* :.:::*.:   *:*:**:**:** ::*  * .:: ::*********K*** :  ***

An  KLPNMEVVGLNFSSATTPDLLIKTFEQYCEYKKTLSGVVMSPNQIGRWLVIFCDEINLPA 2668
Sc  NSSLYDVVGINFSKDTTTEHILSALHRHTNYVTTSKGLTLLPKSDIKNLVLFCDEINLPK 2493
Hs  ALPDMEVVGLNFSSATTPELLLKTFDHYCEYRRTPNGVVLAPVQLGKWLVLFCDEINLPD 2670
      :***:***. ** : ::.:*.:: :*  * .*:.: * .  : **:***E*****

An  PDKYGTQRAISFLRQLVEQNGFWRTSDKTWVSLDRIQFVGACNPPTDAGRTPLAERFLRH 2728
Sc  LDKYGSQNVVLFLRQLMEKQGFWKTPENKWVTIERIHIVGACNPPTDPGRIPMSERFTRH 2553
Hs  MDKYGTQRVISFIRQMVEHGGFYRTSDQTWVKLERIQFVGACNPPTDPGRKPLSHRFLRH 2730
    ****:*..: *:**::*: **::* ::.**.::**::*********** ** *::.** **
```

Figure S2. The spindle-pole body (SPB) localization of the wB-AAA3 dynein on nuclei. The wB-AAA3 dynein is labeled with GFP. Nuclei are labeled with NLS-DsRed and pseudo-colored in blue. A wild-type hypha is shown as a control, in which GFP-dynein mainly forms microtubule plus-end comets. Bar, 5 μm.

Figure S3. Localization of dynactin p150-GFP in wild type, the wA-AAA3 mutant and the wB-AAA3 mutant. Bright-field images are shown below to indicate hyphal shape and position of septum. Hyphal tip is indicated by a yellow arrowhead and septum by a light brown arrow. Bar, 5 μm.

Figure S4. The wB-AAA3 mutation dramatically decreases the speed of dynein motility in vitro. (A) kymographs showing the in vitro movements of wild-type dynein and wB-AAA3 dynein. For wild-type dynein, we used cell extract from a strain containing the *gpdA*-ΔC-*hookA*-S fusion to ensure that GFP-dynein molecules predominantly move toward the minus ends. For wB-AAA3 dynein, we did not include the *gpdA*-ΔC-*hookA*-S fusion because the wB-AAA3 dynein accumulates at the minus ends even in the absence of HookA, and the wB-AAA3, *gpdA*-ΔC-*hookA*-S double mutant is very sick. (B) Velocity of wild-type dynein verses that of the wB-AAA3 dynein (n=26 for wild-type dynein; n=25 for the wB-AAA3 mutant dynein). Note that the wB-AAA3 dynein moves with an extremely low speed. The mean ranks of these two sets of data are significantly different (p<0.0001, unpaired, Mann-Whitney test, Prism 8). (C) Kymographs of the wB-AAA3 dynein movements in 3 mM ATP or 1 mM ATP. All the velocity data from the wB-AAA3 dynein were obtained in the in vitro motility assays with 3 mM ATP, although 1 mM ATP was used for the wild-type dynein. This is because the wB-AAA3 dynein was not motile if 1 mM ATP is used. (D) A kymograph showing the occasional slow movement of a p150-GFP signal toward a septum in vivo. The very bright line corresponds to the position of the septum (note that the wB-AAA3 mutation causes both dynein and dynactin to accumulate at septa).

References:

Abenza, J. F., Galindo, A., Pantazopoulou, A., Gil, C., de los Ríos, V., & Peñalva, M. A. (2010). Aspergillus RabB Rab5 integrates acquisition of degradative identity with the long distance movement of early endosomes. *Mol Biol Cell, 21*(15), 2756-2769. doi:10.1091/mbc.e10-02-0119

Abenza, J. F., Pantazopoulou, A., Rodriguez, J. M., Galindo, A., & Penalva, M. A. (2009). Long-distance movement of Aspergillus nidulans early endosomes on microtubule tracks. *Traffic, 10*(1), 57-75. doi:10.1111/j.1600-0854.2008.00848.x

Alberti-Segui, C., Dietrich, F., Altmann-Jöhl, R., Hoepfner, D., & Philippsen, P. (2001). Cytoplasmic dynein is required to oppose the force that moves nuclei towards the hyphal tip in the filamentous ascomycete Ashbya gossypii. *J Cell Sci, 114*(Pt 5), 975-986.

Amos, L. A. (1989). Brain dynein crossbridges microtubules into bundles. *J Cell Sci, 93 (Pt 1)*, 19-28.

Arimoto, M., Koushika, S. P., Choudhary, B. C., Li, C., Matsumoto, K., & Hisamoto, N. (2011). The Caenorhabditis elegans JIP3 protein UNC-16 functions as an adaptor to link kinesin-1 with cytoplasmic dynein. *J Neurosci, 31*(6), 2216-2224. doi:10.1523/jneurosci.2653-10.2011

Baumann, S., Pohlmann, T., Jungbluth, M., Brachmann, A., & Feldbrugge, M. (2012). Kinesin-3 and dynein mediate microtubule-dependent co-transport of mRNPs and endosomes. *J Cell Sci, 125*(Pt 11), 2740-2752. doi:10.1242/jcs.101212

Baumbach, J., Murthy, A., McClintock, M. A., Dix, C. I., Zalyte, R., Hoang, H. T., & Bullock, S. L. (2017). Lissencephaly-1 is a context-dependent regulator of the human dynein complex. *Elife, 6*. doi:10.7554/eLife.21768

Belyy, V., Schlager, M. A., Foster, H., Reimer, A. E., Carter, A. P., & Yildiz, A. (2016). The mammalian dynein-dynactin complex is a strong opponent to kinesin in a tug-of-war competition. *Nat Cell Biol, 18*(9), 1018-1024. doi:10.1038/ncb3393

Bhabha, G., Cheng, H. C., Zhang, N., Moeller, A., Liao, M., Speir, J. A., . . . Vale, R. D. (2014). Allosteric communication in the dynein motor domain. *Cell, 159*(4), 857-868. doi:10.1016/j.cell.2014.10.018

Bhabha, G., Johnson, G. T., Schroeder, C. M., & Vale, R. D. (2016). How Dynein Moves Along Microtubules. *Trends Biochem Sci, 41*(1), 94-105. doi:10.1016/j.tibs.2015.11.004

Bieger, B. D., Osmani, A. H., Xiang, X., & Egan, M. J. (2021). The spindle pole-body localization of activated cytoplasmic dynein is cell cycle-dependent in Aspergillus nidulans. *Fungal Genet Biol, 148*, 103519. doi:10.1016/j.fgb.2021.103519

Bielska, E., Schuster, M., Roger, Y., Berepiki, A., Soanes, D. M., Talbot, N. J., & Steinberg, G. (2014). Hook is an adapter that coordinates kinesin-3 and dynein cargo attachment on early endosomes. *J Cell Biol, 204*(6), 989-1007. doi:10.1083/jcb.201309022

Burgess, S. A., Walker, M. L., Sakakibara, H., Knight, P. J., & Oiwa, K. (2003). Dynein structure and power stroke. *Nature, 421*(6924), 715-718. doi:10.1038/nature01377

Can, S., Lacey, S., Gur, M., Carter, A. P., & Yildiz, A. (2019). Directionality of dynein is controlled by the angle and length of its stalk. *Nature, 566*(7744), 407-410. doi:10.1038/s41586-019-0914-z

Carter, A. P., Cho, C., Jin, L., & Vale, R. D. (2011). Crystal structure of the dynein motor domain. *Science, 331*(6021), 1159-1165. doi:10.1126/science.1202393

Carter, A. P., Diamant, A. G., & Urnavicius, L. (2016). How dynein and dynactin transport cargos: a structural perspective. *Curr Opin Struct Biol, 37*, 62-70. doi:10.1016/j.sbi.2015.12.003

Carvalho, P., Gupta, M. L., Jr., Hoyt, M. A., & Pellman, D. (2004). Cell cycle control of kinesin-mediated transport of Bik1 (CLIP-170) regulates microtubule stability and dynein activation. *Dev Cell, 6*(6), 815-829. doi:10.1016/j.devcel.2004.05.001

Celestino, R., Henen, M. A., Gama, J. B., Carvalho, C., McCabe, M., Barbosa, D. J., . . . Vögeli, B. (2019). A transient helix in the disordered region of dynein light intermediate chain links the motor to structurally diverse adaptors for cargo transport. *PLoS Biol, 17*(1), e3000100. doi:10.1371/journal.pbio.3000100

Cho, C., Reck-Peterson, S. L., & Vale, R. D. (2008). Regulatory ATPase sites of cytoplasmic dynein affect processivity and force generation. *J Biol Chem, 283*(38), 25839-25845. doi:10.1074/jbc.M802951200

Chowdhury, S., Ketcham, S. A., Schroer, T. A., & Lander, G. C. (2015). Structural organization of the dynein-dynactin complex bound to microtubules. *Nat Struct Mol Biol, 22*(4), 345-347. doi:10.1038/nsmb.2996

DeSantis, M. E., Cianfrocco, M. A., Htet, Z. M., Tran, P. T., Reck-Peterson, S. L., & Leschziner, A. E. (2017). Lis1 Has Two Opposing Modes of Regulating Cytoplasmic Dynein. *Cell, 170*(6), 1197-1208.e1112. doi:10.1016/j.cell.2017.08.037

DeWitt, M. A., Cypranowska, C. A., Cleary, F. B., Belyy, V., & Yildiz, A. (2015). The AAA3 domain of cytoplasmic dynein acts as a switch to facilitate microtubule release. *Nat Struct Mol Biol, 22*(1), 73-80. doi:10.1038/nsmb.2930

Efimov, V. P. (2003). Roles of NUDE and NUDF proteins of Aspergillus nidulans: insights from intracellular localization and overexpression effects. *Mol Biol Cell, 14*(3), 871-888. doi:10.1091/mbc.E02-06-0359

Efimov, V. P., & Morris, N. R. (2000). The LIS1-related NUDF protein of Aspergillus nidulans interacts with the coiled-coil domain of the NUDE/RO11 protein. *J Cell Biol, 150*(3), 681-688. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/10931877

Egan, M. J., Tan, K., & Reck-Peterson, S. L. (2012). Lis1 is an initiation factor for dynein-driven organelle transport. *J Cell Biol, 197*(7), 971-982. doi:10.1083/jcb.201112101

Elshenawy, M. M., Kusakci, E., Volz, S., Baumbach, J., Bullock, S. L., & Yildiz, A. (2020). Lis1 activates dynein motility by modulating its pairing with dynactin. *Nat Cell Biol*. doi:10.1038/s41556-020-0501-4

Etxebeste, O., Villarino, M., Markina-Inarrairaegui, A., Araujo-Bazan, L., & Espeso, E. A. (2013). Cytoplasmic dynamics of the general nuclear import machinery in apically growing syncytial cells. *PLoS One, 8*(12), e85076. doi:10.1371/journal.pone.0085076

Feng, Y., Olson, E. C., Stukenberg, P. T., Flanagan, L. A., Kirschner, M. W., & Walsh, C. A. (2000). LIS1 regulates CNS lamination by interacting with mNudE, a central component of the centrosome. *Neuron, 28*(3), 665-679.

Gee, M. A., Heuser, J. E., & Vallee, R. B. (1997). An extended microtubule-binding structure within the dynein motor domain. *Nature, 390*(6660), 636-639. doi:10.1038/37663

Gibbons, I. R., Gibbons, B. H., Mocz, G., & Asai, D. J. (1991). Multiple nucleotide-binding sites in the sequence of dynein beta heavy chain. *Nature, 352*(6336), 640-643. doi:10.1038/352640a0

Gonçalves, J. C., Quintremil, S., Yi, J., & Vallee, R. B. (2020). Nesprin-2 Recruitment of BicD2 to the Nuclear Envelope Controls Dynein/Kinesin-Mediated Neuronal Migration In Vivo. *Curr Biol, 30*(16), 3116-3129.e3114. doi:10.1016/j.cub.2020.05.091

Grotjahn, D. A., Chowdhury, S., Xu, Y., McKenney, R. J., Schroer, T. A., & Lander, G. C. (2018). Cryo-electron tomography reveals that dynactin recruits a team of dyneins for processive motility. *Nat Struct Mol Biol, 25*(3), 203-207. doi:10.1038/s41594-018-0027-7

Guo, X., Farias, G. G., Mattera, R., & Bonifacino, J. S. (2016). Rab5 and its effector FHF contribute to neuronal polarity through dynein-dependent retrieval of somatodendritic proteins from the axon. *Proc Natl Acad Sci U S A, 113*(36), E5318-5327. doi:10.1073/pnas.1601844113

Gutierrez, P. A., Ackermann, B. E., Vershinin, M., & McKenney, R. J. (2017). Differential effects of the dynein-regulatory factor Lissencephaly-1 on processive dynein-dynactin motility. *J Biol Chem, 292*(29), 12245-12255. doi:10.1074/jbc.M117.790048

Han, G., Liu, B., Zhang, J., Zuo, W., Morris, N. R., & Xiang, X. (2001). The Aspergillus cytoplasmic dynein heavy chain and NUDF localize to microtubule ends and affect microtubule dynamics. *Curr Biol, 11*(9), 719-724. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/11369237

Hernández-González, M., Bravo-Plaza, I., Pinar, M., de Los Ríos, V., Arst, H. N., Jr., & Peñalva, M. A. (2018). Endocytic recycling via the TGN underlies the polarized hyphal mode of life. *PLoS Genet, 14*(4), e1007291. doi:10.1371/journal.pgen.1007291

Higuchi, Y., Ashwin, P., Roger, Y., & Steinberg, G. (2014). Early endosome motility spatially organizes polysome distribution. *J Cell Biol, 204*(3), 343-357. doi:10.1083/jcb.201307164

Hoogenraad, C. C., & Akhmanova, A. (2016). Bicaudal D Family of Motor Adaptors: Linking Dynein Motility to Cargo Binding. *Trends Cell Biol, 26*(5), 327-340. doi:10.1016/j.tcb.2016.01.001

Htet, Z. M., Gillies, J. P., Baker, R. W., Leschziner, A. E., DeSantis, M. E., & Reck-Peterson, S. L. (2020). LIS1 promotes the formation of activated cytoplasmic dynein-1 complexes. *Nat Cell Biol*. doi:10.1038/s41556-020-0506-z

Huang, J., Roberts, A. J., Leschziner, A. E., & Reck-Peterson, S. L. (2012). Lis1 Acts as a "Clutch" between the ATPase and Microtubule-Binding Domains of the Dynein Motor. *Cell, 150*(5), 975-986. doi:10.1016/j.cell.2012.07.022

Jha, R., Roostalu, J., Cade, N. I., Trokter, M., & Surrey, T. (2017). Combinatorial regulation of the balance between dynein microtubule end accumulation and initiation of directed motility. *Embo j, 36*(22), 3387-3404. doi:10.15252/embj.201797077

Kardon, J. R., Reck-Peterson, S. L., & Vale, R. D. (2009). Regulation of the processivity and intracellular localization of Saccharomyces cerevisiae dynein by dynactin. *Proc Natl Acad Sci U S A, 106*(14), 5669-5674. doi:10.1073/pnas.0900976106

Karki, S., & Holzbaur, E. L. (1995). Affinity chromatography demonstrates a direct binding between cytoplasmic dynein and the dynactin complex. *J Biol Chem, 270*(48), 28806-28811. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=7499404

King, S. J., Brown, C. L., Maier, K. C., Quintyne, N. J., & Schroer, T. A. (2003). Analysis of the dynein-dynactin interaction in vitro and in vivo. *Mol Biol Cell, 14*(12), 5089-5097. doi:10.1091/mbc.E03-01-0025

King, S. J., & Schroer, T. A. (2000). Dynactin increases the processivity of the cytoplasmic dynein motor. *Nat Cell Biol, 2*(1), 20-24. doi:10.1038/71338

King, S. M. (2000). AAA domains and organization of the dynein motor unit. *J Cell Sci, 113 (Pt 14)*, 2521-2526.

Kon, T., Nishiura, M., Ohkura, R., Toyoshima, Y. Y., & Sutoh, K. (2004). Distinct functions of nucleotide-binding/hydrolysis sites in the four AAA modules of cytoplasmic dynein. *Biochemistry, 43*(35), 11266-11274. doi:10.1021/bi048985a

Kon, T., Oyama, T., Shimo-Kon, R., Imamula, K., Shima, T., Sutoh, K., & Kurisu, G. (2012). The 2.8 A crystal structure of the dynein motor domain. *Nature, 484*(7394), 345-350. doi:10.1038/nature10955

Kon, T., Sutoh, K., & Kurisu, G. (2011). X-ray structure of a functional full-length dynein motor domain. *Nat Struct Mol Biol, 18*(6), 638-642. doi:10.1038/nsmb.2074

Koonce, M. P., Grissom, P. M., & McIntosh, J. R. (1992). Dynein from Dictyostelium: primary structure comparisons between a cytoplasmic motor enzyme and flagellar dynein. *J Cell Biol, 119*(6), 1597-1604. doi:10.1083/jcb.119.6.1597

Lammers, L. G., & Markus, S. M. (2015). The dynein cortical anchor Num1 activates dynein motility by relieving Pac1/LIS1-mediated inhibition. *J Cell Biol, 211*(2), 309-322. doi:10.1083/jcb.201506119

Lee, W. L., Oberle, J. R., & Cooper, J. A. (2003). The role of the lissencephaly protein Pac1 during nuclear migration in budding yeast. *J Cell Biol, 160*(3), 355-364. doi:10.1083/jcb.200209022

Lenz, J. H., Schuchardt, I., Straube, A., & Steinberg, G. (2006). A dynein loading zone for retrograde endosome motility at microtubule plus-ends. *Embo j, 25*(11), 2275-2286. doi:10.1038/sj.emboj.7601119

Li, J., Lee, W. L., & Cooper, J. A. (2005). NudEL targets dynein to microtubule ends through LIS1. *Nat Cell Biol, 7*(7), 686-690. doi:10.1038/ncb1273

Liang, Y., Yu, W., Li, Y., Yang, Z., Yan, X., Huang, Q., & Zhu, X. (2004). Nudel functions in membrane traffic mainly through association with Lis1 and cytoplasmic dynein. *J Cell Biol, 164*(4), 557-566. doi:10.1083/jcb.200308058

Lin, J., Okada, K., Raytchev, M., Smith, M. C., & Nicastro, D. (2014). Structural mechanism of the dynein power stroke. *Nat Cell Biol, 16*(5), 479-485. doi:10.1038/ncb2939

Liu, B., Xiang, X., & Lee, Y. R. (2003). The requirement of the LC8 dynein light chain for nuclear migration and septum positioning is temperature dependent in Aspergillus nidulans. *Mol Microbiol, 47*(2), 291-301.

Liu, X., Rao, L., & Gennerich, A. (2020). The regulatory function of the AAA4 ATPase domain of cytoplasmic dynein. *Nat Commun, 11*(1), 5952. doi:10.1038/s41467-020-19477-3

Lloyd, T. E., Machamer, J., O'Hara, K., Kim, J. H., Collins, S. E., Wong, M. Y., . . . Kolodkin, A. L. (2012). The p150(Glued) CAP-Gly domain regulates initiation of retrograde transport at synaptic termini. *Neuron, 74*(2), 344-360. doi:10.1016/j.neuron.2012.02.026

Markus, S. M., Marzo, M. G., & McKenney, R. J. (2020). New insights into the mechanism of dynein motor regulation by lissencephaly-1. *Elife, 9*. doi:10.7554/eLife.59737

Markus, S. M., Plevock, K. M., St Germain, B. J., Punch, J. J., Meaden, C. W., & Lee, W. L. (2011). Quantitative analysis of Pac1/LIS1-mediated dynein targeting: Implications for regulation of dynein activity in budding yeast. *Cytoskeleton (Hoboken), 68*(3), 157-174. doi:10.1002/cm.20502

Markus, S. M., Punch, J. J., & Lee, W. L. (2009). Motor- and tail-dependent targeting of dynein to microtubule plus ends and the cell cortex. *Curr Biol, 19*(3), 196-205. doi:10.1016/j.cub.2008.12.047

Marzo, M. G., Griswold, J. M., & Markus, S. M. (2020). Pac1/LIS1 stabilizes an uninhibited conformation of dynein to coordinate its localization and activity. *Nat Cell Biol*. doi:10.1038/s41556-020-0492-1

McKenney, R. J. (2020). LIS1 cracks open dynein. *Nat Cell Biol*. doi:10.1038/s41556-020-0500-5

McKenney, R. J., Huynh, W., Tanenbaum, M. E., Bhabha, G., & Vale, R. D. (2014). Activation of cytoplasmic dynein motility by dynactin-cargo adapter complexes. *Science, 345*(6194), 337-341. doi:10.1126/science.1254198

McKenney, R. J., Vershinin, M., Kunwar, A., Vallee, R. B., & Gross, S. P. (2010). LIS1 and NudE induce a persistent dynein force-producing state. *Cell, 141*(2), 304-314. doi:10.1016/j.cell.2010.02.035

McKenney, R. J., Weil, S. J., Scherer, J., & Vallee, R. B. (2011). Mutually exclusive cytoplasmic dynein regulation by NudE-Lis1 and dynactin. *J Biol Chem, 286*(45), 39615-39622. doi:10.1074/jbc.M111.289017

Moore, J. K., Li, J., & Cooper, J. A. (2008). Dynactin function in mitotic spindle positioning. *Traffic, 9*(4), 510-527. doi:10.1111/j.1600-0854.2008.00710.x

Moughamian, A. J., & Holzbaur, E. L. (2012). Dynactin is required for transport initiation from the distal axon. *Neuron, 74*(2), 331-343. doi:10.1016/j.neuron.2012.02.025

Moughamian, A. J., Osborn, G. E., Lazarus, J. E., Maday, S., & Holzbaur, E. L. (2013). Ordered recruitment of dynactin to the microtubule plus-end is required for efficient initiation of retrograde axonal transport. *J Neurosci, 33*(32), 13190-13203. doi:10.1523/JNEUROSCI.0935-13.2013

Nicholas, M. P., Berger, F., Rao, L., Brenner, S., Cho, C., & Gennerich, A. (2015). Cytoplasmic dynein regulates its attachment to microtubules via nucleotide state-switched mechanosensing at multiple AAA domains. *Proc Natl Acad Sci U S A, 112*(20), 6371-6376. doi:10.1073/pnas.1417422112

Nicholas, M. P., Rao, L., & Gennerich, A. (2014). Covalent immobilization of microtubules on glass surfaces for molecular motor force measurements and other single-molecule assays. *Methods Mol Biol, 1136*, 137-169. doi:10.1007/978-1-4939-0329-0_9

Niekamp, S., Coudray, N., Zhang, N., Vale, R. D., & Bhabha, G. (2019). Coupling of ATPase activity, microtubule binding, and mechanics in the dynein motor domain. *Embo j, 38*(13), e101414. doi:10.15252/embj.2018101414

Niethammer, M., Smith, D. S., Ayala, R., Peng, J., Ko, J., Lee, M. S., . . . Tsai, L. H. (2000). NUDEL is a novel Cdk5 substrate that associates with LIS1 and cytoplasmic dynein. *Neuron, 28*(3), 697-711.

Oakley, B. R., Oakley, C. E., Yoon, Y., & Jung, M. K. (1990). Gamma-tubulin is a component of the spindle pole body that is essential for microtubule function in Aspergillus nidulans. *Cell, 61*(7), 1289-1301.

Olenick, M. A., Dominguez, R., & Holzbaur, E. L. F. (2019). Dynein activator Hook1 is required for trafficking of BDNF-signaling endosomes in neurons. *J Cell Biol, 218*(1), 220-233. doi:10.1083/jcb.201805016

Olenick, M. A., & Holzbaur, E. L. F. (2019). Dynein activators and adaptors at a glance. *J Cell Sci, 132*(6). doi:10.1242/jcs.227132

Olenick, M. A., Tokito, M., Boczkowska, M., Dominguez, R., & Holzbaur, E. L. (2016). Hook Adaptors Induce Unidirectional Processive Motility by Enhancing the Dynein-Dynactin Interaction. *J Biol Chem, 291*(35), 18239-18251. doi:10.1074/jbc.M116.738211

Otamendi, A., Perez-de-Nanclares-Arregi, E., Oiartzabal-Arano, E., Cortese, M. S., Espeso, E. A., & Etxebeste, O. (2019). Developmental regulators FlbE/D orchestrate the polarity site-to-nucleus dynamics of the fungal bZIP transcription factor FlbB. *Cell Mol Life Sci*. doi:10.1007/s00018-019-03121-5

Penalva, M. A., Zhang, J., Xiang, X., & Pantazopoulou, A. (2017). Transport of fungal RAB11 secretory vesicles involves myosin-5, dynein/dynactin/p25, and kinesin-1 and is independent of kinesin-3. *Mol Biol Cell, 28*(7), 947-961. doi:10.1091/mbc.E16-08-0566

Perlson, E., Maday, S., Fu, M. M., Moughamian, A. J., & Holzbaur, E. L. (2010). Retrograde axonal transport: pathways to cell death? *Trends Neurosci, 33*(7), 335-344. doi:10.1016/j.tins.2010.03.006

Pfister, K. K., Shah, P. R., Hummerich, H., Russ, A., Cotton, J., Annuar, A. A., . . . Fisher, E. M. (2006). Genetic analysis of the cytoplasmic dynein subunit families. *PLoS Genet, 2*(1), e1. doi:10.1371/journal.pgen.0020001

Plamann, M., Minke, P. F., Tinsley, J. H., & Bruno, K. S. (1994). Cytoplasmic dynein and actin-related protein Arp1 are required for normal nuclear distribution in filamentous fungi. *J Cell Biol, 127*(1), 139-149. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/7929559

Qiu, R., Zhang, J., & Xiang, X. (2013). Identification of a novel site in the tail of dynein heavy chain important for dynein function in vivo. *J Biol Chem, 288*(4), 2271-2280. doi:10.1074/jbc.M112.412403

Qiu, R., Zhang, J., & Xiang, X. (2018). p25 of the dynactin complex plays a dual role in cargo binding and dynactin regulation. *J Biol Chem, 293*(40), 15606-15619. doi:10.1074/jbc.RA118.004000

Qiu, R., Zhang, J., & Xiang, X. (2019). LIS1 regulates cargo-adapter-mediated activation of dynein by overcoming its autoinhibition in vivo. *J Cell Biol, 218*(11), 3630-3646. doi:10.1083/jcb.201905178

Rao, L., Berger, F., Nicholas, M. P., & Gennerich, A. (2019). Molecular mechanism of cytoplasmic dynein tension sensing. *Nat Commun, 10*(1), 3332. doi:10.1038/s41467-019-11231-8

Reck-Peterson, S. L., Redwine, W. B., Vale, R. D., & Carter, A. P. (2018). The cytoplasmic dynein transport machinery and its many cargoes. *Nat Rev Mol Cell Biol, 19*(6), 382-398. doi:10.1038/s41580-018-0004-3

Reiner, O., Carrozzo, R., Shen, Y., Wehnert, M., Faustinella, F., Dobyns, W. B., . . . Ledbetter, D. H. (1993). Isolation of a Miller-Dieker lissencephaly gene containing G protein beta-subunit-like repeats. *Nature, 364*(6439), 717-721. doi:10.1038/364717a0

Roberts, A. J., Goodman, B. S., & Reck-Peterson, S. L. (2014). Reconstitution of dynein transport to the microtubule plus end by kinesin. *Elife, 3*, e02641. doi:10.7554/eLife.02641

Roberts, A. J., Kon, T., Knight, P. J., Sutoh, K., & Burgess, S. A. (2013). Functions and mechanics of dynein motor proteins. *Nat Rev Mol Cell Biol, 14*(11), 713-726. doi:10.1038/nrm3667

Roberts, A. J., Numata, N., Walker, M. L., Kato, Y. S., Malkova, B., Kon, T., . . . Burgess, S. A. (2009). AAA+ Ring and linker swing mechanism in the dynein motor. *Cell, 136*(3), 485-495. doi:10.1016/j.cell.2008.11.049

Ross, J. L., Wallace, K., Shuman, H., Goldman, Y. E., & Holzbaur, E. L. (2006). Processive bidirectional motion of dynein-dynactin complexes in vitro. *Nat Cell Biol, 8*(6), 562-570. doi:10.1038/ncb1421

Salogiannis, J., Christensen, J. R., Songster, L. D., Aguilar-Maldonado, A., Shukla, N., & Reck-Peterson, S. L. (2021). PxdA interacts with the DipA phosphatase to regulate peroxisome hitchhiking on early endosomes. *Mol Biol Cell, 32*(6), 492-503. doi:10.1091/mbc.E20-08-0559

Salogiannis, J., & Reck-Peterson, S. L. (2017). Hitchhiking: A Non-Canonical Mode of Microtubule-Based Transport. *Trends Cell Biol, 27*(2), 141-150. doi:10.1016/j.tcb.2016.09.005

Sasaki, S., Shionoya, A., Ishida, M., Gambello, M. J., Yingling, J., Wynshaw-Boris, A., & Hirotsune, S. (2000). A LIS1/NUDEL/cytoplasmic dynein heavy chain complex in the developing and adult nervous system. *Neuron, 28*(3), 681-696.

Scherer, J., Yi, J., & Vallee, R. B. (2014). PKA-dependent dynein switching from lysosomes to adenovirus: a novel form of host-virus competition. *J Cell Biol, 205*(2), 163-177. doi:10.1083/jcb.201307116

Scherer, J., Yi, J., & Vallee, R. B. (2020). Role of cytoplasmic dynein and kinesins in adenovirus transport. *FEBS Lett, 594*(12), 1838-1847. doi:10.1002/1873-3468.13777

Schiavo, G., Greensmith, L., Hafezparast, M., & Fisher, E. M. (2013). Cytoplasmic dynein heavy chain: the servant of many masters. *Trends Neurosci, 36*(11), 641-651. doi:10.1016/j.tins.2013.08.001

Schlager, M. A., Hoang, H. T., Urnavicius, L., Bullock, S. L., & Carter, A. P. (2014). In vitro reconstitution of a highly processive recombinant human dynein complex. *Embo j, 33*(17), 1855-1868. doi:10.15252/embj.201488792

Schmidt, H., & Carter, A. P. (2016). Review: Structure and mechanism of the dynein motor ATPase. *Biopolymers, 105*(8), 557-567. doi:10.1002/bip.22856

Schmidt, H., Gleave, E. S., & Carter, A. P. (2012). Insights into dynein motor domain function from a 3.3-A crystal structure. *Nat Struct Mol Biol, 19*(5), 492-497, s491. doi:10.1038/nsmb.2272

Schmidt, H., Zalyte, R., Urnavicius, L., & Carter, A. P. (2015). Structure of human cytoplasmic dynein-2 primed for its power stroke. *Nature, 518*(7539), 435-438. doi:10.1038/nature14023

Schroeder, C. M., & Vale, R. D. (2016). Assembly and activation of dynein-dynactin by the cargo adaptor protein Hook3. *J Cell Biol, 214*(3), 309-318. doi:10.1083/jcb.201604002

Schroer, T. A. (2004). Dynactin. *Annu Rev Cell Dev Biol, 20*, 759-779. doi:10.1146/annurev.cellbio.20.012103.094623

Sheeman, B., Carvalho, P., Sagot, I., Geiser, J., Kho, D., Hoyt, M. A., & Pellman, D. (2003). Determinants of S. cerevisiae dynein localization and activation: implications for the mechanism of spindle positioning. *Curr Biol, 13*(5), 364-372.

Shu, T., Ayala, R., Nguyen, M. D., Xie, Z., Gleeson, J. G., & Tsai, L. H. (2004). Ndel1 operates in a common pathway with LIS1 and cytoplasmic dynein to regulate cortical neuronal positioning. *Neuron, 44*(2), 263-277. doi:10.1016/j.neuron.2004.09.030

Silvanovich, A., Li, M. G., Serr, M., Mische, S., & Hays, T. S. (2003). The third P-loop domain in cytoplasmic dynein heavy chain is essential for dynein motor function and ATP-sensitive microtubule binding. *Mol Biol Cell, 14*(4), 1355-1365. doi:10.1091/mbc.e02-10-0675

Splinter, D., Razafsky, D. S., Schlager, M. A., Serra-Marques, A., Grigoriev, I., Demmers, J., . . . Akhmanova, A. (2012). BICD2, dynactin, and LIS1 cooperate in regulating dynein recruitment to cellular structures. *Mol Biol Cell, 23*(21), 4226-4241. doi:10.1091/mbc.E12-03-0210

Stavoe, A. K. H., & Holzbaur, E. L. F. (2019). Axonal autophagy: Mini-review for autophagy in the CNS. *Neurosci Lett, 697*, 17-23. doi:10.1016/j.neulet.2018.03.025

Stehman, S. A., Chen, Y., McKenney, R. J., & Vallee, R. B. (2007). NudE and NudEL are required for mitotic progression and are involved in dynein recruitment to kinetochores. *J Cell Biol, 178*(4), 583-594. doi:10.1083/jcb.200610112

Steinberg, G., Peñalva, M. A., Riquelme, M., Wösten, H. A., & Harris, S. D. (2017). Cell Biology of Hyphal Growth. *Microbiol Spectr, 5*(2). doi:10.1128/microbiolspec.FUNK-0034-2016

Torisawa, T., Ichikawa, M., Furuta, A., Saito, K., Oiwa, K., Kojima, H., . . . Furuta, K. (2014). Autoinhibition and cooperative activation mechanisms of cytoplasmic dynein. *Nat Cell Biol, 16*(11), 1118-1124. doi:10.1038/ncb3048

Toropova, K., Zou, S., Roberts, A. J., Redwine, W. B., Goodman, B. S., Reck-Peterson, S. L., & Leschziner, A. E. (2014). Lis1 regulates dynein by sterically blocking its mechanochemical cycle. *Elife, 3*. doi:10.7554/eLife.03372

Tripathy, S. K., Weil, S. J., Chen, C., Anand, P., Vallee, R. B., & Gross, S. P. (2014). Autoregulatory mechanism for dynactin control of processive and diffusive dynein transport. *Nat Cell Biol, 16*(12), 1192-1201. doi:10.1038/ncb3063

Tucker, P. A., & Sallai, L. (2007). The AAA+ superfamily--a myriad of motions. *Curr Opin Struct Biol, 17*(6), 641-652. doi:10.1016/j.sbi.2007.09.012

Twelvetrees, A. E., Pernigo, S., Sanger, A., Guedes-Dias, P., Schiavo, G., Steiner, R. A., . . . Holzbaur, E. L. (2016). The Dynamic Localization of Cytoplasmic Dynein in Neurons Is Driven by Kinesin-1. *Neuron, 90*(5), 1000-1015. doi:10.1016/j.neuron.2016.04.046

Urnavicius, L., Lau, C. K., Elshenawy, M. M., Morales-Rios, E., Motz, C., Yildiz, A., & Carter, A. P. (2018). Cryo-EM shows how dynactin recruits two dyneins for faster movement. *Nature, 554*(7691), 202-206. doi:10.1038/nature25462

Urnavicius, L., Zhang, K., Diamant, A. G., Motz, C., Schlager, M. A., Yu, M., . . . Carter, A. P. (2015). The structure of the dynactin complex and its interaction with dynein. *Science, 347*(6229), 1441-1446. doi:10.1126/science.aaa4080

Vaughan, K. T., & Vallee, R. B. (1995). Cytoplasmic dynein binds dynactin through a direct interaction between the intermediate chains and p150Glued. *J Cell Biol, 131*(6 Pt 1), 1507-1516. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=8522607

Vaughan, P. S., Miura, P., Henderson, M., Byrne, B., & Vaughan, K. T. (2002). A role for regulated binding of p150(Glued) to microtubule plus ends in organelle transport. *J Cell Biol, 158*(2), 305-319. doi:10.1083/jcb.200201029

Wang, S., Ketcham, S. A., Schon, A., Goodman, B., Wang, Y., Yates, J., 3rd, . . . Zheng, Y. (2013). Nudel/NudE and Lis1 promote dynein and dynactin interaction in the context of spindle morphogenesis. *Mol Biol Cell, 24*(22), 3522-3533. doi:10.1091/mbc.E13-05-0283

Wang, S., & Zheng, Y. (2011). Identification of a novel dynein binding domain in nudel essential for spindle pole organization in Xenopus egg extract. *J Biol Chem, 286*(1), 587-593. doi:10.1074/jbc.M110.181578

Waring, R. B., May, G. S., & Morris, N. R. (1989). Characterization of an inducible expression system in Aspergillus nidulans using alcA and tubulin-coding genes. *Gene, 79*(1), 119-130. doi:10.1016/0378-1119(89)90097-8

Wedlich-Soldner, R., Straube, A., Friedrich, M. W., & Steinberg, G. (2002). A balance of KIF1A-like kinesin and dynein organizes early endosomes in the fungus Ustilago maydis. *Embo j, 21*(12), 2946-2957. doi:10.1093/emboj/cdf296

Wendler, P., Ciniawsky, S., Kock, M., & Kube, S. (2012). Structure and function of the AAA+ nucleotide binding pocket. *Biochim Biophys Acta, 1823*(1), 2-14. doi:10.1016/j.bbamcr.2011.06.014

Willins, D. A., Liu, B., Xiang, X., & Morris, N. R. (1997). Mutations in the heavy chain of cytoplasmic dynein suppress the nudF nuclear migration mutation of Aspergillus nidulans. *Mol Gen Genet, 255*(2), 194-200. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/9236777

Winey, M., & Bloom, K. (2012). Mitotic spindle form and function. *Genetics, 190*(4), 1197-1224. doi:10.1534/genetics.111.128710

Wynshaw-Boris, A. (2007). Lissencephaly and LIS1: insights into the molecular mechanisms of neuronal migration and development. *Clin Genet, 72*(4), 296-304. doi:10.1111/j.1399-0004.2007.00888.x

Xiang, X. (2018). Nuclear movement in fungi. *Semin Cell Dev Biol, 82*, 3-16. doi:10.1016/j.semcdb.2017.10.024

Xiang, X., Beckwith, S. M., & Morris, N. R. (1994). Cytoplasmic dynein is involved in nuclear migration in Aspergillus nidulans. *Proc Natl Acad Sci U S A, 91*(6), 2100-2104. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/8134356

Xiang, X., Han, G., Winkelmann, D. A., Zuo, W., & Morris, N. R. (2000). Dynamics of cytoplasmic dynein in living cells and the effect of a mutation in the dynactin complex actin-related protein Arp1. *Curr Biol, 10*(10), 603-606. doi:10.1016/s0960-9822(00)00488-7

Xiang, X., Osmani, A. H., Osmani, S. A., Xin, M., & Morris, N. R. (1995a). NudF, a nuclear migration gene in Aspergillus nidulans, is similar to the human LIS-1 gene required for

neuronal migration. *Mol Biol Cell, 6*(3), 297-310. Retrieved from
http://www.ncbi.nlm.nih.gov/pubmed/7612965

Xiang, X., & Qiu, R. (2020). Cargo-Mediated Activation of Cytoplasmic Dynein in vivo. *Front Cell Dev Biol, 8*, 598952. doi:10.3389/fcell.2020.598952

Xiang, X., Roghi, C., & Morris, N. R. (1995b). Characterization and localization of the cytoplasmic dynein heavy chain in Aspergillus nidulans. *Proc Natl Acad Sci U S A, 92*(21), 9890-9894.

Xu, L., Sowa, M. E., Chen, J., Li, X., Gygi, S. P., & Harper, J. W. (2008). An FTS/Hook/p107(FHIP) complex interacts with and promotes endosomal clustering by the homotypic vacuolar protein sorting complex. *Mol Biol Cell, 19*(12), 5059-5071. doi:10.1091/mbc.E08-05-0473

Yan, X., Li, F., Liang, Y., Shen, Y., Zhao, X., Huang, Q., & Zhu, X. (2003). Human Nudel and NudE as regulators of cytoplasmic dynein in poleward protein transport along the mitotic spindle. *Mol Cell Biol, 23*(4), 1239-1250.

Yao, X., Wang, X., & Xiang, X. (2014). FHIP and FTS proteins are critical for dynein-mediated transport of early endosomes in Aspergillus. *Mol Biol Cell, 25*(14), 2181-2189. doi:10.1091/mbc.E14-04-0873

Yao, X., Zhang, J., Zhou, H., Wang, E., & Xiang, X. (2012). In vivo roles of the basic domain of dynactin p150 in microtubule plus-end tracking and dynein function. *Traffic, 13*(3), 375-387. doi:10.1111/j.1600-0854.2011.01312.x

Yeh, T. Y., Quintyne, N. J., Scipioni, B. R., Eckley, D. M., & Schroer, T. A. (2012). Dynactin's pointed-end complex is a cargo-targeting module. *Mol Biol Cell, 23*(19), 3827-3837. doi:10.1091/mbc.E12-07-0496

Zekert, N., & Fischer, R. (2009). The Aspergillus nidulans kinesin-3 UncA motor moves vesicles along a subpopulation of microtubules. *Mol Biol Cell, 20*(2), 673-684. doi:10.1091/mbc.E08-07-0685

Zhang, J., Li, S., Fischer, R., & Xiang, X. (2003). Accumulation of cytoplasmic dynein and dynactin at microtubule plus ends in Aspergillus nidulans is kinesin dependent. *Mol Biol Cell, 14*(4), 1479-1488. doi:10.1091/mbc.E02-08-0516

Zhang, J., Qiu, R., Arst, H. N., Jr., Penalva, M. A., & Xiang, X. (2014). HookA is a novel dynein-early endosome linker critical for cargo movement in vivo. *J Cell Biol, 204*(6), 1009-1026. doi:10.1083/jcb.201308009

Zhang, J., Wang, L., Zhuang, L., Huo, L., Musa, S., Li, S., & Xiang, X. (2008). Arp11 affects dynein-dynactin interaction and is essential for dynein function in Aspergillus nidulans. *Traffic, 9*(7), 1073-1087. doi:10.1111/j.1600-0854.2008.00748.x

Zhang, J., Yao, X., Fischer, L., Abenza, J. F., Penalva, M. A., & Xiang, X. (2011). The p25 subunit of the dynactin complex is required for dynein-early endosome interaction. *J Cell Biol, 193*(7), 1245-1255. doi:10.1083/jcb.201011022

Zhang, J., Zhuang, L., Lee, Y., Abenza, J. F., Penalva, M. A., & Xiang, X. (2010). The microtubule plus-end localization of Aspergillus dynein is important for dynein-early-endosome interaction but not for dynein ATPase activation. *J Cell Sci, 123*(Pt 20), 3596-3604. doi:10.1242/jcs.075259

Zhang, K., Foster, H. E., Rondelet, A., Lacey, S. E., Bahi-Buisson, N., Bird, A. W., & Carter, A. P. (2017a). Cryo-EM Reveals How Human Cytoplasmic Dynein Is Auto-inhibited and Activated. *Cell, 169*(7), 1303-1314.e1318. doi:10.1016/j.cell.2017.05.025

Zhang, Y., Gao, X., Manck, R., Schmid, M., Osmani, A. H., Osmani, S. A., . . . Fischer, R. (2017b). Microtubule-organizing centers of Aspergillus nidulans are anchored at septa by a disordered protein. *Mol Microbiol, 106*(2), 285-303. doi:10.1111/mmi.13763

Zhuang, L., Zhang, J., & Xiang, X. (2007). Point mutations in the stem region and the fourth AAA domain of cytoplasmic dynein heavy chain partially suppress the phenotype of NUDF/LIS1 loss in Aspergillus nidulans. *Genetics, 175*(3), 1185-1196. doi:10.1534/genetics.106.069013

Zylkiewicz, E., Kijanska, M., Choi, W. C., Derewenda, U., Derewenda, Z. S., & Stukenberg, P. T. (2011). The N-terminal coiled-coil of Ndel1 is a regulated scaffold that recruits LIS1 to dynein. *J Cell Biol, 192*(3), 433-445. doi:10.1083/jcb.201011142