Main Manuscript for

Sugar Signaling Induces Dynamic Changes During Meristem Development to Regulate Flowering in Arabidopsis

Magdalena Musialak-Lange[1], Katharina Fiddeke[1,#], Annika Franke[1,#], Friedrich Kragler[1], Christin Abel[1], and Vanessa Wahl[1,*]

Affiliations: [1]Max Planck Institute of Molecular Plant Physiology, Department of Metabolic Networks, Am Mühlenberg 1, Potsdam, 14476, Germany.

[#]K.F. and A.F. contributed equally

[*]**Correspondence**: Vanessa Wahl

Email: vanessa.wahl@mpimp-golm.mpg.de

Abstract

Aerial parts of plants originate from pluripotent stem cells in the shoot apical meristem. Their population is maintained via the maintenance regulators *WUSCHEL* and *CLAVATA3* in a negative feed-back loop. Meristem size is dynamic and undergoes a more than 2-fold expansion upon the transition to reproductive growth. The mechanism controlling this doming is largely unknown, but coinciding increased trehalose 6-phosphate and changed meristem size in overexpressing or knockdown lines of *TREHALOSE PHOSPHATE SYNTHASE1* suggest a participation of sugar signaling. Here we show that *TREHALOSE PHOSPHATE PHOSPHATASEJ* is directly regulated by WUSCHEL. Plants with reduced levels of *TREHALOSE PHOSPHATE PHOSPHATASEJ* in the outer meristem layer are flowering early and its reduction in the late flowering *clavata3* mutant, restores wild-type flowering. This is caused by a reduction of mature miR156 and increased expression of *SQUAMOSA PROMOTER-BINDING PROTEIN-LIKE* genes. We demonstrate that these are important for age pathway-induced flowering, in a negative feed-back loop with WUSCHEL downstream of the trehalose 6-phosphate pathway. In summary, our findings demonstrate a dynamic feed-back regulation between central maintenance and flowering time regulators with sugar signaling.

Keywords: flowering / meristem maintenance / sugar / Trehalose 6-phosphate (T6P) pathway / WUS

1 Introduction

All above ground plant organs such as leaves and flowers originate from the shoot apical
meristem (SAM). The *Arabidopsis thaliana* (Arabidopsis) SAM is organized in three clonally
distinct cell layers: the outer mono cell layers, L1 and L2, and the several cell files comprising L3.
Functionally, the SAM consist of a central zone (CZ), containing undifferentiated cells with a self-
renewing potential, an organizing centre (OC), inducing and maintaining an adequate number of
cells in the CZ, a peripheral zone (PZ) which receives cells from the CZ that are differentiating
and incorporated into leaf or flower primordia, and a rib zone (RZ) producing the stem (Pfeiffer *et*
al, 2017). Communicating position-dependent properties among cells within and between the
different zones is crucial for SAM function. Two non-cell autonomously acting factors regulate
each other in a negative feed-back loop. The homeodomain transcription factor *WUSCHEL*
(*WUS*) is expressed in the OC, and the signalling peptide *CLAVATA3* (*CLV3*) in the CZ. Thus,
mobile WUS transcription factor, secreted from the OC, activates expression of *CLV3* in the CZ
above, while CLV3 peptide restricts the *WUS* expression domain to the OC. Hence, *WUS* and
CLV3 expression domains are considered mutually exclusive and their negative feed-back
regulation as very robust, which can only be modulated within tight limits (Pfeiffer *et al.*, 2017).

Development as a whole and SAM maintenance in particular demands continuous cross talk
between its regulatory processes and the available resources. SAM maintenance and flowering
time are important determinants of crop yield. During floral transition cell division is activated and
the SAM undergoes a dramatic increase in size known as doming (Jacqmard *et al*, 2003;
Kwiatkowska, 2008) (Figure 1A) and marks the time between the vegetative stage (production of
leaves) and the reproductive stage (formation of flowers). Since this transition requires a massive
reorganization of organ development and sufficient energy, it is tightly controlled by
environmental conditions and availability of nutrients (Andres & Coupland, 2012; Olas *et al*, 2019;
Ponnu *et al*, 2020; Srikanth & Schmid, 2011; Wahl *et al*, 2013). Only recently the early cellular
changes were associated with floral regulators that control GA biosynthesis (Kinoshita *et al*,
2020). However, the knowledge on the mechanisms controlling doming is still scarce.

In plants, the sugar phosphate, trehalose 6-phosphate (T6P), it is synthetized from UDP-glucose
and glucose 6-phosphate via TREHALOSE PHOSPHATE SYNTHASE (TPS) and is converted to
trehalose by TREHALOSE PHOSPHATE PHOSPHATESES (TPP) (Fichtner & Lunn, 2021). T6P
serves as a signal for sucrose availability, which is conveyed to downstream metabolic and
growth responses through still largely unknown mechanisms (Fichtner & Lunn, 2021). In
Arabidopsis sugar signalling affects the vegetative phase change, the transition between the
formation of juvenile and adult leaves, via the T6P pathway (Ponnu *et al.*, 2020; Yang *et al*, 2013;
Yu *et al*, 2013). In addition, the T6P pathway is involved in shoot branching (Fichtner *et al*, 2020),
and induces flowering via regulating key flowering genes in leaves and at the SAM (Wahl *et al.*,

2013). In leaves, *TPS1*, coding for the T6P synthesizing enzyme, is necessary and sufficient to
induce *FLOWERING LOCUS T* (*FT*), the florigen, connecting the photoperiod signal with a
physiological signal. In addition, the T6P pathway affects the age-dependent microRNA156
(miR156)/*SQUAMOSA PROMOTER-BINDING PROTEIN-LIKE* (*SPL*) module at the SAM (Wahl
et al., 2013). Increased T6P levels coincide with doming (Wahl *et al.*, 2013), and a spatial change
of the *WUS* expression domain. This suggests a participation of sugar signaling in regulating
SAM dynamics through transient uncoupling of the negative feed-back regulation between WUS
and CLV3. Here we show that WUS directly regulates *TPPJ*. A late flowering phenotype of *clv3* is
caused by increased levels of mature miR156 and decreased expression of *SPL* genes and is
restored when *TPPJ* is downregulated in the outer meristem layer. We further demonstrate a
negative feed-back loop with WUS and SPL4 downstream of the T6P pathway. In summary, we
provide evidence for a dynamic feed-back regulation between central meristem maintenance and
flowering time regulators with sugar signaling.

Results and Discussion

The T6P pathway affects SAM size. To assess whether the morphological changes at the
transition SAM (Figure 1A) involve the T6P pathway, we investigated the effect on SAM
architecture by decreasing *TPS1* by the means of an artificial microRNA in the SAM proper
(*35S:amiRTPS1*; Figure S1) and increasing *TPS1* in the CZ (*CLV3:TPS1*, Figure S2). These
plants have smaller and bigger vegetative and reproductive meristems (Figure 1B,C), resulting in
smaller and bigger plants, respectively (Wahl *et al.*, 2013).

TPS1 knock-out mutants are embryo lethal, likely as a result of insufficient cell cycle activity at
torpedo stage of embryogenesis (Gómez *et al*, 2006). Cell cycle genes were also found to be
highly induced at transition to flowering (Klepikova *et al*, 2015) confirming earlier findings
(Jacqmard *et al.*, 2003). In order to test, whether cell proliferation is the cause of the effect
observed downstream of the T6P pathway, we used specific probes against transcripts of
HISTONE4, CYCLIN D3;1 and *CYCLIN-DEPENDENT KINASE2;1*, three cell cycle marker genes,
for RNA *in situ* hybridization on longitudinal sections through the SAM (Figure S3). The results
show more abundant staining indicating the three cell cycle marker genes in the presence of
elevated T6P levels and less staining, when T6P levels were reduced. This suggests a higher
proliferation rate and an increased number of cells produced to adopt organ-specific cell fates at
the periphery as a plausible reason for decreased SAM size of *TPS1* overexpression and *vice*
versa increased SAM size of *TPS1* knock-down lines (Figure 1B,C).

Spatial changes of *WUS* expression in a transition SAM. In line with these findings, the floral
transition coincides with increased T6P levels (Wahl *et al.*, 2013). To assess whether this is
related to *WUS* or *CLV3* expression in a wild-type SAM, we analyzed their expression in a time
series spanning floral transition (Figure 1D). Throughout the vegetative phase SAM size gradually
increases due to rising cell numbers (Figures 1A, S4) (Kinoshita *et al.*, 2020), reaching its
maximum at floral transition, independent of whether the plants are grown in long days (LD)
(Figure 1A) or in short days (SD) followed by a transfer to LD (Figure S5). An enlarging SAM at
floral transition correlates with a larger *WUS* expression domain expanding into the CZ and the
outer cell layer (L1) (Figure 1D-F).

Cytokinin signaling was previously reported to respond to carbon in seedlings (Leibfried *et al*,
2005; Pfeiffer *et al*, 2016; Snipes *et al*, 2018). Its transient occurrence in L1 may thus explain the
presence of *WUS* transcript in the outer SAM layers. However, cytokinin levels are not altered in
L1 cells as indicated by the synthetic cytokinin reporter *TCSn:GFP* (Zurcher *et al*, 2013) (Figure
S6). Further, while *WUS* expands to L1 for a short period (8-10 DAG), the *CLV3* expression
domain remains expanded post floral transition (Figure 1D,G). This suggests a transient
uncoupling of the negative feed-back regulation between WUS and CLV3, which is re-established
at the reproductive SAM (16 DAG), resembling earlier vegetative SAM expression patterns (6
DAG) (Figure 1D).

TPPJ is directly regulated by WUS. To understand how the T6P pathway might control this
process, we first analyzed expression of the ten genes encoding TREHALOSE PHOSPHATE
PHOSPHATASEs (TPP) (Fichtner & Lunn, 2021). Except for *TPPC* and *TPPD*, which are
expressed below detection limit of RNA *in situ* hybridization, we found that all are expressed in
distinct SAM domains (Figure S7). *TPPJ* strongly increases in the enlarged SAM of *clv3-7* due to
an ectopic expression in L1 and L2 (Figures 2A-C; S8, S9). Notably, in *clv3-7* and *clv3-10*
mutants high levels of *WUS* in the outer SAM layers (Brand *et al*, 2000) coincide with ectopic
expression of *TPPJ* (Figure S9), while *TPPJ* is repressed in a *wus-7* background (Figure S10A).
In addition, a previously published data set shows induced levels of *TPPJ* upon DEX induced
expression of *WUS* in a *35S:WUS-GR* line (Busch *et al*, 2010) (Figure S10 B). To assess whether
ectopic expression of *TPPJ* contributes to the enlarged SAM of the *clv3* mutant, we
overexpressed *TPPJ* in the wild type (*35S:TPPJ*; Figure 2D,E), which resulted in plants with
significantly enlarged SAMs (Figure 2F).

These results imply a direct influence of WUS on *TPPJ*, and is supported by an *in silico* analysis
which predicts multiple, canonical WUS binding sites in the sequence upstream of *TPPJ*
(*TPPJ*[WUS]; Figure 3A) (Leibfried *et al.*, 2005; Lohmann *et al*, 2001; Sloan *et al*, 2020; Yadav *et al*,
2011). To understand if WUS directly controls *TPPJ*, we performed *in vivo* transactivation assays.

These show that WUS activates reporter gene expression when using a 2865bp 5'*TPPJ*
sequence, containing 17 putative *TPPJ*^WUS sites (Figure 3A). Progressive deletion of this
sequence results in a reduction of reporter gene activation, suggesting an additive effect of the
individual *TPPJ*^WUS sites (#1-6; Figure 3A,B). We confirmed direct binding of WUS to three
distinct regions of the *TPPJ* promoter (I, II, and III; Figures 3C-E; S11) by chromatin
immunoprecipitation (ChIP) coupled to PCR using a specific antibody against WUS (Figures S11,
S12), while all other regions did not indicate binding (Figures S11, S13). We observed enrichment
of WUS binding to *TPPJ*^WUS sites up to 0.67% of the input DNA in *clv3-7* and 0.75% of the input
DNA in *clv3-10* apices, both of which express *WUS* at high levels in comparison to input DNA
from wild-type apices, where *WUS* is expressed in only a few cells and input DNA from leaves,
with no WUS (Figures 3C-E; S10). Electrophoretic mobility shift assays (EMSA) confirm specific
in vitro binding of WUS to the investigated sequences (I, II 1-3, III 1-3; Figure 3F; S14). In
summary, we demonstrate that seven out of 17 putative *TPPJ*^WUS sites are directly targeted by
WUS *in vivo*.

TPPJ controls flowering from the outer SAM layer. A *tppj* mutant with a T-DNA insertion in the
6^th exon, which specifically knocks-out *TPPJ* expression (Figure S15), flowers significantly earlier
than the wild type (Table S1). To further assess the role of *TPPJ* at the SAM we used an artificial
miRNA (amiR) approach to downregulate *TPPJ* (*35S:amiRTPPJ*; Figure S16)(Schwab *et al*,
2006). Plants overexpressing either of two versions of an *amiRTPPJ* (V1, V2) flower significantly
earlier in LD and SD (Figure S17; Table S1). *ML1* expression specifically and stably localizes to
L1 in all investigated stages (Figure S18). *ML1:amiRTPPJ* reduces *TPPJ* expression in L1
(Figure S19), has a smaller SAM and flowers significantly earlier in LD and SD (Figures 4A,B;
S20, S21; Table S1), while the level of *TPPJ* reduction is proportional to the acceleration of
flowering (Figures 4B,C; S16; Table S1).

To date there are no reports that mutants in meristem maintenance genes have a flowering
phenotype. We found that *clv3* plants are late flowering (Figure 4C; Table S1). However, when
ML1:amiRTPPJ is introgressed into the *clv3-7* or *clv3-10* background, the late flowering
phenotype is restored to wild-type (Figures 4C; S21; Table S1), suggesting that *TPPJ* expression
in the outer meristem layers is causal for the late flowering phenotype of *clv3*. Hence, the early
flowering of *ML1:amiRTPPJ* in a wild-type background is due to a reduction of *TPPJ* expression
in the outer meristem layer in its endogenous expression domain. In addition, other prominent
morphological defects of *clv3-10* such as the fasciated stem and enormous inflorescence SAM
are visibly reduced in the presence of *ML1:amiRTPPJ* (Figure S21).

WUS regulates the age pathway through *SPL4*. We previously reported that the T6P pathway
influences the age pathway at the SAM (Wahl *et al.*, 2013). We therefore next analyzed mature

miR156, as well as the expression of *SPL3*, *SPL4*, *SPL5*, *SPL9* and *SPL15*, all associated with
floral transition in one or the other way (Hyun *et al*, 2017; Xie *et al*, 2020). We found a strong
reduction of mature miR156 levels correlating with decreased *TPPJ* in *ML1:amiRTPPJ*, and
corresponding increased expression of *SPL3, SPL4, SPL5*, and *SPL9* (Figure 4D). In line,
miR156 was more abundant in *35S:TPPJ*, while the corresponding *SPL*s were downregulated
(Figure 4D). Expression of *SPL15*, previously described as an important integrator of plant age
into the regulation of floral onset in SD (Hyun *et al*, 2016; Wang *et al*, 2009; Yamaguchi *et al*,
2014), was not differentially expressed in the transgenic lines (Figure S22A).

Uncoupling of the WUS/CLV3 feed-back loop occurs downstream of the T6P pathway and
involves age pathway components. Interestingly, a study by Fouracre and Poethig indicates an
active pool of miR156 in the SAM, which is essential for early shoot maturation and affects leaf
identity. In addition, apices of a *wus-5* mutant had decreased miR156 levels and increased *SPL9*
expression, although expression of none of the other SPLs was tested. However, plants with
increased miR156 (*35S:miR156a*) have a larger SAM, while those with reduced miR156
(quadruple *mir156a,c,mir157a,c*) have a smaller SAM, and a sextuple *spl2,9,10,11,13,15*
increased SAM size. This is accompanied with an increased *WUS* expression in the sextuple
mutant, suggesting the potential repression of *WUS* by any or all of the *SPL*s mutated in hextuple
plants (Fouracre & Poethig, 2019). Notably, we found miR156 levels significantly increased in
clv3-10 apices (Figure 5A). In response, *SPL3*, *SPL5*, and *SPL9* were reduced supporting the
late flowering phenotype of the mutant, while *SPL15* was not affected (Figure S22B). In contrast
to what would be expected, we found more *SPL4* transcript in apices of the late flowering *clv3-10*
(Figure 5A).

The contribution of SPL9 and SPL15 to the regulation of the floral onset is well established (Hyun
et al., 2016; Wang *et al.*, 2009; Yamaguchi *et al.*, 2014), but there is some dispute about whether
or not SPL3, SPL4, and SPL5 are important. While overexpression of *SPL3, SPL4*, and *SPL5*
induces flowering (Jung *et al*, 2016; Wu & Poethig, 2006), a triple T-DNA insertion mutant was
previously reported to not display any flowering phenotype (Xu *et al*, 2016). However, a recent
study analyzed a triple deletion mutant generated by CRISPR/*Cas9* in relation to light signaling
and also reported on a delay in flowering time (Xie *et al.*, 2020). Similarly, we generated individual
deletion mutants in the respective SPLs using CRISPR/*Cas9* (Figure S23). As compared to single
spl mutant and wild-type plants, higher order *spl3*, *spl4* and *spl5* deletion mutants are significantly
later flowering in all lines with *spl4* (Figure 5B-D; Table S1). This argues for an important role of
SPL4 in inducing flowering. *SPL4*, similar to *SPL3* and *SPL5,* is induced at the wild-type SAM at
floral transition (Schmid *et al*, 2003). It is expressed in the center of the SAM, in a domain
overlapping with the WUS protein, while *SPL3*, *SPL5*, *SPL9*, and *SPL15* are expressed at the
periphery of the SAM and in the vasculature of young leaves (Daum *et al*, 2014; Hyun *et al.*,

2016; Olas *et al.*, 2019; Wang *et al.*, 2009; Yadav *et al.*, 2011). This denotes a direct regulation of
SPL4 by WUS, which would explain increased *SPL4* levels in the *clv3* mutant (Figure 5A) and the
previously identified partial miR156-dependent regulation of the *SPL*s by the T6P pathway (Wahl
et al., 2013). Indeed, we observed reduced expression of *SPL4* in *wus-7* apices, a weak allele
with a mutation in the WUS DNA binding domain resulting in smaller but maintained SAM tissue
(Graf *et al*, 2010; Lin *et al*, 2016). In contrast, none of the other *SPL*s were reduced (Figures 5E;
S22C) and miR156 levels did not change (Figure 5E). This is unexpected as a recent publication
demonstrated decreased miR156 levels and increased *SPL9* expression in apices of a *wus-5*
mutant (Fouracre & Poethig, 2019). *wus-5* is however a strong *wus* null allele with very little
residual SAM tissue (McElver *et al*, 2001; Sonoda *et al*, 2007), which might explain the difference.
We identified a larger number of potential *SPL4*^{WUS} sites when compared to the other *SPL*s
(Figures 5F; S24). A transactivation assay demonstrated induction of the luciferase reporter
coupled to 5'*SPL4* sequences by WUS, with the shortest fragment covering most of the inductive
effect (Figure 5G). This sequence contains five *SPL4*^{WUS} sites, four of which could be separately
tested by ChIP-PCR. This indeed confirmed direct induction of *SPL4* by WUS (Figures 5H; S25).
However, these results also suggest that additional players downstream of the WUS/CLV3 feed-
back loop are important for the onset of flowering, which cannot be bypassed by an otherwise
inductive SPL4 (Figure 5A).

A feed-back regulation between the T6P pathway and WUS involves the age pathway. An
enlarging SAM at floral transition correlates with a larger *WUS* expression domain expanding into
the CZ and the outer cell layer (L1) (Figure 1D-F). Interestingly, we found reduced *TPS1* levels
(*35S:amiRTPS1*) lead to an enlarged OC domain marked by *WUS* (Figure 6A) that overlaps with
the CZ, marked by *CLV3* expression (Figure 6A). Plants overexpressing *TPS1* (*35S:TPS1*, Figure
S2) display an increased stem cell pool with otherwise little effect on *WUS* expression (Figure
6A). The size of the CZ decreases when expressing *TPS1* under the *CLV3* promoter (Figures 6A;
S2), indicating a smaller stem cell pool in support of a much smaller SAM size of the very early
flowering *CLV3:TPS1* (Wahl *et al.*, 2013).
TPPJ overexpression results in significantly enlarged SAMs (Figure 2F), and importantly,
expands *WUS* expression into the outer SAM layers, while *CLV3* expression seems to be little
affected (Figure 6B,C). Consistent with this, *CLV3* expression increases to much higher levels in
ML1:amiRTPPJ (Figure 6C), suggesting an active role of the T6P pathway in the outer meristem
layer regarding stem cell maintenance. In addition, *WUS* is induced (Figures 6C-D), indicating
uncoupling of the WUS/CLV3 feed-back loop downstream of the T6P pathway.
We next investigate whether this is in part due to the function of the age pathway downstream of
the T6P pathway. While we found *WUS* to be induced (Figure S26A,B), its expression seems to
be reduced in the CZ in the transition SAM of all but *spl3,5* mutant SAMs (Figure S26A). This

cannot be explained by an altered expression of *CLV3*, as we found it generally slightly reduced
in all mutant lines (Figure S26C,D).

Since *WUS* and *SPL4* expression domains are largely overlapping (Daum *et al.*, 2014; Olas *et al.*,
2019; Torti *et al*, 2012; Yadav *et al.*, 2011), we asked whether a feed-back loop, involving direct
regulation of *WUS* by SPL4, would explain the observed expression patterns. We identified one
WUS^{SPL4} site -1067bp upstream of the *WUS* coding sequence (Figure 6E). SPL4 significantly and
specifically reduced reporter gene expression in an Arabidopsis protoplast-based transactivation
assay, indicating that it can repress *WUS in vivo* (Figure 6F,G). EMSA confirmed this result
(Figure 6H). This finding is supported by a promoter study analyzing a series of deletion
constructs of the 5'*WUS* sequence (Baurle & Laux, 2005). Among them one which deleted a
sequence including the *WUS*^{SPL4} site (Δ4, -941/-604 from transcription start site; -1067/-730
upstream of coding sequence). Importantly, this deletion construct led to a stronger activation
while others rather led to a reduction of a GUS reporter, especially when affecting the sequences
close to the transcription start site. This indicates the loss of regulatory sequences which are
targeted by repressors of WUS such as SPL4 (Baurle & Laux, 2005). Taken together, our results
demonstrate a likely transient negative feed-back regulation between SPL4 and WUS.

Conclusion

Taken together, we unraveled dynamic feed-back regulations between sugar signaling, central
meristem maintenance and flowering time regulators that coordinate cell proliferation and
metabolism, providing the basis and energy necessary for doming during floral transition.

Proliferation, self-renewal and differentiation of cells originating from stem cells are the basis for
the plants' remarkable plasticity. Stem cell systems are formed during embryogenesis. The SAM
as primary meristem is retained stably throughout plant development (Pfeiffer *et al.*, 2017). The
initiation of reproductive growth marks a very prominent change in a plant's morphology and
requires large-scale reorganization of SAM architecture and sink-source relationships within all
flowering plants. Within cells, sucrose is an important basis for energy production and biomass.
To prevent starvation, energy-demanding developmental transitions are tightly coordinated with
endogenous sucrose availability through intricate signaling systems (Fernie *et al*, 2020). The
sucrose signaling T6P pathway was previously found to impact on cell cycle regulators (Gómez *et*
al., 2006). These were also demonstrated as differentially expressed at the transition stage
(Klepikova *et al.*, 2015).

We show that in Arabidopsis the negative feed-back loop between the central meristem
maintenance regulators WUS and CLV3, considered as stable until now, uncouples during floral
transition. As a result, the *WUS* expression domain is no longer fixed in the center of the SAM but

stretches into the outer meristem layers demonstrating unreported OC dynamics. WUS directly

regulates *TPPJ*, a component of the T6P pathway, which has an impact on SAM size and thus

seems important for cell proliferation and doming of the SAM during the transition phase. Notably,

organ production is stalled when *WUS* is ectopically expressed from a *CLV3* promoter into the

stem cell niche, and correlates with massive proliferation of meristematic cells (Brand *et al*, 2002).

Although *TPPJ* is expressed at the rim of the PZ, it can be activated by WUS *in vivo*, requiring the

cell non-autonomous function of a mobile WUS (Daum *et al.*, 2014; Yadav *et al.*, 2011). The fact

that WUS does not activate *TPPJ* outside of this domain suggests other factors repressing it in

the center of the SAM. These factors are clearly still active in a *clv3* mutant background, where

the activation of *TPPJ* by WUS is restricted to the outer SAM layers, suggesting factors outside of

this central regulation of the meristem maintenance network (Pfeiffer *et al.*, 2017). It will be

interesting to learn more about its regulation in the future.

We demonstrate that SPL4, which is expressed in a domain overlapping with the central *WUS*

expression domain, directly and negatively regulates *WUS.* In turn, WUS activates *SPL4* in a

likely transient negative feed-back regulation, which we suggest to balance sugar-signaling-

dependent effects on meristem maintenance at transition. In addition, WUS might repress *SPL5*,

as we found slightly but significantly increased expression in *wus-7* apices (Figure 5E). Although

they don't seem to respond in a WUS-dependent manner, it remains to be determined whether

those SPLs expressed in the periphery of the SAM (SPL3, SPL5 and SPL9) are mobile and

therefore independently report on the sucrose status to the central region of the SAM. We

suggest that the combined action of the T6P pathway and SPLs expressed at the SAM

determines the duration of the doming phase.

An important next step will be to understand how SnRK1 and TOR signaling (Baena-González &

Lunn, 2020; Caldana *et al*, 2019; Pfeiffer *et al.*, 2016) fit into the emerging picture of the

mechanisms regulating floral transition in the future. Apart from the regulation of flowering time

(Wahl *et al.*, 2013), it is interesting that the T6P pathway also influences the re-organisation of the

SAM during floral transition. Given the ubiquitous nature of carbohydrate signaling and the large-

scale change in sink-source relationships within plants (Fernie *et al.*, 2020), it will be interesting to

determine if this regulatory mechanism is widely present in the plant kingdom.

Materials and Methods

<u>Plant material and growth conditions</u>

All Arabidopsis plants were of the Columbia accession (Col-0), except for *wus-7*, which is in

the Ler background (Graf *et al.*, 2010). *clv3-7* (Wisman *et al*, 1998), *clv3-10* (Forner *et al*, 2015),

35S:amiRTPS1, *CLV3:TPS1* (Wahl *et al.*, 2013), and *TCSn:GFP* (Zurcher *et al.*, 2013) were

described previously. The *tppj* insertion mutant (At5g65140, GABI_215C08) was obtained from

the Arabidopsis resource stock center (ABRC). Growth chambers were set to 22°C in LD (16h

light/8h dark) or SD (8h dark/16h light) with a light intensity of approximately 160 μmol/m^{-2}s^{-1} and

a relative humidity of 60-65%. Synchronized induction of flowering was performed as described

(Schmid *et al.*, 2003).

Plant phenotyping

Flowering time of on average of 20 plants per genotype was scored as days to flowering,

when shoots were 0.5cm (bolting), and by the total leaf number (TLN), i.e., the sum of rosette leaf

(RLN) and cauline leaf numbers (CLN) (Table S1). Meristem size was measured as the area

between two organ primordia under the meristem summit of a longitudinal middle section through

at least 10 apices from individual plants using the Fiji software version 2.0.0-rc-69/1.52

(Schindelin *et al*, 2012).

Generation of transgenic lines

Plants were transformed (Clough & Bent, 1998), confirmed by PCR and independent,

single-insertion, homozygous T3 plants were used for all studies. Oligonucleotides for cloning and

for genotyping are given in Tables S2 and S3, respectively. For *35S:TPPJ* and *35S:TPS1* lines,

coding sequences of *TPPJ* (At5g65140) and *TPS1* (At1g78580) were cloned via the Gateway®

entry vector *pJLBlue* reverse (Mathieu *et al*, 2007) into a *pGREEN-II*-based destination vector.

Artificial microRNAs targeting *TPPJ* (*ML1:amiRTPPJ* V1 and V2, *35S:amiRTPPJ* V1 and V2)

were designed with the Web MicroRNA Designer (http://wmd3.weigelworld.org/cgi-

bin/webapp.cgi)(Schwab *et al.*, 2006). The *Eco*RI/*Bam*HI fragment was cloned via *pJLBlue*

reverse (Mathieu *et al.*, 2007) into a *pGREEN-II*-based destination vector with either the *ML1* or

35S promoter. *spl3, spl4, spl5* knockout lines were generated with the CRISPR/*Cas9* technology

(Ruf *et al*, 2019; Wang *et al*, 2015). The NGG PAM recognition sites were defined using the

ATUM webtool (https://www.atum.bio) (Figure S23). The *pJF1033* vector, containing a two single

guided RNAs scaffold, was used as a template. *Bsa*I products were cloned into the *pJF1031*

binary vector (Ruf *et al.*, 2019). Plants with homozygous deletions were back-crossed to Col-0.

Homozygous, *Cas9*-free lines were used to generate all higher order mutant lines.

RT-qPCR

Total RNA was isolated by a modified phenol/chloroform method using a TRIzol reagent

(phenol 38% (v/v), guanidine thiocyanate 0.8M ammonium thiocyanate 0.4M, sodium acetate

0.1M pH5.0, glycerol 5% (v/v), EDTA 5mM pH8.0, Na-lauroylsarcosine 0.5% (v/v)), followed by
sodium acetate precipitation. Genomic DNA was removed with DNase I RNase-free
(Ambion™/Thermo Fisher Scientific, Waltham, Massachusetts, US), and cDNA synthesis carried
out using a SuperScript™IV Reverse Transcriptase Kit (Thermo Fisher Scientific, Waltham,
Massachusetts, US) according to the instructions. Mature miR156 stem-loop primers (Table S4)
were added to the cDNA synthesis reaction (1:1 with oligo dT(18) primer) (Varkonyi-Gasic *et al*,
2007).

RT-qPCR was performed with the ABI Prism 7900 HT fast real time PCR system (Applied
Biosystems/Life Technologies, Darmstadt, Germany) using a Power SYBR® Green-PCR Master
Mix (Applied Biosystems/Life Technologies, Darmstadt, Germany) for expression analyses and
ChIP-PCRs. Oligos are listed in Table S4 (RT-qPCR) and S5 (ChIP-PCR). For analysis the SDS
2.4 software (Applied Biosystems/Life Technologies, Darmstadt, Germany) was used
(Czechowski *et al*, 2004). cDNA quality was determined and expression values were calculated
(Wahl *et al.*, 2013).

RNA *in situ* hybridization

Probes were synthesized using the DIG RNA Labeling Kit (Roche, Mannheim, Germany) on
CDSs of the target genes cloned into the *pGEM®-T Easy* vector (Promega, Madison, Wisconsin,
US). Oligonucleotides and construct IDs are listed in Table S6. Wax embedding, sectioning and
RNA *in situ* hybridization, and imaging were performed as described (Wahl *et al.*, 2013).

ChIP-PCR

For ChIP-PCR, 100 apices or 1.5g leaves per replicate (Col-0, *clv3-7* and *clv3-10*) were
fixed in 1% (v/v) formaldehyde buffer (10mM sodium phosphate buffer, pH7; 50mM NaCl; 100mM
sucrose) under vacuum. ChIP was performed (Kaufmann *et al*, 2010) with modifications: Antibody
incubation (anti-WUS; AS11 1759; Agrisera) was extended o/n at 4°C and Agarose beads
(Protein A-Agarose; sc-2001; Santa Cruz Biotechnology) to 6h at 4°C. Immunoprecipitated DNA
was analyzed by RT-qPCR. Ct values of *TPPJ* and *SPL4* promoter regions were normalized to
the Ct value of a region within the *UBQ10* promoter. The % of enrichment was calculated as
relative to expression of the input of the individual regions. ChIP on apices of Col-0, *clv3-7* and
clv3-10 without antibody and ChIP on Col-0, *clv3-7* and *clv3-10* leaves were controls (Figure
S10). Note that in contrast to the input no amplification was observed in the controls (Figure S11,
S25) and the previously published confirmed WUS target sites were detected (Figure
S12)(Leibfried *et al.*, 2005; Yadav *et al.*, 2011). Oligonucleotides are listed in Table S5.

 Transactivation assay

For the effector line, the *WUS* (At2g17950) and *SPL4* (At1g53160) CDS were cloned via the

Gateway® entry vector *pJLBlue* reverse (Mathieu *et al.*, 2007) into the *pMDC32* Gateway® vector

(pMML058) and the *pGW5* (pKK77). For the reporter constructs, designated *5'TPPJ* regions were

*Kpn*I/*Acy*I cloned into the Gateway® *pMDC162* vector to obtain GUS fusions and *5'SPL4* and

5'WUS regions were *Kpn*I/*Spe*I cloned into *pGREEN800II-LUC* (Hellens *et al*, 2005)(Table S2).

35Somega:LUC-NOS was used as transformation control in the *5'TPPJ* assays. Protoplasts were

isolated from 4-week-old plants, transfected (Wu *et al*, 2009) and incubated for 20h at 22°C and

100μmol/m^{-2}s^{-1}. Luciferase activity of the *5'TPPJ, 5'SPL4* and *5'WUS* assays was measured with

a luciferase and dual-luciferase reporter assay system, respectively (Promega, Madison,

Wisconsin, US). GUS activity was determined as described (Yoo *et al*, 2007).

EMSA

The *WUS* CDS without STOP codon was cloned via the Gateway® entry vector *pDONR207*

(pMML059) into a Gateway® destination vector (*pDEST24*). *35S:WUS-GST* (pMML063) was

transformed into *Escherichia coli* Rosetta plysS cells and protein production induced with 1mM

isopropyl β-D-1-thiogalactopyranoside at 30°C o/n. Cell lysis was performed with 1x sonication of

5 sec, 20% power, 4 cycles (Sonoplus Hd 2070 Sonicator, Badelin, Berlin, Germany) in a freshly

prepared buffer (20mM Na-phosphate buffer, pH7.4; 0.5MNaCl; 1mM phenylmethylsulfonyl

fluoride; 1mM Ethylenediaminetetraacetic acid; 1 tablet cOmplete™ Protease Inhibitor Cocktail

(Merck, Darmstadt, Germany) per 10ml buffer). 2μg protein (crude extract, determined by

Pierce^{TM} BCA Protein Assay Kit, Thermo Fisher Scientific, Waltham, Massachusetts, US) was

used for the EMSA. 50bp, double-stranded, 5′-IRDey-682-labeled probes, spanning the putative

TPPJ^{WUS} and *WUS^{SPL4}* sites are listed in Table S7 (5'*TPPJ*) and Table S8 (5'*WUS*). Binding

reactions were carried out with the Odyssey® EMSA Kit (LI-COR®) according to the instructions

with a competitor and/or mutated competitor to probe ratio of 1:200 and visualized with the

Oddysey Infrafed Imaging System (Li-Cor, Lincoln, NE).

Confocal microscopy

SAMs of *a TCSn:GFP* cytokinin reporter line (Zurcher *et al.*, 2013) were imaged with a Leica

SP8 confocal laser scanning system equipped with a M6000B-CS microscopy stage, an Argon

laser (65mV), and a 40x water immersion HCX APO objective as follows: Laser output power

20%; GFP excitation (green, wavelength 488nm), emission detection channel 3 (495-520nm),

gain PMT 800V; plastid auto-fluorescence (blue), emission detection channel 4 (700-800nm),

gain PMT ~500V; scan speed 600Hz in xyz bi-directional scanning mode with a z-stack distance
of approx. 10µm. Offset=0; pixel dimension: 1024×1024. Middle sections of representative SAMs
were extracted from z-stacks using the Fiji software version 2.0.0-rc-69/1.52 (Schindelin *et al.*,
2012).

<u>Statistical consideration</u>

Statistical significance was analyzed by both one-way ANOVA (Analysis of Variance) with
Tukey' Post Hoc HSD (Honestly Significant Difference) based on Tukey-Kramer correction (P
value <0.05) or a two-tailed Student's *t*-test.

Acknowledgments

The authors wish to thank all members of the Wahl group for fruitful discussions, M. Schmid and
P. Wigge for critical reading of the manuscript, D. Walter for input on statistical analyses, K.
Kontbay for providing the SPL4 effector construct, M. Liang and J. Van Dingenen for their support
with the *spl* CRISPR/*Cas9* deletion lines, M. Molochko for sharing the *35Somega:LUC-NOS*
plasmid, J. Forner for providing the *pJF1031* and *pJF1033* plasmids. Work in the Wahl group was
supported by the BMBF (SolaMI, 031B0191), the DFG (SPP1530: WA3639/1-2), and the Max
Planck Society.

Author Contributions

VW conceived and designed the experiments and prepared the figures with contributions from
MML. VW and MML analyzed the data and wrote the manuscript. MML, KF, AK, and VW
performed all essential experiments, i.e. VW generated most of the lines used, except for
ML1:amiRTPPJ (KF), the *spl* CRISPR/Cas9 deletion lines (MML), and all crosses thereof (MML).
MML performed all transactivation assays, ChIP-PCRs, EMSAs and RT-qPCR analyses. MML,
AK, KF, CA, and VW analyzed SAM sizes and scored flowering time. RNA *in situ* hybridizations
were performed by VW, AK, KF and MML. FK and VW took the confocal images. All authors have
read and commented on the text and figures within this manuscript.

Conflict of interest

The authors declare that they don't have any competing commercial interests.

Data Availability Section

Supporting information is provided in a separate file. This includes:

Figures S1 to S26

Tables S1 to S8

SI References

References

Andres F, Coupland G (2012) The genetic basis of flowering responses to seasonal cues. *Nat*
Rev Genet 13: 627-639

Baena-González E, Lunn JE (2020) SnRK1 and trehalose 6-phosphate - two ancient pathways
converge to regulate plant metabolism and growth. *Curr Opin Plant Biol* 55: 52-59

Baurle I, Laux T (2005) Regulation of WUSCHEL transcription in the stem cell niche of the
Arabidopsis shoot meristem. *Plant Cell* 17: 2271-2280

Brand U, Fletcher JC, Hobe M, Meyerowitz EM, Simon R (2000) Dependence of stem cell fate in
Arabidopsis on a feedback loop regulated by CLV3 activity. *Science (New York, NY)* 289: 617-
619

Brand U, Grunewald M, Hobe M, Simon R (2002) Regulation of CLV3 expression by two
homeobox genes in Arabidopsis. *Plant Physiol* 129: 565-575

Busch W, Miotk A, Ariel FD, Zhao Z, Forner J, Daum G, Suzaki T, Schuster C, Schultheiss SJ,
Leibfried A *et al* (2010) Transcriptional control of a plant stem cell niche. *Dev Cell* 18: 849-861

Caldana C, Martins MCM, Mubeen U, Urrea-Castellanos R (2019) The magic 'hammer' of TOR:
the multiple faces of a single pathway in the metabolic regulation of plant growth and
development. *Journal of Experimental Botany* 70: 2217-2225

Clough SJ, Bent AF (1998) Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. *Plant J* 16: 735-743

Czechowski T, Bari RP, Stitt M, Scheible WR, Udvardi MK (2004) Real-time RT-PCR profiling of
over 1400 Arabidopsis transcription factors: unprecedented sensitivity reveals novel root- and
shoot-specific genes. *Plant J* 38: 366-379

Daum G, Medzihradszky A, Suzaki T, Lohmann JU (2014) A mechanistic framework for noncell
autonomous stem cell induction in Arabidopsis. *Proc Natl Acad Sci U S A* 111: 14619-14624

Fernie AR, Bachem CWB, Helariutta Y, Neuhaus HE, Prat S, Ruan YL, Stitt M, Sweetlove LJ,
Tegeder M, Wahl V *et al* (2020) Synchronization of developmental, molecular and metabolic
aspects of source-sink interactions. *Nat Plants* 6: 55-66

Fichtner F, Lunn JE (2021) The Role of Trehalose 6-Phosphate (Tre6P) in Plant Metabolism and
Development. *Annu Rev Plant Biol*

Fichtner F, Olas JJ, Feil R, Watanabe M, Krause U, Hoefgen R, Stitt M, Lunn JE (2020)
Functional Features of TREHALOSE-6-PHOSPHATE SYNTHASE1, an Essential Enzyme in
Arabidopsis. *Plant Cell* 32: 1949-1972

Forner J, Pfeiffer A, Langenecker T, Manavella PA, Lohmann JU (2015) Germline-transmitted
genome editing in Arabidopsis thaliana Using TAL-effector-nucleases. *PLoS One* 10: e0121056

Fouracre JP, Poethig RS (2019) Role for the shoot apical meristem in the specification of juvenile
leaf identity in Arabidopsis. *Proc Natl Acad Sci U S A* 116: 10168-10177

Gómez LD, Baud S, Gilday A, Li Y, Graham IA (2006) Delayed embryo development in the
ARABIDOPSIS TREHALOSE-6-PHOSPHATE SYNTHASE 1 mutant is associated with altered
cell wall structure, decreased cell division and starch accumulation. *Plant J* 46: 69-84

Graf P, Dolzblasz A, Wurschum T, Lenhard M, Pfreundt U, Laux T (2010) MGOUN1 encodes an
Arabidopsis type IB DNA topoisomerase required in stem cell regulation and to maintain
developmentally regulated gene silencing. *Plant Cell* 22: 716-728

Hellens RP, Allan AC, Friel EN, Bolitho K, Grafton K, Templeton MD, Karunairetnam S, Gleave
AP, Laing WA (2005) Transient expression vectors for functional genomics, quantification of
promoter activity and RNA silencing in plants. *Plant Methods* 1: 13

Hyun Y, Richter R, Coupland G (2017) Competence to Flower: Age-Controlled Sensitivity to
Environmental Cues. *Plant Physiol* 173: 36-46

Hyun Y, Richter R, Vincent C, Martinez-Gallegos R, Porri A, Coupland G (2016) Multi-layered
Regulation of SPL15 and Cooperation with SOC1 Integrate Endogenous Flowering Pathways at
the Arabidopsis Shoot Meristem. *Dev Cell* 37: 254-266

Jacqmard A, Gadisseur I, Bernier G (2003) Cell division and morphological changes in the shoot
apex of Arabidopsis thaliana during floral transition. *Ann Bot* 91: 571-576

Jung JH, Lee HJ, Ryu JY, Park CM (2016) SPL3/4/5 Integrate Developmental Aging and
Photoperiodic Signals into the FT-FD Module in Arabidopsis Flowering. *Mol Plant* 9: 1647-1659

Kaufmann K, Muino JM, Osteras M, Farinelli L, Krajewski P, Angenent GC (2010) Chromatin
immunoprecipitation (ChIP) of plant transcription factors followed by sequencing (ChIP-SEQ) or
hybridization to whole genome arrays (ChIP-CHIP). *Nat Protoc* 5: 457-472

Kinoshita A, Vayssieres A, Richter R, Sang Q, Roggen A, van Driel AD, Smith RS, Coupland G
(2020) Regulation of shoot meristem shape by photoperiodic signaling and phytohormones during
floral induction of Arabidopsis. *Elife* 9

Klepikova AV, Logacheva MD, Dmitriev SE, Penin AA (2015) RNA-seq analysis of an apical
meristem time series reveals a critical point in Arabidopsis thaliana flower initiation. *BMC*
Genomics 16: 466

Kwiatkowska D (2008) Flowering and apical meristem growth dynamics. *J Exp Bot* 59: 187-201

Leibfried A, To JP, Busch W, Stehling S, Kehle A, Demar M, Kieber JJ, Lohmann JU (2005)
WUSCHEL controls meristem function by direct regulation of cytokinin-inducible response
regulators. *Nature* 438: 1172-1175

Lin TF, Saiga S, Abe M, Laux T (2016) OBE3 and WUS Interaction in Shoot Meristem Stem Cell
Regulation. *PLoS One* 11: e0155657

Lohmann JU, Hong RL, Hobe M, Busch MA, Parcy F, Simon R, Weigel D (2001) A molecular link
between stem cell regulation and floral patterning in Arabidopsis. *Cell* 105: 793-803

Mathieu J, Warthmann N, Kuttner F, Schmid M (2007) Export of FT protein from phloem
companion cells is sufficient for floral induction in Arabidopsis. *Curr Biol* 17: 1055-1060

McElver J, Tzafrir I, Aux G, Rogers R, Ashby C, Smith K, Thomas C, Schetter A, Zhou Q,
Cushman MA *et al* (2001) Insertional mutagenesis of genes required for seed development in
Arabidopsis thaliana. *Genetics* 159: 1751-1763

Olas JJ, Van Dingenen J, Abel C, Dzialo MA, Feil R, Krapp A, Schlereth A, Wahl V (2019) Nitrate
acts at the Arabidopsis thaliana shoot apical meristem to regulate flowering time. *New Phytol* 223:
814-827

Pfeiffer A, Janocha D, Dong Y, Medzihradszky A, Schone S, Daum G, Suzaki T, Forner J,
Langenecker T, Rempel E *et al* (2016) Integration of light and metabolic signals for stem cell
activation at the shoot apical meristem. *Elife* 5

Pfeiffer A, Wenzl C, Lohmann JU (2017) Beyond flexibility: controlling stem cells in an ever
changing environment. *Curr Opin Plant Biol* 35: 117-123

Ponnu J, Schlereth A, Zacharaki V, Dzialo MA, Abel C, Feil R, Schmid M, Wahl V (2020) The
Trehalose 6-Phosphate Pathway Impacts Vegetative Phase Change in Arabidopsis thaliana.
Plant J

Ruf S, Forner J, Hasse C, Kroop X, Seeger S, Schollbach L, Schadach A, Bock R (2019) High-
efficiency generation of fertile transplastomic Arabidopsis plants. *Nat Plants* 5: 282-289

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden
C, Saalfeld S, Schmid B *et al* (2012) Fiji: an open-source platform for biological-image analysis.
Nat Methods 9: 676-682

Schmid M, Uhlenhaut NH, Godard F, Demar M, Bressan R, Weigel D, Lohmann JU (2003)
Dissection of floral induction pathways using global expression analysis. *Development* 130: 6001-
6012

Schwab R, Ossowski S, Riester M, Warthmann N, Weigel D (2006) Highly specific gene silencing
by artificial microRNAs in Arabidopsis. *Plant Cell* 18: 1121-1133

Sloan J, Hakenjos JP, Gebert M, Ermakova O, Gumiero A, Stier G, Wild K, Sinning I, Lohmann
JU (2020) Structural basis for the complex DNA binding behavior of the plant stem cell regulator
WUSCHEL. *Nat Commun* 11

Snipes SA, Rodriguez K, DeVries AE, Miyawaki KN, Perales M, Xie M, Reddy GV (2018)
Cytokinin stabilizes WUSCHEL by acting on the protein domains required for nuclear enrichment
and transcription. *PLoS Genet* 14: e1007351

Sonoda Y, Yao SG, Sako K, Sato T, Kato W, Ohto MA, Ichikawa T, Matsui M, Yamaguchi J,
Ikeda A (2007) SHA1, a novel RING finger protein, functions in shoot apical meristem
maintenance in Arabidopsis. *Plant J* 50: 586-596

Srikanth A, Schmid M (2011) Regulation of flowering time: all roads lead to Rome. *Cellular and*
molecular life sciences : CMLS 68: 2013-2037

Torti S, Fornara F, Vincent C, Andres F, Nordstrom K, Gobel U, Knoll D, Schoof H, Coupland G
(2012) Analysis of the Arabidopsis shoot meristem transcriptome during floral transition identifies
distinct regulatory patterns and a leucine-rich repeat protein that promotes flowering. *The Plant*
cell 24: 444-462

Varkonyi-Gasic E, Wu R, Wood M, Walton EF, Hellens RP (2007) Protocol: a highly sensitive RT-
PCR method for detection and quantification of microRNAs. *Plant Methods* 3: 12

Wahl V, Ponnu J, Schlereth A, Arrivault S, Langenecker T, Franke A, Feil R, Lunn JE, Stitt M,
Schmid M (2013) Regulation of flowering by trehalose-6-phosphate signaling in Arabidopsis
thaliana. *Science (New York, NY)* 339: 704-707

Wang JW, Czech B, Weigel D (2009) miR156-regulated SPL transcription factors define an
endogenous flowering pathway in Arabidopsis thaliana. *Cell* 138: 738-749

Wang ZP, Xing HL, Dong L, Zhang HY, Han CY, Wang XC, Chen QJ (2015) Egg cell-specific
promoter-controlled CRISPR/Cas9 efficiently generates homozygous mutants for multiple target
genes in Arabidopsis in a single generation. *Genome Biol* 16: 144

Wisman E, Cardon GH, Fransz P, Saedler H (1998) The behaviour of the autonomous maize
transposable element En/Spm in Arabidopsis thaliana allows efficient mutagenesis. *Plant Mol Biol*
37: 989-999

Wu FH, Shen SC, Lee LY, Lee SH, Chan MT, Lin CS (2009) Tape-Arabidopsis Sandwich - a
simpler Arabidopsis protoplast isolation method. *Plant Methods* 5: 16

Wu G, Poethig RS (2006) Temporal regulation of shoot development in Arabidopsis thaliana by
miR156 and its target SPL3. *Development (Cambridge, England)* 133: 3539-3547

Xie Y, Zhou Q, Zhao Y, Li Q, Liu Y, Ma M, Wang B, Shen R, Zheng Z, Wang H (2020) FHY3 and
FAR1 Integrate Light Signals with the miR156-SPL Module-Mediated Aging Pathway to Regulate
Arabidopsis Flowering. *Mol Plant* 13: 483-498

Xu M, Hu T, Zhao J, Park MY, Earley KW, Wu G, Yang L, Poethig RS (2016) Developmental
Functions of miR156-Regulated SQUAMOSA PROMOTER BINDING PROTEIN-LIKE (SPL)
Genes in Arabidopsis thaliana. *PLoS Genet* 12: e1006263

Yadav RK, Perales M, Gruel J, Girke T, Jonsson H, Reddy GV (2011) WUSCHEL protein
movement mediates stem cell homeostasis in the Arabidopsis shoot apex. *Genes Dev* 25: 2025-
2030

Yamaguchi N, Winter CM, Wu MF, Kanno Y, Yamaguchi A, Seo M, Wagner D (2014) Gibberellin
acts positively then negatively to control onset of flower formation in Arabidopsis. *Science (New*
York, NY) 344: 638-641

Yang L, Xu M, Koo Y, He J, Poethig RS (2013) Sugar promotes vegetative phase change in
Arabidopsis thaliana by repressing the expression of MIR156A and MIR156C. *Elife* 2: e00260

Yoo SD, Cho YH, Sheen J (2007) Arabidopsis mesophyll protoplasts: a versatile cell system for
transient gene expression analysis. *Nat Protoc* 2: 1565-1572

Yu S, Cao L, Zhou CM, Zhang TQ, Lian H, Sun Y, Wu J, Huang J, Wang G, Wang JW (2013)
Sugar is an endogenous cue for juvenile-to-adult phase transition in plants. *Elife* 2: e00269

Zurcher E, Tavor-Deslex D, Lituiev D, Enkerli K, Tarr PT, Muller B (2013) A robust and sensitive
synthetic sensor to monitor the transcriptional output of the cytokinin signaling network in planta.
Plant Physiol 161: 1066-1075

All main figures for

Sugar Signaling Induces Dynamic Changes During Meristem Development to Regulate Flowering in Arabidopsis

Magdalena Musialak-Lange[1], Katharina Fiddeke[1,#], Annika Franke[1,#], Friedrich Kragler[1], Christin Abel[1], and Vanessa Wahl[1,*]

Affiliations: [1]Max Planck Institute of Molecular Plant Physiology, Department of Metabolic Networks, Am Mühlenberg 1, Potsdam, 14476, Germany.

[#]K.F. and A.F. contributed equally

[*]**Correspondence**: Vanessa Wahl

Email: vanessa.wahl@mpimp-golm.mpg.de

Synopsis:

Schematic illustration of dynamic feedback-regulations between sugar signaling, meristem maintenance and the age pathway of the flowering network in the transition SAM as compared to the vegetative SAM.
During vegetative growth SAM maintenance is controlled by spatially separated *WUS* and *CLV3* expression in a negative feedback-loop. At floral transition increased activity of the T6P pathway in the SAM uncouples this regulation. This results in an unreported special relocation of *WUS* expression and involves a transient negative feedback-loop between WUS and SPL4.

Figures

Figure 1. The T6P pathway impacts SAM size during development. (*A*) SAM area throughout development. n=15 (*B*) Vegetative and (*C*) inflorescence SAM size of *CLV3:TPS1* and *35S:amiRTPS1* lines. n=10. (*D*) *WUS* and *CLV3* expression by RNA *in situ* hybridization in vegetative (6 and 8 DAG), transition (10 DAG, marked dark grey) and inflorescence SAMs (12 and 16 DAG) of LD-grown Col-0 plants. (*E*) *WUS* expression domain sizes, (*F*) *WUS* expression domain distance to SAM summit, and (*G*) *CLV3* expression domain area, in vegetative, transition (dark grey) and inflorescence SAMs of LD-grown wild-type plants. n>10. Error bars denote s.d.; significance was calculated based on a Student's *t*-test, ***P<0.001. Star indicates SAM summit. Scale bars 25µm.

Figure 2. *WUS* and *CLV3* expression in response to changes of *TPPJ* in the SAM. (*A*, *B*) *TPPJ* expression by RNA *in situ* hybridization on longitudinal sections through inflorescence SAMs of (*A*) Col-0, and (*B*) *clv3-7*, and by RT-qPCR in apices collected from (*C*) *clv3-7*, *clv3-10*, and (*D*) *35S:TPPJ* plants. n=3. (*E*) Expression of *TPPJ* by RNA *in situ* hybridization on longitudinal sections through inflorescence SAMs of Col-0 and *35S:TPPJ.* (*F*) SAM size of plants overexpressing *TPPJ*. n=10. Error bars denote s.d.; significance calculated by one-way ANOVA (C,D) and Student's *t*-test (F), ***P<0.001. Stars indicates SAM summit. Scale bars are 50μm.

Figure 3. WUS directly regulates *TPPJ* in the SAM. (*A*) Overview of *TPPJ* 5' regulatory region with putative *TPPJ^WUS* sites (grey circles, black boxes), position of ChIP-PCR amplicons corresponding to the results shown in (C-E). Boxes marked with I, II, and III indicate *5' TPPJ* regions with in total seven confirmed core *TPPJ^WUS* sites – I: -2795 – -2789 bp, II: -2073 – -1830 bp, and III: -652 – -564 bp. Sequence location and lengths used in (B) indicated with #1-6. (*B*) Protoplast transactivation assay showing activation of the GUS reporter when coupled to the regions indicated in (A), relative to LUC activity. c indicates untransformed control. n=6. (*C-E*) Enrichment of (*C*) region I, (*D*) region II, and (*E*) region III as indicated in (A) measured by ChIP-PCR relative to the input. PB – post binding fraction. n=3. (*F*) EMSA for WUS binding to the indicated regions (A, I-III). Shifted band in the presence of WUS protein (open arrow head), non-shifted fraction (closed arrow head). Error bars denote s.d.; significance based on one-way ANOVA, ***P<0.001. Scale bars are 50μm.

Figure 4. The role of TPPJ in the outer SAM layer. (*A*) Meristem area of Col-0, *ML1:amiRTPPJ* V1 and V2. n=10. (*B,C*) Flowering time of (*B*) *ML1:amiRTPPJ* and (*C*) *clv3-10;ML1:amiRTPPJ* shown as days to bolting, relative to the wild type. V1 and V2 indicate two independent versions of artificial microRNAs designed to target *TPPJ* transcript. n=22. (*D*) Relative expression of *SPL* genes in SD-grown *ML1:amiRTPPJ* and *35S:TPPJ* at 40 days after germination. n=4. Error bars denote s.d.; significance calculated based on one-way ANOVA (D) and Student's *t*-test (A); *$P<0.05$, **$P<0.01$, ***$P<0.001$.

Figure 5. WUS activates *SPL4*. (*A*) Relative expression of mature miRNA and *SPL* genes in SD-grown Col-0 and *clv3-10* apices. n=4 (*B*) Representative pictures of *SPL3-5* single, double and triple CRISPR/*Cas9* deletion mutants in comparison to Col-0 grown in SD. (*C*) Flowering time of deletion mutants displayed in (*B*) provided as days to bolting and (*D*) total leaf numbers. n=25. (*E*) Relative expression of mature miRNA and *SPL* genes in SD-grown Ler-0 and *wus-7* apices (*F*) Overview of *SPL4* 5' regulatory region with putative *SPL4^WUS* sites (grey, yellow and white circles) and position of ChIP-PCR amplicons (grey, blue and yellow boxes) corresponding to the results shown in (G, H). Boxes marked with I, II, III and IV (1&2) indicate *5' SPL4* regions with in total five core *SPL4^WUS* sites – I: -1073 – -1068 bp, II: -880 – -875 bp, III: -697 – -691, IV1: -259 – -252 and IV2: -214 – -209 bp. Sequence location and lengths used in (*G*) indicated with #1-3. (*G*) Protoplast transactivation assay showing activation of the reporter (LUC) when coupled to the regions indicated in (*F*), relative to REN activity. c indicates vector control. n=3. (*H*) Enrichment of regions I, II, III and IV1&2 as indicated in (*F*) measured by ChIP-PCR relative to the input in *clv3-7* and *clv3-10* apices. PB – post binding fraction. n=3. Error bars denote s.d.; significance calculated based on one-way ANOVA (A, G) and Student's *t*-test (D); *P<0.05, **P<0.01, ***P<0.001.

Figure 6. SPL4 negatively feeds back to *WUS*. (*A*,*B*) *WUS* and *CLV3* expression by RNA *in situ* hybridization on (*A*) longitudinal sections through Col-0, *35S:amiRTPS1, 35S:TPS1* and *CLV3:TPS1* and (*B*) Col-0, and *35S:TPPJ* apices of LD-grown plants. (*C*) Relative expression of *WUS* and *CLV3* in SD-grown Col-0, *35S:TPPJ* and *ML1:amiRTPPJ* V1 apices. n=4 (*D*) RNA *in situ* hybridization on longitudinal sections through Col-0, and *35S:amiRTPPJ* apices. n=3. (*E*) Overview of *WUS* 5' regulatory region with putative *WUS^SPL* site corresponding to the results shown in (F,G,H). Grey box indicates *5' WUS* region with the native and mutated core *WUS^SPL* site -1067 – -1063 bp (yellow and blue circles, respectively). Sequence location and lengths used in (F,G) indicated as lines below. (*F,G*) Protoplast transactivation assay showing repression of the reporter (LUC) when coupled to the *5'WUS* region indicated in (E), (*F*) relative to REN activity of the native *5'WUS*, and (*G*) the comparison of the native and the mutated *WUS^SPL* site (yellow and blue circles, respectively), n=3. (*H*) EMSA for SPL4 binding to the *WUS^SPL* site (E). Shifted band in the presence of SPL4 protein (open arrow head), non-shifted fraction (closed arrow head). Please note the shift in the presence of the mutated competitor (+*), indicating specificity of binding. Error bars denote s.d.; significance calculated based on one-way ANOVA; **P<0.01, ***P<0.001. Scale bars: 25µm. Star indicates SAM summit.

Supplementary Information for

Sugar Signaling Induces Dynamic Changes During Meristem Development to Regulate Flowering in Arabidopsis

Magdalena Musialak-Lange[1], Katharina Fiddeke[1,#], Annika Franke[1,#], Friedrich Kragler[1], Christin Abel[1], and Vanessa Wahl[1,*]

Affiliations: [1]Max Planck Institute of Molecular Plant Physiology, Department of Metabolic Networks, Am Mühlenberg 1, Potsdam, 14476, Germany. [#]K.F. and A.F. contributed equally

[*]**Correspondence**: Vanessa Wahl
Email: vanessa.wahl@mpimp-golm.mpg.de

This PDF file includes:

> Figures S1 to S25
> Tables S1 to S8
> SI References

Supplementary Information

Supplementary Figures

Col-0   35S:amiRTPS1 #2   35S:amiRTPS1 #5   35S:amiRTPS1 #6

Fig. S1. ***TPS1* expression at the shoot apical meristem of *35S:amiRTPS1* plants.** Downregulation of *TPS1* by overexpressing an artificial microRNA targeting *TPS1* (*35S:amiRTPS1*) in plants of three independent lines (#2, #5 and #6; (Wahl *et al*, 2013)) as analyzed by RNA *in situ* hybridization using a specific probe against *TPS1*. Pictures depict representative longitudinal middle sections through vegetative apices (8 DAG in LD) of Col-0 and *35S:amiRTPS1*, respectively. Line #6 was used for the experiments in this study. Star indicates SAM summit. Scale bar is 50 µm**.**

Fig. S2. *TPS1* **expression at the shoot apical meristem of** *35S:TPS1* **and** *CLV3:TPS1* **plants.** In comparison to expression in Col-0 *TPS1* was generally found upregulated in plants overexpressing *TPS1* (*35S:TPS1*, #7) and the central zone when expressing it under the control of the *CLAVATA3* (*CLV3*) promoter (*CLV3:TPS1*, #15) as analyzed by RNA *in situ* hybridization using a specific probe against *TPS1*. Pictures depict representative longitudinal middle sections through vegetative apices (8 DAG in LD) of Col-0, *35S:TPS1*, and *CLV3:TPS1*, respectively. Star indicates SAM summit. Scale bars are 50 µm.

Fig. S3. Cell cycle gene expression at the shoot apical meristem of *35S:TPS1, CLV3:TPS1* and *35S:amiRTPS1* plants. RNA *in situ* hybridization using specific probes for *HISTONE H4* (*H4*), *CYCLIN D3;1* (*CYCD3;1*) and *CYCLIN-DEPENDENT KINASE 2;1* (*CDKB2;1*) to illustrate expression in longitudinal middle sections through the SAM of Col-0, *35S:TPS1* (#7), *CLV3:TPS1* (#15) and *35S:amiRTPS1* (#6). Star indicates SAM summit. Scale bars are 50 μm.

Fig. S4. Metrics at the SAM before and at the floral transition. Cell numbers as recorded at 8 (vegetative SAM) and 10 days (transition SAM) after germination (DAG). Significance calculated based on one-way Anova; *P<0.05.

Fig. S5. Shoot apical meristem area of SD-grown and LD-induced plants. Vegetative and inflorescence SAM areas (light grey), and transition SAM area (dark grey) of wild-type plants grown in SD for 30 days and transferred to LD for 3, 5, 7 and 17 days (DAS). Error bars denote s.d.; ***P<0.001 in relation to 0 days after transfer to LD (Student's *t*-test).

Fig. S6. Analysis of the cytokinin signaling reporter TCSn:GFP. Representative confocal laser scanning microscope images of SAMs of (*A*) 6 DAG, (*B*) 8 DAG, (*C*) 10 DAG, and (*D*) 12 DAG of TCSn:GFP cytokinin reporter line (green (Zurcher *et al*, 2013); blue – autofluorescence of chloroplasts). Please note the absence of cytokinin signal in the L1 (between dashed lines) and L2 (cell layers below) in the meristem proper. Star indicates SAM summit, scale bars are 25μm.

Fig. S7. Expression of Arabidopsis *TPP* genes at the shoot apical meristem detected by RNA *in situ* hybridization. The individual panels contain: a longitudinal middle section and a cross section through a vegetative Col-0 apex and a longitudinal middle section through an inflorescence apex. Representative pictures of *TPPC* and *TPPD* were omitted from this figure, since their specific probes gave no signal, indicating that transcripts of *TPPC* and *TPPD* were below detection limit in the tissues and/or stages analyzed. Star indicates SAM summit. Scale bars are 100 μm.

Fig. S8. AtGenExpress expression profiles of the Arabidopsis *TPP* genes. Values for Col-0 vegetative, transition and inflorescence apices (grey scale) and *clv3-7* inflorescence apices (orange) extracted from data deposited for (Schmid *et al*, 2005). All data points were mean-normalized.

Fig. S9. Expression of meristem maintenance genes and *TPPJ* in the *clv3-7* and *clv3-10* mutant SAM. RT-qPCR results of *TPPJ*, *WUS* and *CLV3* expression in SD-grown *clv3-7* and *clv3-10* SAMs, harvested 40 days after germination. Please note that the graph for *TPPJ* is a biological repetition of the graph in Figure 2C. Please also note that *clv3-10* is a TALEN generated mutant with a five nucleotide deletion in the last exon, which renders the transcript inactive to produce a functional protein (Forner *et al*, 2015). This is the cause for increased expression of *CLV3* downstream of increased *WUS* in the *clv3-10* mutant background. Error bars denote s.d.; *$P<0.05$, **$P<0.01$, ***$P<0.001$ (one-way ANOVA).

Fig. S10. *TPPJ* Expression in *WUS* transgenic lines. *(A)* RT-qPCR results of *TPPJ* expression in SD-grown *wus*-7 as compared to wild-type Ler-0 SAMs, harvested 30 days after germination. Error bars denote s.d.; **$P<0.01$, (one-way ANOVA). (*B*) *TPPJ* expression is increased upon WUS induction as shown by a WUS regulation score (WRS). According to Busch and colleagues, the WRS quantifies gene expression in response to a modulation in WUS activity normalized to the average response of all genes. Values extracted form a data set, deposited as a supplemental material of a publication by (Busch *et al*, 2010).

Fig. S11. ChIP-PCR control experiments for Figure 3C-E. (*A*) Overview of *5'TPPJ* intergenic sequence with all investigated putative $TPPJ^{WUS}$ sites (grey circles, black boxes), position of ChIP-PCR amplicons corresponding to the results shown in (B-D). Black framed boxes marked with I, II (1-4), and III (1-3) indicate *5' TPPJ* regions directly bound by WUS – I: -2795 – -2789 bp, II: -2073 – -1830 bp, and III: -652 – -564 bp, as presented in the Figure 3. Grey framed boxes (II5, III3-4, IV1-3, V1, VI1-2) represent *5'TPPJ* regions not directly bound by WUS. (*B-D*) Relative expression of investigated regions (I-III) containing $TPPJ^{WUS}$ elements (I, II1-4, III1-2) in Col-0, *clv3-7* and *clv3-10* (*B*) shoot apex samples without WUS antibody, and (*C*) leaf samples without WUS were used as specificity controls. Please note the amplification in the input and post-binding fractions (PB) and the absence of amplification in the ChIP samples indicating specificity of the antibody used. (*D*) Control overview of all investigated putative $TPPJ^{WUS}$ sites in input and post binding fraction samples. Error bars denote s.d.

Fig. S12. ChIP-PCR on known WUS targets using a specific WUS antibody. Relative to the input enrichment of known *CLV3*[WUS] (Yadav *et al*, 2011) and *ARABIDOPSIS RESPONSE REGULATOR 7* (*ARR7*[WUS]) (Leibfried *et al*, 2005) in *clv3-7* and *clv3-10* as measured by ChIP-PCR. As a negative control a *5'TPPJ* region was used that did not show any WUS binding (unbound region). Error bars denote s.d.

Fig. S13. ChIP-PCR on G-box like WUS binding sites. Relative expression of two G-box like WUS binding regions (CACGTG) (Sloan *et al.*, 2020) present in the *5'TPPJ* intergenic region in input, post binding and ChIP samples of Col-0, *clv3-7* and *clv3-10*. G-box like 1 and G-box like 2 correspond to the regions VI1 and VI2 of Fig. S11A. Please note the amplification in the input and post-binding fractions and the absence of amplification in the ChIP samples. Error bars denote s.d.

Fig. S14. EMSA control experiments. EMSA for WUS binding to the indicated regions (I-III) in Figure 3A as a control for Figure 3F. Shifted band in the presence of WUS protein (open arrow head), non-shifted fraction (closed arrow head). Please note the shifted band in the presence of the mutated competitor (x*, Table S8) indicating specificity of the binding.

Fig. S15. Chracterization of the GABI_215C08 line. *(A)* Schematic illustration of the T-DNA insertion in the 6[th] exon of *TPPJ* including positions of oligos used for genotyping. *(B)* Expression of *TPPs* in LD-grown Col-0 and *tppj* rosettes, harvested without roots at 10 days after germination as measured by RT-qPCR. In this experiment *TPPI* was not detectable. Error bars denote s.d.; *P<0.01; ***P<0.001 (one-way ANOVA).

Fig. S16. Characterization of the *35S:amiRTPPJ* lines. *(A)* Downregulation of *TPPJ* expression in *35S:amiRTPPJ* V1 and V2 rosettes shown by RT-qPCR, relative to Col-0. *(B)* Expression of all *TPPs* in LD-grown Col-0 and *35S:amiRTPPJ* V1 rosettes, harvested without roots at 10 days after germination by RT-qPCR. In this experiment *TPPI* was not detectable. Error bars denote s.d.; **P<0.01 (one-way ANOVA).

Fig. S17. Earlier flowering of *35S:amiRTPPJ* and *tppj* lines. Flowering time of *35S:amiRTPPJ* V1 and V2 and *tppj* T-DNA insertion line as shown by the total leaf numbers and days to bolting, relative to the wild type in LD and in SD. V1 and V2 indicate two independent versions of artificial microRNAs designed to target *TPPJ* transcript. Representative pictures of *35S:amiRTPPJ* V1 line with stronger reduction in *TPPJ* expression, *35S:amiRTPPJ V2* and *tppj* in comparison to Col-0 are depicted. Scale bars are 1 cm. Error bars denote s.d.; significance calculated based on one way Student's *t*-test; *P<0.05, **P<0.01, ***P<0.001.

Fig. S18. ***ML1* expression throughout development**. RNA *in situ* hybridization using a specific probe against *ML1* on longitudinal sections through Col-0 apices from plants grown in LD. 6, 8 DAG depict vegetative SAMs, 10 DAG the transition to flowering, and 12, 16 DAG inflorescence SAMs. Star indicate SAM summit. Scale bars are 100 μm.

Fig. S19. *TPPJ* **expression at the SAM in the background of** *ML1:amiRTPPJ* **plants**. RNA *in situ* hybridization with a specific probe against *TPPJ* did not detect *TPPJ* transcript in L1 (arrow head) on longitudinal sections through SAMs of *ML1:amiRTPPJ* plants (V1). Pictures depict two biological replicates of representative inflorescence SAMs of *ML1:amiRTPPJ* in comparison to Col-0. Star indicate SAM summit. Scale bars are 50 µm.

Fig. S20. Early flowering phenotype of *ML1:amiRTPPJ* lines. Representative pictures of (*A*) the stronger *ML1:amiRTPPJ V1* line and (*B*) *ML1:amiRTPPJ V2* in comparison to Col-0 grown in SD conditions. Scale bars are 1 cm.

Fig. S21. Wild-type like flowering phenotype when *ML1:amiRTPPJ* is introgressed into *clv3-10*. Representative pictures of *ML1:amiRTPPJ V1;clv3-10* next to the parental lines and wild type (Col-0) in LD (upper panel) and SD (lower panel) indicating a wild-type like flowering phenotype. Open and closed arrowheads indicate fasciated stem and enormous SAM, respectively. Note the reduced phenotype of the crosses as compared to the *clv3-10* mutant. Scale bars are 1 cm.

Fig. S22. Expression of *SPL15* at the SAM. *SPL15* does not significantly change at the SAM of (*A*) *ML1:amiRTPPJ* and *35S:TPPJ,* (*B*) *clv3-10*, and *(C) wus-7* compared to its expression in the corresponding wild type. Statistical significance was calculated using one-way ANOVA. Error bars denote s.d.

Fig. S23. *SPL3, SPL4,* and *SPL5* mutant lines. Schematic illustration of the deletions in the *SPL3* (At2g33810)*, SPL4* (At1g53160)*,* and *SPL5* (At3g15270) locus generated by CRISPR/*Cas9*. Positions of the PAM sequence are given relative to the transcription start site (ATG). Sequences of guided RNAs with PAM sites (in red) are given in white boxes. In grey the sequences flanking the deletion are presented. E – exon.

Fig. S24. Potential *SPL^WUS* sites in the 5' intergenic regions of *SPL3*, *SPL4*, *SPL5*, *SPL9*, and *SPL15*. Overview of analyses of *SPL3, SPL4, SPL5, SPL9* and *SPL15* 5' intergenic sequences for the presence of putative core binding sites (white circles), extended binding sites (yellow circles), TGA boxes (green circles) and G-box-like sites (brown circles). Red boxes indicate *5'UTR*s. Black boxes represent putative transcripts annotated in the intergenic regions upstream the ATG of *SPL3* and *SPL5.* The total sequence lengths are given in base pairs (bp).

Fig. S25. ChIP-PCR control experiments. (*A*) Overview of 5' *SPL4* intergenic sequence with all investigated putative *SPL4^WUS^* sites (white, yellow and brown circles), position of ChIP-PCR amplicons corresponding to the results shown in (B and C). Black framed boxes marked with I, II, III, and VI (1&2) indicate *5'SPL4* regions directly bound by WUS – I: -1073 – -1068 bp, II: -880 – -875 bp, III: -697 – -691 bp, and IV: -259 – -209, as presented in Figure 5H. (*B,C*) Relative expression of investigated regions (I-IV) containing *SPL4^WUS^* elements in (*B*) Col-0, *clv3-7* and *clv3-10* shoot apex samples without WUS antibody. (*C*) Control overview of all investigated putative *SPL4^WUS^* sites in input and post binding fraction samples. Error bars denote s.d.

Fig. S26. *WUS* and *CLV3* **expression in apices of SD-grown SPL double and triple CRISPR/*Cas9* deletion mutants**. Expression provided as (*A,C*) RNA *in situ* hybridization and (*B,D*) RT-qPCR. *WUS* (A,B) and *CLV3* (C,D). n=3. Error bars denote s.d.; significance calculated based on one-way ANOVA; *P<0.05. Scale bars: 25μm. Star indicates SAM summit.

Supplementary Tables S1 to S7

Table S1. Flowering times of Col-0, mutants and transgenic plants. DTF, days to flowering (equals bolting time, see before); RLN, rosette leaf number; CLN, cauline leaf number; TLN, total leaf number; n, number of individual plants; p-value of student's *t*-test for DTF with Co-0; LD, long days; SD, short days; NA, not analyzed (leaf number above 200).

	DTF	DTF range	RLN	CLN	TLN	TLN range	n	p-value
Presented in Figure 4 of the main text:								
Experiment 1 (SD)								
Col-0	71.6	69-75	66.4	11.5	77.9	76-82	20	-
ML1:amiRTPPJ V1 #1	55.0	50-58	49.4	12.0	61.3	58-63	20	3.29E-22
ML1:amiRTPPJ V2 #5	64.4	61-66	52.8	9.2	62.0	60-64	20	3.07E-14
Experiment 3 (SD)[*]								
Col-0	82.2	79-86	68.6	11.9	80.5	76-86	22	-
ML1:amiRTPPJ V1 #1	65.3	61-69	56.7	9.9	66.5	63-72	22	1.08E-25
clv3-10	95.8	95-101	NA	NA	NA	NA	26	2.17E-25
clv3-10;ML1:amiRTPPJ V1 #1_1	87.3	84-92	NA	NA	NA	NA	24	4.39E-09
clv3-10;ML1:amiRTPPJ V1 #1_3	85.8	82-91	NA	NA	NA	NA	23	4.05E-05
Only presented as supplemental material:								
Experiment 2 (LD)								
Col-0	22.8	22-25	12.1	3.2	15.3	14-16	20	-
ML1:amiRTPPJ V1 #1	20.0	17-21	9.0	3.1	12.1	10-13	20	4.33E-10
ML1:amiRTPPJ V2 #5	21.0	20-22	10.3	2.9	13.2	10-14	20	9.95E-06
Experiment 3 (SD)[*]								
clv3-7	87.5	85-91	NA	NA	NA	NA	25	1.38E-09
clv3-7;ML1:amiRTPPJ V1 #1_2	74.6	70-78	NA	NA	NA	NA	24	5.50E-22
clv3-7;ML1:amiRTPPJ V1 #1_12	76.9	73-80	NA	NA	NA	NA	24	1.47E-19
Experiment 4 (LD)								
Col-0	26.5	25-29	20.6	5,1	25,7	24-29	31	-
35S:amiRTPPJ V1 #1	24.3	22-26	17.8	4.6	22.3	21-25	32	1.16E-08
35S:amiRTPPJ V1 #2	23.0	21-26	16.8	4.3	21.2	19-23	32	2.24E-11
tppj	25.8	24-28	19.8	4.8	24.6	22-27	35	0.04
Experiment 5 (SD)								
Col-0	65.1	61-68	63.7	10.1	73.7	70-76	30	-
35S:amiRTPPJ V1 #1	60.4	57-62	58.4	9.7	68,1	63-71	30	6.63E-12
35S:amiRTPPJ V1 #2	58.3	56-60	57.8	9.5	67.2	66-69	30	1.47E-17
tppj	59.3	57-62	58.4	9.4	67.8	65-72	30	6.15E-15
Experiment 6 (SD)								
Col-0	72.4	69-75	67.9	9	77.2	74-81	26	-
spl3	73.4	71-75	68.3	10	78.0	75-81	23	0.15
spl4	74.0	71-77	69.2	14	83.0	80-86	26	5.00E-05
spl5	74.0	71-77	68.0	10	77.8	74-82	24	0.45
spl34	77.9	76-80	74.1	17	90.9	87-94	24	5.04E-09
spl35	73.2	71-75	68.4	10	77.9	75-81	24	0.02
spl45	76.8	74-78	73.0	16	76.8	86-93	24	1.25E-11
spl345	84.1	82-87	79.6	19	99.0	94-112	27	2.79E-26

Please note that data marked with an asterisk (*) were obtained from the same experiment.

Table S2. Oligonucleotides for constructs used for the generation of transgenic lines and for transactivation assays.

Gene (AGI)	Oligo	Construct	Sequence (5'→ 3')
TPPJ At5g65140	P-0188 P-0189 P-0190 P-0191	pVW501 *amiRTPPJ V2*	gaTATAATATATCCCCAAGTCGAtctctcttttgtattcc gaTCGACTTGGGGATATATTATAtcaaagagaatcaatga gaTCAACTTGGGGATTTATTATTtcacaggtcgtgatatg gaAATAATAAATCCCCAAGTTGAtctacatatatattcct
TPPJ At5g65140	P-0192 P-0193 P-0194 P-0195	pVW502 *amiRTPPJ V2*	gaTCTAAATGTACTTCACTGCCCtctctcttttgtattcc gaGGGCAGTGAAGTACATTTAGAtcaaagagaatcaatga gaGGACAGTGAAGTAGATTTAGTtcacaggtcgtgatatg gaACTAAATCTACTTCACTGTCCtctacatatatattcct
TPPJ At5g65140	G-8641 G-8642	pVW279 *35S:TPPJ*	ATGGGTTATGATAATGTATGTG TTAAACGTTTGGCTTCTTGACAC
5' TPPJ At5g65140	P-1620 P-1659	pMML049 *5' TPPJ (#1)*	aagggtaccaGAAATACCCTCAAAGCAAATTTGA CCGTGGCGCGCCCGGTCTTggcgtct
	P-1621 P-1659	pMML050 *5' TPPJ (#2)*	attggtaccacaGAGTCCACAAGTTTAACAAACAACAGA CCGTGGCGCGCCCGGTCTTggcgtct
	P-1654 P-1659	pMML051 *5' TPPJ (#3)*	cttggtaccTTGTCATCATTAAGATGGGATGACA CCGTGGCGCGCCCGGTCTTggcgtct
	P-1655 P-1659	pMML052 *5' TPPJ (#4)*	tatggtaccAACGTGGGTGATCATCTCTCG CCGTGGCGCGCCCGGTCTTggcgtct
	P-1657 P-1659	pMML054 *5' TPPJ (#5)*	ctaggtaccGTGATTATTGTCAAGGCACCTTAT CCGTGGCGCGCCCGGTCTTggcgtct
	P-1658 P-1659	pMML055 *5' TPPJ (#6)*	ttaggtaccCCAAACGAAACAAAAACG CCGTGGCGCGCCCGGTCTTggcgtct
5' SPL4 At1g53160	P-2040 P-0243	pMML070 *5'SPL4 (#1)*	ATATGGTACCGCATAAATCCAGATGTTCCTCC ATATACTAGTCCTTGTCCTTGTGATCTCTTACC
	P-2041 P-2043	pMML071 *5'SPL4 (#2)*	ATATGGTACCGCTACGGACAAGGTCCTACG ATATACTAGTCCTTGTCCTTGTGATCTCTTACC
	P-2042 P-2043	pMML072 *5'SPL4 (#3)*	ATATGGTACCGGCAGCTTTTGGCTTTGG ATATACTAGTCCTTGTCCTTGTGATCTCTTACC
5' WUS A**t**2g17950	P-2040 P-0243	pMML067 *5'WUS*	ATATGGTACCGCATAAATCCAGATGTTCCTCC ATATACTAGTCCTTGTCCTTGTGATCTCTTACC
	P-2041 P-2043	pMML088 *5'WUS* mutated*	CATGTTCATAA**TTTT**ACCTGTCTTCACAC GTGTGAAGACAGGT**AAAA**TTATGAACATG
SPL3 At2g33810	P-1334 P-1336	pMML040 *CRISPR Cas9 SPL3*	accaggtctcaattgAATAGAGCTGGGTCCTCGTCgtttttagagctagaaatagcaag tggtggtctctaaacAGCGGAGTTGCAGGAGACGCcaatctcttagtcgactctacc
SPL4 At1g53160	P-1340 P-1341	pMMl043 *CRISPR Cas9 SPL4*	accaggtctcaattgGAAAAAGAAGTCTTACCTTGgttttagagctagaaatagcaag tggtggtctctaaacTTGATCTTTCTTGATTCTGCcaatctcttagtcgactctacc
SPL5 At3g15270	P-1343 P-1345	pMML027 *CRISPR Cas9 SPL4*	accaggtctcaattgTTGTATTCGTGGGAAGATGAgttttagagctagaaatagcaag tggtggtctctaaacGATAGATATTCATTACAGGAcaatctcttagtcgactctacc

* Mutagenesis PCR was performed on the pMML067 plasmid. The resulting mutated promoter was sequenced after.

Table S3. Oligonucleotides used for genotyping.

Transgenic line	Oligo	Sequence (5'→ 3')	Position
VW507 *ML1:amiRTPPJ* (V1)	P-0110 P-0189	AGTAGTATGAATTTTCTCTCAC *gaTCGACTTGGGGATATATTATAtcaaagagaatcaatga*	*ML1 promoter* *miRNA stemloop*
VW508 *ML1:amiRTPPJ* (V2)	P-0110 P-0193	AGTAGTATGAATTTTCTCTCAC *gaGGGCAGTGAAGTACATTTAGAtcaaagagaatcaatga*	*ML1 promoter* *miRNA stemloop*
VW504 *35S:amiRTPPJ* (V1)	P-0107 P-0189	AACCTCCTCGGATTCCATTGCC *gaTCGACTTGGGGATATATTATAtcaaagagaatcaatga*	*CaMV 35S promoter* *miRNA stemloop*
VW505 *35S:amiRTPPJ* (V2)	P-0107 P-0193	AACCTCCTCGGATTCCATTGCC *gaGGGCAGTGAAGTACATTTAGAtcaaagagaatcaatga*	*CaMV 35S promoter* *miRNA stemloop*
VW161 *35S:TPPJ*	P-0107 P-0121	AACCTCCTCGGATTCCATTGCC ATTGTGTTCTTCTTCGGAAGAG	*CaMV 35S promoter* *TPPJ CDS*
VW161 *35S:TPS1*	P-0107 P-0120	AACCTCCTCGGATTCCATTGCC ATTATAAGGCCTAACTTCTTGC	*CaMV 35S promoter* *TPS1 CDS*
tppj GABI_215C08	P-0208 P-1300 P-1302	TGACCAGCCGCGCTGAATGTG GATGCATCGACAAGGTTTATAGCTTTG GTTACAGTTTTCAAATCCGAGTGACTCTA	*GABI Kat LB* *TPPJ forward* *TPPJ reverse*
clv3-10*	P-1386 P-1387	TGATGCTTCTGATCTCACTCAAGC CCATTTGCTCCAACCCATTC	*CLV3 forward* *CLV3 reverse*
clv3-7	P-1519 P-1518 P-1387	CCTGAGGTTGGGCAGAAACA TCAAGCTCATGCTCACGTTCA CCATTTGCTCCAACCCATTC	*En-1 LB* *CLV3 forward* *CLV3 reverse*
spl3	P-1476 P-1113	GTGGTGTCATAGCAT TGGGAATCAAACGTGACTAGG	*5'SPL3* *3'SPL3*
spl4	P-1498 P-1499	GCTTAACCTATTAGAGTTTC TCGTGGCTCTGAAACTTCAG	*5'SPL4* *3'SPL4*
spl5	P-1497 P-1475	GGAAGCAAGCACATCACGTA CGCAAATAAGAATCAAGTCGAGa	*5'SPL4* *3'SPL4*
Cas9	P-1478 P-1480	CGACCCTCATCCACCAGTCGATTAC CCCAGGCTTTACACTTTATGCTTCC	*Cas9 cassette*

*Note that *clv3-10* was genotyped based on sequence analyses.

Table S4. **Oligonucleotides used for RT-qPCR**.

Gene	Oligo	Sequence (5'→3')
TPPJ	P-1300	GATGCATCGACAAGGTTTATAGCTTTG
At5g65140	P-1301	TGAGAAACCTTTAGTTGGGCCTTT
TPPA	P-1493	GATGATCGATAAGGTGCTTTGTTC
At5g51460	P-1494	GCGGTAATGCACAGAGATGC
TPPB	P-1489	CAAGGTAGAAAGGTACTTGAAATCCG
At1g78090	P-1490	ACACCGGCACAACATCATCC
TPPC	P-1487	CAAGGCAGAAAGGTCTTAGAACTCC
At1g22210	P-1488	TCCCTAATGAGTTTAGCAAAAATTCG
TPPD	P-1483	GATGCCGTGATAAGGTTCGTAGAT
At1g35910	P-1484	TGGAAGGTCCTTTGATGTCCA
TPPE	P-1485	CAAGGAAGAAAAGTATTGGAGATTCG
At2g22190	P-1486	ATAGGAAAAACATCGGTACAATTATCA
TPPF	P-1481	ACGCGGGTAGTCATGGAATG
At4g12430	P-1482	CACTTCCTTACCCTGTTGGTCAGT
TPPG	P-1303	GTAAATCCAAATGGATCCCCTGAAG
At4g22590	P-1304	TTACCTCCTCACCCTGTTGGTC
TPPH	P-1495	AGAGTCACTTGGATACGCTAATTGTAC
At4g39770	P-1496	GCCAAGACCTTGTCTTCGTTC
TPPI	P-1482	GCCTTCGCCTCTGTCTCGTA
At5g10100	P-1483	GATGATCGATAAGGTGCTTTGTTC
SPL3	P-1617	GTTTCATGCCAAAGCTCCTC
At2g33810	P-1618	TTTTCCGCCTTCTCTCGTT
SPL4	P-0546	AATGGTCAGGTGGTGATGCAG
At1g53160	P-0547	GCATAGGAAGTGTCATCTCTACCCTT
SPL5	P-0548	CAGCAGGTTTCATGAGCTACCAG
At3g15270	P-0549	CAAAACTGTCACCAGAGATCTTCCTC
SPL9	P-1307	CTTCGCTTTACGAAAATGGTGATG
At2g42200	P-1308	ACTGGCCGCCTCATCACTCT
SPL15	P-1309	CATCTCTTTACGGAAACCCCAATG
At3g57920	P-1310	GCCGCTGCATCACTGATCTT
mir156	P-0342	GCGGCGGTGACAGAAGAGAGT
miRNA universal	P-0343	GTGCAGGGTCCGAGGT
CLV3	P-1386	TGATGCTTCTGATCTCACTCAAGC
At2g27250	P-1387	CCATTTGCTCCAACCCATTC
WUS	P-1382	CCCAGCTTCAATAACGGGAAT
At2g17950	P-1383	ACAGTCTTGTTCCATAGATCCATAGAC
UBQ10	P-0348	CACACTCCACTTGGTCTTGCGT
At4g05320	P-0349	TGGTCTTTCCGGTGAGAGTCTTCA
TUB2	P-0344	GAGCCTTACAACGCTACTCTGTCTGTC
At5g62690	P-0345	ACACCAGACATAGTAGCAGAAATCAAG
SAND	P-0346	AACTCTATGCAGCATTTGATCCACT
At2g28390	P-0347	TGATTGCATATCTTTATCGCCATC
PDF2	P-0350	TAACGTGGCCAAAATGATGC
At1g13320	P-0351	GTTCTCCACAACCGCTTGGT
GAPDH *	P-1125	TCTCGATCTCAATTTCGCAAAA
	P-1126	CGAAACCGTTGATTCCGATTC
At1g13440	P-1127	TTGGTGACAACAGGTCAAGCA
	P-1128	AAACTTGTCGCTCAATGCAATC
miR156 stem-loop **	P-1010	GTCGTATCCAGTGCAGGGTCCGAGGTATTCGCACTGGATACGACG

* Primers used for cDNA quality check according to (Czechowski *et al*, 2004).

** Please note that this oligo was used for cDNA synthesis according to (Varkonyi-Gasic *et al*, 2007).

Table S5. **Oligonucleotides used for ChIP-PCR**.

Gene	Oligo	Region	Sequence (5'→3')
5' TPPJ: At5g65140	P-1624 P-1625	**I**: -3073 bp -2941 bp	GAAATACCCTCAAAGCAAATTTGA AGCTTCTAAAATTCATCAAAACCAA
	P-1628 P-1783	**II (1)**: -2150 bp -1924 bp	GAGTCCACAAGTTTAACAAACAACAGA TCCCATCTTAATGATGACAAACA
	P-1626 P-1782	**II (2)**: -1943 bp -1847 bp	TTGTCATCATTAAGATGGGATGACA TACTGATATATGTTTGCAAAGTTT
	P-1784 P-1627	**II (3+4)**: -1887 bp -1751 bp	TGAATGGAGATAATATAAAACTTTGC TCGAAGACGCCCCTTGTC
	P-1629 P-1630	**II (5)** -1818 bp – 1617 bp	GCCGCTGCTGGTTTTGAG TGTCATTGCATGCATGTGTG
	P-1628 P-1782	**II**: -2150 bp -1847 bp	GAGTCCACAAGTTTAACAAACAACAGA TACTGATATATGTTTGCAAAGTTT
	P-1636 P-1785	**III (1)**: -697 bp -604 bp	CGTGTTCCTATCCATGTTCCT GTATAATGATAAATTTGTAGTAATACC
	P-1786 P-1637	**III (2+3)**: -630 bp -521 bp	GGTATTACTACAAATTTATCATTATAC CCACTTCAACGTTATATCAACCG
	P-1640 P-1641	**III (4)** -826 bp -663 bp	TGTGATTATTGTCAAGGCACCTTAT TTTCATGATTGTATAGGAACATGG
	P-1638 P-1639	**III (5)** -561 bp – 381 bp	CCCCAAACGAAACAAAAACG CCCCGAGCCAAAGGAAAA
	P-1636 P-1637	**III**: -697 bp -521 bp	CGTGTTCCTATCCATGTTCCT CCACTTCAACGTTATATCAACCG
	P-1631 P-1632	**IV (1)**: -1353bp -1167 bp	AACGTGGGTGATCATCTCTCG CGTTCAATAGTTGGCACTAAACCA
	P-1633 P-1635	**IV (2)**: -1184 bp - 891bp	AGTGCCAACTATTGAACGATTG CGATGCATGCAGCCACAC
	P-1633 P-1634	**IV (3)**: -1185 bp – 968 bp	AGTGCCAACTATTGAACGATTG CCATCACGCTCTCGAACTTTCT
	P-1642 P-1643	**V**: -94 bp – 44 bp	TCCTCCTCCTTCCTAGTTTCTCG ATCCCGGTCTTGGCGTCT
	P-2014 P-2015	**VI (1)**: -1513 bp – 1387 bp	CCAAAAGGGGACGCGAG TCGATTCCGGTTTCATCGTT
	P-2016 P-2017	**VI (2)**: -1513 bp – 1387bp	CGAATACGACGTAGACGAATAAGACA AGCCATGGGACCGAGGTT
5' SPL4: At1g53160	P-2109 P-2110	**I**: -1166 bp -1015 bp	CTTATCTGTTTCCTGCTTTGTTGG CAGTGTGTTGTGTGGAGAAGCA
	P-2111 P-2112	**II**: -1005 bp -821 bp	AGGAAGGTAAAAGTGAGAGGCTTTT GGAGGAATTACAGATTGAAAGATGG
	P-2113 P-2114	**III**: -765 bp -593 bp	GAAAGAAATCAAAAGGGTTAGATCGT TTTGGTAGACTCTTATTGGAAATGCA
	P-2115 P-2116	**IV (1&2)**: -425 bp -169 bp	CCTCTTAAAGGCAACCAATGATATC GTGCATGATTACGTTGATGTGC
5' UBQ10: At4g05320	P-1889 P-1890	-582 bp – 306 bp	TCGTGTTTATCAACTCAAAGCACAA TGATCACGGTAGAGAGAATTGAGAGA

Abbreviations: I, **II**, **III (*TPPJ*)** and **I, II, III; IV (*SPL4*)** regions positive in ChIP-PCR, other regions – negative

Table S6. **Oligonucleotides used to prepare constructs for probe synthesis**.

Gene	Construct ID	Oligo	Sequence (5'→3')
WUS: At2g17950	pVW310	P-0148 P-0149	ATGGAGCCGCCACAGCATCAGCATC CTAGTTCAGACGTAGCTCAAGAGAAG
CLV3: At2g27250	pJL102	P-0150 P-0151	TCACTCAGTCACTTTCTCTC GAAAATCATGAGATATAATAG
ML1: At4g21750	pVW510	P-0152 P-0153	ATGTATCATCCAAACATGTTCG TTAGGCTCCGTCGCAGGCCAGAG
TPPA: At5g51460	pVW518	P-0214 G-13892	ATGGACATGAAATCTGGTCACTCGTC TTAACCCATTGATCTCTTCCATGTC
TPPB: At1g78090	pVW516	P-0211 P-0212	ATGACTAACCAGAATGTCATCGTTTC TCACTCTTCTCCCACTGTCTTCCTC
TPPC: At1g22210	pVW515	P-0210 G-13888	ATGAAGATTACGGATATTTCCGGAA TTATTCTCCAAGTGTTTGTTTCTTC
TPPD: At1g35910	pVW257	G-8814 G-8815	ATGACAAACCATAATGCCTTAATC CTATCTTCCTCTTAGTGACATTTG
TPPE: At2g22190	pVW258	G-8816 G-8817	ATGGTTAGATTCATCGAAGAAAATATTAC TCATGCCCCACACCTTGACTGTTTCC
TPPF: At4g12430	pVW227	G-8641 G-8642	ATGGATTTAAACTCAAACCACAAATC TCAAAAACCAGTAGAATTCTTCTCC
TPPG: At4g22590	pVW514	G-8639 G-8640	ATGGATTTGAATATAAACAAGACG TCAAAAACTTGTTTTTGAACTTTC
TPPH: At4g39770	pVW519	P-0215 G-13894	ATGGTTAGATTCATAGAAGAAAACAC TCATGCTCCAGATCTCAATTGTTTCC
TPPI: At5g10100	pVW517	P-0216 P-0213	ATGTCAGCTAGTCAAAACATTGTCG TCACATTCTTGGCTGCATTTGTTTCC
TPPJ: At5g65140	pVW228	G-8641 G-8642	ATGGGTTATGATAATGTATGTG TTAAACGTTTGGCTTCTTGACAC
H4: At2g28740	pAK029	P-0285 P-0286	ATGCTGTTACTTACACTGAGCATG CCAGAAAACCCAGATCTCAAATTG
CYCD3;1: AT4G34160	pVW531	P-0250 P-0251	ATGGCGATTCGGAAGGAGGAAG TTATGGAGTGGCTACGATTGCCC
CDKB2;1 AT1G76540	pVW533	P-0258 P-0259	TGGACGAGGGAGTTATAGCAGTTTC TTAGAGAGAGGACTTTTCTGGCAG

Table S7. **EMSA oligonucleotides (probes) for *5'TPPJ* At5g65140**

Oligo	Modification	Region	Sequence (5'→3')
P-1777 P-1778	5' IR DEY6832	**I**:-2815 bp -2766 bp	ctaaaaattcgttttttttcttaatccggaagataaaaaggttataaaga tctttataaaccttttttatcttccggattaagaaaaaaaacgaatttttag
P-1779 P-1780	none – competitor	**I**:-2815 bp -2766 bp	ctaaaaattcgttttttttcttaatccggaagataaaaaggttataaaga tctttataaaccttttttatcttccggattaagaaaaaaaacgaatttttag
P-2186 P-2187	none – mutated competitor	**I**:-2815 bp -2766 bp	ctaaaaattcgtttttttttcttcctccggaagataaaaaggttataaaga tctttataaacctttttatcttccggaggaagaaaaaaaacgaatttttag
P-1829 P-1730	5' IR DEY682	**II (1)**:-2095 bp -2040 bp	ctattattttttcgctttcaaattaatacacatacaaattaacaaaagta tacttttgttaatttgtatgtgtattaatttgaaagcgaaaaaataatag
P-1841 P-1842	none – competitor	**II (1)**:-2095 bp -2040 bp	ctattattttttcgctttcaaattaatacacatacaaattaacaaaagta tacttttgttaatttgtatgtgtattaatttgaaagcgaaaaaataatag
P-2188 P-2189	none – mutated competitor	**II (1)**:-2095 bp -2040 bp	ctattattttttcgctttcaaattggtacacatacaaattaacaaaagta tacttttgttaatttgtatgtgtaccaatttgaaagcgaaaaaataatag
P-1831 P-1832	5' IR DEY682	**II (2)**:-1922 bp -1878 bp	gacataaaagtcacactgtcaaattaatttgaatatgaatggaga tctccattcatattcaaattaatttgacagtgtgactttttatgtc
P-1843 P-1844	none – competitor	**II (2)**:-1922 bp -1878 bp	gacataaaagtcacactgtcaaattaatttgaatatgaatggaga tctccattcatattcaaattaatttgacagtgtgactttttatgtc
P-2190 P-2191	none – mutated competitor	**II (2)**:-1922 bp -1878 bp	gacataaaagtcacactgtcaaattggtttggtatggatggaga tctccatccataccaaaccaatttgacagtgtgactttttatgtc
P-1833 P-1834	5' IR DEY682	**II (3+4)**:-1863 bp -1814 bp	gcaaacatatatcagtattaattaaatgattaatatgcatttgcagccgc gcggctgcaaatgcatattaatcatttaattaatactgatatatgtttgc
P-1845 P-1846	none – competitor	**II (3+4)**:-1863 bp -1814 bp	gcaaacatatatcagtattaattaaatgattaatatgcatttgcagccgc gcggctgcaaatgcatattaatcatttaattaatactgatatatgtttgc
P-2192 P-2193	none – mutated competitor	**II (3+4)**:-1863 bp -1814 bp	gcaaacatatatcagtattggtaaatgattggtatgcatttgcagccgc gcggctgcaaatgcataccaatcatttaaccaatactgatatatgtttgc
P-1835 P-1836	5' IR DEY682	**III (1)**:-674 bp -625 bp	acaatcatgaaagcttttggttttaatcttttttatttttattttttggtatt aataccaaaaataaaataaaaagattaaaaccaaaagctttcatgattgt
P-1847 p-1848	none – competitor	**III (1)**:-674 bp -625 bp	acaatcatgaaagcttttggttttaatcttttttatttttattttttggtatt aataccaaaaataaaataaaaagattaaaaccaaaagctttcatgattgt
P-2194 P-2195	none – mutated competitor	**III (1)**:-674 bp -625 bp	acaatcatgaaagcttttggtttggtctttttatttttattttttggtatt aataccaaaaataaaataaaaagaccaaaaccaaaagctttcatgattgt
P-1837 P-1838	5' IR DEY682	**III (2)**:-618 bp -569 bp	aatttatcattatacaactgtacataataatagtagtcatcactatacaa ttgtatagtgatgactactattattatgtacagttgtataatgataaatt
P-1849 p-1850	none – competitor	**III (2)**:-618 bp -569 bp	aatttatcattatacaactgtacataataatagtagtcatcactatacaa ttgtatagtgatgactactattattatgtacagttgtataatgataaatt
P-2196 P-2197	none – mutated competitor	**III (2)**:-618 bp -569 bp	aatttatcattatacaactgtacatggtggtagtagtcatcactatacaa ttgtatagtgatgactactaccaccatgtacagttgtataatgataaatt
P-1839 P-1840	5' IR DEY682	**III (3)**:-585 bp -536 bp	tagtcatcactatacaattaatttccccaaacgaaacaaaaacggttgat atcaaccgtttttgtttcgtttggggaaattaattgtatagtgatgacta
P-1851 P-1852	none – competitor	**III (3)**:-585 bp -536 bp	tagtcatcactatacaattaatttccccaaacgaaacaaaaacggttgat atcaaccgtttttgtttcgtttggggaaattaattgtatagtgatgacta
P-2198 P-2199	none – mutated competitor	**III (3)**:-585 bp -536 bp	tagtcatcactatacaattggtttccccaaacgaaacaaaaacggttgat atcaaccgtttttgtttcgtttggggaaaccaattgtatagtgatgacta

Abbreviations: I, **II**, **III** regions positive in ChIP-PCR.

Table S8. **EMSA oligonucleotides (probes) for 5'*WUS* At2g17950**.

Oligo	Modification	Region	Sequence (5'→3')
P-2172 P-2173	5' IR DEY6832	-1088 bp -1038 bp	gaatactcaacatgttcataagtacacctgtcttcacactcgtttcacac gtgtgaaacgagtgtgaagacaggtgtacttatgaacatgttgagtattc
P-2174 P-2175	none – competitor	-1088 bp -1038 bp	gaatactcaacatgttcataagtacacctgtcttcacactcgtttcacac gtgtgaaacgagtgtgaagacaggtgtacttatgaacatgttgagtattc
P-2180 P-2181	none – mutated competitor	-1088 bp -1038 bp	gaatactcaacatgttcataattttacctgtcttcacactcgtttcacac gtgtgaaacgagtgtgaagacaggtaaaattatgaacatgttgagtattc

SI References

Busch W, Miotk A, Ariel FD, Zhao Z, Forner J, Daum G, Suzaki T, Schuster C, Schultheiss SJ, Leibfried A *et al* (2010) Transcriptional control of a plant stem cell niche. *Dev Cell* 18: 849-861

Czechowski T, Bari RP, Stitt M, Scheible WR, Udvardi MK (2004) Real-time RT-PCR profiling of over 1400 Arabidopsis transcription factors: unprecedented sensitivity reveals novel root- and shoot-specific genes. *Plant J* 38: 366-379

Forner J, Pfeiffer A, Langenecker T, Manavella PA, Lohmann JU (2015) Germline-transmitted genome editing in Arabidopsis thaliana Using TAL-effector-nucleases. *PLoS One* 10: e0121056

Leibfried A, To JP, Busch W, Stehling S, Kehle A, Demar M, Kieber JJ, Lohmann JU (2005) WUSCHEL controls meristem function by direct regulation of cytokinin-inducible response regulators. *Nature* 438: 1172-1175

Schmid M, Davison TS, Henz SR, Pape UJ, Demar M, Vingron M, Scholkopf B, Weigel D, Lohmann JU (2005) A gene expression map of Arabidopsis thaliana development. *Nat Genet* 37: 501-506

Varkonyi-Gasic E, Wu R, Wood M, Walton EF, Hellens RP (2007) Protocol: a highly sensitive RT-PCR method for detection and quantification of microRNAs. *Plant Methods* 3: 12

Wahl V, Ponnu J, Schlereth A, Arrivault S, Langenecker T, Franke A, Feil R, Lunn JE, Stitt M, Schmid M (2013) Regulation of flowering by trehalose-6-phosphate signaling in Arabidopsis thaliana. *Science (New York, NY)* 339: 704-707

Yadav RK, Perales M, Gruel J, Girke T, Jonsson H, Reddy GV (2011) WUSCHEL protein movement mediates stem cell homeostasis in the Arabidopsis shoot apex. *Genes Dev* 25: 2025-2030

Zurcher E, Tavor-Deslex D, Lituiev D, Enkerli K, Tarr PT, Muller B (2013) A robust and sensitive synthetic sensor to monitor the transcriptional output of the cytokinin signaling network in planta. *Plant Physiol* 161: 1066-1075