Sugar Signaling Induces Dynamic Changes during Meristem Development in Arabidopsis

Magdalena Musialak-Lange[1], Katharina Fiddeke[1,†,#], Annika Franke[1,†,#], Friedrich Kragler[1], Christin Abel[1], and Vanessa Wahl[1,*]

Affiliations:

[1]Max Planck Institute of Molecular Plant Physiology, Department of Metabolic Networks, Am Mühlenberg 1, Potsdam, 14476, Germany.

[†]Current work addresses: ProBioGen AG, Herbert-Bayer-Str. 8, 13086 Berlin (K.F.) and Takeda GmbH, Lehnizstra. 70-98, 16515 Oranienburg (A.F.)

[#]K.F. and A.F. contributed equally

*Correspondence to: vanessa.wahl@mpimp-golm.mpg.de.

One Sentence Summary:

The increase in meristem size during the floral transition is regulated via a feedback regulation involving sugar signaling.

Aerial parts of plants originate from pluripotent cells in the shoot apical meristem

(SAM). This population of stem cells is maintained via a negative feedback loop involving stable

expression of *WUSCHEL* (*WUS*) and *CLAVATA3*. SAM size is dynamic and undergoes a more than

2-fold expansion upon the transition to reproductive growth. The underlying mechanism

controlling this process is largely unknown, but coinciding increased levels of trehalose 6-

phosphate (T6P) suggest a participation of sugar signaling. Here we show that *TREHALOSE*

PHOSPHATE PHOSPHATASE J, a component of the T6P pathway, is directly regulated by WUS,

and controls SAM expansion at floral transition through WUS. Our findings demonstrate a

dynamic feedback-regulation between central meristem maintenance and flowering time

regulators with sugar signaling.

During the floral transition the *Arabidopsis thaliana* SAM undergoes a dramatic increase in size[1] (Fig. 1a). Since the transition between developmental stages requires a massive reorganization of organ development and sufficient energy, it is tightly controlled by environmental conditions and availability of nutrients[2-5]. Hence, plant development as a whole and SAM maintenance in particular demands continuous cross talk between its regulatory processes and the available resources. The sugar phosphate, trehalose 6-phosphate (T6P), serves as a signal for sucrose availability, which is conveyed to downstream metabolic and growth responses through a still largely unknown mechanism[6]. Also knowledge on the underlying mechanism controlling the morphological processes at the SAM at floral transition is scarce. However, increased levels of T6P coincide with the floral transition and the T6P pathway induces flowering by regulating key flowering genes expressed in leaves and the SAM[4], which suggests a participation of sugar signaling in regulating the SAM size changes throughout the floral transition.

To assess whether the morphological changes at the SAM involve the T6P pathway, we investigated the effect on SAM architecture by decreasing *TPS1*, coding for the T6P synthesizing enzyme, by the means of an artificial microRNA also in the SAM proper (*35S:amiRTPS1,* Fig. S1). We compared these lines with plants with increased *TPS1* in the central zone (*CLV3:TPS1*, Fig. S2). These have smaller and bigger vegetative and reproductive meristems (Fig. 1b, 1c), resulting in smaller and bigger plants, respectively[4]. The homeodomain transcription factor WUS and the signaling peptide CLV3 non-cell autonomously contribute to communication of position-dependent properties among SAM cells in a negative feedback loop and maintain the SAM cell population[7]. We therefore analyzed their expression in the SAM with altered *TPS1*. While the expression of *WUS* in the organizing centre (OC) and *CLV3* in the central zone (CZ) harboring

the stem cells are considered mutually exclusive[7], we found reduced *TPS1* levels

(*35S:amiRTPS1*) lead to an enlarged OC domain marked by *WUS* (Fig. 1d) that overlaps with the

CZ, marked by *CLV3* expression (Fig. 1D). Plants overexpressing *TPS1* (*35S:TPS1*, Fig. S2)

display an increased stem cell pool with otherwise little effect on *WUS* expression (Fig. 1d).

However, the size of the CZ decreases in plants expressing *TPS1* under the *CLV3* promoter (Fig.

1d, Fig. S2), indicating a smaller stem cell pool in support of a much smaller SAM size

phenotype (Fig. 1d) of the very early flowering *CLV3:TPS1* plants[4]. The expression of

HISTONE4, CYCLIN D3;1 and *CYCLIN-DEPENDENT KINASE2;1*, three cell cycle marker

genes (Fig. S3), suggests a higher proliferation rate and an increased number of cells displaced

from the CZ to adopt organ-specific cell fates at the periphery as a plausible reason for the

decreased SAM size of *CLV3:TPS1* and *vice versa* increased SAM size of *35S:amiRTPS1* plants

(Fig. 1b, 1c).

To assess whether the WUS/CLV feedback loop can uncouple in a wild-type SAM

during normal growth, we analyzed *WUS* and *CLV3* expression in wild-type plants in a time

series spanning the floral transition (Fig. 1e). Throughout the vegetative phase SAM size

gradually increases due to rising cell numbers (Fig. 1a; Fig. S4)[8], reaching its maximum at the

floral transition, independent of whether the plants are grown in continuous long days (LD) (Fig.

1a) or in short days (SD) followed by a transfer to LD (Fig. S5). An enlarging SAM at the floral

transition correlates with a larger *WUS* expression domain expanding into the CZ and the outer

cell layer (L1) (Fig. 1e-g). This is supported by larger meristems of the *35S:amiRTPS1* line,

which express *WUS* in the central zone (Fig. 1d). This observation is also consistent with

massive proliferation of meristematic cells detected when *WUS* is expressed from a *CLV3*

promoter in the CZ that resulted in meristems, no longer producing organs[9]. The presence of

WUS transcript in the outer meristem layers might be explained by a transient cytokinin signal in

L1, which induces *WUS* and was reported to respond to carbon in seedlings[10-12]. However,

cytokinin levels seem not to be altered in the L1 layer of cells as indicated by the synthetic

cytokinin reporter *TCSn:GFP*[13] (Fig. S6). Further, while *WUS* expands to L1 for a short period

(8-10 DAG), the *CLV3* expression domain remains expanded post floral transition (Fig. 1e, 1h).

This suggests a transient uncoupling, which is re-established at the reproductive SAM (16 DAG),

resembling earlier vegetative SAM expression patterns (6 DAG) (Fig. 1e), allowing organ

production to resume.

To understand how the T6P pathway might control this process, we first analyzed

expression of the ten genes encoding TREHALOSE PHOSPHATE PHOSPHATASEs (TPP)[6].

Except for *TPPC* and *TPPD*, which are expressed below detection limit, all were expressed in

distinct SAM domains (Fig. S7). *TPPJ* strongly increases in the enlarged SAM of *clv3-7* due to

an ectopic expression in L1 and L2 (Fig. 2a-c, Fig. S8, S9). Notably, in *clv3-7* and *clv3-10*

mutants high levels of *WUS* in the outer SAM layers (Fig S9)[14] coincides with ectopic expression

of *TPPJ* (Fig. S9). To assess whether ectopic expression of *TPPJ* contributes to the enlarged

SAM of the *clv3* mutant, we next overexpressed *TPPJ* (*35S:TPPJ*, Fig. 2d). This results in

significantly enlarged SAMs (Fig. 2e), and importantly, expands the *WUS* expression domain

into L1 (Fig. 2f), consistent with our finding in wild-type plants at the floral transition (Fig. 1e).

Overlapping expression domains of ectopic *TPPJ* (Fig. 2b) and *WUS* in the *clv3* mutant

SAM implies a direct influence of WUS on *TPPJ* expression. This is supported by an *in silico*

analysis which predicts multiple, canonical WUS binding sites in the sequence upstream of the

TPPJ coding sequence (*TPPJ*[WUS], Fig. 3a)[10,15-17]. WUS is a bifunctional transcription factor,

which was found to both activate and repress gene expression[7]. To understand if WUS directly

controls *TPPJ*, we performed *in vivo* transactivation assays. These show that WUS activates

reporter gene expression when using a 2865 bp *TPPJ* 5' sequence, containing 17 putative

TPPJ^WUS sites (Fig. 3a). Progressive deletion of the *TPPJ* 5' sequence results in a reduction of

reporter gene activation, suggesting an additive effect of the individual *TPPJ*^WUS sites (#1-6; Fig.

3a, 3b). We confirmed the direct binding of WUS to three distinct regions of the *TPPJ* promoter

(I, II, and III, Fig. 3c-e, Fig. S10) by Chromatin immunoprecipitation (ChIP) coupled to PCR

using a specific antibody against WUS (Fig. S11), while all other regions did not indicate

binding (Fig. S10, Fig. S12). We observed enrichment of WUS binding to *TPPJ*^WUS elements up

to 0.67% of the input DNA in *clv3-7* and up to 0.75% of the input DNA in *clv3-10* apices, both

of which express *WUS* at high levels in comparison to input DNA from wild-type apices, where

WUS is expressed in only a few cells and input DNA from leaves, with no WUS (Fig. 3c-e; Fig.

S10). Lastly, electrophoretic mobility shift assays (EMSA) confirm *in vitro* binding of WUS to

the investigated sequences (I, II 1-3, III 1-3; Fig. 3f). In summary, we demonstrate that seven out

of 17 putative *TPPJ*^WUS sites are directly targeted by WUS *in vivo*.

To further assess the role of *TPPJ* at the SAM we used an artificial miRNA (*amiRTPPJ*)

approach to downregulate *TPPJ* (*35S:amiRTPPJ*) (Fig. S13). Similar to a *tppj* T-DNA insertion

mutant, plants overexpressing either of two versions of an *amiRTPPJ* (V1, V2) flower

significantly earlier in both LD and SD (*35S:amiRTPPJ*, Fig. S14, Table S1). *ML1* expression

specifically and stably localizes to the epidermis (L1) throughout the investigated developmental

stages (Fig. S15). *ML1:amiRTPPJ* plants reduce *TPPJ* expression in L1 (arrowheads in Fig.

S16), have smaller SAMs and flower significantly earlier in LD and SD (Fig. 4a-c, Fig. S17,

S18, Table S1), while the level of *TPPJ* reduction is proportional to the acceleration of flowering

(Fig. 4b and c, Fig. S13, Table S1). To date there are no reports that mutants in meristem

maintenance genes are affected in their flowering time. We found that plants mutant for *CLV3*

are late flowering (Fig. 4c, Table S1). However, when *ML1:amiRTPPJ* is introgressed into either

the *clv3-7* or *clv3-10* background, the late flowering phenotype of the mutant plants is restored to

wild type (Fig. 4c, Fig. S18, Table S1), suggesting that *TPPJ* expression in the epidermis is also

necessary for the late flowering phenotype of *clv3*. Hence, the early flowering of

ML1:amiRTPPJ in a wild-type background is due to a reduction of *TPPJ* expression in L1 in its

endogenous expression domain. In addition, other prominent morphological defects of *clv3-10*

such as fasciated stems and enormous inflorescence SAMs are visibly reduced in the presence of

ML1:amiRTPPJ (Fig. S18). *CLV3* expression increases to much higher levels in *ML1:amiRTPPJ*

than in plants overexpressing *TPPJ* (Fig. 4d), suggesting an active role of the T6P pathway in the

outer SAM layer with regard to stem cell maintenance. In addition, *WUS* is induced (Fig. 4d, Fig.

S19), indicating uncoupling of the WUS/CLV feedback loop, consistent with our findings with

altered T6P signaling at the SAM (Fig. 1d, Fig. 2f).

Since the T6P pathway influences the age pathway at the SAM [4], we analyzed mature

miR156 as well as the expression of *SPL3*, *SPL4*, *SPL5*, *SPL9* and *SPL15*, all associated with

floral transition in general[18,19] (Fig. S20, Table S1). We found a strong reduction of mature

miR156 levels correlating with lower *TPPJ* expression in *ML1:amiRTPPJ*, and corresponding

increased expression of *SPL3, SPL4, SPL5*, and *SPL9* (Fig. 4e). In line, miR156 abundance was

highly increased in *TPPJ* overexpressing plants, while the corresponding *SPL*s were

downregulated (Fig. 4e). However, expression of *SPL15*, previously described to be mainly

responsible for integrating the aging stimulus into the regulation of the floral onset[20], was not

differentially expressed in either of the transgenic lines (Fig. S21). In addition, we found miR156

levels significantly increased in *clv3-10* mutant apices (Fig. 4f). In response, the levels of *SPL3*,

SPL5, and *SPL9* were reduced in line with the late flowering phenotype of the mutant. In contrast

to what would be expected, we found more *SPL4* in apices of the late flowering *clv3-10* (Fig. 4f).

Higher order lines from *spl3*, *spl4* and *spl5* deletion mutant plants are significantly later

flowering in all lines with *spl4* (Fig. S20, Table S1). This argues for an important role of *SPL4* in

inducing flowering. *SPL4*, similar to *SPL3*, *SPL5* and *SPL9*, is induced at the wild-type SAM at

floral transition[21]. However, it is expressed in the very center of the SAM, in a domain largely

overlapping with the WUS protein domain[5,17,20,22], while *SPL3*, *SPL5*, and *SPL9* are expressed at

the periphery of the SAM and in the vasculature of young leaves[5,20]. This might denote a direct

regulation of *SPL4* by WUS, which would explain increased *SPL4* in the *clv3* mutant

background (Fig. 4f) and the previously identified partially miR156-dependent regulation of the

*SPL*s by the T6P pathway[4]. Indeed, we identified a larger number of potential *SPL4*^{WUS} sites

upstream of the *SPL4* coding sequence when compared to the 5' intergenic regions of the other

*SPL*s (Fig. S22). However, this finding also suggests that additional players downstream of

CLV3 are important for the onset of flowering, which cannot be bypassed by an otherwise

inductive SPL4 (Fig. 4f). Since carbohydrate status regulates timing of the floral transition[4], it is

interesting that T6P signaling also directly influences the re-organisation of the SAM during the

floral transition (Fig. 4g). Given the ubiquitous nature of carbohydrate signaling and the large-

scale change in sink-source relationships within the plants[23], it will be interesting to determine if

this regulatory mechanism is widely present in the plant kingdom.

Methods

Materials and Methods

<u>Plant material and growth conditions</u>

All *Arabidopsis thaliana* plants used for this study are of the Columbia accession (Col-0).

The lines *clv3-7*[24], *clv3-10*[25], *35S:amiRTPS1*, *CLV3:TPS1*[4], and *WUS:WUS:GFP*[22] have been

described previously as indicated. Plants were grown in controlled growth chambers at 22°C in

long day (LD, 16h light/8h dark) or short day (SD, 8h dark/16h light) conditions with a light

intensity of approximately 160 µmol/m^{-2}s^{-1} and a relative humidity of 60-65%. Controlled

induction of flowering was performed using a previously described shift protocol[21].

<u>Phenotypic analysis</u>

Flowering time was defined as days to flowering (DTF), recorded when shoots were 0.5 cm

high (bolting), and by scoring the total leaf number (TLN). The TLN is the sum of rosette leaves

(RLN, rosette leaf number) and cauline leaves (CLN, cauline leaf number) per plant (Table S1).

On average 20 plants of each genotype were analyzed in one experiment. Plants were grown in

the conditions described above, and randomized every second day to avoid position effects.

<u>Generation of transgenic lines</u>

Generally, Col-0 plants were transformed by the floral dip method[26]. The presence of the

transgene was confirmed by PCR and independent, single-insertion, homozygous T3 plants

were selected and used for all further studies. Oligonucleotides used for cloning are provided in

Table S2, those used for genotyping are given in Table S3. To generate the *35S:TPPJ* and

35S:TPS1 lines, the coding sequences of *TPPJ* (At5g65140) and *TPS1* (At1g78580) were amplified

and introduced into the *pGEM®-Teasy* vector (Promega, Madison, Wisconsin, US), sub-cloned

into the Gateway® entry vector pJLBlue reverse[27] (pVW275 and pVW099), and recombined into

a pGREEN-II-based destination vector with 35S promoter (pVW279 – *TPPJ* and pVW161 – *TPS1*)

using the Gateway® LR clonase II Enzyme mix (Invitrogen, Carlsbad, CA). Generation of artificial

microRNAs targeting *TPPJ* (*ML1:amiRTPPJ* V1 and V2, *35S:amiRTPPJ* V1 and V2) was performed

according to a previously published protocol[28] using the WMD3–Web MicroRNA Designer

(http://wmd3.weigelworld.org/cgi-bin/webapp.cgi). The resulting *Eco*RI/*Bam*HI fragment was

sub-cloned into the Gateway® entry vector *pJLBlue* reverse[27] (V1: pVW498, V2: pVW499) and

recombined using the Gateway® LR clonase II Enzyme mix (Invitrogen, Carlsbad, CA) into a

pGREEN-II-based destination vector with either the *ML1* or *35S* promoter (V1: pVW507, V2:

pVW508, V1: pVW504 V2: pVW505). Double mutant lines of *clv3* and *ML1:amiRTPPJ #1 and #2*

were generated through crosses. All lines were propagated to the F4 generation and tested for

homozygosity. To generate the *spl3, spl4, spl5,* knockout lines a modified version of the two

single guided RNAs (2sgRNA) CRISPR/*Cas9* technology was used[29,30]. In brief, the NGG PAM

recognition sites in the *5'UTR* and last exon of *SPL3*, first exon and *3'UTR* of *SPL4* and *5'UTR* and

3'UTR of *SPL5* were defined using the ATUM webtool (https://www.atum.bio). *pJF1033* vector,

containing 2sgRNA scaffold, was used as a template. Resulting *Bsa*I products were restriction

cloned into the *pJF1031* binary vector[30]. Plants with homozygous deletions for *SPL3, SPL4* or

SPL5 were back-crossed to Col-0. Homozygous, *Cas9*-free lines were used for the experiments

and to generate all higher order mutations, i.e. the *spl34, spl35, spl45* double as well as the

spl345 triple mutant lines.

RNA extraction and cDNA synthesis

Total RNA form homogenized rosette or dissected SAM samples was isolated by a

phenol/chloroform extraction method using a modified TRIzol reagent with the following

composition: phenol 38% (v/v), guanidine thiocyanate 0.8 M ammonium thiocyanate 0.4 M,

sodium acetate 0.1 M pH 5.0, glycerol 5% (v/v), EDTA 5 mM pH 8.0, Na-lauroylsarcosine 0.5%

(v/v). This was then followed by a sodium acetate precipitation to improve RNA purity. cDNA

synthesis, preceded with removal of putative genomic DNA contamination with Dnase I Rnase-

free (Ambion™/Thermo Fisher Scientific, Waltham, Massachusetts, US), was carried out using a

SuperScript™IV Reverse Transcriptase Kit (Thermo Fisher Scientific, Waltham, Massachusetts,

US) according to the manufacturer's instruction. For quantification of mature miR156 the

respective stem-loop primers (Table S4) were added to the cDNA synthesis reaction (1:1 with

oligo dT(18) primer) (Thermo Fisher Scientific, Waltham, Massachusetts, US) as previously

described[31].

RT-qPCR analysis

RT-qPCR analyses were performed with the ABI Prism 7900 HT fast real time PCR system

(Applied Biosystems/Life Technologies, Darmstadt, Germany) using a Power SYBR® Green-PCR

Master Mix (Applied Biosystems/Life Technologies, Darmstadt, Germany) in a 10 µl reaction

volume for expression analyses and in a 5 µl volume for ChIP-PCRs. All oligos are listed in Table

S4 and Table S5 (RT-qPCR and ChIP-PCR respectively). All data were analyzed using the SDS 2.4

software (Applied Biosystems/Life Technologies, Darmstadt, Germany) applying the criteria

described by Czechowski and co-workers (2004). cDNA quality was determined with primers

binding to the 3´ and 5´ regions of *GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE* (*GAPDH*,

At1g13440), and samples with Ct *5´GAPDH*/Ct *3´GAPDH* values > 1 were excluded from further

analyses. Expression values were calculated by the comparative Ct method using the reference

gene index (RGI) as previously described[4] and *POLYUBIQUITIN10* (*UBQ10*, At4g05320) for figure

3C and S14[32]. The primer sequences for the reference and analyzed genes are given in the Table

S4.

RNA *in situ* hybridization

Wax embedding was performed using an automated processor and embedding system

(Leica EG1160, Solms, Germany). Sections of 8 µm thickness were prepared using a rotary

microtome (Leica RM2265; Leica, Wetzlar, Germany).

Probes for RNA *in situ* hybridization were synthesized using the DIG RNA Labeling Kit

(Roche, Mannheim, Germany). For this the ORFs of the target genes were cloned into the

pGEM®-T Easy vector (Promega, Madison, Wisconsin, US) as a template according to

manufactures instructions (oligonucleotides and construct IDs listed in the Table S6) and the

sense and antisense RNA probes were synthetized using T7 and SP6 polymerases, respectively.

RNA in situ hybridizations were carried out as previously described[4]. The final sections were

imaged with an Olympus BX-61 microscope equipped with a DC View II digital camera (Olympus

Europa SE & Co, Hamburg, Germany).

Chromatin immunoprecipitation (ChIP)

For ChIP-PCR, 100 apices per biological replicate (Col-0, *clv3-7* and *clv3-10*) were collected

and snap frozen in liquid N_2. Samples were fixed in 1% (v/v) formaldehyde buffer (10 mM

sodium phosphate buffer, pH 7; 50 mM NaCl; 100 mM sucrose) under vacuum. ChIP was

performed as previously described[33] with modifications: Antibody incubation (anti-WUS; AS11

1759; Agrisera) was extended to o/n at 4°C and incubation with Agarose beads (Protein A-

Agarose; sc-2001; Santa Cruz Biotechnology) to 6 h at 4°C. Immunoprecipitated DNA was

analyzed by RT-qPCR assay. Data were analyzed using the SDS 2.4 software (Applied

Biosystems/Life Technologies, Darmstadt, Germany). Ct values of *TPPJ* promoter regions were

normalized to the Ct value of a region within the *UBQ10* promoter. The % of enrichment was

calculated as relative to the input expression of the individual *TPPJ* promoter regions. The

following controls were performed: ChIP on 100 apices of Col-0, *clv3-7* and *clv3-10* omitting

addition of AB and ChIP on 1.5 g of Col-0, *clv3-7* and *clv3-10* leaves (Fig. S. 10). Please note that

in contrast to the input samples there was no amplification in any of the other ChIP control

samples. Oligonucleotides are listed in the Table S5.

Transactivation assay

For the effector construct line, the *WUS* (At2g17950) coding sequence was PCR amplified

using the Phusion High-Fidelity DNA Polymerase (Thermo Fisher Scientific, Waltham,

Massachusetts, US). The resulting fragment was cloned into the pGEM®-Teasy vector (Promega,

Madison, Wisconsin, US) (pVW310), sub-cloned into the Gateway® entry vector pJLBlue

reverse[27] (pMML056) and recombined using the Gateway® LR clonase II Enzyme mix

(Invitrogen, Carlsbad, CA) into a Gateway® destination vector with 35S promoter (pMDC32)

(pMML058). For the reporter gene constructs, designated regions of the 5' TPPJ region were

amplified with specific primers (Table S2) using the Phusion High-Fidelity DNA Polymerase

(Thermo Fisher Scientific, Waltham, Massachusetts, US). The resulting *Kpn*I/*Acy*I fragment was

sub-cloned into the Gateway® entry vector pMDC162 to obtain GUS gene fusions (Table S2).

The 35Somega:LUC-NOS plasmid containing the LUC gene driven by the 35S promoter was used

as an internal control. Col-0 protoplasts were isolated from leaves of 4-week-old plants and

transfected by a modified polyethylene glycol method[34]. The transfected cells were incubated

for 20 h at 22°C in the light (100 µmol/m^{-2}s^{-1}), harvested by centrifugation at 100 g for 2 min,

and then lysed[35]. Luciferase activity was measured with a luciferase assay kit (Promega,

Madison, Wisconsin, US) according to the manufacturer's instruction, and GUS activity was

determined according to a previously described protocol[35].

<u>Electrophoretic mobility shift assay (EMSA)</u>

WUS coding sequence without STOP codon was PCR amplified using the Phusion DNA

polymerase (New England Biolabs). The resulting fragment was sub-cloned into the Gateway®

entry vector pDONR207 (pMML059) and recombined into a Gateway® destination vector with

35S promoter (pDEST24) using the Gateway® LR clonase II Enzyme mix (Invitrogen, Carlsbad,

CA) resulting in pMML063. The plasmid was transformed into the *Escherichia coli* strain Rosetta

plysS and the protein production was induced with 1 mM isopropyl β-D-1-

thiogalactopyranoside at 30°C over-night. Bacterial cell lysis was performed by sonication (1x, 5

sec, 20% power, 4 cycles, Sonoplus Hd 2070 Sonicator, Badelin, Berlin, Germany) preceded by

incubation on ice for 20 min in the freshly prepared protein extraction buffer (20 mM Na-

phosphate buffer, pH 7.4; 0.5 M NaCl; 1 mM phenylmethylsulfonyl fluoride; 1 mM

Ethylenediaminetetraacetic acid; 1 tablet of cOmplete™ Protease Inhibitor Cocktail (Merck,

Darmstadt, Grmany) per 10 ml of the buffer). For the EMSA 2 µg of protein (crude extract,

determined by Pierce™ BCA Protein Assay Kit, Thermo Fisher Scientific, Waltham,

Massachusetts, US) was used. The EMSA was performed using 5'-IRDey-682-labeled, double-

stranded oligos of 50 bp spanning the putative WUS binding sites in the 5' regulatory region of

TPPJ (Table S7). Binding reactions were carried out using the Odyssey® EMSA Kit (LI-COR®)

according to the manufacturer's instruction with a competitor to probe ratio of 1:200. Results

were visualized using an Oddysey Infrafed Imaging System (Li-Cor, Lincoln, NE).

<u>Confocal microscopy</u>

Apices from developmental series of Col-0 TCSn:GFP cytokinin reporter line lines were

excised using a diamond knife leaving a stalk of stem tissue. This stalk was used to fix the

sample for imaging in a droplet of 0.1% agar covered with Perfluorodecalin (F2 Chmicals,

Lancashire, UK) on a glass slide. A Leica SP8 confocal laser scanning system equipped with a

M6000B-CS microscopy stage, an Argon laser (65 mV), and a 40x water immersion HCX APO

objective was used to image SAMs with following settings: Laser output power 20%; GFP

Excitation (green): wavelength 488 nm / emission detection channel 3: 495 - 520 nm, gain PMT

800 V; plastid auto-fluorescence (blue) emission detection channel 4: 700 - 800 nm, gain PMT

~500 V; scan speed 600 Hz in xyz bi-directional scanning mode with a z-stack distance of

approx. 10 μm. Offset = 0; pixel dimension: 1024 × 1024. To visualize the differences in GFP

presence middle sections of representative SAMs were extracted from z-stacks using the Fiji

software package[36], version 2.0.0-rc-69/1.52 max-intensity Z-Projection function.

<u>Statistical consideration</u>

Statistical significance of flowering time and RT-qPCR data was analyzed both, by one-way

ANOVA (Analysis of Variance) with Tukey' Post Hoc HSD (Honestly Significant Difference) based

on Tukey-Kramer correction (P value <0.05) and two-tailed Student's *t*-test. Significance of the

data was marked with asterisks (*) based on the following criteria: P ≤ 0,05: *, P ≤ 0,01: **, P ≤

0,001: ***. The presented statistical data are based on: flowering time experiments – min. 20

plants of each genotype; SAM area measurements – longitudinal middle section through min. 7

individual SAMs; *WUS* and *CLV3* expression domain size and distance to the summit –

longitudinal middle section through min. 6 individual SAMs RT-qPCR – min. three biological

replicates; Transactivation assay – 6 biological replications; ChIP – 3 biological replications.

Data availability

All data supporting the findings of this study are available in the main text or the

Supplementary Information. All biological materials used in this study are available from the

corresponding author on reasonable request. Source data are provided with this paper.

References and Notes

1   Kwiatkowska, D. Flowering and apical meristem growth dynamics. *J Exp Bot* **59**,
187-201, doi:10.1093/jxb/erm290 (2008).
2   Srikanth, A. & Schmid, M. Regulation of flowering time: all roads lead to Rome.
Cellular and molecular life sciences : CMLS **68**, 2013-2037, doi:10.1007/s00018-011-
0673-y (2011).
3   Ponnu, J. *et al.* The Trehalose 6-Phosphate Pathway Impacts Vegetative Phase
Change in Arabidopsis thaliana. *Plant J*, doi:10.1111/tpj.14965 (2020).
4   Wahl, V. *et al.* Regulation of flowering by trehalose-6-phosphate signaling in
Arabidopsis thaliana. *Science (New York, N.Y.)* **339**, 704-707,
doi:10.1126/science.1230406 (2013).
5   Olas, J. J. *et al.* Nitrate acts at the Arabidopsis thaliana shoot apical meristem to
regulate flowering time. *New Phytol* **223**, 814-827, doi:10.1111/nph.15812 (2019).
6   Fichtner, F. & Lunn, J. E. The Role of Trehalose 6-Phosphate (Tre6P) in Plant
Metabolism and Development. *Annu Rev Plant Biol*, doi:10.1146/annurev-arplant-
050718-095929 (2021).

7   Pfeiffer, A., Wenzl, C. & Lohmann, J. U. Beyond flexibility: controlling stem cells in an
ever changing environment. *Curr Opin Plant Biol* **35**, 117-123,
doi:10.1016/j.pbi.2016.11.014 (2017).
8   Kinoshita, A. *et al.* Regulation of shoot meristem shape by photoperiodic signaling
and phytohormones during floral induction of Arabidopsis. *Elife* **9**,
doi:10.7554/eLife.60661 (2020).
9   Brand, U., Grunewald, M., Hobe, M. & Simon, R. Regulation of CLV3 expression by
two homeobox genes in Arabidopsis. *Plant Physiol* **129**, 565-575,
doi:10.1104/pp.001867 (2002).
10   Leibfried, A. *et al.* WUSCHEL controls meristem function by direct regulation of
cytokinin-inducible response regulators. *Nature* **438**, 1172-1175,
doi:10.1038/nature04270 (2005).
11   Snipes, S. A. *et al.* Cytokinin stabilizes WUSCHEL by acting on the protein domains
required for nuclear enrichment and transcription. *PLoS Genet* **14**, e1007351,
doi:10.1371/journal.pgen.1007351 (2018).
12   Pfeiffer, A. *et al.* Integration of light and metabolic signals for stem cell activation at
the shoot apical meristem. *Elife* **5**, doi:10.7554/eLife.17023 (2016).
13   Zurcher, E. *et al.* A robust and sensitive synthetic sensor to monitor the
transcriptional output of the cytokinin signaling network in planta. *Plant Physiol*
161, 1066-1075, doi:10.1104/pp.112.211763 (2013).
14   Brand, U., Fletcher, J. C., Hobe, M., Meyerowitz, E. M. & Simon, R. Dependence of stem
cell fate in Arabidopsis on a feedback loop regulated by CLV3 activity. *Science (New*
York, N.Y.) **289**, 617-619, doi:10.1126/science.289.5479.617 (2000).
15   Lohmann, J. U. *et al.* A molecular link between stem cell regulation and floral
patterning in Arabidopsis. *Cell* **105**, 793-803, doi:10.1016/s0092-8674(01)00384-1
(2001).
16   Sloan, J. *et al.* Structural basis for the complex DNA binding behavior of the plant
stem cell regulator WUSCHEL. *Nat Commun* **11**, doi:ARTN 2223
10.1038/s41467-020-16024-y (2020).
17   Yadav, R. K. *et al.* WUSCHEL protein movement mediates stem cell homeostasis in
the Arabidopsis shoot apex. *Genes Dev* **25**, 2025-2030, doi:10.1101/gad.17258511
(2011).
18   Hyun, Y., Richter, R. & Coupland, G. Competence to Flower: Age-Controlled
Sensitivity to Environmental Cues. *Plant Physiol* **173**, 36-46,
doi:10.1104/pp.16.01523 (2017).
19   Xie, Y. *et al.* FHY3 and FAR1 Integrate Light Signals with the miR156-SPL Module-
Mediated Aging Pathway to Regulate Arabidopsis Flowering. *Mol Plant* **13**, 483-498,
doi:10.1016/j.molp.2020.01.013 (2020).
20   Hyun, Y. *et al.* Multi-layered Regulation of SPL15 and Cooperation with SOC1
Integrate Endogenous Flowering Pathways at the Arabidopsis Shoot Meristem. *Dev*
Cell **37**, 254-266, doi:10.1016/j.devcel.2016.04.001 (2016).
21   Schmid, M. *et al.* Dissection of floral induction pathways using global expression
analysis. *Development* **130**, 6001-6012, doi:10.1242/dev.00842 (2003).
22   Daum, G., Medzihradszky, A., Suzaki, T. & Lohmann, J. U. A mechanistic framework
for noncell autonomous stem cell induction in Arabidopsis. *Proc Natl Acad Sci U S A*
111, 14619-14624, doi:10.1073/pnas.1406446111 (2014).

23   Fernie, A. R. *et al.* Synchronization of developmental, molecular and metabolic
aspects of source-sink interactions. *Nat Plants* **6**, 55-66, doi:10.1038/s41477-020-
0590-x (2020).
24   Wisman, E., Cardon, G. H., Fransz, P. & Saedler, H. The behaviour of the autonomous
maize transposable element En/Spm in Arabidopsis thaliana allows efficient
mutagenesis. *Plant Mol Biol* **37**, 989-999, doi:10.1023/a:1006082009151 (1998).
25   Forner, J., Pfeiffer, A., Langenecker, T., Manavella, P. A. & Lohmann, J. U. Germline-
transmitted genome editing in Arabidopsis thaliana Using TAL-effector-nucleases.
PLoS One **10**, e0121056, doi:10.1371/journal.pone.0121056 (2015).
26   Clough, S. J. & Bent, A. F. Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. *Plant J* **16**, 735-743, doi:10.1046/j.1365-
313x.1998.00343.x (1998).
27   Mathieu, J., Warthmann, N., Kuttner, F. & Schmid, M. Export of FT protein from
phloem companion cells is sufficient for floral induction in Arabidopsis. *Curr Biol* **17**,
1055-1060, doi:10.1016/j.cub.2007.05.009 (2007).
28   Schwab, R. *et al.* Specific effects of microRNAs on the plant transcriptome. *Dev Cell* **8**,
517-527, doi:10.1016/j.devcel.2005.01.018 (2005).
29   Wang, Z. P. *et al.* Egg cell-specific promoter-controlled CRISPR/Cas9 efficiently
generates homozygous mutants for multiple target genes in Arabidopsis in a single
generation. *Genome Biol* **16**, 144, doi:10.1186/s13059-015-0715-0 (2015).
30   Ruf, S. *et al.* High-efficiency generation of fertile transplastomic Arabidopsis plants.
Nat Plants **5**, 282-289, doi:10.1038/s41477-019-0359-2 (2019).
31   Varkonyi-Gasic, E., Wu, R., Wood, M., Walton, E. F. & Hellens, R. P. Protocol: a highly
sensitive RT-PCR method for detection and quantification of microRNAs. *Plant*
Methods **3**, 12, doi:10.1186/1746-4811-3-12 (2007).
32   Czechowski, T., Bari, R. P., Stitt, M., Scheible, W. R. & Udvardi, M. K. Real-time RT-PCR
profiling of over 1400 Arabidopsis transcription factors: unprecedented sensitivity
reveals novel root- and shoot-specific genes. *Plant J* **38**, 366-379,
doi:10.1111/j.1365-313X.2004.02051.x (2004).
33   Kaufmann, K. *et al.* Chromatin immunoprecipitation (ChIP) of plant transcription
factors followed by sequencing (ChIP-SEQ) or hybridization to whole genome arrays
(ChIP-CHIP). *Nat Protoc* **5**, 457-472, doi:10.1038/nprot.2009.244 (2010).
34   Wu, F. H. *et al.* Tape-Arabidopsis Sandwich - a simpler Arabidopsis protoplast
isolation method. *Plant Methods* **5**, 16, doi:10.1186/1746-4811-5-16 (2009).
35   Yoo, S. D., Cho, Y. H. & Sheen, J. Arabidopsis mesophyll protoplasts: a versatile cell
system for transient gene expression analysis. *Nat Protoc* **2**, 1565-1572,
doi:10.1038/nprot.2007.199 (2007).
36   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat*
Methods **9**, 676-682, doi:10.1038/nmeth.2019 (2012).
421

Acknowledgments

The authors wish to thank all members of the Wahl group for fruitful discussions, T. Seibert, M.

Schmid and P. Wigge for critical reading of the manuscript, D. Walter for input on statistical

analyses, K. Kontbay, M. Liang and J. Van Dingenen for their support with the *spl*

CRISPR/*Cas9* deletion lines, M. Molochko for sharing the *35Somega:LUC-NOS* plasmid, J.

Forner for providing the *pJF1031* and *pJF1033* plasmids.

Funding: Work in the Wahl group was supported by the BMBF (SolaMI, 031B0191), the DFG

(SPP1530: WA3639/1-2, 2-1), and the Max Planck Society.

Authors contribution

VW conceived and designed the experiments and prepared the figures with contributions from

MML. VW and MML analyzed the data and wrote the manuscript. MML, KF, AK, and VW

performed all essential experiments, i.e. VW generated most of the lines used, except for

ML1:amiRTPPJ (KF), the *spl* CRISPR/*Cas9* deletion lines (MML), and all crosses thereof

(MML). MML performed all transactivation assays, ChIP-PCRs, EMSAs and qPCR analyses.

MML, AK, KF, CA, and VW analyzed SAM sizes and scored flowering time. RNA *in situ*

hybridizations were performed by VW, AK, KF and MML. FK and VW took the confocal

images. All authors have read and commented on the text and figures within this manuscript.

Competing interests

440   Authors declare no competing interests.

Additional Information

442 Supplementary Information: Figures S1-S22, Tables S1-S7, References (1-3).

 Figures captions

Fig. 1. The T6P pathway impacts SAM size during development. (**a**) SAM area throughout

development. (**b**) Vegetative and (**c**) inflorescence SAM size of *CLV3:TPS1* and *35S:amiRTPS1*

lines. (**d**) *WUS* and *CLV3* expression by RNA *in situ* hybridisation on longitudinal sections

through Col-0, *35S:amiRTPS1, 35S:TPS1* and *CLV3:TPS1* SAMs of LD-grown plants 8 days

after germination. (**e**) *WUS* and *CLV3* expression by RNA *in situ* hybridisation in vegetative (6

and 8 DAG), transition (10 DAG, dark grey) and inflorescence SAMs (12 and 16 DAG) of LD-

grown Col-0 plants. (**f**) *WUS* expression domain sizes, (**g**) *WUS* expression domain distance to

SAM summit, and (**h**) *CLV3* expression domain area, in vegetative, transition (dark grey) and

inflorescence SAMs of LD-grown wild-type plants. (**i**) Schematic picture of *WUS, CLV3* and

TPS1 expression dynamics with arrows indicating direction of increasing expression domain at

transition. Error bars denote s.d.; significance was calculated based on one-way Annova and a

Student's *t*-test, ***$P<0.001$. ☆ indicates SAM summit. Scale bars 25μm.

Fig. 2. *WUS* and *CLV3* expression in response to changes of *TPPJ* in the SAM. (a, **b)** *TPPJ* expression by RNA *in situ* hybridisation on longitudinal sections through inflorescence SAMs of Col-0 (**a**), and *clv3-7* (**b**), and by qRT-PCR in apices collected from *clv3-7*, *clv3-10* (**c**), and *35S:TPPJ* (**d**) plants. (**e, f**) SAM size of plants overexpressing *TPPJ* (**e**), expression of *TPPJ*, *WUS*, *CLV3* by RNA *in situ* hybridization (**f**). Error bars denote s.d.; significance calculated by one-way Anova and Student's *t*-test, ***P<0.001. ☆ indicates SAM summit. Scale bars are 50μm.

Fig. 3. WUS directly regulates *TPPJ* in the SAM. (**a**) Overview of *TPPJ* 5' regulatory region with putative WUS binding sites (●, ■), position of ChIP-PCR amplicons corresponding to the results shown in (c-e). Boxes marked with I, II, and III indicate 5' TPPJ regions with in total seven core WUS binding sites – **I**: -2795 – -2789 bp, **II**: -2073 – -1830 bp, and **III**: -652 – -564 bp. Sequence location and lengths used in (b) indicated with #1-6. (**b**) Protoplast transactivation assay showing activation of the GUS reporter when coupled to the regions indicated in (a), relative to LUC activity. c indicates untransformed control. (**c-e**) Enrichment of (**c**) region I, (**d**) region II, and (**e**) region III as indicated in (a) measured by ChIP-PCR relative to the input. PB – post binding fraction. (**f**) EMSA for WUS binding to the indicated regions (**a**, I-III). Error bars denote s.d.; significance based on one way Anova and Student's *t*-test, ****P*<0.001. Scale bars are 50μm.

Fig. 4. The role of TPPJ in the outer SAM layer. (**a**) Meristem area of Col-0, *ML1:amiRTPPJ*

V1 and V2, (**b-c**) Flowering time of *ML1:amiRTPPJ* (**B**) and *clv3-10;ML1:amiRTPPJ* (**C**)

shown as days to bolting (**a**, **c**) relative to the wild type. V1 and V2 indicate two independent

versions of artificial microRNAs designed to target *TPPJ* transcript. (**D**, **E**) Relative expression

of *SPL* genes, mature miR156 (**d**), and *TPPJ, CLV3*, and *WUS* (**E**) in SD-grown *ML1:amiRTPPJ*

and *35S:TPPJ* at 40 days after germination. (**f**) Relative expression of mature miR156 and *SPL*

genes in SD-grown *clv3-10* at 40 days after germination. (**g**) Schematic model of vegetative and

transition SAM circuits. Error bars denote s.d.; significance calculated based on one way Anova

and Student's *t* test; *$P<0.05$, **$P<0.01$, ***$P<0.001$. ☆ indicates SAM summit.