Peripheral Neuronal Activation Shapes the Microbiome and Alters Gut Physiology

Jessica A. Griffiths[1,2+], Bryan B. Yoo[1+], Peter Thuy-Boun[3++], Victor Cantu[4++], Kelly Weldon[5,6,++], Collin Challis[1], Michael J. Sweredoski[1], Ken Y. Chan[1#], Taren M. Thron[1], Gil Sharon[1], Annie Moradian[1%], Gregory Humphrey[3], Qiyun Zhu[3], Justin Shaffer[4], Dennis W. Wolan[3], Pieter C. Dorrestein[4,5,6], Rob Knight[4,6,7,8], Viviana Gradinaru[1,2], and Sarkis K. Mazmanian[1,2*]

[1]Division of Biology & Biological Engineering, California Institute of Technology, Pasadena, California, USA, 91125
[2]Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD, 20815
[3]Departments of Molecular Medicine and Integrative Structural and Computational Biology, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, CA 92037, USA
[4]Department of Pediatrics, University of California San Diego, San Diego, CA, USA
[5]Collaborative Mass Spectrometry Innovation Center, Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California San Diego, San Diego, CA, USA
[6]UCSD Center for Microbiome Innovation, University of California San Diego, San Diego, CA, USA
[7]Department of Computer Science and Engineering, University of California San Diego, San Diego, CA, USA
[8]Department of Engineering, University of California San Diego, San Diego, CA, USA
+These authors contributed equally
++These authors contributed equally
#Present Address: Stanley Center for Psychiatric Research, Broad Institute, Massachusetts Institute of Technology, Cambridge, MA, 02142, USA
%Present Address: Precision Biomarker Labs, Cedars-Sinai, 8700 Beverly Blvd. Davis 2904, Los Angeles, CA 90048

ORCIDs: 0000-0002-5586-1567 (J.A.G.); 0000-0001-5868-348X (V.G.); 0000-0003-2713-1513 (S.K.M.)

SUMMARY

The gastrointestinal (GI) tract is extensively innervated by both intrinsic neurons of the enteric nervous system (ENS) and extrinsic neurons of the central nervous system (CNS) and peripheral ganglia that together regulate gut motility, secretion, and immunity. The GI tract also harbors a diverse microbiome, but interactions between the ENS and the gut microbes remain poorly understood. Herein, we activated gut-associated neurons in mice to determine effects on intestinal microbial communities and their metabolites, as well as on host physiology. We used recombinant adeno-associated viral vectors with enhanced tropism for the gut, and no targeting to the brain, to chemogenetically activate either choline acetyltransferase (ChAT)-expressing or tyrosine hydroxylase (TH)-expressing neurons in the periphery. Targeted activation of discrete neuronal subtypes distinctively altered the metagenome, fecal metabolome, and mouse and microbial proteomes. The resulting datasets provide a rich resource, revealing broad and previously unknown roles for ChAT$^+$ and TH$^+$ neurons in modulating microbiome structure, and providing evidence for novel ENS functions such as shaping bile acid profiles and regulating fungal colonization of the gut. Further, ChAT$^+$ neuronal activation upregulated transcriptional pathways for muscle cell proliferation, angiogenesis, and muscle development. Physiologically, while mice displayed increased fecal output following activation of gut-associated ChAT$^+$ and TH$^+$ neurons, only ChAT$^+$ neuronal activation resulted in increased colonic migrating motor complexes and diarrhea-like fluid secretion. These findings suggest that specific subsets of peripherally-activated ENS neurons differentially regulate the gut microbiome and GI physiology in mice.

INTRODUCTION

Diverse cell types in the gastrointestinal (GI) tract coordinate physiology within the gut (Furness, 2006) and throughout the body (Rao and Gershon, 2016). The mammalian gut receives and transmits neuronal signals through ~100,000 extrinsic nerve fibers originating from the sympathetic, parasympathetic, and sensory nervous systems (Grundy and Brookes, 2011). The GI tract is also innervated by an extensive network of over 100 million intrinsic neurons organized into two distinct compartments within the GI tract, namely the myenteric plexus and submucosal plexus (Furness et al., 2014). The neurons of the GI tract, composing the enteric nervous system (ENS), have been implicated in processes including digestion (Schneider et al., 2019), immunity (Gabanyi et al., 2016; Muller et al., 2014), and even complex behaviors (Bravo et al., 2011) in mice. Interactions between neurons of the GI tract and other cell types highlight the diverse roles of the ENS. For example, neuronal pathways in the gut regulate nutrient sensation through intestinal enteroendocrine cells (Kaelberer et al., 2018), modulate the epithelial barrier and mucosal immunity (Jarret et al., 2020; Seillet et al., 2020; Talbot et al., 2020), and dynamically interface with the microbiome (Lai et al., 2019; Matheis et al., 2020). Exposure of the ENS to changing diet, microbiome, and xenobiotics creates inputs distinct from those in the central nervous system (CNS), i.e., the brain and spinal cord.

Choline acetyltransferase (ChAT) and tyrosine hydroxylase (TH) are the rate-limiting enzymes in acetylcholine and catecholamine biosynthesis, respectively, and are key chemical mediators of neurotransmission in the brain and the periphery. Acetylcholine is the primary excitatory neurotransmitter of the gut, and cholinergic neurons represent 60% of the ENS, mediating intestinal propulsion and secretion (Nezami & Srinivasan, 2010; Qu et al., 2008). Several studies have established correlations between neuronal activity, abundance, and specific physiological outcomes (Furness, 2012; Hennig et al., 2015; Niesler et al., 2021). For example, age-associated reduction of ChAT[+] neurons in the ENS coincides with constipation and evacuation disorders (Nezami & Srinivasan, 2010; Qu et al., 2008), and clinical studies have shown that anticholinergic drugs cause constipation and cholinergic agonists can cause diarrhea (Lott & Jones, 2023; Monane et al., 1993). In a disease context, cholera toxin induces hypersecretion and sustained activation of submucosal ChAT[+] neurons in mice (Fung et al., 2018; Wang et al., 2022). Although less characterized, TH[+] neurons and dopamine signaling pathways have also been shown to affect GI motility (Shan Li et al., 2006), and TH[+] neuronal damage in individuals with Parkinson's disease (PD) correlates with increased constipation (Baumuratov et al., 2016; McQuade et al., 2021).

Though known to be important for motility and secretomotor function, ChAT[+] and TH[+] neurons have not yet been systematically characterized and interrogated for their roles in GI physiology (Furness, 2012; Mittal et al., 2017). One barrier to modulation of neuronal populations in the ENS is its size: 35-40 cm in mice. To circumvent the need for direct delivery of effectors, we leveraged a systemically-delivered engineered adeno-associated virus (AAV) with enhanced tropism for the ENS and other peripheral ganglia of mice (Chan et al., 2017). Importantly, this vector, AAV-PHP.S, does not transduce the CNS, allowing us to uncouple peripheral activation from brain-to-gut signaling. We find that activating gut-associated ChAT[+] and TH[+] neurons of mice with chemogenetic modulators (Wess et al., 2013) alters the transcriptional and proteomic landscape of the intestines, as well as the gut metagenome and metabolome. Multi-'omic' analyses allow us to characterize detailed and complex host-microbial interactions, and enable prediction of neuronal influences on a number of biological processes in the gut, including providing novel insights into secondary bile acid production and control of fungal populations, among other interesting associations. In addition, we show that activation of gut-associated neurons strikingly impacts GI function, including motility and fluid secretion. Together, this work reveals differential effects of non-brain activation of ChAT[+] and TH[+] neurons in shaping the gut environment and GI physiology and generates rich datasets as a resource for further exploration (https://github.com/mazmanianlab/Griffiths_Yoo_et_al/).

RESULTS

Distinct spatial localization of ChAT[+] and TH[+] neurons in the ENS

Broad ENS morphology has been previously characterized using immunohistochemistry (IHC) (Furness, 2006; Qu et al., 2008; Hamnett et al., 2022). To map neurons in mice with higher resolution, we used recombinant AAVs to fluorescently label enteric neurons *in vivo*, and tissue clearing techniques to enhance visualization of intact GI tissue (Hama et al., 2015; Treweek et al., 2015; Yang et al., 2014). Imaging whole tissue, without the need for sectioning, preserves neuronal architectures over large distances and across both longitudinal and cross-sectional axes. The AAV capsid variant AAV-PHP.S is optimized for systemic delivery in mice (Deverman et al., 2016) and displays increased tropism for the peripheral nervous system (PNS), including the ENS (Chan et al., 2017). To further optimize ENS expression, we replaced the CAG promoter used in (Chan et al., 2017) with the human Synapsin 1 (hSYN1) promoter, which has been shown to restrict gene expression to neurons (Chan et al., 2017) and minimize expression in peripheral targets such as the dorsal root ganglia (DRGs) (Haenraets et al., 2017). To assess

off-target effects, we compared expression of AAV-PHP.S-delivered hSYN1-mNeonGreen to

that of CAG-mNeonGreen in various non-ENS tissues known to affect GI function (Figure S1).

Expression from the hSYN1 construct only sparsely labeled the DRGs and jugular-nodose

ganglia, and did not label neuronal projections in the vagus nerve or dorsal root, unlike the

previously-used CAG construct (Figures S2A and S2B) (Chan et al., 2017). In the CNS, AAV-

PHP.S-hSYN1 did not label neurons in the brain, brainstem, or spinal cord (Figure S3).

We packaged genes encoding fluorescent proteins (tdTomato or mNeonGreen) under control of

the hSYN1 promoter into AAV-PHP.S, delivered them systemically, and found that 90% (±2.6%

SD) of ENS cells labelled with antibodies against $\underline{P}$rotein $\underline{G}$ene $\underline{P}$roduct 9.5 (PGP9.5), a pan-

neuronal protein, co-localized with virally-labelled neurons in the small intestine (SI) and colon

(Figure 1A). A single systemic injection of AAV-PHP.S-hSYN1-mNeonGreen at a dose of 10^{12}

viral genomes (vg) was sufficient to label spatially diverse regions of the ENS, such as ganglia

proximal and distal to the mesentery (Figure S4A). Viral transduction was uniform throughout

the SI and colon, aside from a small (~1.5 cm) section of the medial colon that, for unknown

reasons, was consistently not well transduced and was therefore excluded from further analysis

(Figure S4B).

To explore the general architecture of the ENS, we transduced wild-type mice with a single i.v.

injection of a pool of AAV-PHP.S packaging multiple fluorescent proteins (AAV-PHP.S-hSYN1-

XFP), which broadly labelled enteric neurons in the gut and enabled us to distinguish cells by

distinct colors resulting from stochastic transduction with different combinations of XFPs (Figure

1B). We quantified the number of neurons and ganglia, as well as the ganglion size (i.e., the

number of neurons in each ganglion) in the myenteric and submucosal plexuses of seven

regions of the SI and two regions of the colon (Figures S5A-S5F). Regions were approximately

1 cm in length and the tissue was sampled every 2-3 cm. We saw that in the SI, the numbers of

neurons and ganglia generally increased toward the distal portion of the myenteric plexus, while

the converse was true for the submucosal plexus (i.e., lower numbers in distal than proximal

regions) (Figures S5A and S5B). Additionally, the size of the ganglia (i.e., the number of

neurons per ganglion) increased in the distal region of the SI myenteric plexus, a feature not

observed in the submucosal plexus (Figure S5C). While neuronal numbers were similar in the

proximal and distal regions of the colonic plexuses (Figure S5D), the number of myenteric

ganglia increased (Figure S5E) while the size of each ganglion decreased in the distal colon

(Figure S5F). Interestingly, submucosal neurons in the proximal colon localized to natural folds

in the tissue (Figure 1B, dashed lines in lower second-from-right panel).

To visualize ChAT$^+$ and TH$^+$ neurons, we employed mouse lines in which Cre recombinase

(Cre) is expressed under the control of the respective gene promoter and engineered viral

constructs with the transgene in a double-floxed inverted orientation (DIO) so that the transgene

is flipped and expressed in a Cre-dependent manner. After transducing ChAT-Cre or TH-Cre

mice with AAV-PHP.S-hSYN1-DIO-XFP, we observed that both neuronal populations occupy

spatially distinct layers of the GI tract, with ChAT$^+$ neurons primarily located in the myenteric

plexus and TH$^+$ neurons more abundant in the submucosal plexus (Figure 1C). Quantifying this

effect, we found more ChAT$^+$ than TH$^+$ neurons in all assayed regions of the myenteric plexus

(Figure 1D), although the density of TH$^+$ myenteric neurons increased distally (Figure 1D, 10-

fold increase from SI-1 vs SI-7; 2-fold increase from SI-7 vs SI-10/13/15). In the small intestine,

by contrast, there were more TH$^+$ than ChAT$^+$ submucosal neurons (Figure 1E). In addition to

providing these insights into ENS architecture, this approach for whole tissue imaging without

the need for antibody labeling (which has limited penetration to deeper layers) should be

broadly useful for profiling other neuronal and non-neuronal cell types in the gut.

Activation of gut-associated neurons reshapes the gut microbiome

The unique spatial organization of ChAT$^+$ and TH$^+$ neurons we observed suggests potentially

distinct functions, which we decided to investigate through specific activation of each neuronal

population. First, we examined the specificity of AAV-PHP.S-hSYN1 by staining gut-extrinsic

PNS ganglia for TH, and found no transduction of TH$^+$ cells in the DRGs or jugular-nodose

ganglia (Figures S2B and S2C). ChAT$^+$ neurons are absent in these peripheral ganglia (Figure

S2C) (Jakob et al., 2021; Tavares-Ferreira et al., 2022). Prior research has shown that AAV-

PHP.S-hSYN1 transduces the prevertebral sympathetic ganglia, which are known to innervate

the gut (Challis et al., 2019), but these ganglia also lack ChAT$^+$ neurons (Kaestner et al., 2019).

In fact, the vast majority of neurons in the prevertebral sympathetic ganglia are TH$^+$ (Browning &

Travagli, 2014; Kaestner et al., 2019).Therefore, for the remainder of the manuscript, we will

use the term "gut-associated" to refer to ChAT$^+$ neurons in the ENS, or TH$^+$ neurons in the ENS

plus innervating prevertebral sympathetic ganglia.

For cell-specific neuronal activation, we employed a Cre-dependent genetic construct encoding

an activating '*D*esigner *R*eceptor *E*xclusively *A*ctivated by *D*esigner *D*rugs' (DREADD), named

hM3Dq, which is a modified neurotransmitter receptor designed to induce neuronal activation

when exposed to Compound 21 (C21), a "designer drug" specific to this receptor (Thompson et

al., 2018). We validated functional gene delivery and expression using intestinal explants from a

ChAT-Cre mouse transduced with the activating DREADD and a construct encoding the calcium

sensor GCaMP6f, observing a gradual increase in fluorescence consistent with a calcium

transient following administration of C21 (Figure S5G and Video Supplement 1).

We reasoned that neuronal activation in the gut may impact the composition and community

structure of the gut microbiome. Accordingly, we transduced ChAT-Cre or TH-Cre mice with

either virus carrying the activating DREADD (AAV-PHP.S-hSYN1-DIO-hM3Dq-mRuby2) or a

control virus expressing only the fluorescent reporter protein (AAV-PHP.S-hSYN1-DIO-

mRuby2). We performed shotgun metagenomics on a longitudinal series of fecal samples

collected prior to and following ChAT[+] or TH[+] neuron activation by C21 (on days 2, 6, and 10 of

C21 administration), as well as contents of the terminal cecum collected on day 10 (Figure 2A).

In ChAT[+]-activated mice, Faith's phylogenetic diversity (i.e., alpha-diversity) decreased

dramatically in the day 10 fecal and cecal samples (Figure 2B), with many microbial taxa less

abundant (Figures 2I-2K; Figure S6; Figure S7). In contrast, TH[+]-activated mice displayed

similar phylogenetic diversity to controls throughout the experiment (Figure 2B). Using weighted

UniFrac distances and principal coordinate analysis (PCoA) to determine the composition of

microbial communities (i.e., beta-diversity), we observed a distinction between ChAT[+]-activated

and control animals in both feces and cecal contents, a shift that was absent in samples from

TH[+]-activated mice and controls (Figures 2C-H). Over the experimental time course,

Verrucomicrobia became significantly enriched in ChAT[+]-activated mice (Figure 2I). To explore

differentially abundant bacterial taxa, we used linear discriminant analysis effect size (LEfSe)

(Segata et al., 2011) and generated cladograms depicting the phylogenetic relationships of

differentially abundant taxa (Figures 2J-2M). This analysis revealed that the bacterial species

Akkermansia muciniphila drove the increase in Verrucomicrobia we observed in ChAT[+]-

activated mice (Figures 2N and 2O).

In addition to identifying microbial species, metagenomic analysis can reveal gene families and

pathways that are differentially abundant in the microbiome. ChAT[+]-activated mice, but not TH[+]-

activated mice, showed changes in beta-diversity of both gene families and pathways, with

shifts evident in the cecal contents and feces collected 9 days after activation (Figures 2P-2S).

The most distinguishing features were highly represented in the control group and

downregulated in ChAT+-activated mice and were mainly associated with bacterial processes,

such as nucleotide biosynthesis and metabolism, and protein translation and transport (Figures

2P-2S; Figures S7C and S7D). This downregulation is consistent with the decrease in bacterial

alpha-diversity we observed in ChAT+-activated mice (Figure 2B). We conclude that neuronal

activation actively reshapes the gut microbiome at community, species, and genetic levels, with

considerable differences between the effects of ChAT+ and TH+ neurons.

Neuronal stimulation impacts the gut metabolome

Given the intimate and intertwined mouse and microbial co-metabolism, the changes in the

microbial metagenome we observed in response to neuronal activation led us to predict that

there would also be alterations in the profile of gut metabolites. We therefore performed

untargeted metabolomics using liquid chromatography with tandem mass spectrometry (LC-

MS/MS) to assay molecular changes in cecal contents and feces following neuronal activation in

the gut. In both ChAT+- and TH+-activated neurons, compared to nonactivated controls (no

DREADD), we observed a strong separation of metabolome profiles in cecal samples taken one

hour following the last C21 injection (Figures 3A and 3B). Thus, targeted activation of ChAT+

and TH+ gut associated neurons appears to strongly influence the gut metabolome.

To contextualize these data, we applied the Global Natural Products Social Molecular

Networking (GNPS) tool (Wang et al., 2016), an open-access mass spectrometry repository and

analysis pipeline. GNPS revealed metabolic networks of both annotated and unannotated

molecules in the cecal contents of ChAT+-activated and TH+-activated mice (Figures 3C and

3D), allowing us to identify metabolites with differential abundance between control and

activated samples. Activation of TH+ neurons strongly increased metabolites whose closest

spectral matches were linoelaidic acid (ID: 626), oleanolic acid methyl ester (ID: 378), and

coproporphyrin I (ID: 739). Metabolites that spectrally resembled xanthine (ID: 259), genistein

(ID: 846), and trans-ferulic acid (ID: 707) were decreased upon activation of TH+ neurons (Table

S1).

In both ChAT+-activated and TH+-activated mice, the molecular networks largely consisted of

level 3 annotations (based on the Metabolomics Standards Initiative (MSI) (Sumner et al.,

2007)) of compounds belonging to the bile acid molecular family and their conjugates, as well as

unannotated analogs (Figures 3C-3D). Primary bile acids are chemicals derived from host

(mouse) cholesterol biosynthesis, which are subsequently co-metabolized by gut bacteria into

secondary bile acids (Aries et al., 1969; Sakai et al., 1980). Interestingly, metabolites with a

closest spectral match to the primary bile acid cholic acid (IDs: 108, 114, 215, 219, 221, 224,

259) were significantly enriched in the cecum of ChAT[+]-activated mice (Figures 3D-3F; Table

S1). Additional metabolites that spectrally resemble tauro-conjugated primary bile acids, such

as taurocholic acid (IDs: 234, 248) and taurohyocholic acid (ID: 235), trended upwards.

Conversely, features matching the spectra of secondary bile acids and bile acid metabolites

such as ursodeoxycholic acid (ID: 13), deoxycholic acid (ID: 100), beta-hyodeoxycholic acid

(IDs: 1, 143) and 12-ketodeoxycholic acid (IDs: 19, 138) were decreased in ChAT[+]-activated

mice (Figures 3D-3F). These data suggest that activation of ChAT[+] neurons may modulate,

either directly or indirectly, primary bile acid secretion and/or metabolism to secondary bile

acids, which have been implicated in a number of metabolic and immunologic functions, as

discussed below.

Neuronal subpopulations differentially shape the gut luminal proteome

Proteins from the mouse, gut microbes, and diet converge and interact in the GI tract

(Albenberg and Wu, 2014). We performed untargeted label-free proteomics by LC-MS/MS of

cell-free supernatants of the cecal contents from ChAT[+]-activated and TH[+]-activated mice and

controls collected one hour following the final C21 treatment (see Figure 2A). Consistent with

the increase in cecal bile acid metabolites that we observed in ChAT[+]-activated mice, we report

an increased abundance of Niemann-Pick C1-Like 1 protein (NPC1L1) in the cecum of these

mice (Figure 4A). NPC1L1 is expressed on the apical surface of enterocytes, and is integral to

the absorption of free cholesterol, the precursor of bile acids, from the lumen (Jia et al., 2011).

Goblet cell-related proteins, specifically Mucin-19 (MUC19) and Zymogen granule 16 (ZG16), a

protein localized to secretory granules (Rodríguez-Piñeiro et al., 2013), also trended upwards

following ChAT[+] neuronal activation (Figure 4A). Conversely, one of the most highly

downregulated proteins was an aldehyde dehydrogenase (Q3U367) encoded by the *Aldh9a1*

gene, which is involved in the catalytic conversion of putrescine to gamma-aminobutyric acid

(GABA) (Matsushima et al., 1986). While GABA is the primary inhibitory neurotransmitter in the

CNS, little is known about its role in the ENS. The most significantly upregulated proteins in

cecal contents of ChAT[+]-activated mice were pancreatic digestive enzymes including

chymopasin (CTRL), chymotrypsinogen B1 (CTRB1), and pancreatic lipase related protein 2

(PNLIPRP2) (Figure 4A). Accordingly, network analysis of upregulated proteins revealed that

KEGG pathways associated with digestion represent the majority of the network (Figure 4B).

This is consistent with evidence that cholinergic, viscerofugal neurons send signals from the GI

tract to other organs of the digestive system, including the pancreas (Furness et al., 2014).

Cholinergic innervation of the pancreas plays a significant role in regulating pancreatic

functions, such as the secretion of digestive enzymes and insulin release (Li et al., 2019).

Peripheral activation of TH$^+$ gut-associated neurons also altered the luminal proteome of the

cecum. Notably, 88% (52/59) of the differentially abundant proteins ($p_{adj.}$<0.25) were distinct

from those identified in ChAT$^+$-activated mice. The overall direction of the effect was also

reversed: ~90% of differentially-abundant cecal proteins in TH$^+$-activated mice were upregulated

(53/59), compared to ~18% in ChAT$^+$-activated mice (20/112), suggesting that activation of

distinct neuronal subsets is associated with opposing changes in GI function. We observed

signatures of increased protein-protein interactions in cecal contents of TH$^+$-activated mice,

evidenced by more network nodes and connections (Figure 4D). Filamin B (FLNB) and spectrin

beta chain, non-erythrocytic 1 (SPTBN1) were two of the most significantly enriched proteins

following TH$^+$ neuron activation (Figure 4C). Both are associated with the intestinal brush border

and membrane vesicles (Donowitz et al., 2007; McConnell et al., 2011). Accordingly, coatomer

proteins also trended upward (COPA and COPB2) (Figure 4C) and vesicle-mediated transport

was one of the major protein networks altered (Figure 4D). Other upregulated protein interaction

networks were associated with metabolic pathways, ribosomal activity, and the immune system

(Figure 4D). For example, the immune-related proteins immunoglobulin heavy constant alpha

(IGHA) (in ChAT$^+$-activated), immunoglobulin heavy constant gamma 2C (IGHG2C), and

complement component 3 (in TH$^+$-activated) trended upward (Figures 4A and 4C).

Perhaps the most intriguing observation was the strong depletion of acidic mammalian chitinase

(CHIA) upon activation of TH$^+$ neurons (Figure 4C). Chitin is a natural polysaccharide that is a

major component of fungal cell walls (Latgé, 2007), but intestinal chitinases are poorly studied

in mice. This result prompted us to query the pan-proteomic dataset against a microbial protein

database, which revealed that the decrease in CHIA abundance following TH$^+$ neuron activation

was accompanied by a large bloom in fungal-associated peptides in the microbiome (~59% of

peptides mapped to any microbe) (Figure 4E). In contrast, fungal peptides represented only

~0.4% of enriched peptides in the lumen of ChAT$^+$-activated mice (Figure 4F). Unfortunately, we

were unable to corroborate these proteomic data with metagenomics since the DNA extraction

method we used was not optimized for fungi. However, these findings suggest that the reduced

chitinase production of activated TH$^+$ cells is directly associated with a dramatic increase in

fungal proteins, which, if experimentally validated in future, would represent a novel circuit by

which the gut-associated neurons of mice regulates fungal load in the gut.

Activation of ChAT⁺ and TH⁺ neurons alters the intestinal transcriptome

Given the changes to the gut microbiome, proteome and metabolome that we observed, we

were interested in the tissue-level impact of neuronal activation on the intestinal transcriptome.

We therefore profiled gene expression with QuantSeq, a quantitative 3' mRNA-sequencing

technology, in 1 cm of tissue from the distal SI and proximal colon harvested one hour after the

last C21 injection. Rapid and transient expression of immediate early genes (IEGs) is widely

used as a measure of increased neuronal activity (Wu et al., 2017), and the IEGs *Fos*, *Egr1*,

Jun, and *Klf2* were among the most significantly upregulated transcripts we identified in the SI

and colon of both ChAT⁺- and TH⁺-activated mice (Figures 5A-5D) . These IEGs are also known

to be upregulated during growth and differentiation of highly active cell types such as immune

cells (Bahrami and Drabløs, 2016; Ramirez-Carrozzi et al., 2009), smooth muscle cells (Miano

et al., 1993), and intestinal epithelial cells (Flandez et al., 2008).

In the distal SI, we found similar numbers of differentially-expressed genes (DEGs; $p_{adj.}<0.05$) in

ChAT⁺-activated mice (162 DEGs) and TH⁺-activated mice (165 DEGs) (Figures 5A and 5C).

The direction of regulation differed, however, with ~73% of DEGs upregulated upon ChAT⁺

activation (118 up, 44 down) and ~58% of DEGs downregulated upon TH⁺ activation (69 up, 96

down). IEGs followed this overall pattern, with 29 upregulated in the distal SI of ChAT⁺-activated

mice but only two upregulated in TH⁺-activated mice (Figure 5E), and three (i.e., *Hbegf, Soca3,*

Mcl1) downregulated (Figure 5C). Similar proportions of DEGs were upregulated in the proximal

colon of both ChAT⁺-activated (169 up, 84 down) and TH⁺-activated mice (130 up, 62 down)

(Figures 5B and 5D). Given the enrichment in fungal proteins and reduction in the level of the

CHIA protein in the TH⁺-activated mice, we explored potential immune responses to fungi but

found no obvious inflammatory signals compared to control mice (Table S3).

To gain insight into the cellular functions of DEGs, we used Gene Set Enrichment Analysis

(GSEA) (Figures 5F-I; Table S2). Notably, the most highly enriched gene ontology (GO) term for

the distal SI of ChAT⁺-activated mice was "regulation of smooth muscle cell proliferation" (Figure

5F), whereas in TH⁺-activated mice it was "response to bacteria" (Figure 5H), consistent with

the increase in immune-related responses suggested by our proteomic dataset. In the proximal

colon, we observed similar GO pathways in ChAT⁺-activated and TH⁺-activated mice (Figures

5G and 5I), suggesting that transcriptomic signatures may depend on the context of the

activated neurons. In the SI, ChAT+ neurons predominantly border muscle cells in the myenteric

plexus, while TH+ neurons neighbor epithelial and immune cells which respond to bacteria in the

submucosal plexus of the distal SI (see Figures 1C-1E). In the colon, both neuronal subsets are

abundant in the myenteric plexus (see Figure 1D). In both myenteric and submucosal plexuses,

we saw a wider breadth of pathways upregulated by activation of ChAT+ neurons than TH+

neurons, with ChAT+ neuronal activation impacting diverse cellular functions in the GI tract,

involving endothelial, epithelial, immune, and adipose cells (Figure 5G; Table S2).

Differential functional GI outcomes of activation of ChAT+ and TH+ neurons

Motivated by the complexity of responses we observed following activation of neuronal

populations in the gut, we decided to assay functional GI outcomes. Both ChAT+ and TH+

neuronal populations are known to be important for motility and secretory function (Furness,

2012; Mittal et al., 2017), but they have never been specifically modulated to study GI

physiology in a freely behaving mammal. Activation of either ChAT+ or TH+ gut-associated

neurons resulted in faster whole gut transit time, increased fecal pellet output, and mass of

cecal contents compared to control mice (Figures 6A-C). Fecal pellets from ChAT+-activated,

but not TH+-activated, mice had increased water content, which is consistent with reports in the

literature of involvement of ChAT+ enteric neurons in fluid secretion (Figures 6D-6F) (Fung et

al., 2018; Monane et al., 1993; Nezami & Srinivasan, 2010). This distinction is particularly

notable given the higher concentration of TH+ neurons than ChAT+ neurons in most regions of

the submucosal plexus (see Figure 1), which is largely responsible for fluid secretion and

absorption (Furness, 2006). Daily administration of C21 for 9 days to control mice (no DREADD)

did not cause any obvious health impairment and the mice maintained body weight throughout

the experimental period (Figures S6C and S6D). TH+-activated mice also maintained body

weight, but ChAT+-activated animals experienced slight weight loss that likely reflects the

diarrhea-like phenotype over 9 consecutive days. To further examine gut motility in the absence

of extrinsic innervation, we analyzed propulsive colonic migrating motor complexes (CMMCs) in

an *ex vivo* system. Activation of ChAT+ neurons by C21 administration resulted in more frequent

migration of motor complexes (Figures 6G, 6I, and 6J; Table S4), whereas activation of TH+

neurons had no effect on CMMCs (Figure 6H; Table S4). The discrepancy between the *in vivo*

and *ex vivo* results we observed may be due to activation of TH+ neurons in the sympathetic

prevertebral ganglia that project to the gut (Furness, 2006). Overall, these data reveal that

ChAT[+], but not TH[+], neurons in the gut mediate intestinal fluid balance and *ex vivo* colonic

motility.

DISCUSSION

While early pioneers of neuroscience in the 20[th] century focused on the ENS as a model, more

recent research has centered on the brain, and our understanding of the CNS has outpaced that

of other neuronal systems in the body. As a result, basic knowledge of many aspects of

neuronal architecture and function within the gut remain rudimentary (Fung et al., 2018; Monane

et al., 1993; Nezami & Srinivasan, 2010). Here, using a viral delivery system with enhanced

tropism for the ENS, we mapped the distribution of ChAT[+] and TH[+] neurons across the mouse

GI tract and assayed the complex effects of their peripheral activation on physiology and

function. Although the DREADD-based activation paradigm we use in this study is inherently

artificial, the results reveal strikingly different roles for neuronal populations, with nearly every

feature characterized (spatial distribution, metagenomic, metabolic, transcriptional, and

proteomic profiles, and even physiological output) unique to cell type.

The viral vector we used, AAV-PHP.S, can transduce other neuronal subtypes in the PNS such

as those in the DRGs and, with a strong ubiquitous promoter, induce transgene expression

(Chan et al., 2017). To limit this off-target effect, we utilized a weaker promoter with increased

ENS specificity and focused our analyses on GI tissue and lumen. By thus excluding most

known extrinsic innervation pathways, we uncover cell-type-specific effects of gut-associated

neuronal activation that are independent of signaling from the brain. It was recently shown that a

similar strategy using AAV-PHP.S-hSYN1 in ChAT-Cre mice also labels ChAT[+] neurons in

cardiac ganglia, and activation of cholinergic neurons reduces heart rate and blood pressure

(Rajendran et al., 2019; Roy et al., 2015). Cardiac afferent neurons signal through the vagus

nerve and jugular-nodose ganglia to the brain, and in sensory pathways through the DRGs and

spinal cord (Mohanta et al., 2023). Though there is no known direct route for signaling from

cardiac ganglia to gut-associated neurons, it is possible that the gut may be impacted by an

indirect route involving the CNS. In the future, further refinement of AAVs through directed

evolution may generate serotypes with exclusive tropism for the ENS and allow full separation

of the functions of intrinsic and extrinsic activation of gut-associated neuronal subsets. Since the

hSYN1 promoter we used has been shown to drive expression only in neurons (Finneran et al.,

2021; Kügler et al., 2003), we did not characterize other cell types, such as enteroendocrine

cells (EECs), also called neuropod cells, which were recently shown to form synapses with

enteric neurons and contribute to sensory transmission from the gut to the brain through the

vagus nerve. Since these cells turn over every few days (Moran et al., 2008) and have not been

reported to express TH or ChAT, we think it unlikely that they are a major contributor to the

activation-induced phenotypes we observed, but we cannot completely rule out their

involvement.

Exposure to the external environment charges the intestines with myriad responsibilities

including absorption and digestion of dietary nutrients, exclusion of xenobiotics, protection from

enteric infection, and partnership with the gut microbiome. Deletion of ChAT in enteric neurons

leads to microbiome dysbiosis (Johnson et al., 2018), and we observed differences in the

compositional profile (both metagenomic and proteomic) of the gut microbiome specific to

activation of ChAT+ neurons. Notably, we found an expansion of Verrucomicrobia driven by *A.*

muciniphila, which has been implicated in human diseases such as obesity (Everard et al.,

2013; Plovier et al., 2017), multiple sclerosis (Cekanaviciute et al., 2017; Jangi et al., 2016), and

seizures (Olson et al., 2018). *A. muciniphila* metabolizes host-derived mucus as a nutrient

source (Derrien et al., 2004; Van Herreweghen et al., 2017), consistent with the increase in

luminal mucin proteins and digestive enzymes we observed in ChAT+-activated mice. A

particularly interesting host-microbial interaction emerged from activation of TH+ cells, which led

to a dramatic decrease in anti-fungal chitinase (CHIA) protein expression and a concomitant

bloom in fungi, suggesting that neuronal circuits can regulate fungal populations in the gut. If

validated, this would be a novel demonstration of a host-microbial interaction with implications

for health. Our study does not, however, reveal the mechanism(s) by which ENS activation

reshapes the gut microbial community structure, which may involve altered colonic motility,

changes in mucus production, modulation of mucosal immune responses, and/or shifts in

metabolism and nutrient availability.

The human gut microbiome possesses as much metabolic capacity as the liver; it is therefore

no surprise that changes to both mouse gut physiology and the microbiome have major

influences on the gut metabolome. In a striking example of mutualism, we report widespread

changes to the pool of intestinal bile acids, molecules produced via host-microbial co-

metabolism. Activation of ChAT+ neurons, but not TH+ neurons, impacted expression of

NPC1L1, which is involved in cholesterol transport. In mammals, cholesterol is the substrate for

production of primary bile acids, which are then metabolized exclusively by the gut microbiome

into secondary bile acids. Bile acids play critical roles in fat absorption (de Aguiar Vallim et al.,

2013), gut motility (Kirwan et al., 1975), hormonal signaling (Watanabe et al., 2006) and

immune functions (Fiorucci et al., 2018), and neurological conditions (McMillin and DeMorrow,

2016). Expression of bile salt hydrolases and hydratases increases the fitness of both

commensal and pathogenic bacteria (Begley et al., 2005; Delpino et al., 2007; Hofmann and

Eckmann, 2006; Jones et al., 2008; Sannasiddappa et al., 2017). While additional work is

required to determine how the ENS affects levels and constitution of the bile acid pool,

understanding the processes that regulate synthesis of secondary bile acids may have

implications for organ systems throughout the body.

Our study complements recent single-cell RNAseq studies of the ENS neuronal transcriptome

(Drokhlyansky et al., 2020) by giving us the ability to selectively activate specific enteric neurons

and explore the dynamic interplay between cells of various lineages in the gut. Importantly, the

transcriptomic changes we observed may be a consequence of direct or indirect effects of

neuronal activation. Indeed, induced activation of ChAT+ or TH+ neurons rapidly changed GI

transit and fluid secretion patterns, which are only a fraction of the processes that may feed

back on epithelial or immune cells, altering their gene expression profiles. Further single-cell

analysis may help dissect the roles of the various intestinal cells that collaborate to coordinate

gut functions.

The ENS adapts and responds to incredibly diverse molecular cues from the environment and

must do so throughout the entire length and surface area of the intestines—the largest and most

extensive internal organ, with a rich network of neurons termed the "second brain" (Gershon,

2015). Exposure to molecules from the diet or the microbiome may modulate ENS function,

along with signals from outside the gut such as the circulatory system. Curiously, many

disorders of the brain are also associated with GI symptoms (Bhavsar et al., 2013; Cersosimo et

al., 2013; Del Giudice et al., 1999; Pfeiffer, 2003; Valicenti-McDermott et al., 2008). While

mechanisms linking the gut and the brain, and their consequences for health, are an active area

of study, the impact of neuronal activation within the ENS has largely been unexplored. Herein,

we establish an experimental system that allows controlled activation of intrinsic and extrinsic

neurons of the gut, separated from inputs from the brain, and demonstrate broad changes in the

gut environment and its physiology that differ by activated neuronal population. The extensive

datasets on activation of two major gut-associated neuronal populations that we generated

should serve as a resource for further studies on the interconnected biological systems

governing the complex relationship between gut physiology and the microbiome. Future

deployment of this approach could enable mapping of neuronal connections into and out of the

gut, providing insights into how the ENS networks with tissues throughout the body and

advancing growing research into the many functions of the GI tract, an endeavor with important

consequences for human health.

ACKNOWLEDGMENTS

We thank members of the Mazmanian laboratory for discussions throughout the research project and critical reading of the manuscript, and Dr. Catherine Oikonomou for invaluable manuscript editing. We thank Dr. Andres Collazo and Caltech's Biological Imaging Facility for training and access to microscopy capabilities and the Caltech Proteome Exploration Laboratory for access to LC-MS/MS capabilities. This research was funded in part by Aligning Science Across Parkinson's (ASAP-020495 and ASAP-000375) through the Michael J. Fox Foundation for Parkinson's Research (MJFF). For the purpose of open access, the authors have applied a CC BY 4.0 public copyright license to all Author Accepted Manuscripts arising from this submission. This work was also supported by grants to S.K.M. from the Heritage Medical Research Institute, Emerald Foundation, Caltech Center for Environmental and Microbial Interactions (CEMI), the National Institutes of Health (GM007616 and DK078938), and the Department of Defense (PD160030).

AUTHOR CONTRIBUTIONS

J.A.G., B.B.Y., and S.K.M. designed experiments. J.A.G. and B.B.Y. performed experiments. J.A.G., K.Y.C., and C.C. (supervised by V.G.) helped with AAV-mediated ENS characterization. P.T.B (supervised by D.W.W.), M.J.S., and A.M. performed acquisition and analysis of proteomic data. V.C., G.H., G.S., Q.Z., and J.S. (supervised by R.K.) performed acquisition and analysis of metagenomic data. K.W. (supervised by P.C.D.) performed acquisition and analysis of metabolomic data. T.M.T. assisted in animal-related work. J.A.G., B.B.Y., and S.K.M wrote the manuscript with input from all authors.

DECLARATION OF INTERESTS

The authors declare no financial or other interests directly related to this work

519   **Figure 1**

Figure 1. ChAT⁺ versus TH⁺ Neuronal Distribution in the ENS

(A) Representative images of SI and colon from mice infected with AAV-PHP.S-hSYN1-tdTomato and immunolabelled with the pan-neuronal antibody PGP9.5. Inset shows quantification of the ratio of tdTomato$^+$ cells / PGP9.5$^+$ cells (N=3 mice, each data point represents the average of 3 representative fields).

(B) Representative images of proximal and distal regions of the SI and colon from AAV-PHP.S-hSYN1-XFP infected mice. Dotted lines demarcate the rugae (folds) in the proximal colon.

(C) Representative images of cross-sections, myenteric, and submucosal plexuses in ChAT-Cre and TH-Cre mice infected with AAV-PHP.S-hSYN1-DIO-XFP.

(D-E) Density of neurons in the myenteric plexus and submucosal plexus of ChAT-Cre and TH-Cre mice normalized to the number of crypts (N=3 mice, each data point represents the average of 3 representative fields).

See also Figures S1-S5 and Video Supplement 1

Source Data Figure 1
https://github.com/mazmanianlab/Griffiths_Yoo_et_al/tree/main/ENS%20quantification

538 **Figure S1**

539
540
541

Figure S2. Extrinsic and Intrinsic Innervation of the GI Tract
Neurons of the sympathetic, parasympathetic, and sensory nervous systems extrinsically innervate the gut and have effects on GI function. In the sensory nervous system, sensory information from the gut is transmitted to the dorsal root ganglia (DRGs) and then to the spinal cord. Additionally, projections from the DRGs transmit sensory information to the gut. Other sensory signals arise from the periphery and are transmitted through the vagus nerve into the jugular-nodose ganglia and then into the brain. In the sympathetic nervous system, neurons from the spinal cord synapse onto neurons in the sympathetic prevertebral ganglia, celiac ganglia, superior mesenteric ganglia, and inferior mesenteric ganglia. These neurons then project to the gut. In the parasympathetic nervous system, the vagus nerve transmits signals from the brain to the gut. The intrinsic neurons of the gut, i.e., the enteric nervous system (ENS), reside in the myenteric and submucosal plexuses along the gut. Figure created with BioRender.com.

554   **Figure S2**

Figure S2. Viral Transduction of the Peripheral Nervous System (PNS)

(A) Representative images showing labelling of the dorsal root ganglia (DRGs) by a single injection (10^{12} vg) of either PHP.S-CAG-mNeonGreen or PHP.S-hSYN1-mNeonGreen. Cryosections were immunostained for the neuronal marker NeuN and tyrosine hydroxylase (TH). Plots show quantification of NeuN+ cells and NeuN+ TH+ cells virally labelled by AAV-PHP.S-CAG vs. AAV-PHP.S-hSyn1 in DRGs. (N=3-5 mice, each data point represents a 40 μM medial cross-section of the DRG, **$p<0.01$, determined by Welch's two-tailed t-test)

(B) Representative images showing labelling of the jugular-nodose ganglia by a single injection (10^{12} vg) of either PHP.S-CAG-mNeonGreen or PHP.S-hSYN1-mNeonGreen. Cryosections were immunostained for the neuronal marker NeuN and TH. Plots show quantification of NeuN+ cells and NeuN+ TH+ cells virally labelled by AAV-PHP.S-CAG vs. AAV-PHP.S-hSyn1 in the jugular-nodose ganglia (N=4-5 mice, each data point represents a 40 μM medial cross-section of the jugular-nodose ganglia, **$p<0.01$, determined by Welch's two-tailed t-test).

(C) Representative images showing labelling of the dorsal root ganglia (DRGs), dorsal, and ventral root by a single injection (10^{12} vg) of PHP.S-CAG-mNeonGreen. Cryosections were immunostained for the neuronal marker NeuN and ChAT. ChAT staining shows expected ChAT+ projections in the ventral root, but not the dorsal root or intrinsic neurons of the DRGs.

574 **Figure S3**

575
576

Figure S3. Viral Transduction of the CNS by AAV-PHP.S

(A) Representative images showing labelling of the brain by a single injection (10^{12} vg) of either AAV-PHP.S-CAG-mNeonGreen or AAV-PHP.S-hSYN1-mNeonGreen. Sections were immunostained for the neuronal marker NeuN.

(B, C) The PHP.S-hSYN1 injected mice did not show detectable mNeonGreen+ projections in the (B) brainstem or (C) spinal cord as were seen in the PHP.S-CAG injected mice.

583 **Figure S4**

584
585

Figure S4. Viral Transduction by AAV-PHP.S in the Colon

(A) Tiled, whole mount microscopy showing widespread labelling in the proximal colon (>1 cm of tissue) from a single injection of AAV-PHP.S-hSYN1-mNeonGreen (10^{12} vg). Orange inset highlights a region opposite the mesenteric attachment, and pink inset highlights a central region of the tissue.

(B) Representative images showing labelling patterns along the colon, including a 1.5 cm region that AAV-PHP.S does not transduce well.

     Figure S5

Figure S5. ENS Architecture and ChAT[+] Neuronal Stimulation in the ENS

(A-C) Quantification of neurons in 7 regions from the SI

(D-F) Quantification of neurons in 2 regions from the colon

(In A-F: N=3-4 mice per group, per intestinal region. Data points are averages from 2-3 images per mouse, per region; *: $p<0.05$, **: $p<0.01$, ***: $p<0.001$, ****: $p<0.0001$, determined by 2-way ANOVA with Sidak's multiple comparisons test)

(G) Fluorescence time course of GCaMP6f following C21 administration in an *ex vivo* preparation of ChAT[+] SI infected with hSYN1-DiO-hM3Dq-mRuby2 and CAG-GCaMP6f.

Figure 2

Figure 2. Gut-Associated ChAT+ and TH+ Neuronal Activation Alters the Gut Microbiome

(A) Experimental paradigm: Cre-dependent hM3Dq was virally administered to either ChAT-Cre or TH-Cre mice. After 3-4 weeks of expression, C21 was injected daily for 10 days to induce specific neuronal activation. Feces were sampled the day prior to the first C21 injection and on days 2, 6, and 10 of C21 administration, and tissue and cecal contents were collected one hour after the last injection.

(B) Faith's phylogenetic diversity of feces and cecal contents over 10 days of neuronal activation in ChAT+ and TH+ mice. Feces were collected pre-experiment (1 day before first injection) and on day 2, 6, and 10. Cecal contents were collected at experimental endpoint on day 10.

(**$p<0.01$, ***$p<0.001$ determined by Kruskal-Wallis one-way ANOVA)

(C-H) Weighted UniFrac principle coordinate analysis (PCoA) of Activated vs. Control in ChAT+ and TH+ mice. Statistics performed in QIIME2 as in (Bolyen et al., 2019)

(I) Stacked bar graph showing phylum-level changes in relative abundance on day 6 and day 10 of injection for feces and day 10 for cecal contents

(J-M) Linear discriminant analysis (LDA) Effect Size (LEfSe) of the cecal microbiome. Cladograms showing differential phylogenetic clusters and family-level differences in activated and control (J,K) ChAT+ and (L,M) TH+ mice

(Cutoff: $\log_{10}$(LDA Score) > 2 or < -2)

(N-O) Changes in relative abundance of *Akkermansia muciniphila* in feces and cecal contents of (N) ChAT+ and (O) TH+ mice. (n=11-14 mice per group, per time point; red=Control, green=hM3Dq-Activated, *$p<0.05$, **$p<0.01$, determined by multiple t-tests with Holm-Sidak correction for multiple comparisons)

(P-S) Beta-diversity of bacterial gene families and pathways in the (P,R) cecum and (Q,S) post-9 feces of control and activated mice. The direction and length of arrows indicate their influence in separating control and activated groups. Colors represent gene families and pathways (annotated in Figure S7).

See also Figures S6 and S7

Source Data Figure 2
https://github.com/mazmanianlab/Griffiths_Yoo_et_al/tree/main/metagenomics

Figure S6

Figure S6. Annotated changes in gut bacteria (Related to Figure 2 and Figure 6)

(A,B) Species-levels differences in cecal bacteria after 10 days of C21 injection in (A) ChAT[+] and (B) TH[+] mice

(C,D) Analysis of activation-mediated changes in body weight of (C) ChAT[+] and (D) TH[+] mice over 9 days of C21 administration

(N = 10-11 mice per group; **: $p<0.01$, ***: $p<0.001$, determined by 2-way ANOVA with Sidak's method for multiple comparisons)

643   **Figure S7**

A.

Firmicutes
Clostridia
Erysipelotrichia
Betaproteobacteria
Proteobacteria
Flavobacteriia
Verrucomicrobiae
Verrucomicrobia
Spirochaetia
Spirochaetes

B.

Flavobacteriia
Spirochaetia
Spirochaetes

ChAT-Cre Day 10 Feces							
ID	Microbial Taxon	ID	Microbial Taxon	ID	Microbial Taxon	ID	Microbial Taxon
a	Bifidobacterium	p	Parabacteroides	b4	Dorea	d0	Clostridiales
b	Bifidobacteriaceae	q	Tannerellaceae	b5	Eisenbergiella	d1	Candidatus Stoquefichus
c	Bifidobacteriales	r	Marinilbiliales	b6	Faecalicatena	d2	Faecalitalea
d	Cellulomonas	s	Flavobacteriales	b7	Lachnoclostridium	d3	Erysipelotrichaeceae
e	Cellulomondaceae	w	Enterococcus	b8	Marvinbryantia	d4	Erysipelotrichales
f	Micrococcales	x	Enterococcaceae	b9	Sellimonas	d5	Parasutterella
g	Adlercreutzia	a2	Intestinimonas	c0	Tyzzerella	d6	Sutterellaceae
h	Barnsiella	a3	Pseudoflavonifractor	c1	Lachnospiraceae	d7	Burkholderiales
i	Coprobacter	a5	Clostridium	c2	Oscillibacter	d8	Lawsonia
j	Barnesiellaceae	a6	Clostridiaceae	c3	Oscillospiraceae	d9	Treponema
k	Prevotella	a8	Anaerofustis	c5	Anaerotruncus	e0	Spirochaetaceae
l	Prevotellaceae	a9	Eubacterium	c6	Ruminiclostridium	e1	Spirochaetales
m	Alistipes	b0	Eubacteriaceae	c7	Ruminococcus	e2	Akkermansia
n	Rikenella	b2	Blautia	c8	Ruthenibacterium	e3	Akkermansiaceae
o	Rikenellaceae	b3	Coprococcus	c9	Ruminococaceae	e4	Verrucomicrobiales
						e5	Varicellovirus

TH-Cre Day 6 Feces	
ID	Microbial Taxon
a	Parabacteroides
b	Tannerellaceae
c	Fluviicola
d	Flavobacteriales
e	Streptococcus
f	Streptococcaceae
g	Pseudoflavonifractor
i	Clostridium
j	Blautia
k	Eisenbergiella
l	Clostridiodes

Enriched in:
🔴 Control
🟢 Activated

C.

ChAT-Cre Cecum		
ID	Gene Family	Annotation
1	K02003	putative ABC transport system ATP-binding protein
2	K02518	translation initiation factor IF-1
3	K02948	small subunit ribosomal protein S11
4	K02950	small subunit ribosomal protein S12
5	K02965	small subunit ribosomal protein S19
6	K03205	type IV secretion system protein VirD4
7	K03294	basic amino acid/polyamine antiporter, APA family
8	K03496	chromosome partitioning protein
ID	Gene Pathway	Annotation
1	HISDEG-PWY	L-histidine degradation I
2	ILEUSYN-PWY	L-isoleucine biosynthesis I (from threonine)
3	PWY-6122	5-aminoimidazole ribonucleotide biosynthesis II
4	PWY-6595	superpathway of guanosine nucleotides degradation (plants)
5	PWY-6606	guanosine nucleotides degradation II
6	PWY-7111	pyruvate fermentation to isobutanol (engineered)
7	PWY-7219	adenosine ribonucleotides de novo biosynthesis
8	VALSYN-PWY	L-valine biosynthesis

D.

ChAT-Cre Day 10 Feces		
ID	Gene Family	Annotation
1	K02003	putative ABC transport system ATP-binding protein
2	K02518	translation initiation factor IF-1
3	K02913	large subunit ribosomal protein L33
4	K02946	small subunit ribosomal protein S10
5	K02948	small subunit ribosomal protein S11
6	K02950	small subunit ribosomal protein S12
7	K02965	small subunit ribosomal protein S19
8	K03205	type IV secretion system protein VirD4
ID	Gene Pathway	Annotation
1	ILEUSYN-PWY	L-isoleucine biosynthesis I (from threonine)
2	PWY-6151	S-adenosyl-L-methionine cycle I
3	PWY-6700	queuosine biosynthesis I (de novo)
4	PWY-6859	all-trans-farnesol biosynthesis
5	PWY-7111	pyruvate fermentation to isobutanol
6	PWY-7219	adenosine ribonucleotides de novo biosynthesis
7	PWY0-1296	purine ribonucleosides degradation
8	VALSYN-PWY	L-valine biosynthesis

644
645
646

Figure S7. Gut-Associated Activation-Mediated Changes in the Fecal Microbiome
(Related to Figure 2)
(A) LEfSe cladograms and annotations for feces on Day 10 of C21 administration in ChAT+ mice
(B) LEfSe Cladograms and annotations for feces on Day 6 of C21 administration in TH+ mice
(C-D) Microbial gene family and pathway annotations for the feature arrows in Figures 2P-2S. Colors in
the ID column refer to the colors of the arrows in the main figure.

Figure 3

ChAT⁺ Cecum						
Activated > Control			**Activated < Control**			
ID	GNPS Annotation	Adduct	ID	GNPS Annotation	Adduct	
215	Cholic acid	2M+H	19	Spectral Match to 12-Ketodeoxycholic acid (NIST14)	M+H	
219	Cholic acid	2M+Na	187	hyocholic acid	M-2H2O+H	
218	7-Keto-3.alpha.,12-.alpha.-dihydroxycholanic acid	M+H-3H2O	138	Spectral Match to 12-Ketodeoxycholic acid (NIST14)	M+H	
108	Cholic acid	M-3H2O+H	101	.beta.-Muricholic acid	M+H-3H2O	
221	Cholic acid	3M+H	34	3.alpha.-Hydroxy-7-oxo-5.beta.-cholanic acid	M+H-H2O	
114	Cholic acid	M+Na	20	.alpha.-Hyodeoxycholic acid methyl ester	M+H-CH6O2	
259	Cholic Acid	[M+NH4]+	280	.gamma.-Muricholic acid/hyocholic acid	M+H-H2O	
224	Cholic acid	3M+Na	100	Spectral Match to Deoxycholic acid (NIST14)	2M+H	
235	taurohyocholic acid	M+H	13	Spectral Match to Ursodeoxycholic acid (NIST14)	M+H-H2O	
234	Spectral Match to Taurocholic acid (NIST14)	M+H	143	.beta.-Hyodeoxycholic acid	M+H-H2O	
248	Spectral Match to Taurocholic acid (NIST14)	M+Na	1	.beta.-Hyodeoxycholic acid	M+H-2H2O	
245	taurohyocholic acid	M-2H2O+H				
272	Spectral Match to Cholic acid (NIST14)	M+H-2H2O				
102	.beta.-Muricholic acid	M+H-H2O				
99	hyocholic acid	M-2H2O+H				

Primary Bile Acid
Secondary Bile Acid

Figure 3. Gut-Associated ChAT$^+$ or TH$^+$ Neuronal Activation Alters Host and Microbe-Derived Luminal Metabolites

(A-B) Canberra PCoA of the cell-free, luminal metabolome of cecal contents from control (red) and activated (green) ChAT$^+$ and TH$^+$ mice. Statistical analyses were performed in QIIME2 as in (Bolyen et al., 2019).

(C-D) Metabolic networks constructed from identified cecal metabolites in TH$^+$ and ChAT$^+$ mice. Each node is colored by its upregulation (green) or downregulation (red) in the activated group and is labelled with an ID number corresponding to annotation, mass-to-charge ratio, retention time, fold change, and significance value in Table S1.

(E) Fold changes of specific bile acids identified as upregulated (green bars) or downregulated (red bars) in activated ChAT$^+$ mice

(F) Annotations of bile acids highlighted in (E). Metabolite IDs are colored according to annotation as primary (blue) or secondary (orange) bile acids. Metabolite IDs are specific to each sample (N=12-14 for each group analyzed, *: $p<0.05$, **: $p<0.01$, ***: $p<0.001$, ****: $p<0.0001$)

See also Table S1

 Figure 4

Figure 4. Gut-Associated ChAT⁺ or TH⁺ Neuronal Activation Alters Host and Microbe-Derived Luminal Proteins

(A) Volcano plot of differentially-expressed host proteins identified in the cecal contents of ChAT⁺-activated (N=8) vs. ChAT⁺-control mice (N=9) mice, 1 hour after final C21 administration

(B) STRING network analysis of host proteins that were more abundant in ChAT+-activated mice ($p_{nom.}$< 0.2)

(C) Proteomic volcano plot of TH⁺-activated vs. TH⁺-control mice (N=7 mice per group)

(D) STRING network analysis of upregulated host proteins in TH⁺-activated mice ($p_{nom.}$< 0.2).

(E-F) Unipept metaproteomic analysis of upregulated microbial proteins (fold change > 2, $p_{nom.}$< 0.2) in TH⁺-activated and ChAT⁺-activated mice (N=7-9 mice per group)

Source Data Figure 4A

https://github.com/mazmanianlab/Griffiths_Yoo_et_al/blob/main/proteomics/CHAT_proteomics_volcano.txt

Source Data Figure 4C

https://github.com/mazmanianlab/Griffiths_Yoo_et_al/blob/main/proteomics/TH_proteomics_volcano.txt

Source Data Figures 4E and 4F

https://github.com/mazmanianlab/Griffiths_Yoo_et_al/blob/main/proteomics/metaproteomics/Microbiome_associated_proteins.xlsx

Figure 5

Figure 5. ChAT $^+$ and TH$^+$ Activation-Mediated Transcriptomic Changes
(A-D) Differentially-expressed genes in DREADD-activated vs. control (A) ChAT $^+$ distal SI, (B) ChAT$^+$
proximal colon, (C) TH$^+$ distal SI, (D) TH$^+$ proximal colon
(Dashed vertical lines: Fold Change (FC) = +/- 1.5; dashed horizontal lines: $p_{adj.} < 0.05$. Transcripts of
IEGs are highlighted in green and annotated. N=10 mice per group)
(E) Fold changes of upregulated IEGs ($p_{adj.} < 0.05$) as defined by Wu et al., 2017
(F-I) Gene set enrichment analysis of gene ontology (GO) terms for (F) ChAT$^+$ distal SI, (G) ChAT$^+$
proximal colon, (H) TH$^+$ distal SI, (I) TH$^+$ proximal colon.
See also Table S2 and Table S3
706

707 **Source Data Figure 5**

708 https://github.com/mazmanianlab/Griffiths_Yoo_et_al/tree/main/RNAseq

709   **Figure 6**

Figure 6. GI Physiology Differences in ChAT⁺ vs. TH⁺ mice Following Activation

(A) Activation-mediated changes in whole gut transit time in ChAT⁺ and TH⁺ mice

(B) Activation-mediated changes in fecal pellet output in ChAT⁺ and TH⁺ mice

(C) Activation-mediated changes in normalized cecal content mass in ChAT⁺ and TH⁺ mice

(D-E) Fecal pellet water content in (D) ChAT⁺ and (E) TH⁺ mice over 2 hours following C21 activation

(F) Average fecal pellet water content in ChAT⁺ and TH⁺ mice following activation

(A-F: N = 10-11 mice per group; *: $p < 0.05$, ***: $p < 0.001$, ****: $p < 0.0001$, determined by 2-way ANOVA with Sidak's method for multiple comparisons)

(G-H) Frequency of *ex vivo* CMMCs from (G) ChAT⁺ and (H) TH⁺ mice over 30 minutes following activation

(N = 3-8 mice per group; *: $p < 0.05$, determined by 2-way ANOVA with Sidak's method for multiple comparisons)

(I-J) Heatmaps showing frequency of CMMCs over 400 seconds following activation in *ex vivo* preps from (I) ChAT⁺ control and (J) ChAT⁺ DREADD-administered mice

See also Figure S6 and Table S4

Table S1 - Extended GNPS Annotations of Metabolite Network Nodes (Related to Figure 3).

Extended annotations for networked (bold) MS/MS spectra from luminal cecal contents of activated TH[+] and ChAT[+] mice in Figures 3C and 3D, respectively.

Source Data Table S1

https://github.com/mazmanianlab/Griffiths_Yoo_et_al/blob/main/metabolomics/Table_S1-Extended_GNPS_annotations_of_Metabolite_Network_Notes_related_to_figure_3.xlsx

Table S2 - Gene Set Enrichment Analysis of Gene Ontology (GO) Terms (Related to Figure 5).

Gene set enrichment analysis of gene ontology (GO) terms that are upregulated in the distal SI and proximal colon of activated ChAT[+] and TH[+] mice.

Source Data Table S2

https://github.com/mazmanianlab/Griffiths_Yoo_et_al/blob/main/RNAseq/Table_S2-Gene_Set_Enrichment_Analysis_of_GO_related_to_figure_5.xlsx

Table S3 – Genes Related to T helper responses (Related to Figure 5).

List of genes in the transcriptomics dataset associated with T helper responses.

Source Data Table S3

https://github.com/mazmanianlab/Griffiths_Yoo_et_al/blob/main/RNAseq/Table_S3-Genes_Related_to_T_Helper_Responses.xlsx

Table S4 – Annotations of Colonic Migrating Motor Complexes (Related to Figure 6)

Video annotations of CMMCs in *ex vivo* colonic preparations from ChAT[+] and TH[+] mice at baseline and during activation.

Source Data Table S4

https://github.com/mazmanianlab/Griffiths_Yoo_et_al/blob/main/ex_vivo_motility/Table_S4%E2%80%93Annotations_of_Colonic_Migrating_Motor_Complexes_Related_to_figure_6.xlsx

Video S1 – Calcium Imaging of ChAT[+] activated Gut Neurons (Related to Figure S5)

Video of *in vivo* calcium imaging of GCaMP6f-expressing ChAT[+] activated neurons in the myenteric plexus at 5 Hz following C21 administration.

Source Data Video S1

https://github.com/mazmanianlab/Griffiths_Yoo_et_al/blob/main/ENS%20quantification/Video%20Supplement%201-%20GCaMP6F%20at%205Hz.avi

METHODS

Mice

All mouse experiments were performed in accordance with the NIH Guide for the Care and Use of Laboratory Animals using protocols approved by the Institutional Animal Care and Use Committee at the California Institute of Technology. Mice were fed ad libitum for the entire duration of experiments. Homozygous TH-Cre (gift to V.G. from Ted Ebendal, B6.129X1-Thtm1(cre)Te/Kieg (Lindeberg et al., 2004)) and ChAT-Cre (Jackson Laboratories, Bar Harbor, ME- Stock# 028861, RRID:IMSR_JAX:028861) mice were bred to wild-type mice to yield the male and female heterozygous Cre-mice used for our studies. Wild-type specific pathogen free (SPF) C57BL/6 (Jackson Laboratories, Bar Harbor, ME- Stock #000664, RRID:IMSR_JAX:000664) males and females were used for breeding and experiments.

Virus Production

Virus was produced using the methods described in Challis et al., 2019 and dx.doi.org/10.17504/protocols.io.bzn6p5he. Briefly, human embryonic kidney (HEK293T) cells were triple-transfected with pUCmini-iCAP-AAV-PHP.S, pHelper plasmid, and one of the following pAAV genomes: CAG-NLS-GFP, hSYN1-tdTomato, GFAP-tdTomato, hSYN1-mRuby2, hSYN1-DIO-mRuby2, hSYN1-mNeonGreen, hSYN1-DIO-mNeonGreen, hSYN1-mTurquoise2, hSYN1-DIO-mTurquoise2, hSYN1-DIO-hM3Dq-mRuby2, CAG-GCaMP6f. Cells were grown in DMEM + Glutamax + Pyruvate (Gibco, Gaithersburg, MD- Stock# 10569-010) + 5% FBS + non-essential amino acids (Gibco, Gaithersburg, MD- Stock# 11140-050) + penicillin-streptomycin (Gibco, Gaithersburg, MD- Stock# 15070-063). Virus was precipitated from cells and supernatant with an 8% PEG solution (wt/vol), and purified by ultracentrifugation using 15%, 25%, 40%, 60% stacked iodixanol gradients.

Systemic Delivery of AAV

Mice were anesthetized using 2% isoflurane. Virus was titered to 1×10^{12} vg, resuspended in a volume of 100 µl with sterile PBS, and injected retro-orbitally as described in dx.doi.org/10.17504/protocols.io.bzn6p5he.

Neuronal activation of the GI tract

See dx.doi.org/10.17504/protocols.io.bzp5p5q6. TH-Cre and ChAT-Cre mice were used for these experiments. "Activated" mice were infected with AAV-PHP.S-hSYN1-DIO-hM3Dq-mRuby2 and "Control Mice" were infected with AAV-PHP.S-hSYN1-DIO-mRuby2. This was to control for both AAV-PHP.S-mediated expression and the effects of Compound 21 dihydrochloride (C21) (HelloBio, Princeton, NJ- HB6124). C21 was injected intraperitoneally (i.p.) at a dose of 3 mg/kg for 10 consecutive days to both groups of mice. Mice for timecourse experiments were single-housed in sterile cages with autoclaved water following the first C21 administration. Injections of C21 were administered at the same time every day (10AM). Mice were sacrificed one hour after the day 10 injection.

Tissue Preparation, Immunohistochemistry, Imaging, and Quantification

Procedures are described in dx.doi.org/10.17504/protocols.io.bzp6p5re. 100 mg/kg of pentobarbital (Euthasol - Virbac, Carros, France) was administered i.p., and tissues were perfused with 30 mL of phosphate-buffered saline (PBS) and then cold 4% paraformaldehyde (PFA) in PBS. GI tract was post-fixed in 4% PFA overnight at 4 ºC, and stored in PBS + 0.025% sodium azide. Tissues that underwent subsequent immunohistochemistry were made transparent by the passive CLARITY technique (PACT) (Treweek et al., 2015). Briefly, perfused and fixed tissues were embedded with polymerized 4% (wt/vol) acrylamide, and lipids were eliminated using 8% (wt/vol) SDS solution. Jugular-nodose ganglia and dorsal root ganglia

tissues were cryoprotected in 10% then 30% sucrose in PBS for 1 day each. Tissues were
embedded and flash frozen in OCT and cryostat sectioned into 40 μM sections. Spinal cord and
brain tissues were vibratome sectioned into 50 μM sections. Tissues were blocked in 3%
donkey serum and permeabilized with PBS + 0.3% Triton (PBST). Primary antibodies were
incubated in PBST for 48 hours and washed with PBST for 24 hours (replacing the wash
solution 3 times). Tissues were next incubated in secondary antibodies (and DAPI) for 24 hours
and washed in PBS for 48 hours, intermittently replacing the wash solution with fresh PBS.
Primary antibodies used were rabbit anti-PGP9.5 (1:300; Millipore Cat# AB1761-I,
RRID:AB_2868444), chicken anti-GFAP (1:500, BioLegend Cat# 829401, RRID:AB_2564929),
rabbit anti-tyrosine hydroxylase (1:500, Abcam Cat# ab112, RRID:AB_297840), rabbit anti-
choline acetyltransferase, (1:250, Abcam Cat# ab178850, RRID:AB_2721842), and mouse anti-
NeuN (1:300, Abcam Cat# ab104224, RRID:AB_10711040). Secondary antibodies used were
goat anti-chicken Alexa 647 (Thermo Fisher Scientific Cat# A32933TR, RRID:AB_2866500),
donkey anti-rabbit Alexa 568 (Thermo Fisher Scientific Cat# A10042, RRID:AB_2534017), goat
anti-rabbit Alexa 647 (Thermo Fisher Scientific Cat# A-21245 (also A21245),
RRID:AB_2535813), and goat anti-mouse Alexa 594 (Thermo Fisher Scientific Cat# A-11032,
RRID:AB_2534091). GI tissues imaged for virally-expressed, endogenous fluorescence were
made transparent using a sorbitol-based optical clearing method, Sca*l*eS (Hama et al., 2015).
Tissues were mounted in method-respective mounting media (RIMS and Scales S4) on a glass
slide with a 0.5mm spacer (iSpacer, SunJin Lab Co.). Images were acquired on Zeiss LSM 780
or 880, and microscope, laser settings, contrast, and gamma remained constant across images
that were directly compared. All confocal images were taken with the following objectives: Fluar
5× 0.25 M27 Plan-Apochromat, 10× 0.45 M27 (working distance 2.0 mm) and Plan-Apochromat
25× 0.8 Imm Corr DIC M27 multi-immersion. Neurons in each ganglion were counted by
counting cells that were of a distinct color. Colonic ganglia were defined as distinct if separated
by a width of 3 or more neurons.
837
GCaMP6f Fluorescence in *Ex Vivo* Intestinal Preparations
As described in dx.doi.org/10.17504/protocols.io.bzqap5se, small intestinal tissue was quickly
harvested from ChAT-Cre mice, flushed, and placed in oxygenated (95% O_2, 5% CO_2), ice cold
Krebs-Henseleit solution for 1 hour followed by 15 min at room temperature. A segment was cut
along the mesenteric attachment and pinned flat (mucosa facing down) on a Sylgard-lined
recording chamber (Warner Instruments, PH1) in oxygenated Krebs-Henseleit solution. C21
was added at 10nM and GCaMP6f fluorescence was detected on an upright microscope (Zeiss,
Oberkochen, Germany- Examiner D1).
846
Metagenomics
Procedures are described in dx.doi.org/10.17504/protocols.io.bzqep5te.
Fecal collection
AAV-PHP.S-hSYN1-hM3Dq-mRuby2 (10^{12} vg) was delivered systemically to TH-Cre and ChAT-
Cre mice. 3-4 week after infection, C21 (3 mg/kg) was administered daily for 10 consecutive
days. Fecal pellets were collected in sterile containers one day before the initial C21 dose, and
on day 2, 6, and 10 of injections.
Fecal sample DNA extraction and library preparation
DNA was extracted with the Qiagen MagAttract PowerSoil DNA kit as previously described
(Marotz et al., 2017).This protocol is optimized for an input quantity of 1 ng DNA per reaction.
Prior to library preparation, input DNA was transferred to a 384-well plate and quantified using a
PicoGreen fluorescence assay (ThermoFisher, Inc). Input DNA was then normalized to 1 ng in a
volume of 3.5 μL of molecular-grade water using an Echo 550 acoustic liquid-handling robot
(Labcyte, Inc). Enzyme mixes for fragmentation, end repair and A-tailing, ligation, and PCR
were prepared and added in approximately 1:8 scale volumes using a Mosquito HV

micropipetting robot (TTP Labtech). Fragmentation was performed at 37 °C for 20 min, followed
by end repair and A-tailing at 65 °C for 30 min.
864
Sequencing adapters and barcode indices were added in two steps, following the iTru adapter
protocol (Glenn et al., 2019). Universal "stub" adapter molecules and ligase mix were first added
to the end-repaired DNA using the Mosquito HV robot and ligation performed at 20 °C for 1 h.
Unligated adapters and adapter dimers were then removed using AMPure XP magnetic beads
and a BlueCat purification robot (BlueCat Bio). 7.5 µL magnetic bead solution was added to the
total adapter-ligated sample volume, washed twice with 70% EtOH, and then resuspended in
7 µL molecular-grade water.
872
Next, individual i7 and i5 were added to the adapter-ligated samples using the Echo 550 robot.
Because this liquid handler individually addresses wells, and we used the full set of 384 unique
error-correcting i7 and i5 indices, we were able to generate each plate of 384 libraries without
repeating any barcodes, eliminating the problem of sequence misassignment due to barcode
swapping (Costello et al., 2018; Sinha et al., 2017). To ensure that libraries generated on
different plates could be pooled if necessary, and to safeguard against the possibility of
contamination due to sample carryover between runs, we also iterated the assignment of i7 to i5
indices each run, such that each unique i7:i5 index combination was repeated only once every
147,456 libraries. 4.5 µL of eluted bead-washed ligated samples was added to 5.5 µL of PCR
master mix and PCR-amplified for 15 cycles. The amplified and indexed libraries were then
purified again using magnetic beads and the BlueCat robot, resuspended in 10 µL water, and
9 µL of final purified library transferred to a 384-well plate using the Mosquito HV liquid-handling
robot for library quantitation, sequencing, and storage. 384 samples were then normalized
based on a PicoGreen fluorescence assay.
Shallow shotgun metagenome sequencing and diversity analysis
The Illumina data for each HiSeq lane was uploaded to Qiita, a tool with standardized pipelines
for processing and analyzing metagenomic data (Gonzalez et al., 2018). Adapter sequences
were removed from the reads using the Atropos v.1.1.15 (RRID:SCR_023962,
https://github.com/jdidion/atropos) (Didion et al., 2017) command (from the qp-shogun 0.1.5
pipeline) and the trimmed sequences were downloaded from Qiita. The reads for each sample
were filtered of any potential mouse contamination using Bowtie2 v.2-2.2.3
(RRID:SCR_016368, https://bowtie-bio.sourceforge.net/bowtie2/index.shtml) (Langmead and
Salzberg, 2012). The filtered reads were then aligned to the Web of Life (WoL) reference
phylogeny (Zhu et al., 2019) with Bowtie2 using an adapted SHOGUN pipeline (Hillmann et al.,
2018). The WoL contains 10,575 bacterial and archaeal genomes, with each genome
representing an operational taxonomic unit (OTU). Sequencing reads that did not map to a
single reference genome as well as reads that mapped to multiple genomes were not included
in the analysis. If an OTU had a relative abundance less than 0.01% in a given sample, the OTU
was not included for that sample. Additionally, OTUs with fewer than 5 assigned reads were not
considered. The samples were rarefied to a depth of 12,750 reads and those with fewer than
the rarefaction depth were excluded. The QIIME2 v.2019.7 (RRID:SCR_021258,
https://qiime2.org/) (Bolyen et al., 2019) DEICODE plugin was used to calculate the Aitchison
distances, a compositional beta diversity metric, and perform Robust Aitchison PCA to create
biplots that visualize relationships between features and samples (Martino et al., 2019). The
QIIME2 diversity plugin was used to calculate the other alpha- and beta-diversity metrics used
in this study.
Metagenomics-based functional profiling
The filtered reads were also analyzed using HUMAnN2 v2.8.1 (RRID:SCR_016280,
https://huttenhower.sph.harvard.edu/humann2) (Franzosa et al., 2018) to establish functional
profiles for the samples. HUMAnN2 is a pipeline that begins by using MetaPhlAn 2 to compile

custom databases of reference genomes based on the species detected in a sample (Truong et
al., 2015). HUMAnN2 then maps the filtered reads onto these custom databases and the reads
that do not map to any of the references are then subjected to a translated search against
UniProt Reference Clusters or UniRef (RRID:SCR_010646, https://www.uniprot.org/help/uniref)
(Suzek et al., 2007). Here, the UniRef90 database was used for the translated search and
installed according to the HUMAnN2 documentation. The results from both the search
performed using the custom reference genome database and the search against the UniRef90
database were combined and the gene families identified in each sample were reported in units
of reads per kilobase (RPKs) to account for gene length. HUMAnN2 also compared the gene
families found in a sample with the MetaCyc pathways database (RRID:SCR_007778,
https://metacyc.org/) (Caspi et al., 2018) and output a table reporting the pathway abundances
found in each sample. After rarefying gene family tables to a depth of 166,000 RPKs and using
a depth of 22,600 for pathway abundances, the QIIME2 diversity and DEICODE plugins were
used to calculate alpha- and beta-diversity metrics. The metagenomics data from this study are
available from
(https://github.com/mazmanianlab/Griffiths_Yoo_et_al/tree/main/metagenomics).

Metabolomics
Procedures are described in dx.doi.org/10.17504/protocols.io.bzqfp5tn.
Sample preparation
Frozen cecal samples were transported on dry ice for metabolomics analysis. The samples
were weighed and an extraction solvent (1:1 methanol to water with an internal standard of 1
µM sulfamethazine) was added at a 1:10 milligram to microliter ratio. The samples were then
homogenized using a TissueLyser II (Qiagen) for 5 minutes at 25 Hz followed by a 15 min
centrifugation at 14,000 rpm. 120 µL of supernatant was transferred to a 96-well DeepWell plate
(Eppendorf) and lyophilized using a CentriVap Benchtop Vacuum Concentrator (Labconco) and
stored at -80 ºC. At the time of data acquisition, the lyophilized plates were resuspended in a
1:1 methanol to water solvent spiked with 1 µM of sulfadimethoxine. The plates were vortexed
for 2 minutes, centrifuged at 14,000 rpm for 15 min and 120 µL of the supernatant was
transferred to a 96-well autosampler plate (Eppendorf). Plates were stored at 4 °C prior to
LCMS analysis.
Data acquisition
The untargeted metabolomics analysis was completed using an ultra-high performance liquid
chromatography system (Thermo Dionex Ultimate 3000 UHPLC) coupled to an ultra-high
resolution quadrupole time of flight (qTOF) mass spectrometer (Bruker Daltonics MaXis HD). A
Phenomenex Kinetex column (C18 1.7 µm, 2.1 mm x 50 mm) was used for chromatographic
separation. An injection volume of 5 µL was used for each sample and a flow-rate of 0.500 mL
was used throughout the analysis. The mobile phase consisted of solvent A: 100% LC-MS
grade water spiked with 0.1% formic acid and solvent B: 100% LC-MS grade acetonitrile spiked
with 0.1% formic acid. The chromatographic gradient was as follows: 0.0–1.0 min, 5% B; 1.0–
9.0 min, 5–100% B; 9.0-11.0 min, 100% B; 11.0-11.5 min, 100-5% B; 11.5-12.5 min, 5% B. Data
was collected with electrospray ionization in positive mode, and was saved as .d file folders.
Data processing
The raw .d data files were converted to mzXML format using Bruker Compass DataAnalysis 4.1
software. The resulting .mzXML file, the original .d file folders, and basic prep information sheet
are stored in the UC San Diego MassIVE data repository under the accession number
MSV000084550. For MS1 level feature detection, the open-source software MZmine version
2.51 (RRID:SCR_012040, https://mzmine.github.io/) was used. The parameters used are as
follows: 1) Mass Detection (Centroid, Noise Level MS1 1E3, MS2 1E2); 2) ADAP
Chromatogram Builder (Min Group size in # of scans=3, Group Intensity Threshold= 3E3, Min

Highest Intensity=1E3, m/z tolerance 0.01 m/z or 10.0 ppm); 3) Chromatogram Deconvolution
(Local Minimum Search>Chromatographic Threshold 0.01%, Minimum in RT range 0.50 min,
<Minimum Relative Height 0.01%, Minimum Absolute Height 3E3, Min Ratio of Peak Top/Edge
2, Peak Duration Range 0.05-0.50 min; m/z Calculation Auto, m/z range for MS2 pairing 0.01
Da, and RT Range for MS2 Pairing 0.1 min); Isotopic Peaks Grouper (m/z Tolerance 0.01 m/z
or 10.0 ppm, Retention Time Tolerance 0.3 min, Maximum Charge 4, Representative Ion Most
Intense); Join Aligner (m/z Tolerance 0.01 m/z or 10.0 ppm, Weight for m/z 75, Retention Time
Tolerance 0.3 min, Weight for RT 25); Gap-Filling Peak Finder (Intensity Tolerance 20%, m/z
Tolerance 0.005 m/z or 10.0 ppm, Retention Time Tolerance 0.2 min). The resulting feature
table was saved as a .csv file and .mgf file for use in GNPS and MetaboAnalyst.
Molecular networking and statistical analysis
Molecular networking was performed using the feature networking tool available through the
Global Natural Products Social Molecular Networking (GNPS, RRID:SCR_019012,
https://gnps.ucsd.edu/ProteoSAFe/static/gnps-splash.jsp) portal:
https://gnps.ucsd.edu/ProteoSAFe/index.jsp?params=%7B%22workflow%22:%22FEATURE-
BASED-MOLECULAR-
NETWORKING%22,%22library_on_server%22:%22d.speclibs;%22%7D.
The annotations obtained using this workflow fall under MSI level 2 or 3 and were used for
feature analysis (Sumner et al., 2007). Briefly, level 2 compounds are putatively annotated,
meaning they are not identified using chemical reference standards but rather based on
physical properties and/or spectral similarities to available spectral libraries (publicly available
and purchased NIST17 CID). Level 3 compounds are putatively characterized classes of
compounds identified similarly to level 2 compounds. The feature-based molecular networking
workflow on GNPS (Nothias et al., 2019) was utilized in order to analyze the spectra associated
with the feature tables produced using the open source software Mzmine version 2.51
(RRID:SCR_012040, https://mzmine.github.io/) (Pluskal et al., 2010). The .mgf and .csv outputs
from MZmine v2.51 were used to run the workflow. The GNPS workflow parameters used were
as follows: Precursor Ion Mass 0.02 Da, Fragment Ion Mass Tolerance 0.02 Da, Min Pairs Cos
0.7, Minimum Matched Fragments 6, Maximum Shift Between Precursors 500 Da, Network
TopK 10, Maximum Connected Component Size (Beta) 100, and the files were row sum
normalized. Default parameters were used for the rest of the settings. Visualizations and
statistical analyses were performed using QIIME 2 v.2019.10 (RRID:SCR_021258) (Bolyen et
al., 2019), MetaboAnalyst (RRID:SCR_015539, https://www.metaboanalyst.ca/) and Cytoscape
v3.7.2 (RRID:SCR_003032, https://cytoscape.org/). The metabolomics data from this study are
available from
(https://github.com/mazmanianlab/Griffiths_Yoo_et_al/tree/main/metabolomics).
998
Proteome Preparation
Procedures are described in dx.doi.org/10.17504/protocols.io.bzqcp5sw.
Protein extraction
Mice were sacrificed 1 hour after the final C21 administration and cecal contents were isolated
and resuspended in 400 µl of phosphate buffered solution and centrifuged at 20,000 xg to spin
down cells and lysate. Protein was isolated from the resulting supernatant using Wessel-
Flügge's methanol/chloroform extraction method (Wessel and Flügge, 1984). Briefly, MeOH and
chloroform were added to the samples at a 4:1 and 1:1 ratio, respectively. Next, dH$_2$O was
added at a 3:1 ratio, samples were vortexed and centrifuged at 20,000 xg. Resulting
precipitated protein was collected and washed with MeOH. Precipitated protein was centrifuged
and left to air dry, and stored at -20 ºC until protein digestion.
In-solution protein digestion and desalting
Precipitated protein samples were denatured in 40 µL of 8M Urea (100 mM Tris-HCl pH 8.5). To
reduce disulfide bonds, 1.25 µL of 100 mM Tris(2-carboxyethyl)Phosphine was added and

incubated at room temperature (RT) for 20 minutes. Then 1.8 µL of 250 mM iodoacetamide was
added and incubated at RT in the dark to alkylate cysteines. The first step of digestion was
initiated by adding 1 µL of 0.1 µg/µL of lysyl endopeptidase. After 4 hours of incubation, the urea
concentration was adjusted to 2 M by adding 120 µL of 100 mM Tris-HCl pH 8.5. The second
step of digestion was done by adding 2.5 µL of 2µg/µL trypsin plus 1.6 µL of 100 mM $CaCl_2$ and
incubating overnight in the dark. Formic acid was added to stop trypsin digestion. Digested
peptides were desalted by HPLC using a C8 peptide microtrap (Microm Bioresources),
lyophilized, and diluted to 200 ng/µl in 0.2% formic acid prior to LC-MS/MS analysis.

LC-MS/MS
Samples were analyzed on a Q Exactive HF Orbitrap mass spectrometer coupled to an EASY
nLC 1200 liquid chromatographic system (Thermo Scientific, San Jose, CA). Approximately 200
ng of peptides were loaded on a 50 µm I.D. × 25 cm column with a 10 µm electrospray tip
(PicoFrit from New Objective, Woburn, MA) in-house-packed with ReproSil-Pur C18-AQ 1.9 µm
(Dr. Maisch, Ammerbuch, Germany). Solvent A consisted of 2% MeCN in 0.2% FA and solvent
B consisted of 80% MeCN in 0.2% FA. A non-linear 60 minute gradient from 2% B to 40% B
was used to separate the peptides for analysis. The mass spectrometer was operated in a data-
dependent mode, with MS1 scans collected from 400-1650 m/z at 60,000 resolution and MS/MS
scans collected from 200-2000 m/z at 30,000 resolution. Dynamic exclusion of 45 s was used.
The top 12 most abundant peptides with charge states between 2 and 5 were selected for
fragmentation with normalized collision energy of 28.

Peptide and protein identification
Thermo .raw files were converted to .ms1 and .ms2 files using RawConverter 1.1.0.18 (He et
al., 2015) operating in data dependent mode and selecting for monoisotopic m/z. Tandem mass
spectra (.ms2 files) were identified by a database search method using the Integrated
Proteomics Pipeline 6.5.4 (IP2, Integrated Proteomics Applications, Inc.,
http://www.integratedproteomics.com). Briefly, databases containing forward and reverse
(decoy) (Elias and Gygi, 2007; Peng et al., 2003) peptide sequences were generated from *in*
silico trypsin digestion of either the mouse proteome (UniProt; Oct. 2, 2019) or protein
sequences derived from large comprehensive public repositories (ComPIL 2.0) (Park et al.,
2018). Tandem mass spectra were matched to peptide sequences using the
ProLuCID/SEQUEST (1.4) (Xu et al., 2015; 2006) software package. The validity of spectrum-
peptide matches was assessed using the SEQUEST-defined parameters XCorr (cross-
correlation score) and DeltaCN (normalized difference in cross-correlation scores) in the
DTASelect2 (2.1.4) (Cociorva et al., 2007; Tabb et al., 2002) software package. Search settings
were configured as follows: (1) 5 ppm precursor ion mass tolerance, (2) 10 ppm fragment ion
mass tolerance, (3) 1% peptide false discovery rate, (4) 2 peptide per protein minimum, (5) 600-
6000 Da precursor mass window, (6) 2 differential modifications per peptide maximum
(methionine oxidation: M+15.994915 Da), (7) unlimited static modifications per peptide (cysteine
carbamidomethylation: C+57.02146 Da), and (8) the search space included half- and fully-
tryptic (cleavage C-terminal to K and R residues) peptide candidates with unlimited (mouse
database, custom metagenomic shotgun database) or 2 missed cleavage events (ComPIL 2.0).

Differential analysis of detected proteins using peptide-spectrum matches (spectral
counts)
Detected proteins were grouped by sequence similarity into "clusters" using CD-HIT 4.8.1
(RRID:SCR_007105, http://weizhong-lab.ucsd.edu/cd-hit/) (Fu et al., 2012; Li and Godzik, 2006;
Li et al., 2001) at the following similarity cut-offs: 65%, 75%, 85%, and 95%. The following is an
example command line input: "cd-hit -i fastafile.fasta -o outputfile -c 0.65 -g 1 -d 0". Tandem
mass spectra identified as peptides (peptide spectrum matches, PSMs) were mapped to CD-
HIT generated clusters. PSMs mapping to >1 cluster were discarded. Cluster-PSM tables were
generated and differential analysis was performed in DESeq2 (1.25.13, RRID:SCR_015687,
https://bioconductor.org/packages/release/bioc/html/DESeq2.html) (Love et al., 2014). Briefly,

count data (PSMs) were modeled using the negative binomial distribution, and the mean-
variance relationship was estimated. Variance was estimated using an information sharing
approach whereby a single feature's (or cluster's) variance was estimated by taking into account
variances of other clusters measured in the same experiment. Feature significance calling and
ranking were performed using estimated effect sizes. Multiple testing correction was performed
by the Benjamini-Hochberg method within the DESeq2 package. Volcano plots were generated
in Prism (GraphPad).
Differential analysis of detected proteins using ion intensity (precursor intensity)
Detected proteins were grouped into "clusters" by sequence similarity using CD-HIT 4.8.1 (Fu et
al., 2012; Li and Godzik, 2006; Li et al., 2001) at the following similarity cut-offs: 65%, 75%,
85%, and 95%. The following is an example command line input: "cd-hit -i fastafile.fasta -o
outputfile -c 0.65 -g 1 -d 0". Using the Census software package (Park et al.; 2008) (Integrated
Proteomics Pipeline 6.5.4), peptide ion intensities were calculated from .ms1 files. Peptide ion
intensities were assigned to their parent peptide, then parent peptides were mapped to their
appropriate CD-HIT generated clusters. Ion intensities belonging to parent peptides that
mapped to >1 CD-HIT cluster were discarded. Cluster-ion intensity tables were generated.
1080
Ion intensity data were analyzed using the Differential Enrichment analysis of Proteomics data
DEP package (RRID:SCR_023090,
https://bioconductor.org/packages/release/bioc/html/DEP.html) (Zhang et al., 2018) operating in
R. Intensity values were automatically Log2 transformed in DEP. The cluster list was
subsequently filtered with the 'filter_proteins' function such that clusters with missing values
above a 65% threshold were discarded. Remaining intensities were further transformed by the
'normalize_vsn' function (Huber et al., 2002). Missing data in remaining clusters were imputed
using a mixed approach. Clusters where either the control or treatment group contained only
null entries were classified as 'missing not at random' (MNAR) and imputed with 0 values. All
other groups were treated as 'missing at random' (MAR) and imputed using the maximum
likelihood method ('MLE') (Gatto and Lilley, 2012). Note that for a given cluster, missing values
for treatment groups were imputed separately by treatment group. Differential expression
analyses were performed on filled-in cluster-ion intensity tables using the 'test_diff' function
(Ritchie et al., 2015) and multiple testing correction was performed using the 'add_rejections'
function.
Network analysis using the STRING database
Upregulated proteins with a nominal p-value < 0.2 were searched against protein-protein
interactions in the STRING database (RRID:SCR_005223, http://www.string-db.org) where high
confidence interactions are selected for. Briefly, the STRING database sources protein-protein
interactions from primary databases consisting of genomic context predictions, high-throughput
lab experiments, (conserved) co-expression, automated text mining, and previous knowledge in
databases (Szklarczyk et al., 2019).
Metaproteome analysis using Unipept
Upregulated tryptic, microbial peptide sequences, with fold change and nominal p-value cutoffs
of >2 and <0.2, respectively, were input into Unipept (http://unipept.ugent.be) (Gurdeep Singh et
al., 2019; Mesuere et al., 2015), equating leucine and isoleucine and filtering duplicate peptides.
Briefly, Unipept indexes tryptic peptide sequences from the UniProtKB and details peptides with
NCBI's taxonomic database. Lowest common ancestor was calculated for each tryptic peptide.
The proteomics data from this study are available from
(https://github.com/mazmanianlab/Griffiths_Yoo_et_al/tree/main/proteomics).
1111
3' mRNA sequencing
Procedures are described in dx.doi.org/10.17504/protocols.io.bzqbp5sn.
Tissue collection and RNA extraction

Mice were cervically dislocated and the GI tract was removed. 1 cm of tissue above and below
the cecum were dissected and cleaned to represent tissue from the distal SI and proximal colon,
respectively. Tissue was homogenized in TRIzol (ThermoFisher Scientific, Waltham. MA- Cat.
No. 15596018) solution using a bead-based homogenizing method, and total RNA was
extracted using chloroform per manufacturer's instructions.
Library preparation, sequencing, and analysis
The cDNA libraries were prepared using the QuantSeq 3'mRNA-Seq Library Prep Kit FWD for
Illumina (Lexogen, Greenland, NH) supplemented with UMI (unique molecular index) as per the
manufacturer's instructions. Briefly, total RNA was reverse transcribed using oligo (dT) primers.
The second cDNA strand was synthesized by random priming, in a manner such that DNA
polymerase is efficiently stopped when reaching the next hybridized random primer, so only the
fragment closest to the 3' end gets captured for later indexed adapter ligation and PCR
amplification. UMIs were incorporated to the first 6 bases of each read, followed by 4 bases of
spacer sequences. UMIs are used to eliminate possible PCR duplicates in sequencing datasets
and therefore facilitate unbiased gene expression profiling. The basic principle behind the UMI
deduplication step is to collapse reads with identical mapping coordinates and UMI sequences.
This step helps increase the accuracy of sequencing read counts for downstream analysis of
gene expression levels. The processed libraries were assessed for size distribution and
concentration using the Bioanalyzer High Sensitivity DNA Kit (Agilent Technologies, Santa
Clara, CA- Cat. No. 5067-4626 and -4627). Pooled libraries were diluted to 2 nM in EB buffer
(Qiagen, Hilden, Germany, Cat. No. 19086) and then denatured using the Illumina protocol. The
libraries were pooled and diluted to 2 nM using 10 mM Tris-HCl, pH 8.5 and then denatured
using the Illumina protocol. The denatured libraries were diluted to 10 pM with pre-chilled
hybridization buffer and loaded onto an Illumina MiSeq v3 flow cell for 150 cycles using a single-
read recipe according to the manufacturer's instructions. Single-end 75 bp reads (max 4.5M
reads) were obtained. De-multiplexed sequencing reads were generated using Illumina
BaseSpace.
1142
UMI specific workflows that were developed and distributed by Lexogen were used to extract
reads that are free from PCR artifacts (i.e., deduplication). First, the umi2index tool was used to
add the 6 nucleotide UMI sequence to the identifier of each read and trim the UMI from the start
of each read. This generated a new FASTQ file, which was then processed through trimming
and alignment. Second, after the quality and polyA trimming by BBDuk (Bestus Bioinformaticus
Duk, RRID:SCR_016969, https://jgi.doe.gov/data-and-tools/bbtools/bb-tools-user-guide/bbduk-
guide/) and alignment by HISAT2 (version 2.1.0, RRID:SCR_015530,
https://daehwankimlab.github.io/hisat2/) (Kim et al., 2015), the mapped reads were collapsed
according to the UMI sequence of each read. Reads were collapsed if they had the same
mapping coordinates (CIGAR string) and identical UMI sequences. Collapsing reads in this
manner removes PCR duplicates. Read counts were calculated using HTSeq
(RRID:SCR_005514, https://htseq.readthedocs.io/en/release_0.9.1/) (Anders et al., 2015) by
supplementing Ensembl gene annotation (GRCm38.78). Raw read counts were run through
ShinySeq to obtain differentially expressed genes and downstream gene ontology analyses
(Sundararajan et al., 2019). The RNAseq data from this study are available from
(https://github.com/mazmanianlab/Griffiths_Yoo_et_al/tree/main/RNAseq).
1159
Whole Gut Transit Time, Fecal Water Content, and Fecal Output
Procedures are described in dx.doi.org/10.17504/protocols.io.36wgq3p1xlk5/v1.
6% (w/v) carmine red (Sigma-Aldrich, St. Louis, MO) with 0.5% methylcellulose (Sigma-Aldrich)
was dissolved and autoclaved prior to use. ChAT-Cre and TH-Cre mice were administered C21
(3 mg/kg) intraperitoneally, and subsequently orally gavaged with 150 µL of carmine red
solution. Mice were single-housed with no bedding for the duration of the experiment, and

animals were not fasted beforehand. Over the 5 hours following gavage, the time of expulsion
was recorded for each fecal pellet, and each pellet was collected in pre-weighed, 1.5 mL
microcentrifuge tube. Each pellet collected was checked for the presence of carmine red, and
the time of initial carmine red pellet expulsion was recorded as GI transit time. The mass of
collected fecal pellets was determined, and pellets were left to dry in an 80 ºC oven for 2 days
before weighing the desiccated pellets and calculating the pellets' initial water content. Fecal
output rate for each mouse was calculated as the total number of pellets expelled during the 5
hour time course post-C21 administration divided by the time the last fecal pellet was expelled.
1174
Colonic Migrating Motor Complexes in *Ex Vivo* Intestinal Preparations
Procedures are described in dx.doi.org/10.17504/protocols.io.n92ldm61nl5b/v1.
Intact colons were dissected from cervically-dislocated ChAT-Cre and TH-Cre mice, flushed and
placed in pre-oxygenated (95% O_2, 5% CO_2) Krebs-Henseleit solution at RT. Proximal and
distal ends were cannulated to 2 mm diameter tubes and secured in the center of an organ bath
with continuously oxygenated Krebs-Henseleit solution at 37 ºC. Syringe pumps were
connected to the inlet and outlet tubes to maintain a flow of solution at a rate of 500 µL/min
through the colon. The system was allowed to equilibrate for 30 minutes before recording.
Baseline recordings were taken for 30 minutes, then the Krebs solution in the organ bath was
briefly removed, mixed with C21 to a final concentration of 2 µM, and replaced in the organ
bath. Recordings were taken for another 30 minutes.
1186

REFERENCES

Alavi, S., Mitchell, J.D., Cho, J.Y., Liu, R., Macbeth, J.C., and Hsiao, A. (2020). Interpersonal Gut Microbiome Variation Drives Susceptibility and Resistance to Cholera Infection. Cell *181*, 1533–1546.e13.

Albenberg, L.G., and Wu, G.D. (2014). Diet and the intestinal microbiome: associations, functions, and implications for health and disease. Gastroenterology *146*, 1564–1572.

Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq--a Python framework to work with high-throughput sequencing data. Bioinformatics *31*, 166–169.

Anlauf, M., Schäfer, M.K.H., Eiden, L., and Weihe, E. (2003). Chemical coding of the human gastrointestinal nervous system: cholinergic, VIPergic, and catecholaminergic phenotypes. The Journal of Comparative Neurology *459*, 90–111.

Aries, V., Crowther, J.S., Drasar, B.S., and Hill, M.J. (1969). Degradation of bile salts by human intestinal bacteria. Gut *10*, 575–576.

Bahrami, S., and Drabløs, F. (2016). Gene regulation in the immediate-early response process. Advances in Biological Regulation *62*, 37–49.

Baumuratov, A. S., Antony, P. M. A., Ostaszewski, M., He, F., Salamanca, L., Antunes, L., Weber, J., Longhino, L., Derkinderen, P., Koopman, W. J. H., Diederich, N. J. (2016). Enteric neurons from Parkinson's disease patients display ex vivo aberrations in mitochondrial structure. Scientific Reports *6*.

Begley, M., Sleator, R.D., Gahan, C.G.M., and Hill, C. (2005). Contribution of three bile-associated loci, bsh, pva, and btlB, to gastrointestinal persistence and bile tolerance of Listeria monocytogenes. Infect. Immun. *73*, 894–904.

Benskey, M.J., Kuhn, N.C., Galligan, J.J., Garcia, J., Boye, S.E., Hauswirth, W.W., Mueller, C., Boye, S.L., and Manfredsson, F.P. (2015). Targeted gene delivery to the enteric nervous system using AAV: a comparison across serotypes and capsid mutants. Mol. Ther. *23*, 488–500.

Bhavsar, A.S., Verma, S., Lamba, R., Lall, C.G., Koenigsknecht, V., and Rajesh, A. (2013). Abdominal manifestations of neurologic disorders. Radiographics *33*, 135–153.

Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F., et al. (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol *37*, 852–857.

Bravo, J.A., Forsythe, P., Chew, M.V., Escaravage, E., Savignac, H.M., Dinan, T.G., Bienenstock, J., and Cryan, J.F. (2011). Ingestion of Lactobacillus strain regulates emotional behavior and central GABA receptor expression in a mouse via the vagus nerve. Proceedings of the National Academy of Sciences *108*, 16050–16055.

Browning, K.N. and Travagli R.A. (2014). Central nervous system control of gastrointestinal motility and secretion and modulation of gastrointestinal functions. Compr Physiol. *4*, 1339-68.

Bruel-Jungerman, E., Davis, S., and Laroche, S. (2016). Brain Plasticity Mechanisms and Memory: A Party of Four. Neuroscientist *13*, 492–505.

Carabotti, M., Scirocco, A., Maselli, M.A., and Severi, C. (2015). The gut-brain axis: interactions between enteric microbiota, central and enteric nervous systems. Ann Gastroenterol *28*, 203–209.

Carroll, I.M., Ringel-Kulka, T., Ferrier, L., Wu, M.C., Siddle, J.P., Bueno, L., and Ringel, Y.
(2013). Fecal protease activity is associated with compositional alterations in the intestinal
microbiota. PloS One *8*, e78017.

Caspi, R., Billington, R., Fulcher, C.A., Keseler, I.M., Kothari, A., Krummenacker, M.,
Latendresse, M., Midford, P.E., Ong, Q., Ong, W.K., et al. (2018). The MetaCyc database of
metabolic pathways and enzymes. Nucleic Acids Res *46*, D633–D639.

Cekanaviciute, E., Yoo, B.B., Runia, T.F., Debelius, J.W., Singh, S., Nelson, C.A., Kanner, R.,
Bencosme, Y., Lee, Y.K., Hauser, S.L., et al. (2017). Gut bacteria from multiple sclerosis
patients modulate human T cells and exacerbate symptoms in mouse models. Proc. Natl. Acad.
Sci. U.S.a. *114*, 10713–10718.

Cersosimo, M.G., Raina, G.B., Pecci, C., Pellene, A., Calandra, C.R., Gutiérrez, C., Micheli,
F.E., and Benarroch, E.E. (2013). Gastrointestinal manifestations in Parkinson's disease:
prevalence and occurrence before motor symptoms. J. Neurol. *260*, 1332–1338.

Challis, R.C., Ravindra Kumar, S., Chan, K.Y., Challis, C., Beadle, K., Jang, M.J., Kim, H.M.,
Rajendran, P.S., Tompkins, J.D., Shivkumar, K., et al. (2019). Systemic AAV vectors for
widespread and targeted gene delivery in rodents. Nature Protocols *14*, 379–414.

Chan, K.Y., Jang, M.J., Yoo, B.B., Greenbaum, A., Ravi, N., Wu, W.-L., Sánchez-Guardado, L.,
Lois, C., Mazmanian, S.K., Deverman, B.E., et al. (2017). Engineered AAVs for efficient
noninvasive gene delivery to the central and peripheral nervous systems. Nat. Neurosci. *20*,
1172–1179.

Chen, T.-W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R.,
Kerr, R.A., Orger, M.B., Jayaraman, V., et al. (2013). Ultrasensitive fluorescent proteins for
imaging neuronal activity. Nature *499*, 295–300.

Cociorva, D., L Tabb, D., and Yates, J.R. (2007). Validation of Tandem Mass Spectrometry
Database Search Results Using DTASelect. Current Protocols in Bioinformatics *16*.

Costello, M., Fleharty, M., Abreu, J., Farjoun, Y., Ferriera, S., Holmes, L., Granger, B., Green,
L., Howd, T., Mason, T., et al. (2018). Characterization and remediation of sample index swaps
by non-redundant dual indexing on massively parallel sequencing platforms. BMC Genomics
19, 332–10.

de Aguiar Vallim, T.Q., Tarling, E.J., and Edwards, P.A. (2013). Pleiotropic roles of bile acids in
metabolism. Cell Metabolism *17*, 657–669.

Del Giudice, E., Staiano, A., Capano, G., Romano, A., Florimonte, L., Miele, E., Ciarla, C.,
Campanozzi, A., and Crisanti, A.F. (1999). Gastrointestinal manifestations in children with
cerebral palsy. Brain Dev. *21*, 307–311.

Delpino, M.V., Marchesini, M.I., Estein, S.M., Comerci, D.J., Cassataro, J., Fossati, C.A., and
Baldi, P.C. (2007). A bile salt hydrolase of Brucella abortus contributes to the establishment of a
successful infection through the oral route in mice. Infect. Immun. *75*, 299–305.

Deneux, T., Kaszas, A., Szalay, G., Katona, G., Lakner, T., Grinvald, A., Rózsa, B., and
Vanzetta, I. (2016). Accurate spike estimation from noisy calcium signals for ultrafast three-
dimensional imaging of large neuronal populations in vivo. Nature Communications *7*, 12190–
17.

Derrien, M., Vaughan, E.E., Plugge, C.M., and de Vos, W.M. (2004). Akkermansia muciniphila
gen. nov., sp. nov., a human intestinal mucin-degrading bacterium. International Journal of
Systematic and Evolutionary Microbiology *54*, 1469–1476.

Deverman, B.E., Pravdo, P.L., Simpson, B.P., Kumar, S.R., Chan, K.Y., Banerjee, A., Wu, W.-L., Yang, B., Huber, N., Pasca, S.P., et al. (2016). Cre-dependent selection yields AAV variants for widespread gene transfer to the adult brain. Nat Biotechnol *34*, 204–209.

Didion, J.P., Martin, M., and Collins, F.S. (2017). Atropos: specific, sensitive, and speedy trimming of sequencing reads. PeerJ *5*, e3720.

Donaldson, G.P., Ladinsky, M.S., Yu, K.B., Sanders, J.G., Yoo, B.B., Chou, W.C., Conner, M.E., Earl, A.M., Knight, R., Bjorkman, P.J., et al. (2018). Gut microbiota utilize immunoglobulin A for mucosal colonization. Science *360*, 795–800.

Donaldson, G.P., Lee, S.M., and Mazmanian, S.K. (2015). Gut biogeography of the bacterial microbiota. Nat Rev Micro *14*, 20–32.

Donowitz, M., Singh, S., Salahuddin, F.F., Hogema, B.M., Chen, Y., Gucek, M., Cole, R.N., Ham, A., Zachos, N.C., Kovbasnjuk, O., et al. (2007). Proteome of murine jejunal brush border membrane vesicles. J. Proteome Res. *6*, 4068–4079.

Drokhlyansky, E., Smillie, C.S., Van Wittenberghe, N., Ericsson, M., Griffin, G.K., Eraslan, G., Dionne, D., Cuoco, M.S., Goder-Reiser, M.N., Sharova, T., et al. (2020). The Human and Mouse Enteric Nervous System at Single-Cell Resolution. Cell *182*, 1606–1622.e1623.

Elias, J.E., and Gygi, S.P. (2007). Target-decoy search strategy for increased confidence in large-scale protein identifications by mass spectrometry. Nat. Methods *4*, 207–214.

Everard, A., Belzer, C., Geurts, L., Ouwerkerk, J.P., Druart, C., Bindels, L.B., Guiot, Y., Derrien, M., Muccioli, G.G., Delzenne, N.M., et al. (2013). Cross-talk between Akkermansia muciniphila and intestinal epithelium controls diet-induced obesity. Proc. Natl. Acad. Sci. U.S.a. *110*, 9066–9071.

Finneran, D.J., Njoku, I.P., Flores-Pazarin, D., Ranabothu, M. R., Nash, K.R., Morgan, D., and Gordon, M.N. (2021). Towards Development of Neuron Specific Transduction After Systemic Delivery of Viral Vectors. Frontiers in Neurology *12*, 685802.

Fiorucci, S., Biagioli, M., Zampella, A., and Distrutti, E. (2018). Bile Acids Activated Receptors Regulate Innate Immunity. Frontiers in Immunology *9*, eaao1605–eaao1617.

Flandez, M., Guilmeau, S., Blache, P., and Augenlicht, L.H. (2008). KLF4 regulation in intestinal epithelial cell maturation. Experimental Cell Research *314*, 3712–3723.

Foong, J.P.P., Tough, I.R., Cox, H.M., and Bornstein, J.C. (2014). Properties of cholinergic and non-cholinergic submucosal neurons along the mouse colon. The Journal of Physiology *592*, 777–793.

Franzosa, E.A., McIver, L.J., Rahnavard, G., Thompson, L.R., Schirmer, M., Weingart, G., Lipson, K.S., Knight, R., Caporaso, J.G., Segata, N., et al. (2018). Species-level functional profiling of metagenomes and metatranscriptomes. Nat. Methods *15*, 962–968.

Fried, D.E., and Gulbransen, B.D. (2015). In situ Ca2+ imaging of the enteric nervous system. JoVE.

Fu, L., Niu, B., Zhu, Z., Wu, S., and Li, W. (2012). CD-HIT: accelerated for clustering the next-generation sequencing data. Bioinformatics *28*, 3150–3152.

Fung, C., Koussoulas, K., Unterweger, P., Allen, A. M., Bornstein, J. C., Foong, J. P. P. (2018). Cholinergic Submucosal Neurons Display Increased Excitability Following in Vivo Cholera Toxin Exposure in Mouse Ileum. Sec. Gastrointestinal Sciences 9.

Furness, J.B. (2012). The enteric nervous system and neurogastroenterology. Nature Reviews Gastroenterology and Hepatology *9*, 286–294.

Furness, J.B., Callaghan, B.P., Rivera, L.R., and Cho, H.-J. (2014). The Enteric Nervous System and Gastrointestinal Innervation: Integrated Local and Central Control. In The Enteric Nervous System, (New York, NY: Springer New York), pp. 39–71.

Furness, J.B. (2006). The Enteric Nervous System (Wiley-Blackwell).

Gabanyi, I., Muller, P.A., Feighery, L., Oliveira, T.Y., Costa-Pinto, F.A., and Mucida, D. (2016). Neuro-immune Interactions Drive Tissue Programming in Intestinal Macrophages. Cell *164*, 378–391.

Gatto, L., and Lilley, K.S. (2012). MSnbase-an R/Bioconductor package for isobaric tagged mass spectrometry data visualization, processing and quantitation. Bioinformatics *28*, 288–289.

Gershon, M.D. (2015). The Enteric Nervous System: A Second Brain. Hospital Practice *34*, 31–52.

Glenn, T.C., Nilsen, R.A., Kieran, T.J., Sanders, J.G., Bayona-Vásquez, N.J., Finger, J.W., Pierson, T.W., Bentley, K.E., Hoffberg, S.L., Louha, S., et al. (2019). Adapterama I: universal stubs and primers for 384 unique dual-indexed or 147,456 combinatorially-indexed Illumina libraries (iTru & iNext). PeerJ *7*, e7755.

Gonzalez, A., Navas-Molina, J.A., Kosciolek, T., McDonald, D., Vázquez-Baeza, Y., Ackermann, G., DeReus, J., Janssen, S., Swafford, A.D., Orchanian, S.B., et al. (2018). Qiita: rapid, web-enabled microbiome meta-analysis. Nat. Methods *15*, 796–798.

Gottlieb, A., and Canbay, A. (2019). Why Bile Acids Are So Important in Non-Alcoholic Fatty Liver Disease (NAFLD) Progression. Cells *8*.

Grundy, D., and Brookes, S. (2011). Neural Control of Gastrointestinal Function. Colloquium Series on Integrated Systems Physiology: From Molecule to Function *3*, 1–134.

Gurdeep Singh, R., Tanca, A., Palomba, A., Van der Jeugt, F., Verschaffelt, P., Uzzau, S., Martens, L., Dawyndt, P., and Mesuere, B. (2019). Unipept 4.0: Functional Analysis of Metaproteome Data. J. Proteome Res. *18*, 606–615.

Haenraets, K., Foster, E., Johannssen, H., Kandra, V., Frezel, N., Steffen, T., Jaramillo, V., Paterna, J.-C., Zeilhofer, H.U., and Wildner, H. (2017). Spinal nociceptive circuit analysis with recombinant adeno-associated viruses: the impact of serotypes and promoters. Journal of Neurochemistry *142*, 721–733.

Hama, H., Hioki, H., Namiki, K., Hoshida, T., Kurokawa, H., Ishidate, F., Kaneko, T., Akagi, T., Saito, T., Saido, T., et al. (2015). ScaleS: an optical clearing palette for biological imaging. Nat. Neurosci. *18*, 1518–1529.

Hamnett, R., Dershowitz, L.B., Sampathkumar, V., Wang, Z., Gomez-Frittelli, J., De Andrade, V., Kasthuri, N., Druckmann, S., Kaltschmidt, J.A. (2022). Regional cytoarchitecture of the adult and developing mouse enteric nervous system. Curr. Biol. *32*, 1–10.

Hansen, M.B. (2003). The enteric nervous system II: Gastrointestinal functions. Pharmacol. Toxicol. *92*, 249–257.

He, L., Diedrich, J., Chu, Y.-Y., and Yates, J.R. (2015). Extracting Accurate Precursor Information for Tandem Mass Spectra by RawConverter. Anal. Chem. *87*, 11361–11367.

Hennig, G.W., Gould, T.W., Koh, S.D., Corrigan, R.D., Heredia, D.J., Shonnard, M.C., and Smith, T.K. (2015). Use of Genetically Encoded Calcium Indicators (GECIs) Combined with

Advanced Motion Tracking Techniques to Examine the Behavior of Neurons and Glia in the Enteric Nervous System of the Intact Murine Colon. Front. Cell. Neurosci. *9*, 13819.

Hillmann, B., Al-Ghalith, G.A., Shields-Cutler, R.R., Zhu, Q., Gohl, D.M., Beckman, K.B., Knight, R., and Knights, D. (2018). Evaluating the Information Content of Shallow Shotgun Metagenomics. mSystems *3*, 457.

Hofmann, A.F., and Eckmann, L. (2006). How bile acids confer gut mucosal protection against bacteria. Proceedings of the National Academy of Sciences *103*, 4333–4334.

Huber, W., Heydebreck, von, A., Sültmann, H., Poustka, A., and Vingron, M. (2002). Variance stabilization applied to microarray data calibration and to the quantification of differential expression. Bioinformatics *18 Suppl 1*, S96–S104.

Hyland, N.P., and Cryan, J.F. (2010). A Gut Feeling about GABA: Focus on GABA(B) Receptors. Front Pharmacol *1*, 124.

Ichikawa, R., Takayama, T., Yoneno, K., Kamada, N., Kitazume, M.T., Higuchi, H., Matsuoka, K., Watanabe, M., Itoh, H., Kanai, T., et al. (2012). Bile acids induce monocyte differentiation toward interleukin-12 hypo-producing dendritic cells via a TGR5-dependent pathway. Immunology *136*, 153–162.

Jackson, K.L., Dayton, R.D., Deverman, B.E., and Klein, R.L. (2016). Better Targeting, Better Efficiency for Wide-Scale Neuronal Transduction with the Synapsin Promoter and AAV-PHP.B. Front. Mol. Neurosci. *9*, 116.

Jakob, MO., Kofoed-Branzk, M., Deshpande, D., Murugan, S., and Klose, C.S.N. An integrated view on neuronal subsets in the peripheral nervous system and their role in immunoregulation. Front. Immunol. *12*, 679055.

Jangi, S., Gandhi, R., Cox, L.M., Li, N., Glehn, von, F., Yan, R., Patel, B., Mazzola, M.A., Liu, S., Glanz, B.L., et al. (2016). Alterations of the human gut microbiome in multiple sclerosis. Nature Communications *7*, 12015–11.

Jarret, A., Jackson, R., Duizer, C., Healy, M.E., Zhao, J., Rone, J.M., Bielecki, P., Sefik, E., Roulis, M., Rice, T., et al. (2020). Enteric Nervous System-Derived IL-18 Orchestrates Mucosal Barrier Immunity. Cell *180*, 50–63.e12.

Jia, L., Betters, J.L., and Yu, L. (2011). Niemann-pick C1-like 1 (NPC1L1) protein in intestinal and hepatic cholesterol transport. Annu. Rev. Physiol. *73*, 239–259.

Johnson, C.D., Barlow-Anacker, A.J., Pierre, J.F., Touw, K., Erickson, C.S., Furness, J.B., Epstein, M.L., and Gosain, A. (2018). Deletion of choline acetyltransferase in enteric neurons results in postnatal intestinal dysmotility and dysbiosis. The FASEB Journal *32*, 4744–4752.

Jones, B.V., Begley, M., Hill, C., Gahan, C.G.M., and Marchesi, J.R. (2008). Functional and comparative metagenomic analysis of bile salt hydrolase activity in the human gut microbiome. Proc. Natl. Acad. Sci. U.S.a. *105*, 13580–13585.

Kaelberer, M.M., Buchanan, K.L., Klein, M.E., Barth, B.B., Montoya, M.M., Shen, X., and Bohórquez, D.V. (2018). A gut-brain neural circuit for nutrient sensory transduction. Science *361*, eaat5236.

Kaestner, C.L., Smith, E.H., Peirce, S.G., and Hoover, D.B. (2019). Immunohistochemical analysis of the mouse celiac ganglion: An integrative relay station of the peripheral nervous system. J. Comp. Neurol. *16*, 2742-2760.

Kim, D., Langmead, B., and Salzberg, S.L. (2015). HISAT: a fast spliced aligner with low memory requirements. Nat. Methods *12*, 357–360.

Kirwan, W.O., Smith, A.N., Mitchell, W.D., Falconer, J.D., and Eastwood, M.A. (1975). Bile acids
and colonic motility in the rabbit and the human. Gut *16*, 894–902.

Kotterman, M.A., and Schaffer, D.V. (2014). Engineering adeno-associated viruses for clinical
gene therapy. Nat Rev Genet *15*, 445–451.

Kügler, S., Kilic, E., and Bähr, M. (2003). Human synapsin 1 gene promoter confers highly
neuron-specific long-term transgene expression from an adenoviral vector in the adult rat brain
depending on the transduced area. Gene Ther *10*, 337–347.

Lai, N.Y., Musser, M.A., Pinho-Ribeiro, F.A., Baral, P., Jacobson, A., Ma, P., Potts, D.E., Chen,
Z., Paik, D., Soualhi, S., et al. (2019). Gut-Innervating Nociceptor Neurons Regulate Peyer's
Patch Microfold Cells and SFB Levels to Mediate Salmonella Host Defense. Cell *180*, 33–
49.e22.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat.
Methods *9*, 357–359.

Latgé, J.P. (2007). The cell wall: a carbohydrate armour for the fungal cell. Molecular
Microbiology *66*, 279–290.

Li, W., and Godzik, A. (2006). Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. Bioinformatics *22*, 1658–1659.

Li, W., Jaroszewski, L., and Godzik, A. (2001). Clustering of highly homologous sequences to
reduce the size of large protein databases. Bioinformatics *17*, 282–283.

Li, W., Yu, G., Liu, Y., and Sha, L. (2019). Intrapancreatic Ganglia and Neural Regulation of
Pancreatic Endocrine Secretion. Front. Neurosci *13,* 21.

Li, Z., Chalazonitis, A., Huang, Y.Y., Mann, J.J., Margolis, K.G., Yang, Q.M., Kim, D.O., Cote,
F., Mallet, J., and Gershon, M.D. (2011). Essential Roles of Enteric Neuronal Serotonin in
Gastrointestinal Motility and the Development/Survival of Enteric Dopaminergic Neurons.
Journal of Neuroscience *31*, 8998–9009.

Lindeberg, J., Usoskin, D., Bengtsson, H., Gustafsson, A., Kylberg, A., Söderström, S., and
Ebendal, T. (2004). Transgenic expression of Cre recombinase from the tyrosine hydroxylase
locus. Genesis *40*, 67–73.

Long, S.L., Gahan, C.G.M., and Joyce, S.A. (2017). Interactions between gut bacteria and bile
in health and disease. Mol. Aspects Med. *56*, 54–65.

Lott, E. L. & Jones, E. B. (2022). Cholinergic Toxicity. StatPearls.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 1–21.

Marotz, C., Amir, A., Humphrey, G., Gaffney, J., Gogul, G., and Knight, R. (2017). DNA
extraction for streamlined metagenomics of diverse environmental samples. Biotechniques. *62*,
290-293.

Martino, C., Morton, J.T., Marotz, C.A., Thompson, L.R., Tripathi, A., Knight, R., and Zengler, K.
(2019). A Novel Sparse Compositional Technique Reveals Microbial Perturbations. mSystems
4, 813.

Matheis, F., Muller, P.A., Graves, C.L., Gabanyi, I., Kerner, Z.J., Costa-Borges, D., Ahrends, T.,
Rosenstiel, P., and Mucida, D. (2020). Adrenergic Signaling in Muscularis Macrophages Limits
Infection-Induced Neuronal Loss. Cell *180*, 64–78.e16.

Matsushima, S., Hori, S., and Matsuda, M. (1986). Conversion of 4-aminobutyraldehyde to
gamma-aminobutyric acid in striatum treated with semicarbazide and kainic acid. Neurochem.
Res. *11*, 1313–1319.

McConnell, R.E., Benesh, A.E., Mao, S., Tabb, D.L., and Tyska, M.J. (2011). Proteomic
analysis of the enterocyte brush border. American Journal of Physiology - Gastrointestinal and
Liver Physiology *300*, G914–G926.

McMillin, M., and DeMorrow, S. (2016). Effects of bile acids on neurological function and
disease. Faseb J. *30*, 3658–3668.

McQuade, R. M., Singleton, L. M., Wu, H., Lee, S., Constable, R., Di Natale, M., Ringuet, M. T.,
Berger, J. P., Kauhausen, J., Parish, C. L., Finkelstein, D. I., Furness, J. B., Diwakarla, S.
(2021). The association of enteric neuropathy with gut phenotypes in acute and progressive
models of Parkinson's disease. Scientific Reports *11*.

Mesuere, B., Debyser, G., Aerts, M., Devreese, B., Vandamme, P., and Dawyndt, P. (2015).
The Unipept metaproteomics analysis pipeline. Proteomics *15*, 1437–1442.

Miano, J.M., Miano, J.M., Vlasic, N., Vlasic, N., TOTA, R.R., Tota, R.R., STEMERMAN, M.B.,
and Stemerman, M.B. (1993). Smooth muscle cell immediate-early gene and growth factor
activation follows vascular injury. A putative in vivo mechanism for autocrine growth.
Arterioscler. Thromb. *13*, 211–219.

Mittal, R., Debs, L.H., Patel, A.P., Nguyen, D., Patel, K., O'Connor, G., Grati, M., Mittal, J., Yan,
D., Eshraghi, A.A., et al. (2017). Neurotransmitters: The Critical Modulators Regulating Gut-
Brain Axis. J. Cell. Physiol. *232*, 2359–2372.

Mohanta, S.K., Yin, C., Weber, C., Godinho-Silva, C., Veiga-Fernandes, H., Xu, Q.J., Chang,
R.B., Habenicht, A.J.R. (2023). Cardiovascular Brain Circuits. Circulation Research *11*, 1546-
1565.

Monane, M., Avorn, J., Beers, M.H., Everitt, D.E. (1993). Anticholinergic drug use and bowel
function in nursing home patients. Arch. Intern. Med. *153*, 633-638.

Moran, G.W., Leslie F.C., Levison, S.E., Worthington, J., and McLaughlin, J.T. (2008).
Enteroendocrine cells: neglected players in gastrointestinal disorders? Therap. Adv.
Gastroenterol. *1*, 51-60.

Muller, P.A., Koscsó, B., Rajani, G.M., Stevanovic, K., Berres, M.-L., Hashimoto, D., Mortha, A.,
Leboeuf, M., Li, X.-M., Mucida, D., et al. (2014). Crosstalk between Muscularis Macrophages
and Enteric Neurons Regulates Gastrointestinal Motility. Cell *158*, 1210.

Niesler, B., Kuerten, S., Demir, I.E., and Schäfer, K.H. (2021). Disorders of the enteric nervous
system - a holistic view. Nat Rev Gastroenterol Hepatol. *18*, 393-410.

Nezami, B.G. & Srinivasan, S. (2013). Enteric Nervous System in the Small Intestine:
Pathophysiology and Clinical Implications. Current Gastroenterology Reports *12,* 258-365.

Nishiyama, H., Nagai, T., Kudo, M., Okazaki, Y., Azuma, Y., Watanabe, T., Goto, S., Ogata, H.,
and Sakurai, T. (2018). Supplementation of pancreatic digestive enzymes alters the composition
of intestinal microbiota in mice. Biochem. Biophys. Res. Commun. *495*, 273–279.

Nothias, L.F., Petras, D., Schmid, R., Dührkop, K., Rainer, J., Sarvepalli, A., Protsyuk, I., Ernst,
M., Tsugawa, H., Fleischauer, M., et al. (2019). Feature-based Molecular Networking in the
GNPS Analysis Environment. bioRxiv *38*, 812404.

Olson, C.A., Vuong, H.E., Yano, J.M., Liang, Q.Y., Nusbaum, D.J., and Hsiao, E.Y. (2018). The
Gut Microbiota Mediates the Anti-Seizure Effects of the Ketogenic Diet. Cell *173*, 1728–
1741.e13.

Park, S.K.R., Jung, T., Thuy-Boun, P.S., Wang, A.Y., John R Yates, I., and Wolan, D.W. (2018).
ComPIL 2.0: An Updated Comprehensive Metaproteomics Database. J. Proteome Res. *18*,
616–622.

Park, S.K., Venable, J.D., Xu, T., and Yates, J.R. (2008). A quantitative analysis software tool
for mass spectrometry–based proteomics. Nat. Methods *5*, 319–322.

Peng, J., Elias, J.E., Thoreen, C.C., Licklider, L.J., and Gygi, S.P. (2003). Evaluation of
multidimensional chromatography coupled with tandem mass spectrometry (LC/LC-MS/MS) for
large-scale protein analysis: the yeast proteome. J. Proteome Res. *2*, 43–50.

Pfeiffer, R.F. (2003). Gastrointestinal dysfunction in Parkinson's disease. Lancet Neurol *2*, 107–
116.

Plovier, H., Everard, A., Druart, C., Depommier, C., Van Hul, M., Geurts, L., Chilloux, J.,
Ottman, N., Duparc, T., Lichtenstein, L., et al. (2017). A purified membrane protein from
Akkermansia muciniphila or the pasteurized bacterium improves metabolism in obese and
diabetic mice. Nature Medicine *23*, 107–113.

Pluskal, T., Castillo, S., Villar-Briones, A., and Oresic, M. (2010). MZmine 2: Modular framework
for processing, visualizing, and analyzing mass spectrometry-based molecular profile data.
BMC Bioinformatics *11*.

Qu, Z.D., Thacker, M., Castelucci, P., Bagyánszki, M., Epstein, M.L., Furness, J.B. (2008).
Immunohistochemical analysis of neuron types in the mouse small intestine. Cell and tissue
research *334*, 147–161.

Rajendran, P.S., Challis, R.C., Fowlkes, C.C., Hanna, P., Tompkins, J.D., Jordan, M.C., Hiyari,
S., Gabris-Weber, B.A., Greenbaum, A., Chan, K.Y., et al. (2019). Identification of peripheral
neural circuits that regulate heart rate using optogenetic and viral vector strategies. Nature
Communications 10, 1944.

Rakhilin, N., Barth, B., Choi, J., Muñoz, N.L., Kulkarni, S., Jones, J.S., Small, D.M., Cheng, Y.-
T., Cao, Y., LaVinka, C., et al. (2016). Simultaneous optical and electrical in vivo analysis of the
enteric nervous system. Nature Communications *7*, 31.

Rakhilin, N., Garrett, A., Eom, C.-Y., Chavez, K.R., Small, D.M., Daniel, A.R., Kaelberer, M.M.,
Mejooli, M.A., Huang, Q., Ding, S., et al. (2019). An intravital window to image the colon in real
time. Nature Communications *10*, 5647–10.

Ramirez-Carrozzi, V.R., Braas, D., Bhatt, D.M., Cheng, C.S., Hong, C., Doty, K.R., Black, J.C.,
Hoffmann, A., Carey, M., and Smale, S.T. (2009). A Unifying Model for the Selective Regulation
of Inducible Transcription by CpG Islands and Nucleosome Remodeling. Cell *138*, 114–128.

Rao, M., and Gershon, M.D. (2016). The bowel and beyond: the enteric nervous system in
neurological disorders. Nature Reviews Gastroenterology and Hepatology *13*, 517–528.

Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015). limma
powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic
Acids Res *43*, e47–e47.

Rodríguez-Piñeiro, A.M., Bergström, J.H., Ermund, A., Gustafsson, J.K., Schütte, A.,
Johansson, M.E.V., and Hansson, G.C. (2013). Studies of mucus in mouse stomach, small
intestine, and colon. II. Gastrointestinal mucus proteome reveals Muc2 and Muc5ac

accompanied by a set of core proteins. American Journal of Physiology - Gastrointestinal and
Liver Physiology *305*, G348–G356.

Roy, A., Guatimosim, S., Prado, V.F., Gros, R., Prado, M.A. (2015). Cholinergic activity as a
new target in diseases of the heart. Molecular Medicine *20*, 527–537.

Sakai, K., Makino, T., Kawai, Y., and Mutai, M. (1980). Intestinal microflora and bile acids. Effect
of bile acids on the distribution of microflora and bile acid in the digestive tract of the rat.
Microbiol. Immunol. *24*, 187–196.

Samulski, R.J., and Muzyczka, N. (2014). AAV-Mediated Gene Therapy for Research and
Therapeutic Purposes. Annu Rev Virol *1*, 427–451.

Sang, Q., and Young, H.M. (1998). The identification and chemical coding of cholinergic
neurons in the small and large intestine of the mouse. Anat. Rec. *251*, 185–199.

Sannasiddappa, T.H., Lund, P.A., and Clarke, S.R. (2017). In Vitro Antibacterial Activity of
Unconjugated and Conjugated Bile Salts on Staphylococcus aureus. Front. Microbio. *8*, 1581.

Schneider, S., Wright, C.M., and Heuckeroth, R.O. (2019). Unexpected Roles for the Second
Brain: Enteric Nervous System as Master Regulator of Bowel Function. Annu. Rev. Physiol. *81*,
235–259.

Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W.S., and Huttenhower,
C. (2011). Metagenomic biomarker discovery and explanation. Genome Biol *12*, R60.

Seillet, C., Luong, K., Tellier, J., Jacquelot, N., Shen, R.D., Hickey, P., Wimmer, V.C.,
Whitehead, L., Rogers, K., Smyth, G.K., et al. (2020). The neuropeptide VIP confers anticipatory
mucosal immunity by regulating ILC3 activity. Nat Immunol *21*, 168–177.

Shan Li, Z., Schmauss, C., Cuenca, A., Ratcliffe, E., Gershon, M. D. (2006). Physiological
Modulation of Intestinal Motility by Enteric Dopaminergic Neurons and the D2 Receptor:
Analysis of Dopamine Receptor Expression, Location, Development, and Function in Wild-Type
and Knock-Out Mice. J Neurosci. *26*, 2798-2807.

Sinha, R., Stanley, G., Gulati, G.S., Ezran, C., Travaglini, K.J., Wei, E., Chan, C.K.F., Nabhan,
A.N., Su, T., Morganti, R.M., et al. (2017). Index switching causes "spreading-of-signal" among
multiplexed samples in Illumina HiSeq 4000 DNA sequencing. *29*, 1072–29.

Steele, P.A., Brookes, S.J., and Costa, M. (1991). Immunohistochemical identification of
cholinergic neurons in the myenteric plexus of guinea-pig small intestine. Neuroscience *45*,
227–239.

Sumner, L.W., Amberg, A., Barrett, D., Beale, M.H., Beger, R., Daykin, C.A., Fan, T.W.M.,
Fiehn, O., Goodacre, R., Griffin, J.L., et al. (2007). Proposed minimum reporting standards for
chemical analysis Chemical Analysis Working Group (CAWG) Metabolomics Standards
Initiative (MSI). Metabolomics *3*, 211–221.

Sundararajan, Z., Knoll, R., Hombach, P., Becker, M., Schultze, J.L., and Ulas, T. (2019). Shiny-
Seq: advanced guided transcriptome analysis. BMC Res Notes *12*, 432–435.

Suzek, B.E., Huang, H., McGarvey, P., Mazumder, R., and Wu, C.H. (2007). UniRef:
comprehensive and non-redundant UniProt reference clusters. Bioinformatics *23*, 1282–1288.

Szklarczyk, D., Gable, A.L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M.,
Doncheva, N.T., Morris, J.H., Bork, P., et al. (2019). STRING v11: protein-protein association
networks with increased coverage, supporting functional discovery in genome-wide
experimental datasets. Nucleic Acids Res *47*, D607–D613.

Tabb, D.L., McDonald, W.H., and Yates, J.R. (2002). DTASelect and Contrast: tools for
assembling and comparing protein identifications from shotgun proteomics. J. Proteome Res. *1*,
21–26.

Talbot, J., Hahn, P., Kroehling, L., Nguyen, H., Li, D., and Littman, D.R. (2020). Feeding-
dependent VIP neuron-ILC3 circuit regulates the intestinal barrier. Nature *579*, 575–580.

Tavares-Ferreira, D., Shiers, S., Ray, P.R., Wangzhou, A., Jeevakumar, V., Sankaranarayanan,
I., Cervantes, A.M., Reese, J.C., Chamessian, A., Copits, B.A., Dougherty, P.M., Gereau, R.W.,
Burton, M.D., Dussor, G., and Price, T.J. (2022). Spatial transcriptomics of the dorsal root
ganglia identifies molecular signatures of human nociceptors. Sci. Transl. Med. *14*, eabj8186.

Thompson, K.J., Khajehali, E., Bradley, S.J., Navarrete, J.S., Huang, X.P., Slocum, S., Jin, J.,
Liu, J., Xiong, Y., Olsen, R.H.J., et al. (2018). DREADD Agonist 21 Is an Effective Agonist for
Muscarinic-Based DREADDs in Vitro and in Vivo. ACS Pharmacol Transl Sci *1*, 61–72.

Treweek, J.B., Chan, K.Y., Flytzanis, N.C., Yang, B., Deverman, B.E., Greenbaum, A., Lignell,
A., Xiao, C., Cai, L., Ladinsky, M.S., et al. (2015). Whole-body tissue stabilization and selective
extractions via tissue-hydrogel hybrids for high-resolution intact circuit mapping and
phenotyping. Nature Protocols *10*, 1860–1896.

Truong, D.T., Franzosa, E.A., Tickle, T.L., Scholz, M., Weingart, G., Pasolli, E., Tett, A.,
Huttenhower, C., and Segata, N. (2015). MetaPhlAn2 for enhanced metagenomic taxonomic
profiling. Nat. Methods *12*, 902–903.

Valicenti-McDermott, M.D., McVicar, K., Cohen, H.J., Wershil, B.K., and Shinnar, S. (2008).
Gastrointestinal symptoms in children with an autism spectrum disorder and language
regression. Pediatr. Neurol. *39*, 392–398.

Van Herreweghen, F., Van den Abbeele, P., De Mulder, T., De Weirdt, R., Geirnaert, A.,
Hernandez-Sanabria, E., Vilchez-Vargas, R., Jauregui, R., Pieper, D.H., Belzer, C., et al.
(2017). In vitro colonisation of the distal colon by Akkermansia muciniphila is largely mucin and
pH dependent. Benef Microbes *8*, 81–96.

Vavassori, P., Mencarelli, A., Renga, B., Distrutti, E., and Fiorucci, S. (2009). The bile acid
receptor FXR is a modulator of intestinal innate immunity. J Immunol *183*, 6251–6261.

Walsh, K.T., and Zemper, A.E. (2019). The Enteric Nervous System for Epithelial Researchers:
Basic Anatomy, Techniques, and Interactions With the Epithelium. Cellular and Molecular
Gastroenterology and Hepatology *8*, 369–378.

Wang, H. Foong, J. P. P., Harris, N. L., Bornstein, J. C. (2022). Enteric neuroimmune
interactions coordinate intestinal responses in health and disease. Mucosal Immunology 15, 27-
39.

Wang, M., Carver, J.J., Phelan, V.V., Sanchez, L.M., Garg, N., Peng, Y., Nguyen, D.D.,
Watrous, J., Kapono, C.A., Luzzatto-Knaan, T., et al. (2016). Sharing and community curation of
mass spectrometry data with Global Natural Products Social Molecular Networking. Nat
Biotechnol *34*, 828–837.

Watanabe, M., Houten, S.M., Mataki, C., Christoffolete, M.A., Kim, B.W., Sato, H., Messaddeq,
N., Harney, J.W., Ezaki, O., Kodama, T., et al. (2006). Bile acids induce energy expenditure by
promoting intracellular thyroid hormone activation. Nature *439*, 484–489.

Wess, J., Nakajima, K., and Jain, S. (2013). Novel designer receptors to probe GPCR signaling
and physiology. Trends Pharmacol. Sci. *34*, 385–392.

Wessel, D., and Flügge, U.I. (1984). A method for the quantitative recovery of protein in dilute
solution in the presence of detergents and lipids. Analytical Biochemistry *138*, 141–143.

Whitcomb, D.C., and Lowe, M.E. (2007). Human pancreatic digestive enzymes. Dig. Dis. Sci.
52, 1–17.

Wu, Y.E., Pan, L., Zuo, Y., Li, X., and Hong, W. (2017). Detecting Activated Cell Populations
Using Single-Cell RNA-Seq. Neuron *96*, 313–329.e316.

Xu, T., Park, S.K., Venable, J.D., Wohlschlegel, J.A., Diedrich, J.K., Cociorva, D., Lu, B., Liao,
L., Hewel, J., Han, X., et al. (2015). ProLuCID: An improved SEQUEST-like algorithm with
enhanced sensitivity and specificity. J Proteomics *129*, 16–24.

Xu, T., Venable, J.D., Park, S.K., Cociorva, D., Lu, Q.B., Liao, L., Wohlschlegel, J., Hewel, J.,
Yates, J.R., and Conciorva, D. (2006). ProLuCID, a fast and sensitive tandem mass spectra-
based protein identification program.

Yang, B., Treweek, J.B., Kulkarni, R.P., Deverman, B.E., Chen, C.-K., Lubeck, E., Shah, S.,
Cai, L., and Gradinaru, V. (2014). Single-Cell Phenotyping within Transparent Intact Tissue
through Whole-Body Clearing. Cell *158*, 945–958.

Yano, J.M., Yu, K., Donaldson, G.P., Shastri, G.G., Ann, P., Ma, L., Nagler, C.R., Ismagilov,
R.F., Mazmanian, S.K., and Hsiao, E.Y. (2015). Indigenous Bacteria from the Gut Microbiota
Regulate Host Serotonin Biosynthesis. Cell *161*, 264–276.

Zhang, H., Sparks, J.B., Karyala, S.V., Settlage, R., and Luo, X.M. (2015). Host adaptive
immunity alters gut microbiota. Isme J *9*, 770–781.

Zhang, X., Smits, A.H., van Tilburg, G.B., Ovaa, H., Huber, W., and Vermeulen, M. (2018).
Proteome-wide identification of ubiquitin interactions using UbIA-MS. Nature Protocols *13*, 530–
550.

Zhu, Q., Mai, U., Pfeiffer, W., Janssen, S., Asnicar, F., Sanders, J.G., Belda-Ferre, P., Al-
Ghalith, G.A., Kopylova, E., McDonald, D., et al. (2019). Phylogenomics of 10,575 genomes
reveals evolutionary proximity between domains Bacteria and Archaea. Nature Communications
10, 5477–14.