1 **Formins play important role in *Leishmania* physiology by acting as cytosolic actin bundlers**
2

3 **Renu Kushwaha*, Arunava Seth*, A.S. Jijumon, Reshmi P.B., Drisya Dileep, Rupak**

4 **Datta[#] and Sankar Maiti[#]**

5 Department of Biological Sciences, Indian Institute of Science Education and Research Kolkata,

6 Mohanpur, - 741246, Nadia, West Bengal, India

8 *(contributed equally)

9
[#] Correspondence:
Sankar Maiti (PhD) spm@iiserkol.ac.in
Rupak Datta (PhD) rupakdatta@iiserkol.ac.in
Indian Institute of Science Education and Research Kolkata
Mohanpur, Nadia - 741246
West Bengal, India
16

17 **Keywords:**

18 Leishmania, actin, formin, actin nucleation, actin bundling, SMIFH2

19 **Abbreviations:**
20
21 G-actin, globular actin; F-actin, filamentous actin; SMIFH2, small molecule inhibitor of Formin
22 Homology 2 domains; FH2, formin homology 2 domains; PBS, phosphate buffer saline; WT,
23 wild-type; IC_{50}, 50% inhibitory concentration.
24
25

Abstract

Formins are a highly conserved eukaryotic family of proteins that regulate actin dynamics. They

play important physiological roles in cell adhesion, motility, vesicular trafficking and

cytokinesis. Although sequence analysis of Trypanosomatida genomes predicted multiple

formin-encoding genes, none of them are functionally characterized yet. We report here

experimental identification and functional characterization of two constitutively expressed

formins from the trypanosomatid protozoa *Leishmania major* viz. LmForminA and LmForminB.

These formins exhibited irregular cytosolic distribution that co-localized with actin patches. Co-

sedimentation assay and surface plasmon resonance confirmed that purified LmForminA and B

FH2 domains can bind actin, albeit with differential affinity. Interestingly, both LmForminA and

B FH2 domains were found to be actin bundlers as revealed by low-speed co-sedimentation

assay and TIRF microscopy. LmForminA and B also had actin-nucleating activities, which was

abolished by mutating their conserved Ile residue crucial for actin assembly. The Ile-mutant

formins, however, retained their actin binding and bundling properties. Treatment of *Leishmania*

cells with formin inhibitor SMIFH2 severely perturbed parasite growth and morphology

indicating that Lmformins are physiologically important and may be considered as novel drug

targets.

Introduction

Leishmaniasis is a disease caused by the intracellular protozoan parasite *Leishmania sp*. With

more than 12 million affected people worldwide and 20,000-40,000 annual deaths, leishmaniasis

is one of the most severe neglected tropical diseases (Okwor and Uzonna, 2016). The

unavailability of a vaccine and the emergence of drug-resistant *Leishmania* species calls for the

development of new therapeutic strategies, further emphasizing the need to study the parasite's

cellular mechanisms with greater impetus (Ponte-Sucre et al., 2017). The parasite follows a

digenetic lifecycle, alternating between mammalian hosts (such as humans) and the insect vector

(sandfly). It exists in two distinct forms in the two different hosts, the motile elongated

promastigotes with long flagella in sandfly vector and as non-motile rounded amastigote form,

which resides in host macrophage (Chow et al., 2011; Tsigankov et al., 2014). A crucial event

for entry of the parasite into host cells and subsequent infection and parasite survival in

phagolysosome is cytoskeleton restructuring of the parasite (Frénal and Soldati-Favre, 2009).

The decrease in cell size is probably due to a decrease in surface to volume ratio of the parasite

to minimize contact with harsh parasitophorous vacuole (Sunter and Gull, 2017). The

cytoskeletal restructuring event and range of physiological and cytological changes in

Leishmania physiology are yet to be understood properly (Goyard et al., 2003).

Leishmania cytoskeleton is known to contain actin along with microtubule and intermediate

filaments (Gull, 1999). *Leishmania* actin is present within the cytoplasm, flagella, flagellar

pocket, nucleus and kinetoplast, also present on the nuclear, vacular, and cytoplasmic face of the

plasma membrane. *Leishmania* actin shares 69.8% identity with the mammalian actin

(Sahasrabuddhe et al., 2004). *Leishmania* actin plays a vital role in cellular processes like

microtubule-actin-associated vesicular transport, organelle movement, endocytosis, basal body

separation, and flagellar pocket division (Sahasrabuddhe et al., 2004; Tammana et al., 2010).

Some actin-binding proteins such as profilin, Arp2/3 protein, cofilin, and coronin are reported

in *Leishmania*, which are involved in different cellular functions from cell division to vesicular

trafficking, indicating the importance of actin in the physiology of *Leishmania* (Nayak et al.,

2005; Tammana et al., 2010). However, the functional role of *Leishmania* formins, an important

class of actin-nucleating protein, remains elusive till date (Gupta et al., 2020).

The Formin class of proteins are long multi-domain proteins (Castrillon and Wasserman, 1994;

Zeller et al., 1999). FH1, FH2 and FH3 are formin homology domains present in most of the

formin family of proteins. Formin has a conserved FH2 domain that contains around 400 amino

acid residues (Kovar and Pollard, 2004; Romero et al., 2007). FH2 domain binds at the barbed

end of the actin filament, play a key role in the nucleation and elongation of actin filaments

(Pruyne et al., 2002; Sagot et al., 2002). In addition, FH2 domain binding to the barbed end

protects from complete inhibition of filament elongation by the capping protein (Kovar and

Pollard, 2004). FH1 domain is a polyproline-rich region present proximately to the FH2 domain.

The binding of the profilin-actin complex with the FH1 region promotes the continuous addition

of actin monomer molecules on the growing end of the actin filament (Kovar and Pollard, 2004;

Romero et al., 2007). Furthermore, the FH3 domain plays a role in the regulation and

localization of formin inside the cell (Kato et al., 2001; Petersen et al., 1998).

Phylogenetic analysis of formins revealed that they are a multigene family of proteins (Cvrčková

et al., 2004; Li and Higgs, 2005; Rivero et al., 2005; Wasserman, 1998). Multiple formin genes

are present throughout the eukaryotes, such as *Plasmodium falciparum* (2 genes), *Toxoplasma*

gondii (3 genes), *Caenorhabditis elegans* (6 genes), *Dictyostelium discoideum* (10 genes),

Schizosaccharomyces pombe (3 genes), *Saccharomyces cerevisiae* (2 genes), *Drosophila*

melanogaster (6 genes), and mammals (15 genes) (Chalkia et al., 2008). Recent biochemical and

cellular characterization revealed the necessity of studying these formin families in eukaryotes.

For instance, i*n vivo* studies of *Saccharomyces cerevisiae* formins*,* Bnip, and Bn1p, showed their

role in cytokinetic actin ring and actin cable assembly (Evangelista et al., 2002; Sagot et al.,

2002). Strikingly loss of both the formin genes in *S. cerevisiae* shows lethality (Ozaki-Kuroda et

al., 2001). Biochemical studies showed that both Bni1p and Bnr1p are actin filament nucleators;

In addition, bnr1 is an F-actin bundler; conversely, bni1 is not (Moseley and Goode, 2005).

Some formins from protozoa such as *Chlamydomonas, Entamoeba. histolytica, Toxoplasma,*

and Plasmodium also have been characterized. In *Chlamydomonas reinhardtii* formin, CrFor1

nucleates actin molecules, which helps in the formation of the fertilization tubule (Christensen et

al., 2019). *Entamoeba* formin isoforms Ehformin-1 and -2 bind with the F-actin structures and

involved in cellular processes such as motility, phagocytosis and cell division (Majumder and

Lohia, 2008). *Toxoplasma gondii* formins contribute to the movement and invasion of the

parasite in host cells (Daher et al., 2010; Daher et al., 2012). Recently it has been found that

formin-2 in *Plasmodium falciparum* and *Toxoplasma gondii* play a vital role in apicoplast

segregation (Stortz et al., 2019). Also, it has been demonstrated that *Plasmodium falciparum*

formin-1(PfFormin1) and formin-2 (PfFormin2) helps in actin polymerization, and PfFormin1

has been reported to have a role in invasion (Baum et al., 2008).

Such a diverse role of these formin in parasite groups brings our attention towards a very

primitive organism, *Leishmania*, which was thought to be evolved 80 million years ago (Chalkia

et al., 2008). The flagellated protozoan parasite belongs to the kinetoplastida group (Filardy et

al., 2018). Phylogenetic analysis predicted the presence of two putative formin genes in

Leishmania major (*L. major*) (Chalkia et al., 2008). Bioinformatics data predicted that these

formins might have roles in the flagellum dynamics and intraflagellar mechanism (Vasconcelos

et al., 2008). However, so far, formins from *Leishmania* (Trypanosomatids) have not been

functionally characterized. In this study, we demonstrated that *L. major* expresses two formins,

LmForminA and LmForminB, both at the RNA and protein levels. Both the formin express and

localized as patchy distribution in the cytosol of *Leishmania major*. Both the formin LmForminA

and LmForminB acts as an actin nucleator and also able to bundled actin filament. Inhibition of

Formin activity leads to lethality, which indicates that formin activity is essential for *Leishmania*

survivability.

Experimental method

Parasite culture

L. major promastigotes (strain 5ASKH ATCC) were cultured as described before (Pal et al.,

2015). Cells were grown at 26°C in M199 medium (Gibco) supplemented with 15% fetal bovine

serum (Gibco), 23.5 mM HEPES, 0.2 mM adenine, 150 µg/ml folic acid, 10 µg/ml hemin, 120

U/ml penicillin, 120 µg/ml streptomycin, and 60 µg/ml gentamicin.

Total RNA isolation and RT PCR

Total RNA was isolated from *L. major* promastigotes using TRIzol reagent (Invitrogen)

followed by DNase I (Invitrogen) digestion to remove genomic DNA contaminants using the

manufacturer's protocols. cDNA was synthesized from 1 µg of total DNase treated RNA using

an oligo(dT) primer and verso cDNA synthesis kit (Thermo Scientific) using the manufacturer's

protocol. The LmForminA and LmForminB FH2 domain transcripts were amplified using gene-

specific primers P1/P2 and P3/P4 (Table S1).

Sequence alignment & Plasmid construction

L. major has two predicated formins in the protein domain prediction database (Chalkia et al.,

2008). Nucleic acid sequences of predicted *L. major* formins were obtained from the database at

the   universal   protein   resource   (https://www.uniprot.org/)   and

(https://www.ebi.ac.uk/ena/data/sequence/search). *L.major* formins FH2 domains were aligned

with the other characterized formins. Multiple sequence alignment was done using the software

clustal omega (https://www.ebi.ac.uk/Tools/msa/clustalo/) and plotted using the program ESPript

3.0 (http://espript.ibcp.fr/ESPript/ESPript/) (Gouet et al., 1999). EMBOSS Needle, pairwise

sequence alignment was also done with LmForminA FH2 domain (accession number:

Q4QE97) and LmForminB FH2 domain (Accession number: Q4QAM2).

L. major genomic DNA was used to amplify the gene of interest. LmForminA (695aa-1092aa)

and LmForminB (714aa-1149aa) FH2 domains were cloned into the pET28a expression vector

using the primers P1/P2 and P3/P4 (Table S1). Site-directed mutagenesis was generated by

Quikchange site-directed mutagenesis protocol. Primers P5/P6, P7/P8 (Table S1) used for the

mutation of isoleucine to alanine residue in FH2 domain constructs. Mutant LmForminA and

LmForminB constructs (I777A LmForminA and I802A LmForminB) were confirmed by the

sequencing.

Protein expression/purification studies

LmForminA and LmForminB FH2 domain His-tagged constructs were transformed in the

BL21(DE3) (Stratagene) cells at 37 ºC and induced with 0.5 mM IPTG at 19 ºC for 12 hours

(Harris and Higgs, 2006). Post-harvest cells were resuspended in the lysis buffer (50 mM Tris-Cl

pH 8.0, 100 mM NaCl, 30 mM Imidazole pH 8.0, 1 mM DTT, 0.2% IGEPAL (Sigma-Aldrich),

0.2% Thesit (Sigma-Aldrich), protease inhibitor cocktail followed by sonication (1 minute's

pulse with 5 second interval) and centrifuged at 12,000 rpm for 15 minutes. 50% Ni-NTA beads

(Qiagen) were added with the supernatant, then washed with wash buffer (50 mM Tris-Cl pH

8.0, 100 mM NaCl, 30 mM imidazole pH 8.0) and eluted with elution buffer (50 mM Tris-Cl pH

8.0, 100 mM NaCl, 300Mm imidazole pH 8.0 pH 8.0 and 5% glycerol (Sigma-Aldrich) (Dutta et

al., 2017). Purified protein dialysis in TNEG5 buffer (20 mM Tris pH 8.0, 100 mM NaCl, 1 mM

EDTA, 5% Glycerol) at 4 ºC. Dialyzed protein stored at 4 ºC.

Polyclonal antibody generation

Eight-week-old mice (BALB/c) were injected subcutaneously with 50 µg of recombinant protein

of LmForminA and LmForminB mixed with Complete Freund's Adjuvant (Santa Cruz

Biotechnology). Three booster injections followed initial immunization with Incomplete

Freund's adjuvant at a two-week interval. Seven days after the final injection, serum was

collected and stored at -80ºC with 0.01% sodium azide and 10% glycerol. The animal

experiments were conducted in accordance with CPCSEA, Govt. of India guidelines and

Institutional Animal Ethics Committee approved protocol.

Western blot with WT L. major cells extract

2×10^8 log phase WT cells were pelleted by centrifugation at 1000g for 5 mins at 4ºC. The cells

were washed with 1X PBS and centrifuged at 1000g for 5 mins at 4 ^{0}C. The cells were then

resuspended either in 100 µl of protein loading buffer (78 mM Tris-Cl pH 6.8, 0.25% SDS, 25

mM DTT. 12.5% glycerol) for LmForminA 100 µl of urea lysis buffer (8 M Urea, 5% SDS, 50

mM Tris-Cl 6.8, 25 mM DTT, 0.1 mM EDTA) for LmForminB mixed with an equal volume of

2X protein loading buffer and subsequently boiled for 3 minutes at 100°C with intermittent slow

mixing. SDS-PAGE was immediately performed with the prepared sample. LmForminA and

LmForminB were detected with raised antibodies against each protein at a primary antibody

dilution of 1:1000 for both diluted in TBST buffer (10 mM Tris, 150 mM NaCl, 0.005 % (v/v)

Tween20). Incubation with primary antibody was performed overnight in cold condition with

gentle shaking. The blots were then incubated with HRP-conjugated rabbit anti-mouse secondary

antibody at a dilution of 1:4000 for 2 hours. Finally, blots were developed using SuperSignal

West Pico Chemiluminescence substrate and observed in Chemidoc imaging system (BioRad).

Treatment of L. major cells with SMIFH2

SMIFH2 (EMD Millipore) was freshly dissolved in dimethyl sulfoxide (100% DMSO) to

prepare a 100 mM stock solution (in the dark). According to the experimental requirements,

further dilutions were made in DMSO before addition to the culture medium. *L. major*

promastigotes were grown in a medium containing SMIFH2 at desired concentrations for 24

hours, following which the cells were microscopically counted with a hemocytometer. Cells

were incubated with an equivalent concentration of DMSO (0.1%) and was used as an untreated

control in all the experiments.

Scanning electron microscopy of L. major cells

The samples were prepared for electron microscopy as described before (Pal et al., 2015).

Briefly, 1 x10^7 treated and untreated samples were taken and washed with chilled 1X PBS and

centrifuged after incubation with inhibitor (in different concentrations) or equivalent

concentration of DMSO or PBS for 12 hours. The cells were resuspended and fixed in 200 µl

2.5% glutaraldehyde at 4°C for 1 hour. Cells were then pelleted and washed twice with 1X PBS

and twice with distilled water. Subsequently, the cells were resuspended in 200 µl of osmium

tetroxide solution for 20 minutes at room temperature. Cells were again pelleted and washed

with 1X PBS. The cells were gradually dehydrated by treating with a gradient of ethanol solution

from 30% to 90%. Finally, cells were resuspended in absolute ethanol and spread on a cut piece

of a silicon wafer, and allowed to air dry. Silicon wafer is then placed in a desiccator connected

with a vacuum pump. The wafer was gold coated and visualized using a Zeiss Supra 55VP

scanning electron microscope. The cell length was quantified with ImageJ software, cells were

measured from cell body end to end, excluding the flagella. At least 50 cells were counted for

each experimental set. Unpaired t-test were performed. *P*-values ≤0.05 were considered

statistically significant, and levels of statistical significance are indicated as *P≤0.05, **P<0.01,

***P<0.001, ****P<0.0001.

Immunofluorescence study of L. major cells

2 x10^6 cells were pelleted and washed with 1X PBS and spread on poly-L-Lysine coated sterile

coverslip and incubated at room temperature for 30 mins. The excess culture was removed and

washed with 1X PBS and fixed with 1:1 acetone methanol solution in the dark for 10 minutes.

The cells were washed with 1X PBS and permeabilized with 0.1% Triton X 100 for 2 minutes.

Cells were then washed with 1X PBS and blocked with 0.2% gelatin solution for 10 minutes.

Mouse anti-LmForminA (1:200), mouse anti-ForminB (1:200), rabbit anti-LmCA1 (1:200),

rabbit anti-LmActin (cross-reacts with LmActin) (1:2000) was added to the coverslip and

incubated for 90 minutes. The cells were then washed twice with 1X PBS and incubated with

Alexa 488 goat anti-mouse and Alexa 488 goat anti-mouse antibodies for 90 minutes. The cells

were washed thrice with 1X PBS and finally mounted in anti-fade media containing DAPI

(Vectashield). Cells were visualized using Leica SP8 confocal microscope with a 63X oil

immersion lens. All images were processed in LASX and ImageJ software. 5 stacks from the

central stacks are merged, and maximum intensity images presented in each case.

Actin filament co-sedimentation assay

The rabbit skeletal muscle was used for the preparation of the actin acetone power. G-actin was

purified from actin acetone powder with G-Buffer [5 mM Tris pH 8.0 (Sigma-Aldrich), 0.2 mM

ATP (USB), 0.2 mM $CaCl_2$ (USB) and 0.2 mM DTT (USB)] (Pollard, 1984). RMA (Rabbit

skeletal Actin) polymerized in TEKG5 (30 mM Tris, 1 mM EGTA, 50 mM KCl, 5% Glycerol)

buffer at room temperature for 1 hour, 30 minutes. The actin polymerization was initiated by an

ion mix (20X= 1M KCl, 40 mM $MgCl_2$, 10 mM ATP). Different concentrations of LmForminA

and LmForminB FH2 (0.5 µM, 1 µM, 2 µM, 4 µM) were incubated with F-actin for 30 minutes.

After completion of the incubation, the reaction was transferred to an ultracentrifuge tube,

centrifuged at 310x1000g for 30 minutes in a TLA-100 rotor (Beckman Coulter). The

supernatant fractions were collected separately mixed with 4X sample loading buffer. Pellet

fractions were resuspended in polymerization buffer to make up the volume and mixed with 4X

sample loading buffer. All the samples were boiled, loaded on SDS-PAGE and visualized with

Coomassie (R-250) (SRL) staining (Zimmermann et al., 2016). For the actin-bundling assay, the

experiment was done similarly, except pre-polymerized actin was incubated with LmForminA

and LmForminB for 30 minutes, and the reaction mixture was centrifuged at 9.2 x1000g for 10

minutes at 4 ℃.

The affinity of the LmForminA and LmForminB for F-actin were determined by the high-speed

actin co-sedimentation assay followed by polyacrylamide gel electrophoresis. 4 µM LmForminA

and LmForminB incubated with the different concentrations of F-actin (1 µM, 2 µM, 4 µM, 8

µM). The stained gel was analyzed for densitometry using ImageJ software. The amount of

LmForminA and LmForminB bound to F-actin were plotted against different concentrations of

F-actin. The dissociation constant (K_D) of LmForminA and LmForminB were determined by

non-linear curve fitting.

Surface plasmon resonance

Interaction of actin with *L. major* formins was corroborated by the surface Plasmon resonance

technique (SPR) using a Biacore T200 system. Biacore Series S Sensor Chip NTA (consists of

carboxymethylated dextran with covalently immobilized NTA) was utilized to capture

LmForminA and LmForminB (ligand). 0.5 mM NiCl was used for the binding of poly-Histidine

tagged LmForminA and LmForminB, and G-buffer was used as a running buffer. Different

concentrations of F-actin (0.5 µM, 1 µM, 2 µM, 4 µM) flowed onto the capture LmForminA and

LmForminB in running buffer. The LmForminA and LmForminB interaction with F-actin was

monitored by an increment in the response units. Surface regeneration was conduct by

introducing 800 mM imidazole for 60 seconds. Double referencing (blank of the surface without

a ligand and buffer injections) was also performed to avoid the chance of nonspecific interaction.

Actin nucleation assay by fluorescence spectroscopy

12 µM G-actin stock (RMA, with 10% pyrene label) were prepared. The polymerization

reaction was prepared with 2 µM final concentration with the above stock of $MgCl_2$ and EGTA,

and different concentrations of LmForminA and LmForminB were added. Subsequently, actin

polymerization was initiated by adding the 20X Ion-mixed (40 mM $MgCl_2$, 10 mM ATP, 1 M

KCl) buffer. Actin polymerization was observed by the direct observation of fluorescence at

25ºC (Higgs and Pollard, 1999; Moseley et al., 2006). The fluorescence were observed with

excitation (365 nm) and emission (407 nm) in fluorescence spectrophotometer (QM40, Photon

Technology International, Lawrenceville, NJ) (Moseley et al., 2006). The slopes of fluorescence

curves determined actin assembly rates at lag phase to 50 % of total actin polymerization.

(Christensen et al., 2019).

Barbed end elongation assay by fluorescence spectroscopy

10 µM unlabeled actin was polymerized in F-buffer for 4 hours at room temperature. F-actin was

sheared by passing through the 27-gauge needle for five times used as actin seed. F-buffer, along

with the LmForminA and LmForminB were added to the actin seed and mixed gently. 0.5 µM

final of G-actin (10% pyrene-labelled) was added to actin seed with cut tips, mixed twice by

pipetting up-down, and the final reaction mix was transferred to a fluorometer cuvette. The

fluorescence was measured at 365/407nm and recorded for 1000 seconds (Moseley et al., 2006).

Elongation rates were further determined by the linear fitting of the initial 100 seconds

elongation (Christensen et al., 2019).

In another barbed end elongation experiment capping protein and *L. major* formins were added

simultaneously to the F-actin seed, subsequently added the actin monomers and monitored the

fluorescence of the actin filament elongation.

TIRF Microscopy

G-actin was polymerized to F-actin in polymerization buffer (10 mM Tris-Cl pH 8.0, 0.2 mM

DTT, 0.2 mM ATP, 0.2 mM $CaCl_2$) for 1 hour at room temperature. 500 nM LmForminA and

LmForminB were added to the 2.5 µM actin filaments and were mixed gently. Polymerization

buffer, Alexa-488-phalloidin were added and the reaction diluted in the imaging buffer [20 mM

HEPES (USB), 1 mM EDTA (Sigma), 50 mM KCl (Sigma-Aldrich), and 5% Glycerol]

immediately. Samples were applied on the poly-D-lysine coated 22 mm coverslip and were

imaged by the Olympus TIRF IX83 microscopy using 100×1.49 N.A. objective.

Actin nucleation in the presence of the LmForminA and LmForminB were done in a flow

chamber. 0.5 µM actin was polymerized in the presence and absence of the LmForminA and

LmForminB and actin stained by Alexa-488-phalloidin mixing in the polymerization buffer. The

images were captured after 10 minutes. The number of filaments were quantified over time by

using the ImageJ software.

Results

Two formins are constitutively expressed in L. major

Sequencing of *L. major* genome had predicted the presence of two putative formins, LmForminA

and LmForminB, in chromosome number 17 and 24 (Ivens, 2005). To check for expression of

these formins*,* total RNA was extracted from *L. major* promastigotes, cDNA was synthesized

and used as the template to PCR amplify the 1191 bp and 1305 bp FH2 domains of LmForminA

and LmForminB, respectively (fig. 1A, 1B). To check the expression of formins at the protein

level, we had planned to generate antibodies against LmFormins (LmForminA and LmForminB).

The LmFormins FH2 domains were cloned into the pET28a vector from genomic DNA and

confirmed by sequencing (fig. 1A, S1A). The respective FH2 domains were expressed as N-

terminal His-Tag proteins in *E. coli* Bl21(DE3) and purified with Ni-NTA agarose (fig. S1B).

Purified proteins were used for antibody generation in mice. The antibodies were found to be

sensitive up to 10 ng and 40 ng, respectively, in the given condition (fig. S1C). Western blot for

L. major whole cell lysate with anti-LmForminA antibody showed a distinct band of ~180 kDa

and a few bands between 100-135 kDa (fig. 1C). Higher than the expected molecular weight

(138.38 kDa) of LmForminA, might be due to post-translational modifications. Similarly, the

western blot of *L. major* whole cell lysate with LmForminB antibody detected a single band at

130 kDa (fig. 1D), which is the expected molecular weight of LmForminB (129.19 kDa). Taken

together, our data confirmed that *L. major* cells constitutively express two formins, LmForminA

and LmForminB, both at the RNA level and protein level.

Both LmForminA and LmForminB have patchy cellular localization

After we confirmed the expression of both the formins LmForminA and LmForminB in *L.*

major cells, we tried to determine the intracellular localization of these formins.

Immunofluorescence study with anti-LmForminA antibody showed patchy distribution with a

prominent punctum near the nucleus and kinetoplast (fig 2A). Similarly, LmForminB also

showed a patchy distribution inside the cell. LmForminA puncta partially colocalized with *L.*

major Carbonic anhydrase 1 (LmCA1), a cytosolic protein of the parasite (Pal et al., 2017),

indicating the puncta was cytosolic (fig. 2A). Additionally, the LmForminA puncta also

colocalized with the actin patches suggesting that it might be bound with the actin inside the cell,

thus giving it the distinctly distributed localization (fig. S2). Similarly, LmForminB puncta also

colocalized partially with LmCA1 and with LmActin within the cell (fig. 2A, S2).

Growth of L. major cells was inhibited by formin inhibitor SMIFH2

To check the physiological importance of formins*, L. major* promastigotes were grown in the

presence of SMIFH2, a formin FH2 domain inhibitor (Rizvi et al., 2009). Results presented in

fig. 2D showed a dose-dependent reduction in cell number with an IC_{50} value of 11.86 µM.

Scanning electron microscopy of the SMIFH2-treated *L. major* cells revealed morphological

abnormalities when the cells became shortened with rounded shape in the presence of the

inhibitor (fig. 2B). Furthermore, the length of the parasite was significantly reduced at even at

2.5 µM concentration of SMIFH2, which is about one-fourth of the IC_{50} (11.86 µM) (fig. 2C).

This data suggests that rounded morphology and reduction of cell length were not due to general

stress but might be due to altered cytoskeleton dynamics in *L. major* cells. In other words,

formins seem to play an important role in maintaining cell shape and morphology of *L. major*.

Purified L. major formins bind F-actin in vitro

Both the formins were shown to be colocalized with actin in *L. major* cells (fig. S2). To check

actin interaction with LmFormins actin filaments (F-actin) co-sedimentation assay with purified

LmFormins FH2 domains was performed. Our results showed that LmFormins were sedimented

in the pellet fraction along with actin. However, in control experiments without actin,

LmForminA and LmForminB proteins mostly remained in the supernatant fraction (fig. 3A, B,

respectively). These results indicate that LmForminA and LmForminB protein have F-actin

binding ability in the *in vitro* condition with their FH2 domain similar to other well-characterized

formins (Dutta et al., 2017; Shimada et al., 2004).

The dissociation constant of LmForminA and LmForminB with F-actin were determined by

densitometric analysis of the SDS-PAGE gel of actin filament co-sedimentation assay (fig. S7A,

B). The fraction of bound LmFormins with F-actin curve fitting shows the K_D value of 1.84 µM

and 0.2 µM for LmForminA and LmForminB, respectively (fig. S7C, D). Lower dissociation

constant value for LmForminB compare to LmForminA indicated that LmForminB has a greater

affinity for actin filaments than LmForminA. We reconfirmed the LmFormins interactions with

F-actin using surface plasmon resonance, which is typically used to study biomolecular

interaction in a real-time environment. Different concentrations (0.5 µM, 1 µM, 2 µM, 4 µM) of

sheared F-actin filaments were injected into the flow cell with immobilized LmFormins. The

SPR sensorgram clearly showed the interaction of LmFormins with the F-actin with an increase

in the response unit in a concentration-dependent manner followed by the fast dissociation. (fig.

3C, D). The difference in maximum response units of LmForminA and LmForminB might be

due to their different binding affinity for F-actin.

Actin bundling activity of L. major formins

FH2 domain of formins usually bind to the barbed end of the actin filaments. However,

literature has depicted that some formins also has side binding with actin filament, leading to the

formation of an actin filament bundle (Harris et al., 2004; Michelot et al., 2005). In Bnr1p and

Daam1, the C-terminal domain comprising FH1, FH2, and C-terminal end, are responsible for

actin filament bundling (Barko et al., 2010; Moseley and Goode, 2005). In Arabidopsis formin

AFH1, the FH1 domain is also required along with the FH2 domain for actin-bundling (Michelot

et al., 2005). While in mDia2, FRL1 formin only FH2 domain is a strong actin filament bundler

(Harris et al., 2006). F-actin bundling activity of formins might have very important for the

physiological role in various organisms. However, no actin-bundling activity had been shown for

the formin from the protozoan Kinetoplastida group. It was previously reported that *L. major*

cells contain short F-actin bundles (Sahasrabuddhe et al., 2004). We were thus interested to

know if LmForminA and LmForminB have any role in F-actin bundling. Actin-bundling ability

of purified LmForminA and LmForminB FH2 domains were determined by co-sedimentation

assay at low-speed centrifugation. Polymerized filamentous actin at low-speed centrifugation

remained in the supernatant fraction. Presence of the LmForminA and LmForminB were able to

co-sediment F-actin in the pellet fraction at low-speed centrifugation, indicating *L. major*

formins were able to bundle actin filament (fig. 4A, B) Fascin, a well-characterized actin-

bundling protein was used as a positive control (Yamashiro et al., 1998), which could also bring

down the F-actin in the pellet fraction at low-speed co-sedimentation assay (fig. S5C).

We also used TIRF microscopy to examine the F-actin bundle formation by LmForminA and

LmForminB. Pre-polymerized actin filaments were incubated with the LmFormins, stained with

phalloidin and observed under TIRFM. Long, thick actin bundles appeared in the presence of

LmForminA and LmForminB (fig. 4C). LmFormins induced thick actin filament bundles similar

to positive control Fascin induced actin filament bundle (fig. 4C) compare to the actin filament

control. This results had confirmed that LmForminA and LmForminB able to bundle actin

filament in *In vitro* conditions. More numbers of actin filament bundles were found in the

presence of the LmForminA as compared to LmForminB (fig. 4C). In summary, results showed

that LmForminA has a strong actin-bundling activity, while LmForminB is a weak actin-bundler.

It has been reported that the electrostatic interaction between the FRL1 FH2 or mDia2

FH2 domain with actin is vital for their bundling activity. Salt, ionic strength inlfuence the actin-

bundling activity of mDia2 and FRL1 FH2 domain (Harris et al., 2006). We tested the effects of

ionic strength on the actin-bundling activity of the LmForminA and LmForminB FH2 domain.

We found that with increasing the ionic concentration of KCl (50 mM, 100 mM, 150 mM) actin

bundling activity of the LmForminA was lowered. Actin bundling activity of LmForminA had

reduced at 100 mM concentration of KCl and almost lost at 150 mM KCl concentration (fig.

S9A). Whereas high concentration of KCl had mild effect on actin bundling activity of

LmForminB (fig. S9B).

L. major formins were able to polymerize actin filament

Most of the Formin FH2 domain has actin polymerization activity in *in-vitro* condition, except

FHOD1 FH2, which inhibit the actin polymerization (Schönichen et al., 2013). Actin

polymerization by the formin FH2 domain is also reported in human parasites like *Plasmodium*

falciparum, *Toxoplasma gondii*, where formin plays a vital role in their physiological function.

Plasmodium falciparum use actin-based motility to invade the host cell, which is achieved by

PfFormins. It has been shown that PfFormin1 and PfFormin2 can polymerize chicken muscle

actin in *in-vitro* conditions (Baum et al., 2008). We checked for actin polymerization activity of

L. major formins by spontaneous actin assembly by pyrene fluorescence assay. 2 µM actin

control were shown to polymerize at a basal level, while the addition of LmForminA and

LmForminB effectively increased actin polymerization in a concentration-dependent manner

(fig. 5A, C). Furthermore, the comparative analysis of the assembly rate demonstrated that

LmForminB had a higher rate of actin polymerization than LmForminA (fig. 5B, D). Total

internal reflection fluorescence microscopy (TIRF-M) assays were performed for direct

visualization of actin filament polymerization in presence of the LmForminA and LmForminB

FH2 domain. More actin filaments were observed in the presence of LmForminA and

LmForminB FH2 than in the actin control (fig. 5E). From the above result, it could be

summarized that LmFormins were potent actin nucleators.

L. major formins inhibit actin filaments elongation but antagonize the capping protein

Prior reports have shown that the barbed end of actin filaments captured by formin FH2 had a

control on the barbed end elongation. Formins accelerated barbed end elongation of actin

filaments in the presence of the profilin and reduced barbed end elongation in the absence of

profilin (Patel et al., 2018). Therefore, we performed a barbed end elongation assay to test the

effect of the LmForminA and LmForminB FH2 on actin filaments elongation.

We had found that lower concentration (up to 100 nM) of LmForminA and LmForminB FH2 did

not affect barbed end elongation. In contrast, at higher concentrations, LmForminA and

LmForminB FH2 inhibited actin filament elongation compared to actin control (fig. 6A, B).

LmForminA and LmForminB inhibited the filament elongation by ~63% and ~35%,

respectively, at 1µM concentration. This effect might be because of the actin bundle formation

by LmForminA. Actin bundle formation by LmForminA could lead to a shortage of free barbed

end for elongation. So, the filament's free end was not accessible for the addition of actin

subunits on the barbed end. On the contrary, LmForminB was a weak actin bundler and showed

mild inhibition on filament elongation compared to LmForminA (fig. 6B).

The above observation was confirmed by the TIRF microscopic analysis to understand the effect

of LmForminA FH2 on actin filaments elongation. Bundled actin appeared in the presence of

500nM LmForminA (data not shown). Based on the observation, we speculate that the inhibition

of actin elongation was due to the actin bundle formation in the presence of LmForminA.

Formin FH2 domain has a processive activity for actin filament barbed end and antagonizes

barbed end binding with capping protein. Capping protein known to binds and inhibit elongation

of the barbed end actin filament. mDia1, FRL, and Bni1 FH2 domain protect the barbed ends

from the capping proteins (Harris et al., 2004; Harris et al., 2006) while elongating the filaments.

Therefore, we were interested to see the effect of LmForminA and LmForminB FH2 on the

barbed end elongation in the presence of the capping protein. For this, we performed a barbed

end elongation assay with the FH2 domain of LmForminA, LmForminB vs capping protein. In

the presence of the capping protein, LmFormins FH2 rescued the inhibited actin filaments

elongation (fig. 6E, F). Our result showed that LmFormins bind actin filament barbed end

processively and compete with capping protein for the barbed end binding similar to the other

known characterized formin FH2 domains.

Mutant I777A LmForminA and I802A LmForminB can bind and bundle actin filaments

Isoleucine and lysine are highly conserved amino-acid residues in the formin FH2 domain. These

residues are crucial for actin assembly in most reported formins (Harris et al., 2006; Scott et al.,

2011). In Bni1 and Daam1, mutations of isoleucine to alanine altogether abolish actin assembly

(Lu et al., 2007; Xu et al., 2004)**.** Based on the sequence alignment of LmForminA and

LmForminB with other well-characterized formins (fig. S3), we also found the conserved

isoleucine residues in LmForminA and LmForminB. We generated a point mutation for

conserved isoleucine to alanine in LmForminA and LmForminB FH2 domain at the specific

position (I777A LmForminA and I802A LmForminB) (fig. S3). The F-actin binding abilities of

Mutant LmFormins (I777A LmForminA and I802A LmForminB) were confirmed by the SPR

analysis (fig. S5A, B). The pyrene actin assembly assay was performed to monitor the actin

nucleation activity of the mutant LmFormins. We noticed that the I777A LmForminA had no

actin nucleation activity (fig. 7A). Whereas actin nucleation activity of I802A LmForminB FH2

was reduced significantly (fig. 7C). Rate of actin assembly reduced 2~3 fold for the I802A

LmForminB as compared to WT LmForminB (fig. 7D). In summary, we infer that the conserved

isoleucine residue was crucial for the actin assembly activity of LmForminA and LmForminB.

Mutation in conserved amino acid residues Ile to Ala in frl1, mDia2 results in retention of the

actin-bundling activity (Harris et al., 2006). We were curious to see the behavior of mutant

LmFormins regarding the actin-bundling activity. The result of the low-speed co-sedimentation

assay demonstrated that the actin-bundling activity of the I777A LmForminA (fig. S5D), which

was quite similar to that of WT LmForminA. I802A LmForminB actin-bundling slightly

increased as compared to WT LmForminB (fig. S5E). In another actin-bundling experiment,

actin monomer copolymerized with the mutant LmFormins and WT LmFormins. However,

I802A LmForminB could bundle actin while the WT LmForminB could not (fig. S6C, D). On

the other hand, actin-bundling remained unaffected in the comparison of the WT LmForminA

and I777A LmForminA (fig. S6A, B). Based on the above observation, we speculate that (I-A)

mutation might have increased the side bundling of actin for LmForminB, while it does not have

a similar effect for LmForminA.

We had also performed a barbed end elongation assay to understand the effect of mutant

LmFormins binding with the barbed end and protection from capping protein. Mutant

LmFormins binding with the barbed end has no significant influence on actin filaments

elongation compared to WT LmFormins. The addition of capping protein on actin filaments in

the presence of the mutant LmFormins had significantly less ability to protect the barbed end

from capping protein compared to the WT LmFormins (fig. S8A, B).

Discussions

Formins are essential proteins involved in a range of cellular functions from cell division to

trafficking (Castrillon and Wasserman, 1994; Evangelista et al., 2002; Maas et al., 1990; Sagot et

al., 2002). Recently, formins from parasitic protozoans such as *Plasmodium, Toxoplasma* have

been shown to be very important for infectivity (Baum et al., 2008; Daher et al., 2010; Daher et

al., 2012; McConville et al., 2002; Stortz et al., 2019; Tosetti et al., 2019; Von Dippe and Levy,

1982). Nevertheless, nothing was reported in the case of the Kinetoplasts *Leishmania,* although

the importance of actin and other actin-binding proteins such as Arp2/3 complex, cofilin, profilin

are well reported (Ambaru et al., 2020; Tammana et al., 2010, Gupta et al., 2020).

To uncover the functional aspect of formin, we first treated *L. major* cell with SMIFH2 (Formin

inhibitor). Our data indicate that formin inhibitor SMIFH2 inhibit *L. major* cell growth. Growth

inhibition was observed with SMIFH2, with IC_{50} being 11.86 µM, which is comparable to other

reports such as in fibroblast cells where cytotoxicity was observed at 28 µM of SMIFH2 (Rizvi

et al., 2009), 40 µM in Epithelial ovarian cancer cells (Ziske et al., 2016). In addition to that,

cells appeared rounded when treated with a concentration as low as 2.5 µM of SMIFH2; this

might indicate that the cell rounding is due to perturbation of cytoskeleton dynamics rather than

general stress. It confirms the importance of active formin in *L. major* physiology, which might

be crucial for *Leishmania* survival. In future *Laishmania* formin could be used as drug target, as

formin activity is delicate to *Leishmania* physiology. Searching in the genome database, we

found two putative formins, namely LmForminA and LmForminB (Ivens et al., 2005). We

observed LmFormins expression both at the RNA and protein levels (fig. 1A). Bands at western

blot were observed in the case of LmForminA at higher molecular weight than predicted,

indicating there might be a post-translation modification (fig. 1C) similar observation recorded in

DAAM1, FMNL2 in mammalian cells (Li et al., 2019). However, further studies are needed to

confirm the post-translational modifications. The most characterized formins use RhoA mediated

regulation to regulate their activity. However, presence of the RhoA-GTPase gene had not shown

in literature and genome database of *Leishmania,* which could have been a possible way to

regulate the formin activity in *Leishmania*. A related RhoA-GTPase gene is present in

Trypanosoma but is absent in *L. major* (Abbasi et al., 2011). The absence of the Rho-GTPase

suggests that *L. major* formins activity regulation used a distinct approach from that of present in

mammalian system. As a result, it indicates that post-translation modification, which might play

an important role in regulation of formin activity in *L.major* (Angeles Juanes and Piatti, 2016;

DeWard and Alberts, 2009). The post-translational modification is a well-characterized

phenomenon observed in *Leishmania* (Zilberstein, 2015).

Next, we were interested to see how these LmFormins regulate the actin dynamics *in vivo*. To

see its effect on actin dynamics: first, we have to determine where these formins are localized in

the cell. We had raised LmForminA and LmForminB specific antibodies to study the

localization. LmForminA and LmForminB did not possess any predicted signal peptide for the

secretory pathway suggested that both the formins might be localized in the cytosol. Our

research group's previously studied LmCA (*L. major* carbonic anhydrase) localized in the cytosol

is used as a positive control for conforming to the localization of LmFormins (Pal et al., 2017).

LmFormins localization patches are similar to other actin binding proteins' pattern such as

myosin, coronin (Drubin et al., 1988; Sahasrabuddhe et al., 2009). LmFormins patches were

colocalized with *L. major* actin, indicating that LmFormins binds with the actin cytoskeleton

inside the *L. major* cell (fig. S2).

Next, we were interested to see how LmFormins regulates actin dynamics in *in-vitro* conditions.

FH2 domain is the characteristic feature of the formin family protein, which is the crucial player

for the formin activity in regulating the actin filament polymerization. *L. major* putative formin

genes have one FH1 and FH2 domain (Chalkia et al., 2008). The *L. major* Formins FH2 domain

sequence is aligned with well-characterized formins conserved amino acid with identity around

20.1%-40.9%. Pairwise alignment Our results revealed that LmForminA has 57% similarity

with *Trypanosoma cruzi* ForminA. The characterized formin of *Dictyostelium discoideum*

has 44% similarity with LmForminA and a similarity of 39% with LmForminB (fig. S4).

Our study reveals that LmForminA and LmForminB FH2 domain can interact with actin. Our *in*

vitro studies had shown that the LmFormin-FH2 domain binds with F-actin in a concentration-

dependent manner (fig. 3A, B). The binding affinity of LmForminB shows a strong affinity with

a lower K_D value (0.2 µM) as compared to the LmForminA (K_D=1.84 µM). Lower dissociation

constant of LmForminB showing comparable high affinity similar to K_D that of the FRLα and

FRLβ formins (Harris et al., 2004). but higher affinity than the LmForminA similar to mDia1,

mDia3, and Bni1(2-7µM) (Harris et al., 2004; Li and Higgs, 2003; Shimada et al., 2004).

LmForminA and LmForminB also had a feature to bundle actin filaments under *in-vitro*

condition. The low-speed co-sedimentation assay had confirmed that the LmFormins had actin-

bundling activity. Microscopic observation of F-actin in the presence of the LmForminA and

LmForminB had reconfirmed their actin-bundling feature. Earlier reports already describe some

formins across different eukaryotes having actin filaments bundling activity. These actin-

bundling formins are budding yeast Bnr1p (Moseley and Goode, 2005), *Dictyostelium* ForC

(Junemann et al., 2013), mammalian FRL1, and mDia2 (Harris et al., 2006), *Arabidopsis* AFH1

(Michelot et al., 2005), *Drosophila* FHOD1 and FHOD3 (Patel et al., 2018) and *Drosophila*

Daam1 (Barko et al., 2010). Most recently reported mice Delphilin is a weak actin bundler

(Silkworth et al., 2018). According to the literature, in the protozoan parasite group, only

Toxoplasma gondii formins have actin-bundling activity (Skillman et al., 2012). The above-

characterized actin bundling formins have a common feature: they possess positively charged

amino acids in FH2 domains and have higher PI value. LmForminA FH2 domain able bundle

actin filaments, used in this study has a PI value of 8.31, which is higher than the mDia1 and

Bni1 (PI value 6.37 and 5.64, respectively) could not form an actin bundle. FRL1 and mDia2 (PI

value 8.26 and 8.14, respectively) can firmly form an actin bundle. PI value of LmForminA 8.31

is close to the PI value of mDia2 and FRL1 (Harris et al., 2006). LmForminB, a weak actin

bundler, has a PI value of 7.09, which is higher than the PI value of 5.89 for Delphilin, a weak

actin bundler (Silkworth et al., 2018). As an earlier hypothesis, the net positively charged amino-

acid of the FH2 domain involved in the actin bundle formation while the net negativity charge

FH2 domain could not form actin bundle (Harris et al., 2006). The bundling activity difference in

formins indicates that the positively charged amino acids of the FH2 domain might mediate the

electrostatic interaction with the negatively charged actin filament for bundling. We find, this

presumption is strongly supported by the salt susceptivity of LmFormins bundling activity,

which might reduce the electrostatic interaction are significant for actin-bundling activity (fig.

S9A, B). mDia1 (PI value 6.37) net negatively charged amino-acids at the outer surface would

not form an actin bundle (Harris et al., 2006). In the absence of the crystal structure of the

LmForminA and LmForminB, we could not describe the outer surface amino-acids, but the

LmForminB PI value slightly higher than PI value of mDia1 supports LmForminB has weak

actin bundling activity (Harris et al., 2006).

L. major actin present in the patches (Sahasrabuddhe et al., 2004). It has been reported that

coronin protein is vital for actin-bundling in *Leishmania* (Nayak et al., 2005). Actin bundled

structure formation in *Leishmania* might not be solely coronin dependent. Formins protein might

be playing a pivotal role in actin bundle formation inside the *Leishmania* cell. The physiological

effects of this actin bundle in *Leishmania* are not clear. LmForminA and LmForminB actin

bundling activity differences might point to their different phenotype, localization, and

regulation. *In vitro* biochemical assay based on the pyrene labelled actin assembly shows that

LmFormins increase the actin polymerization in a concentration-dependent manner (fig. 5A, C).

LmForminB is a more potent actin nucleator as compare to LmForminA. The intense nucleation

activity of LmForminB as compare to LmForminA might have some physiological significance

in vivo. However, more studies are needed to confirm this hypothesis.

In the actin assembly rate comparison of, LmForminA with mammalian formin Daam1 exhibits a

similar polymerization rate (fig. 5A). While LmForminB has a three-fold higher polymerization

rate. The important role of actin polymerization in the presence of the formins has already been

reported in some protozoan parasites. In *Toxoplasma gondii* formin, TgFRM1, TgFRM2 &

TgFRM3 is also a potent actin nucleator *in vitro* condition. TgFRM1 and TgFRM2 formin play

an essential role in parasite mobility and host cell invasion (Daher et al., 2010). The malaria

parasite *Plasmodium falciparum* invade host cell by using actin-based motility, which shows that

formin is important in actin polymerization involved in the motility (Baum et al., 2008).

Structural analysis of the Bni1p and mDia1 has shown that conserved amino-acid Ile residue

faces the inside of the FH2 homodimer, this residue of the FH2 domain creates important

interaction with the actin (Xu et al., 2004).

Moreover, a mutation in this residue completely abolishes actin assembly. Investigating the point

mutation (Ile-Ala) in LmFormins, we found I802A LmForminB behave differently compared to

WT LmForminB. This mutation completely abolishes the actin assembly of LmForminA,

similarly Daam1 and *Drosophila* Fhod (Barko et al., 2010; Xu et al., 2004). I802A LmForminB

has a less effect on actin assembly. The Ile-Ala mutant of FRL1 also has a less effect on the actin

assembly (Harris et al., 2006).

(I-A) mutation might affect the flexibility of the FH2 domain and might suppress the switch

from the closed to the open configuration during actin assembly. To understand the actual

mechanism, the resolution of the FH2 domain structure of LmForminA and LmForminB in the

presence of the actin would be necessary.

In the barbed end elongation assay, we find LmFormins reduced the barbed end elongation in the

absence of the profilin like the Drosophila Fhod. LmFormins binds to the barbed and protects

from the capping protein. LmFormins barbed end binding and antagonizing the capping protein

nature Indicates that (1) LmFormins and capping protein might passively compete for the barbed

end during filaments elongation. (2) LmFormins form a processive cap that maintains an

association with the barbed end during filament elongation and replacing the capping protein

from the barbed end. Compared to WT LmFormins, (I-A) mutants have similar actin elongation

activity but have significantly less barbed end protecting activity from the capping protein.

Similarly, Ile to Ala mutation in Drosophila Fhod, FRL1, and mDia1 FH2 also decreases barbed

end elongation (Barko et al., 2010; Harris et al., 2006).

Our study shows the biochemical properties of the LmFormins FH2 domain and its effects on the

actin dynamics. LmFormins FH2 domain overall features are similar to the other Formins.

However, it is still unclear what are the exact roles of LmFormins in *L. major* physiology.

Furthermore, some other important domains, such as RBD, DAD, DID are not predicted in

LmFormins. This indicates that there are most probably differences in regulation of formins in

Leishmania compared to mammalian counterpart.

Acknowledgements:

The authors sincerely thank Mr. Kashinath Sahu, Mr. Jajati Keshari Ray and Mr. Sujoy Bose for

their technical assistance in SEM imaging, mice handling and cell culturing. The authors also

thank Dr. Amogh Sahasrabuddhe of CSIR-CDRI for providing the LdActin antibody. The

authors also thank Dr. Arnab Gupta, Mr. Ruturaj and DBT Welcome imaging facility of IISER

Kolkata for helping with confocal microscopy. The authors also thank Dr. Bidisha Sinha, Dr.

Rinku Kumar Patel, Ms. Madhura Chakraborty and Dr. Arikta Biswas of IISER Kolkata for their

setup and assistance in TIRF microscopy imaging.

Competing interests:

The authors declare that they do not have any conflict of interests.

Author contributions:

Conceived and designed the experiments: SM RD. Performed the experiments: RK AS JAS RPB

DD. Analyzed the data: SM RD RK AS. Wrote the manuscript: SM RD RK AS JAS.

Funding:

R.K. was supported by an individual fellowship from the University Grant Commission, and

A.S. was supported by an individual fellowship from the Council of Scientific and Industrial

Research, Government of India. The work was supported by Department of Biotechnology grant

BT/PR26167/MED/29/1229/2017 to SM and Department of Biotechnology and Department of

Science and Technology grants BT/PR21170/MED/29/1109/2016 and EMR/2017/004506,

respectively, to RD and DST-FIST grant No. SR/FST/LS-II/2017/93(c) awarded to Department

of Biological Sciences, IISER Kolkata.

FIGURE LEGENDS

Figure 1. Expression of the two Formins in *L. major*.

[A] Schematics diagram showing the organization of the predicated FH2 domain of LmForminA

(695-1092 aa) and LmForminB (714-1149 aa) with predicted PCR products of 1191 bp and 1305

bp. [B] Agarose gel: indicating the expression of LmForminA and LmForminB by RT-PCR. The

lanes marked –RT represents negative control with the reverse transcriptase (RT) enzyme. In the

PCR, bands of 1191 bp and 1305 bp were observed for the FH2 domain of LmForminA and

LmForminB. [C] Western blot with anti-LmForminA and anti-LmForminB antibodies with *L.*

major whole cell lysate (WCL).

Figure 2. LmForminA and LmForminB localize in the cytosol and Formin inhibitor

SMIFH2 inhibits the growth of *L. major*.

[A] Confocal imaging of *L. major* cells immunostained with DAPI (blue), anti-LmForminA, and

anti-LmForminB (green) (1:200) anti-LmCA1 (cytosolic marker) (red) (1:200), acquired in

LeicaSP8 confocal microscope. Partial colocalization (yellow) was observed in both cases. Scale

bar represents 10 µm. [B] Scanning electron micrographs of SMIFH2 treated and untreated *L.*

major cells. Cells were treated 12 hours with/without SMIFH2 in respective concentrations.

Images were acquired Zeiss Supra 55VP scanning electron microscope. [C] Cell lengths were

quantified using ImageJ software. Asterisks indicate a significant difference between untreated

with respect to treated cells. **P<01 ****P<0.0001 (Student's *t*-test). [D] Growth kinetics of *L.*

major cells in the presence of formin inhibitor SMIFH2 grown for 24 hours. The cell number

was normalized with respect to DMSO control. Error bar represents the standard deviation from

3 independent experiments.

Figure 3. LmForminA and LmForminB FH2 binds with F-actin.

[A, B] 10% Coomassie-stained SDS-PAGE showing FH2 domains of LmForminA and

LmForminB binds to F-actin. LmForminA and LmForminB incubated with F-actin for binding.

Then co-sedimented at high-speed centrifugation. LmForminA and LmForminB, which bind

with the actin filamnet present in the pellet fraction (P for pellet fraction and S for supernatant

fraction), indicated binding with F-actin. [C, D] Surface plasmon resonance was performed to

see the interaction of LmForminA and LmForminB FH2 with F-actin. LmForminA and

LmForminB FH2 was captured on an NTA chip, and different concentrations of F-actin (0.5, 1,

2, 4 µM) flowed on captured LmForminA and LmForminB FH2. The sensorgram has shown the

interaction phase of LmForminA [C] and LmForminB [D], followed by the dissociation phase.

The interaction has been seen by an increase in response in a concentration-dependent manner.

Figure 4. LmForminA and LmForminB FH2 able to bundled F-actin.

10% Coomassie-stained SDS-PAGE showing F-actin bundling for LmForminA (A) and

LmForminB (B). F-actin were incubated with LmFormins and co-sedimented at a low-speed

centrifugation. Actin bundled in pellet fraction in the presence of the LmForminA and

LmForminB FH2, indicating the bundling nature of LmFormins (P for pellet fraction and S for

supernatant fraction)**.** [C] TIRF microscopic image for direct visualization of the event in actin-

bundling in the presence of the 500 nM LmForminA and LmForminB. LmForminA and

LmForminB protein mixed with 2.5 µM actin polymerized and stained with Alexa-488-

phalloidin. 20 µl diluted reaction taken inflow chamber and imaged capture by the TIRF

Microscopy. Formation of thick actin filament bundle (indicated by arrow) in the presence of the

LmFormins and Fascin. The scale bar is 10 µm. Actin-bundling: Fascin served as a well

characterized actin filament bundling protein as positive control.

Figure 5. LmForminA and LmForminB efficiently accelerate an actin filament assembly.

2 µM rabbit muscle actin (10% pyrene-labeled) used for actin assembly. [A, C] shows actin

assembly in the presence of LmForminA and LmForminB, respectively. Actin assembly

increased in a concentration-dependent manner in the presence of the LmForminA and

LmForminB. [B, D] Actin assembly rate at half-maximal polymerization of actin in the presence

of a different concentration of LmForminA and LmForminB. A.U., arbitrary units. [E] 0.5 µM

Actin used for polymerization in the presence or absence of LmForminA and LmForminB,

respectively. Alexa-488-phalloidin was used for the staining of actin, and the image was

captured by TIRF microscopy. More number of actin filaments were observed in the presence of

LmForminA and LmForminB as compared to the actin control. The scale bar is 10 µm.

Figure 6. Actin filament elongation by *L. major* Formins in the presence of capping

protein.

[A, B] Actin seed was prepared by F-actin passing from the 27-gauge needle five times. Actin

seed mixed with (10%) pyrene labelled G-actin. Actin filament elongation was performed in the

presence of different concentrations of LmForminA and LmForminB. Rate of actin filament

elongation decrease in the presence of LmForminA and LmForminB in a concentration-

dependent manner. [C, D] Rate of elongation were measured as the initial slope over the first 100

seconds for LmForminA and LmForminB. [E, F] Actin filament elongation with various

concentrations of LmForminA and LmForminB in the presence of capping protein respectively.

Figure 7. I-A mutant of LmForminA and LmForminB FH2 binds and bundle actin

filament.

[A] Direct comparison of actin assembly in the presence of the WT and I777A LmForminA.

I777A LmForminA had lost actin nucleation activity. [B] Actin polymerization in the presence

of the WT LmForminB and I802A LmForminB. I802A LmForminB mutant had less actin

nucleation activity compare to the WT LmForminB. [C, D] Rate of actin polymerization had

plotted against LmFormins concentrations respectively.

Supplementary Figure

Figure S1. [A and B] Cloning, purification of LmForminA and LmForminB FH2 domain.

[A] Agarose gel electrophoresis of purified plasmids of LmForminA and LmForminB FH2 domains are shown in lanes 1, 2 along with an empty vector in lane 3. Cloned LmFormins plasmids digestion with BamHI and HindIII leading to expected bands at 1.19kb and 1.3kb are shown in lanes 4, 5. [B] 10% Coomassie-stained SDS-PAGE gel of purified LmForminA and LmForminB protein used in this study.

[C and D] Western blot against purified LmForminA and LmForminB FH2 with raised antibody against LmForminA and LmForminB FH2. Antibodies were generated against purified LmForminA and LmForminB FH2 in mice. Purified protein was loaded SDS-PAGE in gradient 1 µg, 2.5 µg, 5 µg, 10, µg, 20 µg, 40 µg for LmForminA and LmForminB FH2 and shown with gradient bar. Western blot with specific raised antibody against LmForminA and LmForminB FH2.

Figure S2. Co-localization of *L. major* Formins with the actin inside *Leishmania* Cell.
Confocal imaging of L. major cells immunostained with DAPI (blue), Anti-LmForminA, and Anti-LmForminB (green) (1:200) LdActin (cross-reacts with LmActin) (red) (1:2000), acquired in LeicaSP8 confocal microscope. Partial colocalization (yellow) was observed in both cases, indicated with white arrows. Scale bar represents 10 µm.

Figure S3. Multiple sequence alignments of *L. major* Formins FH2 domain with the other well characterized FH2 domain of formins.
Multiple sequence alignments of Formins FH2 domains have shown in the figure. *L. major* formin A (Q4QE97) and formin B (Q4QAM2) amino acid sequence compared to the characterized formin FH2 domain from human Daam1(Q944D1), mouseDia1 (O08808), *Dictyostelium discoideum* (Q54N00), *Arabidopsis thaliana* (Q9SE97), *Saccharomyces cerevisiae* (P41832, P40450), *Trypanosoma cruzi* (Q4D1V1). Strictly conserved residues are boxed in the

red box, and identical residues are boxed in the blue box. Dashes indicate gaps introduced for

optimal alignment.

Figure S4. **Sequence similarity of *L. major* Formins FH2 domain with other well**

characterized Formin FH2 domains.

Pairwise alignment of *L. major* Formins FH2 domain with other well characterized Formin FH2

domains showing sequence similarity and identity.

Figure S5. SPR analysis of interaction between I777A LmForminA and I802A LmForminB

with the F-actin.

[A, B] The sensorgram has shown the interaction phase of I777A LmForminA [A] and I802A

LmForminB [B], followed by the dissociation phase. The interaction has been seen by an

increase in response in a concentration-dependent manner. [C] SDS-PAGE of low speed actin

cosedimentation for well characterized actin bundler Fascin protein with F-actin. [D] SDS-PAGE

of low-speed actin filament co-sedimentation in the presence of different concentration of I777A

LmForminA (1 µM, 2 µM, 4 µM). The amount of actin coming in the pellet in the presence of

I777A LmForminA represent the actin-bundling. [E] SDS-PAGE image of low-speed actin

filament co-sedimentation in the presence of different concentration of I802A LmForminB (0.5

µM, 1 µM, 2 µM). The amount of actin in the pellet in the presence of I802A LmForminB

represent its actin-bundling. (P for pellet fraction and S for supernatant fraction)

Figure S6. Low speed cosedimentation for co-polymerize I777A LmForminA and I802A

LmForminB with G-actin.

[A, B] SDS-PAGE image of low-speed co-sedimentation assay using 5 µM actin with different

concentration of WT LmForminA and I777A LmForminA (1 µM, 2 µM, 4 µM, 6 µM). Actin

was co-polymerized with LmForminA constructs for 1 hour at room temperature in the F-buffer.

The amount of actin coming in the pellet in the presence of WT LmForminA and I777A

LmForminA represent the actin-bundling. [C, D] SDS-PAGE image of low-speed co-

sedimentation assay using 5 µM actin with different concentration of WT LmForminA and

I802A LmForminB (0.5 µM, 1 µM, 2 µM). Actin was co-polymerized with LmForminA

constructs for 1 hour at room temperature in the F-buffer. The amount of actin coming in the

pellet in the presence of WT LmForminA and I802A LmForminB represent the actin-bundling.

(P for pellet fraction and S for supernatant fraction).

Figure S7. Co-sedimentation for *L. major* formins with the F-actin.

[A, B] SDS-PAGE gel of a co-sedimentation assay showing LmForminA and LmForminB binds

F-actin in a concentration-dependent manner. Fraction of LmFormins bound to the F-actin from

an SDS-PAGE gel image was used for the densitometry analysis (P for pellet fraction and S for

supernatant fraction). [C, D] The dissociation constant was determined by the non-linear curve

fitting.

Figure S8. I777A LmForminA and I802A LmForminB FH2 not able to replace capping

protein from filament barbed end.

[A, B] 0.5 µM G-actin (10% pyrene-labelled) with F-actin seed used in actin elongation with

various concentrations of I777A LmForminA and I802A LmForminB. Mutant LmFormins and

capping protein were added to see the effect on actin filamnet elongation.

Figure S9. High salt concentration reduces the actin bundling activity of LmForminA and

LmForminB FH2.

[A, B] Coomassie-stained SDS-PAGE of the low-speed F-actin cosedimentation using 5 μM

polymerized actin and increasing concentrations of LmForminA and LmForminB respectively in

the presence of 50, 100, 150 mM KCl.

References

Abbasi, K., DuBois, K. N., Dacks, J. B. and Field, M. C. (2011). A novel rho-like protein

tbrhp is involved in spindle formation and mitosis in trypanosomes. *PLoS One* **6**, e26890.

Ambaru, B., Gopalsamy, A., Tammana, T. V. S., Subramanya, H. S. and Gupta, C. M.

(2020). Actin sequestering protein, profilin, regulates intracellular vesicle transport in

Leishmania. *Mol. Biochem. Parasitol.* **238**, 111280.

Angeles Juanes, M. and Piatti, S. (2016). Control of formin distribution and actin cable

assembly by the E3 ubiquitin ligases Dma1 and Dma2. *Genetics* **204**, 205–220.

Barko, S., Bugyi, B., Carlier, M. F., Gombos, R., Matusek, T., Mihályand, J. and Nyitrai,

M. (2010). Characterization of the biochemical properties and biological function of the

formin homology domains of Drosophila DAAM. *J. Biol. Chem.* **285**, 13154–13169.

Baum, J., Tonkin, C. J., Paul, A. S., Rug, M., Smith, B. J., Gould, S. B., Richard, D.,

Pollard, T. D. and Cowman, A. F. (2008). A Malaria Parasite Formin Regulates Actin

Polymerization and Localizes to the Parasite-Erythrocyte Moving Junction during Invasion.

Cell Host Microbe **3**, 188–198.

Castrillon, D. H. and Wasserman, S. A. (1994). diaphanous is required for cytokinesis in

Drosophila and shares domains of similarity with the products of the limb deformity gene.

Development **120**, 3367–3377.

Chalkia, D., Nikolaidis, N., Makalowski, W., Klein, J. and Nei, M. (2008). Origins and

evolution of the formin multigene family that is involved in the formation of actin

filaments. *Mol. Biol. Evol.* **25**, 2717–2733.

Chow, C., Cloutier, S., Dumas, C., Chou, M. N. and Papadopoulou, B. (2011). Promastigote

to amastigote differentiation of Leishmania is markedly delayed in the absence of PERK

eIF2alpha kinase-dependent eIF2alpha phosphorylation. *Cell. Microbiol.* **13**, 1059–1077.

Christensen, J. R., Craig, E. W., Glista, M. J., Mueller, D. M., Li, Y., Sees, J. A., Huang, S.,

Suarez, C., Mets, L. J., Kovar, D. R., et al. (2019). Chlamydomonas reinhardtii formin

FOR1 and profilin PRF1 are optimized for acute rapid actin filament assembly. *Mol. Biol.*

Cell **30**, 3123–3135.

Cvrčková, F., Novotný, M., Pícková, D. and Žárský, V. (2004). Formin homology 2 domains

occur in multiple contexts in angiosperms. *BMC Genomics* **5**, 1–18.

Daher, W., Plattner, F., Carlier, M. F. and Soldati-Favre, D. (2010). Concerted action of two

formins in gliding motility and host cell invasion by Toxoplasma gondii. *PLoS Pathog.* **6**,

1001132.

Daher, W., Klages, N., Carlier, M. F. and Soldati-Favre, D. (2012). Molecular

characterization of toxoplasma gondii formin 3, an actin nucleator dispensable for

tachyzoite growth and motility. *Eukaryot. Cell* **11**, 343–352.

DeWard, A. D. and Alberts, A. S. (2009). Ubiquitin-mediated degradation of the formin mDia2

upon completion of cell division. *J. Biol. Chem.* **284**, 20061–20069.

Drubin, D. G., Miller, K. G. and Botstein, D. (1988). Yeast actin-binding proteins: evidence

for a role in morphogenesis. *J. Cell Biol.* **107**, 2551–2561.

Dutta, P., Das, S. and Maiti, S. (2017). Non diaphanous formin delphilin acts as a barbed end

capping protein. *Exp. Cell Res.* **357**, 163–169.

Evangelista, M., Pruyne, D., Amberg, D. C., Boone, C. and Bretscher, A. (2002). Formins
direct Arp2/3-independent actin filament assembly to polarize cell growth in yeast. *Nat.*
Cell Biol. **4**, 260–269.

Filardy, A. A., Guimarães-Pinto, K., Nunes, M. P., Zukeram, K., Fliess, L., Pereira, L.,
Oliveira Nascimento, D., Conde, L. and Morrot, A. (2018). Human Kinetoplastid
Protozoan Infections: Where Are We Going Next? *Front. Immunol.* **9**, 1493.

Frénal, K. and Soldati-Favre, D. (2009). Role of the Parasite and Host Cytoskeleton in
Apicomplexa Parasitism. *Cell Host Microbe* **5**, 602–611.

Gouet, P., Courcelle, E., Stuart, D. I. and Métoz, F. (1999). ESPript: Analysis of multiple
sequence alignments in PostScript. *Bioinformatics* **15**, 305–308.

Goyard, S., Segawa, H., Gordon, J., Showalter, M., Duncan, R., Turco, S. J. and Beverley,
S. M. (2003). An in vitro system for developmental and genetic studies of Leishmania
donovani phosphoglycans. *Mol. Biochem. Parasitol.* **130**, 31–42.

Gull, K. (1999). The cytoskeleton of trypanosomatid parasites. *Annu. Rev. Microbiol.* **53**, 629–
655.

Gupta, C.M., Ambaru, B. and Bajaj, R. (2020). Emerging Functions of Actins and Actin
Binding Proteins in Trypanosomatids. *Front Cell Dev Biol* **8**, 587685.

Harris, E. S. and Higgs, H. N. (2006). Biochemical analysis of mammalian formin effects on
actin dynamics. *Methods Enzymol.* **406**, 190–214.

Harris, E. S., Li, F. and Higgs, H. N. (2004). The Mouse Formin, FRLα, Slows Actin Filament
Barbed End Elongation, Competes with Capping Protein, Accelerates Polymerization from
Monomers, and Severs Filaments. *J. Biol. Chem.* **279**, 20076–20087.

Harris, E. S., Rouiller, I., Hanein, D. and Higgs, H. N. (2006). Mechanistic differences in

actin bundling activity of two mammalian formins, FRL1 and mDia2. *J. Biol. Chem.* **281**,

14383–14392.

Higgs, H. N. and Pollard, T. D. (1999). Regulation of actin polymerization by Arp2/3 complex

and WASp/Scar proteins. *J. Biol. Chem.* **274**, 32531–32534.

Ivens, A. C., Peacock, C. S., Worthey, E. A., Murphy, L., Aggarwal, G., Berriman, M., Sisk,

E., Rajandream, M. A., Adlem, E., Aert, R., et al. (2005). The genome of the

kinetoplastid parasite, Leishmania major. *Science (80-.).* **309**, 436–442.

Junemann, A., Winterhoff, M., Nordholz, B., Rottner, K., Eichinger, L., Gräf, R. and Faix,

J. (2013). ForC lacks canonical formin activity but bundles actin filaments and is required

for multicellular development of Dictyostelium cells. *Eur. J. Cell Biol.* **92**, 201–212.

Kato, T., Watanabe, N., Morishima, Y., Fujita, A., Ishizaki, T. and Narumiya, S. (2001).

Localization of a mammalian homolog of diaphanous, mDia1, to the mitotic spindle in

HeLa cells. *J. Cell Sci.* **114**, 775–784.

Kovar, D. R. and Pollard, T. D. (2004). Insertional assembly of actin filament barbed ends in

association with formins produces piconewton forces. *Proc. Natl. Acad. Sci. U. S. A.* **101**,

14725–14730.

Li, F. and Higgs, H. N. (2003). The mouse formin mDia1 is a potent actin nucleation factor

regulated by autoinhibition. *Curr. Biol.* **13**, 1335–1340.

Li, F. and Higgs, H. N. (2005). Dissecting requirements for auto-inhibition of actin nucleation

by the formin, mDia1. *J. Biol. Chem.* **280**, 6986–6992.

Li, M. Y., Peng, W. H., Wu, C. H., Chang, Y. M., Lin, Y. L., Chang, G. D., Wu, H. C. and

Chen, G. C. (2019). PTPN3 suppresses lung cancer cell invasiveness by counteracting Src-

mediated DAAM1 activation and actin polymerization. *Oncogene* **38**, 7002–7016.

Lu, J., Meng, W., Poy, F., Maiti, S., Goode, B. L. and Eck, M. J. (2007). Structure of the FH2 Domain of Daam1: Implications for Formin Regulation of Actin Assembly. *J. Mol. Biol.* **369**, 1258–1269.

Maas, R. L., Zeller, R., Woychik, R. P., Vogt, T. F. and Leder, P. (1990). Disruption of formin-encoding transcripts in two mutant limb deformity alleles. *Nature* **346**, 853–855.

Majumder, S. and Lohia, A. (2008). Entamoeba histolytica encodes unique formins, a subset of which regulates DNA content and cell division. *Infect. Immun.* **76**, 2368–2378.

McConville, M. J., Mullin, K. A., Ilgoutz, S. C. and Teasdale, R. D. (2002). Secretory Pathway of Trypanosomatid Parasites. *Microbiol. Mol. Biol. Rev.* **66**, 122–154.

Michelot, A., Guérin, C., Huang, S., Ingouff, M., Richard, S., Rodiuc, N., Staiger, C. J. and Blanchoin, L. (2005). The formin homology 1 domain modulates the actin nucleation and bundling activity of arabidopsis FORMIN1. *Plant Cell* **17**, 2296–2313.

Moseley, J. B. and Goode, B. L. (2005). Differential activities and regulation of Saccharomyces cerevisiae formin proteins Bni1 and Bnr1 by Bud6. *J. Biol. Chem.* **280**, 28023–28033.

Moseley, J. B., Maiti, S. and Goode, B. L. (2006). Formin proteins: Purification and measurement of effects on actin assembly. *Methods Enzymol.* **406**, 215–234.

Nayak, R. C., Sahasrabuddhe, A. A., Bajpai, V. K. and Gupta, C. M. (2005). A novel homologue of coronin colocalizes with actin in filament-like structures in Leishmania. *Mol. Biochem. Parasitol.* **143**, 152–164.

Okwor, I. and Uzonna, J. (2016). Social and economic burden of human leishmaniasis. *Am. J. Trop. Med. Hyg.* **94**, 489–493.

Ozaki-Kuroda, K., Yamamoto, Y., Nohara, H., Kinoshita, M., Fujiwara, T., Irie, K. and Takai, Y. (2001). Dynamic Localization and Function of Bni1p at the Sites of Directed

Growth in Saccharomyces cerevisiae . *Mol. Cell. Biol.* **21**, 827–839.

Pal, D. S., Mondal, D. K. and Datta, R. (2015). Identification of metal dithiocarbamates as a

novel class of antileishmanial agents. *Antimicrob. Agents Chemother.* **59**, 2144–2152.

Pal, D. S., Abbasi, M., Mondal, D. K., Varghese, B. A., Paul, R., Singh, S. and Datta, R.

(2017). Interplay between a cytosolic and a cell surface carbonic anhydrase in pH

homeostasis and acid tolerance of Leishmania. *J. Cell Sci.* **130**, 754–766.

Patel, A. A., Oztug Durer, Z. A., Van Loon, A. P., Bremer, K. V. and Quinlan, M. E. (2018).

Drosophila and human FHOD family formin proteins nucleate actin filaments. *J. Biol.*

Chem. **293**, 532–540.

Petersen, J., Nielsen, O., Egel, R. and Hagan, I. M. (1998). FH3, a domain found in formins,

targets the fission yeast formin Fus1 to the projection tip during conjugation. *J. Cell Biol.*

141, 1217–1228.

Pollard, T. D. (1984). Polymerization of ADP-actin. *J. Cell Biol.* **99**, 769–777.

Ponte-Sucre, A., Gamarro, F., Dujardin, J. C., Barrett, M. P., López-Vélez, R., García-

Hernández, R., Pountain, A. W., Mwenechanya, R. and Papadopoulou, B. (2017). Drug

resistance and treatment failure in leishmaniasis: A 21st century challenge. *PLoS Negl.*

Trop. Dis. **11**,.

Pruyne, D., Evangelista, M., Yang, C., Bi, E., Zigmond, S., Bretscher, A. and Boone, C.

(2002). Role of formins in actin assembly: Nucleation and barbed-end association. *Science*

(80-.). **297**, 612–615.

Rivero, F., Muramoto, T., Meyer, A. K., Urushihara, H., Uyeda, T. Q. P. and Kitayama, C.

(2005). A comparative sequence analysis reveals a common GBD/FH3-FH1-FH2-DAD

architecture in formins from Dictyostelium, fungi and metazoa. *BMC Genomics* **6**, 1–16.

Rizvi, S. A., Neidt, E. M., Cui, J., Feiger, Z., Skau, C. T., Gardel, M. L., Kozmin, S. A. and

Kovar, D. R. (2009). Identification and Characterization of a Small Molecule Inhibitor of

Formin-Mediated Actin Assembly. *Chem. Biol.* **16**, 1158–1168.

Romero, S., Didry, D., Larquet, E., Boisset, N., Pantaloni, D. and Carlier, M. F. (2007).

How ATP hydrolysis controls filament assembly from profilin-actin: Implication for formin

processivity. *J. Biol. Chem.* **282**, 8435–8445.

Sagot, I., Klee, S. K. and Pellman, D. (2002). Yeast formins regulate cell polarity by

controlling the assembly of actin cables. *Nat. Cell Biol.* **4**, 42–50.

Sahasrabuddhe, A. A., Bajpai, V. K. and Gupta, C. M. (2004). A novel form of actin in

Leishmania: Molecular characterisation, subcellular localisation and association with

subpellicular microtubules. *Mol. Biochem. Parasitol.* **134**, 105–114.

Sahasrabuddhe, A. A., Nayak, R. C. and Gupta, C. M. (2009). Ancient Leishmania coronin

(CRN12) is involved in microtubule remodeling during cytokinesis. *J. Cell Sci.* **122**, 1691–

1699.

Schönichen, A., Mannherz, H. G., Behrmann, E., Mazur, A. J., Kühn, S., Silván, U.,

Schoenenberger, C. A., Fackler, O. T., Raunser, S., Dehmelt, L., et al. (2013). FHOD1

is a combined actin filament capping and bundling factor that selectively associates with

actin arcs and stress fibers. *J. Cell Sci.* **126**, 1891–1901.

Scott, B. J., Neidt, E. M. and Kovar, D. R. (2011). The functionally distinct fission yeast

formins have specific actin-assembly properties. *Mol. Biol. Cell* **22**, 3826–3839.

Shimada, A., Nyitrai, M., Vetter, I. R., Kühlmann, D., Bugyi, B., Narumiya, S., Geeves, M.

A. and Wittinghofer, A. (2004). The core FH2 domain of diaphanous-related formins is an

elongated actin binding protein that inhibits polymerization. *Mol. Cell* **13**, 511–522.

Silkworth, W. T., Kunes, K. L., Nickel, G. C., Phillips, M. L., Quinlan, M. E. and Vizcarra,

C. L. (2018). The neuron-specific formin Delphilin nucleates nonmuscle actin but does not

enhance elongation. *Mol. Biol. Cell* **29**, 610–621.

Skillman, K. M., Daher, W., Ma, C. I., Soldati-Favre, D. and Sibley, L. D. (2012).

Toxoplasma gondii profilin acts primarily to sequester G-actin while formins efficiently

nucleate actin filament formation in vitro. *Biochemistry* **51**, 2486–2495.

Stortz, J. F., Del Rosario, M., Singer, M., Wilkes, J. M., Meissner, M. and Das, S. (2019).

Formin-2 drives polymerisation of actin filaments enabling segregation of apicoplasts and

cytokinesis in Plasmodium Falciparum. *Elife* **8**,.

Sunter, J. and Gull, K. (2017). Shape, form, function and Leishmania pathogenicity: from

textbook descriptions to biological understanding. *Open Biol.* **7**,.

Tammana, T. V. S., Sahasrabuddhe, A. A., Bajpai, V. K. and Gupta, C. M. (2010).

ADF/cofilin-driven actin dynamics in early events of Leishmania cell division. *J. Cell Sci.* **123**,

1894–1901.

Tosetti, N., Pacheco, N. D. S., Favre, D. S. and Jacot, D. (2019). Three f-actin assembly

centers regulate organelle inheritance, cell-cell communication and motility in toxoplasma

gondii. *Elife* **8**,.

Tsigankov, P., Gherardini, P. F., Helmer-Citterich, M., Späth, G. F., Myler, P. J. and

Zilberstein, D. (2014). Regulation dynamics of leishmania differentiation: Deconvoluting

signals and identifying phosphorylation trends. *Mol. Cell. Proteomics* **13**, 1787–1799.

Vasconcelos, E. J. R., Pacheco, A. C. L., Gouveia, J. J. S., Araújo, F. F., Diniz, M. C.,

Kamimura, M. T., Costa, M. P., Araújo-Filho, R. and Magalhães de Oliveira, D.

(2008). Actin-Interacting Proteins in flagellated pathogenic Leishmania spp.: a genome-

based bioinformatics report on profilins, formins and katanins. *Int. J. Funct. Inform.*

Personal. Med. **1**, 234–252.

Von Dippe, P. and Levy, D. (1982). Analysis of the transport system for inorganic anions in

normal and transformed hepatocytes. *J. Biol. Chem.* **257**, 4381–4385.

Wasserman, S. (1998). FH proteins as cytoskeletal organizers. *Trends Cell Biol.* **8**, 111–115.

Xu, Y., Moseley, J. B., Sagot, I., Poy, F., Pellman, D., Goode, B. L. and Eck, M. J. (2004).

Crystal structures of a formin homology-2 domain reveal a tethered dimer architecture. *Cell*

116, 711–723.

Yamashiro, S., Yamakita, Y., Ono, S. and Matsumura, F. (1998). Fascin, an actin-bundling

protein, induces membrane protrusions and increases cell motility of epithelial cells. *Mol.*

Biol. Cell **9**, 993–1006.

Zeller, R., Haramis, A. G., Zuniga, A., McGuigan, C., Dono, R., Davidson, G., Chabanis, S.

and Gibson, T. (1999). Formin defines a large family of morphoregulatory genes and

functions in establishment of the polarising region. *Cell Tissue Res.* **296**, 85–93.

Zilberstein, D. (2015). Proteomic analysis of posttranslational modifications using iTRAQ in

Leishmania. *Methods Mol. Biol.* **1201**, 261–268.

Zimmermann, D., Morganthaler, A. N., Kovar, D. R. and Suarez, C. (2016). In vitro

biochemical characterization of cytokinesis actin-binding proteins. In *Methods in Molecular*

Biology, pp. 151–179. Humana Press Inc.

Ziske, M. A., Pettee, K. M., Khaing, M. N., Rubinic, K. and Eisenmann, K. M. (2016).

SMIFH2-mediated mDia formin functional inhibition potentiates chemotherapeutic

targeting of human ovarian cancer spheroids. *Biochem. Biophys. Res. Commun.* **472**, 33–39.

993   **Figure 1**

995   **Figure 2**

997   Figure 3

999   Figure 4

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

Figure 5

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

Figure 6

Figure 7

1030

1031   **Table S1**

Table S1. List of the primers used in this study			
Primer	Primer sequence (5'→3')	Restriction site	Purpose
P1	5'-GGTAC<u>GGATCC</u>GCGAGCGGCGCAGCAGCC-3'	BamHI	Cloning of LmForminA into pET28a and to check expression of LmForminA
P2	5-GGCCC<u>AAGCTT</u>TTAGCTCTGCCGCCGCTGCTTG-3	HindIII	
P3	5'- GGTAC<u>GGATCC</u>AAGCCGAAGCCACAGTACACC-3'	BamHI	Cloning of LmForminB into pET28a and to check expression of LmForminB
P4	5'-GAG<u>AAGCTT</u>CTACTGCTCCGTCGCTTCCGG-3'	HindIII	
P5	5'-GTGACCGAAACGTCGGCGCTGTGCTGAAGTTTATTCGGC-3'	-	Site-directed mutagenesis of I777A
P6	5'-GCCGAATAAACTTCAGCACAGCGCCGACGTTTCGGTCAC-3'	-	
P7	5'-CAACGGTTGCAGAACATGGGCGCCGCCCTCAAGCGCGTACAG-3'	-	Site-directed mutagenesis of I802A
P8	5'-CTGTACGCGCGCTTGAGGGCGGCGCCCATGTTCTGCAACCGTTG-3'	-	

1032

1033

1034   **Figure S1**

1035

1036   **Figure S2**

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048 **Figure S3**

1049

1050 **Figure S4**

	LmForminA				LmForminB		
Formin	Species	Identity	Similarity	Formin	Species	Identity	Similarity
TC00	*T. cruzi*	40.9%	57%	TC00	*T. cruzi*	18.1%	34.8%
Bni1	*S.cerevisiae*	23.2%	40.1%	Bni1	*S.cerevisiae*	21.3%	35.3%
Bnr1	*S.cerevisiae*	17.9%	33.5%	Bnr1	*S.cerevisiae*	20.6%	36.9%
cdc12	*S.pombe*	20.1%	37.0%	cdc12	*S.pombe*	19.1%	36.1%
dDia2	*D.discoideum*	24.7%	44.8%	dDia2	*D. discoideum*	20.7%	39.0%
AFH1	*A.thaliana*	22.3%	42.1%	AFH1	*A.thaliana*	24.2%	43.5%
mDia1	*M.musculus*	20.8%	36.1%	mDia1	*M. musculus*	21.8%	39.7
Daam1	*H.sapiens*	22.6%	44.2%	Daam1	*H.sapiens*	21.4%	42.6%

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060   **Figure S7**

1061

1062

1063   **Figure S8**

1064

1065

1066   **Figure S9**

1067