One gene - two proteins: The C-terminus of the prototypical M2 muscarinic

receptor localizes to the mitochondria

Irene Fasciani[a†], Francesco Petragnano[a†], Ziming Wang[b], Ruairidh Edwards[c], Narasimha Telegu[b], Ilaria

Pietrantoni[a], Henrik Zauber[b], Mario Rossi[c], Ulrike Zabel[d], Marlies Grieben[b], Maria Eirini Terzenidou[c],

Jacopo Di Gregorio[a], Cristina Pellegrini[a], Silvano Santini Jr[e], Anna Rita Taddei[f], Barbel Pöhl[b], Stefano

Aringhieri[g], Gabriella Aloisi[a], Francesco Marampon[h], Alexandra Roman[b], Sebastian Diecke[b], Vincenzo

Flati[a], Franco Giorgi[f], Fernanda Amicarelli[e], Andrew B. Tobin[c], Marco Scarselli[g], Kostas Tokatlidis[c],

Martin J Lohse[b,d,i]*, Roberto Maggio[a]*, Paolo Annibale[b]*

[a] Department of Biotechnological and Applied Clinical Sciences, University of L'Aquila, 67100

L'Aquila, Italy.

[b]Max Delbrück Center for Molecular Medicine, 13125 Berlin, Germany.

[c]Centre for Translational Pharmacology, Institute of Molecular, Cell and Systems Biology, College of

Medical, Veterinary and Life Sciences, University of Glasgow, G128QQ Glasgow, UK.

[d]Institute of Pharmacology and Toxicology, University of Würzburg, 97078 Würzburg, Germany.

[e]Department of Life, Health and Environmental Sciences, University of L'Aquila, 67100 L'Aquila,

Italy.

[f]Section of Electron Microscopy, Great Equipment Center, University of Tuscia, Viterbo, Italy.

[g]Department of Translational Research and New Technology in Medicine, University of Pisa, 56126

Pisa, Italy.

[h]Department of Radiotherapy, University of Roma La Sapienza, 00151 Roma, Italy

[i]ISAR Bioscience Institute, 82152 Munich, Germany

[†] equal contribution

*email: m.lohse@mdc-berlin.de;

*email: roberto.maggio@univaq.it;

*email: paolo.annibale@mdc-berlin.de

27
28
29
30

Abstract

Muscarinic acetylcholine receptors are G protein-coupled receptors (GPCRs) activated by the

endogenous neurotransmitter acetylcholine. We show here that the carboxyl terminal fragment of the

muscarinic M_2 receptor, containing the transmembrane regions 6 and 7 (M_2tail), is expressed by virtue

of an internal ribosome entry site. Radioligand binding, biochemical and molecular biology experiments,

combined to advanced fluorescence microscopy reveal that the M_2tail, whose expression is upregulated

in cells undergoing integrated stress response, does not follow the normal route to the plasma membrane,

but is almost exclusively sorted to the mitochondria inner membrane: here it controls oxygen

consumption, cell proliferation and the formation of reactive oxygen species via reduction of oxidative

phosphorylation. Genetic editing of the key methionine where cap-independent translation occurs,

reveals the physiological role of this process in influencing cell proliferation and oxygen consumption.

The expression of the carboxyl-terminal of a GPCR, capable of regulating mitochondrial function,

constitutes a hitherto unknown mechanism that cells may use for controlling their metabolism under

variable environmental conditions, notably unrelated to the canonical signalling function of the GPCR

at the plasma membrane.

Muscarinic receptors include five structurally related receptors, belonging to class A of the G protein-coupled

receptor (GPCR) family[1]. They are comprised of seven trans-membrane domains (TM domains 1-7) connected

by intra and extracellular loops. Of these structural elements, the third intracellular (i3) domain is the one

conferring a coupling specificity for G proteins, β-arrestin recruitment and endocytic internalization [2].

However, only a small portion of the i3 loop is required for receptor function, and for this reason large deletions

of this segment have no effect on receptor activation and internalization [3].

We could previously show that the M_2 receptor can be split at the level of the i3 loop into two segments, each

behaving as an autonomous folding unit capable of targeting the plasma membrane and surprisingly

reconstituting a physiologically active receptor when expressed together in transfected cells [4]. In the context

of these studies, a mutant of the receptor carrying a stop codon in position 228 was surprisingly observed to

retain (albeit reduced) ligand binding in radioligand binding assays [11]. This finding cannot be explained by the

fact that the insertion of the stop codon results in the expression of an N-terminal receptor fragment, constituted

by the TM regions 1-5, since this fragment alone has been shown to be not able to bind muscarinic ligands and

to trigger signal transduction [12]. Therefore, an alternative explanation for the behavior of stop-codon mutant

228 is needed. Since the N-terminal fragment can reconstitute a physiologically active receptor when co-

transfected with the C-terminal fragment, a plausible possibility is that the C-terminal segment of the receptor

can be independently produced and reconstitute an active receptor with the N-terminal portion on the cell

membrane. This scenario opens the interesting question of how the C-terminal fragment can be translated

endogenously as an independent polypeptide.

In eukaryotes, the most common mechanism of translation initiation is the cap-dependent process, which

requires the interaction of several key proteins at the 5'-end (5′ cap) of mRNA molecules. However, in recent

years, it has been demonstrated that the presence of internal ribosome entry site (IRES) sequences, which are

common features of virus replication[5], can mediate cap independent translation of eukaryotic RNA(s) [6]. This

presents the suggestive possibility that a cap-independent translation of the M_2 C-terminal domain indeed

occurs in this mutant, opening a number of important questions about the molecular processes involved in the

synthesis and subcellular behaviour of the C-terminal domain fragment.

GPCRs are normally distributed at the plasma membrane, where, after activation by an agonist and coupling

to G proteins, regulate a number of intracellular effectors. Nevertheless, non-canonical localizations of GPCRs

have also been described, such as in endosomes [7], cell division compartments (centrosomes, spindle midzone

and midbodies)[8], and mitochondria [9] [10], underpinning non-traditional roles for this protein family.

In this work, we therefore set out to explore whether the expression of the C-terminal domain of the M_2

muscarinic receptor, a region containing the TM regions 6 and 7, occurs naturally by a cap-independent

translation mechanism. Combining radio-ligand binding, molecular biology approaches, advanced

fluorescence imaging and genetic editing of human induced pluripotent stem cells (hiPSCs) we have observed

that the C-terminal domain of this receptor is indeed expressed as an independent polypeptide. Surprisingly,

using bimolecular fluorescence complementation, in-vitro mitochondrial import assays and cell metabolism

assays we observe that the C-terminal domain is strongly targeted to the cell mitochondria. Importantly, this

mitochondrial targeting has functional ramifications since both cap-dependent mediated overexpression of the

C-terminal domain alone, as well as cap-independent mediated translation starting from the mRNA of the full

length M_2 receptor, resulted in a sizable reduction of oxidative phosphorylation.

In summary, our study reports the new observation that alternative reading of a GPCR receptor gene can

translate into a fragment of the receptor that serves a cellular function entirely different from canonical GPCR

signalling.

Results

Ligand binding and functional properties of M_2 receptors mutants.

To investigate and distinguish the functional roles of the two known N-terminal (TM1-5) and C-terminal

(TM6-7) domains of the M_2 muscarinic receptor, we expressed C-terminally truncated variants of the receptor

in COS-7 cells. We first observed that the M_2stop228 construct, a receptor mutant bearing a stop codon at the

beginning of the i3 loop in position 228 (but containing the downstream sequence, i.e. leading to the

transcription of an mRNA of the same length as that of the wt M_2R) (Fig. 1a and Supplementary Fig. 1),

exhibited unexpected binding activity [11] albeit at a very low level of expression, i.e. 45 fmol/mg of protein.

Both K_D value of [^{3}H]NMS (N-methylscopolamine, an antagonist) calculated from direct saturation

experiments, and the IC_{50} values, calculated in competition experiments for the agonist carbachol, were in fact

similar to those calculated for the wild type M_2 receptor (Table 1). Furthermore, the M_2stop228 receptor was

found functionally active as it was capable to inhibit adenylyl cyclase [11] and increase ERK phosphorylation

(Supplementary Fig. 2b), even though the extent of response was reduced if compared to the wild type M_2

receptor. On the other hand, if a stop codon is inserted upstream or downstream of position 228, e.g. at positions

196 or 400, no radioligand binding can be observed (Fig. 1a and Supplementary Table 1).

As we demonstrated in an earlier report [12], another receptor fragment whose sequence runs only until position

283 (i.e. comprising only TM1-5) of the corresponding wt receptor, called M_2trunk(1-283) (Supplementary

Fig. 1), did not show any radio-ligand binding activity upon expression in COS-7 cells, despite comprising the

first five trans-membrane regions of the wild type M_2 receptor, as the M_2stop228 construct (Table 1).

Consistently, a construct containing only the M_2 receptor sequence until amino acid 228 (M_2trunk(1-228)), did

not show any radio-ligand binding (**Table 1**).

On the other hand, we observed binding and downstream signaling ability in co-transfection experiments of

the M_2trunk(1-283) and M_2tail(281-466) (**Table 1**), the latter being a construct containing only the sequence

between aminoacid 368 and the end of the M_2 receptor, i.e. comprising TM6 and 7. Therefore, co-expression

of the M_2trunk(1-283) (TM1-5) and M_2tail(368-466) (TM6-7) reconstitutes a functional full length receptor at

the cell membrane (**Fig. 1b**).

These data suggest that, for the M_2stop228 receptor to display binding and functional activity, the receptor

sequence downstream of the stop codon in position 228 (containing TM6 and 7) has (i) to be translated and

(ii) to presumably interact with the first five TM regions of the M_2 receptor. This notion is graphically

summarized in **Fig. 1b**. For this hypothesis to be true, however, the C-terminal portion of the receptor needs

to be expressed despite the presence of stop codons in the i3 loop in our M_2stop228 mutant. Indeed, we

confirmed expression of the C-terminal fragment by fusing an enhanced Green Fluorescent Protein (EGFP) at

the C terminus of several receptor mutants (**Fig. 1c**).

Interestingly, though, while cells expressing the M_2-EGFP displayed the normal, predominant plasma

membrane expression (**Fig. 1c**), cells expressing the M_2stop228-EGFP displayed a visible intracellular green

fluorescence, which is in stark contrast to cells expressing M_2stop400-EGFP, which do not show any

fluorescence, in line with the non-transfected control.

To further analyze the atypical behavior of M_2stop228-EGFP, which by displaying green fluorescence

confirms that the sequence downstream of the stop in position 228 is indeed translated, we generated a frame

shift mutant: we inserted a single guanine nucleotide in the sequence of the M_2-EGFP, changing the methionine

in position 368 into an aspartate, which in turn generates an in-frame stop codon at position 370. We named

the construct, which notably still displays an EGFP signal in transfected cells, M_2fr.sh.-EGFP (**Fig. 1c**). The

observation that the mutant retains EGFP fluorescence is most remarkable, since the EGFP is now out of the

original reading frame and it can only be properly translated by a cap-independent translation mechanism,

starting from a methionine in the new reading frame.

Furthermore, the observation that both M_2stop228-EGFP and M_2fr.sh.-EGFP retain EGFP fluorescence,

whereas M_2stop400-EGFP does not, indicates not only that a cap independent translation mechanism occurs,

but also that the starting methionine for the process must be downstream of stop 228 and upstream of stop 400,

leaving as a leading candidate the third in-frame methionine of the i3 loop. This is in line with the radio-ligand

binding data from co-expression of receptor mutants carrying a double stop codon, the first in position 228 and

the second in positions ranging from 248 to 400, including mutations of the first (M248) and second (M296)

in-frame methionines in the third i3 loop (**Supplementary Table 1, Fig. 1a** and **Supplementary Fig. 2a**),

further pointing to M368 as the most likely site for the cap-independent initiation of translation of the C-

terminal domain.

Immunodetection of the M_2 carboxyl terminal fragment

To further characterize the expression of the C-terminal fragment and verify the hypothesis of cap-independent

translation, we tagged several of the M_2 receptors mutants described above with a myc-tag for immune

detection. The constructs were subcloned in a bicistronic vector backbone, allowing also the expression of the

red fluorescent protein mRuby2 as a fluorescent reporter, in order to verify comparable expression levels.

HEK293 cells were therefore transfected with the wild type M_2 receptor M_2-myc, the C-terminal fragment

starting at the third in-frame methionine M_2tail(368-466)-myc, a mutant of the M_2 receptor with the third in-

frame methionine mutated to an alanine, M_2M368A-myc that as discussed above should affect the cap-

independent translation, and finally the M_2stop400-myc; the constructs were resolved by SDS-PAGE and

assayed by Western blotting, as displayed in **Fig. 2a**. Based on the amino acid sequence, the expected molecular

weights for the full length receptor and the C-terminal fragment (inclusive of the myc tag) are ~75 kDa and

~13 kDa respectively.

Interestingly, M_2tail(368-466) runs with two clear bands ~20 and ~15 kDa (**Fig. 2a**), indicating the presence

of post-translational modifications and protein processing (see **Discussion)** and thus defining the qualitative

pattern expected for the M_2 carboxyl terminal fragment starting at the third in-frame start codon (M368). The

full-length receptor was observed, as expected, at about ~75 kDa together with commonly observed higher

order oligomers[13]. Notably, the M_2 receptor clearly displays the lower MW band pair also seen for M_2tail(368-

466). The strongest confirmation for a cap-independent translation process is found in the M_2M368A: here the

~20 kDa band disappears. In line with what we observed for its EGFP tagged version, the M$_2$stop400 does not

exhibit any band, comparable to the negative control using untransfected cells. The absence of any band further

excludes the possibility of a downstream cap-independent translation site yielding a fragment originating from

the next in frame methionine (406) (Fig. 2a). Altogether these results support the notion that the band at ~20

kDa observed in the wild type M$_2$ receptor lane is the bona-fide tail fragment originated by a cap-independent

translation mechanism, and that the origin of this process is at the third in frame methionine (M368) of the M$_2$

receptor.

We then tested whether a similar phenomenon, namely the translation of the C-terminal portion of the M$_2$

receptor, could be observed endogenously in physiological settings. We therefore prepared protein extracts

from heart, brain and spleen tissues from a line of knock-in mice (see Methods), designed to have expression

of a Chrm2/tdtomato fusion protein driven by the endogenous M$_2$ muscarinic acetylcholine receptor

promoter/enhancer sequences. The heart and brain are known to express sizable levels of the M$_2$ receptor,

whereas the spleen, where no expression of the M$_2$ receptor is reported in the proteomic databases, was used

as a negative control.

The use of a transgene tag was necessary, since antibody against the M$_2$ receptor are notoriously ineffective,

in particular when targeted at the C-terminal portion of the receptor. Instead, the tdtomato fluorescent protein

can be targeted by a specific polyclonal antibody, as shown by the Western blot displayed in Fig. 2b. As

expected, the full-length receptor-tdtomato fusion was observed at ~130 kDa (~75 kDa contributed by the

receptor and ~55 kDa contributed by tdtomato).

Notably, both in the heart and brain, but not in the spleen, of these knock-in animals we observed a band

doublet at ~75 kDa, which corresponds to the sum of the molecular weights of the tail fragment (~20 kDa, see

above and Fig. 2a) and the molecular weight of tdtomato.

Testing bicistronic constructs for the presence of an IRES.

To explain the translation of the carboxyl terminal fragment of the M$_2$ receptor, a mechanism enabling a cap-

independent initiation of protein synthesis should be hypothesized [5,6]. Accordingly, an IRES should be found

downstream of the artificial stop codon 228 and upstream of the third in-frame methionine (codon 368) of the

i3 loop. Before testing this hypothesis, we first ruled out several alternative mechanisms that could lead to the

translation of M2tail, such as stop codon read-through, termination-reinitiation and alternative splicing in the

translation of M2tail, as discussed in the **Supplementary Information** and in **Supplementary Figure 3**).

Then, to test the hypothesis that an IRES is leading to the cap-independent translation of the C-terminal domain,

we prepared a fluorescence reporter construct in which the i3 loop of M_2 (from nucleotide 685 (a.a. 228) to

nucleotide 1101 (a.a. 368)) was inserted between the two fluorescent proteins Sirius and EGFP (**Fig. 3a**). This

construct, named Sirius-M_2i3(417n)-EGFP, would translate Sirius according to the classical cap-dependent

mechanism, and, at the same time, it would drive EGFP translation only if the M_2i3 loop contains an IRES

sequence, effectively acting as a bicistronic expression cassette. We transfected several cell lines with this

construct, all positive for the blue and green fluorescence **(Supplementary Fig. 4)**. HEK293, used throughout

our study, also appeared positive for both the blue and green fluorescence, as shown in the inset of **Fig. 3a**.

To rule out the possibility that EGFP translation from the Sirius-M_2i3(417n)-EGFP construct could be due to

read-through and termination-re-initiation mechanisms, a mRNA hairpin loop was inserted in the sequence

after the stop codon of the Sirius protein. The resulting construct (named Sirius-H-M_2i3(417n)-EGFP) showed

only a slight reduction of the EGFP/Sirius with respect to the value measured for the analogous reporter without

hairpin (~90%) (**Fig. 3a**). In contrast the Sirius-H-EGFP plasmid, where the entire M_2i3 loop after the Sirius

was removed, showed a negligible EGFP/Sirius ratio (below 5%), underlining the importance of the i3 loop

for EGFP expression (**Fig. 3a**). Notably, when the hairpin was inserted within the i3 loop of the full length M_2-

EGFP, upstream of the M368, it was still possible to observe GFP fluorescence (**Supplementary Fig. 5**).

To narrow down the position of the IRES sequence we divided the M_2 i3 loop in three parts and the resulting

nucleotide sequences, 685-699, 685-1071 and 685-699/1072-1101, were inserted downstream of the hairpin

loop in three different constructs (**Fig. 3a** and **Supplementary Fig. 6**).

EGFP fluorescence was not significantly rescued upon insertion of the first 15 nucleotides (685-699) of the i3

loop after the hairpin, nor by the addition of the subsequent 372 nucleotides (685-1071). However, when the

last portion of the M_2 i3 loop was inserted in the construct after the Hairpin, leading to Sirius-H-M_2i3(15n/30n)-

EGFP, ~110 % of the value measured for the Sirius-M_2i3(417n)-EGFP plasmid was recovered (**Fig. 3a**).

These results strongly suggest that the IRES is located in a region of the i3 loop sequence immediately upstream

of the third methionine (M368) (see **Fig. 1a**), as also confirmed from ribosome profiling data from human left

ventricle tissue, displaying increased p-sites as well as overall ribosomal coverage in correspondence of this

region. This conclusion is further supported by the finding that when each of the three in-frame i3loop

methionines (namely M248, M296 and M368) after the stop codon 228 of the M_2stop228 were mutated to an

additional stop codon, only the mutation of M368 to a stop codon proved capable of completely abolishing

[^{3}H]NMS binding (**Supplementary results** and **Supplementary Fig. 1 and Supplementary Table 1**). These

binding results can furthermore rule out read-through and termination re-initiation: if these mechanisms were

responsible for translation of the C-terminal of the M_2stop228, suppression of the first methionine would have

abolished the binding. Finally, the origin of the carboxyl terminal fragment of M_2 at the third in-frame

methionine (M368) is also supported by the key observation that the mutant M_2-M368A does not display the

~20 kDa band for the M_2 C-terminal fragment as compared to the wild type M_2 receptor in the Western blot of

Fig. 2a.

Based on this evidence, we mutated the conserved residues highlighted in yellow in **Supplementary Fig. 6a**,

in order to check for their relevance in recruiting the ribosome on the M_2 i3 loop sequence, as shown in **Fig.**

3b. As discussed in detail in the **Supplementary Results**, we found that the four nucleotides AAGA1090-

1093 next to the methionine 368 play a pivotal role in recruiting the ribosome on the M_2 mRNA and in

translating the M_2 C-terminal fragment.

Generalization to other members of the GPCR superfamily

In order to determine if the observed behavior is specific to the M_2 receptor, or could highlight a feature

common to other members of the GPCR superfamily, we repeated several of our experiments in another GPCR,

the related muscarinic M_3 receptor. The M_3 displays an in-frame methionine at the end of the i3loop, in position

481 (**Supplementary Fig. 6a)**. We then inserted a stop codon in position 273, generating the construct

M_3stop273 that still displayed radio-ligand binding. Several other mutants, mimicking the single and double

stop mutations performed for the M_2 displayed a behavior consistent with M481 being the start point of a cap-

independent translation of the C-terminal also for the M_3 receptor (**Supplementary Tables 2 and 3)**, thus

indicating that IRES-mediated cap-independent translation of the C-terminal domain could be a phenomenon

not restricted solely to the M_2 receptor, but may be shared at least by some other members of the GPCR

superfamily. The presence of an IRES sequence upstream of M481 further appears to be supported by the

observation that 558 nucleotides from the i3 loop of the M_3 receptor (from nucleotide 880 to 1437) were

inserted in the same Sirius-EGFP reporter construct as used for the M_2 (**Supplementary Fig. 6b**). This appears

to be indeed supported by the sequence-alignment of the five muscarinic receptors. The alignment of the last

portion of the M_2 i3 loop (from nucleotide 1072 to 1104) with regions adjacent to the second (M_1), fourth (M_3

and M_4) and first (M_5) in-frame Methionines, clearly shows that some nucleotide residues are conserved

(**Supplementary Fig. 6a**). Overall these data underline that cap-independent translation could be a more

general feature of GPCRs mRNAs.

Localization of the M_2 C-terminal fragment generated by cap-dependent expression under the control

of a viral promoter

After having established that the expression of the M_2 C-terminal domain is regulated by an IRES and that its

translation starts at M368, we went on to study the role it might play in the cell. For this purpose, M_2tail(368-

466) fused to the fluorescent protein EGFP in pcDNA3, that is under the control of a strong CMV promoter,

was expressed in HEK293 cells in order to study the subcellular localization of this domain. Confocal

microscopy demonstrated a striking mitochondrial localization of the M_2tail(368-466)-EGFP construct (**Fig.**

4a). We confirmed this observation by co-staining HEK293 cells with a conventional mitochondrial stain

(Mitotracker Deep Red) and by co-expressing M_2tail(368-466)-EGFP with Tom20-mCherry, a fluorescently

labeled construct targeted to the outer mitochondrial membrane (**Supplementary Fig. 7a**). The observed

mitochondrial localization of the M_2tail(368-466)-EGFP appeared in stark contrast to the canonical cellular

localization of the M_2 receptor, as shown in **Fig. 4b** where the M_2 muscarinic receptor was C-terminally labeled

with the red fluorescent protein mRuby2. Interestingly, very little expression of M_2tail(368-466)-EGFP was

observed on the plasma membrane, as we confirmed by fluorescence correlation spectroscopy[14]

(**Supplementary Fig. 7b-d**). By contrast, when the M_2tail(368-466)-EGFP was co-expressed with M_2stop228,

a modest localization could eventually be detected also on the cell membrane thanks to a reduction in diffusion

coefficient of M_2tail(368-466)-EGFP, suggesting a chaperon activity of the M_2 N-terminal domain (M_2trunk)

(**Supplementary Fig. 7e-g**) and hence explaining the [^{3}H]NMS binding observed in cells transfected with

either M_2stop228 or cotransfected with M_2trunk(1-228) and M_2tail(368-466) (**Table 1**). Likewise, constructs

such as M_2tail(368-466)-mRuby2 (a red monomeric fluorescent protein) and M_2tail(368-466)-tdTomato (a

tandem-dimer red fluorescent protein) also displayed clear mitochondrial localization (**Supplementary Fig.**

7h-i), ruling out bias due to the fluorophore used.

Localization of the M$_2$ C-terminal fragment generated by cap-independent expression under the control

of the i3loop IRES.

We then sought to express the C-terminal fragment of the M$_2$ receptor under the control of the i3loop IRES

rather than a viral promoter, to further exclude the possibility that the observed localization of the M$_2$ carboxyl

terminal fragment were due to overexpression of the construct. HEK293 cells were transfected with the

construct M$_2$-mRuby2-STOP-M$_2$-i3-tail-EGFP which contains the full wild type M$_2$ receptor, mRuby2-labeled

at the C-terminal, fused to the C-terminal part of the M$_2$ receptor starting at nucleotide 685 of the i3 loop,

following the stop codon of mRuby2, and ending with a C-terminal EGFP sequence. Through this experimental

strategy, the expression of M$_2$tail(368-466)-EGFP is regulated by the IRES, and the mechanism of C-terminal

fragment production is therefore close to physiological. The expression of EGFP confirmed that the *in vivo*

production of the M$_2$tail(368-466) is regulated by the IRES within the i3 loop. In addition, the processed

receptor fragment was eventually observed to be predominantly confined to the cytosol and the mitochondrial

network (**Fig. 4c**).

Notably, when the M$_2$-EGFP fusion protein was expressed and the cells were co-stained with Mitotracker, a

faint EGFP signal was also observed in the mitochondrial regions (**Supplementary Fig. 8a**), although the

expression of the full-length receptor masks this effect and makes quantification difficult. On the other hand,

Fig. 4d shows that, when the M$_2$fr.sh-EGFP mutant was expressed in HEK293 cells, the IRES generated C-

terminal fragment of the M$_2$ receptor still localized to the mitochondria. As mentioned above, in this construct

the masking effect of the EGFP fused to the full-length receptor is removed (see also **Supplementary Fig. 1**).

Similar mitochondrial localization was also observed when we transfected HEK293 cells with M$_2$fr.sh-

mRuby2 (**Supplementary Fig. 8b**), ruling out possible effects due to the specific fluorophore used as a tag.

We obtained comparable findings also in COS-7 cells (**Supplementary Fig. 8c**).

Mitochondrial import of the C-terminal fragment of the M$_2$ receptor was further verified by immunoblotting

of protein extracts from purified mitochondria of HEK293 cells transfected with myc-tagged M$_2$ receptor,

M$_2$tail(368-466) and M$_2$M368A. **Fig. 4e** shows that in purified mitochondria (unlike in the cytosolic and

microsomal supernatants originating from the purification process, **Supplementary Fig. 8d**), only the protein

bands associated to the M_2 C-terminal fragment were visible.

Notably, no mitochondrial import was visible for the M_2M368A, confirming that the third in frame methionine

(M368) in the i3 loop of the M_2 receptor is necessary for the C-terminal fragment production and its localization

in the mitochondria.

A consistent result was observed when mitochondria were purified from the heart and spleen of the

Chrm2/tdtomato mice **Fig. 4f** (the cytosolic fraction is shown in **Supplementary Fig. 8e**). A band at ~75 kDa,

matching the molecular weight of the M_2 C-terminal domain, was clearly visible. We conclusively confirmed

that this is the fingerprint associated to the M_2 C-terminal fragment, by purifying mitochondria from HEK293

cells expressing the construct M_2tail(368-466)-tdtomato. **Fig. 4g** (cytosolic fraction is shown in

Supplementary Fig. 8e) shows a ~75 kDa protein band matching closely the bands observed in mitochondria

from mice tissues, in addition to lower molecular weight bands most likely arising from degradation of the

tdtomato tag.

Mitochondrial localization of the M_2tail originating from the wild type M_2 receptor was finally confirmed by

relative abundance estimates of EGFP and M_2 receptor from parallel reaction monitoring (PRM)-based mass

spectrometry measurements **Supplementary Fig. 9**. Samples were prepared by generating mitochondrial and

cytosolic fractions (**Methods**) from HEK293 cells, expressing M_2-EGFP and a set of control constructs,

including M_2tail(368-466)-EGFP, pmito-EGFP (an EGFP fused to the mitochondrial leading sequence of the

cytochrome c oxidase subunit 8A (COX8A)), lifeact-EGFP (a cytosolic protein) and Epidermal Growth Factor

receptor fused to EGFP (EGF-EGFP). A 15-aminoacid long peptide originating from EGFP was screened. The

bands intensities of the cytosolic fraction, were normalized to consider both the overall loading variability as

well as the relative transfection efficiencies among the samples. Notably, we observed a mitochondrial

enrichment of the EGFP derived peptide as well the receptor-specific peptide originating from the M_2 receptor,

in agreement with the positive mitochondrial localization control.

Mechanism of import of the M_2 carboxyl terminal fragment into the mitochondria and localization

within the organelle

To gain further insight into the dynamics of the M_2 C-terminal fragment import within the mitochondria, we

first monitored the kinetics of colocalization between mitochondria and EGFP in cells transfected with

M_2tail(368-466)-EGFP. In cells observed for several hours, starting at 5h after transfection (Fig. 4h), the

construct was found initially expressed in the perinuclear region, likely the endoplasmic reticulum, indicating

that at least the initial step of subcellular trafficking of the M_2tail(368-466)-EGFP is the same as for the full

length receptor[15]. At this time (5h post-transfection), almost no colocalization with the Mitotracker-stained

mitochondria was visible. However, at 6h post-transfection, the colocalization level increased significantly,

and the M_2 C-terminal fragment was either found in the cytosol or in the mitochondria. At 9h post-transfection

the colocalization at the mitochondria was well visible. We quantified the colocalization level over time using

the Manders' coefficients (See **Methods**). The resulting chart, indicates that almost half of the mitochondrial

import occurred within 6h post-transfection.

This value is qualitatively in agreement with what we observed using an in-vitro assay to determine the import

of the M_2 carboxyl terminal into isolated mitochondria (see Methods). To this end, we used mitochondria form

S. cerevisiae, which is a robust and well-studied system for protein import assays. Supplementary Fig. 10a

shows the autoradiography experiment of isolated, intact, mitochondria incubated with 35S-labelled full length

muscarinic M_2 receptor or M_2tail(368-466), made using a reticulocyte in vitro translation system. The data

show that the full-length receptor was not imported into mitochondria. On the other hand, the M_2tail(368-466)

displays a clear mitochondrial import already after 5 minutes and increasing over time, up to 30 minutes.

Interestingly, this import (Supplementary Fig. 10b) appears to be independent upon the transporters of the

outer membrane Tom5, Tom6, Tom7 and Tom70, whereas the import of an inner membrane protein, the

ADP/ATP carrier (AAC), used here as a control, depends on Tom70. Upon extraction with carbonate most of

M_2tail(368-466) was found in the pellet, and therefore presumably integrated in a membrane. Furthermore Blue

native analysis suggests that the mitochondrially imported M_2tail(368-466) is not a monomer but rather it is

part of a larger complex of approximately 180 kDa Supplementary Fig. 10c.

All together these data indicate that the import of the M_2tail(368-466) leads to its localization in one of the

mitochondrial membranes, to form a complex with other native mitochondrial proteins.

To further discriminate if the construct is on the outer or inner mitochondrial membrane, we set up a

fluorescence assay based upon GFP fluorescence complementation[16]. HEK293 cells were first transfected with

two constructs: M_2tail(368-466)-GFP11, in which a short 16 amino acids peptide of GFP was fused at the C-

terminus of M_2tail(368-466), and mito-GFP1-10, in which the COX8A was N-terminally fused to the other

GFP fragment containing the remaining 10 beta strands (See Methods). Supplementary Figure 10d shows

the successful recombination of the two within the mitochondrial matrix. We then applied this approach to the

full length M_2 receptor, by generating the M_2-GFP11 construct and by transfecting it in HEK293 cells. Fig. 4i

shows a green signal from the recombined GFP visible in the mitochondria. All together these data suggest

that the M_2 C-terminal fragment is localized in the inner mitochondrial membrane, with its own C-terminal

domain facing the matrix.

This interpretation appears to be confirmed in cells expressing the M_2stop228-myc receptor and examined by

electron microscopy (Supplementary Fig. 10e) after immune-gold labelling. As shown in Supplementary

Fig. 10f-g, gold particles were observed only in mitochondria, and at this stage they appeared localized either

to the mitochondrial matrix or to the lumen of the cristae (Supplementary Fig. 10h-k).

Functional role of the C-terminal fragment in the mitochondria

The next step of our investigation concerned the obvious question about the functional effects, if any, that the

observed localization of the M_2 C-terminal fragment might have on the mitochondrial activity, and -more

generally- on the energy metabolism of the cell. To investigate this aspect, we measured the extent of oxygen

consumption in COS-7 cells expressing M_2-EGFP, M_2tail(368-466)-EGFP or the sole EGFP as a control.

Initially, we normalized the expression of the three proteins to the amount of EGFP expressed in the cells, then

the metabolic functions were assayed on a Seahorse instrument using cells transfected with normalized

amounts of plasmids (Fig. 5a).

Our data did not show a significant difference in total oxygen consumption capacity between cells transfected

with cytosolic EGFP and the untransfected control. However, a significant decrease of oxygen consumption

was found in cells transfected with M_2-EGFP and even more so with M_2tail(368-466)-EGFP (Fig. 5a). Notably,

the basal oxygen consumption rate (OCR), in starvation-stress conditions, was profoundly decreased in

M_2tail(368-466) cells and to a smaller extent in M_2 cells compared to EGFP and empty control cells (see also

the time-course in **Supplementary Fig. 11a)**. These results are in line with what was observed using a Clarke-

type oxygen electrode (**Supplementary results and Supplementary Fig. 11b and 11c**).

A cell apoptosis assay, based on the detection of activated caspase-3[17], did not show substantial differences

among the tested mutants (**Supplementary Fig. 11d**). The overexposed western blot showed a tiny amount of

activated caspase-3 in most of the transfected COS-7 cells, indicating that the expression of M_2tail(368-466)

in mitochondria, in agreement with the qualitative post-starvation data shown in **Supplementary Fig. 11e**,

does not induce fragmentation of the mitochondrial network and cell apoptosis.

To test whether the C-terminal-M_2 fragment effects on mitochondrial oxygen consumption could alter the

production of Reactive Oxygen Species (ROS), the measurements were combined in a single cell assay.

Remarkably, ROS production (assayed using a fluorescence readout kit, where the fluorochrome becomes

activated in contact with oxidized species) was clearly reduced within and in proximity of mitochondria of

cells expressing a high level of the M_2tail(368-466) (**Fig. 5b**). By regrouping individual mitochondria in classes

according to the level of the ROS marker expression, an actual inverse correlation between the amount of

receptor fragment (Green Fluorescence) and the production of ROS (Red Fluorescence) was revealed.

Cell growth was also found to be affected depending on which of the constructs was transfected. Notably,

transfection of HEK293 cells with the wild type M_2 receptor reduced significantly cell growth compared to the

control, whereas cells transfected with the mutant M_2M368A, that do not express the M_2 C-terminal fragment

(**Fig. 5c**), displayed a significantly higher growth. All cells were incubated with saturating concentration of the

M_2 antagonist atropine throughout, in order to exclude effects on cell growth and proliferation due to the

canonical receptor signaling.

Cell stress upregulates M_2 C-terminal expression and mitochondrial localization

Having established that the C-terminal fragment localizes in the mitochondria and affects the cell oxygen

consumption rate, we looked at the effect of stress on the level of expression of the C-terminal fragment.

Stressed cells attenuate their translational activity via phosphorylation of the initiation factor eIF2α. However,

certain mRNAs may be insensitive to eIF2α phosphorylation level if their translation is regulated by IRES.

Under these conditions their expression is bound to increase rather than to attenuate [18,19].

Therefore, we transfected COS-7 cells with the Sirius-M_2i3(417n)-EGFP reporter construct and stressed by

starvation for three hours by replacing the culture medium with HBSS or PBS. During this time interval, the

green fluorescence was seen to gradually increase up to 1.4 times the basal level observed in HBSS-starved

cells and 3.5 times for PBS-starved cells (Fig. 5d), confirming that the C-terminal fragment expression is

upregulated in response to stress conditions such as cell starvation. The same experiment was reproduced in

single cells, displaying an increase of the average EGFP/Sirius ratio upon HBSS starvation in HEK293 cells

(Fig. 5e).

We further used the GFP fluorescence complementation assay to emphasize the importance of stress in the

pattern of expression of the M_2 gene. HEK293 cells were co-transfected with M_2-GFP11 and Mito-GFP1-10.

HEK293 cells grown in culture medium predominantly expressed the full M_2 receptor protein (Fig. 5f). The

green fluorescence detected in the plasma membrane is explained with the chaperoning effect of M_2-GFP11

on Mito-GFP1-10 before its sorting to the mitochondria. In sharp contrast, when HEK293 cells were first

transfected with Mito-GFP1-10 and, 24h later, transfected with M_2-GFP11 followed by a 6 h starvation in

HBSS, the pattern of expression changed drastically: IRES mediated translation of the M_2tail(368-466)

fragment led to a predominantly mitochondrial green fluorescence signal of the complemented GFP (Fig. 5f).

These results are in line with our observation that the IRES mediated C-terminal fragment localization to the

mitochondria in cells expressing the construct *M_2-mRuby2-STOP-M_2-i3-tail-EGFP* was enhanced upon cell

starvation, and drastically reduced upon mutation of the M368 (**Supplementary Figure 12** and

Supplementary Methods).

This observation was also supported by the lack of complementation of M_2tail-GFP11 with GFP1-10 when

this last fragment was N-terminally fused to the targeting sequence of SMAC, a protein expressed in the

mitochondrial intermembrane space [20] (**Supplementary Fig. 13c**).

Mutation of the M368A via CRISPR/Cas9 affects cell growth and oxygen consumption rates in hiPSCs

We finally asked the question of what would be the effect of abolishing the translocation of the M_2 C-terminal

fragment to the mitochondria, in a physiological and entirely label-free setting. To reach this goal we

genetically engineered (see Methods) human induced pluripotent stem cell (hiPSC), to incorporate in both

alleles of the M_2 gene the M368A mutation, that, based on our data, should be sufficient to significantly impair

the M_2 C-terminal production and mitochondrial translocation. Wild typ (wt)e and M368A hiPSCs were

cultured in pluripotency maintaining medium, and subsequently undirectedly differentiated upon exposure for

seven days to DMEM-F12 medium, as described in the Methods. Fig. 6a shows confocal micrographs for wild

type and M368A hiPSCs cultured in E8 medium respectively, with the M368A hiPSCs displaying a mildly

more spread-out phenotype. We shall note here that the M_2 receptor has been found to be expressed in PSCs

lines, and it has been shown to play a role in stem cell differentiation [21,22]. In this context, however, we were

not interested in the canonical signaling of the receptor, and all experiments were performed with a saturating

concentration of atropine, to block any activity of the receptor. We then evaluated cell proliferation, observing

a significant increase in the proliferation of the M368A clone 72h after seeding Fig. 6b. The observation of an

increased proliferation of the cells harboring the mutation that impairs the production and localization of the

C-terminal fragment in the mitochondria, suggests that the same effect is at play here, as observed in the

overexpression system (Fig. 5a,b,c). These results are supported by extracellular oxygen consumption assays

performed, for both hiPSCs and differentiated hiPSCs (Fig. 6c). Our data suggest that the M368A cells indeed

display an accelerated oxygen consumption rate, and that this effect increases as the cell progressively

differentiates.

For comparison, the assay was also carried on HEK293 overexpressing the wild type M_2 receptor and the

mutant M_2M368A (Supplementary Fig. 11f). We could observe with this assay a reduced oxygen

consumption of the wild type M_2 receptor compared to untransfected cells, in line with what was observed in

Fig. 5a, and an increased oxygen consumption of the M_2M368A expressing cells, in line with what observed

for the hiPSCs.

Discussion

Over the last 10 years, growing evidence has accumulated indicating that GPCRs can be functional and signal

also at cellular locations other than the plasma membrane. These locations can be associated to their canonical

trafficking pathways (such as sorting vesicles and endosomes)[23], but also to non-canonical locations, such as

the nucleus, the nuclear membrane, the Golgi apparatus and the mitochondria (reviewed in Jong et al. BPJ

2018 [24]). Some of these locations are surprising, since many ligands are non-membrane permeable, and it is

still debated how they can reach their target receptors at such intracellular locations.

Interestingly, the specific localization of these receptors appears to go hand in hand with a specific function.

In these cases, activation/inhibition of mitochondrially localized GPCRs have been shown to specifically affect

the organelle function, ranging from oxygen consumption rates to Ca^{2+} handling and even to apoptosis. Notably

Bénárd et al [25] showed that the CB_1R on the outer mitochondrial membranes affects neuronal energy

metabolism; Suofu et al. [10] demonstrated that activation of the MT_1R by a mitochondrial melatonin pool can

affect cytochrome C release. This is particularly interesting, since cytosolic cAMP does not appear to be able

to activate mitochondrial PKA, and hence a mitochondrially localized GPCR signalosome appears to be

required to enable such downstream signaling cascade [26].

However, other non-canonical mechanisms are also possible for the regulation of mitochondrial function. It is

well known that GPCRs are subject to interindividual variability, arising, for example, from mRNA alternative

splicing (i.e. the process which allows the recombination of exons) which leads to different isoforms which

may present distinct signaling and localization properties [27]. Receptor variants with different functional and

localization properties may also occur due to other processes. For example, our comprehension of eukaryotic

translational regulation is facing a rapid expansion, as exemplified by the observation that small open reading

frames (smORFs), also within lncRNAs, are an important class of genes, oftentimes with mitochondrial

localization[28].

In this study, for the first time, we provide evidence that an IRES sequence drives expression of a membrane

receptor fragment and acts as a regulator of its expression levels and function. IRES sequences are distinct

regions of the mRNAs that are capable of recruiting ribosomes in the proximity of an internal initiation codon

and translate transcripts in a cap-independent manner [5,6,29-31]. Here, we show that an IRES sequence is located

within the third intracellular loop of the M_2 muscarinic receptor, where it plays a crucial role in regulating the

expression of its carboxy-terminal domain, which we identified as M_2tail(368-466).

We assessed the IRES-driven expression of the C-terminal fragment through a broad set of constructs, meant

to exclude alternative synthesis mechanisms such as stop-codon read-through, and a diversified palette of

techniques. Radioligand binding data (Table 1) pointed to the necessity of the C-terminal fragment synthesis,

in order to explain the observed downstream signaling in mutants carrying a stop codon in the initial portion

of the i3loop of the M_2 receptor, upstream of M368. Fluorescence microscopy allowed us visualizing the

production of the EGFP-tagged C-terminal fragment in these receptor mutants (Fig. 1d). We further detected

its production by Western blotting on protein extracts from cells; here the C-terminal fragment gives rise to

two bands: the lower of the two (~15 kDa) appears to be proportional to the ~75 kDa band, suggesting that this

may result from subsequent processing or degradation. On the other hand the ~20 kDa band intensity appears

independent of the M_2 full-length band (~75 kDa), as would be expected for a cap-independent translation

mechanism, and this band disappears in the M368A mutant.

We further performed Western blotting from heart and brain tissues from knock-in mice carrying a C-

terminally tagged M_2 receptor (tdtomato) and under the control of the endogenous promoter/enhancer

sequences (Fig. 2a-c).

Mutagenesis of key residues within the i3 loop of the M_2 receptor allowed us to pinpoint the 3rd in-frame

methionine as the most likely candidate for the beginning of the cap-independent translation (Fig. 3). This

observation appears to be supported by the observation of a peak in ribosome coverage around the M368 seen

in ribosome profiling databases (as reviewed in Ingolia et al.[32]), and particularly prominent in those from left

ventricle cardiomyocytes, where M_2 receptor expression is known to be abundant[28].

To our surprise, in contrast to the plasma membrane localization of the wild type, full length M_2 receptor, the

C-terminal fragment was localized mostly in mitochondria. This localization occurred markedly when

overexpressing the C-terminal fragment M_2tail(368-466), but could be also clearly detected when the C-

terminal fragment was expressed under the control of the IRES sequence of the full length receptor (Fig. 4

c,d).

We devised several strategies to detect the localization of the IRES-driven fragment in the mitochondria by

avoiding the cell-wide background signal from the full-length receptor. Fig. 4c,d show two prominent

examples, namely relying on spectrally unmixing the two components, or by sending out of frame the EGFP

at the C-terminus of the full-length receptor. By studying mitochondria purified from HEK293 cells we could

confirm that the mutation of the M368, which we identified as the starting point of the IRES-mediated

translation, is sufficient to suppress any mitochondrial targeting of the M_2 C-terminal fragment, and most

likely, its synthesis altogether (Fig. 4e).

We then addressed the mechanism of mitochondrial import of the C-terminal fragment and its localization

within the organelle. Our data, originating both from live cell import kinetics (Fig.4h) as well as from in-vitro

import into isolated mitochondria (Supplementary Fig. 10), support the notion that the fragment becomes

rapidly imported into mitochondria following translation in the cytosol. Once inside the mitochondria, it

undergoes processing leading to the appearance of a lower molecular weight band, observed both in Western

blots from whole cell and mice tissue protein extracts, as well as from purified mitochondria. The fragment is

found predominantly in a carbonate-insoluble fraction, i.e. a membrane, and it appears to form a high molecular

weight complex of ~180 kDa with other mitochondrial proteins (Supplementary Fig. 10c).

Fluorescence complementation assays (Fig. 4i) and electron microscopy (Supplementary Fig. 10d) suggest

that the M_2 C-terminal fragment predominantly resides in the inner mitochondrial membrane, with its own C-

terminal domain exposed towards the matrix. Future work should clarify the precise import pathway.

Most importantly, our data show that mitochondrial oxygen consumption was reduced by the M_2tail(368-466)

(Fig. 5a,b), and also cell proliferation was affected (Fig. 5c). Under integrated stress response, namely

aminoacid starvation, IRES-mediated production of the C-terminal fragment was increased (Fig. 5d-f). On

these bases, the M_2tail might be needed in mitochondria as a means to reduce mitochondrial basal oxygen

consumption and ATP-linked respiration. An important consequence of reduced respiration is the lower level

of ROS produced in mitochondria, as we observed in agreement with this hypothesis.

This observation was confirmed in physiological, label-free conditions, by genetically editing human induced

pluripotent stem cells in order to suppress the synthesis of the M_2 C-terminal fragment mutating the starting

methionine M368. Increase of cell proliferation of this clone compared to its wild type counterpart, together

with an increase of oxygen consumption rates (Fig. 6), corroborates our hypothesis that the M_2 C-terminal

fragment reduces oxygen consumption upon reaching the mitochondria. The effect was further amplified when

the cells were undirectedly differentiated, and it was consistent with the shift of the cell phenotype towards

oxidative phosphorylation[33].

The mechanism by which the M_2tail(368-466) may reduce oxygen consumption in mitochondria and its

meaning needs to be further explored. Nonetheless, since ROS production is implicated in the pathogenesis of

several diseases[34], we would propose the hypothesis that increased localization of the M_2 C-terminal fragment

in mitochondria, as it occurs under integrated stress response, might have protective effects (Fig. 7).

In broader terms, protection of mitochondria by unconventional import mechanisms to allow localization

within the organelle of specific proteins has been previously observed for cells that undergo stress conditions

[35]. The mechanism of IRES-dependent modulation of translation and consequent mitochondrial targeting of

the translation product of a GPCR may provide another means, completely unrelated to the canonical signaling

function of this class of membrane receptors, of a generic cellular response to overcome different types of

stress by modulating the mitochondrial proteome and adjusting the mitochondrial bioenergetic state.

Tables, Figures and Supplementary Tables and Figures

Table 1. Radioligand binding and activation properties of M_2 receptor mutants. Binding properties and inhibition of

forskolin stimulated adenylyl cyclase by M_2, M_2stop228 and the co-transfected M_2trunk(1-283)/M_2tail(M-281-466) and

M_2trunk(1-228)/M_2tail(368-466). Untransfected COS-7 cells did not show any [^{3}H]NMS specific binding. N.B. = no

specific [^{3}H]NMS binding.

	Binding data			Adenylyl cyclase inhibition assay	
Receptor	Bmax (fmol/mg)	[^{3}H]NMS KD (pM)	Carbachol IC50 (µM)	Carbachol EC50 (µM)	Maximum inhibition of forskolin stimulated cAMP accumulation (%)
M_2	980 ± 60	62 ± 7	2.1 ± 0.6	0.04 ± 0.01	53 ± 5
M_2trunk(1-283)	N.B.				
M_2trunk(1-228)	N.B.				
M_2stop228	45 ± 2	70 ± 10	1.8 ± 0.4	0.13 ± 0.02	27 ± 3
M_2trunk(1-283) +M_2tail(M-281-466)	480 ± 20	79 ± 10	0.7 ± 0.2	0.15 ± 0.03	37 ± 4
M_2trunk(1-228) +M_2tail(368-466)	70 ± 10	72 ± 8	0.6 ± 0.2	0.17 ± 0.03	29 ± 4

M₂wt-EGFP   M₂stop228-EGFP   M₂stop400-EGFP   M₂fr.sh.-EGFP   non transfected

Fig. 1. **Mutations of the M₂ receptor confer unexpected functional and localization properties.** **a**, Snake-like plot of the human muscarinic M₂ receptor: in red are indicated the aminoacids substituted with stop codons in the mutants: M₂stop196, M₂stop228 and M₂stop400 (**Supplementary Fig. 1**). The three in-frame methionines of the i3 loop, 248, 296 and 368, are highlighted in yellow; the translation of the C-terminal fragment starts from methionine 368. **b**, Schematic representation of the gene products that reach the plasma membrane of the wild type M₂, M₂trunk, M₂trunk and M₂tail and M₂stop228 (**Supplementary Fig. 1**). The M₂trunk and M₂tail genes generate the respective protein fragments that, only when co-transfected together, are able to form functional receptor complexes capable to bind ligands. The M₂stop228 gene mutant generates both M₂trunk (in green), when translation starts from the cap site, and M₂tail (in brown) when translation is cap-independent driven. This makes it likely that the truncated fragment of the M₂stop228 receptor may act as a chaperon driving part of the C-terminal-M₂ fragment to the plasma membrane, as supported by the binding and functional data shown by the co-transfection of M₂trunk(1-228) with M₂tail(368-466) (**Table 1**). **c**, Fluorescence micrographs of HEK293 cells respectively expressing M₂-EGFP, M₂stop228-EGFP, M₂fr.sh.-EGFP and M₂stop400-EGFP, together with an untransfected control. Imaging settings are the same for the five images, whereas contrast was adjusted to improve visualization. In the untransfected cells and stop400 panels, the contrast was set to 0. Insets display DIC image of the same field of view (scale bars 10 μm).

Schematic representation of wild type muscarinic M₂ (human) receptors and derived mutants. Each

construct was obtained as described under Methods. The left column indicates the mRNA product from each construct,

the right column the expected protein product. For the last two constructs, above mRNA product, and below the expected

protein product.

M₂ receptor mutants with single and double stop codons. a, Schematic representation of the gene

products M₂stop196 and M₂stop400, that are not functional by themselves but that are rescued by co-transfection with

M₂Trunk(1-283) and M₂Tail(M-281-466), respectively. **b**, Dose response curve of carbachol induced ERK

phosphorylation in HeLa cells transiently transfected with M₂ and M₂stop228 muscarinic receptors. Cells were stimulated

for 5 minutes with the indicated concentration of carbachol. Phosphorylated ERK (P-ERK) was normalized vs total ERK

(Tot ERK). Significance (*p < 0.05, **p < 0.01) was calculated against carbachol 0 μM.

Supplementary Table 1. Ligand binding and functional properties of M₂ receptor mutants with single and double

stop codons. Number of [³H]NMS binding sites for the indicated mutants. Untransfected COS-7 cells did not show any

[³H]NMS specific binding. N.B. = no specific [³H]NMS binding.

Receptor	Bmax (fmol/mg)	Receptor	Bmax (fmol/mg)
M₂stop228	45.3 ± 2.4	M₂stop228/stop368	N.B.
M₂stop196	N.B.	M₂stop228/stop368 + M₂tail(281-466)	370± 20
M₂stop400	N.B.	M₂stop228/stop400	N.B.
M₂stop196+ M₂trunk(1-283)	47 ± 3	M₂stop228/stop400+ M₂tail(281-466)	380 ± 20

M₂stop400+ M₂tail(281-466)	400 ± 30	M₂stop196/stop400+ M₂trunk(1-283)	N.B.
M₂stop196/stop400	N.B.	M₂stop196/stop400+ M₂tail(281-466)	N.B.
M₂stop228/stop248	49 ± 5	M₂stop228/fr.sh.	47 ± 3
M₂stop228/stop296	38 ± 3	M₂stop228/Hairpin	37 ± 2

Supp. Fig. 3. **Lack of alternative splicing in M₂ wild type and M₂stop228 receptors**. mRNAs were extracted by COS-

7 cells transfected with M₂ wild type and M₂stop228, subjected to reverse transcriptase and the resulting cDNA amplified

by PCR with two oligos directed to the 5' and 3' end of the receptors. The gel shows, in both M₂ wild type and M₂stpo228,

a single band of 1401 bp corresponding to the full-size receptor, running at the same level of the band amplified directly

form the M₂ pcD plasmid.

Fig. 2. **Western blot of tagged M₂ receptor mutants in HEK293 cells and in tdtomato knock-in mice. a, left** Graphical

guide of the fragments observed on the gel when using myc-tag. **right**, M₂-myc, M₂tail(368-466)-myc, M₂M368A-myc

and M₂stop400-myc were transiently transfected in HEK293 cells and immunodetected via antimyc primary antibody,

together with an untransfected control. Loading control normalization was performed vs β-actin **b**, **left** Graphical guide

of the fragments observed on the gels when using the tdtomato tag. **right**, Western blot from tissue preparations

originating from the indicated organs of M₂-tdtomato knock-in mice (HT) and wild-type animals (WT). Immunodetection

was performed using an anti-tdtomato primary antibody. Loading control normalization was performed using GAPDH in

the heart, and β-actin in the brain and spleen.

Supp. Fig. 4. **Expression of the bicistronic plasmid Sirius-M₂i3(417n)-EGFP in four different cell lines**

Expression of Sirius was driven by the canonical scanning mechanism of translation initiation, while EGFP expression

was driven by an IRES dependent mechanism. All Sirius-M₂i3(417n)-EGFP transfected cells expressed both the Sirius

and EGFP proteins. On average, $25 \pm 4\%$ of the COS-7 cells transfected with Sirius-M$_2$i3(417n)-EGFP plasmid were blue

fluorescence positive, while $17 \pm 2\%$ were green fluorescent positive, with only a few cells being only green.

<mark>Fig. 3.</mark> **Fluorescence assay of cap-independent translation of EGFP driven by the i3 loop mutants of the muscarinic**

M$_2$ receptor. a, **left** Schematic representation of bicistronic plasmids bearing the M$_2$ i3 loop between the coding regions

of the Sirius and EGFP fluorescent proteins. A guanine nucleotide was originally inserted before the EGFP in the plasmid

Sirius-H-EGFP in order to maintain in frame the sequences of the two fluorescent proteins. In all the other plasmids the

sequence of the two fluorescent proteins is not in frame. Fifteen nucleotides derived from the M$_2$ i3 loop were inserted

between the end of Sirius and the hairpin loop, in order to allow the easy scanning of the ribosome across the stop codon.

right, Relative fluorescence of EGFP compared to Sirius, highlighting the efficiency of cap-independent translation for

the control Sirius-M$_3$i3(417n)-EGFP and the six mutants. The insert shows the difference in the green fluorescence

between two of the mutants. **b**, **left** Mutations of the bicistronic Sirius-H-M$_2$i3(15n/30n)-EGFP plasmids in the region of

the M$_2$ i3 loop from nucleotides 1972 to 2001, in yellow are the modified nucleotides. In Sirius-H-M$_2$i3(15n/30n)mut#6-

EGFP the four nucleotides in green (AAGA) were reinserted in the stretch of 20 T of Sirius-H-M$_2$i3(15n/30n)mut#5-

EGFP. **right** Relative fluorescence of EGFP compared to Sirius, highlighting the efficiency of cap-independent translation

for the control Sirius-H-M$_2$i3(15n/30n)-EGFP and the six mutants illustrated. Significance ($p < 0.01$) was calculated in **a**

against Sirius-M$_2$i3(417n)-EGFP and in **b** against Sirius-H- M$_2$i3(15n/30n)-EGFP. Error bars are s.e.m.

Supp. Fig. 5. Expression of the construct M₂-Hairpin-EGFP in HEK293 cells. (left) Confocal images of M₂-Hairpin-EGFP, (middle) Mitotracker Deep red staining, (right) DIC. Scale bar 5 μm.

Supp. Fig. 6. Alignment of the nucleotide sequence 1072-1104 of the M₂ i3 loop with analogous sequences of the other four muscarinic receptors, and expression of the bicistronic plasmid Sirius-M₃i3(558n)-EGFP. a, The criteria for the alignment are described in the supplementary data. In yellow are highlighted the conserved nucleotides. In bold are the in frame ATG codons. Next to each sequence is indicated the codon number of the in-frame ATG and its distance in nucleotides from the beginning of TMDVI. **b,** Schematic representation of the bicistronic plasmid bearing the i3 loop of the rat muscarinic M₃ receptor (558 nucleotides from nucleotide 880 to nucleotide 1437) between the coding regions of the Sirius and EGFP fluorescent proteins. **c,** EGFP expression in COS-7 cells transfected with the bicistronic plasmid Sirius-M₃i3(558n)-EGFP.

662 <mark>**Supplementary Table 2**</mark>. **Radioligand binding and activation properties of M$_3$ receptor mutants.** Binding

663 characteristics and stimulation of phosphatidylinositol accumulation by M$_3$, M$_3$stop273 and the co-transfected M$_3$trunk(1-

664 272) and M$_3$tail(M-388-586). Untransfected COS-7 cells did not show any [^{3}H]NMS specific binding. N.B. = no specific

665 [^{3}H]NMS binding.

	Binding data			Phosphatidylinositol accumulation assay	
Receptor	Bmax (fmol/mg)	[^{3}H]NMS KD (pM)	Carbachol IC50 (μM)	Carbachol EC50 (μM)	Maximum increase in IP$_1$ level above baseline (%)
M$_3$	1120 ± 71	29 ± 2	59 ± 4	4.2 ± 0.5	183 ± 15
M$_3$trunk(1-272)+ M$_3$tail(388-589)	114 ± 17	23 ± 2	48 ± 5	1.4 ± 0.1	126 ± 12
M$_3$stop 273	52.3 ± 0.5	31.3 ± 2.8	56 ± 2	1.1 ± 0.1	89 ± 13

667 <mark>**Supp. Table 3**</mark>. **Ligand binding of M$_3$ receptor mutants with single and double stop codons.** Number of [^{3}H]NMS

668 binding sites in COS-7 cells transiently transfected with the indicated M$_3$ receptor mutants with single and double stop

669 codons.

Receptor	Bmax (fmol/mg)	Receptor	Bmax (fmol/mg)
M$_3$stop273	54 ± 1	M$_3$stop273/stop503	N.B.
M$_3$stop240	N.B.	M$_3$stop273/stop503 + M$_3$tail(388-589)	120± 15
M$_3$stop503	N.B.	M$_3$stop240/stop503+ M$_3$trunk(1-272)	N.B.
M$_3$stop240+ M$_3$trunk(1-272)	44 ± 3	M$_3$stop240/stop503+ M$_3$tail(388-589)	N.B.
M$_3$stop503+ M$_3$tail(388-589)	135 ± 12	M$_3$stop273/fr.sh.	120 ± 4
M$_3$stop240/stop503	N.B.	M$_3$stop273/Hairpin	30 ± 2

Fig. 4. Localization of fluorescently labeled M₂ mutant constructs: microscopy and western blots analysis. a, Colocalization of M₂tail(368-466)-EGFP (green) together with the mitochondrial targeted dye Mitotracker (magenta) in HEK293 cells. Separate spectral channels are displayed below. **b**, Co-expression of M₂tail(368-466)-EGFP (green) and wild type M₂-mRuby2 (red) in HEK293 cells. Separate spectral channels are displayed below. **c**, Cellular localization of the construct M₂-mRuby2-STOP-M₂-i3-tail-EGFP in HEK293 cells. Expression of the M₂tail(368-466)-EGFP (green) after the STOP codon is an indicator of the endogenous IRES activity. While M₂-mRuby2 localizes correctly and predominantly to the cell plasma membrane, M₂tail(368-466)-EGFP localizes in the cytosol as well as, strongly, to the cell mitochondria. Separate spectral channels are displayed below. **d**, M₂-fr.sh.-EGFP (green) expressed in conjunction with Mitotracker (magenta). The EGFP is expressed despite the creation of an in-frame stop after M368, confirming endogenous cap-independent translation of the M₂tail(368-466). Arrows indicate areas of colocalization with the mitochondria. Separate spectral channels are displayed below. **e**, Western blot (10% TRIS-Glycine PAA gel) of mitochondria purified from Hek293 cells transfected with the M₂-Myc, M₂tail(368-466)-Myc, M₂M368A-Myc, together with an untransfected control are blotted using an anti-myc antibody. Loading control was verified by immunoblotting COXIV. **f**, Western blot (8% TRIS-Glycine PAA gel) of mitochondria purified from tissue preparations originating from the indicated organs of M₂-tdtomato knock-in mice (HT) and wild-type animals (WT), detected using an anti-tdtomato antibody. Please note that the band at 80kDa is aspecific, and is more prominent in the WT lane due to higher loading. **g**, Western blot (8% TRIS-Glycine PAA gel) of mitochondria purified from HEK293 cells transfected with the M₂tail(368-466)-tdtomato, together with an untransfected control. Mitochondrial proteins were blotted using an anti-tdtomato antibody. Loading control was verified by detecting the 15 kDa mitochondrial matrix protein Cox-IV. **h** Time course of M₂tail(368-466)-EGFP import in the mitochondria of a HEK293 cell over a 4h interval, starting 5h after cell transfection, as indicated. Panels, from top to bottom show M₂tail(368-466)-EGFP (green), Mitotracker (magenta) and overlay displaying increasing colocalization over time. Chart at the bottom quantifies colocalization of the M₂tail(368-466)-EGFP

in the mitochondria by calculating the Manders Coefficients (n=7 cells, from 3 independent transfections, error bars are

s.e.m.) (See <mark>Methods</mark>). **i,** Detail of a HEK293 cell expressing the M_2-GFP11 construct together with pmito-GFP1-10. The

M_2-GFP11 was sequentially transfected 24 h after the pmito-GFP1-10 (see <mark>Methods</mark>) Magenta overlay indicates

mitotracker. White signal in the overlay indicates M2tail(368-466) trafficking to the organelle. All confocal micrographs

were obtained by sequential imaging using 488 nm excitation and detection in the 520-600 nm (EGFP), 561 nm excitation

and detection in the 580-650 nm (mRuby2) and at 633 nm and detected in the 650-700 nm range (Mitotracker) using HyD

detectors in Photon Counting Mode. Scale bars are 10 μm throughout.

<mark>Supp. Fig. 7.</mark> **Subcellular localization of the M2tail(368-466). a,** Confocal image displaying the colocalization (top) of

M2tail(368-466)-EGFP (green, middle)) in HEK293 cells, co-expressed with the outer mitochondrial membrane marker

Tom20-mCherry (red, bottom). **b** expression of M2tail-EGFP displaying marked mitochondria and cytosolic localization.

c, Glasbey colorscale representation of panel (**b**) highlighting the low-intensity pixels. **d**, autocorrelation function of the

M2tail(368-466)-EGFP diffusion taken at or in proximity of the basal membrane, yielding a mean diffusion coefficient of

0.11 μm²/s, incompatible with membrane diffusion. **e**, co-expression of M2stop228 and M2tail(368-466)-EGFP.

M2stop228 is unlabeled. M2tail(368-466)-EGFP maintains the localization to the mitochondria observed in cells

expressing M2tail(368-466)-EGFP alone, but it is also localized to the plasma membrane, visible in panel **f**, Glasbey

colorscale representation of panel **e** highlighting the low-intensity pixels. **g**, autocorrelation function of the M2tail(368-

466)-EGFP diffusion taken at or in proximity of the basal membrane, yielding a mean diffusion coefficient of 0.013

μm²/s, compatible with membrane diffusion. **h**, Cellular localization of M2tail-mRuby2, together with Mitotracker deep

red staining of the mitochondrial network and corresponding DIC image. **i,** Cellular localization of M2tail-tdtomato and

corresponding DIC image. All panels originate from Laser Scanning Confocal Microscope sequential acquisitions, with

laser lines 488 nm (EGFP) 561 nm (mRuby2 and tdtomato) and 633 nm (Mitotracker deep red), and corresponding

emission filters in the ranges 520-600 nm, 570-620 nm and 640-750 nm. Scale bars 10 μm.

Supp. Fig. 8. Mitochondrial localization of the M2tail fragment. **a**, Confocal image of M$_2$-EGFP (green) expressed in

conjunction with Mitotracker (magenta) in HEK293 cells. Separate panels below. **b,** Confocal image of HEK293 cell

expressing *M$_2$fr.sh-mRuby2*, together with DIC, below. **c**, Confocal sequential images displaying the localization of

fluorescently labeled M$_2$tail(368-466)-EGFP (green), together with the mitochondrial network (magenta), colocalization

(white) and DIC image in COS-7 cells. Scale bars are 10 μm. Confocal sequential acquisitions were performed with 488

nm excitation and 520-600 nm detection (EGFP), and with 561nm excitation and 600-750 nm detection (mCherry) and

633 nm excitation and 650-750 nm detection (Mitotracker) using HyD detectors in Photon Counting Mode. **d,** Western

blot (10% TRIS-Glycine PAA gel) of the cytosolic and microsomal fractions resulting from the mitochondria purification

of lysates of HEK293 transfected with myc-tagged constructs, as displayed in **Fig. 4e**. **e**, Western blot (8% TRIS-Glycine

PAA gel) of the cytosolic fraction resulting from the mitochondria purification of lysates of HT and WT tissues of a

tdtomato transgene mice, as displayed in **Fig. 4f. f**, Western blot (10% TRIS-Glycine PAA gel) of the cytosolic fraction

resulting from the mitochondria purification of lysates of HEK293 transfected with M$_2$tail(368-466)-tdtomato, as

displayed in **Fig. 4g.**

Supp. Fig. 9 Relative abundance estimates of EGFP and M₂ receptor from PRM based mass spectrometric

measurements. Normalized fragment intensities were calculated for the peptide FSVSGEGEGDATYGK (specific for

EGFP) for cytosolic and pellet fractions and for different constructs. Each preparation was normalized relative to the

cytosolic fraction, if data was available. Median levels of cytosolic, plasma membrane and mitochondrial control markers

are shown in the left panel. For each identified peptide the normalized spectrum contrast angle, indicating the similarity

of the recorded spectrum to the reference spectrum, is indicated below each plot.

Supp. Fig. 10. **In vitro mitochondrial import of M₂tail(368-466) and electron microscopy of M₂stop228. a,** Import

of M2-tail fragment in yeast mitochondria: 35S-labelled and translated in-vitro full length muscarinic M₂ receptor or

M₂tail(368-466) expressing 5 C-terminal Methionines were incubated with isolated yeast mitochondria for various

timepoints in the presence of the inner membrane potential (See Supplementary Methods). Samples were then washed

in breaking buffer and loaded onto a 12% Tris-Tricine gel followed by semi-dry transfer and visualized via a

phosphorimager. 5% of the crude protein was loaded as a control (left lane). To test for equal mitochondrial loading,

αPorin was probed via western blot. The kinetics pattern of M₂tail is what it is expected for a protein imported in

mitochondria: the appearance of the cleaved M₂-tail is indicative of a processing of the fragment once imported in

mitochondria. **b,** 35S-labelled M₂tail(368-466) or AAC (an integral internal mitochondrial membrane protein) were

imported into isolated yeast mitochondria originating from different knock-out lines for outer membrane transporters

(ΔTom). Samples were incubated for 30 minutes, and then washed in breaking buffer. After import, 100 µg of

mitochondria were subject to Carbonate extraction to determine if M₂tail(368-466) is integrated into a lipid bilayer (Pellet)

or loosely associated/soluble (Supernatant). Samples were then loaded onto a 12% Tris-Tricine gel followed by semi-dry

transfer and visualized via a phosphorimager. To test for equal mitochondrial loading anti-αPorin was used via western

blot. **c,** 35S-labelled M₂tail(368-466) and αPorin were imported into isolated wild-type or ΔTom70 mitochondria for the

indicated timepoints. Samples were then washed in breaking buffer and loaded on blue-native page followed by gel drying

and visualized via a phosphorimager. **d,** Confocal images displaying the mitochondrial localization of M₂tail-GFP11 +

M₂tail-GFP1-10, with corresponding DIC image. **e,** TEM micrograph of a COS-7 cell showing a large nucleus, several

mitochondria and vacuoles nearby. Bar: 5 µm. **f-g,** High magnification of portion of the cell cytoplasm showing a few

gold particles confined to the mitochondria. Bar: 500 nm. **h-k,** Gold particles appear associated with the mitochondrial

cristae, highlighted by the dotted line. A total number of 236 gold particles in 40 mitochondria from 19 sections were

detected.

Fig. 5. **Effects of M₂ mutants on mitochondrial function and cell metabolism in transfected HEK293 cells. a**, Basal oxygen consumption upon HEK293 cell stress -following starvation- measured by a Seahorse assay at a constant temperature of 37 °C. **b**, Localization of M₂tail(368-466)-EGFP in HEK293 (green) cells stained with a marker for reactive oxygen species (Cell Rox Deep Red, ThermoFisher). Relationship between green (M₂tail abundance) vs red (amount of ROS produced) fluorescence intensities measured from regions of the mitochondrial network. The data were binned in four groups as a function of the amplitude of Cell Rox Deep Red signal (0-2.5, 2.5-5. 5-7.5 and 7.5-10 intensity counts). **c**, Cell viability assay performed on HEK293 cells transfected with M₂, M₂tail(368-466) and M₂(M368A), compared to an untransfected control. Cells were counted 24h and 72h after transfection in T-25 flasks, exposed throughout to 100 nM atropine. The experiments were performed in two biological replicates, each with two technical replicates. Transfection efficiency (to the right of the chart) was monitored thanks to the mRuby2 fluorescent reporter within a bicistronic cassette within each vector (see Methods). **d**, 96-well plate reader measurement of the increase of EGFP fluorescence relative to Sirius in COS-7 cells transfected with the bicistronic Sirius-M₂i3(417n)-EGFP plasmid and starved for three hours respectively in PBS (left panel) (n= 3 transfections, 16 wells) and HBSS (right panel) (n=1 transfection, 16 wells). **e,** Representative single cell confocal and DIC micrographs of HEK293 cells expressing the Sirius-M₂i3(417n)-EGFP construct in basal conditions, together with scatter plot of the average EGFP and Sirius fluorescence intensity values measured from single cells in basal (red dots) and 3h HBSS starvation (blue dots) conditions. **f**, HEK293 cells were sequentially transfected with pMito-GFP1-10 (24h) and M₂-GFP11 (8h, of which 6h under starvation conditions). (left panels) HEK293 cells grown in full culture medium and (right panels) HEK293 cells starved for 6 h after transfection of M₂-GFP11.Shown are EGFP fluorescence (top, with DIC inset) and Mitotracker (bottom).

Supp. Fig. 11. **Supplemental data on the effect of the M₂tail(368-466) fragment on mitochondrial function and cell metabolism. a,** Representative experiment of Seahorse assay showing the time course of oxygen consumption inhibition by M₂tail(368-466)-EGFP and wild type M₂-EGFP, compared to the same experiments performed in control cells transfected with EGFP. The rate of ATP-linked respiration measured after oligomycin administration is decreased in M₂tail and slightly decreased in M₂ cells as well as compared to the controls. FCCP, an ionophore, was added to each cell type to investigate the cellular bioenergetic reserves that seem to be slightly reduced only in the M₂ tail cells. **b**, Oxygen consumption measured by a Clark type electrode-based-polarographic method at a constant temperature of 37 °C. After

15 minutes of basal respiration, the ATP synthase inhibitor oligomycin was added at a concentration of 17 nM. Oxygen

consumption in the absence and presence of oligomycin is represented by black and red bars, respectively. The blue bars

represent the percentage of mitochondrial oxygen consumption (total minus oligomycin treated) in cells transfected with

each construct. **c**, is a representative experiment showing the time course of inhibition of oxygen consumption by

M_2tail(368-466), compared to the same experiments performed in control cells and cells transfected with the M_2 wild type

construct, upon addition of oligomycin. Results are given as ng-atom of oxygen/min/10^6 cells. **d**, Western blot of caspase-

3 cleavage in COS-7 cells transfected with the indicated M_2 receptor mutants 48h after transfection (top panel, normal

contrast; bottom panel, increased contrast). **e,** Confocal sequential images of M_2tail(368-466)-EGFP (green), Mitotracker

(magenta) and overlay upon 2h of serum starvation. Scale Bars are 10 μm. **f**, Extracellular oxygen consumption assay of

HEK293 cells transfected with the indicated M_2 receptor mutants using a fluorogenic dye quenched by Oxygen. Blank

indicates the dye lifetime measured in wells containing the dye, but no cells.

Supp. Fig. 12. **Functional aspects of mitochondrial localization of the M_2tail(368-466) fragment. a-e**, Mitochondrial

localization of the IRES mediated C-terminal fragment is enhanced upon cellular stress. HEK293 cells transfected with

the super construct M_2-mRuby2-STOP-M_2-i3-tail-EGFP display increased localization of the EGFP labelled portion to

the mitochondria, upon serum starvation for 2 hours and incubation of the cells in HBSS buffer, in presence of

Mitotracker: **a**, is IRES-driven M_2tail(368-466)-EGFP. **b,** Cap dependent M_2-mRuby2 + IRES-driven M_2tail(368-466)-

mRuby2, both derived from the M_2-mRuby2 gene. **c**, Mitotracker Deep Red **d**, Overlay of Mitotracker Deep Red and

EGFP channel. **e**, Overlay between GFP and mRuby2. **f-l**, Localization of the mutant construct M_2(M368A)-mRuby2-

STOP-M_2-i3-tail-EGFP in HEK293 cells after 2 hours incubation in HBSS buffer. The same legend as in **a-e** applies.

Bottom: Schematic structure of the mega constructs M_2(M368A)-mRuby2-STOP-M_2-i3-tail-EGFP. This construct is

similar to the M_2-mRuby2-STOP-M_2-i3-tail-EGFP construct but the methionine 368 in the M_2 sequence has been replaced

with alanine. The asterisks represent the stop codons at the end of mRuby2 and EGFP. EGFP was excited at 488 nm and

fluorescence collected between 500-600 nm; mRuby2 was excited at 561 nm and fluorescence collected between 580-

620 nm. Mitotracker deep red was excited at 633 nm and fluorescence collected between 650-750 nm. Scale bars are 10

µm.

Supp. Fig. 13. **Cellular localization of split GFP constructs in HEK293 cells.** Confocal micrographs of **a,** co-

expression of SMAC-GFP11 and SMAC-GFP1-10 (positive control) and corresponding Mitotracker deep red image. **b,**

mito-GFP11 and mito-GFP1-10 (positive control). **c,** M$_2$tail(368-466)-GFP11 and SMAC-GFP1-10. **d,** M$_2$tail(368-466)-

GFP11 and mito-GFP1-10. **e,** mito-GFP1-10 alone (negative control). EGFP (green) was excited at 488 nm and

fluorescence collected between 500-600 nm. Mitotracker deep red (magenta) was excited at 633 nm and fluorescence

collected between 650-750 nm. Scale bars are 10 µm.

Fig. 6. **Effects of M368A mutation on cell metabolism in hiPSCs a,** Confocal micrographs of wt hiPSCs (left) and

those harboring the M368A mutation within the M$_2$ gene (right), stained with DAPI (cyan) and Mitotracker (red). **b,** Cell

proliferation assay performed for wt hiPSCs and M368A hiPSCs grown in T25 flasks. Cells were counted 24h and 72h

after seeding. **c,** Extracellular oxygen consumption assay, performed on wt and M368A hiPSCs, prior to and after 7 days

undirected differentiation. The lifetime of a commercial dye increases as oxygen in the medium is progressively consumed
by the cells. The data displayed are collected after 60 minutes incubation.

Fig. 7. **Schematic representation of the wild type M₂ receptor gene products, their destination in the cell and their**
function. The M₂ receptor gene give rise to two different transcripts: when translation starts from the canonical cap site
the wild type M₂ receptor is generated (in blue). When translation is driven by the internal ribosome entry site (IRES) the
C-terminal fragment M₂tail is generated (in green). The two proteins are then sorted in two different cell compartments
where they play completely different roles: the M₂ receptor is transported to the plasma membrane where it functions as
a G protein-coupled neurotransmitter receptor and inhibits the adenylyl cyclase and β-arrestin mediated signalling. The
M₂tail is instead transported in the mitochondria where it inhibits mitochondrial respiration and reduces ROS formation.
IRES driven M₂tail translation is probably stimulated in para-physiological and pathological conditions and might have a
protective role in cells expressing muscarinic M₂ receptors.
840

Methods

Statistical analysis. Significance values reported in the graphs, in comparing tabulated values, were determined by a one-tailed Student t-test using Prism 9 (GraphPad Software), and significance values (p-values) are indicated according to the following legend: **** $p<0.0001$; *** $0.0001<p<0.001$; ** $0.001<p<0.01$; * $0.01<p<0.05$. p-values larger than 0.05 are deemed non-significant. Unless otherwise indicated, bar charts and scatter plots indicate average value with s.e.m.

Generation of mutant muscarinic M_2 receptors. Multiple receptor mutants were constructed starting from the human M_2 muscarinic receptor expressed in the pcD vector [36]. The constructs are sketched in **Supplementary Fig. 1**. The two expression plasmids referred as M_2trunk(1-283) (containing transmembrane domains I-V and the N-terminal portion of the third cytoplasmic loop) and M_2tail(281-466) (containing transmembrane domains VI and VII, and the C-terminal portion of the third cytoplasmic loop) were described previously [12].

M_2stop228 – This plasmid was created by substituting codon 228 (CAA) with a stop codon (TAA) at the N-terminal of the i3 loop of the wild type M_2 receptor.

M_2stop228/stop400 – This plasmid was created by inserting an additional stop codon – TGG was substituted with TAA – at the position codon 400 of the TM region VI of the M_2stop228 mutant.

M_2stop400 – This plasmid was created by substituting codon 400 (TGG) with a stop codon (TAA) in the TM region VI of the wild type M_2 receptor.

M_2stop196 – This plasmid was created by substituting codon 196 (TAT) with a stop codon (TAA) in the TM region V of the wild type M_2 receptor.

M_2stop196/stop400 – This plasmid was created by inserting an additional stop codon in the TM region VI of the M_2stop196 mutant, where TGG was substituted with TAA at position of the codon 400.

M_2stop228/fr.sh. – This plasmid was created by inserting four bases (AATT) fifteen nucleotides downstream of the stop codon to create a shift in the correct reading frame of the M_2stop228 mutant.

$M_2stop228/hairpin$ – This construct was created by inserting a 42 bases long sequence

(AGGGGCGCGTGGTGGCGGCTGCAGCCGCCACCACGCGCCCC).   fifteen

nucleotides downstream of the stop codon of the M$_2$stop228 mutant. Upon transcription,

this palindromic sequence has been shown to form a hairpin structure with a ΔG value

of −64 kcal/mol that blocks effectively ribosome scanning [37,38].

$M_2stop228/stop248, M_2stop228/stop296$ and $M_2stop228/stop368$ – These three constructs were obtained

by substituting the three in-frame ATG codons (1. ATG 248, 2. ATG 296 and 3. ATG

368) with the stop codon TAA downstream of the stop228 of the M$_2$stop228 mutant.

$M_2trunk(1-228)$ – This plasmid was created by substituting codon 228 (CAA) with a stop codon (TAA)

at the N-terminal of the i3 loop of the wild type M$_2$ receptor followed by removal of the

downstream sequence.

$M_2tail(368-466)$ – This construct was created by removing all bases of the M$_2$ receptor sequence up to

codon 368 (ATG) of the C-terminal of the i3 loop. The resulting plasmid encodes for a

polypeptide fragment that contains the trans-membrane domains VI and VII along with

the C-terminal portion of the third cytoplasmic loop.

$M2(M368A)$ – In this construct the third in-frame methionine of the i3-loop of the M$_2$ muscarinic

receptor (ATG) was mutated to alanine (GCG).

M_2-Myc, $M_2stop228$-Myc, $M_2stopM368A$-Myc, $M_2tail(368-466)$-Myc and $M_2stop40$-Myc – M$_2$-Myc was

purchased from OriGene. This plasmid encodes the human M$_2$ receptor with a Myc (

tag at the C-terminus of the protein. M$_2$stop228-Myc was obtained by replacing the 0.9

Kb BmtI-PspOMI fragment of the M$_2$stop228 mutant with the corresponding fragment

of M$_2$-Myc. M$_2$stop400-Myc was obtained by substituting the codon 400 (TGG) with a

stop codon (TAA) of the M$_2$-Myc. All the constructs have an additional DDK tag after

the C-Myc. Subsequently, all C-Myc constructs were extracted by PCR and subcloned

into a bicistrionic pVitro2-MCS plasmid (Invivogen), allowing for the expression of a

reporter gene (in our case the red fluorescent protein mRuby2) after an IRES sequence,

in order to check for transfection efficiency.

Generation of bicistronic constructs expressing Sirius and EGFP proteins (Fig. 3a).

Sirius[PacI-M₂i3(685-1101)]EGFP – This bicistronic construct was created by inserting the i3 loop of

the muscarinic M_2 receptor, from nucleotide 685 to nucleotide 1101, between the

ultramarine fluorescent protein (Sirius) and the green fluorescent protein (EGFP). Eight

nucleotides corresponding to the recognition site of the PacI enzyme were inserted

between Sirius and the i3 loop of the wild type M_2 receptor to alter the reading frame

downstream of the Sirius stop codon. Hereafter this plasmid is referred to as Sirius-

M_2i3(417n)-EGFP.

Sirius[PacI-M₂i3(685-699)-Hairpin-G]EGFP – This bicistronic construct was created by inserting a 42

nucleotides hairpin loop (see above) between Sirius and the EGFP fluorescent protein.

The hairpin loop was spaced from the Sirius stop codon by inserting a PacI recognition

sequence and 15 nucleotides of the M_2 i3 loop sequence comprised between nucleotide

685 and nucleotide 699. A G nucleotide was also inserted following the 42 nucleotides

of the hairpin loop and upstream of the initial ATG triplet of the EGFP to restore a

correct reading frame. Throughout the text we referred to this plasmid as Sirius-H-

EGFP.

Sirius[PacI-M₂i3(685-699)-Hairpin-G-PacI-M₂i3(685-1101)]EGFP – This bicistronic construct was

created by adding 417 nucleotides of the i3 loop of M_2 from nucleotide 685 to nucleotide

1101 to the plasmid *Sirius[PacI-M₂i3(685-699)-Hairpin-G]EGFP*. An additional PacI

restriction site was also inserted upstream of this segment of the i3 loop. Throughout

the text we referred to this plasmid as Sirius-H-M₂i3(417n)-EGFP.

Sirius[PacI-M₂i3(685-699)-Hairpin-G-PacI-M₂i3(685-699/1072-1101)]EGFP – This bicistronic

construct was created by deleting from the plasmid *Sirius[PacI-M₂i3(685-699)-Hirpin-*

G-PacI-M₂i3(685-1101)]EGFP 372 nucleotides of the M_2 i3 loop, from nucleotide 700

to nucleotide 1071. Throughout the text we referred to this plasmid as Sirius-H-

M₂i3(15n/30n)-EGFP.

Sirius[PacI-M₂i3(685-699)-Hirpin-G-PacI-M₂i3(685-699)]EGFP – This bicistronic construct was

created by adding 15 nucleotides of the i3 loop of M_2 from nucleotide 685 to nucleotide

699 to the plasmid *Sirius[PacI-M₂i3(685-699)-Hirpin-G]EGFP*. An additional PacI

restriction site was also inserted upstream of this short segment of the i3 loop.

Throughout the text we will refer to this plasmid as Sirius-H-M₂i3(15n)-EGFP.

Sirius[PacI-M₂i3(685-699)-Hirpin-G-PacI-M₂i3(685-1071)]EGFP – This bicistronic construct was

created by adding 387 nucleotides of the i3 loop of M₂ form nucleotide 685 to nucleotide

1071 to the plasmid *Sirius[PacI-M₂i3(685-699)-Hirpin-G]EGFP*. An additional PacI

restriction site was also inserted upstream this long segment of the i3 loop. Throughout

the text we will refer to this plasmid as Sirius-H-M₂i3(387n)-EGFP.

Mutants of the Sirius-H-M₂i3(15n/30n)-EGFP plasmid – Based on the sequence alignment of amine and

adenosine GPCRs, several conserved nucleotides were mutated in order to define the

putative IRES sequence. These mutants are reported in **Fig. 3b**.

The fluorescence intensities of the two spectral bands were quantified by the use of a fluorometer and

the EGFP intensity values normalized to those of Sirius.

Constructs for fluorescence microscopy.

M₂-EGFP – The wild type M₂ muscarinic receptor was cloned into the pEGFP-N1 expression cassette,

between the restriction sites HindIII and XbaI.

M₂-mRuby2 – The wild type M₂ muscarinic receptor was cloned into the mRuby2 vector (Addgene

plasmid #40260).

M₂tail(368-466)-EGFP – The M₂tail(368-466)-EGFP was constructed so that the receptor protein would

start from the third in-frame methionine of the M₂ i3 loop (M368). The EGFP gene was

fused C-terminally following a short restriction site sequence (TCTAGA) of the XbaI

enzyme.

M₂tail(368-466)-mRuby2 – The M₂tail(368-466)-mRuby2 was constructed so that the receptor protein

would start from the third in-frame methionine of the M₂ i3 loop (M368). The mRuby2

gene was fused C-terminally following a short restriction site sequence (TCTAGA) of

the XbaI enzyme.

M$_2$-mRuby2-STOP-M$_2$-i3-tail-EGFP – This mega construct was obtained by fusing two preceding

953   constructs, M$_2$-mRuby2 and M$_2$tail(368-466)-EGFP, in a unique plasmid, but the M$_2$tail

954   started at codon 229 (M$_2$tail(229-466)-EGFP). After the stop codon of mRuby2 and

955   before the codon 229 of M$_2$tail was inserted a short restriction site sequence (TCCGGA)

956   of the BspEI enzyme (Fig. 4).

957   *M$_2$(M368A)-mRuby2-STOP-M$_2$-i3-tail-EGFP* – This mega construct is analogous to *M$_2$-mRuby2-STOP-*

958   *M$_2$-i3-tail-EGFP, but* the third in-frame methionine of the i3-loop of the M$_2$ muscarinic

959   receptor was mutated to alanine (M368A) (**Supplementary Fig. 1**).

960   *M$_2$fr.sh-EGFP* – A single base insertion (G) upstream of nucleotide 1102 of the wild type M2 receptor,

961   just before the third in-frame methionine of the i3-loop (M368), induces a frame shift

962   that, following 2 aminoacids, generates a stop codon (TAA) in the amino acid position

963   370 of the new reading frame. The construct is then fused to EGFP.

964   *M$_2$fr.sh-mRuby2* – As above, but the EGFP is replaced by a mRuby2.

965   *M$_2$-GFP11* – A short peptide 16 aminoacids long from the GFP protein, GFP11, was fused to the C-

966   terminus of the muscarinic M$_2$ receptor. GFP11 was custom syntethised and inserted as

967   a linker between the XbaI and XhoI restriction sites in the plasmid backbone of M$_2$-

968   EGFP.

969   *Mito-GFP1-10* – The mitochondrial targeting sequence of the cytochrome c oxidase subunit 8A

970   (COX8A) was fused to the N-terminal of GFP1-10 fragment of GFP. GFP1-10 was

971   purchased from Addgene as Addgene plasmid 70219.

972   *Mito-GFP11* – The mitochondrial targeting sequence of COX8A was fused to GFP11 fragment. Mito

973   sequence was obtained from Addgene plasmid 23348.

974   *SMAC-GFP1-10* – The mitochondrial targeting sequence of the SMAC protein was fused to the N-

975   terminal of GFP1-10 fragment. SMAC was obtained from Addgene plasmid 40881.

976   *SMAC-GFP11* – The mitochondrial targeting sequence of the SMAC protein was fused to GFP11

977   fragment.

978   *M$_2$tail(368-466)-GFP1-10* – The GFP1-10 fragment was fused to the C-terminus of the muscarinic

979   M$_2$tail(368-466) receptor fragment.

980   *M₂tail(368-466)-GFP11* – The GFP11 fragment was fused to the C-terminus of the muscarinic

981   M₂tail(368-466) receptor fragment.

983   **Constructs for in-vitro mitochondrial import assays**

984   *PSP64-M₂tail(368-466)-5Met* - The vector backbone PSP64 (courtesy of Ruaridh Edwards). The

985   following primers: FWD: ATGATGTAAtctagaggaggcggacgc, REV:

986   CATCATCATccttgtagcgcctatgttcttataatg

987   were used to add five methionines to the C-terminal domain of the M₂tail(368-466)-

988   GFP11 template (see above) using Q5® Site-Directed Mutagenesis Kit (New England

989   Biolabs® Inc.). The construct was then extracted with appropriate restriction enzymes

990   (HindIII and XbaI) and ligated into the vector backbone PSP64 (courtesy of Ruaridh

991   Edwards).

992   *PSP64-M2R-5Met* – The following primers: FWD: ATGATGTAAtctagaggaggcggacgc REV:

993   CATCATCATccttgtagcgcctatgttcttataatg were used to add five methionines to the C-

994   terminal domain of to the M₂-GFP11 template (see above) using Q5® Site-Directed

995   Mutagenesis Kit (New England Biolabs® Inc.). The construct was then extracted with

996   appropriate restriction enzymes (HindIII and XbaI) and ligated into the vector backbone

997   PSP64.

999   All constructs were sequenced either with a Genetic Analyzer 3500 (Applied Biosystem®) or by a

1000   professional sequencing service (LGC Genomics, Berlin).

1001

Reverse transcription. Reverse transcription was performed using a 24-nucleotide antisense oligo

(TTACCTTGTAGCGCCTATGTTCTT) directed against the last 24 nucleotides at the 3' end of the M₂

open reading frame. cDNA amplification was carried out with the same antisense oligo used for reverse

transcription, together with a 24 nucleotides sense oligo (ATGAATAACTCAACAAACTCCTCT)

directed against the first 24 nucleotides at the 5' end of the M₂ open reading frame.

Cell culture and transfection. COS-7 (African green monkey kidney fibroblast-like SV40 transformed

cell line) and CHO (Chinese Hamster Ovary) cells (from American Type Culture Collection) were

grown at 5% CO_2 and 37 °C, in high glucose Dulbecco's modified Eagle's medium (DMEM)

supplemented with 10% fetal bovine serum (FBS), 2 mM L-glutamine, 50 IU/ml penicillin, and 50

mg/ml streptomycin, and non-essential amino acids. HeLa cells (from American Type Culture

Collection) were cultured in the same medium, but with the double amount of L-glutamine. SHSY-5Y

cells were cultured in Roswell Park Memorial Institute (RPMI) medium supplemented with 10% FBS,

2 mM L-glutamine, 50 IU/ml penicillin, and 50 mg/ml streptomycin. Media and reagents used for cell

cultures were from Euroclone, PAN Biotech, Biochrome and Sigma-Aldrich. Cell transfection was

performed using FuGene® transfection reagents (Promega) as recommended by the manufacturer.

Eighty percent confluent cells were washed twice with free-serum medium and the mix of FuGene and

plasmid DNA was added. In every experimental condition 6-8 µg of plasmid DNA per Petri (10 cm^2)

were used. Cells were grown for 48 h post-transfection before use. In co-transfection experiments, half

amount of each plasmid DNA was used. All microscopy experiments were performed with transiently

transfected HEK293 (Human Embryonic Kidney, ECACC 96121229 from Sigma-Aldrich Chemie

GmbH) and COS-7 cells. HEK293 cells were cultured in DMEM (Dulbecco's modified Eagle's medium

PAN biotech, Aidenbach, Germany), supplemented with 4.5 g/L Glucose, 2 mM L-Glutamine, 10%

FCS, 100 units/mL penicillin and 0.1 mg/mL streptomycin and maintained at 37 °C and 5% CO_2. Cells

cultured in 15-cm dishes were split at a 1:36 ratio into 6-well plates containing poly-D-lysine (PDL)-

coated 24 mm glass coverslips. Cells were transfected using Effectene transfection reagent (QIAGEN)

according to the manufacturer's instructions. Cells seeded on PDL-coated coverslips in 6-well plates

were transfected 16 hours after seeding with 0.6 µg plasmid/well.

1029

Fluorescence microscopy. Fluorescence microscopy experiments were performed either on a Leica

SP5 (or SP8) Confocal Microscopes, using HyD photon counting detectors and an Argon/Ion laser (or

white light laser) source for 488 nm/633 nm excitation and a solid-state diode laser (or white light laser)

for 561 nm excitation. A solid-state diode laser was used for 405 nm excitation. A 40x 1.3 NA objective

was used, and the electronic zoom was set to achieve a pixel size of 50 nm. For visualization,

mitochondria of transfected cells were labelled for 30 minutes at 37 C with 50nM MitoTracker® Deep

Red FM (Invitrogen™, Thermo Scientific).

1037

Ligand binding studies. Cells were lysed 48 h post-transfection with ice-cold 2 mM EDTA, 1 nM Na-

HEPES (pH 7.4), scraped with a rubber policeman in the presence of ~4 ml rinsing medium, and

homogenized with a Polytron homogenizer (ULTRA-TURRAX) in binding assay buffer (50 mM Tris-

HCl, pH 7.4, 155 mM NaCl, 0.01 mg/ml bovine serum albumin). Radioligand bindings were carried out

at 37 °C for 3 h in a final volume of 1 ml. Non-specific binding was determined in presence of 1 μM

atropine. For competition binding assays, increasing concentrations of carbachol (from 0.01 μM up to

10 μM) were used in presence of 200 pM [^{3}H]N-Methylscopolamine ([^{3}H]NMS; 78.9 Ci/mmol;

PerkinElmer).

The bound ligand was separated from the unbound ligand using glass-fiber filters (Whatmann, GF/B)

with a Brandel Cell Harvester, and filters were counted with a scintillation β-counter. K_D values of

[^{3}H]NMS were calculated in direct saturation experiments, whereas carbachol Inibitory Concentration

$(IC)_{50}$ values were calculated in competition curves, against 400 pM [^{3}H]NMS, fitted to one binding

models using the iterative, non-linear least-squares regression analysis of OriginPro 7.5 software

(OriginLab Corporation).

Phosphatidylinositol Breakdown Assay. Transfected COS-7 cells were incubated with myo-

[^{3}H]inositol (23 Ci/mmol; PerkinElmer) for 48 h. Immediately before the experiment the cells were

washed twice with phosphate-buffered saline and incubated for 15 min in Eagle's medium containing

10 mM LiCl and 20 mM HEPES. The medium was then replaced by 0.25 ml of the same medium

containing the experimental agents. After a 1-hour incubation at 25 °C the reaction was arrested by the

addition of 0.75 ml of 7.5% (w/v) ice-cold trichloroacetic acid followed by a 30-min incubation on ice.

The trichloroacetic acid was extracted with water-saturated diethyl ether (3 × 4 ml), and levels of IP_1

determined by anion exchange chromatography.

1060

Adenyl Cyclase Assay. COS-7 cells were transfected with plasmid(s) containing the receptor of interest

plus adenyl cyclase V. 24 h after transfection the cells were detached with trypsin and re-cultured in 24-

well plates. Following an additional period of 24 h culture, they were assayed for adenyl cyclase activity.

The assay was performed in triplicate. In brief, cells were incubated in 24-well plates for 2 h with 0.25

ml/well of fresh growth medium containing 5 μCi/ml [^{3}H]adenine (PerkinElmer). The medium was then

replaced with 0.5 ml/well of DMEM containing 20 mM HEPES, pH 7.4, 0.1 mg of bovine serum

albumin, and the phosphodiesterase inhibitors 1-methyl-3-isobutylxanthine (0.5 mM) and Ro-20–1724

(0.5 mM). AC activity was stimulated by the addition of 1 μM forskolin in the presence or absence of

carbachol. After 10 min of incubation at +30 °C the medium was removed, and the reaction terminated

by addition of perchloric acid containing 0.1 mM unlabeled cAMP followed by neutralization with

KOH. The amount of [^{3}H]cAMP formed under these conditions was determined by a two-step column

separation procedure (referenza?).

1073

Immunoblot of muscarinic M$_2$ receptor mutants transfected in HEK293 cells. Cells were seeded in

10 cm plates and transfected after 24h according to manufacturer protocols (Effectene, Qiagen) using 2

μg of plasmid DNA. 48h after transfection cells were washed (2x) in ice cold PBS. 200 μL per plate of

ice-cold lysis buffer were added. Lysis buffer was a RIPA buffer: 50 mM Tris-HCL pH 8, 150 mM

NaCl, 1% TRITON, 0.5% sodium deoxycholate and 0.1% SDS. The following compounds were added

to the buffer to a final 1x dilution: 100x Halt Protease Inhibitor Cocktail (Thermo Fisher), 100x 0.5 M

EDTA and (10x) 10 mM PMSF. Cells were then scraped and the cell lysate transferred to pre-cooled

1.5 mL Eppendorf tubes. The tubes were placed in a Thermomixer at 4 °C, 300 rpm for 30 minutes.

Then the Eppendorf tubes were centrifuged for 30 minutes at 14000 rpm at 4 °C. The supernatant was

then transferred to new Eppendorf tubes. Protein concentration in the cell lysates was quantified by a

BCA assay (Pierce BCA Protein Assay Kit, from Thermo Fisher) according to manufacturer

instructions.

Cell lysates were loaded into 10% polyacrylamide gels. Cell lysates were loaded using Laemmli Buffer

2x (Sigma Aldrich) in equal mass. As a reference marker we loaded 5 uL of PageRuler Prestained

Protein Ladder (Thermo Fisher). Gels were mounted into a Mini Protean Tetra Cell kit (BioRad) using

1x SDS running buffer. Gels were ran using a constant voltage of 90V for approximately 15 minutes,

and afterwards at 130V for 45 minutes. Gels were transferred to PVDF membranes. Wet transfer was

achieved by using the Mini Trans Blot Module (BioRad) in Wet Transfer Buffer (25 mM Tris pH 8.3,

192 mM Glycine and 20% MetOH) at 350 mA for 90 minutes. The immunoblotted membranes were

then blocked in 5% Milk in TBS-T for 1 hour at room temperature. The membrane was then incubated

overnight at 4 °C with primary antibody solution using the following concentrations depending on the

antibody used: 1:1000 anti **β**-actin (13E5) rabbit primary antibody (Cell Signaling), 1:1000 anti myc-

Tag (9B11) mouse primary antibody (Cell Signaling). After overnight incubation, the membranes were

washed 3x 10 minutes in TBS-T and incubated at RT for 1 h with a secondary antibody (1:10000 anti-

mouse IgG/(anti rabbit) HRP-linked antibody from Cell Signaling).

After the incubation with the secondary antibody, the membrane was washed 3x 10 minutes in TBS-T.

Detection was achieved by eliciting HRP luminescence by incubating the membrane with SuperSignal

WestFemto solution (Thermo Fisher) according to the manufacturer's instructions, and imaging the

membrane using a c600 Transilluminator from Azure Biosystems.

1103

Animal Experiments. All animal experiments were carried out according to the German animal welfare act,

considering the guidelines of the National Institute of Health and the 2010/63/EU Directive of the European

Parliament on the protection of animals used for scientific purposes. The animals had free access to food and

water and were kept in individually ventilated cages under a 12h:12h light/dark regime (light from 6:30 am to

6:30 pm), a constant 22±2°C temperature and 55±10% humidity.

The transgenic line B6.Cg-Chrm2tm1.1Hze/J, also known as Chrm2-tdT-D knock-in, was obtained from 'The

Jackson Laboratory' (Stock No: 030330). They are heterozygous animals, designed to have expression of a

Chrm2/tdTomato fusion protein directed by the endogenous M2 muscarinic acetylcholine receptor

promoter/enhancer sequences.

1113

Tissue preparation and immunoblot of endogenous M_2 receptor in Chrm2-tdT-D knock-in mice.

Mouse tissues were weighed out and ice-cold lysis buffer was added to approx. 0.2-0.5mg tissue/ml

buffer (10mM K2HPO4, 150mM NaCl, 5mM EDTA, 5mM EGTA, 1% Triton X-100, 0.2% Na-

deoxycholate. Fresh protease inhibitors PMSF 0.5mM, Na3VO4 0.1mM, NaF 50mM, Jenny's mix

(Soybean trypsin inhibitor 3.2μg/ml, Aprotinin 2μg/ml, Benzamidin 0.5mM) were added to the buffer.

Samples were first homogenized, and then pellet debris were centrifuged at 4°C for 30 minutes at max

speed (tabletop) and the supernatant (soluble proteins) was transferred into fresh tube (pellets = non-

soluble proteins)

Protein concentration were determined with BCA assay. Cell lysates were loaded using Laemmli Buffer

2x (Sigma Aldrich) in equal mass (see above for the immunoblot procedure) into 8% polyacrylamide

gels.

The membrane was then incubated overnight at 4 °C with primary antibody solution (1:1000 anti **β**-

actin (13E5) rabbit primary antibody #4970 from Cell Signaling) (1:1000 anti GAPDH) (1:1000 anti-

TdTomato rabbit primary antibody #632496 from Takara).

1128

Isolation of mitochondrial fractions and immunodetection. Cells were seeded in 15 cm plates and

transfected after 24h according to manufacturer protocols (Effectene, Qiagen) using 5 μg of plasmid

DNA. 48h after transfection cells were harvested and centrifuged at 1000 rpm for 10 minutes,

resuspended in NaCl 0.9% (w/v) and centrifuged at 1000 rpm for 10 minutes.

Mouse tissue were freshly excised and placed in ice cold PBS (1x) . The samples were washed with

NaCl 0.9% (w/v). The homogenization steps were performed with Precellys Evolution Homogenizer, at

4500 rpm for 10 seconds.

The samples were processed according to manufacturer protocols (Qproteome Mitochondria Isolation

Kit, Qiagen). From the different steps were obtained three different fractions: the cytosolic, the

microsomal and the mitochondrial fraction.

The fresh mitochondrial pellet was resuspended in 150μL of ice-cold lysis buffer and the protein

concentration of the three fractions from each sample was quantified with a BCA assay (see above).

Cell lysates were loaded into 12% polyacrylamide gels using an SDS loading buffer (6x solution: 2.28g

Tris-HCl, 2.313 DTT, 6g SDS, 30% Glycerol and 0.03% bromophenol blue in a final volume of 50 mL)

in equal mass (see above for the immunoblot procedure).

The membrane was then incubated overnight at 4 °C with primary antibody solution using the following

concentrations: 1:1000 anti **β**-actin (13E5) rabbit primary antibody (#4970 from Cell Signaling), 1:1000

anti-tdtomato rabbit primary antibody (#632496 from Takara), 1:1000 anti **β**-tubulin mouse primary

antibody, 1:1000 anti GAPDH mouse primary antibody, 1:1000 anti COXIV mouse primary antibody

(#ab33985 from abcam).

1149

Quantification of M$_2$tail(368-466) mitochondrial import kinetics from confocal time sequences.

Before imaging, 2.5×10^4 HEK-TSA cells were seeded in each PDL-coated well of the μ-Slide 8 well

(Ibidi) with DMEM (ScienCell™) supplemented with 10% FBS and 1% P/S overnight, and transfected

with a total of 2 μg C-terminal EGFP tagged M$_2$ tail using Lipofectamine 2000 (Thermo Fisher

Scientific) according to the manufacturer protocol after washing with 200 μL PBS and refreshing with

100 μL DMEM (ScienCell™) supplemented with 10% FBS and 1% P/S. After transfection for five

hours, mitochondria of transfected cells were rapidly labelled with 50nM MitoTracker® Deep Red FM

(Invitrogen™, Thermo Scientific) in FluoroBrite DMEM medium (Thermo Scientific) supplemented

with 1% P/S for 20 minutes after the removal of old culture medium. After mitochondria labeling, the

μ-Slide were washed twice with FluoroBrite DMEM medium (Thermo Scientific) and subsequently

mounted onto the TCS SP8 confocal microscope (Leica) (See imaging). Firstly, the stage of the

microscope was equilibrated in temperature, and the acquisition mode was set to the tile scanning. After

storing the position of each desired transfected cells, the imaging mode was set to XYT and a HC PL

APO CS2 40×/NA1.3 oil immersion objective was selected. Both DIC channel and HyD were utilized

to obtain 512×512 pixels images. Laser wavelengths were set as described in the section Fluorescence

Microscopy above. All settings remained the same throughout the whole measurement. Images were

taken at every 30 minutes over a 5h interval. The images were afterwards analyzed by ImageJ software,

using the Plugin 'JACoP'. Briefly, the degree of colocalization between EGFP-tagged M$_2$tail(368-466)

(CH1) and MitoTracker stained mitochondria (CH2) was quantified by Mander's colocalization

coefficients [39]. The two colocalization coefficients M1 ($M1 = \frac{CHB}{CH1}$) and M2 ($M2 = \frac{CHB}{CH2}$), CHB indicates

overlapping pixels in channel 1 and channel 2. A threshold was determined for individual channel in

order to distinguish specific signals from nonspecific ones. Due to their sensitivities to thresholds, the

Pearson correlation coefficient (PCC)[40], as well as the faster decaying attribute of nonspecific signals,

were also taken into account while estimating thresholds for each channel. For the analysis of each cell,

the threshold for Mitotracker channel was kept constant. Whereas the threshold for EGFP channel was

evaluated based on the PCC value.

1176

Transfection and expression of split-GFP constructs in HEK293 cells. To perform the sequential

split GFP complementation assay illustrated in **Figures 4h-i and 5f**, 3×10^5 HEK-TSA cells were seeded

on a 10% poly-D-lysine (PDL)-coated 25mm coverslip in each well of a Falcon® 6-well flat-bottom

plate (StemCell™) and cultured overnight. Cells were transfected using Effectene (see above) with a

0.4 μg Mito-GFP1-10 plasmid. The cells were then washed once with fresh 2 mL phosphate-buffered

saline (PBS) first and supplemented with 1.5 mL fresh DMEM medium (ScienCell™) supplemented

with 10% FBS and 1% P/S. After adding the transfection mixture, the cells were cultured overnight.

After the first transfection, a second transfection of a 0.6 μg plasmid encoding either the C-terminal

GFP11 tagged M_2tail(368-466) or M_2 receptor was performed using the Effectene Transfection Reagent

(Qiagen). After washing each well with 2 mL PBS, the old medium was exchanged with 1.5 mL low

glucose Minimum Essential Medium (MEM) (Gibco™, Thermo Scientific) supplemented with 1% P/S,

in order to create a stress-like cell starvation condition. After a four hours starvation, the cells were fixed

using 4% polyformaldehyde (PFA)/PBS at 37°C for 30 minutes and washed twice with PBS before

imaging.

1191

Oxygen consumption measured with the Seahorse assay. 5×10^4 COS7 cells were seeded in seahorse

24MW plates with Complete media (DMEM supplemented with 10%FBS and 1X PS) and then

incubated for about 4h at 37C in a cell incubator. After this incubation, cells were washed twice with

Seahorse media supplemented with glutamate (1 mM), pyruvate (1 mM) and glucose (20 mM) and then

incubated for 1h at 37 °C in an incubator without CO2 before being analyzed in the Seahorse XF

Analyzers. In particular, the baseline cell oxygen consumption rate (OCR) was measured at the

beginning of the assay, followed by measurements of ATP linked respiration and maximal respiratory

capacity after the addition of respectively oligomycin (1 μM) and the FCCP protonophore (1 μM).

Finally, rotenone (1 μM), an inhibitor of complex I, and antimycin (1 μM), inhibitor of complex III,

were added to shut down completely the electron transport chain (ETC) function, revealing the non-

mitochondrial respiration.

1203

Generation of M368A hiPSC clone. The human induced pluripotent stem cell (hiPSC) line BIHi005-

A (https://hpscreg.eu/cell-line/BIHi005-A) was used to generate an isogenic iPSC clone (BIHi005-A-

39) with **M368A mutation** using a protocol previously described[41]. In brief, small guide RNA (sgRNA)

targeting the M_2 gene close to the mutation site to be corrected were designed using the

http://crispor.tefor.net/ webpage and the sgRNA (5`- AGTCATCTTCACAATCTTGC-3`) were

synthesized by Integrated DNA technologies (https://eu.idtdna.com). The donor ssODN template was

designed to convert the ATG to GCT and was synthesized as an Ultramar DNA oligo by IDT. (5´-

AAAAGTGACTCATGTACCCCAACTAATACCACCGTGGAGGTAGTGGGGTCTTCAGGTCAG

AATGGAGATGAAAAGCAGAATATTGTAGCCCGCAAGATTGTGAAGGCTACTAAGCAGCC

TGCAAAA-3´). The Ribonucleoprotein (RNP) complexes were prepared by incubating of 1.5μg Cas-

9 protein and 360 ng gRNA for 10 minutes at room temperature. For delivery of RNPs and ssODN

template, 10μl cell suspension containing Cas9 RNPs and ssODN and 1x105 cells were electroporated

using the Neon transfection System (ThermoFisher Scientific). The electroporated cells were plated in

one well of 6 well plate with StemFlex media (ThermoFisher Scientific) supplemented CloneR™

(Stemcell technologies). Three days after the transfection we analyzed the bulk population using the

Sanger sequencing to estimate the editing efficiency. Thereafter, the automated single cell cloning of

the genome edited cell pool was performed as described in protocol [42]. The clones were screened by

SANGER sequencing. The positive confirmed clones were banked and characterized for pluripotency

and karyotype stability as described in [43].

1223

Briefly, for pluripotency assay, single cells were labelled with conjugated antibodies. Surface markers

(SSEA1, SSEA4 and TRA1-60) staining was performed in unfixed cells by incubation with antibodies

diluted in 0.5% BSA/PBS (10 min, 4 °C, dark). Intracellular markers (OCT3/4, NANOG) were stained

by fixation/permeabilization solution (Miltenyi Biotec, 130-093-142) (30 min, 4 °C, dark) followed by

antibody incubation in permeabilization buffer (Miltenyi Biotec) (30 min, 4 °C, dark). Analysis was

done using the MACSQuant® AnalyzerVYB and FlowJo v10.4.

1230

Briefly, for karyotype stability, genomic DNA was isolated using the DNeasy blood and tissue kit

(Qiagen, Valencia, CA, United States) and samples were analyzed using the human Illumina OMNI-

EXPRESS-8v1.6 BeadChip. First, the genotyping was analyzed using GenomeStudio 1 genotyping

module (Illumina). Thereafter, KaryoStudio 1.3 (Illumina) was used to perform automatic normalization

and to identify genomic aberrations utilizing settings to generate B-allele frequency and smoothened

Log R ratio plots for detected regions. The stringency parameters used to detect copy number variations

(CNVs) were set to 75 kb (loss), 100 kb (gain) and CN-LOH (loss of heterozygosity) regions larger than

3 MB.

1239

Culture and handling of hiPSCs cells. hiPSCs were cultured on Geltrex (Thermo Fisher) coated plates

using: Essential 8 medium (Thermo Fisher). The culture was routinely monitored for mycoplasma

contamination using a PCR assay. 10 µM ROCK Inhibitor Y-27632 2HCl (Selleck Chemicals, #SEL-

S1049) was added after cell splitting to promote survival. hiPSC cultures were kept in a humidified

atmosphere of 5% CO_2 at 37°C and 5% oxygen. hiPSCs were undirectedly differentiated to a mixture

of three lineages (ectodermal, endodermal and mesodermal) by replacing Essential 8 medium with

DMEM-F12 medium supplemented with 10% FBS for seven days. 10 µM ROCK Inhibitor Y-27632

2HCl (Selleck Chemicals) was added after cell splitting to promote survival.

1248

Cell proliferation assay with HEK293 cells. $4x10^5$ HEK293 cells 24h after transfection and $4x10^5$

hPSCs were seeded respectively in T25 flask and pre-coated T25 flask, in culture medium with Atropine

at final concentration of 100nM; the medium with Atropine was replaced every day. After 72h from

transfection of the HEK293 and from seeding of the hiPSCs, the medium and the cells were collected

and centrifuged, the pellet was resuspended in the same amount of medium for each pellet; an aliquots

of the resuspension was mixed with the same amount of Trypan blue (ratio 1:1), 10 microliters were

used for counting in a Neubauer chamber.

1256

Fluorescence plate reader assays. HEK293 cells were grown on 6 well plates without coverslips, and,

upon reaching 60-70% confluence, the cells were transfected using Effectene (Quiagen, Venlo

Netherlands) according to the manufacturer's instructions. After 24 h the cells were split and transferred

to 96 well (black plate) at a seeding density of 50,000 cells/well. The 96 well plate was pre-coated with

Poly-D-Lysine for 30 min at RT. The medium was removed and each well was washed 3x with 100µl

HBSS. The cells were finally analyzed by a fluorescence plate reader (Neo2, BioTek, Bad

Friedrichshall) in HBSS.

1264

1265 **Extracellular oxygen consumption.** HEK293 cells 24h after transfection and hPSCs were harvested

1266 and seeded respectively in 96 multiwell black plate and pre-coated (Geltrex A1413302, Thermo Fisher)

1267 96 multiwell black plate, $5x10^4$ cells for well; after incubation O/N at 37 °C and in 5% CO2, the cell

1268 culture medium was replaced by HBSS and the assay were performed according to the manufacturer

1269 protocols (abcam Extracellular Oxygen Consumption Assay).

1270

1271 **Author Contributions**

1272 PA, FP, IF, ZW, MR, IP, RE, HZ, UZ, MET, JDG, SS, ART, SA conduced primary experiments; MG, CP,

1273 GA, FM, BP, AR contributed to the experimental data; NT and SD contributed tools and reagents. PA, IP, IF,

1274 MR, UZ, AT, KT, VF, FA, FG, MS contributed to experimental design and assisted in writing paper, PA, MJL

1275 and RM conceived and lead the study, and wrote the paper.

1276

1277 **Acknowledgments**

1278 We acknowledge funding from European Union's Horizon2020 Marie Skłodowska-Curie Actions (MSCA)

1279 Program under Grant Agreement 641833 (ONCORNET). We are grateful to Dr. Marco Marcia (EMBL) for

1280 insightful discussion.

1281

1282 **Competing Interests**

1283 The authors declare that they have no conflicts of interest with the contents of this article.

1284

1285 **Data Availability Statement**

1286 Source data are provided with this manuscript.

1287

References

Kruse, A. C. *et al.* Muscarinic acetylcholine receptors: novel opportunities for drug development. *Nat Rev Drug Discov* **13**, 549-560, doi:10.1038/nrd4295 (2014).

Fryer, A. D., Christopoulos A. & Nathanson N.M. in *Handb. Exp. Pharmacol.* Vol. 208 1-499 (2012).

Novi, F., Scarselli, M., Corsini, G. U. & Maggio, R. The paired activation of the two components of the muscarinic M3 receptor dimer is required for induction of ERK1/2 phosphorylation. *J Biol Chem* **279**, 7476-7486, doi:10.1074/jbc.M306818200 (2004).

Novi, F., Scarselli, M., Corsini, G. U. & Maggio, R. An unusual form of the association binding kinetics of N-[3H]methylscopolamine to the split muscarinic M2trunk/M2tail receptor. *J Pharmacol Exp Ther* **305**, 786-795, doi:10.1124/jpet.102.045393 (2003).

Jackson, R. J. & Kaminski, A. Internal initiation of translation in eukaryotes: the picornavirus paradigm and beyond. *RNA* **1**, 985-1000 (1995).

Hellen, C. U. & Sarnow, P. Internal ribosome entry sites in eukaryotic mRNA molecules. *Genes Dev* **15**, 1593-1612, doi:10.1101/gad.891101 (2001).

Calebiro, D., Nikolaev, V. O., Persani, L. & Lohse, M. J. Signaling by internalized G-protein-coupled receptors. *Trends in Pharmacological Sciences* **31**, 221-228, doi:https://doi.org/10.1016/j.tips.2010.02.002 (2010).

Zhang, X., Bedigian, A. V., Wang, W. & Eggert, U. S. G protein-coupled receptors participate in cytokinesis. *Cytoskeleton (Hoboken)* **69**, 810-818, doi:10.1002/cm.21055 (2012).

Hebert-Chatelain, E. *et al.* A cannabinoid link between mitochondria and memory. *Nature* **539**, 555-559, doi:10.1038/nature20127 (2016).

Suofu, Y. *et al.* Dual role of mitochondria in producing melatonin and driving GPCR signaling to block cytochrome c release. **114**, E7997-E8006, doi:10.1073/pnas.1705768114 %J Proceedings of the National Academy of Sciences (2017).

Maggio, R. *et al.* Variants of G protein-coupled receptors: a reappraisal of their role in receptor regulation. *Biochem Soc Trans* **44**, 589-594, doi:10.1042/BST20150239 (2016).

Maggio, R., Vogel, Z. & Wess, J. Reconstitution of functional muscarinic receptors by co-expression of amino- and carboxyl-terminal receptor fragments. *FEBS Lett* **319**, 195-200, doi:10.1016/0014-5793(93)80066-4 (1993).

Park, P. S. & Wells, J. W. Monomers and oligomers of the M2 muscarinic cholinergic receptor purified from Sf9 cells. *Biochemistry* **42**, 12960-12971, doi:10.1021/bi034491m (2003).

Bathe-Peters, M., Gmach, P., Annibale, P. & Lohse, M. J. Linescan microscopy data to extract diffusion coefficient of a fluorescent species using a commercial confocal microscope. *Data Brief* **29**, 105063, doi:10.1016/j.dib.2019.105063 (2020).

Dong, C., Filipeanu, C. M., Duvernay, M. T. & Wu, G. Regulation of G protein-coupled receptor export trafficking. *Biochim Biophys Acta* **1768**, 853-870, doi:10.1016/j.bbamem.2006.09.008 (2007).

Kamiyama, D. *et al.* Versatile protein tagging in cells with split fluorescent protein. *Nat Commun* **7**, 11046, doi:10.1038/ncomms11046 (2016).

Bonfili, L. *et al.* Essential amino acid mixtures drive cancer cells to apoptosis through proteasome inhibition and autophagy activation. *FEBS J* **284**, 1726-1737, doi:10.1111/febs.14081 (2017).

Thakor, N. & Holcik, M. IRES-mediated translation of cellular messenger RNA operates in eIF2alpha- independent manner during stress. *Nucleic Acids Res* **40**, 541-552, doi:10.1093/nar/gkr701 (2012).

Holcik, M. & Sonenberg, N. Translational control in stress and apoptosis. *Nat Rev Mol Cell Biol* **6**, 318-327, doi:10.1038/nrm1618 (2005).

| 1337 | 20 | Newman, L. E., Schiavon, C. & Kahn, R. A. Plasmids for variable expression of proteins targeted to the mitochondrial matrix or intermembrane space. *Cell Logist* **6**, e1247939, doi:10.1080/21592799.2016.1247939 (2016). |

20 Newman, L. E., Schiavon, C. & Kahn, R. A. Plasmids for variable expression of proteins
targeted to the mitochondrial matrix or intermembrane space. *Cell Logist* **6**, e1247939,
doi:10.1080/21592799.2016.1247939 (2016).
21 Ishizuka, T., Ozawa, A., Katsuura, M., Nomura, S. & Satoh, Y. Effects of muscarinic
acetylcholine receptor stimulation on the differentiation of mouse induced pluripotent stem
cells into neural progenitor cells. *Clin Exp Pharmacol Physiol* **45**, 1198-1205,
doi:10.1111/1440-1681.12993 (2018).
22 Hoogduijn, M. J., Cheng, A. & Genever, P. G. Functional nicotinic and muscarinic receptors
on mesenchymal stem cells. *Stem Cells Dev* **18**, 103-112, doi:10.1089/scd.2008.0032 (2009).
23 Weinberg, Z. Y. & Puthenveedu, M. A. Regulation of G protein-coupled receptor signaling
by plasma membrane organization and endocytosis. *Traffic* **20**, 121-129,
doi:10.1111/tra.12628 (2019).
24 Jong, Y. I., Harmon, S. K. & O'Malley, K. L. GPCR signalling from within the cell. *Br J*
Pharmacol **175**, 4026-4035, doi:10.1111/bph.14023 (2018).
25 Benard, G. *et al.* Mitochondrial CB(1) receptors regulate neuronal energy metabolism. *Nat*
Neurosci **15**, 558-564, doi:10.1038/nn.3053 (2012).
26 Lefkimmiatis, K., Leronni, D. & Hofer, A. M. The inner and outer compartments of
mitochondria are sites of distinct cAMP/PKA signaling dynamics. *J Cell Biol* **202**, 453-462,
doi:10.1083/jcb.201303159 (2013).
27 Marti-Solano, M. *et al.* Combinatorial expression of GPCR isoforms affects signalling and
drug responses. *Nature* **587**, 650-656, doi:10.1038/s41586-020-2888-2 (2020).
28 van Heesch, S. *et al.* The Translational Landscape of the Human Heart. *Cell* **178**, 242-260
e229, doi:10.1016/j.cell.2019.05.010 (2019).
29 Pestova, T. V., Shatsky, I. N. & Hellen, C. U. Functional dissection of eukaryotic initiation
factor 4F: the 4A subunit and the central domain of the 4G subunit are sufficient to mediate
internal entry of 43S preinitiation complexes. *Mol Cell Biol* **16**, 6870-6878,
doi:10.1128/mcb.16.12.6870 (1996).
30 Jan, E. & Sarnow, P. Factorless ribosome assembly on the internal ribosome entry site of
cricket paralysis virus. *J Mol Biol* **324**, 889-902, doi:10.1016/s0022-2836(02)01099-9 (2002).
31 Fitzgerald, K. D. & Semler, B. L. Bridging IRES elements in mRNAs to the eukaryotic
translation apparatus. *Biochim Biophys Acta* **1789**, 518-528,
doi:10.1016/j.bbagrm.2009.07.004 (2009).
32 Ingolia, N. T., Hussmann, J. A. & Weissman, J. S. Ribosome Profiling: Global Views of
Translation. *Cold Spring Harb Perspect Biol* **11**, doi:10.1101/cshperspect.a032698 (2019).
33 Zhang, J., Nuebel, E., Daley, G. Q., Koehler, C. M. & Teitell, M. A. Metabolic regulation in
pluripotent stem cells during reprogramming and self-renewal. *Cell Stem Cell* **11**, 589-595,
doi:10.1016/j.stem.2012.10.005 (2012).
34 Guo, C., Sun, L., Chen, X. & Zhang, D. Oxidative stress, mitochondrial damage and
neurodegenerative diseases. *Neural Regen Res* **8**, 2003-2014, doi:10.3969/j.issn.1673-
5374.2013.21.009 (2013).
35 Kritsiligkou, P. *et al.* Unconventional Targeting of a Thiol Peroxidase to the Mitochondrial
Intermembrane Space Facilitates Oxidative Protein Folding. *Cell Rep* **18**, 2729-2741,
doi:10.1016/j.celrep.2017.02.053 (2017).
36 Bonner, T. I., Buckley, N. J., Young, A. C. & Brann, M. R. Identification of a family of
muscarinic acetylcholine receptor genes. *Science* **237**, 527-532, doi:10.1126/science.3037705
(1987).
37 Kozak, M. Constraints on reinitiation of translation in mammals. *Nucleic Acids Res* **29**, 5226-
5232, doi:10.1093/nar/29.24.5226 (2001).
38 Kozak, M. Circumstances and mechanisms of inhibition of translation by secondary structure
in eucaryotic mRNAs. *Mol Cell Biol* **9**, 5134-5142, doi:10.1128/mcb.9.11.5134 (1989).

39   Manders, E. M. M., Verbeek, F. J. & Aten, J. A. Measurement of Colocalization of Objects
1388   in Dual-Color Confocal Images. *J Microsc-Oxford* **169**, 375-382, doi:DOI 10.1111/j.1365-
1389   2818.1993.tb03313.x (1993).
1390   40   Manders, E. M. M., Stap, J., Brakenhoff, G. J., Vandriel, R. & Aten, J. A. Dynamics of 3-
1391   Dimensional Replication Patterns during the S-Phase, Analyzed by Double Labeling of DNA
1392   and Confocal Microscopy. *J Cell Sci* **103**, 857-862 (1992).
1393   41   Richardson, C. D., Ray, G. J., DeWitt, M. A., Curie, G. L. & Corn, J. E. Enhancing homology-
1394   directed genome editing by catalytically active and inactive CRISPR-Cas9 using asymmetric
1395   donor DNA. *Nat Biotechnol* **34**, 339-344, doi:10.1038/nbt.3481 (2016).
1396   42   Vallone, V. F. *et al.* Methods for Automated Single Cell Isolation and Sub-Cloning of Human
1397   Pluripotent Stem Cells. *Curr Protoc Stem Cell Biol* **55**, e123, doi:10.1002/cpsc.123 (2020).
1398   43   Metzler, E., Telugu, N., Diecke, S., Spuler, S. & Escobar, H. Generation of two human
1399   induced pluripotent stem cell lines derived from myoblasts (MDCi014-A) and from peripheral
1400   blood mononuclear cells (MDCi014-B) from the same donor. *Stem Cell Res* **48**, 101998,
1401   doi:10.1016/j.scr.2020.101998 (2020).
1402