One gene - two proteins: The C-terminus of the prototypical M2 muscarinic receptor localizes to the mitochondria and regulates cell respiration.

Irene Fasciani[1†], Francesco Petragnano[1†], Ziming Wang[2], Ruairidh Edwards[3], Narasimha Telegu[2], Ilaria Pietrantoni[1], Ulrike Zabel[4], Henrik Zauber[2],Marlies Grieben[2], Maria E. Terzenidou[3], Jacopo Di Gregorio[1], Cristina Pellegrini[1], Silvano Santini Jr[5], Anna R. Taddei[6], Barbel Pöhl[2], Stefano Aringhieri[7], Marco Carli[7], Gabriella Aloisi[1], Francesco Marampon[8], Alexandra Roman[2], Sebastian Diecke[2], Vincenzo Flati[1], Franco Giorgi[7], Fernanda Amicarelli[5], Andrew B. Tobin[3], Marco Scarselli[7], Kostas Tokatlidis[3], Mario Rossi[1], Martin J. Lohse[2,4,9]*, Paolo Annibale[2,10]*, Roberto Maggio[1]*.

[1] Department of Biotechnological and Applied Clinical Sciences, University of L'Aquila, 67100 L'Aquila, Italy.
[2] Max Delbrück Center for Molecular Medicine, 13125 Berlin, Germany.
[3] Centre for Translational Pharmacology, Institute of Molecular, Cell and Systems Biology, College of Medical, Veterinary and Life Sciences, University of Glasgow, G128QQ Glasgow, UK.
[4] Institute of Pharmacology and Toxicology, University of Würzburg, 97078 Würzburg, Germany.
[5] Department of Life, Health and Environmental Sciences, University of L'Aquila, 67100 L'Aquila, Italy.
[6] Section of Electron Microscopy, Great Equipment Center, University of Tuscia, Viterbo, Italy.
[7] Department of Translational Research and New Technology in Medicine, University of Pisa, 56126 Pisa, Italy.
[8] Department of Radiotherapy, University of Roma La Sapienza, 00151 Roma, Italy.
[9] ISAR Bioscience Institute, 82152 Munich, Germany.
[10] School of Physics and Astronomy, University of St Andrews, KY16 9SS St Andrews.
[†] Equal contribution
*email: martin.lohse@isarbioscience.de;
*email: roberto.maggio@univaq.it;
*email: paolo.annibale@mdc-berlin.de, pa53@st-andrews.ac.uk

Abstract

Muscarinic acetylcholine receptors are prototypical G protein-coupled receptors (GPCRs), members of a large family of seven transmembrane receptors that mediates a wide variety of extracellular signals. We show here, for the first time in a GPCR, that the carboxyl terminal fragment of the muscarinic M_2 receptor, containing the transmembrane regions 6 and 7 (M_2tail), is expressed by virtue of an internal ribosome entry site located in the third intracellular loop. Single cell imaging reveals that M_2tail, whose expression is upregulated in cells undergoing integrated stress response, does not follow the normal route to the plasma membrane, but is almost exclusively sorted to the mitochondria inner membrane: here it controls oxygen consumption, cell proliferation and the formation of reactive oxygen species by reducing oxidative phosphorylation. Crispr/Cas9 editing of the key methionine where cap-independent translation begins in human induced pluripotent stem cells, reveals the physiological role of this process in influencing cell proliferation and oxygen consumption at the endogenous level. The expression of the C-terminal domain of a GPCR, capable of regulating mitochondrial function, constitutes a hitherto unknown mechanism notably unrelated to its canonical signalling function as a GPCR at the plasma membrane. This work thus highlights a potential novel mechanism thar cells may use for controlling their metabolism under variable environmental conditions, notably as a negative regulator of cell respiration.

Introduction

Muscarinic receptors include five structurally related receptors, belonging to class A of the G protein-coupled receptor (GPCR) family [1]. They are comprised of seven trans-membrane domains (TM domains 1-7) connected by intra and extracellular loops. Of these structural elements, the third intracellular (i3) domain is the one conferring a coupling specificity for G proteins, β-arrestin recruitment and endocytic internalization [2]. However, only a small portion of the i3 loop is required for receptor function, and for this reason large deletions of this segment have no effect on receptor activation and internalization [3].

GPCRs are normally distributed at the plasma membrane, where, after activation by an agonist and coupling to either G proteins or β-arrestin, regulate a number of intracellular effectors. Nevertheless, non-canonical localizations of GPCRs have also been described, such as in endosomes [4], cell division compartments (centrosomes, spindle midzone and midbodies) [5], and mitochondria [6],[7] underpinning non-traditional roles for this protein family.

Several studies have shown that the GPCR structurally related transmembrane protein bacteriorhodopsin consists of autonomous folding domains that translocate to the membrane and

interact with each other to function as a proton pump [8,9]. Interestingly, it was also shown that fragments of muscarinic receptors can interact to form functionally active chimeric receptors, highlighting the conserved evolutionary ability of domains from these receptor family to interact among each other, as well as with other proteins. Intriguingly, in 2016, we observed that the insertion of a stop codon in the third intracellular loop of the muscarinic M_2 receptor was not able to abolish its function [10], presenting the suggestive possibility that a cap-independent translation of the M_2 C-terminal domain may occur in this mutant, i.e. that the M_2 mRNA may lead to the translation not only of the full length receptor, but also of its C-terminal domain, that we name here M_2tail.

In this work, we therefore set out to explore whether the expression of the M_2tail, a region containing the TM regions 6 and 7, occurs naturally by a cap-independent translation mechanism. Combining radio-ligand binding, molecular biology approaches, advanced fluorescence imaging and Crispr/Cas9 genetic editing of human induced pluripotent stem cells (hiPSCs) we have gathered compelling data indicating a novel translation mechanisms for C-terminal domain of the M_2 muscarinic receptor under control of a IRES sequence within the M_2 mRNA and that the C-terminal domain of this receptor can indeed be expressed as an independent polypeptide.

While in eukaryotes the most common mechanism of translation initiation is the cap-dependent process, which requires the interaction of several key proteins at the 5'-end (5′ cap) of mRNA molecules, recent years have seen growing evidence that internal ribosome entry site (IRES) sequences, which are common features of virus replication [11], can mediate cap independent translation of eukaryotic RNA(s) [12].

Moreover, by using bimolecular fluorescence complementation, in-vitro mitochondrial import and cell metabolism assays we observe that M_2tail is strongly targeted to the cell mitochondria. Importantly, this mitochondrial targeting has functional ramifications since both cap-dependent mediated overexpression of M_2tail, as well as cap-independent mediated translation starting from the mRNA of the full length M_2 receptor, resulted in a sizable reduction of oxidative phosphorylation. In fact, we show that cellular stress favours the activation of this newly discovered IRES-dependent translation of M_2tail and that the fragment interacts with the Complex V of the respiratory chain on the mitochondria inner membrane. The implication of this newly discovered regulatory mechanism of the IRES mediated translation of the M_2 C-terminal domain might have profound physiological and pathophysiological implications.

In summary our study reports the new observation that alternative reading of a GPCR receptor gene can translate into a fragment of the receptor that serves a cellular function entirely different from canonical GPCR signalling. Taken together these data strongly suggest that the M_2 mRNA yields not only a plasma membrane receptor able to transduce many of the acetylcholine functions but M_2tail, a protein fragment that reduces the activity of mitochondria and ROS accumulation and it might represent a hitherto unappreciated step for in-vivo anti-oxidative stress, during for instance reperfusion, tissue damage, and disease-promoted metabolic cellular stress.

Results

Cap independent production of the C-terminus of the M_2 receptor and evidence for an IRES

We began our work by testing our earlier hypothesis that the last two TM regions (TM6-7) of M_2stop228, a M_2 muscarinic receptor mutant bearing a stop codon at the beginning of the i3 loop in position 228 (Fig. S1), may still be translated [10]. We thus generated EGFP C-terminal fusion of the M_2 wt receptor and M_2stop228: Fig. 1A clearly shows that cells expressing M_2stop228-EGFP display a green fluorescence signal. Moreover, the signal displays a predominant intracellular localization with respect to the wt M_2R, which instead localizes -as expected- largely at the plasma membrane. A key observation is that the position of the stop codon upstream of the i3 loop appears to bear a particular relevance, since if a stop is instead inserted downstream of position 228, e.g. at positions 400, right before the first methionine in TM6, no fluorescence can be observed in cells expressing the mutant M_2stop400-EGFP Fig. 1A.

To further appreciate the origin of this unexpected process -the 228 stop codon should abolish any EGFP signal- we performed radioligand binding experiments in cells transfected with this mutant. M_2stop228 showed both K_D value of [^{3}H]NMS (N-methylscopolamine, an antagonist) calculated from direct saturation experiments, and the IC_{50} values, calculated in competition experiments for the agonist carbachol, similar to those calculated for the wild type M_2 receptor (Table 1), albeit displaying a very low level of expression at the plasma membrane, i.e. 45 fmol/mg of protein. Furthermore, M_2stop228 was found functionally active as it was capable to inhibit adenylyl cyclase [10] and increase ERK phosphorylation (Fig. S2A), even though the extent of response was reduced if compared to the wild type M_2 receptor (Table 1). As a control, receptor fragments, translated from mRNAs comprising only the sequence for TM1-5 (i.e. whose sequences run only until position 228 or 283 of the corresponding full length receptor), called M_2trunk(1-228)/M_2trunk(1-283), did not show any radio-ligand binding activity, despite coding for

the same first five trans-membrane regions (TM1-5) that the canonical translation of M$_2$stop228
should also produce (**Fig. 1B** and **Table 1**) [13].

We showed in the past that the last two TMs (TM6-7) of the M$_2$ receptor can reconstitute a
functional receptor with TM1-5, supporting the notion that the explanation of the apparent
paradox must be that M$_2$tail is translated as an independent polypeptide, as graphically
summarized in **Fig. 1B**.

Fig. 1. **The c-terminus of the M$_2$ receptor is expressed by a cap independent translation**
mechanism, originating in the i3loop.

A, Fluorescence micrographs of HEK293 and COS7 cells respectively expressing M$_2$-EGFP, M$_2$stop228-
EGFP, M$_2$stop400-EGFP, together with an untransfected control. Imaging settings are the same for the
five images, whereas contrast was adjusted to improve visualization. In the untransfected cells and
stop400 panels, the contrast was set to 0. Insets display DIC image of the same field of view (scale bars
10 µm). **B**, Schematic representation of the gene products that reach the plasma membrane of the wild
type M$_2$, M$_2$stop228, M$_2$trunk(1-283) and M$_2$fr.sh. (**Fig. S3**). **C**, Fluorescence micrograph of a

representative HEK293 cell expressing M_2fr.sh.-EGFP. Insets display DIC image of the same field of view
(scale bar 10 μm).

	Binding data			Adenylyl cyclase inhibition assay	
Receptor	Bmax (fmol/mg)	[3H]NMS KD (pM)	Carbachol IC50 (μM)	Carbachol EC50 (μM)	Maximum inhibition of forskolin stimulated cAMP accumulation (%)
M_2	980 ± 60	62 ± 7	2.1 ± 0.6	0.04 ± 0.01	53 ± 5
M_2trunk(1-283)	N.B.				
M_2trunk(1-228)	N.B.				
M_2stop228	45 ± 2	70 ± 10	1.8 ± 0.4	0.13 ± 0.02	27 ± 3

Table 1. Radioligand binding and activation properties of M_2 receptor mutants.

Binding properties and inhibition of forskolin stimulated adenylyl cyclase by M_2, M_2stop228, M_2trunk(1-228)
and M_2trunk(1-283). Untransfected COS-7 cells did not show any [3H]NMS specific binding. N.B. = no
specific [3H]NMS binding. Data are averages with s.e.m.

Evidence of an IRES-mediated M_2 C-terminus translation

M_2stop228 mRNA appears thus able to give rise -despite the presence of a stop codon in the i3
loop- to two distinct protein products (a *trunk* TM1-5 and a *tail* TM6-7), in turn able to
reconstitute a functional receptor at the plasma level.

To challenge this hypothesis and gain further insight into the mechanism at play, we generated a
frame shift mutant of the M_2 receptor: we inserted a single guanine nucleotide in the sequence of
the M_2-EGFP, changing the methionine in position 368 into an aspartate, which in turn generates
an in-frame stop codon at position 370 (**Fig. S1 and Fig. S3**). The observation that cells
transfected with M_2fr.sh.-EGFP retain green fluorescence is most remarkable (**Fig. 1C**), since the
EGFP is now out of the original reading frame and it can only be properly translated by a *cap-*
independent translation mechanism, starting from a methionine in the new reading frame.

Furthermore, the observation that both M_2stop228-EGFP and M_2fr.sh.-EGFP retain green
fluorescence, whereas M_2stop400-EGFP does not, indicates not only that a cap independent
translation mechanism occurs, but also that the starting methionine for the process must be
downstream of stop 228 and upstream of stop 400, leaving as a leading candidate one of the in-
frame methionines of the i3 loop.

To narrow this down, we employed radio-ligand binding data from expression of receptor mutants carrying a double stop codon, the first in position 228 and the second in positions ranging from 248 to 400, including mutations of the first (M248) and second (M296) in-frame methionines in the third i3 loop (Table S1, Fig. S3). The observation that mutation of the first two methionines does not affect B_{max} significantly points to M368 as the most likely site for the cap-independent initiation of translation of the M_2tail, that we now term more specifically M_2tail(368-466). These binding results can furthermore rule out read-through and termination re-initiation: if these mechanisms were responsible for translation of the C-terminal of the M_2stop228, suppression of the first methionine would have abolished the binding. We shall further note that cells expressing M_2stop228 showed only one single M_2 specific cDNA band supporting the concept that the M_2tail(368-466) was produced from the same M_2-mRNA molecule by a cap independent mechanism instead of, for instance, by RNA splicing variants (Supplementary Materials and Methods and Fig. S2B).

IRES position within the M_2 coding sequence

To further validate the results obtained working with full length M_2 receptor mutants and screen for what nucleotides are important for the IRES activity, we additionally prepared a fluorescence reporter construct in which the i3 loop of M_2 was inserted between the two fluorescent proteins Sirius and EGFP (Fig. 2A). The sequence for Sirius is followed by a stop codon, thus making Sirius-M_2i3(417n)-EGFP translate Sirius according to the classical cap-dependent mechanism, reporting on transfection efficiency, and, at the same time, drive EGFP translation only if the M_2i3 loop contains an IRES sequence, effectively acting as a bicistronic expression cassette. We transfected several cell lines with this construct, all positive for the blue and green fluorescence (Fig. S4A). HEK293 cells, used throughout our study, appeared positive for both the blue and green fluorescence, as shown in the inset of Fig. 2A and in line with the results reported in Fig. 1A. To further rule out the possibility that EGFP translation from the Sirius-M_2i3(417n)-EGFP construct could be due to read-through and termination-re-initiation mechanisms, a mRNA hairpin loop was inserted in the sequence after the stop codon of the Sirius protein (Sirius-H-EGFP), abolishing completely EGFP signal.

To narrow down the position of the IRES sequence we divided the M_2 i3 loop in three parts and the resulting nucleotide sequences, 685-699, 685-1071 and 685-699/1072-1101, were inserted downstream of the hairpin loop in three different constructs (Fig. 2A and Fig. S3).

Fig. 2. **Localization of the IRES within the i3 loop the muscarinic M$_2$ receptor using a fluorescence reporter assay.**

A, **left** Schematic representation of bicistronic plasmids bearing the M$_2$ i3 loop between the coding regions of the Sirius and EGFP fluorescent proteins. A guanine nucleotide was originally inserted before the EGFP in the plasmid Sirius-H-EGFP in order to maintain in frame the sequences of the two fluorescent proteins. In all the other plasmids the sequence of the two fluorescent proteins is not in frame. Fifteen nucleotides derived from the M$_2$ i3 loop were inserted between the end of Sirius and the hairpin loop, in order to allow the easy scanning of the ribosome across the stop codon. **right**, IRES activity, highlighting the efficiency of cap-independent translation for the control Sirius-M$_2$i3(417n)-EGFP and the six mutants. The insert shows the difference in the green fluorescence between two of the mutants. Significance values reported in the graphs were determined by a one-tailed Student t-test, p-values: ** $0.001 < p < 0.01$. **B,** Expression of the construct M$_2$-Hairpin-EGFP in HEK293 cells: (top) Confocal images of M$_2$-Hairpin-EGFP, Mitotracker Deep red staining and DIC (bottom). Scale bar 5 μm. **C**, **left** Mutations of the bicistronic Sirius-H-M$_2$i3(15n/30n)-EGFP plasmids in the region of the M$_2$ i3 loop from nucleotides 1972 to 2001, in yellow are the modified nucleotides. In Sirius-H-M$_2$i3(15n/30n)mut#6-EGFP the four nucleotides in green (AAGA) were reinserted in the stretch of 20 T of Sirius-H-M$_2$i3(15n/30n)mut#5-EGFP. **right** IRES activity, highlighting the efficiency of cap-independent translation for the control Sirius-H-M$_2$i3(15n/30n)-EGFP and the six mutants illustrated. Significance ($p < 0.01$) was calculated in **A** against Sirius-M$_2$i3(417n)-EGFP and in **B** against Sirius-H-M$_2$i3(15n/30n)-EGFP. Error bars are s.e.m. Significance values reported in the graphs were determined by a one-tailed Student t-test, p-values: ** $0.001 < p < 0.01$.

However, when the full i3loop was placed after the hairpin, green fluorescence was fully recovered and the resulting construct (Sirius-H-M_2i3(417n)-EGFP) showed only a slight reduction of the EGFP/Sirius with respect to the value measured for the analogous reporter without hairpin (~90%) (**Fig. 2A**). Finally, when the last portion of the M_2 i3 loop was inserted in the construct after the Hairpin, leading to Sirius-H-M_2i3(15n/30n)-EGFP, ~110 % of the value measured for the Sirius-M_2i3(417n)-EGFP plasmid was recovered (**Fig. 2A**). Notably, when the hairpin was inserted within the i3 loop of the full length M_2-EGFP, upstream of the M368, it was still possible to observe green fluorescence in transfected cells (**Fig. 2B**).

These results confirm that the IRES is located in a region of the i3 loop sequence immediately upstream of the third methionine (M368). This is as also confirmed from ribosome profiling data from human left ventricle tissue, displaying increased p-sites as well as overall ribosomal coverage in correspondence of this region (**Fig. S4B**). Based on this evidence, we mutated the conserved residues highlighted in yellow in **Fig. S4C**, in order to check for their relevance in recruiting the ribosome on the M_2 i3 loop sequence, as shown in **Fig. 2C**. As discussed in detail in the **Supplementary Materials**, we found that the four nucleotides AAGA1090-1093 next to the methionine 368 play a pivotal role in recruiting the ribosome on the M_2 mRNA and in translating the M_2 C-terminal fragment.

Generalization to other members of the GPCR superfamily

In order to determine if the observed behavior is specific to the M_2 receptor, or could highlight a feature common to other members of the GPCR superfamily, we repeated several of our experiments in another GPCR, the related muscarinic M_3 receptor. The M_3 displays an in-frame methionine at the end of the i3loop, in position 481. We then inserted a stop codon in position 273, generating the construct M_3stop273 that still displayed radio-ligand binding and functional activity (**Table S2**). Several other mutants, mimicking the single and double stop mutations performed for the M_2 displayed a behavior consistent with M481 being the start point of a cap-independent translation of the C-terminal also for the M_3 receptor (**Table S2**), thus indicating that IRES-mediated cap-independent translation of the C-terminal domain could be a phenomenon not restricted solely to the M_2 receptor, but may be shared at least by some other members of the GPCR superfamily. The presence of an IRES sequence upstream of M481 further appears to be supported by the observation of GFP fluorescence when 558 nucleotides from the i3 loop of the M_3 receptor (from nucleotide 880 to 1437) were inserted in the same Sirius-EGFP reporter construct as used for the M_2 (**Fig. S5B**). This appears to be indeed supported by the sequence-alignment of the five muscarinic receptors. The alignment of the last portion of the M_2 i3 loop (from nucleotide 1072 to 1104) with regions adjacent to the second (M_1), fourth (M_3 and M_4) and

first (M_5) in-frame Methionines, clearly shows that some nucleotide residues are conserved (<mark>Fig. S4C</mark>). Overall these data underline that cap-independent translation could be a more general feature of GPCRs mRNAs.

Mitochondrial localization of the M_2 C-terminal fragment generated by cap-dependent expression under the control of a viral promoter

After having established that the expression of the M_2 C-terminal domain is regulated by an IRES and that its translation starts with M368, we went on to study the role it might play in the cell. For this purpose, we fused M_2tail(368-466) to EGFP in pcDNA3, that is under the control of a strong CMV promoter, and we expressed it in HEK293 cells in order to study its subcellular localization. Confocal microscopy demonstrated a striking mitochondrial localization of the M_2tail(368-466)-EGFP construct (<mark>Fig. 3A</mark>).

We confirmed this observation by co-staining HEK293 cells with a conventional mitochondrial stain and by co-expressing M_2tail(368-466)-EGFP with Tom20-mCherry, a marker for the outer mitochondrial membrane (<mark>Fig. S6A</mark>). The observed mitochondrial localization of M_2tail(368-466)-EGFP appeared in stark contrast to the canonical cellular localization of the M_2 receptor, as shown in <mark>Fig. 3b</mark> where the M_2 muscarinic receptor was C-terminally labeled with the red fluorescent protein mRuby2. In line with the radioligand binding data shown in <mark>Table 1</mark>, very little expression of M_2tail(368-466)-EGFP was observed on the plasma membrane, as we confirmed by fluorescence correlation spectroscopy (FCS) [14](<mark>Figs S6B-D</mark>). By contrast, when the M_2tail(368-466)-EGFP was co-expressed with M_2stop228, an increased localization could be seen on the cell membrane, suggesting a chaperon activity of the M_2 N-terminal domain (M_2trunk) (<mark>Figs S6E-G</mark>). Likewise, constructs such as M_2tail(368-466)-mRuby2 (a red monomeric fluorescent protein) and M_2tail(368-466)-tdTomato (a tandem-dimer red fluorescent protein) also displayed clear mitochondrial localization (<mark>Figs S6H-I</mark>), ruling out bias due to the fluorophore used. Mitochondrial import of the C-terminal fragment of the M_2 receptor was further verified by immunoblotting of protein extracts from purified mitochondria of HEK293 cells transfected with myc-tagged M_2 receptor, M_2tail(368-466) and M_2M368A, a mutant where the key methionine for the cap-independent translation of the C-terminus is changed to an Alanine. <mark>Fig. 3C</mark> shows that in purified mitochondria (unlike in the cytosolic and microsomal supernatants originating from the purification process, <mark>Fig. S6J</mark>), only the protein bands associated to the M_2 C-terminal fragment were visible. Notably, no mitochondrial import was visible for the M_2M368A, confirming that the third in-frame methionine (M368) in the i3 loop of the M_2 receptor is necessary for the C-terminal fragment production and its localization to the mitochondria. The slight offset between the lower bands in the M_2 wt and M_2tail(368-466) lanes matches exactly the different length of the linker used to fuse the myc-tag in the two constructs.

Fig. 3. M$_2$tail(368-466) predominantly localizes to the cell mitochondria

A, Colocalization of M$_2$tail(368-466)-EGFP (green) together with the mitochondrial targeted dye Mitotracker (magenta) in HEK293 cells. Separate spectral channels are displayed below. **B**, Co-expression of M$_2$tail(368-466)-EGFP (green) and wild type M$_2$-mRuby2 (red) in HEK293 cells. Separate spectral channels are displayed below. **C**, Western blot (10% TRIS-Glycine PAA gel) of mitochondria purified from Hek293 cells transfected with the M$_2$-myc, M$_2$tail(368-466)-myc, M$_2$M368A-myc, together with an untransfected control are blotted using an anti-myc antibody. Loading control was verified by immunoblotting COXIV. **D**, Cellular localization of the construct M$_2$-mRuby2-STOP-M$_2$-i3-tail-EGFP in HEK293 cells. Expression of the M$_2$tail(368-466)-EGFP (green) after the STOP codon is an indicator of the endogenous IRES activity. While M$_2$-mRuby2 localizes correctly and predominantly to the cell plasma membrane, M$_2$tail(368-466)-EGFP localizes in the cytosol as well as, strongly, to the cell mitochondria. Separate spectral channels are displayed below. **E**, M$_2$-fr.sh.-EGFP (green) expressed in conjunction with Mitotracker (magenta). The EGFP is expressed despite the creation of an in-frame stop after M368, confirming endogenous cap-independent translation of the M$_2$tail(368-466). Arrows indicate areas of colocalization with the mitochondria. Separate spectral channels are displayed below.

Localization of the M$_2$ C-terminal fragment generated by cap-independent expression under the control of the i3loop IRES

We then sought to express the M$_2$tail under the control of the i3loop IRES rather than a viral promoter, to further exclude the possibility that the observed mitochondrial localization of the M$_2$ carboxyl terminal fragment were due to overexpression of the construct. HEK293 cells were transfected with the construct M$_2$-mRuby2-STOP-M$_2$-i3-tail-EGFP which contains the full wild type M$_2$ receptor, labeled at the C-terminal with mRuby2, additionally fused to the C-terminal part of the M$_2$ receptor starting at nucleotide 685 of the i3 loop, following the stop codon of mRuby2, and ending with a C-terminal EGFP sequence (Fig. S3). Through this experimental strategy, the expression of M$_2$tail(368-466)-EGFP is regulated by the IRES, and the mechanism of C-terminal fragment production is therefore close to physiological. The expression of EGFP confirmed that the *in vivo* production of the M$_2$tail(368-466) is regulated by the IRES within the i3 loop. In

addition, the processed receptor fragment was eventually observed to be predominantly confined to the cytosol and the mitochondrial network (Fig. 3D).

Notably, when the M_2-EGFP fusion protein was expressed and the cells were co-stained with Mitotracker, a faint EGFP signal was also observed in the mitochondrial regions (Fig. S7A), although the expression of the full-length receptor masks this effect and makes quantification difficult. On the other hand, Fig. 3E shows that, when the M_2fr.sh-EGFP mutant was expressed in HEK293 cells, the IRES-generated C-terminal fragment of the M_2 receptor localized clearly to the mitochondria. As mentioned above, in this construct the masking effect of the EGFP fused to the full-length receptor is removed (see also Fig. S3). Similar mitochondrial localization was also observed when we transfected HEK293 cells with M_2fr.sh-mRuby2 (Fig. S7B), ruling out possible effects due to the specific fluorophore used as a tag. We obtained comparable findings also in COS-7 cells (Fig. S7C).

Mechanism M_2tail import into the mitochondria and localization within the organelle

To gain further insight into the dynamics and mechanisms of the M_2 C-terminal fragment import within the mitochondria, we first monitored the kinetics of colocalization between mitochondria and EGFP in cells transfected with M_2tail(368-466)-EGFP. In cells observed for several hours, starting at 5h after transfection (Fig. 4A), the construct was found initially expressed in the perinuclear region, likely the endoplasmic reticulum, indicating that at least the initial step of subcellular trafficking of the M_2tail(368-466)-EGFP is the same as for the full length receptor [15]. At this time (5h post-transfection), almost no colocalization with the Mitotracker-stained mitochondria was visible. However, at 6h post-transfection, the colocalization level increased significantly, and M_2tail(368-466)-EGFP was either found in the cytosol or in the mitochondria. At 9h post-transfection the colocalization at the mitochondria was well visible (see Materials and Methods) and we could observe that almost half of the mitochondrial import occurred within 6h post-transfection (Fig. 4B).

This value is qualitatively in agreement with what we observed using an in-vitro assay to determine the import of the M_2 carboxyl terminal into isolated mitochondria (see Materials and Methods). To this end, we used mitochondria form *S. cerevisiae*, which is a robust and well-studied system for protein import assays. Fig. 4C shows the autoradiography experiment of isolated, intact, mitochondria incubated with 35S-labelled full length muscarinic M_2 receptor or M_2tail(368-466), made using a reticulocyte in vitro translation system. The data show that the full-length receptor was not imported into mitochondria. On the other hand, the M_2tail(368-466) displays a clear mitochondrial import already after 5 minutes and increasing over time, up to 30 minutes. Interestingly, this import appears to be independent upon the transporters of the outer membrane Tom5, Tom6, Tom7 and Tom70 (Fig. S8A), whereas the import of an inner membrane protein, the ADP/ATP carrier (AAC), used here as a control, depends on Tom70.

Upon extraction with carbonate most of M_2tail(368-466) was found in the pellet, and therefore
presumably integrated in mitochondrial membranes. Furthermore *Blue native* analysis suggests
that the mitochondrially imported M_2tail(368-466) is not a monomer but rather it is part of a larger
complex of approximately 180 kDa (Fig. S8A).

All together these data indicate that the import of the M_2tail(368-466) leads to its localization in
one of the mitochondrial membranes, to form a complex with other native mitochondrial proteins.

To further discriminate if the construct is on the outer or inner mitochondrial membrane, we set
up a fluorescence assay based upon GFP fluorescence complementation [16]. HEK293 cells
were first transfected with two constructs: M_2tail(368-466)-GFP11, in which a short 16 amino
acids peptide of GFP was fused at the C-terminus of M_2tail(368-466), and mito-GFP1-10, in
which the COX8A was N-terminally fused to the other GFP fragment containing the remaining 10
beta strands (See Materials and Methods). Figure S8B shows the successful recombination of
the two within the mitochondrial matrix, and associated controls (Figure S8C-G). We then
applied this approach to the full length M_2 receptor, by generating the M_2-GFP11 construct and by
transfecting it in HEK293 cells. Fig. 4D shows a green signal from the recombined GFP visible in
the mitochondria. All together these data suggest that the M_2 C-terminal fragment is localized in
the inner mitochondrial membrane, with its own C-terminal domain facing the matrix. This
observation was also supported by the lack of complementation of M_2tail-GFP11 with GFP1-10
when this last fragment was N-terminally fused to the targeting sequence of SMAC, a protein
expressed in the mitochondrial intermembrane space [17](Fig. S8E). This interpretation is also
finally confirmed by transmission electron microscopy imaging of thin sections from cells
expressing the M_2stop228 C-terminally labeled using a myc-tag after immune-gold labelling (Fig.
4E). As shown in Figs 4F-G, gold particles were observed only in mitochondria, and at this stage
they appeared localized either to the mitochondrial matrix or to the lumen of the cristae (Figs 4H-
K).

Cell stress upregulates IRES-mediated M_2tail expression and drives mitochondrial
localization

Having established that M_2tail localizes in the inner mitochondrial membrane we then moved to
look at the effect of stress on the level of expression of the C-terminal fragment. Stressed cells
are known to attenuate their translational activity via phosphorylation of the initiation factor eIF2α.
However, certain mRNAs may be insensitive to eIF2α phosphorylation level if their translation is
regulated by IRES, and under these conditions their expression is bound to increase rather than
to attenuate [18,19].

Therefore, we transfected COS-7 cells with the Sirius-M_2i3(417n)-EGFP reporter construct and
stressed by starvation for three hours by replacing the culture medium with HBSS or phosphate

saline buffer (PBS). Notably, PBS has a poorer nutrient content than the HBSS media, thus causing a more pronounced starvation. During this time interval, the green fluorescence was seen to gradually increase up to 1.4 times the basal level observed in HBSS-starved cells and 3.5 times for PBS-starved cells (Fig. 5A), confirming that the C-terminal fragment expression is upregulated in response to stress conditions such as cell starvation. The same experiment was reproduced in single cells, displaying an increase of the average EGFP/Sirius ratio (slope), and thus of IRES-mediated translation, upon HBSS starvation in HEK293 cells (Fig. 5B). We further used the GFP fluorescence complementation assay to emphasize the importance of stress in the pattern of expression of the M_2 gene. HEK293 cells were co-transfected with M_2-GFP11 and Mito-GFP1-10. HEK293 cells grown in culture medium predominantly expressed the full M_2 receptor protein (Fig. 5C).

Fig. 4. Mitochondrial import and organelle localization of M_2tail

A, Time course of M$_2$tail(368-466)-EGFP import in the mitochondria of a HEK293 cell over a 4h interval, starting 5h after cell transfection, as indicated. Panels, from top to bottom show M$_2$tail(368-466)-EGFP (green), Mitotracker (magenta) and overlay displaying increasing colocalization over time. **B,** Quantification of the colocalization of the M$_2$tail(368-466)-EGFP in the mitochondria by calculating the Manders Coefficients (n=7 cells, from 3 independent transfections, error bars are s.e.m.) (**Materials and Methods**). **C**, Import of protein M$_2$tail(368-466) in isolated yeast mitochondria: ^{35}S-labelled and translated in-vitro full length muscarinic M$_2$ receptor or M$_2$tail(368-466) expressing 5 C-terminal Methionines were incubated with isolated yeast mitochondria for various timepoints in the presence of the inner membrane potential (**Materials and Methods**). To test for equal mitochondrial loading, αPorin was probed via western blot. The kinetics pattern of M$_2$tail is what it is expected for a protein imported in mitochondria: the appearance of the cleaved M$_2$-tail is indicative of a post translational modifications of the fragment once imported in mitochondria. **D,** Detail of a HEK293 cell expressing the M$_2$-GFP11 construct together with mito-GFP1-10. The M$_2$-GFP11 was sequentially transfected 24 h after the mito-GFP1-10 (**Materials and Methods**) Magenta overlay indicates mitotracker. White signal in the overlay indicates M$_2$tail(368-466) trafficking to the organelle. All confocal micrographs were obtained by sequential imaging using 488 nm excitation and detection in the 520-600 nm (EGFP), 561 nm excitation and detection in the 580-650 nm (mRuby2) and at 633 nm and detected in the 650-700 nm range (Mitotracker) using HyD detectors in Photon Counting Mode. Scale bars are 10 µm throughout. **E,** TEM micrograph of a COS-7 cell showing a large nucleus, several mitochondria and vacuoles nearby. Bar: 5 µm. **F-G**, High magnification of portion of the cell cytoplasm showing a few gold particles confined to the mitochondria. Bar: 500 nm. **H-K,** Gold particles appear associated with the mitochondrial cristae, highlighted by the dotted line. A total number of 236 gold particles in 40 mitochondria from 19 sections were detected.

The green fluorescence detected at the plasma membrane is explained with the chaperoning effect of M$_2$-GFP11 on Mito-GFP1-10 before its sorting to the mitochondria. In sharp contrast, when HEK293 cells were first transfected with Mito-GFP1-10 and, 24h later, transfected with M$_2$-GFP11 followed by a 6 h starvation in HBSS, the pattern of expression changed drastically: IRES mediated translation of the M$_2$tail(368-466) fragment led to a predominantly mitochondrial green fluorescence signal of the complemented GFP (**Fig. 5D**). These results are in line with our observation that the IRES mediated C-terminal fragment localization to the mitochondria in cells expressing the construct M$_2$-mRuby2-STOP-M$_2$-i3-tail-EGFP was enhanced upon cell starvation, and drastically reduced upon mutation of the M368 (**Figure S9** and **Supplementary Materials**).

Fig. 5 Effect of integrated stress response on IRES activity and M$_2$tail(368-466) localization

A, 96-well plate reader measurement of the increase of EGFP fluorescence relative to Sirius in COS-7

cells transfected with the bicistronic Sirius-M$_2$i3(417n)-EGFP plasmid and starved for three hours

respectively in PBS (left panel) (n= 3 transfections, 16 wells) and HBSS (right panel) (n=1 transfection, 16

wells). Significance values reported in the graphs were determined by a one-tailed Student t-test, p-values:

**** $p<0.0001$; ** $0.001<p<0.01$. **B,** Single cell measurement of EGFP vs SIRIUS expression in HEK293

cells, in basal conditions and upon starvation. Lines are linear fits (note log-log scale). **C**, HEK293 cells

were sequentially transfected with pMito-GFP1-10 (24h) and M$_2$-GFP11 (8h, of which 6h under starvation

conditions) and then grown in full culture medium or **D,** starved for 6 h after transfection of M$_2$-GFP11.

Shown are EGFP fluorescence (left, with DIC inset), Mitotracker (center) and overlay (right). Scale bars are

10 μm.

Functional role of the C-terminal fragment in the mitochondria

The next step of our investigation concerned the obvious question about the functional effects

that the observed localization of the M$_2$ C-terminal fragment might have on the mitochondrial

activity, and -more generally- on the energy metabolism of the cell. To investigate this aspect, we measured the extent of oxygen consumption in COS-7 cells expressing M_2-EGFP, M_2tail(368-466)-EGFP or the sole EGFP as a control. Initially, we normalized the expression of the three proteins to the amount of EGFP expressed in the cells, then the metabolic functions were assayed on a Seahorse instrument using cells transfected with normalized amounts of plasmids (Fig. 6A).

Our data did not show a significant difference in total oxygen consumption capacity between cells transfected with cytosolic EGFP and the untransfected control. However, a significant decrease of oxygen consumption was found in cells transfected with M_2-EGFP and even more so with M_2tail(368-466)-EGFP (Fig. 6A). Notably, the basal oxygen consumption rate (OCR) in starvation-stress conditions was profoundly decreased in M_2tail(368-466) cells. This occurred to a smaller extent also in M_2 cells compared to EGFP-transfected and empty control cells (see also the time-course in Fig. S10A). These results are in line with what was observed using a Clarke-type oxygen electrode (Supplementary Materials and Figs S10B-C).

Notably, there was no difference in the number of apoptotic cells among the different cell groups as shown by the caspase-3 activation assays of apoptosis (17) (Fig. S10D-E). To test whether the M_2tail effects on mitochondrial oxygen consumption could alter the production of Reactive Oxygen Species (ROS), the measurements were combined in a single cell assay. Remarkably, ROS production (assayed using a fluorescence readout kit, where the fluorochrome becomes activated in contact with oxidized species) was clearly reduced within and in proximity of mitochondria of cells expressing a high level of the M_2tail(368-466) (Fig. 6B). By regrouping individual mitochondria in classes according to the level of the ROS marker expression, an actual inverse correlation between the amount of receptor fragment (Green Fluorescence) and the production of ROS (Red Fluorescence) was revealed.

Cell growth was also found to be affected depending on which of the constructs was transfected. Notably, transfection of HEK293 cells with full length M_2 receptor reduced significantly cell growth compared to the control, whereas cells transfected with the mutant M_2M368A, that do not express the M_2 C-terminal fragment (Fig. 6C), displayed a significantly higher growth. All cells were incubated with saturating concentration of the M_2 antagonist atropine throughout, in order to exclude effects on cell growth and proliferation due to the canonical receptor signaling.

Fig. 6. Effects of M₂tail expression, or absence thereof, on mitochondrial function and cell metabolism in transfected HEK293 cells.

A, Basal oxygen consumption upon HEK293 cell stress -following starvation- measured by a Seahorse assay at a constant temperature of 37 °C. Significance values reported in the graphs were determined by a one-tailed Student t-test, p-values: **** $p<0.0001$; *** $0.0001<p<0.001$;* $0.01<p<0.05$. **B**, Localization of M₂tail(368-466)-EGFP in HEK293 (green) cells stained with a marker for reactive oxygen species (Cell Rox Deep Red, ThermoFisher). Relationship between green (M₂tail abundance) vs red (amount of ROS

produced) fluorescence intensities measured from regions of the mitochondrial network. The data were
binned in four groups as a function of the amplitude of Cell Rox Deep Red signal (0-2.5, 2.5-5. 5-7.5 and
7.5-10 intensity counts). **C**, Cell viability assay performed on HEK293 cells transfected with M_2 and
M_2M368A, compared to an untransfected control. Cells were counted 24h and 72h after transfection in T-
25 flasks, exposed throughout to 100 nM atropine. The experiments were performed in two biological
replicates, each with two technical replicates. Transfection efficiency (to the right of the chart) was
monitored thanks to the mRuby2 fluorescent reporter within a bicistronic cassette within each vector (see
Materials and Methods). Significance values reported in the graphs were determined by a one-tailed
Student t-test, p-values: ** $0.001 < p < 0.01$; * $0.01 < p < 0.05$. **D**, FRET efficiency plot of the interaction
between M_2tail(368-466)-mTurquoise2 and F_0F_1ATP Synthetase (CV-mCitrine), COX8a(pMito)-mCitrine
and SMAC-mCitrine, as determined by acceptor photobleaching measurement in a confocal microscope.
Each dot represents a ROI within a cell. At least 3 cells from two separate transfections were imaged for
each pair.
519
These functional data are further supported by Fluorescence Resonance Energy Transfer
(FRET) data aimed at investigating the molecular interactions of the mTurquoise2 labeled
M_2tail(368-466) with mitochondrial proteins. Based on the oxygen consumption data (**Fig. 6A** and
Fig. S10A) we chose as primary target the F_0F_1 ATP Synthetase complex, also indicated as
complex V (CV), and we labeled the γ subunit with mCitrine, a fluorescent acceptor. The γ
subunit is located towards the mitochondrial matrix. We further monitored interactions with an
intermembrane space protein (SMAC) and a protein localized to the mitochondrial matrix
(COX8a), both fused to the mCitrine fluorescent acceptor. As clearly illustrated in **Fig. 6D**, FRET
efficiency, an indicator of molecular interaction, is significantly larger for M_2tail(368-466)-CV pair
than both the intermembrane space SMAC and the matrix COX8a. The latter result is particularly
relevant, as it indicates a specific interaction of M_2tail(368-466) with CV.
531

Relevance of C-terminal expression and mitochondrial translocation in a physiological, endogenous setting

The results provided in the previous sections convincingly demonstrate the cap-independent
translation of the M_2 C-terminal fragment, its upregulation and mitochondrial localization upon
stress response, and the subsequent effect in regulating oxidative phosphorylation by a direct
interaction with complex CV of the respiratory chain. To test the physiological relevance of these
findings we first prepared protein extracts from heart, brain and spleen tissues from a line of
knock-in mice (see **Materials and Methods**), designed to have expression of a Chrm2/tdtomato
fusion protein driven by the endogenous M_2 muscarinic acetylcholine receptor
promoter/enhancer sequences. The heart and brain are known to express sizable levels of the
M_2 receptor, whereas the spleen, where no expression of the M_2 receptor is reported in the
proteomic databases, was used as a negative control.

Fig. 7. IRES-mediated M₂tail(368-466) expression occurs in endogenous, physiological settings and its suppression affects cell metabolism

A, Western blot from tissue preparations originating from the indicated organs of M₂-tdtomato knock-in mice (HT) and wild-type animals (WT). Immunodetection was performed using an anti-tdtomato primary antibody. Loading control normalization was performed using GAPDH in the heart, and ‑actin in the brain and spleen. **B**, Western blot (8% TRIS-Glycine PAA gel) of mitochondria purified from tissue preparations originating from the indicated organs of M₂-tdtomato knock-in mice (HT) and wild-type animals (WT), detected using an anti-tdtomato antibody. **C,** Confocal micrographs of wt hiPSCs (left) and Crispr/Cas9 clone harboring the M368A mutation within the M₂ gene (right), stained with DAPI (cyan) and Mitotracker (red). **D**, Cell proliferation assay performed for wt hiPSCs and M368A hiPSCs grown in T25 flasks., exposed to Atropine. Cells were counted 24h and 72h after seeding. Significance values reported in the graphs were determined by a one-tailed Student t-test, p-values: ** 0.001<p<0.01. **E**, Extracellular oxygen consumption assay, performed on wt and M368A hiPSCs, prior to and after 7 days undirected

differentiation under Atropine exposure. The lifetime of a commercial dye increases as oxygen in the medium is progressively consumed by the cells. The data displayed are collected after 60 minutes incubation. Significance values reported in the graphs were determined by a one-tailed Student t-test, p-values: *** 0.0001<p<0.001; ** 0.001<p<0.01.

The use of a transgene tag was necessary, since antibody against the M_2 receptor are notoriously ineffective, in particular when targeted at the C-terminal portion of the receptor. Instead, the tdtomato fluorescent protein can be targeted by a specific polyclonal antibody, as shown by the Western blot displayed in Fig. 7A. As expected, the full-length receptor-tdtomato fusion was observed at ~130 kDa (~75 kDa contributed by the receptor and ~55 kDa contributed by tdtomato). Notably we observed again (both in the heart and brain, but not in the spleen) a specific band doublet, this time at ~75 kDa, which corresponds to the sum of the molecular weights of the tail fragment added to the molecular weight of tdtomato. When mitochondria were purified from the heart and spleen of the Chrm2/tdtomato mice Fig. 7B the band at ~75 kDa was substantially enriched, matching the expected molecular weight of the M_2tail. These data strongly suggest that the M_2tail is expressed endogenously and it is sorted to the mitochondria.

We finally asked the question of what would be the effect of abolishing the translocation of the M_2 C-terminal fragment to the mitochondria, in a physiological and entirely label-free setting. Notably, human induced pluripotent stem cells (hiPSC) combined with genome editing approach are state-of-the art tools to investigate the effect of gene mutations in physiological and label-free setting. Therefore, we incorporated by Crispr/Cas9 the homozygous M368A mutation in hiPSCs (Materials and Methods). Based on what we reported so far, in particular our data of (Fig. 3C), this mutation should be sufficient to significantly impair the M_2tail production and mitochondrial translocation. Wild type (wt) and M368A hiPSCs were cultured in pluripotency maintaining medium, and subsequently undirectedly differentiated upon exposure for seven days to DMEM-F12 medium, as described in the Materials and Methods. Fig. 7C shows confocal micrographs for wild type and M368A hiPSCs cultured in E8 medium respectively, with the M368A hiPSCs displaying a mildly more spread-out phenotype. We shall note here that the M_2 receptor has been found to be expressed in PSCs lines, and it has been shown to play a role in stem cell differentiation [21,22](21,22). In this context, however, we were not interested in the canonical signaling of the receptor, and all experiments were performed with a saturating concentration of atropine, to block any possible effect of the canonical signaling of the M_2 receptor. We then evaluated cell proliferation, observing a significant increase in the proliferation of the M368A clone 72h after seeding Fig. 7D. The observation that cells harboring the mutation that suppresses M_2tail display an increased proliferation, suggests that the same effect is at play here, as observed in the overexpression system (Fig. 6C). These results are supported by extracellular oxygen consumption assays performed, for both hiPSCs and differentiated hiPSCs

(**Fig. 7E**): the M368A cells indeed display an accelerated oxygen consumption rate, and that this effect increases as the cells progressively differentiate.

For comparison, the assay was also carried on HEK293 overexpressing the wild type M_2 receptor and the mutant M_2M368A (**Fig. S10F**). We shall further note here that signaling properties of M_2 M368A with respect to the wild type M_2 receptor are comparable, as displayed by pERK immunoblotting as well as FRET G_i activation assays performed in cells overexpressing either the wild type M_2 or the M_2 M368A (**Fig. S10 G,H**).

Discussion

Over the last 10 years, growing evidence has accumulated indicating that GPCRs can be functional and signal also at cellular locations other than the plasma membrane. These locations can be associated to their canonical trafficking pathways (such as sorting vesicles and endosomes) [23], but also to non-canonical locations, such as the nucleus, the nuclear membrane, the Golgi apparatus and the mitochondria (reviewed in Jong et al. BPJ 2018 [24]). Some of these locations are surprising, since many ligands are non-membrane permeable, and it is still debated how they can reach their target receptors at such intracellular locations.

Interestingly, the specific localization of these receptors appears to go hand in hand with a specific function. In these cases, activation/inhibition of mitochondrially localized GPCRs have been shown to specifically affect the organelle function, ranging from oxygen consumption rates to Ca^{2+} handling and even to apoptosis. Notably Benárd et al [25] showed that the CB_1R on the outer mitochondrial membranes affects neuronal energy metabolism; Suofu et al. [7] demonstrated that activation of the MT_1R by a mitochondrial melatonin pool can affect cytochrome C release. This is particularly interesting, since cytosolic cAMP does not appear to be able to activate mitochondrial PKA, and hence a mitochondrially localized GPCR signalosome appears to be required to enable such downstream signaling cascade [26].

However, other non-canonical mechanisms are also possible for the regulation of mitochondrial function. It is well known that GPCRs are subject to interindividual variability, arising, for example, from mRNA alternative splicing (i.e. the process which allows the recombination of exons) which leads to different isoforms which may present distinct signaling and localization properties [27]. Receptor variants with different functional and localization properties may also occur due to other processes. For example, our comprehension of eukaryotic translational regulation is facing a rapid expansion, as exemplified by the observation that small open reading frames (smORFs), also within lncRNAs, are an important class of genes, oftentimes with mitochondrial localization [28].

In this study, for the first time, we provide evidence that an IRES sequence drives expression of a membrane receptor fragment and acts as a regulator of its expression levels and function. IRES sequences are distinct regions of the mRNAs that are capable of recruiting ribosomes in the proximity of an internal initiation codon and translate transcripts in a cap-independent manner [11,12,29–31]. Here, we show that an IRES sequence is located within the third intracellular loop of the M_2 muscarinic receptor, where it plays a crucial role in regulating the expression of its carboxy-terminal domain, which we identified as M_2tail(368-466).

We assessed the IRES-driven expression of the C-terminal fragment through a broad set of constructs, meant to exclude alternative synthesis mechanisms such as stop-codon read-through, and a diversified palette of techniques. Radioligand binding data (Table 1) pointed to the necessity of the C-terminal fragment synthesis, in order to explain the observed downstream signaling in mutants carrying a stop codon in the initial portion of the i3loop of the M_2 receptor, upstream of M368. Fluorescence microscopy allowed us visualizing the production of the EGFP-tagged C-terminal fragment in these receptor mutants (Fig. 1).

Mutagenesis of key residues within the i3 loop of the M_2 receptor allowed us to pinpoint the 3rd in-frame methionine as the most likely candidate for the beginning of the cap-independent translation (Fig. 2). This observation appears to be supported by the observation of a peak in ribosome coverage around the M368 seen in ribosome profiling databases (as reviewed in Ingolia et al. [33]), and particularly prominent in those from left ventricle cardiomyocytes, where M_2 receptor expression is known to be abundant [28].

To our surprise, in contrast to the plasma membrane localization of the wild type, full length M_2 receptor, the C-terminal fragment was localized mostly in mitochondria. This localization occurred markedly when overexpressing the C-terminal fragment M_2tail(368-466), but could be also clearly detected when the C-terminal fragment was expressed under the control of the IRES sequence of the full-length receptor (Fig. 3).

We devised several strategies to detect the localization of the IRES-driven fragment in the mitochondria by avoiding the cell-wide background signal from the full-length receptor. Fig. 3D,E show two prominent examples, namely relying on spectrally unmixing the two components, or by sending out of frame the EGFP at the C-terminus of the full-length receptor. By studying mitochondria purified from HEK293 cells we could confirm that the mutation of the M368, which we identified as the starting point of the IRES-mediated translation, is sufficient to suppress any mitochondrial targeting of the M_2 C-terminal fragment, and most likely, its synthesis altogether (Fig. 3C).

We then addressed the mechanism of mitochondrial import of the C-terminal fragment and its localization within the organelle (Fig.4). Our data, originating both from live cell import kinetics as well as from in-vitro import into isolated mitochondria (Fig.4A,B), support the notion that the fragment becomes rapidly imported into mitochondria following translation in the cytosol. Once inside the mitochondria, it undergoes processing leading to the appearance of a lower molecular weight band, observed both in Western blots from whole cell and mice tissue protein extracts, as well as from purified mitochondria. The fragment is found predominantly in a carbonate-insoluble fraction, i.e. a membrane, and it appears to form a high molecular weight complex of ~180 kDa with other mitochondrial proteins (Fig.4C).

Fluorescence complementation assays (Fig. 4D) and electron microscopy (Fig. 4E-H) suggest that the M_2 C-terminal fragment predominantly resides in the inner mitochondrial membrane, with its own C-terminal domain exposed towards the matrix. Future work should clarify the precise import pathway.

Under integrated stress response, namely aminoacid starvation, IRES-mediated production of the C-terminal fragment was increased (Fig. 5A-B), with a marked increase in mitochondrial localization (Fig. 5C). Most importantly, our data show that mitochondrial oxygen consumption was reduced by the M_2tail(368-466) (Fig. 6A,B), and also cell proliferation was affected (Fig. 6C). On these bases, the M_2tail might be needed in mitochondria as a means to reduce mitochondrial basal oxygen consumption and ATP-linked respiration. An important consequence of reduced respiration is the lower level of ROS produced in mitochondria, as we observed in agreement with this hypothesis Fig. 6B.

This observation was confirmed in physiological conditions by Western blots in a knock-in mice carrying a tag at the C-terminus of the endogenous M_2 receptor. These could show mitochondrial enrichment of a band corresponding to the molecular weight of the C-terminal domain (Fig. 7A). **Moreover,** the physiological relevance of our observations was tested in an entirely label-free assay, by Crispr/Cas9 editing of pluripotent stem cells in order to suppress the synthesis of the M_2 C-terminal fragment. By mutating the IRES starting methionine M368, we observed an increase of cell proliferation of this clone compared to its wild type counterpart, together with an increase of oxygen consumption rates (Fig. 7B-D). Overall this corroborates our hypothesis that the M_2 C-terminal fragment reduces oxygen consumption upon reaching the mitochondria. The effect was further amplified when the cells were undirectedly differentiated, and it was consistent with the shift of the cell phenotype towards oxidative phosphorylation [34].

The mechanism by which the M_2tail(368-466) may reduce oxygen consumption in mitochondria and its meaning needs to be further explored. Our FRET data (**Fig. 6D**) indicate clearly that the M_2 C-terminal fragment has a specific interaction with the F_0F_1 ATP-synthase complex, supporting the concept that it inhibits the activity of this enzyme, thus leading to the observed reduction in oxygen consumption rates. Future work should clarify the precise import pathway and the inhibitory effect of M_2tail(368-466) on F_0F_1 ATP-synthase. Nonetheless, since ROS production is implicated in the pathogenesis of several diseases [35], we would propose the hypothesis that increased localization of the M_2 C-terminal fragment in mitochondria, as it occurs under integrated stress response, might have protective effects (**Fig. 8**).

Fig. 8. Schematic representation of the wild type M_2 receptor gene products, their destination in the cell and their function.

The M_2 receptor gene gives rise to one transcript but two different translation outputs: when translation starts from the canonical cap site the wild type M2 receptor is generated (full length M_2 receptor). When translation is driven by the internal ribosome entry site (IRES) the C-terminal fragment is generated (M_2 tail). The two proteins are then sorted in two different cell compartments where they play completely different roles: the M_2 receptor signals canonically at the plasma membrane as a G protein-coupled receptor. The M_2tail is instead sorted to the mitochondria.

In normal conditions the activity of the IRES is low, and only a small amount of M_2tail is produced in the cell (left). However, upon cell stress conditions (right), M_2tail IRES-mediated translation is upregulated and M_2tail in the mitochondria interacts with the ATP synthase and reduces oxidative phosphorylation and ROS formation. Created with BioRender.

In broader terms, protection of mitochondria by unconventional import mechanisms to allow localization within the organelle of specific proteins has been previously observed for cells that

730 undergo stress conditions (36) [36]. The mechanism of IRES-dependent modulation of
731 translation and consequent mitochondrial targeting of the translation product of a GPCR may
732 provide another means, completely unrelated to the canonical signaling function of this class of
733 membrane receptors, of a generic cellular response to overcome different types of stress by
734 modulating the mitochondrial proteome and adjusting the mitochondrial bioenergetic state.
735

Materials and Methods

Generation of mutant muscarinic M$_2$ receptors for radioligand binding studies, Western blotting, immunohistochemistry and fluorescence microscopy

Multiple receptor mutants were constructed starting from the human M$_2$ muscarinic receptor expressed in the pcD vector [37]. The constructs are sketched in **Fig. S3** and their cloning is described in detail in the Supplementary Methods.

Reverse transcription

Reverse transcription was performed using a 24-nucleotide antisense oligo (TTACCTTGTAGCGCCTATGTTCTT) directed against the last 24 nucleotides at the 3' end of the M$_2$ open reading frame. cDNA amplification was carried out with the same antisense oligo used for reverse transcription, together with a 24 nucleotides sense oligo (ATGAATAACTCAACAAACTCCTCT) directed against the first 24 nucleotides at the 5' end of the M$_2$ open reading frame.

Cell culture and transfection

COS-7 (African green monkey kidney fibroblast-like SV40 transformed cell line) and CHO (Chinese Hamster Ovary) cells (from American Type Culture Collection) were grown at 5% CO$_2$ and 37 °C, in high glucose Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS), 2 mM L-glutamine, 50 IU/ml penicillin, and 50 mg/ml streptomycin, and non-essential amino acids. HeLa cells (from American Type Culture Collection) were cultured in the same medium, but with the double amount of L-glutamine. SHSY-5Y cells were cultured in Roswell Park Memorial Institute (RPMI) medium supplemented with 10% FBS, 2 mM L-glutamine, 50 IU/ml penicillin, and 50 mg/ml streptomycin. Media and reagents used for cell cultures were from Euroclone, PAN Biotech, Biochrome and Sigma-Aldrich. Cell transfection was performed using FuGene® transfection reagents (Promega) as recommended by the manufacturer. Eighty percent confluent cells were washed twice with free-serum medium and the mix of FuGene and plasmid DNA was added. In every experimental condition 6-8 μg of plasmid DNA per Petri (10 cm^2) were used. Cells were grown for 48 h post-transfection before use. In co-transfection experiments, half amount of each plasmid DNA was used. All microscopy experiments were performed with transiently transfected HEK293 (Human Embryonic Kidney, ECACC 96121229 from Sigma-Aldrich Chemie GmbH) and COS-7 cells. HEK293 cells were cultured in DMEM (Dulbecco's modified Eagle's medium PAN biotech, Aidenbach, Germany), supplemented with 4.5 g/L Glucose, 2 mM L-Glutamine, 10% FCS,

100 units/mL penicillin and 0.1 mg/mL streptomycin and maintained at 37 °C and 5% CO_2. Cells cultured in 15-cm dishes were split at a 1:36 ratio into 6-well plates containing poly-D-lysine (PDL)-coated 24 mm glass coverslips. Cells were transfected using Effectene transfection reagent (QIAGEN) according to the manufacturer's instructions. Cells seeded on PDL-coated coverslips in 6-well plates were transfected 16 hours after seeding with 0.6 µg plasmid/well.

Fluorescence microscopy

Fluorescence microscopy experiments were performed either on a Leica SP5 (or SP8) Confocal Microscopes, using HyD photon counting detectors and an Argon/Ion laser (or white light laser) source for 488 nm/633 nm excitation and a solid-state diode laser (or white light laser) for 561 nm excitation. A solid-state diode laser was used for 405 nm excitation. A 40x 1.3 NA objective was used, and the electronic zoom was set to achieve a pixel size of 50 nm. For visualization, mitochondria of transfected cells were labelled for 30 minutes at 37 C with 50nM MitoTracker® Deep Red FM (Invitrogen™, Thermo Scientific).

Ligand binding studies

Cells were lysed 48 h post-transfection with ice-cold 2 mM EDTA, 1 nM Na-HEPES (pH 7.4), scraped with a rubber policeman in the presence of ~4 ml rinsing medium, and homogenized with a Polytron homogenizer (ULTRA-TURRAX) in binding assay buffer (50 mM Tris-HCl, pH 7.4, 155 mM NaCl, 0.01 mg/ml bovine serum albumin). Radioligand bindings were carried out at 37 °C for 3 h in a final volume of 1 ml. Non-specific binding was determined in presence of 1 µM atropine. For competition binding assays, increasing concentrations of carbachol (from 0.01 µM up to 10 µM) were used in presence of 200 pM [^{3}H]N-Methylscopolamine ([^{3}H]NMS; 78.9 Ci/mmol; PerkinElmer).

The bound ligand was separated from the unbound ligand using glass-fiber filters (Whatmann, GF/B) with a Brandel Cell Harvester, and filters were counted with a scintillation β-counter. K_D values of [^{3}H]NMS were calculated in direct saturation experiments, whereas carbachol Inibitory Concentration $(IC)_{50}$ values were calculated in competition curves, against 400 pM [^{3}H]NMS, fitted to one binding models using the iterative, non-linear least-squares regression analysis of OriginPro 7.5 software (OriginLab Corporation).

Phosphatidylinositol Breakdown Assay

Transfected COS-7 cells were incubated with myo-[^{3}H]inositol (23 Ci/mmol; PerkinElmer) for 48 h. Immediately before the experiment the cells were washed twice with phosphate-buffered saline and incubated for 15 min in Eagle's medium containing 10 mM LiCl and 20

mM HEPES. The medium was then replaced by 0.25 ml of the same medium containing the experimental agents. After a 1-hour incubation at 25 °C the reaction was arrested by the addition of 0.75 ml of 7.5% (w/v) ice-cold trichloroacetic acid followed by a 30-min incubation on ice. The trichloroacetic acid was extracted with water-saturated diethyl ether (3 × 4 ml), and levels of IP_1 determined by anion exchange chromatography.

Adenyl Cyclase Assay

COS-7 cells were transfected with plasmid(s) containing the receptor of interest plus adenyl cyclase V. 24 h after transfection the cells were detached with trypsin and re-cultured in 24-well plates. Following an additional period of 24 h culture, they were assayed for adenyl cyclase activity. The assay was performed in triplicate**.** In brief, cells were incubated in 24-well plates for 2 h with 0.25 ml/well of fresh growth medium containing 5 µCi/ml [^{3}H]adenine (PerkinElmer). The medium was then replaced with 0.5 ml/well of DMEM containing 20 mM HEPES, pH 7.4, 0.1 mg of bovine serum albumin, and the phosphodiesterase inhibitors 1-methyl-3-isobutylxanthine (0.5 mM) and Ro-20–1724 (0.5 mM). AC activity was stimulated by the addition of 1 µM forskolin in the presence or absence of carbachol. After 10 min of incubation at +30 °C the medium was removed, and the reaction terminated by addition of perchloric acid containing 0.1 mM unlabeled cAMP followed by neutralization with KOH. The amount of [^{3}H]cAMP formed under these conditions was determined by a two-step column separation procedure (3) [3].

Immunoblot of muscarinic M$_2$ receptor mutants transfected in HEK293 cells

Cells were seeded in 10 cm plates and transfected after 24h according to manufacturer protocols (Effectene, Qiagen) using 2 µg of plasmid DNA. 48h after transfection cells were washed (2x) in ice cold PBS. 200 µL per plate of ice-cold lysis buffer were added. Lysis buffer was a RIPA buffer: 50 mM Tris-HCL pH 8, 150 mM NaCl, 1% TRITON, 0.5% sodium deoxycholate and 0.1% SDS. The following compounds were added to the buffer to a final 1x dilution: 100x Halt Protease Inhibitor Cocktail (Thermo Fisher), 100x 0.5 M EDTA and (10x) 10 mM PMSF. Cells were then scraped and the cell lysate transferred to pre-cooled 1.5 mL Eppendorf tubes. The tubes were placed in a Thermomixer at 4 °C, 300 rpm for 30 minutes. Then the Eppendorf tubes were centrifuged for 30 minutes at 14000 rpm at 4 °C. The supernatant was then transferred to new Eppendorf tubes. Protein concentration in the cell lysates was quantified by a BCA assay (Pierce BCA Protein Assay Kit, from Thermo Fisher) according to manufacturer instructions.

Cell lysates were loaded into 10% polyacrylamide gels. Cell lysates were loaded using Laemmli Buffer 2x (Sigma Aldrich) in equal mass. As a reference marker we loaded 5 uL of PageRuler Prestained Protein Ladder (Thermo Fisher). Gels were mounted into a Mini

Protean Tetra Cell kit (BioRad) using 1x SDS running buffer. Gels were ran using a constant voltage of 90V for approximately 15 minutes, and afterwards at 130V for 45 minutes. Gels were transferred to PVDF membranes. Wet transfer was achieved by using the Mini Trans Blot Module (BioRad) in Wet Transfer Buffer (25 mM Tris pH 8.3, 192 mM Glycine and 20% MetOH) at 350 mA for 90 minutes. The immunoblotted membranes were then blocked in 5% Milk in TBS-T for 1 hour at room temperature. The membrane was then incubated overnight at 4 °C with primary antibody solution using the following concentrations depending on the antibody used: 1:1000 anti β-actin (13E5) rabbit primary antibody (Cell Signaling), 1:1000 anti myc-Tag (9B11) mouse primary antibody (Cell Signaling). After overnight incubation, the membranes were washed 3x 10 minutes in TBS-T and incubated at RT for 1 h with a secondary antibody (1:10000 anti-mouse IgG/(anti rabbit) HRP-linked antibody from Cell Signaling).

After the incubation with the secondary antibody, the membrane was washed 3x 10 minutes in TBS-T. Detection was achieved by eliciting HRP luminescence by incubating the membrane with SuperSignal WestFemto solution (Thermo Fisher) according to the manufacturer's instructions, and imaging the membrane using a c600 Transilluminator from Azure Biosystems.

Animal Experiments

All animal experiments were carried out according to the German animal welfare act, considering the guidelines of the National Institute of Health and the 2010/63/EU Directive of the European Parliament on the protection of animals used for scientific purposes. The animals had free access to food and water and were kept in individually ventilated cages under a 12h:12h light/dark regime (light from 6:30 am to 6:30 pm), a constant $22\pm2°C$ temperature and $55\pm10\%$ humidity.

The transgenic line B6.Cg-Chrm2tm1.1Hze/J, also known as Chrm2-tdT-D knock-in, was obtained from 'The Jackson Laboratory' (Stock No: 030330). They are heterozygous animals, designed to have expression of a Chrm2/tdTomato fusion protein directed by the endogenous M2 muscarinic acetylcholine receptor promoter/enhancer sequences.

Tissue preparation and immunoblot of endogenous M$_2$ receptor in Chrm2-tdT-D knock-in mice

Mouse tissues were weighed out and ice-cold lysis buffer was added to approx. 0.2-0.5mg tissue/ml buffer (10mM K2HPO4, 150mM NaCl, 5mM EDTA, 5mM EGTA, 1% Triton X-100, 0.2% Na-deoxycholate. Fresh protease inhibitors PMSF 0.5mM, Na3VO4 0.1mM, NaF 50mM, Jenny's mix (Soybean trypsin inhibitor 3.2µg/ml, Aprotinin 2µg/ml, Benzamidin 0.5mM) were added to the buffer.

Samples were first homogenized, and then pellet debris were centrifuged at 4°C for 30 minutes at max speed (tabletop) and the supernatant (soluble proteins) was transferred into fresh tube (pellets = non-soluble proteins)

Protein concentration were determined with BCA assay. Cell lysates were loaded using Laemmli Buffer 2x (Sigma Aldrich) in equal mass (see above for the immunoblot procedure) into 8% polyacrylamide gels.

The membrane was then incubated overnight at 4 °C with primary antibody solution (1:1000 anti β-actin (13E5) rabbit primary antibody #4970 from Cell Signaling) (1:1000 anti GAPDH) (1:1000 anti-TdTomato rabbit primary antibody #632496 from Takara).

Isolation of mitochondrial fractions and immunodetection

Cells were seeded in 15 cm plates and transfected after 24h according to manufacturer protocols (Effectene, Qiagen) using 5 µg of plasmid DNA. 48h after transfection cells were harvested and centrifuged at 1000 rpm for 10 minutes, resuspended in NaCl 0.9% (w/v) and centrifuged at 1000 rpm for 10 minutes.

Mouse tissues were freshly excised and placed in ice cold PBS (1x). The samples were washed with NaCl 0.9% (w/v). The homogenization steps were performed with Precellys Evolution Homogenizer, at 4500 rpm for 10 seconds.

The samples were processed according to manufacturer protocols (Qproteome Mitochondria Isolation Kit, Qiagen). From the different steps were obtained three different fractions: the cytosolic, the microsomal and the mitochondrial fraction.

The fresh mitochondrial pellet was resuspended in 150µL of ice-cold lysis buffer and the protein concentration of the three fractions from each sample was quantified with a BCA assay (see above).

Cell lysates were loaded into 12% polyacrylamide gels using an SDS loading buffer (6x solution: 2.28g Tris-HCl, 2.313 DTT, 6g SDS, 30% Glycerol and 0.03% bromophenol blue in a final volume of 50 mL) in equal mass (see above for the immunoblot procedure).

The membrane was then incubated overnight at 4 °C with primary antibody solution using the following concentrations: 1:1000 anti β-actin (13E5) rabbit primary antibody (#4970 from Cell Signaling), 1:1000 anti-tdtomato rabbit primary antibody (#632496 from Takara), 1:1000 anti β-tubulin mouse primary antibody, 1:1000 anti GAPDH mouse primary antibody, 1:1000 anti COXIV mouse primary antibody (#ab33985 from abcam).

Quantification of M_2tail(368-466) mitochondrial import kinetics from confocal time sequences Before imaging, 2.5×10^4 HEK-TSA cells were seeded in each PDL-coated well of the µ-Slide 8 well (Ibidi) with DMEM (ScienCell™) supplemented with 10% FBS and 1% P/S overnight, and transfected with a total of 2 µg C-terminal EGFP tagged M_2 tail using

Lipofectamine 2000 (Thermo Fisher Scientific) according to the manufacturer protocol after washing with 200 µL PBS and refreshing with 100 µL DMEM (ScienCell™) supplemented with 10% FBS and 1% P/S. After transfection for five hours, mitochondria of transfected cells were rapidly labelled with 50nM MitoTracker® Deep Red FM (Invitrogen™, Thermo Scientific) in FluoroBrite DMEM medium (Thermo Scientific) supplemented with 1% P/S for 20 minutes after the removal of old culture medium. After mitochondria labeling, the µ-Slide were washed twice with FluoroBrite DMEM medium (Thermo Scientific) and subsequently mounted onto the TCS SP8 confocal microscope (Leica) (See imaging). Firstly, the stage of the microscope was equilibrated in temperature, and the acquisition mode was set to the tile scanning. After storing the position of each desired transfected cells, the imaging mode was set to XYT and a HC PL APO CS2 40×/NA1.3 oil immersion objective was selected. Both DIC channel and HyD were utilized to obtain 512×512 pixels images. Laser wavelengths were set as described in the section Fluorescence Microscopy above. All settings remained the same throughout the whole measurement. Images were taken at every 30 minutes over a 5h interval. The images were afterwards analyzed by ImageJ software, using the Plugin 'JACoP'. Briefly, the degree of colocalization between EGFP-tagged M_2tail(368-466) (CH1) and MitoTracker stained mitochondria (CH2) was quantified by Mander's colocalization coefficients (40) [38]. The two colocalization coefficients M1 ($M1 = \frac{CHB}{CH1}$) and M2 ($M2 = \frac{CHB}{CH2}$), CHB indicates overlapping pixels in channel 1 and channel 2. A threshold was determined for individual channel in order to distinguish specific signals from nonspecific ones. Due to their sensitivities to thresholds, the Pearson correlation coefficient (PCC)(41) [39], as well as the faster decaying attribute of nonspecific signals, were also taken into account while estimating thresholds for each channel. For the analysis of each cell, the threshold for Mitotracker channel was kept constant. Whereas the threshold for EGFP channel was evaluated based on the PCC value.

Transfection and expression of split-GFP constructs in HEK293 cells

To perform the sequential split GFP complementation assay illustrated in <mark>Figs 4D and 5C-D</mark>, 3×10⁵ HEK-TSA cells were seeded on a 10% poly-D-lysine (PDL)-coated 25mm coverslip in each well of a Falcon® 6-well flat-bottom plate (StemCell™) and cultured overnight. Cells were transfected using Effectene (see above) with a 0.4 µg Mito-GFP1-10 plasmid. The cells were then washed once with fresh 2 mL phosphate-buffered saline (PBS) first and supplemented with 1.5 mL fresh DMEM medium (ScienCell™) supplemented with 10% FBS and 1% P/S. After adding the transfection mixture, the cells were cultured overnight. After the first transfection, a second transfection of a 0.6 µg plasmid encoding either the C-terminal GFP11 tagged M_2tail(368-466) or M_2 receptor was performed using the Effectene Transfection Reagent (Qiagen). After washing each well

with 2 mL PBS, the old medium was exchanged with 1.5 mL low glucose Minimum Essential Medium (MEM) (Gibco™, Thermo Scientific) supplemented with 1% P/S, in order to create a stress-like cell starvation condition. After a four hours starvation, the cells were fixed using 4% polyformaldehyde (PFA)/PBS at 37℃ for 30 minutes and washed twice with PBS before imaging.

Oxygen consumption measured with the Seahorse assay

$5x10^4$ COS7 cells were seeded in seahorse 24MW plates with Complete media (DMEM supplemented with 10%FBS and 1X PS) and then incubated for about 4h at 37C in a cell incubator. After this incubation, cells were washed twice with Seahorse media supplemented with glutamate (1 mM), pyruvate (1 mM) and glucose (20 mM) and then incubated for 1h at 37 °C in an incubator without CO2 before being analyzed in the Seahorse XF Analyzers. In particular, the baseline cell oxygen consumption rate (OCR) was measured at the beginning of the assay, followed by measurements of ATP linked respiration and maximal respiratory capacity after the addition of respectively oligomycine (1 μM) and the FCCP protonophore (1 μM). Finally, rotenone (1 μM), an inhibitor of complex I, and antimycin (1 μM), inhibitor of complex III, were added to shut down completely the electron transport chain (ETC) function, revealing the non-mitochondrial respiration.

Extracellular oxygen consumption

HEK293 cells 24h after transfection and hPSCs were harvested and seeded respectively in 96 multiwell black plate and pre-coated (Geltrex A1413302, Thermo Fisher) 96 multiwell black plate, $5x10^4$ cells for well; after incubation O/N at 37 °C and in 5% CO2, the cell culture medium was replaced by HBSS and the assay were performed according to the manufacturer protocols (abcam Extracellular Oxygen Consumption Assay).

Generation of M368A hiPSC clone

The human induced pluripotent stem cell (hiPSC) line BIHi005-A (https://hpscreg.eu/cell-line/BIHi005-A) was used to generate an isogenic iPSC clone (BIHi005-A-39) with M368A mutation using a protocol previously described (42) [40]. In brief, small guide RNA (sgRNA) targeting the M_2 gene close to the mutation site to be corrected were designed using the http://crispor.tefor.net/ webpage and the sgRNA (5`- AGTCATCTTCACAATCTTGC-3`) were synthesized by Integrated DNA technologies (https://eu.idtdna.com). The donor ssODN template was designed to convert the ATG to GCT and was synthesized as an Ultramar DNA oligo by IDT. (5´-AAAAGTGACTCATGTACCCCAACTAATACCACCGTGGAGGTAGTGGGGTCTTCAGGT

CAGAATGGAGATGAAAAGCAGAATATTGTAGCCC<u>GCAAGATTGTGAAGGCTACTAAGC</u>
AGCCTGCAAAA-3´). The Ribonucleoprotein (RNP) complexes were prepared by incubating of 1.5µg Cas-9 protein and 360 ng gRNA for 10 minutes at room temperature. For delivery of RNPs and ssODN template, 10µl cell suspension containing Cas9 RNPs and ssODN and 1x105 cells were electroporated using the Neon transfection System (ThermoFisher Scientific). The electroporated cells were plated in one well of 6 well plate with StemFlex media (ThermoFisher Scientific) supplemented CloneR™ (Stemcell technologies). Three days after the transfection we analyzed the bulk population using the Sanger sequencing to estimate the editing efficiency. Thereafter, the automated single cell cloning of the genome edited cell pool was performed as described in protocol (43) [41]. The clones were screened by SANGER sequencing. The positive confirmed clones were banked and characterized for pluripotency and karyotype stability as described in (44) [42]. Briefly, for pluripotency assay, single cells were labelled with conjugated antibodies. Surface markers (SSEA1, SSEA4 and TRA1-60) staining was performed in unfixed cells by incubation with antibodies diluted in 0.5% BSA/PBS (10 min, 4 °C, dark). Intracellular markers (OCT3/4, NANOG) were stained by fixation/permeabilization solution (Miltenyi Biotec, 130-093-142) (30 min, 4 °C, dark) followed by antibody incubation in permeabilization buffer (Miltenyi Biotec) (30 min, 4 °C, dark). Analysis was done using the MACSQuant® AnalyzerVYB and FlowJo v10.4.

Briefly, for karyotype stability, genomic DNA was isolated using the DNeasy blood and tissue kit (Qiagen, Valencia, CA, United States) and samples were analyzed using the human Illumina OMNI-EXPRESS-8v1.6 BeadChip. First, the genotyping was analyzed using GenomeStudio 1 genotyping module (Illumina). Thereafter, KaryoStudio 1.3 (Illumina) was used to perform automatic normalization and to identify genomic aberrations utilizing settings to generate B-allele frequency and smoothened Log R ratio plots for detected regions. The stringency parameters used to detect copy number variations (CNVs) were set to 75 kb (loss), 100 kb (gain) and CN-LOH (loss of heterozygosity) regions larger than 3 MB.

Culture and handling of hiPSCs cells

hiPSCs were cultured on Geltrex (Thermo Fisher) coated plates using: Essential 8 medium (Thermo Fisher). The culture was routinely monitored for mycoplasma contamination using a PCR assay. 10 µM ROCK Inhibitor Y-27632 2HCl (Selleck Chemicals, #SEL-S1049) was added after cell splitting to promote survival. hiPSC cultures were kept in a humidified atmosphere of 5% CO_2 at 37°C and 5% oxygen. hiPSCs were undirectedly differentiated to a mixture of three lineages (ectodermal, endodermal and mesodermal) by replacing Essential 8 medium with DMEM-F12 medium supplemented with 10% FBS for seven days.

10 µM ROCK Inhibitor Y-27632 2HCl (Selleck Chemicals) was added after cell splitting to

032 promote survival.

Cell proliferation assay with HEK293 cells

$4x10^5$ HEK293 cells 24h after transfection and $4x10^5$ hPSCs were seeded respectively in

T25 flask and pre-coated T25 flask, in culture medium with Atropine at final concentration

of 100nM; the medium with Atropine was replaced every day. After 72h from transfection of

the HEK293 and from seeding of the hiPSCs, the medium and the cells were collected and

centrifuged, the pellet was resuspended in the same amount of medium for each pellet; an

aliquot of the resuspension was mixed with the same amount of Trypan blue (ratio 1:1), 10

microliters were used for counting in a Neubauer chamber.

Fluorescence plate reader assays

HEK293 cells were grown on 6 well plates without coverslips, and, upon reaching 60-70%

confluence, the cells were transfected using Effectene (Quiagen, Venlo Netherlands)

according to the manufacturer's instructions. After 24 h the cells were split and transferred

to 96 well (black plate) at a seeding density of 50,000 cells/well. The 96 well plate was pre-

coated with Poly-D-Lysine for 30 min at RT. The medium was removed and each well was

washed 3x with 100µl HBSS. The cells were finally analyzed by a fluorescence plate

reader (Neo2, BioTek, Bad Friedrichshall) in HBSS.

Statistical analysis

Significance values reported in the graphs, in comparing tabulated values, were

determined by a one-tailed Student t-test using Prism 9 (GraphPad Software), and

significance values (p-values) are indicated according to the following legend: ****

$p<0.0001$; *** $0.0001<p<0.001$; ** $0.001<p<0.01$; * $0.01<p<0.05$. p-values larger than 0.05

are deemed non-significant. Unless otherwise indicated, bar charts and scatter plots

indicate average value with s.e.m.

Acknowledgments

We are grateful to Dr. Marco Marcia (EMBL) for insightful discussion. I.P acknowledges a short term exchange grant from theDeutscher Akademischer Austauschdienst (DAAD). F.P. Acknowledges support from the University of L'Aquila towards international mobility.

Funding:

European Union's Horizon2020 Marie Skłodowska-Curie Actions (MSCA) Program under Grant Agreement 641833 (ONCORNET). This project was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through Project 421152132 SFB1423 subproject C03 (P.A. and M.J.L.)

Author contributions:

Conduced primary experiments: FP, IF, PA, ZW, MR, IP, RE, HZ, UZ, MET, JDG, SS, ART, SA, MC

Contributed to the experimental data: MG, CP, GA, FM, BP, AR

Contributed tools and reagents: NT and SD

Contributed to experimental design and assisted in writing paper: PA, IP, IF, MR, UZ, AT, KT, VF, FA, FG, MS

Conceived and lead the study, and wrote the paper: PA, MJL and RM

Competing interests:

The authors declare that they have no conflicts of interest with the contents of this article.

Data and materials availability:

Source data will be provided with this manuscript.

References

1. Kruse AC, Kobilka BK, Gautam D, Sexton PM, Christopoulos A, Wess J. Muscarinic acetylcholine receptors: novel opportunities for drug development. Nat Rev Drug Discov. 2014;13: 549–560. doi:10.1038/nrd4295

2. Fryer AD, Christopoulos A, Nathanson NM, editors. Muscarinic Receptors. Berlin, Heidelberg: Springer Berlin Heidelberg; 2012. doi:10.1007/978-3-642-23274-9

3. Novi F, Scarselli M, Corsini GU, Maggio R. The Paired Activation of the Two Components of the Muscarinic M3 Receptor Dimer Is Required for Induction of ERK1/2 Phosphorylation. Journal of Biological Chemistry. 2004;279: 7476–7486. doi:10.1074/jbc.M306818200

4. Calebiro D, Nikolaev VO, Persani L, Lohse MJ. Signaling by internalized G-protein-coupled receptors. Trends Pharmacol Sci. 2010;31: 221–228. doi:10.1016/j.tips.2010.02.002

5. Zhang X, Bedigian A v., Wang W, Eggert US. G protein-coupled receptors participate in cytokinesis. Cytoskeleton. 2012;69: 810–818. doi:10.1002/cm.21055

6. Hebert-Chatelain E, Desprez T, Serrat R, Bellocchio L, Soria-Gomez E, Busquets-Garcia A, et al. A cannabinoid link between mitochondria and memory. Nature. 2016;539: 555–559. doi:10.1038/nature20127

7. Suofu Y, Li W, Jean-Alphonse FG, Jia J, Khattar NK, Li J, et al. Dual role of mitochondria in producing melatonin and driving GPCR signaling to block cytochrome c release. Proceedings of the National Academy of Sciences. 2017;114. doi:10.1073/pnas.1705768114

8. Popot J-L, de Vitry C. On the Microassembly of Integral Membrane Proteins. Annu Rev Biophys Biophys Chem. 1990;19: 369–403. doi:10.1146/annurev.bb.19.060190.002101

9. Popot JL, Engelman DM. Membrane protein folding and oligomerization: the two-stage model. Biochemistry. 1990;29: 4031–4037. doi:10.1021/bi00469a001

10. Maggio R, Fasciani I, Rossi M, di Gregorio J, Pietrantoni I, Puca V, et al. Variants of G protein-coupled receptors: a reappraisal of their role in receptor regulation. Biochem Soc Trans. 2016;44: 589–594. doi:10.1042/BST20150239

11. Jackson RJ, Kaminski A. Internal initiation of translation in eukaryotes: the picornavirus paradigm and beyond. RNA. 1995;1: 985–1000.

12. Hellen CUT, Sarnow P. Internal ribosome entry sites in eukaryotic mRNA molecules. Genes Dev. 2001;15: 1593–1612. doi:10.1101/gad.891101

13. Maggio R, Vogel Z, Wess J. Reconstitution of functional muscarinic receptors by co-expression of amino- and carboxyl-terminal receptor fragments. FEBS Lett. 1993;319: 195–200. doi:10.1016/0014-5793(93)80066-4

14. Bathe-Peters M, Gmach P, Annibale P, Lohse MJ. Linescan microscopy data to extract diffusion coefficient of a fluorescent species using a commercial confocal microscope. Data Brief. 2020;29: 105063. doi:10.1016/j.dib.2019.105063

15. Dong C, Filipeanu CM, Duvernay MT, Wu G. Regulation of G protein-coupled receptor export trafficking. Biochimica et Biophysica Acta (BBA) - Biomembranes. 2007;1768: 853–870. doi:10.1016/j.bbamem.2006.09.008

16. Kamiyama D, Sekine S, Barsi-Rhyne B, Hu J, Chen B, Gilbert LA, et al. Versatile protein tagging in cells with split fluorescent protein. Nat Commun. 2016;7: 11046. doi:10.1038/ncomms11046

17. Newman LE, Schiavon C, Kahn RA. Plasmids for variable expression of proteins targeted to the mitochondrial matrix or intermembrane space. Cell Logist. 2016;6: e1247939. doi:10.1080/21592799.2016.1247939

18. Thakor N, Holcik M. IRES-mediated translation of cellular messenger RNA operates in eIF2α- independent manner during stress. Nucleic Acids Res. 2012;40: 541–552. doi:10.1093/nar/gkr701

19. Holcik M, Sonenberg N. Translational control in stress and apoptosis. Nat Rev Mol Cell Biol. 2005;6: 318–327. doi:10.1038/nrm1618

20. Bonfili L, Cecarini V, Cuccioloni M, Angeletti M, Flati V, Corsetti G, et al. Essential amino acid mixtures drive cancer cells to apoptosis through proteasome inhibition and autophagy activation. FEBS J. 2017;284: 1726–1737. doi:10.1111/febs.14081

21. Ishizuka T, Ozawa A, Katsuura M, Nomura S, Satoh Y. Effects of muscarinic acetylcholine receptor stimulation on the differentiation of mouse induced pluripotent stem cells into neural progenitor cells. Clin Exp Pharmacol Physiol. 2018;45: 1198–1205. doi:10.1111/1440-1681.12993

22. Hoogduijn MJ, Cheng A, Genever PG. Functional Nicotinic and Muscarinic Receptors on Mesenchymal Stem Cells. Stem Cells Dev. 2009;18: 103–112. doi:10.1089/scd.2008.0032

23. Weinberg ZY, Puthenveedu MA. Regulation of G protein-coupled receptor signaling by plasma membrane organization and endocytosis. Traffic. 2019;20: 121–129. doi:10.1111/tra.12628

24. Jong Y-JI, Harmon SK, O'Malley KL. GPCR signalling from within the cell. Br J Pharmacol. 2018;175: 4026–4035. doi:10.1111/bph.14023

25. Bénard G, Massa F, Puente N, Lourenço J, Bellocchio L, Soria-Gómez E, et al. Mitochondrial CB1 receptors regulate neuronal energy metabolism. Nat Neurosci. 2012;15: 558–564. doi:10.1038/nn.3053

26. Lefkimmiatis K, Leronni D, Hofer AM. The inner and outer compartments of mitochondria are sites of distinct cAMP/PKA signaling dynamics. Journal of Cell Biology. 2013;202: 453–462. doi:10.1083/jcb.201303159

27. Marti-Solano M, Crilly SE, Malinverni D, Munk C, Harris M, Pearce A, et al. Combinatorial expression of GPCR isoforms affects signalling and drug responses. Nature. 2020;587: 650–656. doi:10.1038/s41586-020-2888-2

28. van Heesch S, Witte F, Schneider-Lunitz V, Schulz JF, Adami E, Faber AB, et al. The Translational Landscape of the Human Heart. Cell. 2019;178: 242-260.e29. doi:10.1016/j.cell.2019.05.010

29. Pestova T v, Shatsky IN, Hellen CU. Functional dissection of eukaryotic initiation factor 4F: the 4A subunit and the central domain of the 4G subunit are sufficient to mediate internal entry of 43S preinitiation complexes. Mol Cell Biol. 1996;16: 6870–6878. doi:10.1128/MCB.16.12.6870

30. Jan E, Sarnow P. Factorless Ribosome Assembly on the Internal Ribosome Entry Site of Cricket Paralysis Virus. J Mol Biol. 2002;324: 889–902. doi:10.1016/S0022-2836(02)01099-9

31. Fitzgerald KD, Semler BL. Bridging IRES elements in mRNAs to the eukaryotic translation apparatus. Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms. 2009;1789: 518–528. doi:10.1016/j.bbagrm.2009.07.004

32. Davies CW, Vidal SE, Phu L, Sudhamsu J, Hinkle TB, Chan Rosenberg S, et al. Antibody toolkit reveals N-terminally ubiquitinated substrates of UBE2W. Nat Commun. 2021;12: 4608. doi:10.1038/s41467-021-24669-6

33. Ingolia NT, Hussmann JA, Weissman JS. Ribosome Profiling: Global Views of Translation. Cold Spring Harb Perspect Biol. 2019;11: a032698. doi:10.1101/cshperspect.a032698

34. Zhang J, Nuebel E, Daley GQ, Koehler CM, Teitell MA. Metabolic Regulation in Pluripotent Stem Cells during Reprogramming and Self-Renewal. Cell Stem Cell. 2012;11: 589–595. doi:10.1016/j.stem.2012.10.005

35. Guo C, Sun L, Chen X, Zhang D. Oxidative stress, mitochondrial damage and neurodegenerative diseases. Neural Regen Res. 2013;8: 2003–14. doi:10.3969/j.issn.1673-5374.2013.21.009

36. Kritsiligkou P, Chatzi A, Charalampous G, Mironov A, Grant CM, Tokatlidis K. Unconventional Targeting of a Thiol Peroxidase to the Mitochondrial Intermembrane Space Facilitates Oxidative Protein Folding. Cell Rep. 2017;18: 2729–2741. doi:10.1016/j.celrep.2017.02.053

37. Bonner TI, Buckley NJ, Young AC, Brann MR. Identification of a Family of Muscarinic Acetylcholine Receptor Genes. Science (1979). 1987;237: 527–532. doi:10.1126/science.3037705

38. MANDERS EMM, VERBEEK FJ, ATEN JA. Measurement of co-localization of objects in dual-colour confocal images. J Microsc. 1993;169: 375–382. doi:10.1111/j.1365-2818.1993.tb03313.x

39. Manders EM, Stap J, Brakenhoff GJ, van Driel R, Aten JA. Dynamics of three-dimensional replication patterns during the S-phase, analysed by double labelling of DNA and confocal microscopy. J Cell Sci. 1992;103: 857–862. doi:10.1242/jcs.103.3.857

40. Richardson CD, Ray GJ, DeWitt MA, Curie GL, Corn JE. Enhancing homology-directed genome editing by catalytically active and inactive CRISPR-Cas9 using asymmetric donor DNA. Nat Biotechnol. 2016;34: 339–344. doi:10.1038/nbt.3481

41. Vallone VF, Telugu NS, Fischer I, Miller D, Schommer S, Diecke S, et al. Methods for Automated Single Cell Isolation and Sub-Cloning of Human Pluripotent Stem Cells. Curr Protoc Stem Cell Biol. 2020;55. doi:10.1002/cpsc.123

42. Metzler E, Telugu N, Diecke S, Spuler S, Escobar H. Generation of two human induced pluripotent stem cell lines derived from myoblasts (MDCi014-A) and from peripheral blood mononuclear cells (MDCi014-B) from the same donor. Stem Cell Res. 2020;48: 101998. doi:10.1016/j.scr.2020.101998

A

M$_2$wt-EGFP

M$_2$stop400-EGFP

non transfected

C

M$_2$fr.sh.-EGFP

B

M$_2$ wt

M$_2$Stop228

M$_2$Stop400

M$_2$TrunC
(1-466/383)

M$_2$ fr.sh.

originating
from the same gene

originating
from the same gene

C-terminus present, functional receptor

C-terminus absent, non-functional receptor

C-terminus present

normal conditions

stress conditions

Extracellular space

Cell membrane

full length
M₂ receptor

canonical
signaling

ROS

ADP + P → ATP

ATP synthase

5' G P P P M₂ receptor mRNA IRES AAAAAA₁₅₀₋₂₅₀ 3'

M₂ tail

Cell nucleus

ROS

ADP + P → ATP

5' G P P P M₂ receptor mRNA IRES AAAAAA₁₅₀₋₂₅₀ 3'