***In vitro* reconstitution of Sgk3 activation by phosphatidylinositol-3-phosphate**

Daniel Pokorny[1,2], Linda Truebestein[1,2], Kaelin D. Fleming[3], John E. Burke[3,4] and

Thomas A. Leonard[1,2,5*]

[1]Department of Structural and Computational Biology, Max Perutz Labs, Campus

Vienna Biocenter 5, 1030 Vienna, Austria

[2]Department of Medical Biochemistry, Medical University of Vienna, 1090 Vienna,

Austria

[3]Department of Biochemistry and Microbiology, University of Victoria, Victoria, BC,

Canada V8W 2Y2

[4]Department of Biochemistry and Molecular Biology, The University of British

Columbia, Vancouver, British Columbia V6T 1Z3, Canada

[5]Correspondence: thomas.leonard@meduniwien.ac.at

Summary

Serum- and glucocorticoid-regulated kinase 3 (Sgk3) is activated by the phospholipid

phosphatidylinositol-3-phosphate (PI3P) downstream of growth factor signaling and

by Vps34-mediated PI3P production on endosomes. Upregulation of Sgk3 activity has

recently been linked to a number of human cancers. Here, we show that Sgk3 is

regulated by a combination of phosphorylation and allosteric activation by PI3P. We

demonstrate that PI3P binding induces large conformational changes in Sgk3

associated with its activation, and that the PI3P binding pocket of the PX domain of

Sgk3 is sequestered in its inactive conformation. Finally, we reconstituted Sgk3

activation via Vps34-mediated PI3P synthesis on phosphatidylinositol liposomes *in*

vitro. In addition to defining the mechanism of Sgk3 activation by PI3P, our findings

open up potential therapeutic avenues in allosteric inhibitor development to target

Sgk3 in cancer.

Introduction

Cellular membranes are hotspots at which numerous signaling pathways controlling various aspects of cell physiology and pathology converge. One of the most intensively studied pathways downstream of extracellular growth factors is the phosphoinositide 3-kinase (PI3K) pathway. PI3K activation by receptor tyrosine kinases, G-protein-coupled receptors, or RAS leads to the generation of various phosphoinositide lipids, that serve as a platform for the recruitment and activation of effector molecules (Martini *et al*, 2014). Dysregulation of PI3K signaling is frequently observed in cancer, diabetes, neurodegenerative and cardiac disorders, and other diseases (Fruman & Rommel, 2014; Fruman *et al*, 2017), making the pathway a prominent target for pharmacological inhibition (Marone *et al*, 2008).

Many important downstream effectors of PI3K are protein kinases. Protein phosphorylation is a ubiquitous post-translational modification that regulates cell signaling and is carried out by the arsenal of more than 500 protein kinases in the human proteome (Manning, 2002). Since the mechanism of phosphotransfer catalyzed by protein kinases is highly conserved (Endicott *et al*, 2012), individual kinases have evolved unique regulatory mechanisms to ensure that they can faithfully respond to different, often very transient, lipid-based signaling cues (Leonard & Hurley, 2011).

Serum- and glucocorticoid-regulated kinase 3 (Sgk3) [also known as cytokine-independent survival kinase (CISK)] is a serine/threonine protein kinase that is activated downstream of growth factors (Liu *et al*, 2000). Like the closely related

kinase Akt, it belongs to the AGC kinase family and is activated by phosphorylation of

a hydrophobic motif in its C-terminal tail by mTORC2 (Bago *et al*, 2016) and

subsequent PDK1-dependent phosphorylation of its activation loop (Kobayashi &

Cohen, 1999; Kobayashi *et al*, 1999). A third site, called the turn motif, also in the tail,

regulates the stability of Akt and protein kinase C (PKC) (Ikenoue *et al*, 2008;

Facchinetti *et al*, 2008), but its phosphorylation has not been reported in Sgk3.

Distinct from the activation of Akt by PIP_3 or $PI(3,4)P_2$, Sgk3 depends on the lipid

phosphatidylinositol-3-phosphate (PI3P) for its activation (Virbasius *et al*, 2001).

Sgk3 is recruited specifically to endosomes via its PI3P-binding phox homology (PX)

domain (Tessier & Woodgett, 2006; Xing *et al*, 2004), which promotes

phosphorylation of its activation loop and hydrophobic motif by PDK1 and mTORC2

respectively (Tessier & Woodgett, 2006; Bago *et al*, 2014, 2016). Two major pathways

drive the production of PI3P on endosomes: downstream of phosphatidylinositol-

3,4,5-trisphosphate (PIP_3) production by class I PI3K at the plasma membrane,

phosphatidylinositol-3,4-bisphosphate [$PI(3,4)P_2$] is produced by the action of SH2-

containing inositol 5' phosphatase (SHIP2) (Liu *et al*, 2018; Goulden *et al*, 2019).

Phosphatidylinositol-3-phosphate (PI3P) is generated either from $PI(3,4)P_2$ by

inositol polyphosphate 4-phosphatases (INPP4) (Ivetac *et al*, 2005) or, de novo, from

phosphatidylinositol by the class III PI3K Vps34 (Backer, 2008). Both pathways have

been shown to contribute to Sgk3 activation (Bago *et al*, 2014; Malik *et al*, 2018).

Sgk isoforms are transcriptionally induced by serum and glucocorticoid

hormones (Webster *et al*, 1993) and are involved, among others, in the regulation of

ion channels, transcription factors, enzymatic activities and have been implicated in

hair growth defects and renal dysfunction (Lang *et al*, 2006). Recently, Sgk3 has been

shown to play a role in different cancers in an Akt-independent manner (Gasser *et al*,

2014; Bago *et al*, 2016; Vasudevan *et al*, 2009; Wang *et al*, 2017), highlighting the

importance of Sgk3 as an effector downstream of PI3K signaling. Despite its

physiological and pathological relevance, however, relatively little is known about the

precise mechanism of Sgk3 regulation. In this manuscript, we provide clear evidence

for the allosteric activation of Sgk3 by PI3P, highlighting the role of the PX domain

and PI3P in the regulation of Sgk3 enzymatic activity. Our findings have obvious

implications for the spatial and temporal control of Sgk3 activity as well as for the

design of Sgk3-targeted therapeutics.

Results

Sgk3 localizes specifically to PI3P-containing endosomes

The Sgk3 PX domain (residues 1-124) has previously been shown to bind to PI3P, and is required for localization at early endosomes (Xu *et al*, 2001; Chandra *et al*, 2019; Virbasius *et al*, 2001; Bago *et al*, 2014). Serum-starved HeLa cells transfected with mCherry-Sgk3 show a punctate endosomal membrane localization (Fig. 1A), which is lost upon 20 min treatment with the PI3K inhibitor wortmannin (Fig. 1B). The membrane localization of Sgk3 is strictly dependent on its PX domain, since a binding site mutant of Sgk3 (Sgk3[R90A]) fails to localize to endosomes in the same cells (Fig. 1C-D). This is further illustrated by a representative cross section intensity profile (Fig. 1E-F). These data confirm both the specificity of Sgk3 for PI3P *in vivo* and the endosomal localization of Sgk3 to endosomes.

In vitro **characterization of recombinant Sgk3**

The phosphorylation state of recombinant human Sgk3 co-expressed with PDK1 and purified from baculovirus-infected insect cells was characterized by tandem mass spectrometry (Supplementary Figure 1A). Sub-stoichiometric (~20%) activation loop (Thr320) phosphorylation was observed; however, phosphorylated peptides were not observed for either the turn motif (Ser461) or the hydrophobic motif (Ser486), despite purification with phosphatase inhibitors known to preserve phosphorylation at these sites in Akt1 (manuscript in revision). Turn motif phosphorylation has previously been reported for Sgk1 (Chen *et al*, 2009) and other

AGC kinases, in particular Akt and PKC, in which it plays an important role in

regulating protein stability and degradation (Oh *et al*, 2010; Facchinetti *et al*, 2008;

Ikenoue *et al*, 2008; Hauge *et al*, 2007). Since both the turn motif and the basic patch

to which it binds on the kinase domain are conserved, it seemed logical to assume

that Sgk3 should also be modified at this site. Insects, however, do not express an

obvious homolog of Sgk3, raising the possibility that the kinase responsible for turn

motif phosphorylation of Sgk3 is also not present in insect cells. We therefore

expressed human Sgk3 in HEK293 cells and characterized its phosphorylation state

by tandem mass spectrometry. Surprisingly, phosphorylated peptides were again not

observed for either the turn motif or hydrophobic motif (Supplementary Figure 1B).

It is therefore an open question as to whether Sgk3 is indeed modified on the turn

motif and under what cellular conditions.

To prepare site-specifically phosphorylated Sgk3 for in vitro kinase assays,

both wild-type Sgk3 (Sgk3^{WT}) and kinase-dead Sgk3 (Sgk3^{D286A}) were

dephosphorylated with lambda phosphatase and re-phosphorylated with

recombinant PDK1. Specific modification of Sgk3 with a single phosphate on its

activation loop (Thr320) was confirmed by mass spectrometry (Supplementary

Figure 1C-D). The purification of recombinant Sgk3 was challenging, due to solubility

issues under certain purification conditions. However, careful optimization of the

purification protocol enabled us to purify Sgk3 in a detergent-free buffer with pH and

ionic strength, close to physiological values (see Materials and Methods). We

performed size-exclusion chromatography coupled to multi-angle light scattering to

determine its precise mass and monodispersity. Sgk3 exhibited a single peak with a

measured mass of 57.3 kDa, in close agreement with its actual mass reported by mass

spectrometry and a polydispersity value of 1.001 (Supplementary Figure 1E). To

ensure that the Sgk3 used in subsequent assays was monodisperse and remained so

for the duration of the assays, we measured the particle size distribution of Sgk3 by

dynamic light scattering at 0 and 60 min post-freeze-thaw under kinase assay buffer

conditions. Sgk3 exhibited near identical mass distributions centered around a

particle size of 3 nm at 0 and 60 min at 20°C (Supplementary Figure 1F).

140 Sgk3 is specifically and allosterically activated by PI3P

Previous studies have shown that loss of PI3P binding decreases phosphorylation of

Sgk3 on its activation loop and hydrophobic motif (Tessier & Woodgett, 2006), as well

as downstream substrate phosphorylation (Bago *et al*, 2016). However, it is not

known whether PI3P binding serves to allosterically activate Sgk3 in addition to its

known role in promoting phosphorylation of the activation loop and hydrophobic

motif (Bago *et al*, 2014). Previous studies have established that the closely-related

kinase Akt depends on the binding of PI(3,4,5)P$_3$ or PI(3,4)P$_2$ for full activity (Ebner

et al, 2017; Lučić *et al*, 2018). We therefore investigated whether PI3P is simply a

membrane anchor to target Sgk3 to endosomes, or whether PI3P also allosterically

governs Sgk3 activity.

Having established that our recombinant Sgk3 is well-behaved during our

assays, we assessed the influence of PI3P on kinase activity. Dephosphorylated Sgk3

exhibited undetectable kinase activity in vitro and could not be stimulated with PI3P

(Supplementary Figure 1G). To test whether PI3P allosterically activates Sgk3, we bound PDK1-phosphorylated Sgk3^{WT} to liposomes with increasing PI3P concentrations and assayed its kinase activity against Crosstide, a glycogen synthase kinase 3β (GSK3β)-derived peptide substrate of Sgk3 (Dai *et al*), using a radiometric kinase assay (Hastie *et al*, 2006). We observed robust, PI3P-dependent activation of Sgk3, which correlated with the fraction of protein bound to the liposomes (Fig. 2A). Both the activity and binding curves can be fitted with a one site binding model characterized by nearly identical binding affinities of 24.6±10.8 μM and 29.3±8.0 μM, respectively. No signal was observed in the control without Crosstide, thereby ruling out autophosphorylation of Sgk3. Basal kinase activity against Crosstide seen in the absence of PI3P can be explained by the high degree of activation loop phosphorylation (Fig. S1C). To confirm that the observed activity comes exclusively from Sgk3, we mutated D286 in the catalytic loop to alanine to create a kinase-dead Sgk3. While this protein shows almost identical PI3P binding to Sgk3^{WT}, neither basal nor PI3P-stimulated Crosstide phosphorylation was observed (Fig. 2B).

We next reconstituted Sgk3 activation by PI3P in vitro using a coupled Vps34-Sgk3 kinase activity assay. Vps34 is the only class III PI3K expressed in mammalian cells and generates PI3P from PI on endosomes and other endomembrane structures, including autophagosomes (Hurley & Young, 2017; Backer, 2016). We prepared liposomes containing 10 mol % PI as a substrate for Vps34 and monitored both Vps34 and Sgk3 activity simultaneously in a radiometric kinase assay (Fig. 2C). We could detect a time-dependent production of PI3P in the liposomes and the consequent activation of Sgk3 as a function of increasing PI3P (Fig. 2D). This demonstrates that

Sgk3 is allosterically activated by the product of class III PI3K activity, highlighting

that full Sgk3 activity requires membranes enriched in PI3P.

HDX-MS reveals conformational changes in Sgk3 upon membrane binding

Having demonstrated that Sgk3 is allosterically activated by PI3P, we next set out to

investigate the structural basis of PI3P-mediated activation. In previous studies on

Akt, we have shown that PI(3,4,5)P_3 or PI(3,4)P_2 binding relieves pleckstrin

homology (PH) domain-induced steric occlusion of the kinase domain catalytic cleft,

leading to Akt activation (Ebner *et al*, 2017; Lučić *et al*, 2018). To verify whether this

is also the case for Sgk3, we performed hydrogen-deuterium exchange mass

spectrometry (HDX-MS) on full-length Sgk3 and Sgk3 PX domain in the presence of

vesicles containing 0 or 5 mol % PI3P. Binding of the isolated PX domain to PI3P

vesicles led to dramatically lower exchange rates for peptides across the domain,

including the PI3P binding site, indicating that PI3P binding is captured by HDX-MS

analysis (Fig. 3A). We next compared the HDX rates between full-length Sgk3^{WT} and

the isolated PX domain to determine whether they interact with each other. We

observed dramatic deprotection of the PX domain in the absence of the kinase domain

of Sgk3 (Fig. 3B), indicating that the PX domain is sequestered in an intramolecular

interaction in the full-length protein. Finally, we monitored HDX changes in full-

length Sgk3 upon PI3P binding. Protection of the PX domain upon membrane binding

was also observed, accompanied by deprotection in several areas of the kinase

domain. These sites form a contiguous network starting with β5, β7 and β8 strands

in the N-lobe and propagating to the C-lobe via the αG and αI helices (Fig. 3C).

Additional deprotection was observed in residues 468-476 localized in the C-terminal

tail. It was not possible to delineate a clear surface of interaction between the PX and

kinase domains from the changes in exchange rates in the kinase domain, but the data,

taken together, provide unambiguous evidence for an intramolecular PX-kinase

domain interaction and a large conformational change upon PI3P binding.

Autoinhibition of Sgk3 impairs binding to PI3P

Many AGC kinases that are activated by lipid second messengers exhibit a closed

conformation of their autoinhibited state in which the binding site for the lipid

headgroup is sequestered in an intramolecular interaction. To investigate whether

the autoinhibited conformation of Sgk3 might also sequester the PI3P-binding site of

its PX domain in intramolecular contacts, we compared the binding affinity for PI3P-

containing liposomes of full-length Sgk3 and its isolated PX domain. To ensure

identical conditions, we incubated both proteins with PI3P liposomes in a single

reaction, pelleted the liposomes, resolved the proteins in the supernatant and pellet

fraction by SDS-PAGE, and quantified their abundance by Coomassie stain

densitometry. To compensate for the lower staining sensitivity of the PX domain, we

used an N-terminally FITC-labelled version of this construct and quantified its

fluorescence intensity in the gel before Coomassie staining. We observed an

approximately 5-fold higher affinity to PI3P of the isolated PX domain over the full-

length protein (Fig. 4A), confirming that the PI3P binding site is sequestered by the kinase domain in the autoinhibited state.

To substantiate this finding in vivo, we expressed mCherry-Sgk3^{WT} and eGFP-Sgk3^{PX} in HeLa cells and imaged their localization in serum-starved cells by confocal microscopy. While serum starvation of the cells overnight did not reduce PI3P levels sufficiently to dissociate Sgk3 from endosomes (Fig. 1A), it was clear that the isolated Sgk3 PX domain associates almost exclusively with endosomes (Fig. 4B), while full-length Sgk3 exhibits a predominantly cytosolic distribution (Fig. 4C). A representative cross-section intensity trace for both channels reveals that the bulk of full length Sgk3 is uniformly distributed in the cytoplasm of cells, while the PX domain is exclusively found on endosomes (Figure 4D-E). These findings confirm that the PI3P binding pocket is sequestered in the autoinhibited conformation of Sgk3.

Discussion

Sgk3 is dependent on both PI3P and phosphorylation for its activity. We have shown here that Sgk3 exhibits negligible activity in the absence of activation loop phosphorylation and that PI3P significantly increases the basal activity of pre-phosphorylated Sgk3. Like other AGC kinases regulated by phospholipids, including Akt and PKC, the PI3P binding pocket of the PX domain of Sgk3 is sequestered in an autoinhibitory, intramolecular conformation. This both necessitates PI3P for activity and simultaneously raises the threshold level of PI3P that must be reached in the cell in order to activate Sgk3. The structure of Akt1 reveals an intramolecular interaction

between its PH and kinase domains that obscures the PIP_3 binding site (manuscript

in revision), a feature which impairs its binding to PIP_3 in vitro and in vivo (Ebner *et*

al, 2017). Similarly, the diacylglycerol (DAG) binding site of the C1b domain of PKCβII

is also blocked by intramolecular contacts that impair its membrane translocation in

response to phorbol ester treatment (Leonard *et al*, 2011). More generally, the

regulatory C1 domains of all PKC isozymes, as well as those of the PKD family, are

engaged in autoinhibitory intramolecular interactions that impair their binding to

membranes (Oancea & Meyer, 1998; Lučić *et al*, 2016; Elsner *et al*, 2019).

Mechanistically, Sgk3 exhibits many of the features previously described for Akt,

though given the structural divergence between the PX domain of Sgk3 and the PH

domain of Akt, the atomistic details of autoinhibition are likely to be different. Sgk3 is

the only Sgk to comprise a full PX domain, although Sgk1 has been shown to possess

a short 'PX-like' domain in its N-terminus that mediates lipid binding (Pao *et al*, 2007).

Importantly, activation loop phosphorylation by PDK1 is insufficient to override the

requirement for PI3P for full Sgk3 activity, a property also shared by Akt (manuscript

in revision).

Whilst not the major focus of this study, it is worth noting that turn motif

phosphorylation of Sgk3 could not be observed in Sgk3 expressed either in insect or

human cells. This finding was unexpected, given the prominent role of turn motif

phosphorylation in other AGC kinases. Phosphorylation of the turn motif helps to

anchor the C-terminal tail of the kinase domain and increase the affinity of the

phosphorylated hydrophobic motif to its cognate hydrophobic pocket in the N-lobe

via local concentration, which leads to the overall stabilization of the kinase fold

(Hauge *et al*, 2007). While turn motif phosphorylation has been reported in all PKC

isoforms, Akt, Sgk1, and PKN (Hornbeck *et al*, 2004), whether it plays a role in Sgk3

regulation will require further investigation. Finally, it should also be noted that while

hydrophobic motif phosphorylation has been reported to increase the catalytic

activity of Sgk3 (Bago *et al*, 2014), our study does not address its contribution to Sgk3

activation.

Evidence suggests that PI3P binding by Sgk3 also promotes PDK1-dependent

phosphorylation of its activation loop. This mechanism of activation loop exposure

upon membrane binding is well established for Akt (Alessi *et al*, 1997; Stokoe *et al*,

1997). The requirement of PI3P for Sgk3 activity is therefore three-fold: in the first

place, it is required to target Sgk3 to membranes where it can encounter active PDK1;

secondly, PI3P binding triggers a conformational change that renders it a substrate

for PDK1; thirdly, and importantly, it is required to prevent autoinhibition of the

kinase domain by the PX domain. Studies on Akt have also implicated ATP in

stabilizing a phosphatase-resistant conformation of the kinase domain (Lu *et al*, 2015;

Lin *et al*, 2012; Chan *et al*, 2011), a mechanism which we recently showed to be

dependent on binding to membrane-embedded PIP$_3$ (Lučić *et al*, 2018). While such a

mechanism has not been explicitly shown for Sgk3, the homology between the kinase

domains and the mechanisms by which Akt and Sgk3 are activated suggests that this

is also likely to be the case for Sgk3 bound to PI3P-rich endosomes.

Our findings suggest that PI3P and phosphorylation are both required for Sgk3

activity, which makes Sgk3 a coincidence detector of both PI3P and PI3K signaling.

According to this model, PI3P-decorated endosomes lacking PDK1 would not serve as

signaling hubs for Sgk3. In this respect, it is worth noting that PDK1 does not bind to,

nor can it be activated by, PI3P. PDK1 binds with almost equal affinity to both

PI(3,4,5)P$_3$ (1.6 nM) and PI(3,4)P$_2$ (5.2 nM) (Currie *et al*, 1999), which targets it to

both the plasma membrane, which is abundant in PIP$_3$, and endomembranes, which

are decorated with PI(3,4)P$_2$ and heavily implicated in Akt signaling (Liu *et al*, 2018;

Ebner *et al*, 2017; Schenck *et al*, 2008; Braccini *et al*, 2015). PI3P is generally not

observed at the plasma membrane. Presumably, PDK1 and Sgk3 cooperate to drive

Sgk3 signaling on endosomes containing both PI(3,4)P$_2$ and PI3P (Figure 5). Indeed,

PDK1 has also been reported to associate with endosomes dependent on Sgk3

hydrophobic motif phosphorylation, where it inhibits ligand-induced degradation of

the chemokine receptor CXCR4 (Slagsvold *et al*, 2006). Sgk3 activity could therefore

be restricted to a subpopulation of endosomes with a specific phosphoinositide

composition. Interestingly, a recent large scale analysis of human PX domains

revealed the presence of a secondary lipid binding site in the PX domain of Sgk3 which

exhibits low-affinity PI(3,4)P$_2$ binding (Chandra *et al*, 2019). The functional

importance of the second lipid-binding site is still unclear, though one could easily

imagine that it might serve to either fine-tune the lipid binding specificity or increase

binding affinity via an avidity effect (Carlton & Cullen, 2005).

A similar model for PI(3,4)P$_2$-driven Akt activity on endomembranes has

recently been proposed to explain the transduction of growth factor signaling by Akt

at subcellular locations distal to the plasma membrane. In this model, both PDK1 and

Akt are targeted to PI(3,4)P$_2$-containing endomembranes, where they cooperate to

drive Akt signaling in the interior of the cell. Quantitative lipid imaging has revealed

that endosomal pools of PI(3,4)P$_2$ are signaling hubs for Akt (Liu *et al*, 2018). Like

Sgk3, Akt activity depends strictly on phosphorylation and the binding of its PH

domain to either PIP$_3$ or PI(3,4)P$_2$, thereby confining its activity to membranes

enriched in these signaling lipids (Ebner *et al*, 2017; Lučić *et al*, 2018; Siess & Leonard,

2019).

Such a model of localized Sgk3 activity may provide an explanation for the

reported overlap in Akt and Sgk3 substrate specificity (Kobayashi *et al*, 1999; Tessier

& Woodgett, 2006; Brunet *et al*, 2001; Murray *et al*, 2005). Notwithstanding the

similarity in their kinase domains, it is clear that their activity profiles would overlap

in such a model, since the PI(3,4)P$_2$ that drives PDK1 and Akt activation must be

present in membranes that support PI3P- and PDK1-mediated Sgk3 activation.

Indeed, a recent phosphoproteomic study identified several Sgk3-specific substrates

localized on endosomes (Malik *et al*, 2019). The reported role of Sgk3 in driving

human cancers in the absence of Akt hyperactivation is then also reconcilable with

this model and reinforces the potential value of Sgk3 as a therapeutic target either as

a monotherapy or in combination with Akt inhibitors.

The chemical toolkit for Sgk3 inhibition is still limited due to similarities with

other AGC kinases, such as Akt, and indeed with Sgk1 and Sgk2. However, inhibitor

14h has been reported to target all three Sgk isoforms without inhibiting Akt (Bago

et al, 2016) and, more recently, an Sgk3-specific PROTAC degrader has been

described (Tovell *et al*, 2019). Much effort has been aimed at developing allosteric

inhibitors of Akt that specifically target the autoinhibited conformation of the kinase

(Wu *et al*, 2010; Lapierre *et al*, 2016; Weisner *et al*, 2019), although despite high oral

bioavailability and specificity, none have yet made it to phase III clinical trials (Landel

et al, 2020). Nevertheless, the identification of an autoinhibitory intramolecular

interface in Sgk3, analogous but not homologous to Akt, presents opportunities for

the development of Sgk3-specific inhibitors that may have clinical value in the

treatment of cancers in which Sgk3 signaling is observed to be upregulated. Such

studies will undoubtedly benefit from the high-resolution structure determination of

full-length, autoinhibited Sgk3.

Acknowledgements

Intact mass analyses and phosphorylation mapping experiments were performed by

Dorothea Anrather and Thomas Gossenreiter from the Mass Spectrometry Facility at

the Max Perutz Labs using instruments of the Vienna BioCenter Core Facilities

(VBCF). We thank Justyna Sawa-Makarska and Sascha Martens (Max Perutz Labs) for

their generous gift of baculovirus for the expression of the Vps34 complex. This work

was supported by the Austrian Science Fund (FWF) grants P28135, P30584 and

P33066 to T.L. L.T. was supported by a Hertha Firnberg Postdoctoral Fellowship from

the Austrian Science Fund (FWF T915). J.E.B. is supported by a Michael Smith

Foundation for Health Research (MSFHR) Scholar award (17686), and an operating

grant from the Cancer Research Society (CRS-24368).

Author Contributions

D.P. purified all proteins and performed all biochemical experiments. L.T. performed all fluorescence microscopy experiments. K.F. and J.B. performed all HDX-MS experiments and analyzed the corresponding data. T.L. conceived the project.

Declaration of Interests

The authors declare no competing interests.

Materials and methods

Fluorescence microscopy

For localization studies of SGK3 in live cells, HeLa cells were co-transfected with mCherry-SGK3 (wt) and either eGFP-SGK3^{PX} (residues 1-124) or eGFP-SGK3^{R90A}. Cells were transfected using Turbofect transfection reagent according to the manufacturer's protocol. Cells were serum starved overnight and changed into HBSS for live imaging. Images were acquired 24 hours after transfection on a Zeiss LSM710 or LSM 700 confocal microscope equipped with a Zeiss 63x oil immersion objective (NA 1.4) and 488 and 561 nm lasers. Image processing and analysis were done using ImageJ 1.51j. All adjustments were applied uniformly across the entire image; no nonlinear adjustments were used.

Protein expression and purification

Sgk3 PX domain (residues 1-124) was cloned into pHis-parallel bacterial expression vector and transformed into *E. c*oli BL21 STAR electrocompetent cells. For large-scale expression, the cells were grown in LB medium containing 100 μg/ml ampicillin at 37°C/180 RPM shaking, until OD$_{600}$ reached 0.6. The temperature was lowered to 20°C, expression was induced by the addition of 0.1 mM IPTG and cells were grown overnight. Cells were harvested by centrifugation at 4000g/20 min, the pellet was flash-frozen in LN2 and stored at -80°C for further use. Pellet corresponding to 1 l of

culture was resuspended in 100 ml PX lysis buffer: 50 mM Tris pH 7.5, 150 mM NaCl,

20 mM imidazole, 1% glycerol, 1 mM TCEP, 1 mM PMSF, 1 mM benzamidine

hydrochloride , 2 mM $MgCl_2$, 1x protease inhibitor cocktail (made in-house: 100 µM

bestatin, 14 µM E-64, 10 µM pepstatin A, and 1 µM phosphoramidon), 5 µl Denarase.

The cell suspension was sonicated on ice, spun at 38 724 g, 4°C for 30 min. The soluble

fraction was loaded on a HisTrap FF 5 ml column (Cytiva) equilibrated in NiA buffer:

50 mM Tris pH 7.5, 150 mM Nacl, 20 mM imidazole, 1% glycerol, 1 mM TCEP. The

column was washed with 20 column volumes (CVs) of NiA, 10 CVs of high salt wash

buffer (NiA + 500 mM NaCl), an additional 10 CVs of NiA, and the bound protein eluted

with 0.02-1 M imidazole gradient over 10 CVs. Fractions containing His-tagged PX

domain were pooled and diluted with SA buffer in (50 mM Hepes pH 8.0, 1% glycerol,

1 mM TCEP). The His tag was cleaved with TEV protease (made in-house) overnight

at 4°C. The cleaved protein was loaded on HiTrap SP HP 1 ml cation exchange column

(Cytiva) equilibrated in SA buffer. The column was washed with 10 CVs of SA and

eluted with a 0-1 M NaCl gradient over 20 CVs. Fractions containing the PX domain

were diluted with SA to adjust NaCl concentration to ~150 mM and were either used

for Sortase labeling (see below) or concentrated to 20 mg/ml, snap-frozen in LN2 and

stored at -80°C for further use.

Purified PX domain was specifically labeled on its N-terminal glycine with a

FITC-LPETGG peptide (Genscript) in a Sortase-mediated reaction according to the

protocol described in (Theile *et al*, 2013). Briefly, the PX domain was incubated with

a 10-fold excess of FITC-LPETGG peptide in the presence of His-tagged Sortase A

(prepared in-house) and 5 mM $CaCl_2$ at 4°C overnight. Sortase was removed by NiNTA

affinity chromatography and the labeled PX domain purified from free label by size

exclusion chromatography on a Superdex 75 10/300 column equilibrated in 20 mM

Tris pH 8, 150 mM NaCl, 1% glycerol, 20 mM imidazole, 1 mM TCEP. Fractions

containing FITC-labeled PX domain were pooled, concentrated to 0.8 mg/ml, snap-

frozen in LN2 and stored at -80°C.

The human Sgk3 gene was codon optimized for expression in insect cells

(Genscript) and fused to the C-terminus of His_{10}/StrepII-tagged eGFP (A206K

monomeric variant) to aid expression of soluble Sgk3. Both wild-type and kinase-

dead (D286A) Sgk3 were expressed in baculvirus-infected Sf9 insect cells grown in

ESF 921 medium (Expression Systems) at 27°C. Cells were harvested 4-5 days post-

infection. Wild-type and D286A constructs were purified under identical conditions.

Briefly, the pellet of 0.5 l Sf9 cells was resuspended in 50 ml Sgk3 lysis buffer: 50 mM

Hepes pH 7.4, 50 mM NaCl, 1% glycerol, 1 mM TCEP, 1 mM PMSF, 1 mM benzamidine,

1 mM $MnCl_2$, 0.5 mM EDTA, 1x protease inhibitor cocktail, 0.1% CHAPS, 5 ul Denarase,

10 mM DTT, 1x protease inhibitor cocktail, 5 µl Denarase, 80 nM lambda phosphatase

(made in-house). The lysate was passed through Dounce homogenizer several times

and spun at 38 724 g (18 000 RPM), 4°C for 30 min. The soluble fraction was loaded

onto a StrepTrap HP 5 ml column (Cytiva) equilibrated in 50 mM Hepes 7.4, 50 mM

NaCl, 1% glycerol, 1 mM TCEP. The column was washed with 10 CVs of binding buffer

and the protein eluted with 2.5 mM d-desthiobiotin in binding buffer. Fractions

containing EGFP-Sgk3 were pooled and dephosphorylated with lambda phosphatase

in 50 mM Hepes pH 7.4, 150 mM NaCl, 1% glycerol, 10 mM DTT, 1 mM $MnCl_2$, 0.5 mM

EDTA, 0.5 µM lambda phosphatase (made in-house). Dephosphorylation was carried

out for 30 min at room temperature and 4°C overnight together with TEV cleavage of

the His/StrepII tag. To separate His-tagged lambda phosphatase from Sgk3, the

reaction mixture was loaded on HisTrap FF 1 ml column equilibrated in 20 mM Tris

pH 7.4, 150 mM NaCl, 1% glycerol, 1 mM TCEP. Dephosphorylated and cleaved Sgk3

was washed from the column with 20 mM imidazole, while lambda phosphatase

remained bound. Sgk3 was then site-specifically in vitro re-phosphorylated by GST-

PDK1 under the following conditions: 50 mM Tris pH 7.4, 150 mM NaCl, 1% glycerol,

1 mM TCEP, 1 mM ATP, 2 mM $MgSO_4$, 1:10 PDK1:Sgk3. The reaction was carried out

at 4°C for 3 h, then stopped by addition of 10 mM EDTA. GST-PDK1 was separated by

addition of glutathione-sepharose 4B beads (Cytiva) and incubation at 4°C overnight

with gentle rotation. Supernatant containing rephosphorylated Sgk3 was

concentrated in 30 kDa MWCO spin concentrator and injected onto Superdex 200

Increase 10/300 GL column (Cytiva) equilibrated in 20 mM Tris pH 7.4, 150 mM NaCl,

1% glycerol, 1 mM TCEP. Fractions containing Sgk3 were pooled, concentrated, snap-

frozen in LN2 and stored at -80°C for further use.

Recombinant baculovirus encoding Vps34 complex I (Atg6, protein A-tagged

Atg14, Vps15 and Vps34) (Sawa-Makarska *et al*, 2020) was used to infect 3 l of Sf9

insect cell culture. Cells were harvested 4-5 days post-infection. Pellets were lysed in

300 ml of Vps34 lysis buffer: 50 mM Tris pH 8.8, 300 mM Nacl, 1 mM DTT, 0.5 %

CHAPS, 1 mM $MgCl_2$, 0.5 mM PMSF, 1x protease inhibitor cocktail, 15 ul Denarase. The

lysate was passed several times through a Dounce homogenizer and spun at 38 724 g

(18 000 RPM), 4°C for 30 min. The soluble fraction was bound to IgG-Sepharose beads

(Cytiva) for 1 h at 4°C with gentle rotation. Resin was collected in a gravity flow

column and washed 3 x 50 ml wash buffer 1 (50 mM Tris pH 8, 300 mM NaCl, 1 mM

DTT, 0.5% CHAPS, 0.5 mM PMSF), 2 x 50 ml wash buffer 2 (50 mM Tris pH 8, 300 mM

NaCl, 1 mM DT, 2% glycerol, 0.25 % CHAPS). Finally, beads were resuspended in wash

buffer 2 and bound protein complex cleaved with TEV at 4°C overnight. The

supernatant containing the full Vps34 complex was concentrated and injected onto a

Superdex 200 Increase 10/300 GL column equilibrated in 50 mM Tris pH 8, 150 mM

NaCl, 1 mM TCEP, 2% glycerol. Fractions containing the complete complex of Atg6,

Atg14, Vps15 and Vps34 were pooled, concentrated to 0.9 mg/ml, snap-frozen in LN2

and stored at -80°C for further use.

Intact mass spectrometry

Intact protein samples were diluted in H_2O and up to 100 ng protein were loaded on

an XBridge Protein BEH C4 column (2.5 µm particle size, dimensions 2.1 mm X 150

mm; Waters) using a Dionex Ultimate 3000 HPLC system (Thermo Fisher Scientific)

with a working temperature of 50 °C, 0.1% formic acid (FA) as solvent A, 80%

acetonitrile, 0.08% FA as solvent B. Proteins were separated with a 6 min step

gradient from 10 to 80% solvent B at a flow rate of 300 µL/min and analysed on a

Synapt G2-Si coupled via a ZSpray ESI source (Waters). Data were recorded with

MassLynx V 4.1 (Waters) and analyzed using the MaxEnt 1 process to reconstruct

the uncharged average protein mass.

Tandem mass spectrometry

The reduced protein was denatured in 4 M urea, 50 mM ammonium bicarbonate

(ABC) and alkylated with 20 mM iodoacetamide for 30 min at room temperature in

the dark. The sample was diluted with 50 mM ABC down to 0.5 M urea and then

digested overnight using sequencing-grade endoproteinase GluC (Roche) at 25 °C.

The digestion was stopped by adding trifluoroacetic acid to a final concentration of

1% and the peptides were desalted using custom-made C18 stagetips (Rappsilber *et

al*, 2007). The peptides were separated on an Ultimate 3000 RSLC nano-flow

chromatography system (Thermo Fisher Scientific), using a pre-column for sample

loading (PepMapAcclaim C18, 2 cm × 0.1 mm, 5 μm) and a C18 analytical column

(PepMapAcclaim C18, 50 cm × 0.75 mm, 2 μm, both Dionex-Thermo Fisher Scientific),

applying a linear gradient from 2 to 35% solvent B (80% acetonitrile, 0.08% FA acid;

solvent A: 0.1% FA) at a flow rate of 230 nl/min over 60 min. Eluting peptides were

analysed on a Q Exactive HF or HF-X Orbitrap mass spectrometer equipped with a

Proxeon nanospray source (Thermo Fisher Scientific), operated in data-dependent

mode. Survey scans were obtained in a scan range of 375–1500 m/z, at a resolution

of 60000 at 200 m/z and an AGC target value of 3E6. The 8 most intense ions were

selected with an isolation width of 1.6 Da, fragmented in the HCD cell at 28% collision

energy and the spectra recorded at a target value of 1E5 and a resolution of 30000.

Peptides with a charge of +1 were excluded from fragmentation, the peptide match

and exclude isotope features were enabled and selected precursors were dynamically

excluded from repeated sampling. Raw data were processed using the MaxQuant

software package (version 1.6.0.16, http://www.maxquant.org/) (Cox & Mann, 2008)

and searched against a custom database containing the sequences of the Sgk3

constructs in the protein background of the expression system (either Spodoptera

spp. protein sequences available in Uniprot plus the Bombyx mori reference

proteome (www.uniprot.org) or the human proteome) plus the sequences of common

contaminants. The search was performed with full trypsin specificity and a maximum

of two missed cleavages, C-terminal cleavage to glutamate and aspartate for the GluC

digests or unspecific cleavage for pepsin digests. Carbamidomethylation of cysteine

residues was set as fixed, oxidation of methionine, protein N-terminal acetylation and

phosphorylation of serine, threonine and tyrosine as variable modifications—all

other parameters were set to default. Results were filtered at a false discovery rate of

10% at the peptide and protein level. Spectra of phosphorylated Sgk3 peptides were

validated manually. All proteomics data have been deposited in the PRIDE repository

(Perez-Riverol *et al*, 2019).

Size exclusion chromatography with multiangle light scattering (SEC-MALS)

80 μl of Sgk3 at 0.9 mg/ml was injected onto Superdex 200 Increase 10/300 GL

column (Cytiva) equilibrated in 20 mM Tris pH 8, 150 mM NaCl, 1% glycerol, 1 mM

TCEP. Separation was carried out using HPLC Agilent Technologies 1260 infinity with

a flow rate of 0.5 ml/min operated at room temperature. On-line light scattering of

690 nm laser was recorded on a miniDawn Treos detector (Wyatt Technology Corp.).

Shodex RI-101 (Shodex) refractive index detector was used to measure protein

concentration on-line. Data was analyzed using the ASTRA software (Wyatt

Technology Corp.).

Dynamic light scattering (DLS)

Autocorrelation curves of freshly thawed and centrifuged 7.5 µM Sgk3 in buffer

containing 20 mM Hepes pH 7.4, 150 mM NaCl, 1 % glycerol, 1 mM TCEP, 1 mM ATP,

2 mM $MgCl_2$, 100 µM Crosstide were recorded at 0 and 60 min of incubation at room

temperature. Measurements were performed on a DynaPro NanoStar instrument

(Wyatt Technology Corp.). Each measurement is an average of 10 acquisitions of 10 s

each.

Sucrose-loaded vesicles (SLVs)

Cholesterol, DOPC, DOPE, DOPS and liver PI were dissolved in chloroform, PI3P di C16

was dissolved in chloroform:methanol:water 1:2:0.8, PI(4,5)P_2 was dissolved in

chloroform:methanol:water 20:13:3. For liposomes with synthetic PI3P the lipids

were mixed in the following molar ratio: 20% cholesterol, 30 % DOPC, 15% DOPS,

35% DOPE. For liposomes containing PI3P, DOPC was substituted with 0-4% PI3P.

Liver PI liposomes contained 10% PI at the expense of DOPC. Lipid mixtures in

borosilicate glass tubes were evaporated with nitrogen to form a thin film and then

rehydrated by vortexing for 1 min with sucrose buffer (20 mM Hepes pH 7.4, 300 mM

sucrose). Resuspended lipids were subjected to 4 cycles of freezing in LN2 and

thawing in a sonicator bath. One volume of gel filtration buffer was added and

liposomes were pelleted in Beckman Coulter Optima MAX-XP Ultracentrifuge in TLA

100 rotor operated at 50 000 rpm/20°C for 30 min. The supernatant was discarded,

liposome pellets were resuspended in gel filtration buffer to a final lipid

concentration of 1 mM and stored for a maximum of two days at 4°C before use.

Radiometric kinase assays

Sgk3 kinase activity was monitored as the incorporation of radioactive phosphate

from γ^{32}P-ATP (Hartmann Analytic) into Crosstide (GRPRTSSFAEG), a validated Sgk3

substrate peptide. Sgk3 was mixed with PI3P-containing liposomes and incubated for

30 min at RT. 5 µl of liposomes-Sgk3 mixture was added to 5 ul of substrate master

mix to obtain following conditions: 20 mM Tris pH 8, 150 mM NaCl, 1% glycerol, 1

mM TCEP, 50 nM Sgk3, 100 µM Crosstide, 1 mM ATP (spiked 1:10 with γ^{32}P-ATP), 2

mM MgCl$_2$. The reaction was left to proceed for 15 min at room temperature and

stopped by addition of 2 µl of 0.5 M EDTA. 10 ul of the reaction was then spotted onto

a nitrocellulose membrane (0.2 um pore size), washed 5 x 3 min with 75 mM H$_3$PO$_4$

and the washed membrane inserted into a 20 ml polyethylene scintillation vial

containing 5 ml deionized H$_2$O. Cerenkov radiation was measured in a Tri-Carb 4910

TR liquid scintillator (Perkin Elmer). Datapoints were fitted with Michaelis-Menten

equation modified with a constant to reflect the basal enzymatic activity in 0% PI3P

condition:

559
$$V = \frac{V_{max}[PI3P]}{k_1 + [PI3P]} + const$$

Liposome pelleting assay

1 µM Sgk3^{WT} or Sgk3^{PX} was mixed 1:1 with liposomes, incubated at 20°C for 30 min

and spun at 9800g for 30 min at 20°C. Supernatant was collected and the liposome

pellet resuspended in gel filtration buffer. Equal volumes of supernatant and pellet

fractions were separated by SDS-PAGE and quantified by Coomassie densitometry. In

the comparative binding assay with full-length Sgk3 and Sgk3^{PX} in the same reaction,

FITC-labelled PX domain was quantified by fluorescence at 448 nm. Binding curves

were fitted with a one-site binding model.

Vps34-Sgk3 coupled kinase assay

To monitor PI3P production by Vps34 and Crosstide phosphorylation by Sgk3 in a

single reaction, the components were mixed at following concentrations: 50 nM

Vps34 complex, 50 nM Sgk3, 500 µM Biotin-Crosstide, 0.5 mM ATP (spiked 1:10 with

γ^{32}P-ATP), 0.5 mM MgCl$_2$, 2 mM MnCl$_2$, 1 mM EGTA, 10% liver PI liposomes. The

reaction was carried out at 25°C and gentle shaking. 15 µl samples were mixed with

5 µl 4x quench buffer (20 mM Tris pH 7.4, 150 mM NaCl, 10 mM EDTA, 10 mM EGTA,

1% glycerol, 1 mM TCEP) at indicated time points. The quenched reactions were spun

for 5 min at 20800 g/20°C. The supernatant was mixed 1:1 with NeutrAvidin (Thermo

Scientific) at 2 mg/ml in water to sequester the biotinylated Crosstide and 5 ul were

spotted onto 0.45 um nitrocellulose membrane. The membrane was washed 5x3 min

with TBS (20 mM Tris pH 7.6, 137 mM NaCl), briefly dried, exposed in storage

phosphor screen overnight and imaged with an Amersham Typhoon

phosphorimager. The intensity of the spots was quantified in ImageJ. The pellets

containing liposomes with PI3^{32}P were washed 4 times with wash buffer (20 mM Tris

pH 7.4, 150 mM NaCl, 2.5 mM EDTA, 2.5 mM EGTA, 1% glycerol, 1 mM TCEP),

transferred into PCR tube and inserted into 20 ml polyethylene scintillation vials

containing 5 ml MilliQ water. PI3P synthesis was measured by liquid scintillation.

HDX-MS

Lipid Vesicle Preparation

Lipid films containing 20% cholesterol, 25% DOPC, 15% DOPS, 35% DOPE, and 5%

PI3P were dried and stored under argon for shipping. The dried films were

resuspended to 5 mM in gel filtration buffer (20 mM Tris pH 7.4, 150 mM NaCl, 1%

glycerol, 1 mM TCEP) by vortexing. The resuspended lipid vesicle solutions were

sonicated for 10 minutes, subjected to ten freeze-thaw cycles, and sonicated again for

5 minutes. The vesicles were then snap-frozen in liquid nitrogen and stored at -80°C

HDX-MS sample preparation

HDX reactions were conducted in a final reaction volume of 25 µL with a final Sgk3

concentration of 13 µM (for both FL and PX Sgk3). Prior to the addition of deuterated

solvent, 1.5 µL of FL or PX Sgk3 was allowed to incubate with either 2 µL of 5 mM 0%

PI3P vesicles or 2 µL 5 mM 5% PI3P vescles. After the two minute incubation period,

21.5 µL D2O buffer (20mM pH 7 HEPES, 100mM NaCl, 0.5 mM TCEP, 94% D2O) was

added with a final %D2O of 81.5% (v/v). The reaction was allowed to proceed for 3s,

30s, 300s, or 3000s at 18°C before being quenched with ice cold acidic quench buffer.

All conditions and timepoints were created and run in triplicate. Samples were flash

frozen in liquid nitrogen and stored at -80°C until injection onto the ultra-

performance liquid chromatography system for proteolytic cleavage, peptide

separation, and injection onto a QTOF for mass analysis.

Protein digestion and MS/MS data collection

Protein samples were rapidly thawed and injected onto an integrated fluidics system

containing a HDx-3 PAL liquid handling robot and climate-controlled

chromatography system (LEAP Technologies), a Dionex Ultimate 3000 UHPLC

system, as well as an Impact HD QTOF Mass spectrometer (Bruker). The protein was

run over either one immobilized pepsin column at at 10°C (Trajan; ProDx protease

column, 2.1 mm x 30 mm PDX.PP01-F32) at 200 mL/min for 3 minutes. The resulting

peptides were collected and desalted on a C18 trap column (Acquity UPLC BEH C18

1.7mm column (2.1 x 5 mm); Waters 186003975). The trap was subsequently eluted

in line with a C18 reverse-phase separation column (Acquity 1.7 mm particle, 100 x 1

mm^2 C18 UPLC column, Waters 186002352), using a gradient of 5-36% B (Buffer A

0.1% formic acid; Buffer B 100% acetonitrile) over 16 minutes. Mass spectrometry

experiments acquired over a mass range from 150 to 2200 m/z using an electrospray

ionization source operated at a temperature of 200C and a spray voltage of 4.5 kV.

Peptide identification

Peptides were identified from the non-deuterated samples of FL and PX Sgk3 using

data-dependent acquisition following tandem MS/MS experiments (0.5 s precursor

scan from 150-2000 m/z; twelve 0.25 s fragment scans from 150-2000 m/z). MS/MS

datasets were analysed using PEAKS7 (PEAKS), and peptide identification was carried out by using a false discovery based approach, with a threshold set to 1% using a database of purified proteins and known contaminants found in Sf9 cells (Dobbs *et al*, 2020). The search parameters were set with a precursor tolerance of 20 ppm, fragment mass error 0.02 Da, charge states from 1-8, leading to a selection criterion of peptides that had a -10logP score of 21.4.

Mass Analysis of Peptide Centroids and Measurement of Deuterium Incorporation

HD-Examiner Software (Sierra Analytics) was used to automatically calculate the level of deuterium incorporation into each peptide. All peptides were manually inspected for correct charge state, correct retention time, appropriate selection of isotopic distribution, etc. Deuteration levels were calculated using the centroid of the experimental isotope clusters. Results are presented as relative levels of deuterium incorporation and the only control for back exchange was the level of deuterium present in the buffer (81.5%). Differences in exchange in a peptide were considered significant if they met all three of the following criteria: : $\geq$7% change in exchange, $\geq$0.5 Da difference in exchange, and a p value <0.01 using a two tailed student t-test. All compared samples were set within the same experiment. To allow for visualization of differences across all peptides, we utilized number of deuteron difference (#D) plots. These plots show the total difference in deuterium incorporation over the entire H/D exchange time course, with each point indicating a single peptide. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository (Perez-Riverol *et al*,

2019) with the dataset identifier PXD025327. The data analysis statistics for all HDX-

MS experiments are in Supplementary Table 1 according to the guidelines of Masson

et al (Masson *et al*, 2019).

References

Alessi DR, Deak M, Casamayor A, Caudwell FB, Morrice N, Norman DG, Gaffney P,

Reese CB, MacDougall CN, Harbison D, *et al* (1997) 3-Phosphoinositide-

dependent protein kinase-1 (PDK1): structural and functional homology with

the Drosophila DSTPK61 kinase. *Curr Biol* 7: 776–89

Backer JM (2008) The regulation and function of Class III PI3Ks: novel roles for

Vps34. *Biochem J* 410: 1–17

Backer JM (2016) The intricate regulation and complex functions of the Class III

phosphoinositide 3-kinase Vps34. *Biochem J* 473: 2251–2271

Bago R, Malik N, Munson MJ, Prescott AR, Davies P, Sommer E, Shpiro N, Ward R,

Cross D, Ganley IG, *et al* (2014) Characterization of VPS34-IN1, a selective

inhibitor of Vps34, reveals that the phosphatidylinositol 3-phosphate-binding

SGK3 protein kinase is a downstream target of class III phosphoinositide 3-

kinase. *Biochem J* 463: 413–27

Bago R, Sommer E, Castel P, Crafter C, Bailey FP, Shpiro N, Baselga J, Cross D, Eyers

PA & Alessi DR (2016) The hVps34-SGK3 pathway alleviates sustained

PI3K/Akt inhibition by stimulating mTORC1 and tumour growth. *EMBO J* 35:

1902–22

Braccini L, Ciraolo E, Campa CC, Perino A, Longo DL, Tibolla G, Pregnolato M, Cao Y,

Tassone B, Damilano F, *et al* (2015) PI3K-C2γ is a Rab5 effector selectively

controlling endosomal Akt2 activation downstream of insulin signalling. *Nat*

Commun 6: 7400

Brunet A, Park J, Tran H, Hu LS, Hemmings BA & Greenberg ME (2001) Protein

kinase SGK mediates survival signals by phosphorylating the forkhead

transcription factor FKHRL1 (FOXO3a). *Mol Cell Biol* 21: 952–65

Carlton JG & Cullen PJ (2005) Coincidence detection in phosphoinositide signaling.

Trends Cell Biol 15: 540–547

Chan TO, Zhang J, Rodeck U, Pascal JM, Armen RS, Spring M, Dumitru CD, Myers V, Li

X, Cheung JY, *et al* (2011) Resistance of Akt kinases to dephosphorylation

through ATP-dependent conformational plasticity. *Proc Natl Acad Sci U S A* 108:

E1120-7

Chandra M, Chin YK-Y, Mas C, Feathers JR, Paul B, Datta S, Chen K-E, Jia X, Yang Z,

Norwood SJ, *et al* (2019) Classification of the human phox homology (PX)

domains based on their phosphoinositide binding specificities. *Nat Commun* 10:

1528

Chen W, Chen Y, Xu B, Juang Y-C, Stippec S, Zhao Y & Cobb MH (2009) Regulation of

a third conserved phosphorylation site in SGK1. *J Biol Chem* 284: 3453–60

Cox J & Mann M (2008) MaxQuant enables high peptide identification rates,

individualized p.p.b.-range mass accuracies and proteome-wide protein

quantification. *Nat Biotechnol* 26: 1367–72

Currie RA, Walker KS, Gray A, Deak M, Casamayor A, Downes CP, Cohen P, Alessi DR

& Lucocq J (1999) Role of phosphatidylinositol 3,4,5-trisphosphate in

regulating the activity and localization of 3-phosphoinositide-dependent

protein kinase-1. *Biochem J* 337 (Pt 3: 575–83

Dai F, Yu L, He H, Chen Y, Yu J, Yang Y, Xu Y, Ling W & Zhao S Human serum and

glucocorticoid-inducible kinase-like kinase (SGKL) phosphorylates glycogen

syntheses kinase 3 beta (GSK-3b) at serine-9 through direct interaction

Dobbs JM, Jenkins ML & Burke JE (2020) Escherichia coli and Sf9 Contaminant

Databases to Increase Efficiency of Tandem Mass Spectrometry Peptide

Identification in Structural Mass Spectrometry Experiments. *J Am Soc Mass*

Spectrom 31: 2202–2209

Ebner M, Lučić I, Leonard TA & Yudushkin I (2017) PI(3,4,5)P3 Engagement

Restricts Akt Activity to Cellular Membranes. *Mol Cell* 65: 416-431.e6

Elsner DJ, Siess KM, Gossenreiter T, Hartl M & Leonard TA (2019) A ubiquitin-like

domain controls protein kinase D dimerization and activation by trans-

autophosphorylation. *J Biol Chem* 294: 14422–14441

Endicott JA, Noble MEM & Johnson LN (2012) The structural basis for control of

eukaryotic protein kinases. *Annu Rev Biochem* 81: 587–613

Facchinetti V, Ouyang W, Wei H, Soto N, Lazorchak A, Gould C, Lowry C, Newton AC,

Mao Y, Miao RQ, *et al* (2008) The mammalian target of rapamycin complex 2

controls folding and stability of Akt and protein kinase C. *EMBO J* 27: 1932–43

Fruman DA, Chiu H, Hopkins BD, Bagrodia S, Cantley LC & Abraham RT (2017) The

PI3K Pathway in Human Disease. *Cell* 170: 605–635

Fruman DA & Rommel C (2014) PI3K and cancer: lessons, challenges and

opportunities. *Nat Rev Drug Discov* 13: 140–156

Gasser JA, Inuzuka H, Lau AW, Wei W, Beroukhim R & Toker A (2014) SGK3

mediates INPP4B-dependent PI3K signaling in breast cancer. *Mol Cell* 56: 595–

607

Goulden BD, Pacheco J, Dull A, Zewe JP, Deiters A & Hammond GR V (2019) A high-

avidity biosensor reveals plasma membrane PI(3,4)P2 is predominantly a class

I PI3K signaling product. *J Cell Biol* 218: 1066–1079

Hastie CJ, McLauchlan HJ & Cohen P (2006) Assay of protein kinases using

radiolabeled ATP: A protocol. *Nat Protoc* 1: 968–971

Hauge C, Antal TL, Hirschberg D, Doehn U, Thorup K, Idrissova L, Hansen K, Jensen

ON, Jørgensen TJ, Biondi RM, *et al* (2007) Mechanism for activation of the

growth factor-activated AGC kinases by turn motif phosphorylation. *EMBO J* 26:

2251–2261

Hornbeck P V., Chabra I, Kornhauser JM, Skrzypek E & Zhang B (2004) PhosphoSite:

A bioinformatics resource dedicated to physiological protein phosphorylation.

Proteomics 4: 1551–61

Hurley JH & Young LN (2017) Mechanisms of Autophagy Initiation. *Annu Rev*

Biochem 86: 225–244

Ikenoue T, Inoki K, Yang Q, Zhou X & Guan K-L (2008) Essential function of TORC2 in

PKC and Akt turn motif phosphorylation, maturation and signalling. *EMBO J* 27:

1919–1931

Ivetac I, Munday AD, Kisseleva M V., Zhang X-M, Luff S, Tiganis T, Whisstock JC, Rowe

T, Majerus PW & Mitchell CA (2005) The type Ialpha inositol polyphosphate 4-

phosphatase generates and terminates phosphoinositide 3-kinase signals on

endosomes and the plasma membrane. *Mol Biol Cell* 16: 2218–33

Kobayashi T & Cohen P (1999) Activation of serum- and glucocorticoid-regulated

protein kinase by agonists that activate phosphatidylinositide 3-kinase is

mediated by 3-phosphoinositide-dependent protein kinase-1 (PDK1) and

PDK2. *Biochem J* 339 (Pt 2: 319–28

Kobayashi T, Deak M, Morrice N & Cohen P (1999) Characterization of the structure

and regulation of two novel isoforms of serum- and glucocorticoid-induced

protein kinase. *Biochem J* 344 Pt 1: 189–97

Landel I, Quambusch L, Depta L & Rauh D (2020) Spotlight on AKT: Current

Therapeutic Challenges. *ACS Med Chem Lett* 11: 225–227

Lang F, Bohmer C, Palmada M, Seebohm G, Strutz-Seebohm N & Vallon V (2006)

(Patho)physiological Significance of the Serum- and Glucocorticoid-Inducible

Kinase Isoforms. *Physiol Rev* 86: 1151–1178

Lapierre J-M, Eathiraj S, Vensel D, Liu Y, Bull CO, Cornell-Kennon S, Iimura S,

Kelleher EW, Kizer DE, Koerner S, *et al* (2016) Discovery of 3-(3-(4-(1-

Aminocyclobutyl)phenyl)-5-phenyl-3H-imidazo[4,5-b]pyridin-2-yl)pyridin-2-

amine (ARQ 092): An Orally Bioavailable, Selective, and Potent Allosteric AKT

Inhibitor. *J Med Chem* 59: 6455–69

Leonard TA & Hurley JH (2011) Regulation of protein kinases by lipids. *Curr Opin*

Struct Biol 21: 785–791

Leonard TA, Róycki B, Saidi LF, Hummer G & Hurley JH (2011) Crystal structure and

allosteric activation of protein kinase C βiI. *Cell* 144: 55–66

Lin K, Lin J, Wu W-I, Ballard J, Lee BB, Gloor SL, Vigers GPA, Morales TH, Friedman

LS, Skelton N, *et al* (2012) An ATP-site on-off switch that restricts phosphatase

accessibility of Akt. *Sci Signal* 5: ra37

Liu D, Yang X & Songyang Z (2000) Identification of CISK, a new member of the SGK

kinase family that promotes IL-3-dependent survival. *Curr Biol* 10: 1233–6

Liu S-L, Wang Z-G, Hu Y, Xin Y, Singaram I, Gorai S, Zhou X, Shim Y, Min J-H, Gong L-

W, *et al* (2018) Quantitative Lipid Imaging Reveals a New Signaling Function of

Phosphatidylinositol-3,4-Bisphophate: Isoform- and Site-Specific Activation of

Akt. *Mol Cell* 71: 1092-1104.e5

Lu S, Deng R, Jiang H, Song H, Li S, Shen Q, Huang W, Nussinov R, Yu J & Zhang J

(2015) The Mechanism of ATP-Dependent Allosteric Protection of Akt Kinase

Phosphorylation. *Structure* 23: 1725–1734

Lučić I, Rathinaswamy MK, Truebestein L, Hamelin DJ, Burke JE & Leonard TA

(2018) Conformational sampling of membranes by Akt controls its activation

and inactivation. *Proc Natl Acad Sci U S A* 115: E3940–E3949

Lučić I, Truebestein L & Leonard TA (2016) Novel Features of DAG-Activated PKC

Isozymes Reveal a Conserved 3-D Architecture. *J Mol Biol* 428: 121–141

Malik N, Macartney T, Hornberger A, Anderson KE, Tovell H, Prescott AR & Alessi DR

(2018) Mechanism of activation of SGK3 by growth factors via the Class 1 and

Class 3 PI3Ks. *Biochem J* 475: 117–135

Malik N, Nirujogi RS, Peltier J, Macartney T, Wightman M, Prescott AR, Gourlay R,

Trost M, Alessi DR & Karapetsas A (2019) Phosphoproteomics reveals that the

hVPS34 regulated SGK3 kinase specifically phosphorylates endosomal proteins

including Syntaxin-7, Syntaxin-12, RFIP4 and WDR44. *Biochem J* 476: 3081–

3107

Manning G (2002) The Protein Kinase Complement of the Human Genome. *Science*

(80-) 298: 1912–1934

Marone R, Cmiljanovic V, Giese B & Wymann MP (2008) Targeting phosphoinositide

3-kinase-Moving towards therapy. *Biochim Biophys Acta - Proteins Proteomics*

1784: 159–185 doi:10.1016/j.bbapap.2007.10.003 [PREPRINT]

Martini M, De Santis MC, Braccini L, Gulluni F & Hirsch E (2014) PI3K/AKT signaling

pathway and cancer: an updated review. *Ann Med* 46: 372–383

Masson GR, Burke JE, Ahn NG, Anand GS, Borchers C, Brier S, Bou-Assaf GM, Engen

JR, Englander SW, Faber J, *et al* (2019) Recommendations for performing,

interpreting and reporting hydrogen deuterium exchange mass spectrometry

(HDX-MS) experiments. *Nat Methods* 16: 595–602

Murray JT, Cummings LA, Bloomberg GB & Cohen P (2005) Identification of different

specificity requirements between SGK1 and PKBα. *FEBS Lett* 579: 991–994

Oancea E & Meyer T (1998) Protein kinase C as a molecular machine for decoding

calcium and diacylglycerol signals. *Cell* 95: 307–18

Oh WJ, Wu C, Kim SJ, Facchinetti V, Julien L-A, Finlan M, Roux PP, Su B & Jacinto E

(2010) mTORC2 can associate with ribosomes to promote cotranslational

phosphorylation and stability of nascent Akt polypeptide. *EMBO J* 29: 3939–

3951

Pao AC, McCormick JA, Li H, Siu J, Govaerts C, Bhalla V, Soundararajan R & Pearce D

(2007) NH 2 terminus of serum and glucocorticoid-regulated kinase 1 binds to

phosphoinositides and is essential for isoform-specific physiological functions.

Am J Physiol Physiol 292: F1741–F1750

Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ,

Inuganti A, Griss J, Mayer G, Eisenacher M, *et al* (2019) The PRIDE database and

related tools and resources in 2019: improving support for quantification data.

Nucleic Acids Res 47: D442–D450

Rappsilber J, Mann M & Ishihama Y (2007) Protocol for micro-purification,

enrichment, pre-fractionation and storage of peptides for proteomics using

StageTips. *Nat Protoc* 2: 1896–906

Sawa-Makarska J, Baumann V, Coudevylle N, von Bülow S, Nogellova V, Abert C,

Schuschnig M, Graef M, Hummer G & Martens S (2020) Reconstition of

autophagosome nucleation defines Atg9 vesicles as seeds for membrane

formation. *Science* 369

Schenck A, Goto-Silva L, Collinet C, Rhinn M, Giner A, Habermann B, Brand M &

Zerial M (2008) The endosomal protein Appl1 mediates Akt substrate

specificity and cell survival in vertebrate development. *Cell* 133: 486–97

Siess KM & Leonard TA (2019) Lipid-dependent Akt-ivity: where, when, and how.

Biochem Soc Trans 47: 897–908

Slagsvold T, Marchese A, Brech A & Stenmark H (2006) CISK attenuates degradation

of the chemokine receptor CXCR4 via the ubiquitin ligase AIP4. *EMBO J* 25:

3738–49

Song Y, DiMaio F, Yu-Ruei Wang R, Kim D, Miles C, Brunette T, Thompson J & Baker

D Technical Advance High-Resolution Comparative Modeling with RosettaCM.

Stokoe D, Stephens LR, Copeland T, Gaffney PR, Reese CB, Painter GF, Holmes AB,

McCormick F & Hawkins PT (1997) Dual role of phosphatidylinositol-3,4,5-

trisphosphate in the activation of protein kinase B. *Science* 277: 567–70

Tessier M & Woodgett JR (2006) Role of the Phox homology domain and

phosphorylation in activation of serum and glucocorticoid-regulated kinase-3. *J*

Biol Chem 281: 23978–89

Theile CS, Witte MD, Blom AEM, Kundrat L, Ploegh HL & Guimaraes CP (2013) Site-

specific N-terminal labeling of proteins using sortase-mediated reactions. *Nat*

Protoc 8: 1800–1807

Tovell H, Testa A, Zhou H, Shpiro N, Crafter C, Ciulli A & Alessi DR (2019) Design and

Characterization of SGK3-PROTAC1, an Isoform Specific SGK3 Kinase PROTAC

Degrader. *ACS Chem Biol*: acschembio.9b00505

Vasudevan KM, Barbie DA, Davies MA, Rabinovsky R, McNear CJ, Kim JJ, Hennessy

BT, Tseng H, Pochanard P, Kim SY, *et al* (2009) AKT-independent signaling

downstream of oncogenic PIK3CA mutations in human cancer. *Cancer Cell* 16:

21–32

Virbasius J V, Song X, Pomerleau DP, Zhan Y, Zhou GW & Czech MP (2001) Activation

of the Akt-related cytokine-independent survival kinase requires interaction of

its phox domain with endosomal phosphatidylinositol 3-phosphate. *Proc Natl*

Acad Sci U S A 98: 12908–13

Wang Y, Zhou D, Phung S, Warden C, Rashid R, Chan N & Chen S (2017) SGK3

sustains ERα signaling and drives acquired aromatase inhibitor resistance

through maintaining endoplasmic reticulum homeostasis. *Proc Natl Acad Sci U S*

A 114: E1500–E1508

Webster MK, Goya L, Ge Y, Maiyar AC & Firestone GL (1993) Characterization of sgk,

a novel member of the serine/threonine protein kinase gene family which is

transcriptionally induced by glucocorticoids and serum. *Mol Cell Biol* 13: 2031–

2040

Weisner J, Landel I, Reintjes C, Uhlenbrock N, Trajkovic-Arsic M, Dienstbier N,

Hardick J, Ladigan S, Lindemann M, Smith S, *et al* (2019) Preclinical Efficacy of

Covalent-Allosteric AKT Inhibitor Borussertib in Combination with Trametinib

in KRAS-mutant Pancreatic and Colorectal Cancer. *Cancer Res* 79:

canres.2861.2018

Wu W-I, Voegtli WC, Sturgis HL, Dizon FP, Vigers GPA & Brandhuber BJ (2010)

Crystal structure of human AKT1 with an allosteric inhibitor reveals a new

mode of kinase inhibition. *PLoS One* 5: e12913

Xing Y, Liu D, Zhang R, Joachimiak A, Songyang Z & Xu W (2004) Structural basis of

membrane targeting by the Phox homology domain of cytokine-independent

survival kinase (CISK-PX). *J Biol Chem* 279: 30662–9

Xu J, Liu D, Gill G & Songyang Z (2001) Regulation of cytokine-independent survival

kinase (CISK) by the phox homology domain and phosphoinositides. *J Cell Biol*

154: 699–705

Figure 1. Sgk3 localizes specifically to PI3P-containing endosomes

A) HeLa cells were transfected with mCherry-Sgk3^{WT} and serum starved

overnight.

B) Starved cells were treated with 100 nM wortmannin for 20 min.

C-E) Cells were co-transfected with mCherry-Sgk3^{WT} and EGFP-Sgk3^{R90A} mutant.

F) Fluorescence intensity profile of mCherry Sgk3^{WT} and EGFP-Sgk3^{R90A} along the

white line shown in C-E.

Figure 2. Sgk3 is specifically and allosterically activated by PI3P

A) Sgk3^{WT} kinase activity was assayed in the presence (closed black square) or

absence (closed red square) of 100 μM Crosstide substrate with 0-4 mol %

PI3P liposomes, or with 4 mol % PI(4,5)P$_2$ liposomes (open black square) as a

negative control. In parallel, binding of Sgk3 to 0-4 % PI3P liposomes (full blue

circles) or 4% PI(4,5)P$_2$ (open blue circle) as a negative control was measured.

Dotted black line indicates basal Sgk3^{WT} kinase activity of unbound protein,

dotted red line indicates no-substrate control. Kinase activity was read out

radiometrically by liquid scintillation and liposome binding by Coomassie

densitometry. Error bars are the standard deviation of three independent

experiments.

B) Same as A) with Sgk3^{D286A} kinase-dead mutant.

C) Schematic of Vps34-Sgk3 coupled kinase assay. Phosphorylation of Crosstide

was assayed by phosphorimaging. PI3P synthesis was measured by liquid

scintillation.

D) Vps34-Sgk3 coupled kinase assay. Crosstide phosphorylation was measured

in the presence (full blue line) or absence (dashed blue line) of 50 nM Vps34.

At the same time, PI3P synthesis was measured in the presence (full green

line) or absence (dashed green line) of 50 nM Vps34. Error bars are the

standard deviation of three independent experiments.

Figure 3. Sgk3 is autoinhibited by its PX domain.

A) Comparison of deuterium incorporation in Sgk3 PX domain with 0% vs 5%

PI3P liposomes. Plot: difference in number of deuterons incorporated as a

function of peptide centroid over the PX domain. Negative numbers indicate

protection (blue). Peptides showing significant deuterium exchange

differences (>7%, >0.5 Da, and p<0.01) were mapped on the structure of the

human Sgk3 PX domain (PDB:6EDX). The binding site for PI3P is indicated by

superimposing the structure of the PX domain of p40phox in complex with

PI3P (PDB: 1H6H) and rendering the PI3P in spheres.

B) Comparison of deuterium incorporation in Sgk3^{WT} vs. Sgk3 PX domain in the

presence of 0% PI3P liposomes (unbound). Plot: difference in number of

deuterons incorporated as a function of peptide centroid over the PX domain.

Positive numbers indicate domain exposure (red).

C) Comparison of deuterium incorporation in Sgk3^{WT} unbound (0% PI3P) vs

bound (5% PI3P) for the PX domain (left) and the kinase domain (right). A

homology model of Sgk3 kinase domain (Rosetta) (Song *et al*) was used to map

the changes. Color coding expressing the magnitude of changes in deuterium

incorporation is on bottom right. All measurements were done in triplicates,

error bars are the standard deviations.

Figure 4. Autoinhibition of Sgk3 impairs binding to PI3P

A) Binding of Sgk3^{WT} full-length (blue squares) and FITC-labelled Sgk3 PX

domain (black squares) to 0-4 mol % PI3P liposomes was monitored in single

reaction using fluorescence (FITX-PX) or Coomassie densitometry (full-length

Sgk3) in 15% SDS-PAGE gels. Curves were fitted with a one binding site model.

Error bars are the standard deviations of three independent experiments.

B-D) Distribution of EGFP-PX domain (B) or mCherry-Sgk3^{WT} (full-length) (C) in

serum-starved HeLa cells.

E) Quantification of fluorescence intensity of both constructs along the white line

shown in images B-D.

Figure 5. Model for the allosteric activation of Sgk3 by PI3P

Cytosolic Sgk3 is allosterically inhibited by its PX domain, which is relieved by

PI3P binding (1). This is followed by phosphorylation of the Sgk3 activation loop

by PDK1 as a consequence of its association with endosomal $PI(3,4)P_2$ (2).

Phosphorylation activates Sgk3 (3), thereby permitting downstream substrate

phosphorylation (4).

Figure 1. Sgk3 localizes specifically to PI3P-containing endosomes.

Figure 2. Sgk3 is specifically and allosterically activated by PI3P.

Figure 3. Sgk3 is autoinhibited by its PX domain.

Figure 4. Autoinhibition of Sgk3 impairs binding to PI3P.

Figure 5. Model for the allosteric activation of Sgk3 by PI3P.