Divergent Acyl Carrier Protein Mediates Mitochondrial Fe-S Cluster Biosynthesis in Malaria Parasites

Seyi Falekun[1], Jaime Sepulveda[1], Yasaman Jami-Alahmadi[2], Hahnbeom Park[3], James A. Wohlschlegel[2], and Paul A. Sigala[1*]

[1]Department of Biochemistry, University of Utah School of Medicine, Salt Lake City, UT, 84112, [2]Department of Biological Chemistry, University of California, Los Angeles, Los Angeles, CA, 90095, [3]Department of Biochemistry, University of Washington, Seattle, WA, 98195

[*]Corresponding author: Paul A. Sigala
University of Utah, Dept. of Biochemistry
15 N. Medical Dr. East, Room 4100
Salt Lake City, UT 84112
801-587-4874
p.sigala@biochem.utah.edu

Preprint server:

Classification: Biological Sciences, Biochemistry

Keywords: malaria, Plasmodium, mitochondria, acyl carrier protein, metabolic adaption, iron-sulfur clusters

Abstract

Plasmodium falciparum malaria parasites are early-diverging eukaryotes with many unusual metabolic adaptations. Understanding these adaptations will give insight into parasite evolution and unveil new parasite-specific drug targets. In contrast to human cells, the *Plasmodium* mitochondrion lacks type II fatty acid biosynthesis (FASII) enzymes yet curiously retains a divergent acyl carrier protein (mACP) incapable of modification by a 4-phosphopantetheine (Ppant) group required for canonical ACP function as the scaffold for fatty acid synthesis. We report that ligand-dependent knockdown of mACP expression is lethal to parasites, indicating an essential FASII-independent function. Decyl-ubiquinone rescues parasites temporarily from this lethal phenotype, suggesting a dominant dysfunction of the mitochondrial electron transport chain (ETC) followed by broader defects beyond the ETC. Biochemical studies reveal that *Plasmodium* mACP binds and stabilizes the Isd11-Nfs1 cysteine desulfurase complex required for Fe-S cluster biosynthesis, and mACP knockdown causes loss of both Nfs1 and the Rieske Fe-S cluster protein in ETC Complex III. This work identifies Ppant-independent mACP as an essential mitochondrial adaptation in *Plasmodium* malaria parasites that appears to be a shared metabolic feature of *Apicomplexan* pathogens, including *Toxoplasma* and *Babesia*. This parasite-specific adaptation highlights the ancient, fundamental role of ACP in mitochondrial Fe-S cluster biogenesis and reveals an evolutionary driving force to retain this interaction with ACP independent of its eponymous function in fatty acid synthesis.

Significance Statement

Plasmodium malaria parasites are single-celled eukaryotes that evolved unusual metabolic adaptations. Parasites require a mitochondrion for blood-stage viability, but essential functions beyond the electron transport chain are sparsely understood. Unlike human cells, the *Plasmodium* mitochondrion lacks fatty acid synthesis enzymes but retains a divergent acyl carrier protein (mACP) incapable of tethering acyl groups. Nevertheless, mACP is essential for parasite viability by binding and stabilizing the core mitochondrial Fe-S cluster biogenesis complex via a divergent molecular interface lacking an acyl-pantetheine group that contrasts with humans. This discovery unveils an essential mitochondrial adaptation in *Plasmodium* shared by other human-infecting parasites and suggests a primordial ACP function in Fe-S cluster biogenesis independent of its canonical role as a scaffold for fatty acid synthesis.

Malaria is an ancient scourge of humanity and remains a pressing global health challenge, especially in tropical Africa where hundreds of thousands of people die from malaria each year. *Plasmodium falciparum* malaria parasites are single-celled eukaryotes that evolved under unique selective pressures with unusual metabolic adaptations compared to human cells and well-studied model organisms such as yeast. Understanding the unique biochemical pathways that specialize parasites for growth within human red blood cells will shed light on their evolutionary divergence from other eukaryotes and unveil new parasite-specific targets for development of novel antimalarial therapies.

P. falciparum retains an essential mitochondrion required for biosynthesis of pyrimidines, acetyl-CoA, and Fe-S clusters needed outside the organelle (1-4). Although the parasite mitochondrion also contains enzymes involved in the citric acid (TCA) cycle and biosynthesis of ATP and heme, these pathways are dispensable for blood-stage parasites, which can scavenge host heme and obtain sufficient ATP from cytoplasmic glycolysis (5-9). In addition to these pathways, most eukaryotes, including mammals and yeast, also contain a mitochondrial type-II fatty acid biosynthesis (FASII) pathway that generates the octanoate precursor of the lipoic acid cofactor used by several mitochondrial dehydrogenases (10). In contrast to human and yeast cells, FASII enzymes in *P. falciparum* are absent from the mitochondrion and targeted instead to the apicoplast organelle (Table S1) (11). Although critical lipoate-dependent enzymes are present in the parasite mitochondrion (2), prior work has shown that these enzymes utilize scavenged lipoate obtained from the red blood cell rather than de novo synthesis (12).

The acyl carrier protein (ACP) is a key component of FASII and contains a strictly conserved Ser residue modified by a 4-phosphopantetheine (4-Ppant) group whose terminal thiol tethers the nascent acyl chain during fatty acid initiation, modification, and elongation (13).

Consistent with FASII targeting to the apicoplast in *P. falciparum*, this organelle features a well-studied ACP homolog (PF3D7_0208500, aACP) that retains canonical ACP features, including the conserved Ser modified by a 4-Ppant group required for FASII function (11, 14). Although apicoplast FASII activity is essential for *P. falciparum* growth within mosquitoes and the human liver, this pathway is dispensable for blood-stage parasites which can scavenge short-chain fatty acids from the host (11, 15, 16).

Despite loss of mitochondrial FASII enzymes, *P. falciparum* has curiously retained a second ACP homolog (PF3D7_1208300) that is annotated as a mitochondrial ACP (mACP) but has not been directly studied. We noted that recent genome-wide knock-out studies in *P. berghei* and *P. falciparum* both reported that this gene was refractory to disruption, suggesting an essential, FASII-independent function (17, 18). We became interested to unravel what this essential function might be. Recent studies in yeast and human cells have identified an expanding network of mitochondrial ACP interactions beyond FASII that includes roles in respiratory chain assembly, Fe-S cluster biogenesis, and mitochondrial ribosomal translation (19-23). However, in all cases these critical interactions in yeast and/or humans are linked to mitochondrial FASII function and the Ppant group of ACP (22, 24-26).

We localized mACP to the *P. falciparum* mitochondrion and set out to understand its FASII-independent function in this organelle. Using conditional knockdown and immunoprecipitation studies, we discovered that mACP is essential for parasite viability and plays a critical role in binding and stabilizing the Isd11-Nfs1 cysteine desulfurase complex required for mitochondrial Fe-S cluster biogenesis. Unlike mACP interactions in yeast and humans, *P. falciparum* mACP binds to Isd11 via a divergent molecular interface that does not involve a 4-Ppant group. This work unveils a new molecular paradigm for essential mACP function without a

Ppant group, underscores the critical and conserved role of mACP in mitochondrial Fe-S cluster biogenesis, and highlights a *Plasmodium*-specific adaptation suitable for exploration as a metabolic vulnerability for parasite-directed antimalarial therapy.

RESULTS

Mitochondrial ACP is essential for *P. falciparum* despite loss of FASII in this organelle. The *P. falciparum* genome encodes two ACP homologs that include the well-studied protein targeted to the apicoplast (aACP, PF3D7_0208500) and a second homolog annotated as a mitochondrial ACP (mACP, PF3D7_1208300). Unlike the apicoplast ACP, which retains the conserved Ser for 4-Ppant attachment, mACP has curiously replaced this Ser with a Phe residue that cannot be modified by a 4-Ppant group (Fig. 1A and Fig. S1A). This Ser-to-Phe substitution in mACP is consistent with the loss of mitochondrial FASII and the ACP-modifying phosphopantetheine transferase enzyme (Table S1) and suggested a non-canonical function for mACP.

To test if mACP localized to the parasite mitochondrion, we created a Dd2 *P. falciparum* line that episomally expressed mACP fused to a C-terminal dual hemagglutinin (HA) tag. Immunofluorescence analysis of this mACP-HA$_2$ line revealed strong co-localization between mACP-HA$_2$ and the mitochondrial marker, HSP60 (Fig. 1B and Fig. S2). On the basis of this colocalization, we conclude that mACP is indeed targeted to the parasite mitochondrion, as predicted. Full-length mACP-HA$_2$ has a predicted molecular weight (MW) ~21 kDa, but SDS-PAGE and western blot (WB) analysis of immunoprecipitated mACP-HA$_2$ indicated that this protein migrated with an apparent molecular weight ~15 kDa (Fig. 1C). This lower size suggested that mACP was likely to be post-translationally processed upon mitochondrial import, which would be consistent with known processing of mitochondrial ACP in yeast and humans (27, 28).

We extended our WB analysis to include mACP-HA$_2$ that was heterologously expressed in *E. coli*, where proteolytic processing is not expected. The recombinant, full-length mACP-HA$_2$ expressed in bacteria migrated with an apparent MW close to 21 kDa, as expected. Based on sequence homology, the known ACP-processing sites in yeast and humans (27, 28), and mass spectrometry analysis of parasite-expressed mACP (Fig. S1B), we identified Leu-51 as the possible N-terminus of mature mACP in *P. falciparum*. We therefore cloned and bacterially expressed a truncated mACP-HA$_2$ construct beginning with Leu-51. This truncated mACP protein co-migrated by SDS-PAGE with mACP-HA$_2$ expressed in parasites (Fig. 1C), suggesting that Leu-51 is at or near the N-terminus of mature mACP in parasites. We conclude that mACP is proteolytically processed upon import into the *P. falciparum* mitochondrion.

To directly test if mACP is essential for blood-stage parasite growth and viability, we used CRISPR/Cas9 to tag the mACP gene in Dd2 parasites to encode a C-terminal HA-FLAG epitope tag and the aptamer/TetR-DOZI system that enables anhydrotetracycline (aTc)-dependent control of protein expression (Fig. 1D) (29). Correct integration into the mACP locus was confirmed by PCR analysis of polyclonal and clonal parasite lines (Fig. S3). To evaluate mACP knockdown and its impact on parasite growth, we synchronized parasites to the ring stage, split these parasites into two equal populations ±aTc, and monitored parasite growth over multiple 48-hour growth cycles. Parasites grew indistinguishably ±aTc for the first 3 days. However, parasites grown without aTc displayed a major growth defect on day 4 in the third intraerythrocytic lifecycle (Fig. 1E), similar to prior studies with this knockdown system (29). Blood-smear analysis on day 5 indicated wide-spread parasite death (Fig. S4). Western blot analysis of parasite samples harvested in the second cycle, after three days of growth -aTc, indicated robust knockdown of mACP expression relative

to parasites grown +aTc (Fig. 1E). We conclude that mACP is essential for *P. falciparum* parasites during blood-stage growth.

Mitochondrial ACP binds the Isd11-Nfs1 complex required for Fe-S cluster biogenesis.
Recent studies in yeast and human cells have revealed that mitochondrial ACP binds to a variety of small, 3-helical adapter proteins that bear a conserved Leu-Tyr-Arg (LYR) sequence motif on their N-terminal helix (23, 30). The Leu mediates an intramolecular helical-helical contact while the Tyr and Arg side-chains interact with conserved acidic residues on ACP (24, 26, 31). These LYR-motif proteins mediate diverse mitochondrial processes that include respiratory chain assembly, Fe-S cluster biogenesis, and ribosomal translation (19, 20, 22, 25). Given the strong conservation of mitochondrial LYR proteins and ACP interactions across eukaryotes, we reasoned that the essential function of *P. falciparum* mACP might involve interaction with a conserved LYR protein.

Using the sequences of known LYR proteins from yeast and humans as bait (30), we conducted a BLAST search of the *P. falciparum* genome to look for homologs. We identified Isd11 (PF3D7_1311000) as the only LYR-protein homolog retained by parasites (Table S2). *P. falciparum* Isd11 is 29% identical to human Isd11 (also known as LYRM4), including conservation of the LYR sequence motif near the N-terminus (Fig. 2A). The parasite mACP retains the conserved Glu residues (5 and 11 amino acids C-terminal to the modified Ser in canonical ACPs) known to interact with the Tyr and Arg residues of the LYR motif on Isd11 (Fig. 1A) (24, 26). In yeast and humans, Isd11 directly binds and stabilizes Nfs1, a mitochondrial cysteine desulfurase required for biogenesis of Fe-S clusters needed within and outside mitochondria (32). Mitochondrial ACP in these organisms directly binds Isd11 to stabilize the mACP-Isd11-Nfs1

complex (Fig. 2B), with loss of ACP and/or Isd11 resulting in Nfs1 instability and defective Fe-S cluster biogenesis (19, 33). *P. falciparum* retains a mitochondrial Nfs1 homolog (PF3D7_0727200) that is predicted from genome-wide knock-out studies to be essential for Fe-S cluster biogenesis (18, 34). Collectively, conservation of Isd11 and Nfs1 in *P. falciparum*, the conserved role for ACP in stabilizing the mACP-Isd11-Nfs1 complex in other eukaryotes, and retention of the LYR sequence motif in *P. falciparum* Isd11 strongly suggested that mACP was likely to interact with Isd11 to form the mACP-Isd11-Nfs1 complex in parasites.

To identify protein interaction partners of mACP, we used anti-HA immunoprecipitation (IP) to isolate mACP-HA$_2$ from parasites followed by tandem mass spectrometry (MS) to identify parasite proteins that co-purified with mACP. In multiple independent pull-downs, we identified Nfs1 as strongly enriched in the mACP-HA$_2$ sample compared to a negative control sample (anti-HA IP of aACP-HA$_2$) (Fig. 2C). Anti-Nfs1 IP/WB studies described below further support Nfs1 pull-down with mACP. We did not observe peptides corresponding to Isd11 in mACP-HA$_2$ IP/MS samples. However, its absence is not unexpected given the short Isd11 length (87 residues) that leads to comparatively few tryptic peptides.

To directly test for mACP interaction with Isd11, we transfected the mACP-HA$_2$/Dd2 line with an episome encoding expression of Isd11 with a C-terminal GFP tag that was previously used to localize Isd11 to the mitochondrion (34). Using this Dd2 line expressing mACP-HA$_2$ and Isd11-GFP, we performed reciprocal IP/WB experiments that confirmed stable interaction of these two proteins in parasites (Fig. 2D and Fig. S5A). This interaction appeared to be specific, as mACP-HA$_2$ did not co-IP with the abundant mitochondrial chaperone, HSP60, or with GFP-tagged cytochrome c$_1$ (Fig. 2E and Fig. S5B & S5C). Because these IP experiments cannot distinguish whether mACP binding to Isd11 is direct or mediated by other parasite proteins, we turned to

heterologous studies in *E. coli*. Reciprocal IP experiments with lysates from bacteria co-expressing *P. falciparum* mACP-HA$_2$ and Isd11-His$_6$ confirmed stable interaction of these proteins in the absence of other parasite-specific factors (Fig. 2F). In experiments with Isd11 LYR mutants, however, this interaction was strongly reduced (YR/AA) or eliminated (LYR/AAA), despite robust expression of all proteins (Fig. 2F and Fig. S5D). Clarified lysates from bacteria co-expressing truncated (Δ2-50) mACP-HA$_2$ and Isd11-His$_6$ were also fractionated by passage over Ni-NTA resin to selectively pull down His-tagged Isd11 and interactors followed by size-exclusion chromatography to separate individual complexes. Analysis by Coomassie-stained SDS-PAGE and tandem mass spectrometry indicated robust isolation of a 1:1 complex containing Isd11 and mACP (Fig. 2G and Fig. S6). We conclude that mACP directly binds to Isd11 and that this interaction requires the LYR motif.

Yeast and human Isd11 retain a conserved Arg six residues upstream of the LYR motif (Fig. 2A) that electrostatically stabilizes the negatively charged oxygen atoms of the Ppant group in the ACP-Isd11 complex (24, 26, 31), and mutation of this Arg ablates binding of human Isd11 to ACP (23). Because parasite mACP lacks the negatively charged Ppant group and has replaced it with a hydrophobic Phe, retention of Arg-6 by *P. falciparum* Isd11 would be predicted to destabilize binding of these two proteins. The parasite Isd11, however, has replaced Arg-6 with an Ile residue (Fig. 2A) expected to more favorably interact with the hydrophobic surface of mACP created by Ser-to-Phe substitution. These sequence features suggest that parasite Isd11 has co-evolved with mACP to optimize formation of the mACP-Isd11 complex in *P. falciparum* in the absence of FASII and an acyl-Ppant group on mACP. Consistent with this model, parasite Isd11 did not appear to interact with *E. coli* ACP, which retains the acyl-Ppant modification, in contrast to human Isd11 that robustly binds bacterial ACP when expressed in *E. coli* (24, 26, 35).

Structural modeling of the mACP-Isd11 interface. Association of Isd11 and mACP in yeast and human cells depends critically on the acyl-Ppant group of mACP (Fig. 3A), with little or no association observed in these cells between Isd11 and the mACP Ser-to-Ala mutant that lacks a Ppant group (19, 20, 23). As an initial step to understand the divergent molecular features of the *P. falciparum* mACP-Isd11 interface that stabilize complexation in the absence of an acyl-Ppant group, we generated a homology model of this complex using published structures and then carried out energy optimization and refinement using the Rosetta software (36). The resulting low-energy structural model suggested that the unusual Phe113 residue of mACP partitions into a hydrophobic pocket at the mACP-Isd11 interface composed of the aliphatic side-chain methylene groups of Lys10 and Arg14 and the phenyl ring of Tyr13 of Isd11 as well as Val117 on mACP (Fig. 3B).

To test if the divergent Phe113 residue of mACP contributes substantially to its association with Isd11, we heterologously co-expressed Isd11-His$_6$ and Δ2-50 mACP-HA$_2$ (WT or the F113A mutant) in *E. coli* and observed robust expression of all proteins in bacterial lysates. We then used Ni-NTA to selectively pull down His-tagged Isd11 and observed that WT mACP-HA$_2$ but not the F113A mutant co-purified with Isd11 (Fig. 3C and S5E). These results indicate that mutation of Phe113 to Ala substantially weakens the mACP-Isd11 interaction, supporting the conclusion that Phe113 contributes to the mACP-Isd11 binding interface in the mACP-Isd11-Nfs1 complex as suggested by structural modeling (Fig. 3B).

To test the functional impact of the Phe113Ala mACP mutation on parasite growth, we transfected our mACP knockdown parasite line with episomes encoding either WT or Phe113Ala mACP. Parasites expressing a second-copy of WT mACP showed identical growth in the presence or absence of aTc. In contrast, parasites that episomally expressed the Phe113Ala mACP mutant

showed strongly diminished growth upon aTc removal and loss of endogenous mACP (Fig. 3D). These results support the conclusion that the Phe residue of mACP plays a key role in stabilizing the mACP-Isd11 association required for Nfs1 stability and function.

Loss of mACP destabilizes Nfs1 and the Rieske protein in ETC complex III. Nfs1 stability in other eukaryotes depends critically on its association with Isd11 and ACP, with loss of ACP resulting in degradation of Nfs1 and lethal defects in Fe-S cluster biogenesis (19, 24, 26, 33, 35). Loss of Nfs1 upon mACP knockdown would therefore be sufficient to explain mACP essentiality in *P. falciparum*. To test the impact of mACP knockdown on Nfs1 protein levels in parasites, we used a commercial anti-Nfs1 antibody (Abcam 229829) raised against a 250-amino acid region of human Nfs1 that is 55% identical to *P. falciparum* Nfs1 and which selectively recognized recombinant *P. falciparum* Nfs1 expressed in *E. coli* (Fig. S5F). We grew our mACP aptamer/TetR-DOZI parasites ±aTc, harvested them at the end of the second intraerythrocytic growth cycle before parasite death, and analyzed expression of Nfs1 by western blot using this Nfs1 antibody.

In parasites grown +aTc, the anti-Nfs1 antibody recognized a band at ~55 kDa (Fig. 4A) that is slightly lower than the ~63 kDa band recognized by this antibody for full-length Nfs1 expressed in *E. coli* (Fig. S5C). This observation suggests that Nfs1 is processed upon mitochondrial import in *P. falciparum,* which is consistent with observed processing of Nfs1 in yeast (27). Western blot analysis of parasites grown -aTc failed to detect this 55 kDa band, strongly suggesting that Nfs1 stability is tightly coupled to mACP expression and that loss of mACP results in Nfs1 degradation (Fig. 4A). Loss of Nfs1 upon mACP knockdown suggests that parasite death

is due to dysfunction in one or more Fe-S cluster-dependent pathways. We set out to understand which pathway defect(s) might explain parasite death.

The mitochondrial iron-sulfur cluster (ISC) biogenesis pathway has been shown in yeast and other organisms to supply Fe-S clusters to various mitochondrial proteins and to produce a key sulfur-containing intermediate that is exported out of the organelle and used for maturation of Fe-S clusters in the cytoplasm and delivery to client proteins (33, 37). Multiple Fe-S cluster proteins function within the *P. falciparum* mitochondrion (38). Aconitase (PF3D7_1342100) and succinate dehydrogenase (PF3D7_1212800) are dispensable for blood-stage parasite growth (7). Ferredoxin (PF3D7_1214600) is expected to be essential (18), but its main role is to provide electrons for the ISC pathway (39). Class I fumarate hydratase (PF3D7_0927300), which functions in the TCA cycle but may also contribute to purine scavenging, was refractory to disruption in *P. falciparum* (7) but successfully deleted in *P. berghei* in a mouse strain-dependent manner (40). The Rieske protein (PF3D7_1439400), which is an essential component of ETC complex III (41), is the only known mitochondrial Fe-S cluster protein that has an unequivocally essential function apart from the ISC pathway. We therefore focused on understanding the effect of mACP knockdown on the Rieske protein, as a prior study in yeast showed that mACP knockdown resulted in loss of Rieske (19).

To probe the status of Rieske upon mACP knockdown in parasites, we transfected the mACP-aptamer/TetR-DOZI parasites with an episome encoding Rieske-HA$_2$. A band corresponding to the epitope-tagged Rieske was detected ~43 kDa in western blot analysis of parasites grown +aTc. This band, however, was undetectable in parasites grown 72 hours in -aTc conditions to downregulate mACP expression (Fig. 4B). These results indicate that knockdown of

mACP and subsequent degradation of Nfs1 result in loss of the Fe-S cluster-dependent Rieske protein.

Loss of mACP causes ETC failure and sensitizes parasites to mitochondrial depolarization by proguanil. Since the Rieske protein is an essential electron-transfer component of ETC complex III (41), we reasoned that loss of mACP and Rieske likely results in ETC failure. In blood-stage parasites, which rely on glycolysis rather than oxidative phosphorylation for ATP synthesis, the essential function of the mitochondrial ETC is to oxidatively recycle the ubiquinone cofactor used by several dehydrogenases, of which dihydroorotate dehydrogenase is most critical (1, 42). Prior work has shown that exogenous addition of the soluble ubiquinone analog, decyl-ubiquinone (dQ), rescues parasites from ETC dysfunction caused by the ETC complex III inhibitor, atovaquone (42). We posited that it might be possible to rescue parasites with dQ from mACP knockdown if the dominant cause of parasite death were ETC failure.

As a positive control and to provide a basis for comparison, we first confirmed that 15 µM exogenous dQ substantially rescued parasites over multiple intraerythrocytic cycles from growth inhibition by 100 nM atovaquone (Fig. 5A), as previously reported (42). To test dQ rescue of parasites upon mACP knockdown, we synchronized the mACP-aptamer/TetR-DOZI parasites and monitored their growth ±aTc and ±dQ for several intraerythrocytic cycles. As before, -aTc parasites grew normally for three days but failed to expand into the third growth cycle on day four (Fig. 5A). Addition of dQ rescued parasite growth in the third growth cycle in -aTc conditions, similar to dQ rescue of parasites from atovaquone. However, in contrast to indefinite parasite rescue from atovaquone, dQ only rescued the third-cycle growth of parasites cultured -aTc, which failed to expand further and began to die off on day six in the fourth intraerythrocytic cycle (Fig.

5A). On the basis of dQ rescue, we conclude that ETC failure is the immediate cause of parasite death upon mACP knockdown. However, the inability of dQ to rescue parasites beyond the third cycle indicates that additional dysfunctions beyond the mitochondrial ETC contribute to parasite death on a longer timescale. These additional defects may involve mitochondrial fumarate hydratase and/or essential Fe-S cluster proteins in the cytoplasm and nucleus that depend on Nfs1 function (see Discussion below).

To further unravel the effect of mACP knockdown on ETC function, we next tested if loss of mACP sensitized parasites to mitochondrial depolarization by proguanil. The ETC couples ubiquinone recycling to proton translocation from the mitochondrial matrix to the inner membrane space to establish a transmembrane electrochemical potential. Prior work has shown that parasites retain a second proton-pumping mechanism that also contributes to transmembrane potential and which is inhibited by proguanil (1, 43). Inhibition of ETC function kills parasites due to defective ubiquinone recycling but does not substantially depolarize mitochondria due to activity by this second, proguanil-sensitive pathway. However, ETC dysfunction plus proguanil treatment blocks both pathways for proton pumping and causes mitochondrial depolarization, which can be visualized by a failure to concentrate charged dyes like Mitotracker within the mitochondrion that leads to dispersed dye accumulation in the cytoplasm (1, 44).

If mACP knockdown and subsequent loss of Rieske caused general ETC failure, we predicted that parasite treatment with sub-lethal proguanil would negate the ability of dQ to rescue parasite growth from mACP knockdown in the third cycle and would cause mitochondrial depolarization. We repeated the prior growth assay ±aTc and ±dQ but also included 1 µM proguanil. This proguanil concentration alone had no effect on parasite growth, as previously reported (1). However, when proguanil was combined with growth -aTc, dQ was unable to rescue

parasite growth in the third cycle (Fig. 5B). Microscopy analysis of parasites on day three of the growth assay revealed that proguanil treatment selectively prevented mitochondrial accumulation of Mitotracker Red in parasites grown -aTc, strongly suggesting mitochondrial depolarization (Fig. 5C and 5D, and Fig. S7). This observation strongly supports the model that impaired ubiquinone recycling due to Rieske loss and ETC dysfunction is the immediate cause of parasite death upon mACP knockdown.

DISCUSSION

The *P. falciparum* mitochondrion is a major antimalarial drug target, but nearly all organelle-specific inhibitors target cytochrome b in ETC complex III or dihydroorotate dehydrogenase (DHOD) that depends on complex III function (45, 46). Multiple metabolic pathways operate within the mitochondrion (3), but many are dispensable for blood-stage parasites and few essential functions beyond DHOD and ETC complexes III and IV have been identified. Iron-sulfur cluster biosynthesis is an ancient, essential mitochondrial function that has been well studied in yeast and mammalian cells but is sparsely studied in parasites (34). We have identified a divergent and essential nexus between Fe-S cluster biogenesis and an evolutionary vestige of type-II fatty acid synthesis in the *P. falciparum* mitochondrion.

New molecular paradigm for essential ACP function without an acyl-Ppant group. Most eukaryotic cells, including fungi, plants, and animals, retain a mitochondrial FASII pathway in which the acyl-ACP intermediate has been shown to critically mediate respiratory chain assembly, Fe-S cluster biogenesis, and ribosomal translation by binding to LYR-protein assembly factors (23, 30, 47). These interactions and their dependence on ACP acylation have been proposed to

constitute a regulatory feedback mechanism that couples the availability of acetyl-CoA for FASII activity to respiratory chain assembly for oxidative phosphorylation and ATP synthesis (20). Biochemical coupling of these pathways, however, has not been retained in *Plasmodium*, which has lost has lost mitochondrial FASII enzymes but retains a divergent ACP homolog incapable of 4-Ppant modification or acylation. Nevertheless, we have shown that mACP remains essential to parasites for Fe-S cluster biogenesis by binding Isd11 and stabilizing the mACP-Isd11-Nfs1 cysteine desulfurase complex via a novel interface. *P. falciparum* thus provides a new molecular paradigm for essential ACP function without an acyl-Ppant group. This discovery emphasizes the ancient, fundamental role of ACP in mitochondrial Fe-S cluster biogenesis and suggests an evolutionary driving force to retain mACP interaction with the Isd11-Nfs1 complex independent ACP's scaffolding role in fatty acid synthesis.

Prior in vitro work has provided evidence that activity of purified Isd11-Nfs1 does not strictly require association with ACP (26, 33). However, Nfs1 is unstable in its native mitochondrial context in the absence of ACP, as shown herein for *P. falciparum* and in prior studies of yeast (19, 48). Coupling Nfs1 stability to ACP acylation in eukaryotes that retain mitochondrial FASII has been proposed as a mechanism to up- and down-regulate Fe-S cluster biogenesis congruent with nutrient availability and cellular Fe-S cluster needs for respiration and growth (33). This mechanism, however, cannot explain mitochondrial retention of an acylation-incompetent ACP homolog in *Plasmodium*. ACP may therefore play additional functional and/or regulatory roles that are essential for mitochondrial Fe-S cluster biogenesis and utilization in cells, perhaps involving metal-ion binding and sensing by ACP (e.g., iron or zinc) (31, 49) or interactions with other mitochondrial networks akin to broad ACP interactions observed in *E. coli* (50). In this regard it is interesting to note that anaerobic eukaryotic parasites like *Giardia* and *Cryptosporidium*

lack a mitochondrion but retain a primitive mitosome for Fe-S cluster biogenesis that lacks ACP and Isd11 homologs in this compartment, where Nfs1 appears to function autonomously (51, 52).

Implications for parasites. Loss of mitochondrial FASII but retention of a Ppant-independent mACP lacking the conserved Ser appears to be widespread among Apicomplexan parasites, including *Toxoplasma* and *Babesia*. Outside the phylum Apicomplexa, the only genomic evidence we found for a divergent ACP lacking the conserved Ser was in the related photosynthetic chromerid, *Vitrella brassicaformis,* but not in its algal cousin, *Chromera velia*. Thus, the adaptation of mitochondrial ACP to function without a Ppant group or acylation likely occurred on a similar evolutionary time-frame as the loss of plastid photosynthesis that accompanied the appearance of Apicomplexan parasitism (53).

Our study indicates that mitochondrial ACP function without an acyl-Ppant group is a parasite-specific adaptation that differentiates Apicomplexan parasites from humans. Nevertheless, like yeast and human cells (19, 23), we observed that loss of ACP destabilizes Nfs1 to block Fe-S cluster assembly, impairs the stability and function of the key Rieske subunit of ETC complex III, and leads to parasite death (Fig. 6). In yeast and mammalian cells, mACP plays a second role in Rieske maturation by binding to Mzm1, an LYR-protein chaperone that stabilizes Rieske upon import into the mitochondrial matrix to receive its Fe-S cluster for subsequent insertion into ETC complex III by the AAA-ATPase, BCS1 (23, 54). *P. falciparum*, however, appears to lack an Mzm1 homolog (Table S2), suggesting that parasites have evolved compensatory mechanisms for Rieske maturation in the absence of this ACP-dependent chaperone.

Exogenous dQ rescued parasite growth upon mACP knockdown for one intraerythrocytic cycle but did not rescue growth on a longer time-scale. This short-term rescue indicates that respiratory failure is the immediate cause of *P. falciparum* death upon loss of mACP (1, 42). However, defects in essential processes beyond the mitochondrial ETC contribute to parasite death on longer timescales. These defects may involve mitochondrial fumarate hydratase (7) and/or essential Fe-S cluster proteins in the cytosol and nucleus that depend on Nfs1 activity, including Rli1 required for cytoplasmic ribosome assembly and Rad3 and Pri2 that are essential for nuclear DNA metabolism (55). In yeast, the mitochondrial transporter Atm1 exports a key sulfur-containing intermediate required for cytosolic Fe-S cluster assembly (37, 52). In *P. falciparum*, the Mdr2 transporter (PF3D7_1447900) is 34% identical to yeast Atm1 and has been proposed to perform an analogous function in cytosolic Fe-S cluster assembly (3). However, Mdr2 appears to be dispensable for blood-stage parasites (56), and its possible mitochondrial-export role in cytoplasmic Fe-S cluster biogenesis remains untested.

The mitochondrial ETC is essential for malaria parasite viability in all life-cycle stages (57). The unique molecular features of *P. falciparum* mACP and its interaction with Isd11 that underpin essential functions of the ETC and broader Fe-S cluster utilization suggest the possibility of targeting this complex for antimalarial therapy. We have on-going structural and biochemical studies to assess if parasite Isd11 retains a vestigial acyl-pantetheine binding pocket and to test if parasite-specific features of the mACP-Isd11 interaction can be selectively disrupted via small-molecule inhibitors that mimic the acyl-pantetheine group and/or broader hydrophobic features of this protein-protein interface that are distinct from the human complex. On the basis of molecular conservation in other Apicomplexan parasites, such inhibitors would likely function against other human pathogens such as *Toxoplasma* and *Babesia*.

MATERIALS AND METHODS

A complete description of all experimental methods is given in SI Appendix.

Parasite growth assays: All parasite studies were performed with *P. falciparum* Dd2 parasites cultured in RPMI-1640 medium as previously described (58). CRISPR/Cas9 was used to tag the mACP gene in Dd2 parasites to encode a C-terminal HA-FLAG epitope tag and the aptamer/TetR-DOZI system (29), whose integration was validated by PCR. Growth assays were performed ±aTc by synchronizing parasite by 5% D-sorbitol, monitoring culture parasitemia by acridine orange staining and flow cytometry of ≥2 biological replicate samples, and reporting the parasitemia average and standard deviation. Decyl-ubiquinone was used in chemical rescue experiments at 15 µM final concentration and was added ±1 µM proguanil at the beginning of growth assays.

Western blot analyses: Parasite sample were harvested by 0.05% saponin treatment. For whole-parasite experiments, saponin pellets were dissolved in 60 µl SDS sample buffer containing beta-mercaptoethanol. For immunoprecipitation experiments, parasite pellets from 50-mL cultures were lysed in 1 mL 1% triton or digitonin and clarified by centrifugation. Lysate supernatants were incubated with 30 µl of pre-equilibrated Pierce anti-HA magnetic beads, washed 3X with Tris-buffered saline + 0.05% Tween-20, eluted with 8M urea, and diluted with 5X SDS sample buffer. For bacteria expression studies, BL21/DE3 *E. coli* were transformed with pET plasmids encoding cloned *P. falciparum* mACP-HA$_2$ (full-length or the Δ2-50 truncation, WT or F113A mutant), Isd11-His$_6$ (WT, YR/AA, or LYR/AAA mutants), or full-length Nfs1-HA$_2$. Protein expression was induced by 1 mM IPTG, and bacterial pellets were lysed in PBS by sonication and clarified by centrifugation. Ni-NTA resin was used to pull-down His$_6$-tagged proteins from clarified lysates,

and proteins were eluted in 500 mM imidazole. To purify Isd11-containing complexes, the imidazole eluate was run over an S-100 size-exclusion column on an AKTA FPLC system. Lysates or eluates were diluted into SDS sample buffer.

Samples were fractionated by SDS-polyacrylamide gel electrophoresis (PAGE) using 10% acrylamide gels run at 120 V in the BIO-RAD mini-PROTEAN electrophoresis system and transferred to nitrocellulose. Membranes were probed with 1:1000 dilutions of Roche rat anti-HA monoclonal 3F10 antibody, Abcam goat anti-GFP polyclonal antibody ab5450, Abcam rabbit anti-Nfs1 polyclonal antibody ab229829, Novus anti *P. falciparum* Hsp60 polyclonal antibody NBP2-12734, and/or rabbit anti-EF1α or EF1β polyclonal antibody (59). Blots were probed with 1:5000 dilutions of fluorophore-conjugated secondary antibodies and imaged on a Licor Odyssey system.

Microscopy: For live-cell experiments, parasites nuclei were visualized by staining with Hoechst 33342, and mitochondrial polarization was monitored by incubating parasites with 10 nM MitoTracker Red CMXROS for 15 minutes prior to wash-out and imaging. 40-50 total parasites for each condition in two independent experiments were scored for focal or dispersed MitoTracker signal, and cell percentages were analyzed by two-tailed unpaired t-test in Graphpad Prism. For IFA studies, the parasite mitochondrion was visualized using the Novus anti-HSP60 antibody and AlexaFluor 647-conjugated goat anti-rabbit 2˚ antibody (Invitrogen Life Technologies A21244), and mACP-HA$_2$ was visualized with the rat anti-HA 3F10 primary antibody and FITC-conjugated donkey anti-rat 2˚ antibody (Invitrogen Life Technologies A18746). Images were taken on an EVOS M5000 imaging system or Zeiss 880 laser-scanning confocal microscope fitted with an Airyscan detector. Fiji/ImageJ was used to process and analyze images. All image adjustments, including contrast and brightness, were made on a linear scale.

Sequence analysis: Sequence similarity searches of the *P. falciparum* genome for protein homologs of *E. coli* FASII proteins and known LYR proteins from yeast and humans (30) were performed by BLASTP analysis as implemented at the Plasmodium Genomics Resource webpage (www.plasmodb.org, release 48). Only protein hits with e-values <0.01 were retained.

Structural modeling: A homology model of *P. falciparum* mACP bound to Isd11 was created using the InterPred modeling interface (60) and the previously determined structure (24) of *E. coli* ACP bound to human Isd11 (PDB 5USR, chains L and D). This structural model was independently refined using Rosetta Dual Relax (61), with and without restraining refinement to template coordinates, and evaluated for Rosetta energy (62). The unrestrained refinement resulted in the final model of lowest calculated energy, and this model was used for further analysis. MacPyMOL version 1.8 was used for structural visualization.

ACKNOWLEDGEMENTS

We thank Greg Ducker, Chris Hill, Roland Lill, Sara Nowinski, Jared Rutter, Dennis Winge, and members of the Sigala lab for helpful discussions and thank Sean Prigge for helpful discussions and the plasmid encoding *P. falciparum* Isd11-GFP. We thank Rebecca Marvin for assistance with making the *P. falciparum* cytochrome c_1-GFP expression plasmid, Megan Okada for assistance with graphical figure schemes and the design of integration vectors, and Sandra Osburn for assistance with recombinant protein mass spectrometry. This work was funded by a pilot grant to P.A.S. from the Utah Center for Iron and Heme Disorders (funded by NIH grant U54 DK110858) and NIH grants to P.A.S. (R35 GM133764) and J.A.W. (R01 GM089778). P.A.S. holds a Career

Award at the Scientific Interface from the Burroughs Wellcome Fund and a Pew Biomedical Scholarship from the Pew Charitable Trusts. J.S. was funded in part by NIH T32 GM122740. S.F. was supported by the African American Doctoral Scholars Initiative at the University of Utah. Microscopy, flow cytometry, protein mass spectrometry, and DNA synthesis and sequencing were performed using core facilities at the University of Utah. Mass spectrometry core facilities at the University of Utah were supported by an NIH instrument grant (S10 OD018210).

REFERENCES

1. H. J. Painter, J. M. Morrisey, M. W. Mather, A. B. Vaidya, Specific role of mitochondrial electron transport in blood-stage *Plasmodium falciparum*. *Nature* **446**, 88-91 (2007).
2. H. Jhun, M. S. Walters, S. T. Prigge, Using Lipoamidase as a Novel Probe To Interrogate the Importance of Lipoylation in Plasmodium falciparum. *mBio* **9** (2018).
3. G. G. van Dooren, L. M. Stimmler, G. I. McFadden, Metabolic maps and functions of the Plasmodium mitochondrion. *FEMS Microbiol. Rev.* **30**, 596-630 (2006).
4. R. D. Oppenheim *et al.*, BCKDH: the missing link in apicomplexan mitochondrial metabolism is required for full virulence of Toxoplasma gondii and Plasmodium berghei. *PLoS Pathog.* **10**, e1004263 (2014).
5. H. Ke *et al.*, The heme biosynthesis pathway is essential for Plasmodium falciparum development in mosquito stage but not in blood stages. *J. Biol. Chem.* **289**, 34827-34837 (2014).
6. V. A. Nagaraj *et al.*, Malaria parasite-synthesized heme is essential in the mosquito and liver stages and complements host heme in the blood stages of infection. *PLoS Pathog.* **9**, e1003522 (2013).
7. H. Ke *et al.*, Genetic investigation of tricarboxylic acid metabolism during the Plasmodium falciparum life cycle. *Cell Rep.* **11**, 164-174 (2015).
8. A. Sturm, V. Mollard, A. Cozijnsen, C. D. Goodman, G. I. McFadden, Mitochondrial ATP synthase is dispensable in blood-stage Plasmodium berghei rodent malaria but essential in the mosquito phase. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 10216-10223 (2015).
9. J. I. MacRae *et al.*, Mitochondrial metabolism of sexual and asexual blood stages of the malaria parasite Plasmodium falciparum. *BMC Biol.* **11**, 67-77 (2013).
10. J. K. Hiltunen *et al.*, Mitochondrial fatty acid synthesis type II: more than just fatty acids. *J. Biol. Chem.* **284**, 9011-9015 (2009).
11. M. J. Shears, C. Y. Botte, G. I. McFadden, Fatty acid metabolism in the Plasmodium apicoplast: Drugs, doubts and knockouts. *Mol. Biochem. Parasitol.* **199**, 34-50 (2015).
12. M. Allary, J. Z. Lu, L. Zhu, S. T. Prigge, Scavenging of the cofactor lipoate is essential for the survival of the malaria parasite Plasmodium falciparum. *Mol. Microbiol.* **63**, 1331-1344 (2007).

13.   D. I. Chan, H. J. Vogel, Current understanding of fatty acid biosynthesis and the acyl carrier protein. *Biochem. J.* **430**, 1-19 (2010).

14.   J. R. Gallagher, S. T. Prigge, Plasmodium falciparum acyl carrier protein crystal structures in disulfide-linked and reduced states and their prevalence during blood stage growth. *Proteins* **78**, 575-588 (2010).

15.   B. C. van Schaijk *et al.*, Type II fatty acid biosynthesis is essential for Plasmodium falciparum sporozoite development in the midgut of Anopheles mosquitoes. *Eukaryot. Cell* **13**, 550-559 (2014).

16.   M. Yu *et al.*, The Fatty Acid Biosynthesis Enzyme Fabl Plays a Key Role in the Development of Liver-Stage Malarial Parasites. *Cell Host Microbe* **4**, 567-578 (2008).

17.   M. Zhang *et al.*, Uncovering the essential genes of the human malaria parasite Plasmodium falciparum by saturation mutagenesis. *Science* **360** (2018).

18.   A. R. Gomes *et al.*, A genome-scale vector resource enables high-throughput reverse genetic screening in a malaria parasite. *Cell Host Microbe* **17**, 404-413 (2015).

19.   J. G. Van Vranken *et al.*, The mitochondrial acyl carrier protein (ACP) coordinates mitochondrial fatty acid synthesis with iron sulfur cluster biogenesis. *Elife* **5** (2016).

20.   J. G. Van Vranken *et al.*, ACP Acylation Is an Acetyl-CoA-Dependent Modification Required for Electron Transport Chain Assembly. *Mol. Cell* **71**, 567-580 e564 (2018).

21.   S. M. Nowinski *et al.*, Mitochondrial fatty acid synthesis coordinates oxidative metabolism in mammalian mitochondria. *Elife* **9** (2020).

22.   A. Brown *et al.*, Structures of the human mitochondrial ribosome in native states of assembly. *Nat. Struct. Mol. Biol.* **24**, 866-869 (2017).

23.   J. D. Majmudar *et al.*, 4'-Phosphopantetheine and long acyl chain-dependent interactions are integral to human mitochondrial acyl carrier protein function. *Med. Chem. Comm.* **10**, 209-220 (2019).

24.   S. A. Cory *et al.*, Structure of human Fe-S assembly subcomplex reveals unexpected cysteine desulfurase architecture and acyl-ACP-ISD11 interactions. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E5325-E5334 (2017).

25.   H. Angerer *et al.*, The LYR protein subunit NB4M/NDUFA6 of mitochondrial complex I anchors an acyl carrier protein and is essential for catalytic activity. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 5207-5212 (2014).

26.   M. T. Boniecki, S. A. Freibert, U. Muhlenhoff, R. Lill, M. Cygler, Structure and functional dynamics of the mitochondrial Fe/S cluster synthesis complex. *Nat. Commun.* **8**, 1287 (2017).

27.   F. N. Vogtle *et al.*, Global analysis of the mitochondrial N-proteome identifies a processing peptidase critical for protein stability. *Cell* **139**, 428-439 (2009).

28.   A. S. Vaca Jacome *et al.*, N-terminome analysis of the human mitochondrial proteome. *Proteomics* **15**, 2519-2524 (2015).

29.   S. M. Ganesan, A. Falla, S. J. Goldfless, A. S. Nasamu, J. C. Niles, Synthetic RNA-protein modules integrated with native translation mechanisms to control gene expression in malaria parasites. *Nat. Commun.* **7**, 10727 (2016).

30.   H. Angerer, Eukaryotic LYR Proteins Interact with Mitochondrial Protein Complexes. *Biology (Basel)* **4**, 133-150 (2015).

31.   M. G. Herrera *et al.*, Structure of the Human ACP-ISD11 Heterodimer. *Biochemistry* **58**, 4596-4609 (2019).

32.   N. Wiedemann *et al.*, Essential role of Isd11 in mitochondrial iron-sulfur cluster synthesis on Isu scaffold proteins. *EMBO J.* **25**, 184-195 (2006).

33.   R. Lill, S. A. Freibert, Mechanisms of Mitochondrial Iron-Sulfur Protein Biogenesis. *Annu. Rev. Biochem.* **89**, 471-499 (2020).

34.   J. E. Gisselberg, T. A. Dellibovi-Ragheb, K. A. Matthews, G. Bosch, S. T. Prigge, The suf iron-sulfur cluster synthesis pathway is required for apicoplast maintenance in malaria parasites. *PLoS Path.* **9**, e1003655 (2013).

35.   K. Cai, R. O. Frederick, M. Tonelli, J. L. Markley, Mitochondrial Cysteine Desulfurase and ISD11 Coexpressed in Escherichia coli Yield Complex Containing Acyl Carrier Protein. *ACS Chem. Biol.* **12**, 918-921 (2017).

36.   H. Park, S. Ovchinnikov, D. E. Kim, F. DiMaio, D. Baker, Protein homology model refinement by large-scale energy optimization. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 3054-3059 (2018).

37.   G. Kispal, P. Csere, C. Prohl, R. Lill, The mitochondrial proteins Atm1p and Nfs1p are essential for biogenesis of cytosolic Fe/S proteins. *EMBO J.* **18**, 3981-3989 (1999).

38.   J. M. Haussig, K. Matuschewski, T. W. Kooij, Identification of vital and dispensable sulfur utilization factors in the Plasmodium apicoplast. *PLoS One* **9**, e89718 (2014).

39.   H. Lange, A. Kaut, G. Kispal, R. Lill, A mitochondrial ferredoxin is essential for biogenesis of cellular iron-sulfur proteins. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 1050-1055 (2000).

40.   V. Jayaraman, A. Suryavanshi, P. Kalale, J. Kunala, H. Balaram, Biochemical characterization and essentiality of Plasmodium fumarate hydratase. *J. Biol. Chem.* **293**, 5878-5894 (2018).

41.   Z. Zhang *et al.*, Electron transfer by domain movement in cytochrome bc1. *Nature* **392**, 677-684 (1998).

42.   H. J. Ke *et al.*, Variation among Plasmodium falciparum Strains in Their Reliance on Mitochondrial Electron Transport Chain Function. *Eukaryot. Cell* **10**, 1053-1061 (2011).

43.   T. S. Skinner-Adams *et al.*, Cyclization-blocked proguanil as a strategy to improve the antimalarial activity of atovaquone. *Commun. Biol.* **2**, 166 (2019).

44.   H. Ke, S. Dass, J. M. Morrisey, M. W. Mather, A. B. Vaidya, The mitochondrial ribosomal protein L13 is critical for the structural and functional integrity of the mitochondrion in Plasmodium falciparum. *J. Biol. Chem.* **293**, 8128-8137 (2018).

45.   Y. Antonova-Koch *et al.*, Open-source discovery of chemical leads for next-generation chemoprotective antimalarials. *Science* **362** (2018).

46.   C. D. Goodman, H. D. Buchanan, G. I. McFadden, Is the Mitochondrion a Good Malaria Drug Target? *Trends Parasitol.* **33**, 185-193 (2017).

47.   S. M. Nowinski, J. G. Van Vranken, K. K. Dove, J. Rutter, Impact of Mitochondrial Fatty Acid Synthesis on Mitochondrial Biogenesis. *Curr. Biol.* **28**, R1212-R1219 (2018).

48.   A. C. Adam, C. Bornhovd, H. Prokisch, W. Neupert, K. Hell, The Nfs1 interacting protein Isd11 has an essential role in Fe/S cluster biogenesis in mitochondria. *EMBO J.* **25**, 174-183 (2006).

49.   X. Qiu, C. A. Janson, Structure of apo acyl carrier protein and a proposal to engineer protein crystallization through metal ions. *Acta Crystallogr. D Biol. Crystallogr.* **60**, 1545-1554 (2004).

50. D. Gully, D. Moinier, L. Loiseau, E. Bouveret, New partners of acyl carrier protein detected in Escherichia coli by tandem affinity purification. *FEBS Lett.* **548**, 90-96 (2003).

51. T. A. Richards, M. van der Giezen, Evolution of the Isd11-IscS complex reveals a single alpha-proteobacterial endosymbiosis for all eukaryotes. *Mol. Biol. Evol.* **23**, 1341-1344 (2006).

52. J. J. Braymer, S. A. Freibert, M. Rakwalska-Bange, R. Lill, Mechanistic concepts of iron-sulfur protein biogenesis in Biology. *Biochim. Biophys. Acta Mol. Cell. Res.* **1868**, 118863 (2021).

53. J. Janouskovec *et al.*, Factors mediating plastid dependency and the origins of parasitism in apicomplexans and their close relatives. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 10200-10207 (2015).

54. A. Atkinson *et al.*, The LYR protein Mzm1 functions in the insertion of the Rieske Fe/S protein in yeast mitochondria. *Mol. Cell. Biol.* **31**, 3988-3996 (2011).

55. T. A. Dellibovi-Ragheb, J. E. Gisselberg, S. T. Prigge, Parasites FeS up: iron-sulfur cluster biogenesis in eukaryotic pathogens. *PLoS Pathog.* **9**, e1003227 (2013).

56. M. van der Velden, S. R. Rijpma, F. G. Russel, R. W. Sauerwein, J. B. Koenderink, PfMDR2 and PfMDR5 are dispensable for Plasmodium falciparum asexual parasite multiplication but change in vitro susceptibility to anti-malarial drugs. *Malar. J.* **14**, 76 (2015).

57. M. Delves *et al.*, The activities of current antimalarial drugs on the life cycle stages of Plasmodium: a comparative study with human and rodent parasites. *PLoS Med.* **9**, e1001169 (2012).

58. M. Okada, P. Guo, S. A. Nalder, P. A. Sigala, Doxycycline has distinct apicoplast-specific mechanisms of antimalarial activity. *Elife* **9** (2020).

59. C. B. Mamoun, D. E. Goldberg, Plasmodium protein phosphatase 2C dephosphorylates translation elongation factor 1beta and inhibits its PKC-mediated nucleotide exchange activity in vitro. *Mol. Microbiol.* **39**, 973-981 (2001).

60. C. Mirabello, B. Wallner, InterPred: A pipeline to identify and model protein-protein interactions. *Proteins* **85**, 1159-1170 (2017).

61. P. Conway, M. D. Tyka, F. DiMaio, D. E. Konerding, D. Baker, Relaxation of backbone bond geometry improves protein energy landscape modeling. *Protein Sci.* **23**, 47-55 (2014).

62. H. Park *et al.*, Simultaneous Optimization of Biomolecular Energy Functions on Features from Small Molecules and Macromolecules. *J. Chem. Theory Comput.* **12**, 6201-6212 (2016).

Fig. 1. Mitochondrial ACP is essential for blood-stage *P. falciparum*. (A) Partial sequence alignment of acyl carrier protein homologs from *E. coli* and the *P. falciparum* apicoplast (aACP) and mitochondrion (mACP). The conserved Ser residue that is modified by a 4-Ppant group is in red, the divergent Phe residue of mACP is in purple, and conserved Glu residues are in blue. (B) Immunofluorescence microscopy of fixed Dd2 parasites episomally expressing mACP-HA$_2$ and stained with DAPI (nucleus, blue), anti-HA (green), and anti-HSP60 (mitochondrion, red) antibodies (BF = bright field). (C) Anti-HA western blot (WB) analysis of cellular lysates of *E. coli* expressing full-length (FL) or truncated (Δ2-50) mACP-HA$_2$ and Dd2 *P. falciparum* parasites episomally expressing full-length mACP-HA$_2$. The left and right images are from different WB experiments (10% gels) that were aligned by molecular weight (MW) markers. (D) Schematic depiction of protein translational regulation using the aptamer/TetR-DOZI system. Normal protein expression occurs in the presence but not absence of anhydrotetracycline (aTc). (E) Continuous growth assay of synchronous mACP-HA/FLAG-aptamer/TetR-DOZI parasites in the presence or absence of aTc. Data points and error bars are the average and standard deviation from two biological replicates. Inset is an anti-HA and anti-EF1α western blot of parasite lysates harvested on day 3 of the continuous growth assay.

Fig. 2. Mitochondrial ACP binds the Isd11-Nfs1 complex. (A) Partial sequence alignment of human and *P. falciparum* Isd11. The conserved LYR sequence motif is in red, the conserved Phe residue is in blue, and residue 6 is in green. (B) X-ray structural model of the mACP-Isd11-Nfs1 complex (PDB entry 5USR). For simplicity, only a single copy of each protein is shown, rather than the functional dimer (26). (C) Table of spectral counts for *P. falciparum* Nfs1 detected by tandem mass spectrometry for anti-HA immunoprecipitation studies of lysates from Dd2 parasites episomally expressing mACP-HA$_2$ or aACP-HA$_2$. Parasites were lysed in either Triton X-100 or digitonin. Similar spectral counts for mACP and aACP bait proteins were detected in each experiment (Table S3). (D) Anti-HA or anti-GFP co-immunoprecipitation/western blot studies of Dd2 parasites episomally expressing mACP-HA$_2$ with or without Isd11-GFP. (E) Anti-HA, anti-HSP60, or anti-GFP co-IP/WB studies of Dd2 parasites episomally expressing mACP-HA$_2$ with or without cyt c$_1$-GFP. (F) Anti-HA co-IP or nickel-nitrilotriacetic acid (Ni-NTA) pulldown and WB studies of *E. coli* bacteria recombinantly expressing full-length mACP-HA$_2$ and Isd11-His$_6$ (WT or LYR/AAA mutant) and probed with anti-HA and anti-His$_6$ antibodies. All lanes are from the same blot and were processed identically but were cropped from non-contiguous lanes. (G) Coomassie-stained SDS-PAGE gel of recombinant His$_6$-Isd11 and Δ2-50 mACP-HA$_2$ purified from *E. coli* by Ni-NTA pulldown of His$_6$-Isd11 and size-exclusion chromatography (SEC). Full images for all WB experiments are shown in Fig. S5.

Fig. 3. Structural modeling and functional tests of the divergent Phe113 residue of parasite mACP. (A) X-ray crystallographic structure of human Isd11 bound to *E. coli* ACP highlighting the central role of the acyl-Ppant group of ACP and R6 residue of Isd11 in stabilizing this complex (PDB entry 5USR). (B) Rosetta-based, energy-minimized structural model of the *P. falciparum* mACP-Isd11 interface. (C) Nickel-nitrilotriacetic acid (Ni-NTA) pulldown and WB studies of *E. coli* bacteria recombinantly expressing Δ2-50 mACP-HA$_2$ (WT or F113A mutant) and Isd11-His$_6$ and probed with anti-HA and anti-His$_6$ antibodies. All lanes are from the same blot and were processed identically but were cropped from non-contiguous lanes. The uncropped WB is shown in Fig. S5E. (D) Continuous growth assays of synchronous mACP-aptamer/TetR-DOZI parasites episomally expressing WT or F113A mACP-HA$_2$ and grown in the presence or absence of aTc. Data points and error bars are the average and standard deviation from two biological replicates. The presence of an HA epitope on both the endogenous and episomal mACP prevented determination of endogenous mACP expression differences ±aTc by WB.

Fig. 4. Loss of mACP destabilizes Nfs1 and the Rieske protein. Western blot analysis of lysates from mACP-aptamer/TetR-DOZI parasites episomally expressing Rieske-HA$_2$ that were grown ±aTc for 3 days and probed for EF1α and (A) Nfs1 or (B) HA.

Fig. 5. Loss of mACP causes ETC failure and sensitizes parasites to proguanil. Continuous growth assays of synchronous mACP-aptamer/TetR-DOZI parasites grown (A) ±aTc, ±atovaquone (Atov, 100 nM), ±decyl-ubiquinone (dQ, 15 μM), and (B) ±proguanil (Prog, 1 μM). Data points and error bars are the average and standard deviation from 2-3 biological replicates. (C) Fluorescence microscopy images of live mACP-aptamer/TetR-DOZI Dd2 parasites cultured 3 days ±aTc, 2 days ±5 μM proguanil, and stained with Hoechst or Mitotracker Red (10 nM). (D) Statistical analysis of MitoTracker signal for 40-50 total parasites from each condition in panel C from two independent experiments. For clarity, error bars are not shown but standard errors of the mean were ≤10% in all cases. Cell percentage differences were analyzed by two-tailed unpaired t-test (p values in parentheses, ns = not significant).

Fig. 6. Schematic model for the impact of mACP knockdown (-aTc) on Fe-S cluster metabolism in *P. falciparum* parasites. Based on studies in yeast (37), eukaryotic mitochondria are proposed to export a sulfur-containing intermediate (X-S) of uncertain identity that is required for Fe-S cluster assembly and utilization in the cytoplasm and nucleus. For simplicity, the mACP/Isd11/Nfs1 complex is shown with only one monomer for each protein, rather than the functional dimer (26). Illustration by Megan Okada.