Quantification and image-derived phenotyping of retinal ganglion cell nuclei in the *nee* mouse model of congenital glaucoma

Carly J. van der Heide[1], Kacie J. Meyer[1], Hannah E. Mercer[1], Adam Hedberg-Buenz[1,3], Michael G. Anderson[1,2,3*]

[1]Department of Molecular Physiology and Biophysics, University of Iowa, Iowa City, IA

[2] Department of Ophthalmology and Visual Sciences, University of Iowa, Iowa City, IA

[3]VA Center for the Prevention and Treatment of Visual Loss, Iowa City VA Health Care System, Iowa City, IA

*Corresponding author:

Dr. Michael G. Anderson, Department of Molecular Physiology and Biophysics, 3123 Medical Education and Research Facility, 375 Newton Road, Iowa City, IA 52242, (319) 355-7839 (telephone), (319) 335-7330 (FAX), michael-g-anderson@uiowa.edu

ABSTRACT

The *nee* mouse model exhibits characteristic features of congenital glaucoma, a

common cause of childhood blindness. The current study of *nee* mice had two components.

First, the time course of neurodegeneration in *nee* retinal flat-mounts was studied over time

using a retinal ganglion cell (RGC)-marker, BRN3A; a pan-nuclear marker, TO-PRO-3; and H&E

staining. Based on segmentation of nuclei using ImageJ and RetFM-J, this analysis identified a

rapid loss of BRN3A$^+$ nuclei from 4–15 weeks of age, with the first statistically significant

difference in average density compared to age-matched controls detected in 8-week-old cohorts

(49% reduction in *nee*). Consistent with a model of glaucoma, no reductions in BRN3A$^-$ nuclei

were detected, but the combined analysis indicated that some RGCs lost BRN3A marker

expression prior to actual cell loss. These results have a practical application in the design of

experiments using *nee* mice to study mechanisms or potential therapies for congenital

glaucoma. The second component of the study pertains to a discovery-based analysis of the

large amount of image data with 748,782 segmented retinal nuclei. Using the automatedly

collected region of interest feature data captured by ImageJ, we tested whether RGC density of

glaucomatous mice was significantly correlated to average nuclear area, perimeter, Feret

diameter, or MinFeret diameter. These results pointed to two events influencing nuclear size.

For variations in RGC density above approximately 3,000 nuclei/mm^2 apparent spreading was

observed, in which BRN3A$^-$ nuclei—regardless of genotype—became slightly larger as RGC

density decreased. This same spreading occurred in BRN3A$^+$ nuclei of wild-type mice. For

variation in RGC density below 3,000 nuclei/mm^2, which only occurred in glaucomatous *nee*

mutants, BRN3A$^+$ nuclei became smaller as disease was progressively severe. These

observations have relevance to defining RGCs of relatively higher sensitivity to glaucomatous

cell death and the nuclear dynamics occurring during their demise.

1. Introduction

Mice homozygous for the spontaneously arising $Sh3pxd2b^{nee}$ mutation (i.e. *nee* mice) exhibit anterior segment dysgenesis and features of severe congenital glaucoma. In two different genetic backgrounds, the original C57BL/10-related and a congenic C57BL/6J background, *nee* mice develop high intraocular pressure (IOP) followed by retinal ganglion cell (RGC) loss which is profound by 3 months of age (Daniel et al., 2019; Mao et al., 2011). Along with congenital glaucoma, *nee* mice exhibit craniofacial, cardiac and skeletal abnormalities that recapitulate Frank-Ter Haar syndrome (Mao et al., 2009), a rare human disease caused by recessive mutations in the homologous gene, *SH3PXD2B* (Iqbal et al., 2010). Thus, *nee* mice directly model a known cause of glaucoma in humans. The manifestation of glaucoma in *nee* mice is also notable among animal models of glaucoma because the disease is spontaneous, severe, and fully penetrant (Mao et al., 2011), which are all experimental advantages.

While RGC loss in *nee* mice is known to be relatively severe and early onset, refining the precise time-course of RGC loss in *nee* could uncover new insights and is a prerequisite needed to inform the design of many future experiments, such as utilizing the *nee* model in tests of neuroprotective agents. To achieve this, we have characterized the progression of RGC death in young *nee* mice using ImageJ in a high-throughput image analysis technique that quantified thousands of cell nuclei in the inner retina of each mouse retina. In addition to providing counts of RGC nuclei, our image analysis data included information about the size and shape of nuclei. In a discovery-based approach, we used these data to explore whether RGC density of glaucomatous mice was significantly correlated to average nuclear area, perimeter, Feret diameter, or MinFeret diameter. Combined, the results identify ages past 8 weeks as having statistically detectable differences in RGC density, identify a caveat regarding loss of RGC marker expression prior to RGC cell loss in some cells of *nee* mice, and point to

50   the possible ways in which automatedly collected image data can be used to discover new

51   disease-associated phenotypes.

53   **2. Materials and methods**

55   *2.1 Animal Husbandry*

57   Generation of N10 congenic B6.*Sh3pxd2b^{nee}* mice has been previously described

(Daniel et al., 2019). The *nee* strain was maintained in heterozygote X heterozygote

intercrosses, with littermates homozygous for the mutant (abbreviated throughout as "*nee*") or

wild-type (abbreviated throughout as "WT") alleles primarily utilized in this study. As indicated in

the text, the SD-OCT imaging study utilized both WT and heterozygous mice (referred to as

"non-mutant", abbreviated throughout as "non-mut") to increase the sample size of controls.

Mice were housed at the University of Iowa Animal Research Animal Facility, where they were

maintained on a 4% fat NIH 31 diet provided ad libitum, housed in cages containing dry bedding

(Cellu-dri; Shepherd Specialty Papers, Kalamazoo, MI), and kept in a 21⁰C environment with a

12-h light: 12-h dark cycle. All experiments included both male and female mice. All mice were

treated in accordance with the Association for Research in Vision and Ophthalmology

Statement for the Use of Animals in Ophthalmic and Vision Research. All experimental

protocols were approved by the Institutional Animal Care and Use Committee of the University

of Iowa.

2.2 Preparation of retinal flat-mounts for RGC quantification

74   Mice were euthanized using CO_2 asphyxiation followed by cervical dislocation and eyes

were immediately placed in cold 4% paraformaldehyde for 10 minutes. Posterior cups were

dissected in PBS and replaced in cold 4% paraformaldehyde for 1 hour. Posterior cups were

washed with PBS, and permeabilized with 0.3% Triton-X 100 in PBS (PBST) overnight at 37°C.

All subsequent steps were carried out at room temperature unless otherwise specified. Posterior

cups were incubated in 3% H_2O_2-1% $NaHPO_4$ for 2 minutes and then retinas were dissected in

PBS. Retinas were replaced in 3% H_2O_2-1% $NaHPO_4$ for 1 hour. Retinas were washed with

PBST three times for 15 minutes, and then further permeabilized with PBST at -80°C for 15

minutes. Retinas were washed again with PBST twice for 15 minutes, then blocked with 2%

normal donkey serum in PBST overnight at 4°C. Retinas were then incubated with anti-BRN3A

antibody (C-20, 1:200; Santa Cruz Biotechnology, Dallas, TX) in PBS with 2% normal donkey

serum, 1% Triton-X 100, and 1% DMSO for two overnights at 4°C. Retinas were washed for 5

minutes in PBS with 2% Triton X-100, then washed further with PBST four times for 15 minutes.

Then retinas were incubated with a donkey anti-goat Alexa 488-conjugated secondary antibody

(A11055, 1:200; Life Technologies, Madison, WI) in PBS with 5% normal donkey serum, 1%

Triton-X 100, and 1% DMSO for two hours. Retinas were washed twice for 15 minutes in PBST,

then incubated in 1 μM TO-PRO-3 (T3605; Thermo Fisher Scientific, Waltham, MA) in PBS for

15 minutes, and washed again with PBST three times for 15 minutes. Retinas were then

washed briefly with PBS and mounted using Aqua-Mount (Lerner, Pittsburgh, PA). Individual

retinas that were studied by both immunolabeling and hematoxylin/eosin (H&E) staining were

labeled as described above, followed by soaking off coverslips by submersion of the slide in

distilled water. H&E staining was performed under the following conditions: hematoxylin (1

minute), rinse in double-distilled water, acid alcohol (3 seconds), rinse in water, 80% ethanol (1

minute), eosin (2 seconds), 95% ethanol (30 seconds), 100% ethanol (30 seconds), and

xylenes (twice for 30 seconds each time). MM 24 mounting media (Leica, Wetzlar, Germany)

was used to apply coverslips to slides.

2.3 Microscopy and quantification of nuclei in the inner retina

Immunolabeled retinas were photographed using a Zeiss LSM710 confocal microscope

(Carl Zeiss, Oberkochen, Germany). Images were collected at ~538X total magnification

(1024x1024 px, 0.18 mm^2 image area) from non-overlapping fields at each of three zones of

eccentricity (n = 12 images total: 4 central, 4 mid-peripheral, 4 peripheral). Photos were

captured in a single plane of focus at the level of the RGC layer. Images were processed in

ImageJ with the Enhance Local Contrast (CLAHE) process (blocksize 125, histogram bins 256,

maximum set to 3 for BRN3A-labeled images and maximum set to 10 for TO-PRO3-labeled

images), the Subtract Background tool with rolling ball radius set to 35 pixels for BRN3A and 20

pixels for TO-PRO-3, followed by the Smooth tool. The TO-PRO-3 images were additionally

processed with Gaussian blur (sigma set to 0.5) and the Enhance contrast process with 10%

saturated pixels and application of histogram equalization. Images were then converted to

binary using Huang thresholding. Binary images were further processed using the Open,

Watershed, and Fill Holes functions. To count BRN3A$^+$ nuclei, the Analyze Particles function

was applied to the BRN3A images with particle size set to 38–150 μm^2 and circularity 0.55–1.

The segmentation of BRN3A$^+$ nuclei was saved as a region of interest and subtracted from the

TO-PRO-3 images. Then, the Analyze Particles function was applied to TO-PRO-3 images with

size set to 20–150 μm^2 and circularity of 0.4–1 to quantify the BRN3A$^-$, TO-PRO-3$^+$ population.

Individual retinas that were studied by both immunolabeling and H&E staining were

photographed and quantified as described above, followed by H&E staining and imaging with a

light microscope (BX52; Olympus, Tokyo, Japan) equipped with a digital camera (DP72;

Olympus, Tokyo, Japan). H&E-stained nuclei were quantified with RetFM-J as previously

described (Hedberg-Buenz et al., 2016c).

2.4 Optical coherence tomography

Mice received an intraperitoneal injection of a mixture of ketamine/xylazine (87.5 mg/kg

Ketamine (VetaKet®, AKORN, Lake Forest, IL)/12.5 mg/kg Xylazine (Anased, Lloyd

Laboratories®, Shenandoah, IA). Upon anesthesia, eyes were hydrated with balanced salt

solution (BSS; Alcon Laboratories, Fort Worth, TX). To obtain retinal images, the tear film was

wicked away and eyes were imaged with a Bioptigen spectral domain optical coherence

tomographer (SD-OCT; Bioptigen, USA) using the mouse retinal bore. The volume intensity

projection was centered on the optic nerve. Following imaging, eyes were hydrated with

lubricant ophthalmic ointment (Artificial Tears, AKORN, Lake Forest, IL) and mice were provided

supplemental indirect warmth for anesthesia recovery. Retinal ganglion cell complex (RGCC;

inner limiting membrane to the innermost border of the inner nuclear layer) thickness was

measured from retinal images within the Bioptigen InVivoVue Clinic software four times per eye

using vertical angle-locked B-scan calipers.

2.5 Statistics

Nuclear quantification data were analyzed using 2-way ANOVA and corrected for

multiple comparisons using Sidak's multiple comparisons test with a single pooled variance.

Correlations were analyzed for significance using linear regression. Data comparing nuclear

area and retinal area were analyzed by multiple two-tailed Student's *t*-tests and corrected for

multiple comparisons using the two-stage step-up method of Benjamini, Krieger and Yekutieli.

The power calculations noted in the Discussion was performed using the Piface Java applet by

Russell V. Lenth (Lenth, 2006) with an analysis for two-sample *t*-test (pooled or Satterthwaite).

For the BRN3A[+] cell density, the assumptions entered are sigma1 = 794.2 (standard deviation

of *nee* BRN3A[+] cell density at 8 weeks old), sigma2 = 162.0 (standard deviation of WT BRN3A[+]

cell density at 8 weeks old), alpha = 0.05, true difference of means = 870.2 (50% of the BRN3A[+]

cell loss in *nee* mice at 8 weeks old compared to WT littermates), power = 0.8981, and solving

for sample size. For the retinal ganglion cell complex (RGCC) thicknesses measured by OCT,

the assumption entered are sigma1 = 14.3 (standard deviation of *nee* RGCC thickness at 8

weeks old), sigma2 = 1.7 (standard deviation of non-mut RGCC thickness at 8 weeks old), true

difference of means = 11.7 (50% of the RGCC thickness decrease in *nee* mice at 8 weeks old

compared to non-mut littermates), power = 0.9019, and solving for sample size.

3. Results

3.1 Severe RGC loss in nee mice by 8 weeks of age

To characterize the early progression of congenital glaucoma in *nee* mice, we quantified

RGC loss in *nee* and WT littermates from 4–15 weeks of age. Immunolabeling of retinal flat-

mounts with an anti-BRN3A antibody showed a progressive loss of BRN3A$^+$ RGCs (Fig. 1A). At

4 weeks of age, quantified data showed a significant loss of BRN3A$^+$ nuclei in all retinas by 8

weeks of age (Fig. 1B), whereby *nee* mice exhibited a 49% reduction in the average density of

BRN3A$^+$ nuclei (1,801.2 ± 794.2 nuclei/mm^2) compared to WT littermates (3,541.3 ± 162.0

nuclei/mm^2; p < 0.05, 2-way ANOVA with Sidak's multiple comparisons correction). BRN3A$^+$

RGC densities in *nee* mice younger than 8 weeks of age were highly variable, with some eyes

severely affected while others were not different from WT littermates.

To further evaluate *nee* mice as a model of glaucoma, we tested whether non-RGC cell

types die in *nee* retinas by quantifying nuclei labeled with TO-PRO-3 but negative for BRN3A

(BRN3A$^-$). There was no reduction in the density of BRN3A$^-$ nuclei of *nee* mice in any age

group (Fig. 1C). Oddly, *nee* mice exhibited a 16% *higher* density of BRN3A$^-$ nuclei at 15 weeks

(5,253.6 ± 176.0 nuclei/mm^2) compared to WT littermates (4,393.4 ± 142.3 nuclei/mm^2 p =

0.0004, 2-way ANOVA with Sidak's multiple comparisons correction). Suspecting that the

increased density of BRN3A⁻ nuclei observed in *nee* mice was due to a loss of marker

expression, not cell loss, these retinas were stained with H&E and nuclei of all cell types were

counted (Hedberg-Buenz et al., 2016c). In *nee* mice aged 4–8 weeks, there was a 21%

reduction in the density of BRN3A⁺ nuclei (2,909 ± 817 nuclei/mm^2) compared to WT (3,686 ±

265 nuclei/mm^2; $p < 0.01$; 2-way ANOVA with Sidak's multiple comparisons correction), but the

density of total nuclei stained by H&E exhibited no corresponding cell loss (7,675 ± 587 in *nee*

vs 7,797 ± 228 nuclei/mm^2 in WT; Fig. 1D), a discrepancy that suggests loss of only BRN3A

immunoreactivity. Slightly older *nee* mice (12–15 weeks old) exhibited a 62.3% loss of BRN3A⁺

RGCs (1,299 ± 873 in *nee* vs 3,444 ± 120 nuclei/mm^2 in WT; $p < 0.0001$; 2-way ANOVA with

Sidak's multiple comparisons correction) along with a 18.5% decrease in total cells stained with

H&E (6,373 ± 646 in *nee* vs 7,816 ± 287 nuclei/mm^2 in WT). Because RGCs constitute

approximately 41–50% of total RGC-layer cells in C57BL/6J mice (Jeon et al., 1998; Schlamp et

al., 2013), if only RGCs were lost, an 18.5% decrease in total cells would correspond to a 37–

45% decrease in RGCs. The RGC loss we detected using BRN3A immunolabeling is 17–25%

higher than the H&E estimate, a difference similar to the 21% discrepancy we detected in the

younger cohort. Conclusively, we estimate that ~20% of RGCs in glaucomatous *nee* mice stop

expressing BRN3A prior to actual RGC loss.

Non-invasive imaging of the retina was performed in 8-week-old glaucomatous *nee* mice

using SD-OCT. Corresponding to the severe loss of RGCs, the RGC complex is significantly

thinner in 8-week-old *nee* mice (n = 6) compared to non-mutant (HET and WT genotypes

combined; n = 5) littermates (32.9 ± 14.3 μm vs. 56.3 ± 1.7 μm; $p < 0.01$; Student's two-tailed *t*-

test; Fig. 2). We also observed differences in the appearance of optic nerve head depth and the

peripapillary retina. Retinas of *nee* mice typically had an overall normal lamination, though 4/21

retinas examined at 8 weeks of age had lamination abnormalities and/or retinal rosettes.

3.2 Size characteristics of surviving BRN3A$^+$ RGCs decrease with increasing severity of

glaucomatous damage in nee mice

207   Using the large amount of systematically collected feature data for the 748,782

segmented retinal nuclei underlying our study of *nee* mice, we examined whether any of the

feature parameters automatedly measured by ImageJ correlated to disease stage. Region of

interest feature data captured by ImageJ was averaged for each retina and tested for correlation

to BRN3A$^+$ cell density (Fig.). Three populations were considered: WT controls, *nee* retinas

with no or mild disease (*nee^{norm}*; within 2 SD of WT density of BRN3A$^+$ nuclei), and *nee* retinas

with moderate to severe disease (*nee$^{mod-sev}$*; < 2 SD of WT density of BRN3A$^+$ nuclei).

214   Among BRN3A$^+$ RGCs (Fig. A-D), WT retinas showed a significant correlation between

average nuclear area and density (Fig. A) in which average nuclear size slightly increased as

density decreased across the normal range of densities (3,238-4,203 nuclei/mm^2; mean =

3,577, 1 SD = 241.5). Contrasting this, the RGCs of *nee$^{mod-sev}$* retinas showed an opposite trend

that was also significant, in which nuclei became progressively smaller on average as density

fell during disease. RGCs of *nee^{norm}* retinas had no detectable correlation between nuclear area

and density. Similar trends and statistical associations were detected in plots of BRN3A$^+$ density

versus nuclear perimeter (Fig. B), Feret diameter (Fig. C), and MinFeret diameter (Fig. D).

Among BRN3A$^-$ nuclei (Fig. E-H), WT retinas again showed a significant correlation between

average nuclear area and density (Fig. E) in which BRN3A$^-$ nuclei became slightly larger with

decreasing density of BRN3A$^+$ nuclei. Unlike RGCs, the BRN3A$^-$ population of nuclei in *nee^{mod-}*

^{sev} retinas showed no significant correlation to density. Also differing from RGCs, the BRN3A$^-$

population of nuclei in *nee^{norm}* retinas mirrored the trends of WT cells, with average nucleus size

slightly increasing as density decreased. Similar trends and statistical associations were

detected in plots of BRN3A$^-$ density versus nuclear perimeter (Fig. F), Feret diameter (Fig. G),

and MinFeret diameter (Fig. H), with the sole exception of one association between BRN3A$^-$

density and MinFeret diameter in *nee^mod-sev* retinas that met statistical significance with a slim

margin (p = 0.049). In sum, these results point to two events influencing nuclear size. In almost

all analyses (regardless of genotype or marker) there was a slight enlargement of nuclei as

RGC density decreased. At approximately 3,000 nuclei/mm^2 there was an inflection point in the

trend lines, below which only the population of viable RGC nuclei, but not BRN3A$^-$ cells, had

nuclei that became smaller as disease was progressively severe.

3.3 RGCs with small nuclei account for a greater proportion of the surviving population in severe

stages of nee glaucoma

We next sought to explore what phenomena might be causing the average nuclear size

to be decreasing through disease progression. First, we analyzed the population distribution of

BRN3A$^+$ nuclear size in *nee^mod-sev* mice compared to WT littermates (Fig. 4). When all retinas

were grouped together, the average size of BRN3A$^+$ nuclei in *nee^mod-sev* retinas was 6.8% larger

(84.5 ± 21.5μm^2) than WT (79.1 ± 18.3μm^2; p < 1E10^{-15}; Fig. 4A) presumably because many

retinas with less severe RGC loss had an average nuclear size that was larger than WT (Fig.

4A, B). However, a limitation of combining data in this way was that the healthiest *nee^mod-sev*

retinas, those with the greatest number of cells, were overrepresented. Thus, instead of

studying a size distribution for all retinas of a genotype together, we next analyzed distributions

for each retina individually.

Retinas from WT mice consistently exhibited a broad distribution of BRN3A$^+$ nuclear size

with a preponderance of small nuclei but a long right-hand tail (Fig. 4C, E). In these retinas, the

average BRN3A$^+$ nuclear size was larger than the median in every case. An example of such a

retina is indicated with a *black arrowhead* in Fig. 4 B&C, in this instance a retina with 3,238

BRN3A$^+$ nuclei/mm^2 (Fig. 4B) having a long right-hand tail (Fig. 4C), with average and median

nuclear sizes of 80.9 ± 18.1 μm^2 and 78.1 μm^2, respectively.

In contrast, the size distributions were highly variable among *nee*^{mod-sev} retinas. Some

nee^{mod-sev} eyes with comparatively modest RGC loss had a rightward shift in the entire

histogram, whereby there was an increase in both the median and the mean relative to WT, with

minimal change in the shape of the curve. An example is illustrated by the *red arrowhead* in Fig.

4 B&C. If the greater average nuclear size were caused by a preferential loss of smaller nuclei,

the left part of the curve would decrease, and the right side would reciprocally increase (note the

Y axis represents the percentage of total BRN3A$^+$ nuclei). Because the shape of the histogram

of this modestly affected eye only changed minimally while the entire curve was shifted

rightward, the increased average size was not a result of preferential loss of the smallest cells

and was more likely reflective of nuclear enlargement broadly occurring in most RGCs

regardless of their initial size. In severe disease, the shape of the histogram was markedly

different. Severely affected *nee*^{mod-sev} eyes had fewer large nuclei and a larger proportion of

smaller RGCs. An example is illustrated by the *red asterisk* in Fig. 4 B&C (in this instance,

combining data from two severely affected retinas, from a single mouse, with very similar

BRN3A$^+$ densities—so that a larger number of nuclei, similar to the other comparisons, could be

considered). The right-hand tail of the curve was attenuated and there was a corresponding

reciprocal peak in the left-hand tail of the curve, representing an increased proportion of small

RGC nuclei relative to the total population. In this case, the smaller average nuclear size in

severely affected eyes reflected a greater proportion of small nuclei in the surviving RGC

population.

To test whether these differences in the nuclear size distribution were specific to RGCs,

we plotted histograms of BRN3A$^-$ nuclear size quantified from the same eyes and did not

observe as pronounced differences in histogram shape or position of the major peak (Fig. 4D).

As expected for a population likely consisting mostly of displaced amacrine cells (Drager and

Olsen, 1981; Janssen et al., 2013), the average size of BRN3A$^-$ nuclei was smaller than

BRN3A$^+$ nuclei.

To broadly visualize how the shape of the nuclear size distributions related to disease

state, distributions were analyzed individually for all retinas and graphed in a box and whisker

plot (Fig. 4E). In this representation, every retina was normalized to its median to allow for

comparisons of the shape of the distribution. In WT retinas, the mean value was consistently

greater than the median, reflecting the long right-hand tail of the distribution. In contrast, many

nee^mod-sev retinas exhibited a smaller difference between the median and the mean, where the

mean was displaced leftward towards the median, suggesting an attenuated right-hand tail of

the curve and a greater proportion of smaller nuclei compared to the WT cell population. In

some *nee^mod-sev* retinas, the mean was displaced even farther to the left beyond the median,

signifying a longer left-tailed distribution with proportionately more RGC nuclei of small size. The

greater proportions of small nuclei in *nee^mod-sev* retinas was also indicated by a decrease in the

average of the 10th, 25th and 90th percentile values of *nee^mod-sev* retinas compared to WT.

4. Discussion

In this study with glaucomatous *nee* mice, ImageJ was used to quantify some traditional

phenotypes commonly needed in studies of glaucoma (density of variously defined cell

populations over time), as well as some novel phenotypes (size metrics for automatedly

detected regions of interest) whose relevance to glaucoma was unknown. The methodology for

RGC nuclei quantification has some elements of advancement within it, but the focus here has

been on its application.

The first component of the study was to study the time course of neurodegeneration in

nee retinal flat-mounts over time. These experiments indicated that *nee* mice have a 49%

reduction in BRN3A$^+$ RGCs by 8 weeks of age. No eyes escaped disease; every retina from

mice aged 8 weeks or older was affected. However, some cases of severely affected retinas

were already detectable at 4 weeks of age (our youngest cohort). While the early onset and

complete penetrance of glaucoma are advantage of the *nee* model for testing neuroprotective

therapies, it is important to recognize that the model does have some heterogeneity in age of

onset and treatments would presumably need to be initiated early. Based on the current

findings, initiation of treatments in newborn mice no later than 4 weeks of age appears

warranted, and in some cases, treating pregnant/nursing mothers may be advantageous. From

the BRN3A$^+$ cell densities measured in 8-week-old WT and *nee* mice, we can predict that a

neuroprotection of 50% of the RGCs otherwise expected to lose BRN3A marker expression

and/or die from glaucoma should be detectable with 90% power (α = 0.05) using a two-tailed *t*-

test with cohorts of 11 samples per treatment group. From the RGCC thickness measured in 8-

week-old non-mut and *nee* mice, we can predict that RGCC thickness increases of 50% of the

thickness lost in *nee* mice should be detectable with 90% power (α = 0.05) using a two-tailed *t*-

test with cohorts of 18 mice per sample group. Thus, tests of neuroprotective agents for *nee*-

mediated early onset glaucoma are feasible for this model by quantifying BRN3A$^+$ cell density

and/or using OCT to measure RGCC thickness, with BRN3A$^+$ cell density requiring fewer mice

and directly quantifying RGC loss. Since a subset of RGCs in the *nee* model lose BRN3A

expression prior to cell death, it is important to note that this power calculation is specific to use

of BRN3A to mark RGCs. Use of other RGC markers may give different results depending on if

or when RGCs lose marker expression prior to cell death. Quantification of cells that are

nuclear-marker positive and RGC-marker negative, as well as use of H&E to count all retinal

nuclei, can help parse loss of marker expression versus cell death as we have done here for the

BRN3A RGC marker. Measurement of the RGCC via OCT requires more mice per cohort and is

an indirect measurement of RGC loss but is a non-invasive assay that has the benefit of

providing same-animal longitudinal data. Therefore, our current data should aid in the design of

meaningful future experiments utilizing the *nee* model of early onset glaucoma.

An additional insight gained through these experiments was that RGC quantification data

obtained from *nee* using immunolabeling techniques should be interpreted with caution. In our

study, we detected an appreciable loss of immunoreactivity in RGCs. Approximately 20% of

RGCs in *nee* mice aged 4–15 weeks lost BRN3A expression prior to cell death. This finding

aligns with previous reports of reduced gene expression in models of acute RGC injury (Huang

et al., 2006; Schlamp et al., 2001) and in DBA/2J mice (Soto et al., 2008). Accordingly, the 49%

reduction in RGC density detected in the 8-week-old *nee* cohort was probably an overestimate

of RGC loss—assuming 20% of the nuclei likely lost BRN3A immunoreactivity, the true degree

of RGC death was likely closer to 30%. Importantly, no loss of BRN3A$^-$ nuclei was observed in

nee retinas. Sparing of non-RGC types is supportive of *nee* as a model of glaucoma, as

opposed to other types of retinal neurodegeneration not specific to RGCs. Many studies with

mouse models of glaucoma do not report the propensity for neuronal damage to non-RGC cell-

types in the inner retina, such as displaced amacrine cells (Pang and Clark, 2020), but the

specificity of RGC loss observed in *nee* mice do distinguish it from models based upon NMDA

excitotoxicity (Hama et al., 2008; Siliprandi et al., 1992) or retinal ischemia (Dijk and Kamphuis,

2004; Ju and Kim, 2011) in which profound loss of other cell-types is known to occur.

Several mouse models relevant to early onset developmental forms of glaucoma have

been described (Akula et al., 2020; Chang et al., 2001; Chen and Gage, 2016; Ishibashi et al.,

1999; Iwao et al., 2009; Kroeber et al., 2010; Labelle-Dumais et al., 2020; Lachke et al., 2011;

Smith et al., 2000; Thomson et al., 2017; Tolman et al., 2021). Among them, many studies have

focused on phenotypes of the anterior segment and there have been relatively few in depth

studies of RGC damage. One exception, a detailed study by Daniel et. al. (Daniel et al., 2019),

utilized *nee* mice to study the susceptibilities of 4 different RGC sub-types to glaucomatous

damage. The overall levels of RGC damage in the study by Daniel et al. were assessed with

NeuN immunolabeling (manual counting of 6–8 retinal flat-mounts per age; 8 sample regions

per retina, each 0.09 mm^2; 4 central and 4 peripheral) and detected a steady age-related

decline in RGC density with statistically significant differences as early as 30 days of age (2,349

± 126 cell/mm^2 in wild-type, which reduced to 1,785 ± 161 cells/mm^2 by day 30 in *nee*

homozygotes). In comparison to the current findings, which had different goals and used

different methodology (BRN3A immunolabeling; automated segmentation of 3–7 retinal flat-

mounts per age; 12 sample regions per retina, each 0.18 mm^2; 4 central, 4 mid-peripheral, and

4 peripheral), both studies detected an early onset glaucomatous loss of RGCs but the study by

Daniel detected a statistically distinguishable loss earlier. Several factors likely contributed to

these slightly different findings. First, the current study used BRN3A immunolabeling, which we

found is lost in some RGCs prior to actual cell death, perhaps confounding an ability to detect

differences. Second, there were notable differences in total RGC densities, even among wild-

type controls between the two studies (~2,400 cells/mm^2, Daniel; ~3,600 cells/mm^2, current

study), hinting that the study by Daniel had a greater emphasis on peripheral sites where RGC

densities are lowest.

The second component of the current study involved a discovery-based analysis of the

large amount of automatedly collected region of interest feature data captured by ImageJ. At the

outset, it was unclear whether subtle differences in these "phenotypes" would add any

information to understanding of glaucoma. At least hundreds, if not thousands, of previous

studies have undoubtedly qualitatively observed RGC nuclei and a few have measured nuclear

areas. In studying a much larger number of nuclei than practical manually, at a micron-level

which might surpass the ability of human observers to notice trends, the broadest question of

the current analysis was "would image-based phenotyping detect anything new?". The results

did detect significant trends, pointing to two events influencing nuclear size. In almost all

analyses (regardless of genotype or marker) there was a slight enlargement of nuclei as RGC

density decreased. At approximately 3,000 nuclei/mm^2 there was an inflection point in the trend

lines, below which only the population of viable RGC nuclei, but not BRN3A$^-$ cells, had nuclei

that became smaller as disease was progressively severe.

One event influencing nuclear size appears to be a passive "spreading out" of nuclei as

space allows. Spreading appears to have a predominant influence on nuclear area in retinas

with densities above 3000 nuclei/mm^2. For most size-related parameters of BRN3A$^+$ and

BRN3A$^-$ nuclei, among both WT and *nee*^{norm} retinas, there was a strong negative correlation

between nuclear size and cell density, whereby average nucleus size increased with decreasing

cell density. We speculate that under normal physiological conditions, neurons generally tend to

be larger in retinas with lower cell densities because space is less confined. The only

exceptions to this finding were a single lack of significance for BRN3A$^+$ nuclei in WT retinas for

MinFeret diameter, and all four of the comparisons for BRN3A$^+$ nuclei in *nee*^{norm} retinas. Among

the *nee*^{norm} retinas, the lack of association might be explained by simultaneously competing

events, a passive spreading out as RGCs begin to be lost that tends to increase average

nuclear size, offset by a preferential loss of larger nuclei that tends to decrease average nuclear

size. The phenomenon of spreading appears to also be present in an anatomical context, where

RGC size is known to vary with respect to eccentricity. RGCs are significantly larger in the

peripheral retina, an area of relatively low cell density compared to the central retina, where

RGCs are smaller and more numerous (Drager and Olsen, 1981; Urcola et al., 2006). RGC

enlargement in some stages of disease has been previously observed in multiple studies (Davis

et al., 2020; Hedberg-Buenz et al., 2016a; Moore and Thanos, 1996; Urcola et al., 2006). For

example, among surviving RGCs in a rat model of chronic IOP elevation induced by microbead

injection (Urcola et al., 2006), in which a loss of 27.2% of RGCs after 24 weeks led to a 11.9%

increase in RGC somal area—a subtle changes in size in response to modest changes in RGC

density similar to those observed here. Urcola et al. described this as "hypertrophy". Because

RGC somal and nuclear areas have previously been shown to be positively correlated (Davis et

al., 2020; Janssen et al., 2013), it seems likely that the "hypertrophy" observed by Urcola et al,

and "spreading" described here, are the same phenomenon. Davis et al. (Davis et al., 2020)

observed RGC nuclear enlargement in the early stages following partial optic nerve transection

and hypothesized that it might involve metabolic responses of the injured RGCs. Because we

have observed spreading in not only glaucomatous RGCs, but also in non-RGCs of mice with

glaucoma and in RGCs of healthy WT mice, we propose a more passive response—though

future studies to study this phenomenon are needed.

A second event influencing nuclear size appears to be a preferential loss of larger RGC

nuclei through progressive stages of glaucoma. For only BRN3A$^+$ nuclei of *nee^{mod-sev}* retinas,

there was a strong positive correlation between RGC nuclear size, whereby average nucleus

size decreased with decreasing cell density. This phenomenon was specific to RGCs and not

detected among BRN3A$^-$ nuclei. In examining size distribution plots, the RGC population in

glaucomatous retinas was markedly different from WT retinas and highly dependent on disease

stage. Healthy WT retinas consistently exhibited a right-skewed distribution of BRN3A$^+$ nuclear

size, in which the average RGC size was always greater than the median, and the right-hand

side of the distribution curve had a long tail. In retinas with modest glaucoma, there was

rightward displacement of the entire distribution curve, reflecting an indiscriminate enlargement

of RGC nuclei of all size. In retinas with advanced glaucoma there was an attenuation of the

right-hand tail of the size distribution, in some cases becoming a left-skewed distribution, and

there was often a second peak in the left-hand tail, indicating an enlarged fraction of small RGC

nuclei.

We speculate that the biological basis for greater proportions of small RGC nuclei in

eyes with severe disease may involve nuclear atrophy and/or size-selective vulnerability to

glaucoma. Some aspects of apoptosis are rapid events, with multiple studies identifying active

apoptosis in only <1–15% of RGCs at any given time (reviewed in (Cordeiro et al., 2011); thus,

many apoptotic events are likely too transient to significantly impact the population average for

nuclear size. In contrast, nuclear atrophy driven by histone deacetylase activity is a sustained

event that can precede BAX-dependent apoptosis in RGCs (Janssen et al., 2013; Schmitt et al.,

2016). In mouse models of optic nerve crush, nuclear atrophy appears within 24 hours, is

maximal at 5 days, and persistent thereafter (Janssen et al., 2013). In comparison to other

events occurring with optic nerve crush, nuclear atrophy precedes both apoptotic nuclear

fragmentation, which typically begins at days 5–7 post-injury, and cell death, which typically

begins after day 10 post-injury (Li et al., 1999). Thus, nuclear atrophy is a sustained event that

could contribute to our current observations with *nee* mice. Alternatively, average RGC size

could decrease with progressive disease because the largest cells are preferentially dying.

Preferential vulnerability of larger RGCs has been a long-standing hypothesis in glaucoma

(Glovinsky et al., 1991; Vickers et al., 1995; Wang et al., 2020), and in mice, has been

supported by an enrichment of small RGCs in old DBA/2J mice with late-onset glaucoma

(Buckingham et al., 2008; Hedberg-Buenz et al., 2016a). The current results establish that this

is also the case for *nee* mice with early onset congenital glaucoma.

There are caveats to consider in the interpretation of these data. First, immunolabeling

was used to quantify only BRN3A$^+$ RGCs. Accordingly, the data most specifically relate to the

surviving RGC population that still expresses BRN3A. Although *Brn3a* appears to be transcribed

in all sub-classes of RGCs (Rheaume et al., 2018), most studies find that immunolabeling fails

to identify some (~3–15%) RGCs (Davis et al., 2020; Galindo-Romero et al., 2011; Rodriguez et

al., 2014). Thus, RGCs that either lost, or never had, detectable BRN3A were potentially

confounding. For example, many melanopsin containing RGCs are not well-labeled by BRN3A

(Sanchez-Migallon et al., 2018; Valiente-Soriano et al., 2014). Because melanopsin-containing

RGCs are relatively rare (~2–3% of the total RGC population)(Valiente-Soriano et al., 2014),

these cells likely had a miniscule impact on our overall findings. Non-RGC cell-types were only

studied in aggregate, not with additional specific markers. This may be of relevance because a

subtle loss of displaced amacrine cells that are coupled to RGCs via gap junctions has been

reported in a microbead-induced model of glaucoma (Akopian et al., 2019). Our current

experimental design, lacking specific markers of displaced amacrine cells and complicated by

RGCs losing BRN3A expression, was not well-matched for detection of this event—though it

would be an interesting area for future work using *nee* mice. Second, with 40+ sub-types of

RGCs now molecularly recognized (Rheaume et al., 2018; Tran et al., 2019), it is probable that

nuclear size may be a susceptibility-factor for many RGCs, but other sub-type specific factors

could also play a role (Tran et al., 2019; Wang et al., 2020). Melanopsin-containing RGCs,

which are somewhat resistant to multiple forms of damage (Cui et al., 2015; Daniel et al., 2019;

Jakobs et al., 2005; Li et al., 2006; Sanchez-Migallon et al., 2018), are often described as an

exception to the "larger = more susceptible" hypothesis, perhaps incorrectly, as their soma are

actually small to average in size and only their dendritic field is comparatively large (Berson et

al., 2010; Coombs et al., 2006; Xiao et al., 2021).

In sum, this study has quantified the severe loss of RGCs in the *nee* mouse model of

congenital glaucoma and used a large amount of simultaneously collected image data related to

nuclear size to phenotype glaucomatous mice in a new way. The results point to competing

events influencing nuclear size, with spreading of multiple cell types occurring with modest

decreases in RGC density and a preferential loss of large RGC nuclei in moderate to severe

cases of RGC loss due to glaucoma. Given the progress in refining approaches using

automated image analysis to quantify RGC soma (Danias et al., 2006; Dordea et al., 2016;

Guymer et al., 2020; Hedberg-Buenz et al., 2016a; Hedberg-Buenz et al., 2016b; Masin et al.,

2021; Salinas-Navarro et al., 2009) and axons (Bosco et al., 2015; Mysona et al., 2020;

Reynaud et al., 2012; Ritch et al., 2020; Teixeira et al., 2014; Zarei et al., 2016) it appears that

the approach that we, and others (Davis et al., 2020), have begun to utilize to phenotype RGCs

in these automated ways may make additional discoveries. Although studies to date have

analyzed only a small number of parameters that were all related to nuclear size, future

experiments could likewise analyze many additional features, some of which may also be

beyond the practical limit for humans to have been led to suspect as important based only on

visual observation.

ACKNOWLEDGEMENTS

This work was supported in part by Merit Review Award (I01 RX001481) from the U.S.

Department of Veterans Affairs RR&D Service and an NIH/NEI Center Support Grant to the

University of Iowa (P30 EY025580). AHB was supported by Training Grant Award

(T32DK112751). The contents do not represent the views of the U.S. Department of Veterans

Affairs or the U.S. Government.

REFERENCES

Akopian, A., Kumar, S., Ramakrishnan, H., Viswanathan, S., Bloomfield, S.A., 2019. Amacrine
cells coupled to ganglion cells via gap junctions are highly vulnerable in glaucomatous mouse
retinas. The Journal of comparative neurology 527, 159-173.
Akula, M., Taiyab, A., Deschamps, P., Yee, S., Ball, A.K., Williams, T., West-Mays, J.A., 2020.
AP-2beta is required for formation of the murine trabecular meshwork and Schlemm's canal.
Exp Eye Res 195, 108042.
Berson, D.M., Castrucci, A.M., Provencio, I., 2010. Morphology and mosaics of melanopsin-
expressing retinal ganglion cell types in mice. The Journal of comparative neurology 518, 2405-
2422.
Bosco, A., Romero, C.O., Breen, K.T., Chagovetz, A.A., Steele, M.R., Ambati, B.K., Vetter,
M.L., 2015. Neurodegeneration severity can be predicted from early microglia alterations
monitored in vivo in a mouse model of chronic glaucoma. Dis Model Mech 8, 443-455.
Buckingham, B.P., Inman, D.M., Lambert, W., Oglesby, E., Calkins, D.J., Steele, M.R., Vetter,
M.L., Marsh-Armstrong, N., Horner, P.J., 2008. Progressive ganglion cell degeneration
precedes neuronal loss in a mouse model of glaucoma. Journal of Neuroscience 28, 2735-
2744.
Chang, B., Smith, R.S., Peters, M., Savinova, O.V., Hawes, N.L., Zabaleta, A., Nusinowitz, S.,
Martin, J.E., Davisson, M.L., Cepko, C.L., Hogan, B.L., John, S.W., 2001. Haploinsufficient
Bmp4 ocular phenotypes include anterior segment dysgenesis with elevated intraocular
pressure. BMC Genet 2, 18.
Chen, L., Gage, P.J., 2016. Heterozygous Pitx2 Null Mice Accurately Recapitulate the Ocular
Features of Axenfeld-Rieger Syndrome and Congenital Glaucoma. Invest Ophthalmol Vis Sci
57, 5023-5030.
Coombs, J., van der List, D., Wang, G.Y., Chalupa, L.M., 2006. Morphological properties of
mouse retinal ganglion cells. Neuroscience 140, 123-136.
Cordeiro, M.F., Migdal, C., Bloom, P., Fitzke, F.W., Moss, S.E., 2011. Imaging apoptosis in the
eye. Eye (Lond) 25, 545-553.
Cui, Q., Ren, C., Sollars, P.J., Pickard, G.E., So, K.F., 2015. The injury resistant ability of
melanopsin-expressing intrinsically photosensitive retinal ganglion cells. Neuroscience 284,
845-853.
Danias, J., Shen, F., Kavalarakis, M., Chen, B., Goldblum, D., Lee, K., Zamora, M.-F., Su, Y.,
Podos, S.M., Mittag, T., 2006. Characterization of retinal damage in the episcleral vein
cauterization rat glaucoma model. Exp Eye Res 82, 219-228.
Daniel, S., Meyer, K.J., Clark, A.F., Anderson, M.G., McDowell, C.M., 2019. Effect of ocular
hypertension on the pattern of retinal ganglion cell subtype loss in a mouse model of early-onset
glaucoma. Exp Eye Res 185, 107703.

Davis, B.M., Guo, L., Ravindran, N., Shamsher, E., Baekelandt, V., Mitchell, H., Bharath, A.A., De Groef, L., Cordeiro, M.F., 2020. Dynamic changes in cell size and corresponding cell fate after optic nerve injury. Sci Rep 10, 21683.

Dijk, F., Kamphuis, W., 2004. An immunocytochemical study on specific amacrine cell subpopulations in the rat retina after ischemia. Brain Res 1026, 205-217.

Dordea, A.C., Bray, M.-A., Allen, K., Logan, D.J., Fei, F., Malhotra, R., Gregory, M.S., Carpenter, A.E., Buys, E.S., 2016. An open-source computational tool to automatically quantify immunolabeled retinal ganglion cells. Exp Eye Res 147, 50-56.

Drager, U.C., Olsen, J.F., 1981. Ganglion cell distribution in the retina of the mouse. Invest Ophthalmol Vis Sci 20, 285-293.

Galindo-Romero, C., Aviles-Trigueros, M., Jimenez-Lopez, M., Valiente-Soriano, F.J., Salinas-Navarro, M., Nadal-Nicolas, F., Villegas-Perez, M.P., Vidal-Sanz, M., Agudo-Barriuso, M., 2011. Axotomy-induced retinal ganglion cell death in adult mice: quantitative and topographic time course analyses. Exp Eye Res 92, 377-387.

Glovinsky, Y., Quigley, H.A., Dunkelberger, G.R., 1991. Retinal ganglion cell loss is size dependent in experimental glaucoma. Invest Ophthalmol Vis Sci 32, 484-491.

Guymer, C., Damp, L., Chidlow, G., Wood, J., Tang, Y.F., Casson, R., 2020. Software for quantifying and batch processing images of brn3a and RBPMS immunolabelled retinal ganglion cells in retinal wholemounts. Translational vision science & technology 9, 28-28.

Hama, Y., Katsuki, H., Suminaka, C., Kume, T., Akaike, A., 2008. Chloride-dependent acute excitotoxicity in adult rat retinal ganglion cells. Neuropharmacology 55, 677-686.

Hedberg-Buenz, A., Christopher, M.A., Lewis, C.J., Fernandes, K.A., Dutca, L.M., Wang, K., Scheetz, T.E., Abràmoff, M.D., Libby, R.T., Garvin, M.K., 2016a. Quantitative measurement of retinal ganglion cell populations via histology-based random forest classification. Exp Eye Res 146, 370-385.

Hedberg-Buenz, A., Christopher, M.A., Lewis, C.J., Meyer, K.J., Rudd, D.S., Dutca, L.M., Wang, K., Garvin, M.K., Scheetz, T.E., Abràmoff, M.D., 2016b. RetFM-J, an ImageJ-based module for automated counting and quantifying features of nuclei in retinal whole-mounts. Exp Eye Res 146, 386-392.

Hedberg-Buenz, A., Christopher, M.A., Lewis, C.J., Meyer, K.J., Rudd, D.S., Dutca, L.M., Wang, K., Garvin, M.K., Scheetz, T.E., Abramoff, M.D., Harper, M.M., Anderson, M.G., 2016c. RetFM-J, an ImageJ-based module for automated counting and quantifying features of nuclei in retinal whole-mounts. Exp Eye Res 146, 386-392.

Huang, W., Fileta, J., Guo, Y., Grosskreutz, C.L., 2006. Downregulation of Thy1 in retinal ganglion cells in experimental glaucoma. Curr Eye Res 31, 265-271.

Iqbal, Z., Cejudo-Martin, P., de Brouwer, A., van der Zwaag, B., Ruiz-Lozano, P., Scimia, M.C., Lindsey, J.D., Weinreb, R., Albrecht, B., Megarbane, A., Alanay, Y., Ben-Neriah, Z., Amenduni, M., Artuso, R., Veltman, J.A., van Beusekom, E., Oudakker, A., Millan, J.L., Hennekam, R., Hamel, B., Courtneidge, S.A., van Bokhoven, H., 2010. Disruption of the podosome adaptor protein TKS4 (SH3PXD2B) causes the skeletal dysplasia, eye, and cardiac abnormalities of Frank-Ter Haar Syndrome. Am J Hum Genet 86, 254-261.

Ishibashi, K., Yamamoto, H., Hatano, M., Koizumi, T., Yamamoto, M., Tokuhisa, T., 1999. Enlargement of the globe with ocular malformations in c-Myc transgenic mice. Jpn J Ophthalmol 43, 201-208.

Iwao, K., Inatani, M., Matsumoto, Y., Ogata-Iwao, M., Takihara, Y., Irie, F., Yamaguchi, Y., Okinami, S., Tanihara, H., 2009. Heparan sulfate deficiency leads to Peters anomaly in mice by disturbing neural crest TGF-beta2 signaling. J Clin Invest 119, 1997-2008.

Jakobs, T.C., Libby, R.T., Ben, Y., John, S.W., Masland, R.H., 2005. Retinal ganglion cell degeneration is topological but not cell type specific in DBA/2J mice. The Journal of cell biology 171, 313-325.

Janssen, K.T., Mac Nair, C.E., Dietz, J.A., Schlamp, C.L., Nickells, R.W., 2013. Nuclear atrophy
of retinal ganglion cells precedes the bax-dependent stage of apoptosis. Invest Ophthalmol Vis
Sci 54, 1805-1815.
Jeon, C.J., Strettoi, E., Masland, R.H., 1998. The major cell populations of the mouse retina. J
Neurosci 18, 8936-8946.
Ju, W.K., Kim, K.Y., 2011. Measuring glutamate receptor activation-induced apoptotic cell death
in ischemic rat retina using the TUNEL assay. Methods Mol Biol 740, 149-156.
Kroeber, M., Davis, N., Holzmann, S., Kritzenberger, M., Shelah-Goraly, M., Ofri, R., Ashery-
Padan, R., Tamm, E.R., 2010. Reduced expression of Pax6 in lens and cornea of mutant mice
leads to failure of chamber angle development and juvenile glaucoma. Hum Mol Genet 19,
3332-3342.
Labelle-Dumais, C., Pyatla, G., Paylakhi, S., Tolman, N.G., Hameed, S., Seymens, Y., Dang,
E., Mandal, A.K., Senthil, S., Khanna, R.C., Kabra, M., Kaur, I., John, S.W.M., Chakrabarti, S.,
Nair, K.S., 2020. Loss of PRSS56 function leads to ocular angle defects and increased
susceptibility to high intraocular pressure. Dis Model Mech 13.
Lachke, S.A., Alkuraya, F.S., Kneeland, S.C., Ohn, T., Aboukhalil, A., Howell, G.R., Saadi, I.,
Cavallesco, R., Yue, Y., Tsai, A.C., Nair, K.S., Cosma, M.I., Smith, R.S., Hodges, E., Alfadhli,
S.M., Al-Hajeri, A., Shamseldin, H.E., Behbehani, A., Hannon, G.J., Bulyk, M.L., Drack, A.V.,
Anderson, P.J., John, S.W., Maas, R.L., 2011. Mutations in the RNA granule component
TDRD7 cause cataract and glaucoma. Science 331, 1571-1576.
Li, R.S., Chen, B.Y., Tay, D.K., Chan, H.H., Pu, M.L., So, K.F., 2006. Melanopsin-expressing
retinal ganglion cells are more injury-resistant in a chronic ocular hypertension model. Invest
Ophthalmol Vis Sci 47, 2951-2958.
Li, Y., Schlamp, C.L., Nickells, R.W., 1999. Experimental induction of retinal ganglion cell death
in adult mice. Invest Ophthalmol Vis Sci 40, 1004-1008.
Mao, M., Hedberg-Buenz, A., Koehn, D., John, S.W., Anderson, M.G., 2011. Anterior segment
dysgenesis and early-onset glaucoma in nee mice with mutation of Sh3pxd2b. Invest
Ophthalmol Vis Sci 52, 2679-2688.
Mao, M., Thedens, D.R., Chang, B., Harris, B.S., Zheng, Q.Y., Johnson, K.R., Donahue, L.R.,
Anderson, M.G., 2009. The podosomal-adaptor protein SH3PXD2B is essential for normal
postnatal development. Mamm Genome 20, 462-475.
Masin, L., Claes, M., Bergmans, S., Cools, L., Andries, L., Davis, B.M., Moons, L., De Groef, L.,
2021. A novel retinal ganglion cell quantification tool based on deep learning. Scientific reports
11, 1-13.
Moore, S., Thanos, S., 1996. Differential increases in rat retinal ganglion cell size with various
methods of optic nerve lesion. Neurosci Lett 207, 117-120.
Mysona, B.A., Segar, S., Hernandez, C., Kim, C., Zhao, J., Mysona, D., Bollinger, K.E., 2020.
QuPath Automated Analysis of Optic Nerve Degeneration in Brown Norway Rats. Transl Vis Sci
Technol 9, 22.
Pang, I.H., Clark, A.F., 2020. Inducible rodent models of glaucoma. Prog Retin Eye Res 75,
100799.
Reynaud, J., Cull, G., Wang, L., Fortune, B., Gardiner, S., Burgoyne, C.F., Cioffi, G.A., 2012.
Automated quantification of optic nerve axons in primate glaucomatous and normal eyes--
method and comparison to semi-automated manual quantification. Invest Ophthalmol Vis Sci
53, 2951-2959.
Rheaume, B.A., Jereen, A., Bolisetty, M., Sajid, M.S., Yang, Y., Renna, K., Sun, L., Robson, P.,
Trakhtenberg, E.F., 2018. Single cell transcriptome profiling of retinal ganglion cells identifies
cellular subtypes. Nat Commun 9, 2759.
Ritch, M.D., Hannon, B.G., Read, A.T., Feola, A.J., Cull, G.A., Reynaud, J., Morrison, J.C.,
Burgoyne, C.F., Pardue, M.T., Ethier, C.R., 2020. Axonet: A deep learning-based tool to count
retinal ganglion cell axons. Scientific reports 10, 1-13.

Rodriguez, A.R., de Sevilla Muller, L.P., Brecha, N.C., 2014. The RNA binding protein RBPMS
is a selective marker of ganglion cells in the mammalian retina. The Journal of comparative
neurology 522, 1411-1443.
Salinas-Navarro, M., Mayor-Torroglosa, S., Jimenez-Lopez, M., Avilés-Trigueros, M., Holmes,
T., Lund, R., Villegas-Pérez, M., Vidal-Sanz, M., 2009. A computerized analysis of the entire
retinal ganglion cell population and its spatial distribution in adult rats. Vision research 49, 115-
126.
Sanchez-Migallon, M.C., Valiente-Soriano, F.J., Nadal-Nicolas, F.M., Di Pierdomenico, J., Vidal-
Sanz, M., Agudo-Barriuso, M., 2018. Survival of melanopsin expressing retinal ganglion cells
long term after optic nerve trauma in mice. Exp Eye Res 174, 93-97.
Schlamp, C.L., Johnson, E.C., Li, Y., Morrison, J.C., Nickells, R.W., 2001. Changes in Thy1
gene expression associated with damaged retinal ganglion cells. Mol Vis 7, 192-201.
Schlamp, C.L., Montgomery, A.D., Mac Nair, C.E., Schuart, C., Willmer, D.J., Nickells, R.W.,
2013. Evaluation of the percentage of ganglion cells in the ganglion cell layer of the rodent
retina. Mol Vis 19, 1387-1396.
Schmitt, H.M., Schlamp, C.L., Nickells, R.W., 2016. Role of HDACs in optic nerve damage-
induced nuclear atrophy of retinal ganglion cells. Neurosci Lett 625, 11-15.
Siliprandi, R., Canella, R., Carmignoto, G., Schiavo, N., Zanellato, A., Zanoni, R., Vantini, G.,
1992. N-methyl-D-aspartate-induced neurotoxicity in the adult rat retina. Visual neuroscience 8,
567-573.
Smith, R.S., Zabaleta, A., Kume, T., Savinova, O.V., Kidson, S.H., Martin, J.E., Nishimura, D.Y.,
Alward, W.L., Hogan, B.L., John, S.W., 2000. Haploinsufficiency of the transcription factors
FOXC1 and FOXC2 results in aberrant ocular development. Hum Mol Genet 9, 1021-1032.
Soto, I., Oglesby, E., Buckingham, B.P., Son, J.L., Roberson, E.D., Steele, M.R., Inman, D.M.,
Vetter, M.L., Horner, P.J., Marsh-Armstrong, N., 2008. Retinal ganglion cells downregulate gene
expression and lose their axons within the optic nerve head in a mouse glaucoma model. J
Neurosci 28, 548-561.
Teixeira, L.B., Buhr, K.A., Bowie, O., Duke, F.D., Nork, T.M., Dubielzig, R.R., McLellan, G.J.,
2014. Quantifying optic nerve axons in a cat glaucoma model by a semi-automated targeted
counting method. Mol Vis 20, 376-385.
Thomson, B.R., Souma, T., Tompson, S.W., Onay, T., Kizhatil, K., Siggs, O.M., Feng, L.,
Whisenhunt, K.N., Yanovitch, T.L., Kalaydjieva, L., Azmanov, D.N., Finzi, S., Tanna, C.E.,
Hewitt, A.W., Mackey, D.A., Bradfield, Y.S., Souzeau, E., Javadiyan, S., Wiggs, J.L., Pasutto,
F., Liu, X., John, S.W., Craig, J.E., Jin, J., Young, T.L., Quaggin, S.E., 2017. Angiopoietin-1 is
required for Schlemm's canal development in mice and humans. J Clin Invest 127, 4421-4436.
Tolman, N.G., Balasubramanian, R., Macalinao, D.G., Kearney, A.L., MacNicoll, K.H.,
Montgomery, C.L., de Vries, W.N., Jackson, I.J., Cross, S.H., Kizhatil, K., Nair, K.S., John,
S.W.M., 2021. Genetic background modifies vulnerability to glaucoma-related phenotypes in
Lmx1b mutant mice. Dis Model Mech 14.
Tran, N.M., Shekhar, K., Whitney, I.E., Jacobi, A., Benhar, I., Hong, G., Yan, W., Adiconis, X.,
Arnold, M.E., Lee, J.M., Levin, J.Z., Lin, D., Wang, C., Lieber, C.M., Regev, A., He, Z., Sanes,
J.R., 2019. Single-Cell Profiles of Retinal Ganglion Cells Differing in Resilience to Injury Reveal
Neuroprotective Genes. Neuron 104, 1039-1055 e1012.
Urcola, J.H., Hernandez, M., Vecino, E., 2006. Three experimental glaucoma models in rats:
comparison of the effects of intraocular pressure elevation on retinal ganglion cell size and
death. Exp Eye Res 83, 429-437.
Valiente-Soriano, F.J., Garcia-Ayuso, D., Ortin-Martinez, A., Jimenez-Lopez, M., Galindo-
Romero, C., Villegas-Perez, M.P., Agudo-Barriuso, M., Vugler, A.A., Vidal-Sanz, M., 2014.
Distribution of melanopsin positive neurons in pigmented and albino mice: evidence for
melanopsin interneurons in the mouse retina. Front Neuroanat 8, 131.

Vickers, J.C., Schumer, R.A., Podos, S.M., Wang, R.F., Riederer, B.M., Morrison, J.H., 1995.
Differential vulnerability of neurochemically identified subpopulations of retinal neurons in a
monkey model of glaucoma. Brain Res 680, 23-35.
Wang, A.Y., Lee, P.Y., Bui, B.V., Jobling, A.I., Greferath, U., Brandli, A., Dixon, M.A., Findlay,
Q., Fletcher, E.L., Vessey, K.A., 2020. Potential mechanisms of retinal ganglion cell type-
specific vulnerability in glaucoma. Clin Exp Optom 103, 562-571.
Xiao, J., Lin, X., Qu, J., Zhang, J., 2021. Morphological and functional diversity of intrinsically
photosensitive retinal ganglion cells. Synapse, e22200.
Zarei, K., Scheetz, T.E., Christopher, M., Miller, K., Hedberg-Buenz, A., Tandon, A., Anderson,
M.G., Fingert, J.H., Abràmoff, M.D., 2016. Automated axon bounting in rodent optic nerve
sections with AxonJ. Scientific Reports 6, 26559.

Fig. 1. Early loss of BRN3A$^+$ RGCs in *nee* mice. (A) Representative images showing retinal
flat-mounts with BRN3A immunolabeling of RGC nuclei (*green*) and all TO-PRO-3 labeled
nuclei (*blue*) in *nee* and WT mice. Confocal images taken from the peripheral region of retinas;
scale bar = 50 µm. **(B)** Density plots of BRN3A$^+$ RGC nuclei in *nee* (*red*) and WT (*black*) retinas
showed a significant loss in RGC density in 8-week-old and older *nee* mice. Note, some eyes in
the 4- and 6-week groups were severely affected. **(C)** Average density of BRN3A$^-$ nuclei in *nee*

retinas had an upward trend with increasing age, with a statistically significant difference at 15
weeks of age. **(D)** Plots of nuclear densities with data from B and C binned by age and
additionally showing density of H&E-detected nuclei. There was no difference in density of H&E
labeled nuclei between *nee* and WT retinas in the 4–8-week bin, even though these same
retinas had a significant reduction in BRN3A$^+$ nuclei; among *nee* retinas in the 12–15-week bin,
the loss of BRN3A$^+$ nuclei, off-set by an increase in BRN3A$^-$ nuclei, approximated the net
change in nuclei detected with H&E. n.s., not significant; *p < 0.05; **p < 0.01; ***p < 0.001;
****p < 0.0001.

Fig. 2. Thinning of retinal ganglion cell complex in 8-week-old *nee* mice. (A)
Representative image showing retinal optical coherence tomography from an 8-week-old WT
mouse. **(B)** Representative image showing retinal optical coherence tomography from an 8-
week-old *nee* mouse. Note differences in the optic nerve head (ONH) region, the peripapillary
retina (PPR) and the retinal ganglion cell complex (RGCC). **(C)** Plot of retinal ganglion cell
complex thickness showing a statistically significant difference in 8-week-old non-mutant (WT
and HET genotypes combined) and *nee* mice; **p < 0.01.

Fig. 3. Correlations between automatedly collected nuclei density and size metrics for BRN3A⁺ and BRN3A⁻ **nuclei.** Average BRN3A⁺ RGC nuclear **(A)** area, **(B)** perimeter, **(C)** Feret

diameter, and **(D)** MinFeret diameter plotted as a function of average density for individual *nee*
and WT retinas. The BRN3A$^+$ RGC nuclei of *nee*$^{mod-sev}$ retinas consistently showed significant
trends that nuclei size characteristics decrease with increasing severity of glaucomatous
damage. Average BRN3A$^-$ nuclear **(E)** area, **(F)** perimeter, **(G)** Feret diameter, and **(H)** MinFeret
diameter plotted as a function of average density for individual *nee* and WT retinas. Unlike
BRN3A$^+$ RGC nuclei, the nuclei of *nee*$^{mod-sev}$ retinas did not robustly show trends that nuclear
size characteristics change during glaucoma. WT *(black circles)*; *nee*$^{mod-sev}$ (*red squares*),
diseased *nee* retinas with BRN3A$^+$ nuclei density < two SD compared to WT; *nee*^{norm} (*red
circles*), healthy *nee* retinas with BRN3A$^+$ nuclei density comparable to WT.

Fig. 4. Population distributions of nuclei size in *nee^mod-sev* retinas. (A) Average area of nuclei with data from all retinas
aggregated by genotype. **(B)** Average area of nuclei with each retina individually plotted. Surviving BRN3A+ RGC nuclei in *nee^mod-sev*
retinas (*red*) were on average larger than WT (*black*). Note that many retinas with relatively modest RGC loss had an average
nuclear size that was larger than WT; as glaucoma progressed the average size decreased. **(C)** Representative distribution curves
for BRN3A+ nuclei, showing a WT retina (*black arrowhead*), a *nee* retina with modest disease (*red arrowhead*), and a *nee* retina with
severe disease (*red asterisk*, in this instance data were combined from both retinas of a *nee* mouse in which both retinas had similar
damage, increasing the number of nuclei analyzed to a number similar to the other examples). The same representative retinas are
similarly indicated in panels B, C, and E. Note that in modest disease, the distribution curve maintained a similar shape compared to
WT, but was shifted to the right, indicating nuclear enlargement broadly occurring in most RGCs regardless of their initial size. In
severe disease, the right-hand tail of the curve was attenuated and there was a corresponding reciprocal peak in the left-hand portion

of the curve, indicating an increased proportion of small RGC nuclei relative to the total population. **(D)** Distribution curves for

BRN3A⁻ nuclei. **(E)** Box and whisker plot for each WT (*grey*) and *nee^mod-sev* (*red*) retina. To highlight differences in the shape of the

distribution, data are normalized to the median value of nucleus size for each retina. Mean (*black dot*), median (*solid line within each*

box), and percentile values (10th, 25th, 75th and 90th; *vertical lines*, *black* for WT and *red* for *nee^mod-sev*) relative to the median. The

average range of the distribution between the 10th and 90th percentiles (*shaded background*) was displaced leftward in *nee^mod-sev*

relative to WT.