Potentially translated sequences determine protein-coding potential of RNAs

in cellular organisms

Yusuke Suenaga[1,‡]*, Mamoru Kato[2,‡], Momoko Nagai[2], Kazuma Nakatani[1,3,4], Hiroyuki

Kogashi[1], Miho Kobatake[1], Takashi Makino[5]

[1]Department of Molecular Carcinogenesis, Chiba Cancer Centre Research Institute,

666-2 Nitona, Chuo-ku, Chiba 260-8717, Japan

[2]Division of Bioinformatics, National Cancer Centre Research Institute, 5-1-1 Tsukiji,

Chuo-ku, Tokyo 104-0045, Japan

[3]Department of Molecular Biology and Oncology, Chiba University School of

Medicine, 666-2 Nitona, Chuo-ku, Chiba 260-8717, Japan

[4]Innovative Medicine CHIBA Doctoral WISE Program

[5]Laboratory of Evolutionary Genomics, Graduate School of Life Sciences, Tohoku

University, Aobayama Campus 6-3, Aramaki Aza-Aoba, Aoba-ku, Sendai 980-8578,

Japan

[‡]These authors equally contributed to this work

*Correspondence to:

Yusuke Suenaga

E-mail: ysuenaga@chiba-cc.jp

Abstract

RNA sequence characteristics determine whether their transcripts are coding or

noncoding. Recent studies have shown that, paradoxical to the definition of noncoding

RNA, several long noncoding RNAs (lncRNAs) translate functional peptides/proteins.

However, the characteristics of RNA sequences that distinguish such newly identified

coding transcripts from lncRNAs remain largely unknown. In this study, we found that

potentially translated sequences in RNAs determine the protein-coding potential of

RNAs in cellular organisms. We defined the potentially translated island (PTI) score as

the fraction of the length of the longest potentially translated region among all regions.

To analyze its relationship with protein-coding potential, we calculated the PTI scores

in 3.4 million RNA transcripts from 100 cellular organisms, including 5 bacteria, 10

archaea, and 85 eukaryotes, as well as 105 positive-sense single-strand RNA virus

genomes. In bacteria and archaea, coding and noncoding transcripts exclusively

presented high and low PTI scores, respectively, whereas those of eukaryotic coding

and noncoding transcripts showed relatively broader distributions. The relationship

between the PTI score and protein-coding potential was sigmoidal in most eukaryotes;

however, it was linear passing through the origin in three distinct eutherian lineages,

including humans. The RNA sequences of virus genomes appeared to adapt to

translation systems of host organisms by maximizing protein-coding potential in host

cells. Hence, the PTIs determined the protein-coding potential of RNAs in cellular

organisms. Additionally, coding and noncoding RNA do not exhibit dichotomous

44 sequence characteristics in eukaryotes, instead they exhibit a gradient of protein-coding

45 potential.

Introduction

Recent advances in sequencing technology have revealed that most of the eukaryotic

genome is transcribed, resulting in production of noncoding RNAs (Okazaki et al. 2002;

Djebali et al. 2012; Ulitsky and Bartel 2013; Kopp et al. 2018). Noncoding RNAs that

are more than 200 nucleotides in length are referred to as long noncoding RNAs

(lncRNAs) and are not translated into proteins (Ulitsky and Bartel 2013; Kopp et al.

2018). LncRNAs have been reported to function in various biological phenomena, such

as the regulation of transcription, modulation of protein or RNA functions, and

organization of the nuclear architecture (Ulitsky and Bartel 2013; Kopp et al. 2018).

However, paradoxical to its definition, a large fraction of lncRNAs are associated with

ribosomes and translated into peptides (Frith et al. 2006; Ingolia et al. 2011; Bazzini et

al. 2014; Ingolia et al. 2014; Ruiz-Orera et al. 2014). Peptides translated from

transcripts annotated as lncRNAs have been shown to have biological functions in

various eukaryotes (Li and Liu 2019). Moreover, known protein-coding genes, such as

TP53, act as functional RNAs (Candeias 2011; Kloc et al. 2011). Detection of these

RNAs with binary functions has blurred the distinction between coding and noncoding

RNAs, and the characteristics of RNA sequences that explain the continuity between

noncoding and coding transcripts remain unclear.

During evolution, new genes originated from pre-existing genes via gene duplication or

from non-genic regions via the generation of new open reading frames (ORFs) (Ohno

1970; Chen et al. 2013; Zhang and Long 2014; McLysaght and Guerzoni 2015;

MyLysaght and Hurst 2016; Holland et al. 2017). Newly evolved genes from non-genic

regions are known as *de novo* genes (Begun et al. 2006; Levine et al. 2006; Begun et al.

2007; Knowles and McLysagtht 2009; Li et al. 2009; Toll-Riera et al. 2009; Li et al.

2010), and regulate phenotypes and diseases (McLysaght and Guerzoni 2015; Chen et

al. 2013; Zhang and Long 2014), such as brain function and carcinogenesis in humans

(Li et al. 2010; Suenaga et al. 2014). LncRNAs serve as sources of newly evolving gene

products (Ruiz-Orera et al. 2014), including *de novo* evolved proteins. Not only ORFs

exposed to natural selection, neutrally evolving ORFs are also translated from lncRNAs

to stably express peptides (Ruiz-Orera et al. 2018), providing a foundation for the

development of new functional peptides/proteins. High levels of lncRNA expression

(Ruiz-Orera et al. 2018), hexamer frequencies of ORFs (Sun et al. 2013; Wang et al.

2013; Ruiz-Orera et al. 2014), and high flexibility of peptides (Wilson et al. 2017) have

been proposed as determinants of coding potential; however, the molecular mechanisms

of how lncRNAs evolve into new coding transcripts remain unclear (Van Oss and

Carvunis 2019).

This study, therefore, sought to identify a new indicator to determine the protein-coding

potential of RNAs. Within an RNA sequence, we termed sequence segments that start

with AUG (the base sequence of the start codon) and end with UAG, UGA, or UAA

(base sequences of stop codons) as potentially translated islands (PTIs). First, we

defined this indicator using PTI lengths and subsequently examined associations

between the indicator and protein-coding potential. We also present analyses of more

than 3.4 million transcripts in 100 cellular organisms belonging to all three domains of

life to investigate the evolution of the relationship between the PTI score and protein-

coding potential. Finally, we examined whether virus RNA genomes have differentially

evolved to maximize the protein-coding potential in different host organisms.

Results

Coding transcripts show higher PTI scores in humans and mice

We defined PTI score as described in the Materials and Methods and illustrated in

Figure 1A to B, and analyzed human transcripts registered in the nucleotide database of

the National Centre for Biotechnology Information for coding (RefSeq ID starting with

NM) and noncoding (RefSeq ID starting with NR) RNA transcripts. Sample PTI score

calculation is presented in Supplementary Figure 1. The nucleic acid sequences and IDs

were downloaded from Table browser (https://genome.ucsc.edu/cgi-bin/hgTables) after

setting the track tab as "RefSeq Genes." A total of 50,052 coding (NM) and 13,550

noncoding (NR) RNAs were registered in 2018 (Supplementary Table 1). To analyze

putative lncRNAs with protein-coding potential, we excluded small RNAs (shorter than

200 bp) or RNAs with a short primary PTI (pPTI; shorter than 20 amino acids) from the

NR transcripts, focusing on the remaining 12,827 transcripts.

We analyzed relative frequencies of NM and NR transcripts, designated as $f(x)$ and $g(x)$

respectively (Figure 1C), where x indicates PTI score. See Materials and Methods for

the definition of $f(x)$ and $g(x)$.

In human transcripts, $g(x)$ showed a distribution that shifted to the left with an apex of

0.15; in contrast, distribution of $f(x)$ shifted to the right with an apex of 0.55 (Figure 1C,

upper panel). As a control, we used randomly generated nucleic acid sequences where

A/T/G/C bases were randomly assigned with equal probabilities. In the controls, the

relative frequencies of PTI scores were shifted to the left in both coding and noncoding

transcripts (Figure 1C, bottom panel). The controls that randomly shuffled the original

sequence without affecting the number of A/T/C/G bases in each transcript also had

relative frequencies of PTI scores shifted to the left in both coding and noncoding

transcripts (Supplementary Figure 2A). Moreover, similar results were obtained using a

data set from the Ensembl database (Supplementary Figure 2B). We also calculated the

PTI scores of mouse transcripts registered in RefSeq and Ensembl and found that the

distribution of $f(x)$ was shifted to the right with an apex of 0.55 (Supplementary Figure

2C), similar to that of human transcripts. These results suggest that the sequences, not

lengths, of the coding transcripts increase PTI scores in mice and humans.

PTI scores correlate with protein-coding potential in humans and mice

Next, we examined the relationship between the PTI score and protein-coding potential

of genes. Protein-coding potential $F(x)$ is defined in the Materials and Methods and a

sample $F(0.15)$ calculation in human transcripts is presented in Figure 1D. This result

indicates that any given human RNA transcript with a calculated PTI score of 0.15 has a

protein-coding potential $F(x)$ of 0.183. $F(x)$ was correlated with PTI score at PTI scores

$\leq$ 0.65 (Figure 1E and Supplementary Figure 3A). The protein-coding potential of

sequences in RefSeq data slightly decreased after peaking at 0.65 (Figure 1E) whereas

that of sequences in the Ensembl data remained high (Supplementary Figure 3A). $F(x)$

of human transcripts was well approximated by the linear regression as follows:

Based on Ensembl data

$F(x) = 1.301x + 0.0072$ ($x \leq 0.65$), $R^2 = 0.984$

Based on RefSeq data

$F(x) = 1.313x + 0.0189$ ($x \leq 0.65$), $R^2 = 0.990$

The intercepts were near zero, and slope was approximately 1.3. By using these

formulas, we can calculate the protein-coding potential $F(x)$ of any given human

transcript with a PTI score $\leq$ 0.65. For example, the protein-coding potential $F(x)$ of

NCYM was calculated as 0.746 or 0.765 based on Ensembl or RefSeq databases,

respectively (Supplementary Figure 1C). In contrast, $F(x)$ in the controls had no positive

correlation with PTI scores (Figure 1E, bottom panel, and Supplementary Figure 3A).

Similar results were obtained for mouse transcripts (Supplementary Figures 3B). $F(x)$ of

mouse transcripts (PTI score $\leq$ 0.65) was approximated as follows.

Based on Ensembl data

$F(x) = 1.142x + 0.067$, $R^2 = 0.982$

Based on RefSeq data

$F(x) = 1.482x - 0.061$, $R^2 = 0.990$

In both human and mouse transcripts, the PTI score correlated linearly with the protein-

coding potential at PTI scores ≤ 0.65. Moreover, when the PTI score limit approached

0, the probability of the transcript being a coding RNA was 0.

PTIs affect the protein-coding potential predicted by Ka/Ks

To examine the relationship between the PTI score and natural selection in the

prediction of protein-coding potential, we calculated the ratio of nonsynonymous (Ka)

to synonymous (Ks) values by comparing human transcripts with syntenic genome

regions of chimpanzee or mouse (Figure 1F). Transcripts were selected based on the

syntenically conserved regions, 44,593 (vs. chimp) and 14,016 (vs. mouse). The results

revealed linear relationships between the $F(x)$ and PTI score in the conserved transcripts

(Figure 1F, left panels). As predicted, coding transcripts contained transcripts with

$Ka/Ks < 0.5$ at a higher frequency than noncoding transcripts, with the largest difference

observed when the PTI score was > 0.9, and the smallest difference at PTI scores of

approximately 0.45 (Figure 1F, right panels).

Therefore, noncoding transcripts showing both negative selection ($Ka/Ks < 0.5$) and the

highest PTI score may include new coding transcript candidates.

We have listed 23 such transcripts (Supplementary Table 2), which include four

transcript variants of a previously identified lncRNA that encodes a tumor suppressive

small peptide, HOXB-AS3 (Huang et al 2017).

Characterization of high PTI score lncRNAs in humans

Next, we investigated whether the PTI score is useful for identifying coding RNAs

among NR transcripts. From the 7,144 transcripts registered as noncoding genes until

2015, we excluded small RNAs (< 200 nucleotides) and those with short primary PTIs

(< 20 amino acids). Among the remaining 6,617 NR genes, 219 were reassigned as NM

over the past 3 years (Supplementary Table 3), including the previously identified *de*

novo gene *MYCNOS/NCYM* (Suenaga et al. 2014). The percentage of reclassification

increased among NR transcripts with high PTI scores (Figure 1G). Thus, a high PTI

score is a useful indicator of coding transcripts. NR transcripts with high protein-coding

potential ($0.6 \leq$ PTI score < 0.8) were then extracted, and the domain structure of the

pPTI amino acid sequence was estimated using BLASTP (See the Materials and

Methods for details). A total of 217 transcripts showed a putative domain structure(s) in

pPTI, whereas 310 transcripts showed none (Supplementary Table 4). Transcripts with a

domain structure are often derived from transcript variants, pseudogenes, or

readthrough of coding genes; transcripts without domain structures are often derived

from antisense RNA or long intergenic noncoding RNA (lincRNA) (Table 1).

We next examined the functions of genes generating NR transcripts with high coding

probabilities (0.6 $\leq$ PTI score < 0.8). We divided the NR transcripts into those with or

without putative domains to investigate candidates of novel coding genes, either

originating from pre-existing genes or created from non-genic regions, respectively.

Analysis using the Database for Annotation, Visualization, and Integrated Discovery

(DAVID) Functional Annotation Tool (Huang et al. 2009a, 2009b) showed that NR

transcripts without domain structures are derived from original genes related to

transcriptional regulation, multicellular organismal process, and developmental process

(Supplementary Table 5; see Materials and Methods for the definition of original

genes). Among the target genes of transcription factors, NMYC, TGIF, and ZIC2 were

ranked in the top three and are all necessary for forebrain development (Supplementary

Table 5) (Brown et al. 1998; Gripp et al. 2000; van Bokhoven et al. 2005). We observed

that NR transcripts with domain structures originated from genes that undergo

alternative splicing which are related to organelle function and are expressed in various

cancers, including respiratory tract tumors, gastrointestinal tumors, retinoblastomas, and

medulloblastomas (Supplementary Table 6).

Relationship between PTI score and relative frequencies of coding/noncoding

transcripts in 100 cellular organisms

To analyze the relationship between PTI scores and protein-coding potential in a broad

lineage of cellular organisms, we selected 100 organisms, consisting of 5 bacteria, 10

archaea and 85 eukaryotes (Supplementary Table 1), and calculated the PTI score for a

total of more than 3.4 million transcripts (Supplementary Table 1). Phylogenic trees of

the cellular organisms are presented on a logarithmic time scale along with the number

of species in each lineage used in the analyses (Figure 2). To examine the evolutionary

conservation of the linear relationship between PTI score and protein-coding potential

in humans and mice, we selected a relatively large number of species (36) from

mammals. The species with fewer than three lncRNAs were not used to calculate $g(x)$

and were not included in the histograms illustrating the relationship with PTI score

(Figure 3). In all organisms, the relative frequency of coding transcripts $f(x)$ was shifted

to the right (higher PTI score) compared to random or random shuffling controls (Figure

3, Figure 4, Supplementary Figure 4 and Supplementary Figure 5).

In bacteria and archaea, $f(x)$ and $g(x)$ exclusively presented high and low PTI scores,

respectively, indicating a clear boundary between the coding transcripts and lncRNAs in

terms of PTI scores (Figure 3 and Supplementary Figure 4). In addition, the highest

frequency of coding transcript $f(x)$ presenting a PTI score was 0.75 in all examined

bacteria (Figure 3) and ≥ 0.75 in archaea (Supplementary Figure 4). Among eukaryotes,

unicellular organisms and non-vertebrates showed the highest frequency of coding

transcripts at 0.65 or 0.75 (Figure 3), while most vertebrates showed highest values $\leq$

0.65 (Figure 3 and 4). In addition, the $f(x)$ distribution in vertebrates was broad and

shifted to the left (lower PTI scores) compared to bacteria and archaea (Figure 3 and 4).

In sharp contrast to $f(x)$, the relative frequency of lncRNAs $g(x)$ was shifted to the right

(higher PTI scores) in eukaryotes, including *G. lamblia* that belongs to the earliest

diverging eukaryotic lineage and lacks mitochondria (Figure 3). Since the distribution

of $f(x)$ in the Excavata, including *G. lamblia,* showed a similar pattern to bacteria, the

right shift of $g(x)$ seems to be an earlier event than the left shift of $f(x)$ in the evolution

of eukaryotes. Collectively, the right and left shifts of $f(x)$ and $g(x)$ contribute to

blurring the boundary between coding and noncoding transcripts in eukaryotes.

Relationship between the PTI score and protein-coding potential

The overlapping of relative frequencies in $f(x)$ and $g(x)$ led us to examine the

relationship between PTI score and protein-coding potential $F(x)$ in eukaryotes. To

avoid misleading data obtained by small sampling numbers, we selected 32 species with

more than 1000 lncRNAs that contain pPTIs for calculation of $F(x)$ (Figure 5 and

Supplementary Figure 6). In humans and mice, the relationship between PTI score and

$F(x)$ was approximated with the linear function passing through the origin of the PTI

score ≤ 0.65. Therefore, we used linear approximation in the $F(x)$ of 32 species and

divided them into two groups, linear and sigmoidal, based on the shape and formula of

the approximated function (Figure 5). We defined the linear group as $R^2 > 0.9$ with an

absolute value of the intercept < 0.1; the sigmoidal group had a slope > 1.0 and intercept

< -0.1. In *U. americanus*, *C. canadensis*, and *G. gorilla*, fewer than five lncRNAs

exhibited PTI scores of 0.05; thus, we eliminated the $F(0.05)$ in these species for the

approximation by linear function (indicated with asterisks in Figure 5). The five species

that did not fit in the linear or sigmoidal group were characterized by high $F(x)$ in low

PTI scores, and belonged to plants (*Z. mays*), reptiles (*A. Carolinensis*) and mammals

(*O. anatinu*, *S. boliviensis*, and *G. gorilla*) (Figure 5). In these species, PTI scores

showed weaker association with protein-coding potential. Sigmoidal relationships were

observed in 18/32 species, while linear relationships were apparent in nine species

within three mammalian lineages, Cetartiodactyla, Rodentia and Primates. Since the

sigmoidal relationship was the most broadly observed across the examined lineages, this

relationship appears to be of an ancestral type.

Characteristics of RNA virus genomes in human and bacterial cells

In sharp contrast to the coding transcripts of bacteria and archaea, the PTI score in

coding transcripts of eukaryotes exhibited overlap with that of noncoding RNA by its

broad distribution of low PTI scores. To investigate the molecular mechanism

underlying the distinct distribution of coding transcripts between bacteria and

eukaryotes, we analyzed genome sequences of RNA viruses that infect human or

bacterial cells. Positive-sense single-strand RNAs, (+) ssRNAs, are parts of virus

genomes that generate mRNAs and are translated into viral proteins via the host

translation system. Therefore, efficient translation in host cells contributes to the

replication of the (+) ssRNA viruses. We speculated that PTIs, other than bona-fide

ORFs, affect the coding potential of the viral genome in host cells. In a viral genome,

multiple bona-fide ORFs exist. We thus extended the concept of PTIs (Figure 6A) to the

multiple ORFs in a viral RNA genome and set the viral ORF (vORF) score (see

Materials and Methods for the definition of the vORF score).

Among the positive-sense ssRNA viruses registered in the NCBI database, 198 were

human viruses and 13 were bacteriophages. We eliminated the viruses that translate

virus proteins by exceptional translation mechanisms such as ribosome frameshifting,

alternative initiation sites, ribosome slippage, and RNA editing, and focused on the

remaining 95 human viruses including 9 retroviruses (Supplementary Table 7) and 10

bacteriophages (Supplementary Table 8). The relative frequency of the human viruses

and bacteriophages showed distinct peaks at PTI scores of 0.65 and 0.75, respectively

(Figure 6B). These values correspond to the PTI scores of the highest protein-coding

potential in humans (Figure 1E and Supplementary Figure 3A) and the highest

frequency of coding transcripts in bacteria (Figure 3), respectively. In addition, the

relative frequency of human viruses showed broader distribution of low PTI scores

compared to bacteriophages, particularly in human retroviruses (Figure 6B). Therefore,

the RNA genomes of viruses appear to have sequence characteristics that maximize

protein-coding potential in host cells.

Discussion

Here, we showed that the PTI is associated with protein-coding potential in cellular

organisms. In bacteria and archaea, the distribution of noncoding and coding transcripts

separately presented at low and high PTI scores, whereas they were merged in

eukaryotes. The overlapping distribution of noncoding and coding RNA in eukaryotes is

caused by the right and left distribution shifts of noncoding and coding transcripts,

respectively.

The right shifts in the distribution of noncoding RNA occurred for *G. lamblia*, one of

the earliest diverging eukaryotes that contain two nuclei and lack mitochondria,

peroxisomes, and a typical Golgi apparatus (Ankarklev et al. 2010; Bartelt et al. 2015;

Buret et al. 2020), and were commonly observed in all examined eukaryotes. The most

prominent difference between eukaryotes and bacteria/archaea is the existence of nuclei

in cells, as this is the definition of eukaryotes. Moreover, functional noncoding RNAs,

by definition, should not be translated by ribosomes in cells. However, in bacteria and

archaea, newly transcribed RNAs are immediately bound by ribosomes (Miller et al.

1970; French et al. 2007) and do not have the chance to escape from translation.

Alternatively, in eukaryotes, the existence of nuclei prevents the immediate binding of

lncRNAs by ribosomes and cytosolic translocation is required for translation. Therefore,

the eukaryotic lncRNAs have a chance to function in the nucleus regardless of PTI

scores or translation efficiencies. This nuclear localization likely promotes

diversification of RNA sequences, resulting in the right shift in the distribution of

noncoding RNAs. Thus, the pervasive transcription of the genome seems to be

advantageous for eukaryotes to create new functional nuclear RNAs while being

disadvantageous for bacteria/archaea by increasing the risk of transcription of a high

PTI score transcript, leading to immediate translation of wasteful and/or toxic proteins

(Monsellier et al. 2007).

The left shifts observed in distribution of coding RNAs were not observed in *G. lamblia*

with its peak at 0.75, however, the majority of fungi, plants and animals, including

humans, exhibited this shift with peaks at PTI scores ≤ 0.65. Thus, the left shift of

coding transcripts likely occurred after the right shift of noncoding transcripts in

evolution of eukaryotes.

Relative frequencies of positive-sense ssRNA viruses exhibit sharp peaks at vORF

scores of 0.75 in bacteriophages and 0.65 in human viruses, indicating the adaptation of

RNA viruses to host cells by maximizing the protein-coding potential of their genomes.

Immediately after viral infection, the viral (+) ssRNA genomes, save for those of

retroviruses, are used as templates for translation in host cytosol. Thus, the distinct

translation systems between humans and bacteria likely affected the left shift in the viral

genome peak, as well as the left shift in coding RNA distribution in eukaryotes. In a

retrovirus, reverse transcriptase produces double-stranded DNA using the viral genome

as a template, which is then inserted into the host genome. The viral genome is

subsequently transcribed within the nucleus and the mRNA is transported to the

cytoplasm where protein products become translated in a manner similar to host

proteins. Therefore, the relatively lower vORF score distribution in human retrovirus

genomes is likely caused by the nuclear localization of the RNA genome, avoiding

immediate binding by ribosomes.

In eukaryotes, we calculated protein-coding potential $F(x)$ based on the overlapped

relative frequencies of noncoding $g(x)$ and coding transcripts $f(x)$. The relationship

between protein-coding potential and PTI score was divided into three groups,

sigmoidal, linear, and others. Among them, switch-type sigmoidal relationships seem to

be of the ancestral-type based on their conservation in eukaryotes, and to emerge after

the all-or-none-type relationships in bacteria and archaea. Meanwhile, the linear group

showed relatively high protein-coding potential at low or intermediate PTI scores,

further blurring the boundary between noncoding and coding transcripts. For example,

the PTI score of human *TP53* (NM_000546.5) is 0.476, and the $F(x)$ calculated by the

linear function is 0.644. The PTI score of *TP53* indicates that the transcript has a high

probability of functioning as both a coding and noncoding RNA, explaining the

bifunction of *TP53* (Candeias 2011; Kloc et al., 2011). Further studies are required to

explain the molecular mechanisms underlying the existence of these three relationships

in eukaryotes.

The PTI score is a value uniquely calculated based on any given RNA sequence,

whereas the Ka/Ks value requires comparison with a species and varies across different

species. Therefore, the Ka/Ks value cannot be uniquely calculated from a given RNA

sequence. Hence, PTI scores, but not Ka/Ks values, exhibit species-specific

relationships with unique distributions of coding potential. Due to the advantages of the

PTI score, we can investigate evolutionary changes in the relationship between PTI

score and coding potential. Furthermore, Ka/Ks < 0.5 did not show different relative

frequencies of noncoding and coding transcripts at a PTI score of approximately 0.45,

suggesting that Ka/Ks does not predict the coding potential at the PTI score and that the

prediction is dependent on the PTI score. In addition, to calculate Ka/Ks, orthologous

transcripts or genomes should be conserved among species. Therefore, Ka/Ks cannot be

applied for coding prediction of species-specific transcripts such as *NCYM*. On the other

hand, by using approximate functions, we can calculate the coding potential $F(x)$ of any

given transcript with a PTI score $\leq$ 0.65 in the nine species classified to linear group.

Human NR transcripts with high PTI scores have been reclassified as coding genes over

the past 3 years, including the human *de novo* gene *NCYM* (Suenaga et al. 2014;

Suenaga et al. 2020). Because *de novo* gene products have no known domain structures,

high PTI score-NR transcripts without putative domains, may be good candidates as

novel *de novo* genes in eukaryotes. *NCYM* is an antisense gene of *MYCN* whose protein

product stabilizes MYCN (Suenaga et al. 2014). MYCN directly stimulates NCYM and

OCT4 transcription, whereas OCT4 induces MYCN (Kaneko et al. 2015). This

functional interplay forms a positive feedback loop for these genes to induce each other

in human neuroblastomas (Suenaga et al. 2014; Islam et al. 2015; Kaneko et al. 2015;

Shoji et al. 2015). Functional annotation of NR genes without putative domains was

related to transcriptional regulation, and the target genes of transcription factors,

including MYCN, TGIF, and ZIC2, were enriched. As *de novo* emergence of *NCYM*

occurred in homininae, NCYM-mediated MYCN activation may modulate human *de*

novo gene births during evolution, regulating the transcription of *MYCN* target genes.

Both TGIF and ZIC2 are causative genes of holoprosencephaly, a disorder caused by a

failure in embryonic forebrain development (Brown et al. 1998; Gripp et al. 2000),

whereas *MYCN* is the causative gene of Feingold syndrome and megalencephaly

syndrome, which are associated with a reduced and increased brain size, respectively

(van Bokhoven et al. 2005; Kato et al. 2019). Thus, the present study also provides a list

of candidate human *de novo* genes possibly involved in brain development and brain-

related diseases.

In conclusion, we identified a novel determinant of protein-coding potential in cellular

organisms. The relationship between PTI score and protein-coding potential revealed

that the boundary between coding and noncoding transcripts in bacteria and archaea is

blurred in eukaryotes. Therefore, when a eukaryotic transcript possesses a moderate PTI

score, bifunctional characterization as both a coding and noncoding transcript may be

essential for a full understanding of the biological roles of RNAs.

Materials and Methods

Potentially translated islands

Definition

PTIs are defined as sequence segments beginning at AUG and ending with any of UAA,

UAG, or UGA in the 5′ to 3′ direction within an RNA sequence in all three possible

reading frames (Figure 1A).

Example

As an example, the PTIs in the human *de novo* gene *NCYM* (Suenaga et al. 2014) are

identified using the cDNA sequence (Supplementary Figure 1A) and are shown as bold

characters (Supplementary Figure 1B). Further information is included in

Supplementary Notes.

The length of PTI and primary/secondary PTIs

Definition

The length of PTI is the length of the amino acid sequence excluding the stop codon and

is represented by l (Figure 1A). In an RNA sequence, the longest PTI is designated as

the primary PTI (pPTI), whereas the others are termed secondary PTIs (sPTIs). The

lengths of a pPTI and sPTI are described as l_{pPTI} and l_{sPTI}, respectively (Figure 1A).

Example

Based on these definitions, the shortest PTI is "AUGUAA," "AUGUAG," or

"AUGUGA," and its amino acid length is 1. For example, the NCYM transcript has a

pPTI with a length of 109 at frame 1; three sPTIs with lengths of 69, 8 and 6,

respectively, at frame 2; and no PTIs at frame 3 (Supplementary Figure 1B and C).

Characteristics

Therefore, the following relationship between the lengths of pPTI and sPTI is held:

422
$$1 \leq l_{\text{sPTI}} \leq l_{\text{pPTI}} \tag{1}$$

424 **PTI score**

425 *Definition*

426 The definition of PTI score is motivated by our hypothetical concept that translation of

427 pPTI is limited by alternate competing sPTIs. We defined the PTI score (Figure 1A)

428 according to Equation 2 and 3:

429
$$\sum_{i=1}^{n} l_{\text{sPTI}i} = l_{\text{sPTI1}} + l_{\text{sPTI2}} + \cdots + l_{\text{sPTI}k} + \cdots + l_{\text{sPTI}n} \tag{2}$$

430
$$\text{PTI score} = \frac{l_{\text{pPTI}}}{l_{\text{pPTI}} + \sum_{i=1}^{n} l_{\text{sPTI}i}} \tag{3}$$

where, $l_{\text{pPTI}} + \sum_{i=1}^{n} l_{\text{sPTI}i}$ represents the sum of all PTI lengths.

Example

If an RNA sequence has only one PTI, the PTI score is 1 (Figure 1B). An RNA

sequence with many sPTIs tends to have a score close to 0 (Figure 1B). If the sum of all

sPTI length is equal to pPTI length, the PTI score is 0.5 (Figure 1B). The PTI score of

the NCYM transcript is calculated as 0.568 (Supplementary Figure 1C). Further

information is included in Supplementary Notes.

Characteristics

Therefore, the range of the PTI score is as follows:

$$0 < \text{PTI score} \leq 1 \tag{4}$$

Relative frequencies *f(x)* and *g(x)*

Definition

We defined the relative frequencies of coding and noncoding transcripts, *f(x)* and *g(x)*,

as follows (Figure 1C):

$$f(x) = \frac{NM(x)}{TNM} \tag{5}$$

$$g(x) = \frac{NR(x)}{TNR} \tag{6}$$

where, TNM and TNR represent the total number of coding and noncoding

transcripts, respectively, excluding the transcripts lacking PTIs. $NM(x)$ or $NR(x)$ is the

number of coding or noncoding transcripts having PTI score $= x$, respectively.

To define coding/non-coding transcripts with a PTI score of x, we made histograms

divided by ten classes, and the median values of the classes were used to represent the

PTI score (Figure 1C). Therefore, in Equation 5 and 6, the PTI score x is restricted as

follows:

$$x = 0.05, 0.15, 0.25, 0.35, 0.45, 0.55, 0.65, 0.75, 0.85, \text{ or } 0.95 \tag{7}$$

Characteristics

Thus, the $f(x)$ and $g(x)$ follow Equation 8 to 11:

$$0 \leq f(x) \leq 1 \qquad (8)$$

$$0 \leq g(x) \leq 1 \qquad (9)$$

$$\sum_x f(x) = 1 \qquad (10)$$

$$\sum_x g(x) = 1 \qquad (11)$$

Protein-coding potential $F(x)$

Definition

The protein-coding potential, $F(x)$, was calculated according to Equation 12:

$$F(x) = \frac{f(x)}{f(x)+g(x)} \qquad (12)$$

Example

As an example, $F(0.15)$ in human transcripts is depicted in Figure 1D. $F(0.15)$ is

calculated using Equation (12):

$$f(0.15) = 0.060$$

$$g(0.15) = 0.268$$

$$F(0.15) = \frac{f(0.15)}{f(0.15)+g(0.15)} = \frac{0.060}{0.060+0.268} = 0.18292 = 0.183$$

Identification of human noncoding transcripts with high protein-coding potential

To identify the noncoding transcripts with high protein-coding potential, NR transcripts

with high $F(x)$ ($0.6 \leq x < 0.8$) were identified from total NR transcripts registered in

the nucleotide database of the National Centre for Biotechnology Information. NR

transcripts shorter than 200 nucleotides or with pPTI encoding putative peptides with

fewer than 20 amino acids were excluded. The amino acid sequences of pPTIs in these

transcripts were subjected to a BLASTP search to examine the presence of putative

domain structures. In the BLASTP search, non-redundant protein sequences (nr) were

applied as the search set, and Quick BLASTP (accelerated protein-protein BLAST) was

used as the algorithm. In the search results, putative conserved domains or the message

"No putative conserved domains have been detected" were shown in the Graphical

Summary tab. CDSEARCH/cdd was the database employed to search conserved

domain structures using the default settings; low complexity filter: no; composition

based adjustment: yes; E-value threshold: 0.01; maximum number of hits: 500. Based

on the results, transcripts with or without putative conserved domain structure were

indicated as + or -, respectively.

Functional annotation of original genes

Original genes are defined as the genes referred in the official gene name of NR

transcripts, including sense genes for antisense transcripts, homologous genes for

pseudogenes, coding genes for noncoding transcript variants, as well as read through,

divergent, or intronic transcripts. For lincRNA, miRNA host gene, small nuclear RNA,

and other lncRNA, the official gene symbol was used for the annotation. This

information was manually checked using information available in the nucleotide

database. The DAVID program (https://www.david.ncifcrf.gov) was used to identify

enriched molecular functions and pathways related to the original genes. Q-values (P-

values adjusted for false discovery rate) were calculated using the Benjamini–Hochberg

method in DAVID.

Ratio of nonsynonymous (Ka) to synonymous (Ks) nucleotide substitution rates

To identify orthologous regions between human transcripts and chimpanzee/mouse

genomes, BLAT v. 36 (Kent 2002) was conducted using human transcript sequences

with the estimated PTI score against chimpanzee (PtRV2) and mouse (GRCm38.p6)

genomic sequences defined in the NCBI database. We defined the blat best-hit genomic

regions of chimpanzee/mouse as orthologs for each human transcript. The human–

chimpanzee (or human–mouse) sequences were aligned for each exon region and the

sequences were combined for each transcript. Only orthologous sequence pairs more

than 60 bp in length (encoding > 20 amino acid sequences) were extracted.

Nonsynonymous (Ka) and synonymous (Ks) nucleotide substitution rates were

estimated by using the method described by Yang and Nielsen (Yang and Nielsen 2000)

as implemented in PAML version 4.8a (Yang 1997). Note that transcripts with high Ka

(> 1) or high Ks (> 1) were excluded from our datasets as outliers. Finally, we estimated

both Ka and Ks for 47,228 NM human–chimpanzee, 14,116 NM human–mouse, 8,810

NR human–chimpanzee, and 1,561 NR human–mouse pairs.

Relative frequency of negatively selected genes

We defined the relative frequencies of negatively selected genes $h(x)$ in coding and

noncoding transcripts (Figure 1F), as shown in Equation 13:

$$h(x) = \frac{Nns(x)}{TNor(x)} \qquad (13)$$

where, $TNor(x)$ represents the total number of coding or noncoding transcripts with

orthologous sequences at PTI score $= x$. $Nns(x)$ is the number of coding or noncoding

transcripts with Ka/Ks < 0.5 at PTI score $= x$. The PTI score x is restricted as shown in

Equation 7.

Phylogenic tree

TimeTree (Hedges et al. 2006) was used to draw the phylogenic tree using the official

names of species.

Selection of viruses and identification of vORF

The complete genomes of positive-sense single-strand RNA viruses of human or

bacteria (Supplementary table 7 and 8) were collected from the NCBI Virus (Hatcher et

al. 2017). Based on the information of described in the Nucleotide database, viral ORFs

were identified and sum of vORF lengths $\sum_{i=1}^{n} l_{vORFi}$ was manually calculated. We

eliminated the viruses that translate virus proteins after splicing or by exceptional

translation mechanisms such as ribosome frameshifting, alternative initiation sites,

ribosome slippage, and RNA editing.

vORF score

Definition

vORF score was calculated according to Equation 14 to 16:

545
$$\sum_{i=1}^{n} l_{vORFi} = l_{vORF1} + l_{vORF2} + \cdots + l_{vORFk} + \cdots + l_{vORFn} \qquad (14)$$

546
$$\sum_{i=1}^{n} l_{sPTIi} = l_{sPTI1} + l_{sPTI2} + \dots + l_{sPTIk} + \dots + l_{sPTIn} \qquad (15)$$

547
$$\text{vORF score} = \frac{\sum_{i=1}^{n} l_{vORFi}}{\sum_{i=1}^{n} l_{vORFi} + \sum_{i=1}^{n} l_{sPTIi}} \qquad (16)$$

where, l_{vORFi} represents the length of bona-fide ORFs and $\sum_{i=1}^{n} l_{sPTIi}$ is the sum of the

lengths of secondary PTI lengths. $\sum_{i=1}^{n} l_{vORFi} + \sum_{i=1}^{n} l_{sPTIi}$ represents the sum of the

length of all PTIs including all ORFs.

Statistical analyses

Statistical analyses were performed using Excel and R software (R Project for Statistical

Computing, Vienna, Austria).

Data availability

1. Source data for statistical analyses and figures (10 example datasets):

https://figshare.com/s/498cb340a075284b2dbf

2. Code associated with generating and analyzing these tables:

https://figshare.com/s/0f1ed0954d5bd620eb59

Funding

This work was partially supported by the Yasuda Medical Foundation and Interstellar

Initiative (grant number 18jm0610006h0001 to Y.S.) from the Japan Agency for

Medical Research and Development.

Acknowledgments

We thank Sana Yokoi, Akira Nakagawara, Yoshitaka Hippo, Yohko Yamaguchi for

helpful comments, Isao Kurosaka and Miayamoto Joe Philips for assistance in

calculating the PTI scores and Mayuko Seto, Ryo Ohtaka and Manami Shinbori for

assistance in checking putative domain structure in BLASTP.

Author contributions

Y.S. conceived and developed the research plan; Y.S., M.K., M.N., K.N., H.K., M.K.,

and T.M. analyzed the data; and Y.S., M.K., and T.M. wrote the manuscript.

Additional information

Supplementary information is available.

Competing interests

The authors declare no competing financial interests.

References

Ankarklev J, Jerlström-Hultqvist J, Ringqvist E, Troell K, Svärd SG. 2010. Behind

the smile: cell biology and disease mechanisms of Giardia species. *Nat Rev*

Microbiol. 8:413–422.

Bailey TL, Grimmond SM. 2006. The abundance of short proteins in the

mammalian proteome. *PLoS Genet*. 2, e52.

Bartelt LA, Sartor RB. 2015. Advances in understanding Giardia: determinants and

mechanisms of chronic sequelae. *F1000Prime Rep*. 7:62.

Bazzini AA, Johnstone TG, Christiano R, Mackowiak SD, Obermayer B, Fleming

ES, Vejnar CE, Lee MT, Rajewsky N, Walther TC, et al. 2014. Identification of

small ORFs in vertebrates using ribosome footprinting and evolutionary

conservation. *EMBO J*. 33:981–993.

Begun DJ, Lindfors HA, Thompson ME, Holloway AK. 2006. Recently evolved

genes identified from Drosophila yakuba and D. erecta accessory gland expressed

sequence tags. *Genetics*. 172:1675–1681.

Begun DJ, Lindfors HA, Kern AD, Jones CD. 2007. Evidence for de novo evolution

of testis-expressed genes in the drosophila yakuba/drosophila erecta clade. *Genetics*.

176:1131–1137.

Brown SA, Warburton D, Brown LY, Yu CY, Roeder ER, Stengel-Rutkowski S,

Hennekam RC, Muenke M. 1998. Holoprosencephaly due to mutations in ZIC2, a

homologue of Drosophila odd-paired. *Nat Genet.* 20:180–183.

Buret AG, Cacciò SM, Favennec L, Svärd S. 2020. Update on Giardia: Highlights

from the seventh International Giardia and Cryptosporidium Conference. Mise à

jour sur Giardia et la giardiase: faits saillants de la Septième Conférence

Internationale sur Giardia et Cryptosporidium. *Parasite* 27:49.

Candeias MM. 2011. The can and can't dos of p53 RNA. *Biochimie* 93:1962–1965.

Chen S, Krinsky BH, Long M. 2013. New genes as drivers of phenotypic evolution.

New genes as drivers of phenotypic evolution. *Nat Rev Genet*. 14:645–660.

Djebali S, Davis CA, Merkel A, Dobin A, Lassermann T, Mortazavi A, Tanzer A,

Lagarde J, Lin W, Schlesinger F, Xue C, et al. 2012. Landscape of transcription in

human cells. *Nature* 489:101–108.

French SL, Santangelo TJ, Beyer AL, Reeve JN. 2007. Transcription and translation

are coupled in Archaea. *Mol Biol Evol*. 24:893–895.

Frith MC, Forrest AR, Nourbakhsh E, Pang KC, Kawai J, Carninci P, Hayashizaki

Y, Ingolia NT, Lareau LF, Weissman JS. 2011. Ribosome profiling of mouse

embryonic stem cells reveals the complexity and dynamics of mammalian

proteomes. *Cell* 147:789–802.

Gripp KW, Wotton D, Edwards MC, Roessler E, Ades L, Meinecke P, Richieri-

Costa A, Zackai EH, Massagué J, Muenke M, et al. 2000. Mutations in TGIF cause

holoprosencephaly and link NODAL signalling to human neural axis determination.

Nat Genet. 25**:**205–208.

Hatcher EL, Zhdanov SA, Bao Y, Blinkova O, Nawrocki EP, Ostapchuck Y,

Schäffer AA, Brister JR. 2017. Virus Variation Resource - improved response to

emergent viral outbreaks. *Nucleic Acids Res*. 45:D482-D490.

Hedges SB, Dudley J, Kumar S. 2006. TimeTree: A public knowledge-base of

divergence times among organisms. *Bioinformatics*. 22: 2971-2972.

Holland PW, Marlétaz F, Maeso I, Dunwell TL, Paps J. 2017. New genes from old:

asymmetric divergence of gene duplicates and the evolution of development. *Philos

Trans R Soc Lond B Biol Sci*. 372:20150480.

Huang DW, Sherman BT, Lempicki RA. 2009a. Systematic and integrative analysis

of large gene lists using DAVID bioinformatics resources. *Nat Protoc.* 4:44–57.

Huang DW, Sherman BT, Lempicki RA. 2009b. Bioinformatics enrichment tools:

paths toward the comprehensive functional analysis of large gene lists. *Nucleic*

Acids Res. 37:1–13.

Huang JZ, Chen M, Chen D, Gao X-C, Zhu S, Huang H, Hu M, Zhu H, Yan G-R.

2017. A peptide encoded by a putative lncRNA HOXB-AS3 suppresses colon cancer

growth. *Mol Cell* 68: 171–184.

Ingolia NT, Lareau LF, Weissman JS. 2011. Ribosome profiling of mouse

embryonic stem cells reveals the complexity and dynamics of mammalian

proteomes. Cell. 147: 789-802.

Ingolia NT. 2014. Ribosome profiling: new views of translation, from single codons

to genome scale. *Nat Rev Genet*. 15:205–213.

Islam SM, Suenaga Y, Takatori A, Ueda Y, Kaneko Y, Kawana H, Itami M, Ohira

M, Yokoi S, Nakagawara A. 2015. Sendai virus-mediated expression of

reprogramming factors promotes plasticity of human neuroblastoma cells.

106:1351–1361.

Kaneko Y, Suenaga Y, Islam SM, Matsumoto D, Nakamura Y, Ohira M, Yokoi S,

Nakagawara A. 2015. Functional interplay between MYCN, NCYM and OCT4

promotes aggressiveness of human neuroblastomas. *Cancer Sci.* 106:840–847.

Kato K, Miya F, Hamada N, Negishi Y, Narumi-Kishimoto Y, Ozawa H, Ito H,

Hori I, Hattori A, Okamoto N, et al. 2019. *MYCN* de novo gain-of-function mutation

in a patient with a novel megalencephaly syndrome. *J Med Genet.* 56:388-395.

Kent WJ. 2002. BLAT--the BLAST-like alignment tool. *Genome Res*. 12:656–664.

Kloc M, Foreman V, Reddy SA. 2011. Binary function of mRNA. *Biochimie*

93:1955–1961.

Knowles DG, McLysaght A. 2009. Recent de novo origin of human protein-coding

genes. *Genome Res*. 19:1752–1759.

Kopp F, Mendell JT. 2018. Functional classification and experimental dissection of

long noncoding RNAs. *Cell*. 172:393–407.

Levine MT, Jones CD, Kern AD, Lindfors HA, Begun DJ. 2006. Novel genes

derived from noncoding DNA in drosophila melanogaster are frequently X-linked

and exhibit testis-biased expression. *Proc Natl Acad Sci U S A*. 103:9935–9939.

Li L, Foster CM, Gan Q, Nettleton D, James MG, Myers AM, Wurtele ES. 2009.

Identification of the novel protein QQS as a component of the starch metabolic

network in Arabidopsis leaves. *Plant J*. 58:485–498.

Li D, Dong Y, Jiang Y, Jiang H, Cai J, Wang W. 2010. A de novo originated gene

depresses budding yeast mating pathway and is repressed by the protein encoded by

its antisense strand. *Cell Res*. 20:408–420.

Li C-Y, Zhang Y, Wang Z, Zhang Y, Cao C, Zhang PW, Lu SJ, Li X, Yu Q, Zheng

X, et al. 2010. A human-specific de novo protein-coding gene associated with

human brain functions. *PLoS Comput Biol*. 6:e1000734.

Li J, Liu C. 2019. Coding or noncoding, the converging concepts of RNAs. *Front*

Genet. 10:496.

McLysaght A, Guerzoni D. 2015. New genes from non-coding sequence: the role of

de novo protein-coding genes in eukaryotic evolutionary innovation. *Philos Trans R*

Soc Lond B Biol Sci. 370:20140332.

McLysaght A, Hurst LD. 2016. Open questions in the study of de novo genes: what,

how and why. *Nat Rev Genet* 17:567–578.

Miller OL Jr, Hamkalo BA, Thomas CA Jr. 1970. Visualization of bacterial genes in

action. *Science* 169:392–395.

Monsellier E, Chiti F. 2007. Prevention of amyloid-like aggregation as a driving

force of protein evolution. *EMBO Rep*. 8:737–742.

Neme R, Tautz D. 2014. Evolution: dynamics of de novo gene emergence. *Curr*

Biol. 24:R238–R240.

Ohno S. 1970. Evolution by gene duplication. New York: Springer-Verlag.

Okazaki Y, Furuno M, Kasukawa T, Adachi J, Bono H, Kondo S, Nikaido I, Osato

N, Saito R, Suzuki H, et al.; FANTOM Consortium; RIKEN Genome Exploration

Research Group Phase I & II Team. 2002. Analysis of the mouse transcriptome

based on functional annotation of 60,770 full-length cDNAs. *Nature*. 420:563–573.

Ruiz-Orera J, Messeguer X, Subirana JA, Alba MM. 2014. Long non-coding RNAs

as a source of new peptides. *Elife* 3:e03523.

Ruiz-Orera J, Verdaguer-Grau P, Villanueva-Cañas JL, Messeguer X, Albà MM.

2018. Translation of neutrally evolving peptides provides a basis for de novo gene

evolution. *Nat Ecol Evol*. 2:890–896.

Shoji W, Suenaga Y, Kaneko Y, Islam SM, Alagu J, Yokoi S, Nio M, Nakagawara

A. 2015. NCYM promotes calpain-mediated Myc-nick production in human

MYCN-amplified neuroblastoma cells. *Biochem Biophys Res Commun.* 461:501–

506.

Suenaga Y, Islam SM, Alagu J, Kaneko Y, Kato M, Tanaka Y, Kawana H, Hossain

S, Matsumoto D, Yamamoto M, et al. 2014. *NCYM*, a Cis-antisense gene of *MYCN*,

encodes a de novo evolved protein that inhibits GSK3β resulting in the stabilization

of MYCN in human neuroblastomas. *PLoS Genet.* 10:e1003996.

Suenaga Y, Nakatani K, Nakagawara A. 2020. De novo evolved gene product

NCYM in the pathogenesis and clinical outcome of human neuroblastomas and

other cancers. *Jpn J Clin Oncol*. 50: 839-846.

Sun L, Luo H, Bu D, Zhao G, Yu K, Zhang C, Liu Y, Chen R, Zhao Y. 2013.

Utilizing sequence intrinsic composition to classify protein-coding and long non-

coding transcripts. *Nucleic Acids Res*. 41:e166.

Toll-Riera M, Bosch N, Bellora N, Castelo R, Armengol L, Estivill X, Albà MM.

2009 Origin of primate orphan genes: a comparative genomics approach. *Mol Biol*

Evol. 26:603–612.

Ulitsky I, Bartel DP. 2013. lincRNAs: genomics, evolution, and mechanisms. *Cell*.

154:26–46.

van Bokhoven H, Celli J, van Reeuwijk J, Rinne T, Glaudemans B, van Beusekom

E, Rieu P, Newbury-Ecob RA, Chiang C, Brunner HG. 2005. MYCN

haploinsufficiency is associated with reduced brain size and intestinal atresias in

Feingold syndrome. *Nat Genet.* 37:465–467.

Van Oss SB, Carvunis AR. 2019. De novo gene birth. *PLoS Genet.* 15:e1008160.

Wang L, Park HJ, Dasari S, Wang S, Kocher JP, Li W. 2013. CPAT: Coding-

Potential Assessment Tool using an alignment-free logistic regression model.

Nucleic Acids Res. 41:e74.

Wilson BA, Foy SG, Neme R, Masel J. 2017. Young genes are highly disordered as

predicted by the preadaptation hypothesis of de novo gene birth. *Nat Ecol Evol*.

1:0146.

Yang Z, Nielsen R. 2000. Estimating synonymous and nonsynonymous substitution

rates under realistic evolutionary models. *Mol Biol Evol.* 17:32–43.

Yang Z. 1997. PAML: a program package for phylogenetic analysis by maximum

likelihood. Comput Appl Biosci. 13:555–556

Zhang YE, Long M. 2014. New genes contribute to genetic and phenotypic

novelties in human evolution. *Curr Opin Genet Dev*. 29:90–96.

Table 1. Number of original transcripts that produced NR transcripts with high coding

frequency (0.6 $\leq$ PTI score < 0.8)

Transcript	Domain		Total	P-value
	With	Without		
Antisense	4	61	65	7.79E-08
LincRNA	3	65	68	7.60E-09
Pseudogene	50	17	67	4.32E-07
Readthrough	7	0	7	6.00E-03
Transcript variant of coding gene	146	35	181	1.05E-19
Divergent	0	2	2	N.S.
Intronic	0	6	6	N.S.
Small nuclear RNA	0	3	3	N.S.
miRNA host gene	0	3	3	N.S.
Other lncRNA	7	118	125	1.12E-13
Total	217	310	527	

P-values were calculated by Yate's continuity correction. N.S., not significant.

Figure legends

Figure 1. Potentially translated island (PTI) score predicts protein-coding potential

of human transcripts. (A) Conceptual explanation of PTIs in an RNA in the three

frames and the definition of PTI score. Black and white rectangles indicate primary and

secondary PTIs, respectively. The primary PTI is the longest PTI, while secondary PTIs

are all others. l is the length of PTIs. (B) Visual image of the PTIs in the RNAs at low,

medium (0.5), and high (1) PTI scores. (C) Histogram of relative frequency of PTI

scores in coding f(x) and noncoding g(x) transcripts (upper) and in random controls

(bottom). (D) Visual explanation of F(x) at the PTI score of 0.15. (E) The PTI score

correlates with protein-coding potential, $F(x)$, at PTI scores $\leq$ 0.65 (upper) and those

in random controls (bottom). (F) Relationship between PTI score and $F(x)$ in syntenic

human transcript to chimpanzee (upper left) or mouse (bottom left). The relative

frequency of transcripts with negative selection $h(x)$ were plotted for each PTI score

(upper and bottom right). See the Methods for the definition of $h(x)$. The transcripts are

syntenic to the genome of chimpanzee (upper right) or mouse (bottom right).The open

circles indicate NR transcripts, and the closed circles indicate NM transcripts. (G)

Relationship between PTI scores and percentage of NR transcripts re-registered as NM

in the past 3 years. N.D., not detected.

Figure 2. Phylogenetic tree for the cellular organisms. The numbers of species are

indicated in each lineage. The lineage of five species including one archaea

(*Nitrososphaera viennensis* EN76), two fungi (*Puccinia graminis f. sp. Tritici* and

Pyricularia oryzae), and two animals (*Strongylocentrotus purpuratus* and *Lingula*

anatine) are unknown, and thus, excluded from the figure.

Figure 3. **Relationship between PTI score and relative frequencies of coding and**

noncoding transcripts from bacteria to mammals. Histogram of $f(x)$ (white) or $g(x)$

(black) in observed data (left) and in nucleic-acid–scrambled controls (right). PTI scores

with highest $f(x)$ are presented in the histograms.

Figure 4. Relationship between PTI score and relative frequencies of coding f(x)

and noncoding transcripts in Primates (A), Glires (B), and Laurasiatheria (C).

Figure 5. Relationship between PTI score and protein-coding potential F(x) in 32

eukaryotes. Phylogenetic tree of the 32 species (left), the dot plots and shape and

formula of approximate functions. L, S, and O indicate linear (in red), sigmoidal (in

black) and other (in blue) functions. Less than five lncRNAs had a PTI score of 0.05 in

U. americanus, *C. canadensis*, and *G. gorilla*. Therefore, we eliminated the $F(0.05)$ in

these species for linear function approximations (asterisks).

Figure 6. Adaptation of (+) ssRNA viruses to host cells by maximizing protein-

coding potential of the RNA genomes. (A) Conceptual explanation of the length of

sPTIs and bona-fide viral ORFs in a genome of (+) ssRNA virus and the definition of

viral ORF (vORF) score. Black and white rectangles indicate viral ORFs and secondary

PTIs, respectively. l is the length of the ORFs and PTIs. (B) Histograms of relative

frequencies of human (+) ssRNA viruses (red) and bacteriophages (black).

Supplementary figure legends

Supplementary Figure 1. PTIs of NCYM and an example of PTI score calculation.

(A) cDNA sequence of *NCYM* transcript. (B) Translated amino acid sequence of *NCYM*

in 3 frames in 5′ to 3′ (sense) direction. Red characters indicate a primary PTI and blue

characters indicate secondary PTIs. Stop codons are shown as asterisks. (C) Calculation

of PTI score and *F(x)* in the *NCYM* transcripts. The length of pPTI is 109, and the sum

of sPTI lengths is 83. Therefore, the PTI score is 0.568.

Supplementary Figure 2. Relative frequency of coding and noncoding transcripts

in human and mouse PTI scores. (A) Histogram of PTI score relative frequency in

coding *f(x)* and noncoding *g(x)* human transcripts with random shuffling controls using

human data sets from RefSeq. (B) Relative frequency of coding *f(x)* and noncoding *g(x)*

transcripts calculated using human data sets from Ensembl. (B) Relative frequency of

coding *f(x)* and noncoding *g(x)* transcripts calculated using mouse data sets from

RefSeq (upper panels) or Ensembl (lower panels).

Supplementary Figure 3. PTI score correlates with protein-coding potential, *F(x)*,

at PTI scores $\leq$ 0.65 in human and mouse transcripts. (A) Relationship between

PTI score and $F(x)$ in a human data set from Ensemble and random controls (center).

Random shuffling controls (right) were generated from a human data set from both

Ensemble and RefSeq. (B) Relationship between PTI score and $F(x)$ in mouse

transcripts using data sets from RefSeq (upper panels) or Ensembl (lower panels).

Supplementary Figure 4. Relationship between PTI score and relative frequencies

of coding and noncoding transcripts in archaea. Phylogenetic tree for 9 archaeal

species and histogram of $f(x)$ (white) or $g(x)$ (black) in the data (left) and in nucleic-

acid–scrambled controls (right). PTI scores with highest $f(x)$ are indicated in the

histograms. The lineage of one archaea (*Nitrososphaera viennensis* EN76) is unknown

and thus excluded from the phylogenic tree.

Supplementary Figure 5. Relationship between PTI score and relative frequencies

of coding and noncoding transcripts in plants. Phylogenetic tree for 12 plants and

histogram of $f(x)$ (white) or $g(x)$ (black) in the data (left) and in nucleic-acid–scrambled

controls (right). PTI scores with highest $f(x)$ are indicated.

Supplementary Figure 6. Relationship between PTI score and protein-coding

potential $F(x)$ in 32 eukaryotes. Relationship between PTI score and $F(x)$ in data sets

from Ensemble (left, used in Figure 5) and those in random controls (right). Mouse and

human data are identical to Supplementary Figure 3. Shape of approximate functions

are shown as L, S, or O, indicating linear (in red), sigmoidal (in black), and other (in

blue) functions, respectively. The number of lncRNAs at PTI score 0.05 were < 5 in *U.*

americanus, *C. canadensis*, and *G. gorilla*. Therefore, we eliminated the *F*(0.05) in

these species for the approximation by linear functions (asterisks).

Supplementary Table 1. Cellular organisms with the official name, taxonomy ID,

lineage information, and number of coding or noncoding transcripts.

Supplementary Table 2. Twenty-three human noncoding transcripts showing both

negative selection (*K*a/*K*s < 0.5) and the highest PTI score.

Supplementary Table 3. Human NR transcripts reassigned as NM over the past

three years.

Supplementary Table 4. Human NR transcripts with high protein-coding potential

(0.6 ≤ PTI score < 0.8).

Supplementary Table 5. Funtional annotation of human NR transcripts with high

protein-coding potential and without putative domain structure(s).

Supplementary Table 6. Functional annotation of human NR transcripts with high

protein-coding potential and with putative domain structure(s).

Supplementary Table 7. Positive-sense single-stranded human viruses with official

name, taxonomy ID, lineage information, as well as genome length and sequence.

Supplementary Table 8. Positive-sense single-stranded bacteriophages with official

name, taxonomy ID, lineage information, source information, and genome length

and sequence.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6