Phosphorus stress induces the synthesis of novel glycolipids in *Pseudomonas*

***aeruginosa* that confer protection against a last-resort antibiotic**

Running title: ecophysiology of lipid renovation in *Pseudomonas aeruginosa*

Rebekah A. Jones[1,2], Holly Shropshire[2], Caimeng Zhao[3], Andrew Murphy[2], Ian Lidbury[2,4],

Tao Wei[3], David J Scanlan[2], Yin Chen[2]

[1]MRC Doctoral Training Partnership, University of Warwick, Coventry, CV4 7AL, UK.

[2]School of Life Sciences, University of Warwick, Coventry, CV7, 7AL, UK

[3]School of Food and Biological Engineering, Zhengzhou University of Light Industry,

Zhengzhou, 450002, China

[4]Department of Animal and Plant Sciences, University of Sheffield, Sheffield, S10 1AE,

UK

Correspondence to Yin Chen, (email, y.chen.25@warwick.ac.uk, phone 00 44 24

76528976)

Classification: *Integrated genomics and post-genomics approaches in microbial ecology*

Abstract

Pseudomonas aeruginosa is a nosocomial pathogen with a prevalence in immunocompromised individuals and is particularly abundant in the lung microbiome of cystic fibrosis patients. A clinically important adaptation for bacterial pathogens during infection is their ability to survive and proliferate under phosphorus limited growth conditions. Here, we demonstrate that *P. aeruginosa* adapts to P-limitation by substituting membrane glycerophospholipids with sugar-containing glycolipids through a lipid renovation pathway involving a phospholipase and two glycosyltransferases. Combining bacterial genetics and multi-omics (proteomics, lipidomics and metatranscriptomic analyses), we show that the surrogate glycolipids monoglucosyldiacylglycerol and glucuronic acid-diacylglycerol are synthesised through the action of a new phospholipase (PA3219) and two glycosyltransferases (PA3218 and PA0842). Comparative genomic analyses revealed that this pathway is strictly conserved in all *P. aeruginosa* strains isolated from a range of clinical and environmental settings and actively expressed in the metatranscriptome of cystic fibrosis patients. Importantly, this phospholipid-to-glycolipid transition comes with significant ecophysiological consequence in terms of antibiotic sensitivity. Mutants defective in glycolipid synthesis survive poorly when challenged with polymyxin B, a last-resort antibiotic for treating multi-drug resistant *P. aeruginosa*. Thus, we demonstrate an intriguing link between adaptation to environmental stress (nutrient availability) and antibiotic resistance, mediated through membrane lipid renovation that is an important new facet in our understanding of the ecophysiology of this bacterium in the lung microbiome of cystic fibrosis patients.

Introduction

P. aeruginosa is a significant nosocomial pathogen in intensive care units causing pneumonia, surgical wound site infections and sepsis (1-2). It is now recognised as a leading cause of morbidity and mortality in chronically infected cystic fibrosis (CF) patients and immunocompromised individuals due to the surge of carbapenem resistant strains, a key group of first line antibiotics for treating *P. aeruginosa* infections (3). For these drug-resistant *P. aeruginosa* strains, a viable but not ideal treatment option are polymyxins, considered to be last resort antibiotics. Although polymyxins are active against *P. aeruginosa*, their use was originally discontinued due to concerns over toxicity (4). Indeed, *P. aeruginosa* has started to develop mechanisms of resistance to polymyxins due to an increase in their use globally. These primarily include modifications to the lipopolysaccharide (LPS) layer of the outer membrane through the addition of 4-amino-4-deoxy-L-arabinose (L-Ara4N) or phosphoethanolamine (pEtN) (5-6). These changes perturb the electrostatic interaction between cationic polymyxins and the normally negatively charged LPS.

Glycerophospholipids, such as phosphatidylglycerol (PG) and phosphatidylethanolamine (PE), are the major lipids forming the membrane lipid bilayer in bacteria, archaea and eukaryotes (7-11). They play a fundamental role in the evolution of the cell and it is widely accepted that the last universal common ancestor possessed a phospholipid membrane (12-13). Although it is uncertain why evolution selected glycerophospholipids as the building blocks for maintaining cellular membranes (13), it is known that organisms can alter their membrane lipid composition in response to nutrient stress or environmental changes (7, 14). Previous studies have firmly established the link between nutrient stress, particularly phosphorus availability, and the expression of a variety of virulence factors in *P. aeruginosa* (15-19). However, it is unclear whether adaptation to phosphorus limitation

in this bacterium causes a change in membrane lipid composition and, if so, whether lipid

remodelling comes with unforeseen ecophysiological consequences. Using a synthesis of

multi-omics approaches, here we show that *P. aeruginosa* produces surrogate glycolipids

to replace phospholipids in response to phosphorus limitation. These glycoglycerolipids

have not previously been reported in *P. aeruginosa*. This lipid renovation pathway is

strictly conserved in all *P. aeruginosa* strains isolated from a range of clinical settings and

actively expressed in the metatranscriptome of cystic fibrosis patients. Importantly, such

a phospholipid-to-glycolipid transition comes with a significant consequence in antibiotic

sensitivity, in that glycolipids confer protection when challenged with the antimicrobial

peptide polymyxin B. As such, glycolipid-mediated resistance to polymyxin B represents

a new resistance mechanism that is quite different from the previously documented

modification of LPS (5-6). This work highlights how the physiological adaptation of

Pseudomonas aeruginosa to phosphorus limitation can mediate a physiological response

that may have profound implications for the survival of the bacteria in the lung microbiome.

Materials and methods

Cultivation of *P. aeruginosa* and mutants

P. aeruginosa strain PAO1 was obtained from the DSMZ culture collection (Germany) and

routinely cultured in lysogeny broth (LB). A defined medium previously outlined for

Pseudomonas species to control phosphate levels was also used (20). This modified

minimal media A comprised: Na-succinate 20 mM, NaCl 200 mg L^{-1}, NH$_4$Cl 450 mg L^{-1},

CaCl$_2$ 200 mg L^{-1}, KCl 200 mg L^{-1}, MgCl$_2$ 450 mg L^{-1}, with trace metals FeCl$_2$ 10 mg L^{-1}

and MnCl$_2$ 10 mg L^{-1}, with 10 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid

(HEPES) buffer used at pH 7. Na$_2$HPO$_4$ was then added to a final concentration of 50 µM

(low P) or 1 mM (high P). An intermediate phosphate source of 400 µM Na$_2$HPO$_4$ was

used for overnight cultures in some experiments to prevent any excess storage of

phosphate that could hamper results. All components were filter sterilised using 0.22 μm

pore-size filters, and made up using double deionised H_2O. Mutants were obtained from

the *P. aeruginosa* strain PAO1 transposon mutant library at the University of Washington,

and confirmed using PCR and subsequent sequencing.

Alkaline phosphatase assay

Alkaline phosphatase activity was monitored as a measure of Pi stress. Liquid *P.*

aeruginosa culture samples were incubated with 10 mM *para*-nitrophenol phosphate

(*p*NPP) to a final concentration of 1 mM. Yellow-*p*NP supernatant was measured in

triplicate at 407 nm (BioRad iMark microplate reader). Readings were normalised using

both a Tris-only incubation control and further by bacterial density (optical density reading

at 600 nm (OD_{600})).

Over-expression of Agt1 and Agt2 in *E. coli*

P. aeruginosa genes PA3218 (*agt1*) and PA0842 (*agt2*) were codon optimised for *E. coli*

and chemically synthesised (GenScript) into plasmid pET-28a(+). *E. coli* BLR(DE3)

competent cells were thawed for 5 minutes before incubation with 10 ng pET-28a_Agt

plasmid, and placed on ice for 5 minutes. Cells were then subjected to heat shock at 42°C

for 30 seconds, placed back on ice for 2 minutes. Recovery SOC media was added, with

samples incubated at 37°C shaking, for 1 hour. Transformed cells were then plated onto

kanamycin-LB agar and grown overnight at 37°C. To harvest cells for lipid extraction,

single colonies were picked to grow in small volume LB-Kan to 0.6 OD_{600} before induction

with 0.4 mM IPTG overnight at 25°C. 1 mL samples were then pelleted at 10,000 x *g* for

5 minutes. Pellets were stored at -80°C until lipid extraction and subsequent analysis on

HPLC-MS.

To purify the Agt1 and Agt2 proteins from recombinant *E. coli*, IPTG was added to

a final concentration of 0.5 mM once the cultures reached an OD_{600} of 0.6. After a further

12 h of growth at 30°C, cells were harvested by centrifugation and resuspended in buffer

A containing 50 mM Tris-HCl, pH 7.9, 50 mM NaCl. Cells were disrupted by sonication

and 1% (w/v) triton X-100 was then added and the cells were then incubated for 2.5 hr at

4°C. The cells were then centrifuged at 12,000 × *g* for 20 min, and the soluble fraction

was loaded onto a nickel column (GE Healthcare, USA) pre-equilibrated with buffer A. The

recombinant Agt1 and Agt2 enzymes were eluted with an elution buffer (20 mM Tris-HCl,

pH 7.9, 500 mM NaCl, 300 mM imidazole) and dialyzed overnight into buffer A to remove

imidazole. For further purification, the samples were dialyzed overnight into buffer B

containing 50 mM Tris-HCl, pH 7.9, 200 mM NaCl, concentrated by ultrafiltration using a

30-kDa membrane (Millipore), and loaded onto a Superdex 200 (16/60) gel filtration

column (GE Healthcare, USA), which was pre-equilibrated with buffer B (50 mM Tris-HCl,

pH 7.9, 200 mM NaCl). The fraction size was 0.5 ml, and the flowrate was 0.5 ml/min.

Purified protein was analysed by SDS-PAGE, and protein concentrations were determined

using the Bradford assay.

Membrane lipid extraction and HPLC-MS analysis

Intact polar membrane lipids were extracted using a modified version of the typically used

Folch extraction method (21, 22). Liquid *P. aeruginosa* cultures growing in high and low

phosphate modified minimal medium A were sampled after 8 hrs, collecting the equivalent

of 0.5 OD_{600} into a 2 mL glass chromacol vial (Thermo Scientific), pelleted at 4°C, 4,000

rpm for 15 minutes. For lipid extraction, a ratio of 500:300:1000 µL of

methanol:water:chloroform (all LC-MS grade) was used. The lipid fraction was collected

from the lower phase using a glass Pasteur pipette. This chloroform extract was then dried

under a stream of nitrogen (Techne sample concentrator) and resuspended in 1 mL 95%

(v/v) acetonitrile (HPLC grade): 5% (w/v) ammonium acetate (10 mM, pH 9.2) for analysis.

Extracted lipid samples were analysed using an UltiMate 3000 HPLC (Thermo Scientific)

system coupled to an AmazonSL quadrupole ion trap (Bruker) mass spectrometer (MS),

using electrospray ionisation. Hydrophilic interaction chromatography (HILIC) using a

BEH amide XP column (Waters) was utilised to separate lipid classes based on their head

group (35). The column chamber was maintained at 30°C and the samples passed

through at a 150 µL min^{-1} flow rate. The mobile phase of acetonitrile:ammonium acetate

(pH 9.2) was used to elute the sample in a 15 minute per sample gradient, from 95% to

28% ammonium acetate. The lipid d17:1/12:0 sphingosylphosphoethanolamine (Sigma-

Aldrich, 50 nM) was added to the samples and used as an internal standard. Tandem MS

(or MS^n) was used to fragment the intact lipids for identification. The data were analysed

using the Bruker Compass software package (DataAnalysis and QuantAnalysis).

Enzyme activity assays

The glycosyltransferase activity of Agt1 and Agt2 was measured using uridine

diphosphate (UDP)-glucose or UDP-glucuronic acid and 0.1 mM C16:0/C18:1

diacylglycerol (DAG) as the substrate. 2.0 µM purified enzyme was used in 10 mM

Tricine/KOH buffer, pH 8.5 with 2 mM dithiothreitol. The resulting mixture (500 µl) was

incubated at 30°C for 60 min with constant shaking at 200 rpm. The lipid products were

extracted using the Folch method as described above. The lipid extracts were further

analysed by LC-MS for the identification of MGDG/GADG though MS^n fragmentation and

for the quantification of DAG against standards. The K_m and V_{max} values were calculated

using Michaelis-Menten plots with various concentrations of UDP-sugars (0.1 to 1.0 mM)

in three replicates.

Antibiotic sensitivity assays

P. aeruginosa cultures were grown to an OD_{600} of 0.6 in high or low phosphate minimal

media A (see above). Cultures were then diluted 1:100 in prewarmed minimal media A

containing 4 µg mL^{-1} polymyxin B sulfate (Sigma). Samples were incubated at 37°C, 180

rpm, and assayed for survivors at specified time points by serial dilution plating onto LB.

E. coli cultures containing pET-28a-Agt1 or pET-28a-Agt2 were grown to an OD_{600} of 0.6

in LB broth, and the expression of Agt1 and Agt2 was induced by incubation with 0.4 mM

IPTG overnight at 25°C. A negative control of *E. coli* containing the pET-28a vector only

was also set up. Overnight cultures were diluted 1:100 in prewarmed LB broth containing

0.4 mM IPTG and 20 µg mL^{-1} polymyxin B sulfate (Sigma). Samples were incubated at

37°C, 180 rpm, and assayed for survivors at specified time points by serial dilution plating

onto LB agar + kanamycin 25 µg mL^{-1}.

Comparative proteomic analysis

P. aeruginosa PAO1 WT (1 mM phosphate, 50 µM phosphate) and PlcP mutant (50 µM

phosphate) cell pellets in three biological replicates were resuspended in LDS (lithium

dodecyl sulfate) sample buffer containing 1% β-mercaptoethanol before lysing at 95°C

and vortexing. 30 µL of each sample were run on NuPAGE 10% Bis-Tris protein gel

(Invitrogen) for a short time before staining with SafeStain (Thermo Fisher) and excising

the whole protein band. In-gel proteins were de-stained using 50% (v/v) ethanol, 50 mM

ammonium bicarbonate (ABC), before being reduced and alkylated for 5 min at 70°C using

10 mM TCEP (tris(2-carboxyethyl)phosphine) and 40 mM CAA (2-chloroacetamide),

respectively. After washing with 50% (v/v) ethanol 50 mM ABC, peptides were lysed

overnight using trypsin. Finally, peptides were extracted by sonication in a water bath (10

min at room temperature), concentrated using a Speed-Vac (50 mins) and resuspended

in 2.5% acetonitrile 0.05% formic acid. Extracted peptides were analysed by nanoLC-ESI-

MS/MS using the Ultimate 3000/Orbitrap Fusion instrumentation (Thermo Scientific). The

UniProt proteome for *P. aeruginosa* strain PAO1 was used for peptide analysis. Further

data analysis was carried out using MaxQuant and Perseus software as described

previously; peptides without triplicate measures were filtered out (23). Comparative

proteomics data are presented in Supplementary Tables S1 and S3.

Phylogenomics and metatranscriptomics analyses

The protein sequences of PA3219, PA3218 and PA0842 were used to search genome

sequences of *Pseudomonas* clades in the JGI IMG genome portal

(https://img.jgi.doe.gov/). Note that the PA3218 protein is incorrectly annotated in the

genome of PAO1. The putative glycosyltransferase located immediately downstream of

PA3219 was manually inspected by aligning to the corresponding gene (PA14_22600) in

the genome of *P. aeruginosa* PA14. To identify PA3218 in misannotated *P. aeruginosa*

genomes, the nucleotide sequence immediately downstream of PA3219 was aligned with

agt1 in strain PA14, (locus tag PA14_22600). The phylogeny of *Pseudomonas* clades was

determined using the nucleotide sequences of six housekeeping genes (*rpoB, rpoD,*

dnaE, recA, atpD, gyrB) retrieved from each genome using IQ-Tree with the parameters -

m TEST -bb 1000 -alrt 1000. The most suitable model was chosen by the software.

Evolutionary distances were inferred using maximum-likelihood analysis. Relationships

were visualised using the online platform the Interactive Tree of Life viewer

(https://itol.embl.de/). The conserved Pho box sequence was predicted using the MEME

server (24).

The metatranscriptomics datasets of sputum samples obtained from a CF patient

7-days (SRX5145606) and 8-days (SRX5145605) before death (25), CF patient G

(SRR6833349) from Denmark (26) and a patient (SRR6833340) with chronic wound

infection (26) were retrieved from the short reads archive (SRA) database

(https://www.ncbi.nlm.nih.gov/sra). The reads were downloaded using fastq-dump and

mapped using the BBMap aligner as described previously (27). Briefly, the SRA reads

were mapped to the genome sequence of *P. aeruginosa* PAO1 using a stringent cut-off

of minid=0.97. Relative abundance data were compared using RPKM (reads per kilobase

of transcript, per million mapped reads). The list of RPKM abundance of individual genes

of *P. aeruginosa* PAO1 is shown in **Suppl. Table S4**.

Results and discussion

P. aeruginosa **produces novel glycolipids in response to Pi stress**

To determine changes in the membrane lipidome in response to P-stress, the model *P.*

aeruginosa strain PAO1 was grown in minimal medium under high (1 mM) or low Pi (50

µM) conditions (**Figure 1a**). The latter condition elicited strong alkaline phosphatase

activity, measured through the liberation of *para*-nitrophenol (*p*NP) from *para*-nitrophenol

phosphate (*p*NPP) (**Figure 1b**), this being a strong indication that cells were P-stressed.

Analysis of membrane lipid profiles using high performance liquid chromatography

coupled to mass spectrometry (HPLC-MS) revealed the presence of several new lipids

under Pi stress conditions (**Figure 1c**). Thus, during Pi-replete growth (1 mM phosphate),

the lipidome is dominated by two glycerophospholipids: phosphatidylglycerol (PG, eluted

at 6.8 min) and phosphatidylethanolamine (PE, eluted at 12.2 min). During Pi-stress a lipid

species with mass to charge ratio (*m/z*) of 623 and 649 were also found, with MS

fragmentation resulting in a 131 *m/z* peak, a diagnostic ion for the amino-acid containing

ornithine lipid. This is consistent with previous reports of ornithine lipids in the *P.*

aeruginosa membrane in response to Pi stress (28-29).

Further to ornithine lipids, three unknown lipids eluting at 7.7 min, 8.7 min and 9.8

min, were only present under Pi stress conditions (**Figure 1c**). Using several rounds of

MS fragmentation (MS^n), with a quadrupole ion trap MS, fragmentation patterns

characteristic of glycolipids were found for all three peaks. For each peak of interest, the most predominant lipid masses of 774.7 *m/z*, 786.8 *m/z* and 788.6 *m/z* were analysed by MS^n in positive ionisation mode (**Figure 1d**). In each case, an initial head group was lost leaving a significant signal of 595.6 *m/z*, the mass of the glycolipid building block diacylglycerol (DAG). Further fragmentation leads to the loss of either fatty acyl chain from DAG, leaving monoacylglycerols of 313.2 *m/z* and 339.3 *m/z*. Two monoacylglycerols with different masses are produced as a result of the original lipid containing 16:0 and 18:1 fatty acids (313.2 *m/z* and 339.3 *m/z* monoacylglycerols, respectively). To further elucidate the identity of the peaks, a search for a neutral loss of a polar head group was carried out. Thus, the intact masses of 774.7 *m/z* and 788.6 *m/z* in positive ionisation mode leads to the loss of a head group of -179 and -193 *m/z*, which corresponds to a hexose- and a glucuronate- group, respectively (**Figure 1d**), suggesting the occurrence of novel monoglucosyldiacylglycerol (MGDG) and glucuronic acid diacylglycerol (GADG) glycolipids in *P. aeruginosa*. The third glycolipid peak at 8.7 min remains an unknown lipid with intact mass of 786.8 *m/z* (hereafter designated as a putative unknown glycolipid, UGL). Together, these data confirm the production of new glycolipids in *P. aeruginosa* in response to Pi stress.

Comparative proteomics uncover the lipid renovation pathway in *P. aeruginosa*

To determine the proteomic response of *P. aeruginosa* to phosphorus limitation, and identify the genes involved in glycolipid formation, strain PAO1 was cultivated under high and low Pi conditions for 8 hours and the cellular proteome then analysed. A total of 2844 proteins were detected, 175 of which were found to be differentially regulated by Pi availability (**Figure 2b, Suppl. Table S1**). In line with previous transcriptomic studies of strain PAO1 (18), major phosphorus acquisition mechanisms were highly expressed

under Pi stress conditions, *e.g.* the Pi-specific transporter PstSCAB, the two-component

regulator PhoBR (**Suppl. Table S1**) (30).

Comparative proteomics also identified several genes which are likely important

for membrane lipid remodelling (**Figure 2a**) including PA3219 (4.6-fold increase under Pi-

depleted conditions, FDR<0.01), encoding a putative phospholipase C protein, and

PA0842 (4-fold increase under Pi-depleted conditions, FDR<0.01), encoding a putative

glycosyltransferase (**Figure 2b**). PA3219 has 47% protein sequence identity to PlcP from

Phaeobacter sp. MED193 and 46% identity to PlcP from *Sinorhizobium meliloti* (25-27).

In these bacteria, PlcP is essential in the lipid remodelling pathway for the formation of the

diacylglycerol (DAG) backbone, representing the essential intermediate for the production

of glycolipids (31-32). In *P. aeruginosa* PAO1, PA3219 appears to form an operon with

PA3218, a putative glycosyltransferase likely under the control of the PhoBR two

component system, as a highly conserved Pho box sequence was recognisable in the

promoter region (**Figure 2c**). PA3218 (hereafter referred to as Agt1) has 41% protein

sequence identity to the Agt of *Phaeobacter* sp. MED193. PA0842 showed 35% identity

to the Agt of *Phaeobacter* sp. MED193 and a Pho box sequence is also found in its

promoter region. This corroborates the finding that the PA0842 protein (hereafter referred

to as Agt2) was significantly upregulated under Pi-depleted conditions (**Figure 2b**). In

summary, comparative proteomic analysis suggests that *P. aeruginosa* PAO1 adopts this

PlcP-Agt lipid remodelling pathway for the production of glycolipids in response to Pi-

stress (**Figure 2a**).

The PlcP-Agt mediated lipid renovation pathway is strictly conserved in *P.*

***aeruginosa* and actively transcribed in the metatranscriptomes of cystic fibrosis**

patients

To uncover how widespread this predicted PlcP-Agt lipid remodelling pathway is amongst

the genus *Pseudomonas,* including *P. aeruginosa* strains, we conducted a thorough

comparative genomics analysis of these lipid renovating loci. PlcP-Agt is strictly conserved

in all 770 genome-sequenced *P. aeruginosa* strains in the IMG/M database, including all

three-previously recognised *P. aeruginosa* lineages (33-34), group 1 represented by strain

PAO1, group 2 represented by strain PA14 and group 3 represented by strain PA7 (**Figure**

3, Suppl. Table S2). Indeed, this remodelling pathway is prevalent in many *Pseudomonas*

groups, including the plant pathogen *Pseudomonas syringae*. To investigate whether the

PlcP-Agt lipid remodelling pathway is involved in host-pathogen interactions, we analysed

metatranscriptomic datasets from cystic fibrosis patients, where *P. aeruginosa* is known

to be prevalent in the fatal exacerbation period before patient death (25). To the best of

our knowledge, only two studies have reported the metatranscriptome of the bacterial

community present in CF sputum (25, 26). Indeed, *phoBR* and *pstS* are amongst the most

highly expressed genes, confirming previous observations that *P. aeruginosa* is Pi-limited

during human airway epithelia infection (36, 37). Interestingly, the alkaline phosphatase

phoA (38) was highly expressed in sputum from CF patients but not from wound samples

which was also dominated by *P. aeruginosa*. Importantly, the transcripts of *P. aeruginosa*

agt1/plcP/agt2 are highly expressed in CF sputum during the fatal exacerbation period

before death (**Figure 2d**). Therefore, our phylogenomic and metatranscriptomic analyses

suggest that not only is the PlcP-Agt lipid remodelling pathway strictly conserved and

prevalent in *P. aeruginosa*, but also the corresponding genes are also highly expressed

during CF patient infection, suggesting a potential role for lipid renovation in host-

pathogen interactions.

Experimental validation of the lipid renovation pathway for glycolipid formation in

P. aeruginosa

To validate the function of these two putative glycosyltransferases (Agt1, Agt2) in the

formation of glycolipids, we synthesized the codon-optimized genes (PA3218 and

PA0842, respectively) for recombinant expression in *Escherichia coli*. The total lipidomes

from the recombinant *E. coli* strains were then analysed by HPLC-MS to determine the

presence of glycolipids in a gain-of-function assay. Expression of *P. aeruginosa* Agt1

(PA3218) was sufficient for the production of MGDG (eluted at 7.7 min) in *E. coli*,

confirmed through MS^n fragmentation (**Figure 4a**). No UGL nor GADG was observed in

the lipidome of this Agt1-overexpressing *E. coli* strain. Expressing Agt2 (PA0842) from *P.*

aeruginosa in *E. coli* was sufficient for the accumulation of the GADG glycolipid (eluted at

9.8 min), also confirmed through the MS^n fragmentation pattern (**Figure 4b**). Equally, no

UGL nor MGDG was observed in the Agt2-overexpressing *E. coli* strain. Production of

these glycolipids was not observed in the same strain of *E. coli* transformed with an empty

vector control (pET28a). It is therefore likely that UGL production is carried out by another

glycosyltransferase, the identity of which remains to be discovered.

To confirm the role of Agt1 and Agt2 in the production of MGDG and GADG, we

purified Agt1 and Agt2 from recombinant *E. coli* (**Figure 4c**) and carried out enzyme

assays using UDP-glucose and UDP-glucuronic acid as the sugar donor and DAG as the

acceptor. Agt1 can only accept UDP-glucose as the substrate with an affinity of K_m=298.1

± 9.5 μM (**Figure 4c**, middle panel) and produced MGDG as expected (**Figure 4d**, left

panel). Similarly, Agt2 can use UDP-glucuronic acid as the substrate (K_m= 373.0 ± 12.9

μM (**Figure 4c**, right panel), producing the GADG lipid (**Figure 4d**, right panel).

Interestingly, the purified Agt2 enzyme can also use UDP-glucose to some extent with a

K_m of 480 μM (data not shown) although the corresponding lipid MGDG was not observed

in the lipid extract from the lipidome of the recombinant host *E. coli* (**Figure 4b**).

To further confirm the role of these genes in *P. aeruginosa* glycolipid biosynthesis

we analysed the lipidomes of mutants in *ΔplcP*, *Δagt1* and *Δagt2* in strain PAO1 (**Figure**

5a, b). Differences were analysed by searching for the intact masses of the glycolipids

MGDG and GADG: 774.6 *m/z* and 788.6 *m/z* in positive ionisation mode with an

ammonium adduct, respectively. As expected, under Pi stress MGDG is no longer

produced in the *Δagt1* mutant and similarly GADG is no longer produced in the *Δagt2*

mutant (**Figure 5a**). In the *ΔplcP* mutant, no MGDG was found and the GADG lipid was

significantly reduced but not entirely abolished (**Figure 5b**). The small amount of GADG

produced in the *ΔplcP* mutant suggests that an alternative supply of DAG (independent of

the degradation of phospholipids by PlcP) is available in this mutant. Nevertheless,

lipidome analyses of the *ΔplcP*, *Δagt1* and *Δagt2* mutants strongly supports the key role

of this PlcP-Agt pathway (**Figure 2A**) in lipid renovation in *P. aeruginosa*.

The protective role of glycolipids to antibiotic resistance in *Pseudomonas*

aeruginosa

To assess whether the growth of the glycolipid-deficient mutants (*ΔplcP*, *Δagt1*, *Δagt2*)

was affected by Pi stress, we grew the mutants in the defined minimal medium under high

and low Pi conditions. However, no significant difference in growth rates was found (Suppl.

Figure S1). The presence of glycolipids in the membrane may, however, have a profound

impact on the functioning of the membrane during Pi stress. For example, PG is an anionic

lipid with net negative charges whereas MGDG has a neutral charged sugar group.

Although a PG-to-GADG substitution may not necessarily change membrane charge (35),

it may affect membrane curvature and the packing density of lipids. Thus, subsequent

knock-on effects in membrane function might be expected (10). We therefore set out to

investigate whether membrane lipid composition may have an impact on antibiotic

resistance in *P. aeruginosa*. As cationic antimicrobial peptides directly interact with

bacterial cell membranes, we focused on the impact of lipid remodelling on the killing

activity of polymyxin B. We conducted the analyses under P-depleted conditions, since

Pi-stress is clinically important, already known to induce the expression of virulence

factors (15, 17, 18, 29), and our own analysis confirmed that an array of genes involved

in phosphate acquisition and lipid remodelling in *P. aeruginosa* are indeed highly

expressed in sputum samples from lung microbiome of CF patients (**Figure 2d**).

Polymyxins represent the drug-of-last resort for effectively treating carbapenem-resistant

P. aeruginosa infections (3, 39).

To test the sensitivity of the mutants in the PlcP-Agt pathway to polymyxin B, we

compared WT and mutants using kill curve analyses as the typically used disk diffusion

method does not work efficiently for cationic antimicrobials (40). Indeed, there was a

significant decrease in the survival of all three PAO1 glycolipid synthesis mutants (*ΔplcP*,

Δagt1 and *Δagt2*) compared to the wild type when challenged with polymyxin B,

suggesting a protective role of glycolipids in polymyxin B resistance (**Figure 5c**). Such a

protective role of glycolipids in polymyxin B resistance was not observed for other

antibiotics, including ciprofloxacin, gentamicin, ceftazidime and meropenem (data not

shown). *P. aeruginosa* is known to enhance its resistance to polymyxins through

decoration of its lipopolysaccharide (LPS) layer using either 4-amino-4-deoxy-L-arabinose

(L-Ara4N) by *arnB* (5), or the addition of phosphoethanolamine (pEtN) by *eptA* (6). It is

thought that these changes perturb the electrostatic interaction between the cationic

polymyxin B and the normally negatively charged LPS. To investigate whether these

mechanisms play a role in the glycolipid-deficient mutants, we conducted a comparative

proteomics analysis of the *ΔplcP* mutant and WT under Pi depleted conditions, which

revealed only a small number of differentially expressed proteins (**Suppl. Table S3**). The

majority of these differentially expressed proteins are uncharacterised. However,

importantly, LPS modification enzymes previously found to confer antimicrobial peptide

resistance, such as ArnB and EptA, were not differentially expressed between the WT and

ΔplcP mutant. Therefore, our data suggests that it is the glycolipids that are the major

contributor to increased polymyxin B resistance, which constitutes a new biological

mechanism for polymyxin resistance. To this end, we tested the resistance to polymyxin

B of recombinant *E. coli* strains overexpressing *P. aeruginosa* Agt1 and Agt2, that produce

MGDG and GADG, respectively (**Figure 4a, 4b**). Indeed, in this gain-of-function assay,

both Agt1 and Agt2 overexpressing *E. coli* strains had enhanced resistance to polymyxin

B compared to the empty vector control (**Figure 5d**), supporting the protective role of

these glycolipids to antimicrobial peptides.

To conclude, we present here the discovery of novel glycolipids produced in *P.*

aeruginosa during adaption to phosphorus stress. This lipid renovation pathway is strictly

conserved in all *P. aeruginosa* isolates to date and highly expressed in the

metatranscriptome of CF patients, suggesting a key role of lipid remodelling in the

ecophysiology of this bacterium. Interestingly, lipid remodelling as a response to survive

phosphorus stress in turn comes with trade-offs in terms of antibiotic resistance; these

glycolipids may protect the bacterium from insult by cationic antimicrobial peptides,

highlighting a new resistance mechanism to polymyxin B which has been previously

overlooked. It remains to be seen whether the altered susceptibility to polymyxin B is the

sole trade-off following lipid remodelling of phospholipids to glycolipids. After all, evolution

appears to have selected phospholipids as the dominant lipids in the last universal

common ancestor (12).

Acknowledgements

This work was funded by an MRC Doctoral Training Partnership studentship in

Interdisciplinary Biomedical Research (MR/J003964/1) awarded to RAJ and by a Royal

Society International Exchanges 2017 Cost Share (China) award (IEC\NSFC\170213; grant

agreement no. 170213). AM and YC are supported by a European Research Council (ERC)

award under the European Union's Horizon 2020 research and innovation programme

(grant agreement no. 726116). We also thank the Proteomics Research Technology

Platform, University of Warwick, UK for their contribution.

Competing interest

The authors declare no competing interests.

References

1. Murray, T. S., Egan, M., & Kazmierczak, B. I. (2007). *Pseudomonas aeruginosa* chronic colonization in cystic fibrosis patients. *Curr. Opin. Pediatr.,* 19: 83-88.

2. Gaynes, R., Edwards, J. R., & System, N. N. I. S. (2005). Overview of nosocomial infections caused by Gram-negative bacilli. *Clin. Infect. Dis.*, 41:848–854.

3. Hawkey PM, Livermore DM (2012). Carbapenem antibiotics for serious infections. *BMJ*. 344:e3236.

4. Landman, D., Georgescu, C., Martin, D. A., & Quale, J. (2008). Polymyxins revisited. *Clin. Microbiol. Rev*. 21:449–465.

5. Chung, E. S., Lee, J. Y., Rhee, J. Y., & Ko, K. S. (2017). Colistin resistance in *Pseudomonas aeruginosa* that is not linked to *arnB. J. Med. Microbiol.*, 66:833–841.

6. Nowicki, E. M., O'Brien, J. P., Brodbelt, J. S., & Trent, M. S. (2015). Extracellular zinc induces phosphoethanolamine addition to *Pseudomonas aeruginosa* lipid A via the ColRS two-component system. *Mol. Microbiol. 97*:166–178.

7. Parsons JB, Rock CO. (2013) Bacterial lipids: metabolism and membrane homeostasis. *Prog. Lipid Res.* 52:249-276.

8. Zhang, Y.-M., & Rock, C. O. (2008). Membrane lipid homeostasis in bacteria. *Nat. Rev. Microbiol.* 6:222–233.

9. van Meer G, Voelker DR, Feigenson GW. (2008) Membrane lipids: where they are and how they behave. *Nat. Rev. Mol. Cell Biol.* 9:112-124.

10. Harayama T, Riezman H. (2018) Understanding the diversity of membrane lipid composition. *Nat. Rev. Mol. Cell Biol.* 19:281-296.

11. May, K.L., & Silhavy T.J. (2017) Making a membrane on the other side of the wall. *Biochim. et Biophy. Acta*. 1862:1386-1393.

12. Lombard J, López-García P, Moreira D. (2012) The early evolution of lipid membranes and the three domains of life. *Nat. Rev. Microbiol.* 10:507-515.

13. Peretó J, López-García P, Moreira D. (2004) Ancestral lipid biosynthesis and early membrane evolution. *Trends Biochem. Sci*. 29:469-477.

14. Sahonero-Canavesi DX, López-Lara IM, Geiger O (2019) Membrane lipid degradation and lipid cycles in microbes. In *Aerobic Utilization of Hydrocarbons, Oils, and Lipids*, 10.1007/978-3-319-50418-6_38

15. Lamarche MG, Wanner BL, Crépin S, Harel J. (2008) The phosphate regulon and bacterial virulence: a regulatory network connecting phosphate homeostasis and pathogenesis. *FEMS Microbiol. Rev.* 32:461-473.

16. Long, J., Zaborina, O., Holbrook, C., Zaborin, A., & Alverdy, J. (2008). Depletion of
intestinal phosphate after operative injury activates the virulence of *P. aeruginosa*
causing lethal gut-derived sepsis. *Surgery*, *144*:189–197.

17. Francis VI, Stevenson EC, Porter SL. (2017) Two-component systems required for
virulence in *Pseudomonas aeruginosa. FEMS Microbiol. Lett.* 364(11). doi:
10.1093/femsle/fnx104.

18. Bains, M., Fernández, L., & Hancock, R. E. W. (2012). Phosphate starvation
promotes swarming motility and cytotoxicity of *Pseudomonas aeruginosa*. *Appl.*
Environ. Microbiol., *78*:6762–6768.

19. Son, M. S., Matthews, W. J., Kang, Y., Nguyen, D. T., & Hoang, T. T. (2007). *In*
vivo evidence of *Pseudomonas aeruginosa* nutrient acquisition and pathogenesis
in the lungs of cystic fibrosis patients. *Infect. Immun.* 75:5313–5324.

20. Lidbury, I. D. E. A., Murphy, A. R. J., Scanlan, D. J., Bending, G. D., Jones, A. M.
E., Moore, J. D., et al., (2016). Comparative genomic, proteomic and exoproteomic
analyses of three *Pseudomonas* strains reveals novel insights into the phosphorus
scavenging capabilities of soil bacteria. *Environ. Microbiol.* 18:3535–3549.

21. Sebastián, M., Smith, A. F., González, J. M., Fredricks, H. F., Van Mooy, B.,
Koblížek, M., et al., (2016). Lipid remodelling is a widespread strategy in marine
heterotrophic bacteria upon phosphorus deficiency. *ISME J.* 10:968–978.

22. Smith AF, Rihtman B, Stirrup R, Silvano E, Mausz MA, Scanlan DJ, Chen Y. (2019)
Elucidation of glutamine lipid biosynthesis in marine bacteria reveals its importance
under phosphorus deplete growth in *Rhodobacteraceae. ISME J.* 13:39-49.

23. Shropshire H, Jones RA, Aguilo-Ferretjans MM, Scanlan DJ, Chen Y. (2021)
Proteomics insights into the *Burkholderia cenocepacia* phosphorus stress
response. doi.org/10.1111/1462-2920.15451.

24. Bailey TL, Elkan C (1994) Fitting a mixture model by expectation maximization to
discover motifs in biopolymers. *Proceedings of the Second International*
Conference on Intelligent Systems for Molecular Biology, 28-36, AAAI Press,
Menlo Park, Californias.

25. Cobián Güemes AG, Lim YW, Quinn RA, Conrad DJ, Benler S, Maughan H,
Edwards R, et al. (2019) Cystic fibrosis rapid response: translating multi-omics data
into clinically relevant information. *mBio.* 16;10(2):e00431-19.

26. Cornforth DM, Dees JL, Ibberson CB, Huse HK, Mathiesen IH, Kirketerp-Moller K,
et al., (2018) *Pseudomonas aeruginosa* trascriptome during human infection. *Proc.*
Nat. Acad. Sci. USA 115:E5125-5134.

27. Jones, H.J., Krober, E., Stephenson, J., Mausz M.A., Jameson, E., Millard, A., Purdy, K.J., Chen, Y. (2019) A new family of uncultivated bacteria involved in methanogenesis from the ubiquitous osmolyte glycine betaine in coastal saltmarsh sediments. *Microbiome*. 7, 120. doi.org/10.1186/s40168-019-0732-4

28. Kim, S. K., Park, S. J., Li, X. H., Choi, Y. S., Im, D. S., & Lee, J. H. (2018). Bacterial ornithine lipid, a surrogate membrane lipid under phosphate-limiting conditions, plays important roles in bacterial persistence and interaction with host. *Environ. Microbiol. 20*:3992–4008.

29. Lewenza, S., Falsafi, R., Bains, M., Rohs, P., Stupak, J., Sprott, G. D., & Hancock, R. E. W. (2011). The *olsA* gene mediates the synthesis of an ornithine lipid in *Pseudomonas aeruginosa* during growth under phosphate-limiting conditions, but is not involved in antimicrobial peptide susceptibility. *FEMS Microbiol. Lett.* 320: 95–102.

30. Wilton M, Halverson TW, Charron-Mazenod L, Parkins MD, Lewnza S. (2018) Secreted phosphatase and deoxyribonuclease are required by *Pseudomonas aeruginosa* to defend against neutrophil extracellular traps. *Infect. Immun* 86:e00403-18.

31. Wei, T., Quareshy, M., Zhang, Y. Z., Scanlan, D. J., & Chen, Y. (2018). Manganese is essential for PlcP metallophosphoesterase activity involved in lipid remodeling in abundant marine heterotrophic bacteria. *Appl. Environ. Microbiol. 84,* e01109-18.

32. Zavaleta-Pastor, M., Sohlenkamp, C., Gao, J.-L., Guan, Z., Zaheer, R., Finan, T. M., et al., (2010). *Sinorhizobium meliloti* phospholipase C required for lipid remodeling during phosphorus limitation. *Proc. Nat. Acad. Sci. USA107*:302–307.

33. Freschi L, Jeukens J, Kukavica-Ibrulj I, Boyle B, Dupont MJ, Laroche J, et al., (2015) Clinical utilization of genomics data produced by the international *Pseudomonas aeruginosa* consortium. *Front Microbiol.* 29;6:1036. doi: 10.3389/fmicb.2015.01036.

34. Ozer EA, Nnah E, Didelot X, Whitaker RJ, Hauser AR. (2019) The population structure of *Pseudomonas aeruginosa* is characterized by genetic isolation of *exoU+* and *exoS+* lineages. *Genome Bio.l Evol.* 11:1780-1796.

35. Diercks H, Semeniuk A, Gisch N, Moll H, Duda KA, Hölzl G. (2015) Accumulation of novel glycolipids and ornithine lipids in *Mesorhizobium loti* under phosphate deprivation. *J. Bacteriol.* 197:497-509.

36. Chugani, SA and Greenberg, EP (2007) The influence of human respiratory

epithelia on *Pseudomonas aeruginosa* gene expression. *Microb. Pathog.* 42: 29–
35.

37. Frisk, A, Schurr, JR, Wang, G, Bertucci, DC, Marrero, L, Hwang, SH, et al. (2004)
Transcriptome analysis of *Pseudomonas aeruginosa* after interaction with human
airway epithelial cells. *Infect. Immun.* 72: 5433–5438.

38. Filloux A, Bally M, Soscia C, Murgier M, Lazdunski A (1988) Phsphate regulation
in *Pseudomonas aeruginosa*: Clonning of the alkaline phosphatase gene and
identification of *phoB* and *phoR*-like gnes. *Mol. Gen. Genet* . 212(3):510-513.

39. Poirel L, Jayol A, Nordmann P. (2017) Polymyxins: antibacterial activity,
susceptibility testing, and resistance mechanisms encoded by plasmids or
chromosomes. *Clin. Microbiol. Rev*. 30:557-596.

40. Ezadi, F., Ardebili A., Mirnead R (2019) Antimicrobial susceptibility testing for
polymyxins: challenges, issues, and recommendations. *J. Clin. Microbiol. 57,*
e01390-18.

41. Monds RD, Newell PD, Schwartzman JA, O'Toole GA. (2006) Conservation of the
Pho regulon in *Pseudomonas fluorescens* Pf0-1. *Appl. Environ. Microbiol*.
72:1910-1924.

42. Semeniuk A, Sohlenkamp C, Duda K, Hölzl G. (2014) A bifunctional
glycosyltransferase from *Agrobacterium tumefaciens* synthesizes monoglucosyl
and glucuronosyl diacylglycerol under phosphate deprivation. *J. Biol. Chem.*
289(14):10104-10114.

43. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von
Haeseler A, et al. (2020) IQ-TREE 2: New models and efficient methods for
phylogenetic inference in the genomic era. *Mol. Biol. Evol*. 37:1530–1534.

 Figure legends

Figure 1 Lipidomics analysis uncovers novel glycolipid formation in *Pseudomonas*

***aeruginosa* strain PAO1 in response to phosphorus limitation.**

a) Growth of strain PAO1 WT in minimal medium A containing 1 mM phosphate (+Pi, blue)

or 50 µM phosphate (-Pi, black) over 12 hours. Data are the average of 3 independent

replicates.

b) Liberation of *para*-nitrophenol (*p*NP) from *para*-nitrophenol phosphate (*p*NPP) through

alkaline phosphatase activity, under Pi-repleted (1 mM, black) and Pi-depleted (50 µM,

yellow) conditions. Error bars represent the standard deviation of three independent

replicates.

c) Representative chromatograms in negative ionisation mode of the *P. aeruginosa*

lipidome when grown under phosphorus stress (-Pi, black) compared to growth under

phosphorus sufficient conditions (+Pi, orange). PG, phosphatidylglycerol; PE,

phosphatidylethanolamine, OL, ornithine lipids. Lower panel: extracted ion

chromatograms of three new glycolipid species in *P. aeruginosa* which are only produced

during Pi-limitation (black, 1 mM; orange, 50 µM). MGDG: monoglucosyldiacylglycerol,

GADG: glucuronic acid-diacylglycerol and UGL: unconfirmed glycolipid.

d) Mass spectrometry fragmentation spectra of three glycolipid species present under Pi

stress in *P. aeruginosa*, at retention times of 7.7 (*m/z* 774.7), 8.7 (*m/z* 786.7) and 9.8 (*m/z*

788.6) minutes, respectively. Each spectrum depicts an intact lipid mass with an

ammonium (NH_4^+) adduct exhibiting neutral loss of a head group, yielding diacylglycerol

(DAG) (595 *m/z*). Further fragmentation yields monoacylglycerols (MAG) with C16:0 or

C18:1 fatty acyl chains.

pathway responsible for glycolipid formation in *Pseudomonas aeruginosa* strain

PAO1.

a) The proposed pathway for lipid remodelling through the PlcP-Agt pathway. PlcP

degrades membrane phospholipids such as PG, to generate diacylglycerol (DAG)

intermediates for the formation of MGDG and GADG through the activity of

glycosyltransferases, using either UDP-glucose or UDP-glucuronate as the co-substrate

(42).

b) Volcano plot depicting differentially expressed proteins when comparing Pi-repleted

and Pi-depleted conditions. Significantly upregulated proteins when under Pi stress are

shown in red (left), and those that are significantly upregulated when Pi is sufficient are in

green (right). Significance was accepted when the false discovery rate (FDR) was < 0.05,

and a fold change ≥ 2.

c) Genomic organisation of predicted lipid remodelling genes in *P. aeruginosa*.

Glycosyltransferases (orange) PA3218 (Agt1) and PA0842 (Agt2) are predicted to be

involved in glycolipid synthesis. PA3219 is predicted to be PlcP in *P. aeruginosa*.

Predicted Pho box sequences in the promoter regions (represented in blue boxes) of each

glycosyltransferase operon from *P. aeruginosa* strains representing the PAO1 clade, the

PA7 clade and the PA14 clade are shown. The black dots represent residues which are

conserved in the Pho box consensus CTGTCATNNNNCTGTCAT (41).

d) Metatranscriptomic analysis of PlcP-Agt lipid remodelling genes in sputum samples

from a cystic fibrosis patient 7-days (CF_D-7) and 8-days (CF_D-8) before death (25) and

a Danish CF patient (CF_Person G) (26) as well as a wound sample from a burns patient

from the USA (Burn patient) (26). Relative abundance is expressed as RPKM (reads per

kilobase of transcript, per million mapped reads). PhoA (PA3296) encodes an alkaline

phosphatase (38).

Figure 3 Occurrence of *plcP-agt* genes in major *Pseudomonas* groups. The

phylogeny of *Pseudomonas* clades was determined using the nucleotide sequences of six

housekeeping genes (*rpoB, rpoD, dnaE, recA, atpD, gyrB*) retrieved from each genome

using IQ-Tree (43). The filled colour indicates the presence of the genes in the genomes

whereas a blank indicates the absence of the corresponding gene in the genomes. The

two-component system PhoBR (black circles) is found in all genomes and the PlcP-

Agt1/Agt2 are strictly conserved in all 770 genome-sequenced *P. aeruginosa* strains that

form three clades represented by strain PA14, PA01 and PA7 respectively. Bootstrap

values >75% are shown.

Figure 4, **Characterization of glycolipid formation from recombinant Agt1 and Agt2.**

a) Extracted ion chromatogram of the MGDG lipid from recombinant *E. coli* expressing

Agt1. An empty vector control is also shown (red line). The identity of MGDG is further

validated using mass spectrometry fragmentation showing the neutral loss of 179

corresponding to the loss of glucose and the formation of monoacylglycerols (MAG) with

C16:0 or C18:1 (*m/z* 313.2, 339.3).

b) Extracted ion chromatogram of the GADG lipid from recombinant *E. coli* expressing

Agt2. An empty vector control is also shown (red line). The identity of GADG is further

validated using mass spectrometry fragmentation showing the neutral loss of 193

corresponding to the loss of glucose and the formation of monoacylglycerols (MAG) with

C16:0 or C18:1 (*m/z* 313.2, 339.2).

c) Purified Agt1 and Agt2 protein from recombinant *E. coli* (left panel) and Michaelis

Menten kinetics of Agt1 towards UDP-glucose (middle panel) and Agt2 towards UDP-

glucuronic acid (right panel) as substrate, respectively.

d) Mass spectrometry identification of MGDG and GADG produced from purified Agt1 and

Agt2 using DAG and UDP-glucose and UDP-glucuronic acid as the substrate,

respectively.

Figure 5 Glycolipid formation in *P. aeruginosa* and mutants under Pi stress,

showing a protective role of glycolipids to polymyxin B.

a-b) Relative abundance of the glycolipid MGDG (**a**) and GADG (**b**) in *P. aeruginosa*

mutants *plcP* (purple trace), *agt1* (blue trace) and *agt2* (green, trace) compared to the wild

type (WT). Cells were cultivated under low Pi conditions (50 µM) and a representative

extracted ion chromatogram of MGDG/GADG is shown between the WT, (black trace) and

each mutant. The right most panel shows the abundance of MGDG or GADG calculated

relative to an internal lipid standard d17:1/12:0 sphingosylphosphoethanolamine (Sigma-

Aldrich) in the wild-type and mutant strains of *P. aeruginosa*. Values are calculated from

three biological replicates and the error bars denote standard deviation. MGDG,

monoglucosyldiacylglycerol, GADG, glucuronic acid-diacylglycerol.

c) Survival of glycolipid remodelling mutants Δ*plcP* (purple), Δ*agt1* (green) and Δ*agt2*

(blue) when challenged with 4 µg mL^{-1} polymyxin B compared to WT under Pi stress

(black). All experiments were conducted under Pi stress conditions and the results are the

average of three biological replicates; error bars denote standard deviation.

d) Survival of glycolipid producing *Escherichia coli* when challenged with 20 µg mL^{-1}

polymyxin B. All experiments were conducted in three replicates and error bars denote

667 standard deviation. Black, *E. coli* containing the empty vector pET28a; green, *E. coli*

668 containing plasmid pET28a-Agt1; blue, *E. coli* containing plasmid pET28a-Agt2.

Supplementary Figures and Tables

Suppl. Figure S1 Growth of the *P. aeruginosa* mutants in the minimal medium under Pi repleted and Pi depleted conditions.

Suppl. Table S1 Proteomic analysis of differentially expressed proteins in the wild-type *P. aeruginosa* PAO1 in response to different Pi levels (1 mM versus 50 μM).

Suppl. Table S2 Protein BLAST identification of locus tags homologous to *agt1* and *agt2* glycolipid synthesis genes in all genome sequenced *P. aeruginosa* strains at the JGI IMG database.

Suppl. Table S3 Proteomic analysis of wild type *P. aeruginosa* PAO1 versus the Δ*plcP* mutant grown at 50 μM Pi.

Suppl. Table S4 Whole genome mapping to *Pseudomonas aeruginosa* PAO1 of metatranscriptomic datasets (SRX5145605, SRX5145606, SRR6833349) from sputum samples taken from cystic fibrosis patients and a patient with chronic wound infection (SRR6833340). **S4A** (SRX5145605)**, S4B** (SRX5145606), **S4C** (SRR6833349), **S4D** (SRR6833340)**.** RPKM, reads per kilobase of transcript per million mapped reads; FPKM, fragments per kilobase of transcript per million mapped reads.

(a) MGDG in *P. aeruginosa* PAO1 and mutants

(b) GADG in *P. aeruginosa* PAO1 and mutants

(c)

(d)