1

Resolving Between Novelty and Homology in the Rapidly Evolving Phallus of Drosophila

5 Running Title: **Defining homology in a rapidly evolving tissue**

10 Gavin R. Rice[1], Jean R. David[2], Nicolas Gompel[3], Amir Yassin[2,4], Mark Rebeiz[1*]

[1] Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA, USA
[2] Laboratoire Evolution, Génomes, Comportement, Ecologie (EGCE), UMR 9191, CNRS,IRD,
Univ.Paris-Sud, Université Paris-Saclay, cedex, France
[3] Ludwig-Maximilians Universität München, Fakultät für Biologie, Biozentrum, Grosshaderner
Strasse 2, 82152 Planegg-Martinsried, Germany
[4] Institut de Systématique, Evolution et Biodiversité, UMR7205, Centre National de la
Recherche Scientifique, MNHN, Sorbonne Université, EPHE, Université des Antilles, 57 rue
Cuvier, 75005 Paris, France
* Correspondence: rebeiz@pitt.edu

Abstract

The genitalia present some of the most rapidly evolving anatomical structures in the animal

kingdom, possessing a variety of parts that can distinguish recently diverged species. In the

Drosophila melanogaster group, the phallus is adorned with several processes, pointed

outgrowths, that are similar in size and shape between species. However, the complex three-

dimensional nature of the phallus can obscure the exact connection points of each process.

Previous descriptions based upon adult morphology have primarily assigned phallic processes

by their approximate positions in the phallus and have remained largely agnostic regarding

their homology relationships. In the absence of clearly identified homology, it can be

challenging to model when each structure first evolved. Here, we employ a comparative

developmental analysis of these processes in eight members of the *melanogaster* species group

to precisely identify the tissue from which each process forms. Our results indicate that adult

phallic processes arise from three pupal primordia in all species. We found that in some cases

the same primordia generate homologous structures whereas in other cases, different

primordia produce phenotypically similar but remarkably non-homologous structures. This

suggests that the same gene regulatory network may have been redeployed to different

primordia to induce phenotypically similar traits. Our results highlight how traits diversify and

can be redeployed, even at short evolutionary scales.

Key Words: Homology, Drosophila, Genitalia, rapid evolution

Introduction:

Most studies of developmental evolution depend on our ability to precisely compare the same body parts in different species or populations. Rigorously establishing such homology relationships allows us to identify novel traits, which are often defined by their lack of homology (Reviewed in Moczek, 2008; G. Wagner, 2007). Many of the current model systems for the study of novelty focus on traits that arose in the distant past (Bruce & Patel, 2020; Clark-Hachtel & Tomoyasu, 2020; Emlen, Szafran, Corley, & Dworkin, 2006; Hinman, Nguyen, Cameron, & Davidson, 2003), making the investigation of their molecular origins difficult. These traits likely arose through a multitude of genetic changes and exist in organisms that are less amenable to genetic studies. Recently evolved traits found in the rapidly evolving tissues of model organisms can provide qualitative changes in morphology produced by genomes that are easily compared and modified. However, rapidly evolving anatomical structures pose a distinct challenge. When differences between the anatomy of two species are numerous and exaggerated, it can be difficult to disentangle which structures are ancestral and which represent newly derived novelties. Thus, while macroevolutionary novelties often appear as clear discontinuities in the evolutionary record, the more molecularly tractable micro- and mesoevolutionary novelties require us to consider their relationships in a rich and complicated comparative context (Abouheif, 2008). Overcoming this challenge is thus critical to develop a genetic portrait of morphological novelty.

Most assertions of homology are defined through establishing that the structure in question connects to an unambiguously homologous tissue in both species (Moczek, 2008). Contentious claims of homology often revolve around the question of whether a set of traits

are formed by the same cells or tissues. These assertions can be strengthened through

developmental analysis, where the primordium that initially forms the trait in question can be

determined (Tanaka, Barmina, & Kopp, 2009). This type of analysis is especially important in

complex three-dimensional traits, as resolution in the X, Y, and Z axes may be required. The

high spatial resolution of confocal microscopy generates three-dimensional renderings of entire

body parts, allowing us to define the position of structures relative to tissues that have

straightforward homology assignments (Klaus, Kulasekera, & Schawaroch, 2003). Many

developing tissues progressively become more complex over developmental time. The

formation of specific traits is often established only after the tissue that encompasses that trait

is identifiable, providing clear anchor points in a conserved tissue to establish homology. Thus,

developmental trajectories provide a rich context in which to disentangle ambiguous

relationships among rapidly evolving structures.

The terminalia (genitalia and analia) of drosophilids host an extensive assortment of

rapidly evolving morphological structures. Variation of terminal structures is often one of the

most reliable ways to differentiate species of *Drosophila* based on adult morphology (Bock &

Wheeler, 1972; Hsu, 1949; Markow & O'Grady, 2006; Okada, 1954). The male genital structures

are often divided into two major compartments: the periphallic parts surrounding the anus,

which mostly play a role in grasping the external surface of the female genitalia (Acebes, Cobb,

& Ferveur, 2003; Jagadeeshan & Singh, 2006; Kamimura & Mitsumoto, 2011; Masly &

Kamimura, 2014; Mattei, Riccio, Avilaa, Wolfner, & Denlinger, 2015; Robertson, 1988; Yassin &

Orgogozo, 2013), and the phallic parts (Figure 1), which comprise the intromittent organ. While

the homology of the periphallic organs has always been relatively straightforward, the phallus

itself has posed several challenges in this regard. In particular, the homology of the various

phallic processes, pointed outgrowths, remains controversial (Figure 2, Supplementary videos)

(reviewed in Rice et al., 2019). These outgrowths have been implicated in sexual conflict

between males and females, and in some species have been shown to physically interact with

corresponding pockets in the female genitalia (Kamimura, 2016; Muto, Kamimura, Tanaka, &

Takahashi, 2018; Yassin & Orgogozo, 2013) or induce copulatory wounds (Kamimura, 2007).

Male seminal proteins can enter the female circulatory system through these copulatory

wounds, and are associated with increased ovulation and reduced remating rates (Avila, Sirot,

Laflamme, Rubinstein, & Wolfner, 2011; Kamimura, 2010). To better understand how genital

morphology may coevolve we must better establish which homologous tissues have been

modified in each sex.

Two major sources have contributed to confusion regarding homology of phallic

processes. The first is the relationship of the postgonal sheath (referred to as aedeagal sheath

in Rice et al., 2019) with respect to these processes (Figure 2 H-I). Several authors consider the

postgonal sheath and postgonal processes (referred to as postgonites in Rice et al., 2019) of *D.*

melanogaster as substructures of a unified tissue that was usually referred to as the "posterior

parameres" (Bock & Wheeler, 1972; Okada, 1954; Tsacas, Bocquet, Daguzan, & Mercier, 1971).

While others designated the postgonal sheath as a separate tissue from the postgonal

processes (Al Sayad & Yassin, 2019; Bächli, Vilela, Andersson Escher, & Saura, 2004; Lachaise et

al., 2004). The three-dimensional nature and the presence of transparent cuticle has made it

difficult to determine the precise connection points of the processes to the tissues of the

phallus. Determining whether these processes were formed by a single or separate primordium

would help resolve this discordance. The second source of confusion is in regard to the

nomenclature used to compare the phallic processes in different members of the *melanogaster*

species group. The term "basal process" has been used to refer to a number of pointed

outgrowths that are attached to different phallic tissues in different species (Kamimura, 2007,

2010, 2016; Kamimura & Mitsumoto, 2011, 2012a, 2012b; Kamimura & Polak, 2011). Such a

designation implies a concept of homology independent of the exact anatomical position.

Yassin & Orgogozo (2013) sought to provide distinct terms, such as "spurs" and "hooks" for

outgrowths emanating from the same tissue, implying the potential for non-homology. Building

upon our recent revision of the male terminalia nomenclature of *D. melanogaster* (Rice et al.,

2019), developmental studies presented here allow us to provide a more detailed, homology-

informed nomenclature for these structures.

In this study, we characterized both the adult morphology and the development of the

pupal genitalia in five members of the *melanogaster* species subgroup and three outgroup

members from the larger *melanogaster* species group. This analysis allows us to determine

whether processes are homologous or of different origins. Tracing the development of the

phallus by confocal microscopy showed that all processes arise from three distinct pupal

primordia in all species. We found both that several similarly shaped processes arise from

distinct primordia, whereas in other cases, distinct processes arise from different parts of the

same primordium. In light of these analyses, we refined the identity and terminology of the

phallic processes and identify distinct homology groups. We map these different morphologies

on previously established phylogenies and identified multiple gain, loss, and homoplastic events

in the history of these diverse structures. Thus, our results demonstrate how developmental

approaches can resolve unclear relationships among rapidly evolving structures.

Figure 1:

Figure 1: The *D. melanogaster* pupal phallus is produced by three primordia

A) Left: A schematic representation of the adult phallus of *D. melanogaster,* the median gonocoxite outlined in dark blue, the pregonites highlighted in light blue, the postgonal sheath in light purple, the dorsal postgonal process in magenta, the ventral postgonal process in violet, and the aedeagus in light green. Right: the adult phallus of a *yw;+:+* line of *D. melanogaster*. **B)** Left: A schematic representation of the early developing pupal genitalia of *D. melanogaster*. The primordia of developing phallus with the ventral primordium in blue, the dorsolateral primordium in purple, and the central primordium in green. Right: Developing pupal phallus of a *D. melanogaster* arm-GFP transgenic line. Apical cellular junctions are shown, highlighting the overall morphology. Blue arrows indicate the position of the pregonal bristles. Aedeagal sheath is an alternative term for postgonal sheath, ventral postgonite is an alternative term for ventral postgonal process, and dorsal postgonite is an alternative term for dorsal postgonal process.

Figure 2

Figure 2: The rapidly evolving phallus is composed of three main components

A) Phylogeny for eight species of the *melanogaster* species group based on Obbard et al., 2012
with nodes that contain the *melanogaster* species subgroup and *melanogaster* species group

indicated by arrows. **B)** A schematic breakdown of the adult phalluses of each species. **C)** Light microscopy images of the whole adult phallus for each species. Image stacks that show the relative position of each part can be found in Supplementary videos **D)** Schematic representation of the ventral portion of the phallus (dark blue) which contains the pregonites (light blue) and processes (filled in dark blue) in *D. erecta* and *D. ananassae*. Light blue asterisks designate the position of the pregonites. **E**, **G**, **I**: Light microscopy images of microdissections of the adult phallus (here in lateral view, distal end pointing downward) separated in the ventral, central, and dorsolateral portions. **E)** Microdissections of the ventral portion processes shows the processes of *D. erecta* and *D. ananassae* are connected to the pregonites. Light blue asterisks designate the position of the pregonites. **F)** Schematic representation of the central portion of the phallus (light green), which contains processes (filled in light green) in *D. santomea, D. teissieri, D. orena,* and *D. malerkotliana*. **G)** Microdissections of the aedeagus confirms that the processes are physically attached to aedeagus. The aedeagus of the *D. malerkotliana* (dashed line) is translucent with only the process sclerotized. **H)** Schematic representation of the dorsolateral portion of the phallus (light purple), which contains two pairs of processes (filled in light purple) in *D. melanogaster*, and one pair in *D. santomea, D. teissieri, D. erecta, and D. orena*. **I)** Microdissection confirms that the processes are physically attached to postgonal sheath. In *D. santomea*, *D. teissieri*, *D. erecta*, and *D. malerkotliana* portions of the anterior postgonal sheath are translucent (outlined with dashed lines). The image of the *D. orena* dorsolateral portion was created by copying and mirroring one side of the structure, as it was difficult to flatten intact for imaging. Aedeagal sheath is an alternative term for postgonal sheath.

Materials and Methods:

Drosophila strains:

To study the evolution of the phallic processes in the *melanogaster* species subgroup we used

the following subset of described species, representing all major complexes: *D. santomea*

(Lachaise et al., 2000), *D. teissieri* (Tsacas, 1971), *D. orena* (Tsacas & David, 1978), *D. erecta*

(Tsacas & Lachaise, 1974), *D. melanogaster* (Meigen, 1830) from the *melanogaster* species

subgroup and the following outgroup species *D. biarmipes* (Malloch, 1924) from the *suzukii*

subgroup, *D. ananassae* (Doleschall, 1858), and *D. malerkotliana* (Parshad & Paika, 1964) from

the *ananassae* subgroup. Previous work has investigated the function of the copulatory

anatomy of all species we analyzed (Kamimura, 2007, 2016; Muto et al., 2018; Yassin &

Orgogozo, 2013). Stocks were obtained from both the National Drosophila Species Stock Center

at Cornell (*D. santomea* (14021-0271.01), *D. teissieri* (14021-0257.01), *D. orena* (14021-

0245.01), *D. erecta* (14021-0224.01), *D. biarmipes* (14023-0361.09), *D. ananassae* (14024-

0371.13), the Bloomington Drosophila Stock Center *D. melanogaster* armadillo-GFP, arm-GFP,

(Bloomington stock number #8556), and from the lab of Dr. Thomas Williams, *D. malerkotliana*.

Sample collection, dissection, and fixation:

Male white pre-pupae were collected at room temperature and incubated in a petri dish

containing a moistened Kimwipe at 25°C prior to dissection. After incubation, pupae were

impaled in their anterior region and immobilized within a glass dissecting well containing

Phosphate Buffered Saline (PBS). The posterior tip of the pupa (20–40% of pupal length) was

separated and washed with a P200 pipette to flush the pupal terminalia into solution. Samples

were then collected in PBS with 0.1% Triton-X-100 (PBT) and 4% paraformaldehyde (PFA, E.M.S.

Scientific) on ice, and multiple samples were collected in the same tube. Samples were then

fixed in PBT + 4% PFA at room temperature for 30 min, washed three times in PBT at room

temperature, and stored at 4°C.

Immunohistochemistry and microscopy:

After fixation developing pupal genitalia of all species except *D. melanogaster* were stained

with rat anti-E-cadherin, 1:100 in PBT (DSHB Cat# DCAD2,RRID:AB_528120) overnight at 4°C,

followed by an overnight at 4°C incubation with anti-rat 488, 1:200 (Invitrogen, Carlsbad, CA) to

visualize apical cell junctions. For *D. melanogaster*, an *armadillo*-GFP tagged line (Bloomington

stock number #8556) was used to visualize apical cell junctions (Huang et al., 2012).

Fluorescently labeled samples were mounted in glycerol mounting solution (80% glycerol, .1M

Tris, pH 8.0) on microscope slides coated with poly-L-lysine (Thermo Fisher Scientific #86010).

Samples for all species except *D. melanogaster* were imaged at 20X on a Leica TCS SP8 confocal

microscope. *D. melanogaster* samples were imaged at either 20X or 40X on an Olympus

Fluoview 1000. As the imaged structures are three-dimensional in nature, we used the program

MorphoGraphX (de Reuille et al., 2015) to render and manipulate images in three-dimensions.

This allowed us to rotate the samples to better present the most informative perspectives of

the various phallic structures.

For light microscopy of adult phallic microdissections, samples were mounted in PVA

Mounting Medium (BioQuip) until fully cleared and imaged at 20X magnification on a Leica DM

2000 with a Leica DFC450C camera and the resulting images were enhanced using Adobe

Photoshop. For light microscopy images and videos of whole phallus samples, genitalia were

dissected in water, cleared overnight in 10% KOH at RT. Mounted in a drop of Dimethyl

Hydantoin Formaldehyde (Steedman, 1958) on a coverslip and oriented using 2 mounting

needles before the resin hardens. Coverslip were positioned on a microscope slide, the hard

drop facing away from the microscope lens. Images were acquired on Ti2-Eclipse Nikon

microscope equipped with a 20x plan apochromatic lens and a 5.5 M sCMOS camera (PCO,

Kelheim, Germany). Each preparation was imaged as a z-stack (z-step = 2 µm). The stacks are

presented as raw images. Stacks of images were also projected into single extended depth-of-

field images using Helicon Focus software (HeliconSoft) and the resulting projections were

enhanced using Adobe Photoshop.

Establishing designated early, middle, and late timepoints:

We used confocal microscopy to establish a time course of the developing phallus (Figures S1-

S3). To compare the development of the phallus of these species, we needed to examine

whether all analyzed species develop at the same rate after pupal formation. Due to the large-

scale changes seen in the phallus of these species we used two stable features found outside of

the phallus to calibrate developmental timing. In all analyzed species, the epandrial ventral lobe

(lateral plate) and surstylus (clasper) first appear as a single continuous structure early in

development but then separate from each other as development progresses (Figure S2). We

use the beginning of this separation as a landmark for the "early" developmental timepoint. We

also used the midpoint of this progression to approximate the "mid" timepoint. This

intermediate timepoint is useful in showing which tissue the phallic processes protrude from

during development. In all species, the adult cerci (anal plates) directly abut against one

another but during "early" and "mid" development, these structures are separated from one

another by a large gap (Figure S3). We designate "late" timepoint as directly preceding the

closing of this gap between the cerci.

Results:

Unpigmented cuticle reveals undescribed connection points in the phallus

In order to better understand how the processes surrounding the aedeagus are physically connected to the neighboring tissues of the phallus, we imaged whole (Figure 2 B,C) and micro-dissected adult phalluses in eight members of the *melanogaster* species group (Figure 2 D-I). The phallus of each species can be broken into three discrete parts. The ventral portion (Figure 2 D,E) contains the pregonites, an outgrowth that contains three bristles, and the median gonocoxite (the central section of the shield shaped hypandrium). The central portion (Figure 2 F,G) contains the aedeagus, through which sperm is transferred. The dorsolateral portion contains the postgonal sheath (referred to as aedeagal sheath in (Rice et al., 2019)), a flat sheet that wraps around the aedeagus and the pair of processes known as the postgonal processes (referred to as postgonites in Rice et al., 2019) (Figure 2 H,I). Analysis of these dissections support the designation of the postgonal sheath and postgonal processes as a single tissue (Bock & Wheeler, 1972; Okada, 1954; Tsacas et al., 1971). Furthermore, we found that certain species had processes connected to different portions of the phallus—ventral portion (*D. erecta*, *D. ananassae*), central portion (*D. santomea*, *D. teissieri*, *D. orena*, and *D. malerkotliana*), dorsolateral portion (all members of the *melanogaster* subgroup).

While imaging, we observed that parts of the postgonal sheath in the *melanogaster* species subgroup and *D. malerkotliana* were partially translucent, and only easily visible after microdissection. It is this translucent tissue of the postgonal sheath that physically connects to the postgonal processes in *D. melanogaster*. These observations highlight that, due to their transparency, determining the exact connection points between the processes and the rest of

the phallus can be difficult to visualize by traditional light microscopy approaches. To test

whether the different connection points of the phallic processes observed in the adult reflect

separate homology groups we investigated whether these phallic processes were initially

produced by the same or different primordia during development.

Phallic structures develop from three primordia in D. melanogaster

To date, the morphogenesis of the three-dimensional adult phallic structures from the

epithelium of the larval genital disc has been investigated only in *D. melanogaster* (Ahmad &

Baker, 2002; Epper, 1983). Additionally, using surgical fragmentation of the larval genital disc,

Bryant & Hsei, 1977 provided a fate map for the different adult structures. They showed that

the phallus is situated at the subcenter of the symmetrical imaginal disc and is surrounded on

each side by a primordium that will produce the medium gonocoxite and pregonites. However,

the sequence and timing of the appearance of the various substructures of the phallus during

development, remains unknown. By finding the key points in development where substructures

first emerge, we can determine the primordium from which each process initially form.

Early in *D. melanogaster* pupal development (see timepoint designation in the Materials

and Methods) the phallus is separated into three primordia: ventral, dorsolateral and central

(Figure 1B). As the pupal phallus continues to develop from this point, the ventral primordia

form a pair of processes (Figure 1A). This pair develops into the small processes known as the

pregonites that can be recognized from the presence of the minute bristles (Figure 1B arrows).

The dorsolateral primordia produce two processes (one dorsal and one ventral) (Figure 3B).

These processes ultimately develop into the ventral and dorsal postgonal processes (referred to

as ventral postgonite and dorsal postgonite in Rice et al., 2019) (Figure 1A). The remaining parts

of the dorsolateral tissue develop into the large flaps of the postgonal sheath (Figure 1A). The

central primordium of *D. melanogaster* develops into the aedeagus and lacks a process (Figure

1A).

The three primordia are conserved across species

Several studies have analyzed pupal development of the terminalia in species outside of

D. melanogaster, but did not investigate phallic structures (Glassford et al., 2015; Hagen et al.,

2019; Smith, Davidson, & Rebeiz, 2019). To determine whether the features of phallic

development observed in *D. melanogaster* are conserved in members of the *melanogaster*

species group, we produced a developmental time course for the remaining seven species

studied here (Figure S1-S3). We found that all adult phallic organs develop from three

primordia similar to the ones described in *D. melanogaster*. Nonetheless, significant

interspecific differences were observed regarding the timing of development (Figure S1). As we

only used one strain per species, we cannot comment if these are particular properties of the

strains/laboratory conditions we used or are general differences between the species. We

found that most species had early, mid, and late developmental timepoints within a six-hour

window of each other, while *D. orena* and *D. teissieri* showed particularly divergent

developmental timing (Figure S1-S3).

Different processes emerge from different primordia

The developmental analysis of the eight species used in this study allowed us to test

whether the phallic processes seen in the adults of each species were produced by the same

primordia. We began by investigating the ventral primordium (Figure 3, Figure S4), which

develops into the pregonites in all analyzed species. While the size of the pregonites varies

between species, during mid-development (Figure 4B) we can identify recognizable outgrowths

from the ventral primordium, consistent with a highly conserved developmental trajectory.

Interestingly, an additional pregonal process is found in two distantly-related species, *D. erecta*

and *D. ananassae*. Both *D. erecta* and *D. ananassae,* produce two processes from their ventral

primordia, a large pregonal medial process and a second smaller lateral process which contains

the three pregonal bristle cells and overall resembles the pregonites of other species. To

determine whether this additional process was produced by duplication or fission of the

pregonite we inspected early pupal timepoints. We found that initially a single process is

formed (Figure 3A), which during mid-development asymmetrically splits along the medial-

lateral axis to form the distinct lobe-like pregonal medial process (Figure 3B). These asymmetric

projections then extend in late development to form the larger pregonal medial process and

smaller pregonite (Figure 3C). Thus, although the ventral primordium produces the pregonite in

all species we examined, in *D. erecta* and *D. ananassae* the ventral primordium is split into the

pregonite and a pregonal medial process.

The dorsolateral primordium (Figure 4, Figure S5) showed a number of large

evolutionary changes within the *melanogaster* species group. We found that no species, other

than *D. melanogaster*, develop the ventral process that forms the ventral postgonal process

(Figure 4). By contrast, all members of the *melanogaster* species subgroup form dorsal

postgonal processes. Outside of the *melanogaster* subgroup, we did not find any modifications

of the dorsolateral primordium, which develops into a single thin, strongly sclerotized structure

in those species that resembles the postgonal sheath of *D. melanogaster*. However, the size,

and shape of these homologous structures significantly differ among species (Figure 2 H,I),

ranging from the flat rod-like processes in *D. biarmipes*, to the strongly pointed sinuate

processes in *D. ananassae*, and the minute, transparent sclerites in *D. malerkotliana*.

While the central primordium (Figure 5, Figure S6) forms a simple aedeagus that lacks

processes in *D. melanogaster*, we note processes which develop in *D. santomea*, *D. teissieri*, *D.*

orena and *D. malerkotliana*. Early in development, the central primordia of all species analyzed

are similar in size and shape (Figure 5A). However, during mid-development, in *D. santomea*, *D.*

teissieri, and *D. orena*, the ventral side of the central primordium elongates to form a process

(Figure 5B). The process of *D. teissieri* and *D. orena* splits along the ventral midline to form a

pair of processes, while in *D. santomea*, it forms one rounded structure. These processes

further elongate in late development to more closely resemble the size and shape of their adult

counterparts (Figure 5C).

Okada, 1954 and Bock & Wheeler, 1972 suggested that the aedeagus in the

melanogaster species group were of two types: fused like in *D. ananassae* and split like in *D.*

malerkotliana. Indeed, we did not observe any process in the central primordium of *D.*

ananassae, whereas a pair of processes develops in *D. malerkotliana*. Kamimura, 2007

suggested that the aedeagus of *D. malerkotliana* has degenerated and was replaced by a pair of

lateral processes. During early development, the central primordium of *D. malerkotliana* is

similar to all other analyzed species (Figure 5A). However, by mid-development, the lateral

sides of the central primordium extend, forming a pair of processes, while the medial-dorsal

and medial-ventral sides of the central primordium fail to extend (Figure 5B). Late in

development, the proximal-dorsal side of the lateral process constricts, conferring a hook like

shape (Figure 5C). As this substructure is produced from the lateral portions of the central

primordium and not from the ventral portion, it is likely non-homologous to the aedeagal

ventral processes of the *yakuba* and *erecta* complexes. We therefore propose the term

aedeagal lateral process for this substructure of *D. malerkotliana*.

Figure 3:

Figure 3: Processes developing from the ventral primordium are found in all members of the
melanogaster species group.
377
A-C) Signal from the apical cellular junctions was used to highlight the overall morphology of
developing pupal genitalia. *D. teissieri*, *D. orena*, *D. erecta*, *D. biarmipes*, and *D. ananassae* were
stained for ECAD while apical cell junctions were visualized in *D. melanogaster* by detecting
arm-GFP (see methods). **A)** Early in development, a pair of processes, that will form the
pregonites, can be visualized in the ventral primordia in all species shown (light blue). **B)** By
mid-development, large processes can be found in all species except *D. melanogaster*. In *D.*
erecta and *D. ananassae*, the pregonite is split into a large pregonal medial process and a small
lateral bristle-bearing process. **C)** By late development, the pregonites have extended to their
full adult size and shape. The pregonites are connected to the medial-ventral portion of the
median gonocoxite, see Figure S4. **D)** Schematic representations of the pregonites (blue) and
pregonal medial process (teal) showing their approximate size, number, and connections to the
medial gonocoxite (outlined in black). All images have the same axes, V (Ventral), A (Anterior),
M (Medial) and are the same scale.

Figure 4:

Figure 4: Processes produced by the dorsolateral primordium are found only in the *melanogaster* subgroup.

A-C) Signal from the apical cellular junctions was used to highlight the overall morphology of developing pupal genitalia. *D. santomea*, *D. teissieri*, *D. orena*, and *D. erecta*, were stained for ECAD while *D. melanogaster* samples used arm-GFP. **A)** Early in development, the dorsolateral primordium is a smooth lobe like structure in all analyzed species. **B)** By mid-development, all shown species form processes in the dorsal portion of the dorsolateral primordium (violet). *D. melanogaster* also forms an additional pair of processes in the ventral portion of dorsolateral primordium (magenta). **C)** By late development, the dorsal and ventral processes have extended to a long thin shape. Both the ventral and dorsal processes are connected to the medial-anterior part of the postgonal sheath which is formed by the remaining tissue of the dorsolateral primordium. **D)** Schematic representations of the dorsal (violet) and ventral (magenta) postgonal process showing where they connect to the postgonal sheath (outlined in black). All images have the same axes, V (Ventral), A (Anterior), M (Medial) and are the same scale. Ventral postgonite is an alternative term for ventral postgonal process, and dorsal postgonite is an alternative term for dorsal postgonite.

Figure 5:

412 Figure 5: Processes developing from the central primordium are found in the *yakuba/erecta*
413 and *bipectinata* complexes.
414
A-C) Signal from the apical cellular junctions was used to highlight the overall morphology of
developing pupal genitalia. *D. santomea*, *D. teissieri*, *D. orena*, and *D. erecta*, were stained for
ECAD. **A)** Early in development, the central primordium forms a flat donut-shaped structure in
all species shown. **B)** By mid-development, the ventral portion of the aedeagus is extended in *D.*
santomea, *D. teissieri*, and *D. orena* in what will form the aedeagal ventral process (dark green).
In *D. malerkotliana* the lateral edges of the central primordium extend anteriorly in what will
form the aedeagal lateral process (yellow-green). **C)** By late development, the aedeagal ventral
process and the aedeagal lateral process further extend from the aedeagus. **D)** Schematic
representations of the aedeagal ventral process (dark green) and aedeagal lateral process
(yellow-green) showing where they connect to the aedeagus (outlined in black). All images have
the same axes, V (Ventral), A (Anterior), M (Medial) and are the same scale.

     Figure 6:

428   Figure 6: Model of the evolutionary of phallic processes in the *melanogaster* species group
429
A,C,E) Phylogeny of the 8 analyzed species based on Obbard et al., 2012. **A)** Parsimony suggests
that the pregonites originated outside of the *D. melanogaster* species group. **B)** Light blue
represents the pregonites and teal represents the pregonal medial process. **C)** Parsimony
suggests that the ventral postgonites originated in the *melanogaster* complex (*D. melanogaster*,
D. simulans, *D. mauritiana*, *D. sechellia*) and that the dorsal postgonal process originated in the
melanogaster subgroup. **D)** Violet represents the ventral postgonal process, magenta
represents the dorsal postgonal process. E) Parsimony suggests that the aedeagal ventral
process originated at the base of the *erecta* and *yakuba* complexes. Additionally, parsimony
suggests that the aedeagal lateral process originated in the bipectinate complex. **F)** Dark green
represents the aedeagal ventral process and yellow-green represents the aedeagal lateral
process. Ventral postgonite is an alternative term for ventral postgonal process, and dorsal
postgonite is an alternative term for dorsal postgonite.

Discussion:

The rapid evolution of morphological structures is an attractive subject for study, as it

allows us to glimpse at the molecular and genetic causes of dramatically remodeled and

restructured anatomical forms. Here, we examined some of the most rapidly evolving

morphologies of *Drosophila melanogaster* and its close relatives. Despite decades of research,

many of the intricate phallic processes have eluded our ability to clearly classify their homology

relationships. By studying the developmental trajectories of these processes in multiple species,

we have better defined their physical connections, and clarified which structures most likely

share ancestry. This research highlights the distinct challenges in studying novelties at

mesoevolutionary scales.

Classification and nomenclature of rapidly evolving phallic structures

Our results suggest that the great diversity of the phallic structures of the eight species

studied here cluster into three homology groups corresponding to the three pupal primordia,

leading us to propose revised naming conventions. First, our developmental analysis supports

the notion, initially suggested by Okada, 1954, that the weakly sclerotized postgonal sheath and

strongly sclerotized postgonal processes in *D. melanogaster*, are both parts of the same tissue,

which *Drosophila* systematists called the "posterior paramere" e.g. (Bock & Wheeler, 1972;

Tsacas et al., 1971). Because the term "posterior paramere" is itself synonymous to the term

"postgonite" in Dipteran systematics (Tsacas et al., 1971; van Emden & Hennig, 1970), we

suggest using the term "postgonite" to encompass the combined tissue produced by the

dorsolateral primordia in species of the *melanogaster* group (including both the postgonal

sheath and the processes), and the term "postgonal processes" to designate the strongly

sclerotized branches emerging from this tissue in the *melanogaster* subgroup. Second, our

results also show that the structures previously called the "basal processes" (Kamimura, 2007,

2010, 2016; Kamimura & Mitsumoto, 2011, 2012a, 2012b; Kamimura & Polak, 2011), develop

from different primordia and are therefore most likely non-homologous. We suggest therefore

to give them distinct names that directly relate to the tissues that produce them: aedeagal

ventral process in species of the *yakuba* complex (synonymous to Yassin & Orgogozo, 2013

phallic spur) and *D. orena* (synonymous to Yassin & Orgogozo, 2013 phallic hook), the pregonal

medial process in *D. erecta* and *D. ananassae,* and aedeagal lateral process in *D. malerkotliana*

and species of the *bipectinata* complex (Table 1).

Evolution of the phallic structures

Mapping character states over robust phylogenies provide the opportunity to

distinguish novel from recurrent (homoplastic) states as well as derived states (synapomorphic)

from ancestral (symplesiomorphic) ones (Figure 6). For example, our demonstration of the

development of an additional pregonal process in *D. erecta* and *D. ananassae* is likely recurrent,

as illustrations from Bock & Wheeler, 1972 suggest that this configuration of the pregonites

might have recurrently evolved in this clade. Reversals to ancestral states through secondary

losses represent another mechanism of recurrent evolution. The lack of the aedeagal ventral

processes in *D. erecta*, is more likely due to loss rather than an independent gain of the

aedeagal ventral process in *D. orena*.

In the *melanogaster* subgroup, all species contain a strongly-sclerotized dorsal

postgonal process which develops as a localized extension within the dorsolateral primordium.

The development of a strongly-sclerotized ventral postgonal process is a definitive novelty in *D.*

melanogaster and allied species of the *melanogaster* complex. Although we did not find

structures resembling the dorsal postgonal processes in members outside of the *melanogaster*

species subgroup that we studied here, polarization remains difficult. Indeed, Okada, 1954 and

Bock & Wheeler, 1972 reported the presence of "basal processes of the posterior parameres"

in multiple members of the *melanogaster* species group. Similarly, Bächli et al., 2004 illustrated

the presence of "ribbon-shaped process" in several members of the *obscura* group which is

sister to the *melanogaster* species group. Further taxonomic sampling and better phylogenetic

resolution of those clades are required to draw a more complete picture of the evolution of the

postgonal differentiation outside the *melanogaster* subgroup. The novel structures described

here may present an excellent model to study the molecular mechanisms producing novelty.

Although we do not address the function of the phallic processes, other research groups

have associated these with sexual conflict. The various processes of the phallus have been

implicated in copulatory wounding of the female (Kamimura, 2007, 2016; Muto et al., 2018;

Yassin & Orgogozo, 2013). Furthermore, studies have found that some of the phallic processes

pivot from pointing posteriorly to pointing laterally, when the phallus is everted during

copulation, thus directing how they interact with the female reproductive tract (Kamimura,

2010). The ability to pivot during copulation correlates with the homology groups we have

found in this study. The ventral and dorsal postgonal processes, and pregonites pivot during

copulation while the aedeagal ventral process does not change orientation. This may be due to

the direct connection of the aedeagal ventral process to the aedeagus. Surprisingly the

aedeagal lateral process, which is also directly connected to the aedeagus, pivots laterally

during copulation, which may only be possible due to the loss of aedeagal sclerotization,

making the tissue between the aedeagal lateral processes flexible. Co-evolution between the

phallic processes and the female genitalia has been suggested and several novel modifications

of the female genitalia have been identified (Kamimura, 2007; Yassin & Orgogozo, 2013). A

developmental analysis of the female genitalia of these species along with three-dimensional

analysis of copulating flies similar to studies in *D. melanogaster* (Mattei et al., 2015; Shao et al.,

2019) would provide vital context for the potential co-evolution of novel male and female

structures.

Developmental mechanisms underlying phallic evolution

A major challenge in the evo-devo field has been to identify the molecular mechanisms

driving morphological novelty (Linz, Hu, & Moczek, 2020; Moczek, 2008; Rebeiz, Patel, &

Hinman, 2015; G. P. Wagner & Lynch, 2010). While macroevolutionary novelties have been the

focus of coarse-grained molecular study (Bruce & Patel, 2020; Clark-Hachtel & Tomoyasu, 2020;

Emlen et al., 2006; Hinman et al., 2003; Prud'Homme et al., 2011), much hope has been placed

on rapidly diverging structures in molecularly amenable systems (Rebeiz & Tsiantis, 2017). Our

work highlights distinct underexplored challenges to interpreting and advancing these model

systems. The ambiguous ancestry of similar parts which appear in different locations causes us

to consider multiple models to explain their emergence. These parts may arise by parallelism –

a predisposition to drive similar new structures by co-opting the same networks (Abouheif,

2008). Alternately, it is entirely possible that these structures are indeed ancestral but have

undergone massive tissue reorganizations to reposition their attachment points. Such

repositioning could be caused by moving the location of a critical signal or transcription factor

within the tissues. Alternately, these structures could be specified before the discernable

tissues of the phallus are separated, and their migration could be caused by differences in

tissue folding. Under this scenario, we would anticipate that critical tissue patterning regulators

of these processes are activated before these tissues become discernable. Finally, it is entirely

possible that completely different networks account for the appearance of these unique

structures. Developmental genetic analysis of the genes that produce the phallic processes

described above will aid us in distinguishing these models. Recent work has identified several

genes that are spatially restricted to the pregonites and postgonal processes of *D.*

melanogaster providing an ideal set of candidates to examine (Vincent et al., 2019). Thus, we

envision that detailed mechanisms of parallelism, repositioning, and novelty will emerge from

studying systems where both network architecture is accessible, and genetic manipulations can

be introduced to test the sufficiency of these mechanisms to produce these novel

morphological structures.

Acknowledgments:

We would like to thank Deepak Dharmadhikari for his help with imaging, the Cornell Species

and Bloomington Stock Centers for providing fly strains used in this study, Ben Vincent and the

Rebeiz lab for their comments on the project and manuscript, Virginie Courtier-Orgogozo,

Masanori Toda, Yoshitaka Kamimura and the Terminalia consortium for their insights on this

project. We would also like to thank TaxoDros and the Japan Drosophila Database for their

work cataloging resources for the original species descriptions for those analyzed.

Tables:

New nomenclature	Previous terminology
postgonite	posterior paramere (Bock & Wheeler, 1972; Tsacas et al., 1971)
dorsal postgonal process	dorsal postgonite (Rice et al., 2019; Vincent et al., 2019) dorsal branch (Kamimura, 2010, 2016; Kamimura & Mitsumoto, 2011) dorsal paramere (Bryant & Hsei, 1977)
ventral postgonal process	ventral postgonite (Rice et al., 2019; Vincent et al., 2019) ventral branch (Kamimura, 2010; Kamimura & Mitsumoto, 2011) ventral paramere (Bryant & Hsei, 1977)
aedeagal ventral process	phallic spur (Yassin & Orgogozo, 2013) phallic hook (Yassin & Orgogozo, 2013) ventral branch (Kamimura, 2012, 2016; Kamimura & Mitsumoto, 2012b, 2012a; A. E. Peluffo et al., 2015; A. Peluffo et al., 2021)
aedeagal lateral process	basal process (Kamimura, 2007; Kamimura & Polak, 2011)
pregonal medial process	basal process (Kamimura, 2007), ventral branch (Kamimura, 2016)

553 Table 1: Table of correspondence between terms previously used in publications and proposed
554 nomenclature.

 Supplemental figures:

556 **Figure S1:**

558 ⬤ early timepoint   ⬤ mid timepoint   ⬤ late timepoint
559 Figure S1:

561 A summary of our designations of early (beige) mid (brown) and late (orange) developmental
562 timepoints for each species.

Figure S2:

565   Figure S2:
566
567   Full ECAD time course of ventral genitalia across the *melanogaster* species group.
568   Developing pupal genitalia of *D. santomea*, *D. teissieri*, *D. orena*, *D. erecta*, *D. biarmipes*, *D.*
569   *ananassae*, and *D. malerkotliana* stained for ECAD, apical cellular junctions, highlighting the
570   overall morphology. Note that *D. melanogaster* samples use a transgenic line arm-GFP that also
571   labels the apical cell junctions.

Figure S3:

574   Figure S3:
575

576   Full ECAD time course of dorsal genitalia across *melanogaster* species group.
577   Developing pupal genitalia of *D. santomea*, *D. teissieri*, *D. orena*, *D. erecta*, *D. biarmipes*, *D.*
578   *ananassae*, and *D. malerkotliana* stained for ECAD, apical cellular junctions, highlighting the
579   overall morphology. Note that *D. melanogaster* samples use a transgenic line arm-GFP that also
580   labels the apical cell junctions.

Figure S4:

582
583
584   Figure S4:

Developing ventral primordia of all analyzed species. For *D. melanogaster* an arm-GFP line of
was used to highlight the apical cellular junctions. **A-C)** The median gonocoxite is highlighted in
dark blue, the pregonite is highlighted in with light blue, and the pregonal medial process is
highlighted in teal **D)** Schematic representation of the median gonocoxite (dark blue), the
pregonite (light blue), pregonal medial process (teal). Note that *D. melanogaster* samples use a
transgenic line arm-GFP that also labels the apical cell junctions, while all other samples are
stained for E-cadherin.

Figure S5:

Figure S5:

Developing dorsolateral primordia of all analyzed species. For *D. melanogaster* an arm-GFP line of was used to highlight the apical cellular junctions. **A-C)** The postgonal sheath is highlighted in light purple, the ventral postgonal process is highlighted in with violet, and the dorsal postgonal process is highlighted in magenta **D)** Schematic representations of the postgonal sheath (light purple), the ventral postgonal process (violet), and dorsal postgonal process (magenta). Aedeagal sheath is an alternative term for postgonal sheath, ventral postgonite is an alternative term for ventral postgonal process, and dorsal postgonite is an alternative term for dorsal postgonal process. Note that *D. melanogaster* samples use a transgenic line arm-GFP that also labels the apical cell junctions, while all other samples are stained for E-cadherin.

Figure S6:

Figure S6:

Developing central primordia of all analyzed species. **A-C)** The aedeagus is highlighted in light
green, the aedeagal ventral process is highlighted in with dark green and the aedeagal lateral
process is highlighted in yellow-green. **D)** Cartoon representations of the aedeagus (light green)
aedeagal ventral process (dark green) and aedeagal lateral process (yellow-green). Note that *D.*
melanogaster samples use a transgenic line arm-GFP that also labels the apical cell junctions,
while all other samples are stained for E-cadherin.

References:

Abouheif, E. (2008). Parallelism as the pattern and process of mesoevolution. *Evolution and*
Development, *10*(1), 3–5. https://doi.org/10.1111/j.1525-142X.2007.00208.x
Acebes, A., Cobb, M., & Ferveur, J. F. (2003). Species-specific effects of single sensillum ablation
on mating position in Drosophila. *Journal of Experimental Biology*, *206*(17), 3095–3100.
https://doi.org/10.1242/jeb.00522
Ahmad, S. M., & Baker, B. S. (2002). Sex-Specific Deployment of FGF Signaling in Drosophila
Recruits Mesodermal Cells into the Male Genital Imaginal Disc, *109*, 651–661.
Al Sayad, S., & Yassin, A. (2019). Quantifying the extent of morphological homoplasy: A
phylogenetic analysis of 490 characters in Drosophila . *Evolution Letters*, *3*(3), 286–298.
https://doi.org/10.1002/evl3.115
Avila, F. W., Sirot, L. K., Laflamme, B. A., Rubinstein, C. D., & Wolfner, M. F. (2011). Insect
seminal fluid proteins: Identification and function. *Annual Review of Entomology*, *56*, 21–
40. https://doi.org/10.1146/annurev-ento-120709-144823
Bächli, G., Vilela, C., Andersson Escher, S., & Saura, A. (2004). *The Drosophilidae (Diptera) of 252*
Fennoscandia and Denmark. Fauna Entomologica Scandinavica. Brill.
Bock, I. R., & Wheeler, M. R. (1972). The Drosophila melanogaster species group. *The University*
of Texas Publication, *VII*(7213), 1–102.
Bruce, H. S., & Patel, N. H. (2020). Knockout of crustacean leg patterning genes suggests that
insect wings and body walls evolved from ancient leg segments. *Nature Ecology and*
Evolution, *4*(12), 1703–1712. https://doi.org/10.1038/s41559-020-01349-0
Bryant, P. J., & Hsei, B. W. (1977). Pattern formation in asymmetrical and symmetrical imaginal
discs of Drosophila melanogaster. *American Zoologist*, *17*(3), 595–611.
https://doi.org/10.1093/icb/17.3.595
Clark-Hachtel, C. M., & Tomoyasu, Y. (2020). Two sets of candidate crustacean wing
homologues and their implication for the origin of insect wings. *Nature Ecology and*
Evolution, *4*(12), 1694–1702. https://doi.org/10.1038/s41559-020-1257-8
de Reuille, P. B., Routier-Kierzkowska, A. L., Kierzkowski, D., Bassel, G. W., Schüpbach, T.,
Tauriello, G., … Smith, R. S. (2015). MorphoGraphX: A platform for quantifying
morphogenesis in 4D. *ELife*, *4*(MAY). https://doi.org/10.7554/eLife.05864
Doleschall, C. L. (1858). Derde bijdrage tot de kennis der Dipteren fauna van Nederlandsch
Indie. *Natuurkundig Tijdschrift Voor Nederlandsch Indie*, *17*, 73–128.
Emlen, D. J., Szafran, Q., Corley, L. S., & Dworkin, I. (2006). Insulin signaling and limb-patterning:
Candidate pathways for the origin and evolutionary diversification of beetle "horns."
Heredity, *97*(3), 179–191. https://doi.org/10.1038/sj.hdy.6800868
Epper, F. (1983). The Evagination of the Genital Imaginal Discs of Drosophila melanogaster II.
Morphogenesis of the Intersexual Genital Disc of the Mutant doublesex-dominant (dsx D).
Roux's Archives of Developmental Biology, *192*, 280–284.
Glassford, W. J., Johnson, W. C., Dall, N. R., Smith, S. J., Liu, Y., Boll, W., … Rebeiz, M. (2015). Co-
option of an Ancestral Hox-Regulated Network Underlies a Recently Evolved
Morphological Novelty Article. *Developmental Cell*, *34*(5), 520–531.
https://doi.org/10.1016/j.devcel.2015.08.005
Hagen, J. F. D., Mendes, C. C., Blogg, A., Payne, A., Tanaka, K. M., Gaspar, P., … Nunes, M. D. S.

(2019). Tartan underlies the evolution of Drosophila male genital morphology. *Proceedings*
of the National Academy of Sciences of the United States of America, *116*(38), 19025–
19030. https://doi.org/10.1073/pnas.1909829116
Hinman, V. F., Nguyen, A. T., Cameron, R. A., & Davidson, E. H. (2003). Developmental gene
regulatory network architecture across 500 million years of echinoderm evolution.
Proceedings of the National Academy of Sciences of the United States of America, *100*(23),
13356–13361. https://doi.org/10.1073/pnas.2235868100
Hsu, T. C. (1949). The external genital apparatus of male Drosophilidae in relation to
systematics. *The University of Texas Publication*, *4920*, 80–142.
Huang, J., Huang, L., Chen, Y. J., Austin, E., Devor, C. E., Roegiers, F., & Hong, Y. (2012).
Differential regulation of adherens junction dynamics during apical-basal polarization.
Development, *139*(3), 4001–4013. https://doi.org/10.1242/jcs.086694
Jagadeeshan, S., & Singh, R. S. (2006). A time-sequence functional analysis of mating behaviour
and genital coupling in Drosophila: Role of cryptic female choice and male sex-drive in the
evolution of male genitalia. *Journal of Evolutionary Biology*, *19*(4), 1058–1070.
https://doi.org/10.1111/j.1420-9101.2006.01099.x
Kamimura, Y. (2007). Twin intromittent organs of Drosophila for traumatic insemination.
Biology Letters, *3*(4), 401–404. https://doi.org/10.1098/rsbl.2007.0192
Kamimura, Y. (2010). Copulation anatomy of Drosophila melanogaster (Diptera: Drosophilidae):
Wound-making organs and their possible roles. *Zoomorphology*, *129*(3), 163–174.
https://doi.org/10.1007/s00435-010-0109-5
Kamimura, Y. (2012). Correlated evolutionary changes in Drosophila female genitalia reduce the
possible infection risk caused by male copulatory wounding. *Behavioral Ecology and*
Sociobiology, *66*(8), 1107–1114. https://doi.org/10.1007/s00265-012-1361-0
Kamimura, Y. (2016). Significance of constraints on genital coevolution: Why do female
Drosophila appear to cooperate with males by accepting harmful matings? *Evolution;*
International Journal of Organic Evolution, *70*(7), 1674–1683.
https://doi.org/10.1111/evo.12955
Kamimura, Y., & Mitsumoto, H. (2011). Comparative copulation anatomy of the Drosophila
melanogaster species complex (Diptera: Drosophilidae). *Entomological Science*, *14*(4),
399–410. https://doi.org/10.1111/j.1479-8298.2011.00467.x
Kamimura, Y., & Mitsumoto, H. (2012a). Genital coupling and copulatory wounding in
drosophila teissieri (diptera: Drosophilidae). *Canadian Journal of Zoology*, *90*(12), 1437–
1440. https://doi.org/10.1139/cjz-2012-0186
Kamimura, Y., & Mitsumoto, H. (2012b). Lock-and-key structural isolation between sibling
Drosophila species. *Entomological Science*, *15*(2), 197–201.
https://doi.org/10.1111/j.1479-8298.2011.00490.x
Kamimura, Y., & Polak, M. (2011). Does surgical manipulation of drosophila intromittent organs
affect insemination success? *Proceedings of the Royal Society B: Biological Sciences*,
278(1707), 815–816. https://doi.org/10.1098/rspb.2010.2431
Klaus, A. V., Kulasekera, V. L., & Schawaroch, V. (2003). Three-dimensional visualization of
insect morphology using confocal laser scanning microscopy. *Journal of Microscopy*,
212(2), 107–121. https://doi.org/10.1046/j.1365-2818.2003.01235.x
Lachaise, D., Capy, P., Cariou, M.-L., Joly, D., Lemeunier, F., & David, J. R. (2004). Nine relatives

from one African ancestor: population biology and evolution of the Drosophila
melanogaster subgroup species. In R. S. Singh, M. K. Uyenoyama, & S. K. Jain (Eds.), *The*
Evolution of Population Biology (pp. 315–344). Cambridge University Press.
https://doi.org/10.1017/cbo9780511542619.019
Lachaise, D., Harry, M., Solignac, M., Lemeunier, F., Benassi, V., & Cariou, M. L. (2000).
Evolutionary novelties in islands: Drosophila santomea, a new melanogaster sister species
from Sao Tome. *Proceedings of the Royal Society B: Biological Sciences*, *267*(1452), 1487–
1495. https://doi.org/10.1098/rspb.2000.1169
Linz, D. M., Hu, Y., & Moczek, A. P. (2020). From descent with modification to the origins of
novelty. *Zoology*, *143*(August), 125836. https://doi.org/10.1016/j.zool.2020.125836
Malloch, J. R. (1924). Two Drosophilidae from Coimbatore. *Memoirs of the Department of*
Agriculture in India. Entomological Series, *8*(6), 63–65.
Markow, T. A., & O'Grady, P. M. (2006). *Drosophila: A Guide to Species Identification and Use*.
Academic Press. London.
Masly, J. P., & Kamimura, Y. (2014). Asymmetric mismatch in strain-specific genital morphology
causes increased harm to drosophila females. *Evolution*, *68*(8), 2401–2411.
https://doi.org/10.1111/evo.12436
Mattei, A. L., Riccio, M. L., Avilaa, F. W., Wolfner, M. F., & Denlinger, D. L. (2015). Integrated 3D
view of postmating responses by the Drosophila melanogaster female reproductive tract,
obtained by micro-computed tomography scanning. *Proceedings of the National Academy*
of Sciences of the United States of America, *112*(27), 8475–8480.
https://doi.org/10.1073/pnas.1505797112
Meigen, J. W. (1830). *Systematische Beschreibung der bekannten europäischen zweiflugeligen*
Insekten. Schulze.
Moczek, A. P. (2008). On the origins of novelty in development and evolution. *BioEssays*, *30*(5),
432–447. https://doi.org/10.1002/bies.20754
Muto, L., Kamimura, Y., Tanaka, K. M., & Takahashi, A. (2018). An innovative ovipositor for
niche exploitation impacts genital coevolution between sexes in a fruit-damaging
Drosophila. *Proceedings of the Royal Society B: Biological Sciences*.
https://doi.org/10.1098/rspb.2018.1635
Obbard, D. J., MacLennan, J., Kim, K. W., Rambaut, A., O'Grady, P. M., & Jiggins, F. M. (2012).
Estimating divergence dates and substitution rates in the drosophila phylogeny. *Molecular*
Biology and Evolution, *29*(11), 3459–3473. https://doi.org/10.1093/molbev/mss150
Okada, T. (1954). Comparative morphology of the drosophilid flies. I. Phallic organs of the
melanogaster group. *Kontyu*, *22*, 36–46.
Parshad, R., & Paika, I. J. (1964). Drosophilid survey of India. II. Taxonomy and cytology of the
subgenus Sophophora (Drosophila). *Research Bulletin of the Panjab University. Science.*,
15, 225–252.
Peluffo, A. E., Nuez, I., Debat, V., Savisaar, R., Stern, D. L., & Orgogozo, V. (2015). A major locus
controls a genital shape difference involved in reproductive isolation between Drosophila
yakuba and Drosophila santomea. *G3: Genes, Genomes, Genetics*, *5*(12), 2893–2901.
https://doi.org/10.1534/g3.115.023481
Peluffo, A., Hamdani, M., Vargas-Valderrama, A., David, J., Mallard, F., Graner, F., & Courtier-
Orgogozo, V. (2021). A morphological trait involved in reproductive isolation between

Drosophila sister species is sensitive to temperature. *BioRx*.
https://doi.org/10.1101/2020.01.20.911826

Prud'Homme, B., Minervino, C., Hocine, M., Cande, J. D., Aouane, A., Dufour, H. D., … Gompel,
N. (2011). Body plan innovation in treehoppers through the evolution of an extra wing-like
appendage. *Nature*, *473*(7345), 83–86. https://doi.org/10.1038/nature09977

Rebeiz, M., Patel, N. H., & Hinman, V. F. (2015). Unraveling the Tangled Skein: The Evolution of
Transcriptional Regulatory Networks in Development. *Annual Review of Genomics and*
Human Genetics, *16*(1), 103–131. https://doi.org/10.1146/annurev-genom-091212-
153423

Rebeiz, M., & Tsiantis, M. (2017). Enhancer evolution and the origins of morphological novelty.
Current Opinion in Genetics and Development, *45*, 115–123.
https://doi.org/10.1016/j.gde.2017.04.006

Rice, G., David, J. R., Kamimura, Y., Masly, J. P., Alistair, P., Nagy, O., … Yassin, A. (2019). A
standardized nomenclature and atlas of the male terminalia of Drosophila melanogaster.
Fly, *13*(1–4), 51–64. https://doi.org/10.1080/19336934.2019.1653733

Robertson, H. (1988). Mating Asymmetries and Phylogeny in the Drosophila melanogaster
Species Complex, *42*.

Shao, L., Chung, P., Wong, A., Siwanowicz, I., Kent, C. F., Long, X., & Heberlein, U. (2019). A
Neural Circuit Encoding the Experience of Copulation in Female Drosophila. *Neuron*,
102(5), 1025-1036.E6. https://doi.org/10.1016/j.neuron.2019.04.009

Smith, S. J., Davidson, L., & Rebeiz, M. (2019). Expansion of apical extracellular matrix underlies
the morphogenesis of a recently evolved structure. *BioRxiv*.
https://doi.org/doi.org/10.1101/686089

Steedman, H. F. (1958). Dimethyl Hydantoin Formaldehyde: A new Water-soluble Resin for Use
as a Mounting Medium. *Journal of Cell Science*, *99*(4), 451–452.
https://doi.org/10.1163/187529266X00220

Tanaka, K., Barmina, O., & Kopp, A. (2009). Distinct developmental mechanisms underlie the
evolutionary diversification of Drosophila sex combs. *Proceedings of the National Academy*
of Sciences, *106*(12), 4764–4769. https://doi.org/10.1073/pnas.0807875106

Tsacas, L. (1971). Drosophila teissieri, nouvelle espece africaine du groupe melanogaster et
note sur deux autres especes nouvelles pour l'Afrique (Dipt. Drosophilidae). *Bulletin de La*
Société Entomologique de France, *76*, 35–45.

Tsacas, L., Bocquet, C., Daguzan, M., & Mercier, A. (1971). Comparaison des genitalia males de
Drosophila melanogaster, de Drosophila simulans et de leurs hybrids. *Annales de La*
Société Entomologique de France, *7*, 75–93. Retrieved from
https://ci.nii.ac.jp/naid/10030580910/

Tsacas, L., & David, J. (1978). Une septieme espece appartenant au sous-groupe Drosophila
melanogaster Meigen: Drosophila orena spec. nov. du Cameroun. (Diptera: Drosophilidae).
Beiträge Zur Entomologie, *28*, 179–182.

Tsacas, L., & Lachaise, D. (1974). Quatre nouvelles especes de la Cote-d'Ivoire du genre
Drosophila, groupe melanogaster, et discussion de l'origine du sous-groupe melanogaster
(Diptera: Drosophilidae). *Annales de l'Université d'Abidjan Série E: Ecologie*, *7*, 193–211.

van Emden, F., & Hennig, W. (1970). *Taxonomists' glossary of genitalia of insects*. (S. L. Tuxen,
Ed.) (2nd ed.). Munksgaard, Copenhagen.

Vincent, B. J., Rice, G. R., Wong, G. M., Glassford, W. J., Downs, K. I., Shastay, J. L., … Rebeiz, M.
(2019). An atlas of transcription factors expressed in the Drosophila melanogaster pupal
terminalia. *G3: Genes|Genomes|Genetics*, *9*(December), 3961–3972.
https://doi.org/10.1101/677260
Wagner, G. (2007). The developmental genetics of homology. *Nature Reviews Genetics*, *8*(6),
473–479. https://doi.org/10.1038/nrg2099
Wagner, G. P., & Lynch, V. J. (2010). Evolutionary novelties. *Current Biology*, *20*(2), 48–52.
https://doi.org/10.1016/j.cub.2009.11.010
Yassin, A., & Orgogozo, V. (2013). Coevolution between Male and Female Genitalia in the
Drosophila melanogaster Species Subgroup. *PLoS ONE*, *8*(2).
https://doi.org/10.1371/journal.pone.0057158
804