Interacting phenotypes and the coevolutionary process: Interspecific indirect genetic effects alter coevolutionary dynamics

Running title: Trait interactions and coevolution

Stephen P. De Lisle[1,2], Daniel I. Bolnick[1], Edmund D. Brodie III[3], Allen J. Moore[4], and Joel W. McGlothlin[5]

Corresponding author: Stephen P. De Lisle   Email: stephen.de_lisle@biol.lu.se

1.   Department of Ecology & Evolutionary Biology
        University of Connecticut
        75 N. Eagleville Road
        Storrs, Connecticut, USA 06269
2   Present address:
        Evolutionary Ecology Unit, Department of Biology
        Lund University
        Solvegatan 37, Lund, Sweden
3   Department of Biology and Mountain Lake Biological Station
        University of Virginia
        485 McCormick Road
        Charlottesville, VA 22904 USA
4   Department of Entomology
        University of Georgia
        Athens, GA 30602 USA
5   Department of Biological Sciences
        Virginia Tech
        2125 Derring Hall, 926 West Campus Drive
        Blacksburg, Virginia, USA 24060

Author Contributions All authors contributed to all aspects of the manuscript.

Acknowledgements Funding was provided by grants from the Royal Swedish Academy of Sciences and Swedish Research Council to S. De Lisle (VR registration number 2019-03706), the University of Connecticut and the NIAID (1R01AI123659-01A1) to D. Bolnick, and the National Science Foundation (DEB 1457463) to J. McGlothlin.

Data Accessibility No data to be archived

1 **Interacting phenotypes and the coevolutionary process: Interspecific**
2 **indirect genetic effects alter coevolutionary dynamics**
3

Abstract Coevolution occurs when species interact to influence one another's fitness,

resulting in reciprocal evolutionary change. In many coevolving lineages, trait expression in

one species is modified by the genotypes and phenotypes of the other, forming feedback

loops reminiscent of models of intraspecific social evolution. Here, we adapt the theory of

within-species social evolution, characterized by indirect genetic effects and social selection

imposed by interacting individuals, to the case of interspecific interactions. In a trait-based

model, we derive general expressions for multivariate evolutionary change in two species and

the expected between-species covariance in evolutionary change when selection varies across

space. We show that reciprocal interspecific indirect genetic effects can dominate the

coevolutionary process and drive patterns of correlated evolution beyond what is expected

from direct selection alone. In extreme cases, interspecific indirect genetic effects can lead to

coevolution when selection does not covary between species or even when one species lacks

genetic variance. Moreover, our model indicates that interspecific indirect genetic effects

may interact in complex ways with cross-species selection to determine the course of

coevolution. Importantly, our model makes empirically testable predictions for how different

forms of reciprocal interactions contribute to the coevolutionary process.

Key words: Coevolution, cross-species selection, interspecific indirect genetic effects,

quantitative genetics, species interactions

Introduction

Coevolution occurs when interacting lineages evolve reciprocally in response to one another

(Janzen 1980, Thompson 1982). Although the concept of coevolution may be applied to

lineages that share genes, such as males and females of the same species (Arnqvist and Rowe

2002), it was originally invoked to explain patterns of correlated evolution between

interacting species (Ehrlich and Raven 1964). In some cases, coevolution can result in tightly

integrated mutualisms or spectacular arms races that drive the evolution of exceptional

phenotypes (Brodie et al. 2002, Pellmyr 2003, Johnson and Anderson 2010). Yet even

beyond these striking cases, coevolution is likely important for a wide range of interacting

lineages, including consumers and their resources, hosts and their pathogens, competitors,

and mutualists (Thompson 1982, 1994). Although coevolution has clearly played a major

role in the origins of diversity, much is still unknown about when and how species

interactions generate reciprocal evolutionary change.

Theoretical models of coevolution typically focus on the fitness effects of trait

interactions between coevolving species and how selection imposed by one species manifests

evolutionary change in an interacting species (Nuismer 2017). The interaction between

species in coevolving lineages is often intimate, with one species spending a greater part of

its life cycle in close contact with the other. Thus, coevolution bears a striking resemblance

to intraspecific social evolution (Stearns 2012). Like social evolution, coevolution often

includes interacting or extended phenotypes, which arise when traits can only be understood

within the context of interactions with others (Dawkins 1982, Moore et al. 1997).

In quantitative genetic models of intraspecific social evolution, social interactants

influence one another via two pathways, each of which has a counterpart in coevolutionary

theory. First, the phenotype of one individual may cause fitness effects in a social partner,

leading to a form of selection known as social selection (West-Eberhard 1979, West-

Eberhard 1983, West-Eberhard 1984, Wolf et al. 1999). At the heart of all coevolutionary

models is a form of reciprocal fitness interaction that resembles social selection, where the

fitness of individuals in one species is influenced by traits in an interacting species (Brodie

and Ridenhour 2003, Ridenhour 2005, Nuismer 2017). Second, models of social evolution

may also include indirect genetic effects, which occur when the phenotype of one individual

depends on the genotype of an interacting partner (Moore et al. 1997, Wolf et al. 1998). A

classic example of indirect genetic effects is maternal effects, in which offspring phenotype is

a function of both their own genes (a direct genetic effect) and maternal phenotypes such as

litter size and provisioning (an indirect genetic effect) (Kirkpatrick and Lande 1989,

Mousseau and Fox 1998, McAdam et al. 2002). When indirect genetic effects are reciprocal,

feedback effects may inflate the genetic variance available for response to selection,

drastically accelerating the rate of evolution (Moore et al. 1997). Although most models of

indirect genetic effects do not extend beyond species boundaries, indirect genetic effects may

also be common in species interactions between mutualists, competitors, parasites and hosts,

and predators and prey. Phenotypic plasticity in response to an interacting species is

common (Agrawal 2001), and when these influences on trait expression have a genetic basis

they may represent interspecific indirect genetic effects (IIGEs; Shuster et al. 2006).

Although IIGEs have received some attention in the context of community genetics (Shuster

et al. 2006, Witham et al. 2020), their potential role in driving trait coevolution has been

mostly unexplored.

To date, most explorations of IIGEs have been studies providing empirical support for

their likely existence and their contribution to trait variation. Examples of interspecific

phenotypic manipulation are common in nature (Table 1), and many of these cases can be

argued to be putative cases of IIGEs. One possible example occurs in arbuscular

mycorrhizae, where the genotypes of fungal mutualists can alter root traits in the plants they

Table 1. Coevolutionary interactions where IIGEs may be prevalent.

Interaction	IIGE synonym	Diffuse/individual interaction	Example References
host-parasite	host manipulation	both	(Thomas et al. 2012)
predator-prey	trait-mediated indirect interaction	both	(Peacor and Werner 2001)
plant-animal	inducible direct defense host-plant manipulations	both	(Weis and Abrahamson 1986, Chen 2008)
plant-microbe	joint trait microbially mediated trait	diffuse	(Friesen et al. 2011, O'Brien et al. 2021)
host-microbiome	host control host-microbiome interaction	diffuse	(Stappenbeck and Virgin 2016, Foster et al. 2017)
consumer-resource	toxin sequestration	both	(Züst et al. 2018)

inhabit (Gianinazzi-Pearson et al. 2007). In host-parasite systems, parasite manipulation of

host traits (such as behavior) and reciprocal host manipulation of parasite traits (such as

growth rate, via immune response) are key features of species interactions (Thomas et al.

2012). Importantly, in many host-parasite systems, both the host and the parasite experience

sustained interactions with a small number of individuals of the other species, often over key

periods of the life history. For example, helminth parasites excrete a variety of

immunomodulatory products that suppress or misdirect the immune system of their

individual host (Damian 1997, Schmid-Hempel 2008, Oladiran and Belosevic 2012). Thus,

host immune response to infection is controlled by the genotype of both the host and parasite

(e.g., Barribeau 2014). As a specific example, some populations of threespine stickleback

(*Gasterosteus aculeatus*) initiate a strong immune response to infection by the cestode

Schistocephalus solidus (Fig. 1A) involving granulocyte proliferation and fibrosis, which

effectively suppress cestode growth and viability (Weber et al. 2017). Other stickleback

populations do not exhibit this response and allow rapid cestode growth, perhaps representing

a tolerance strategy. Cestode growth is thus an indirect genetic effect of its host's genotype.

Conversely, the cestode has been shown to secrete compounds that suppress this host

response (Scharsack et al. 2004, 2007, 2013) and down-regulate sticklebacks' pro-fibrotic

gene expression (Fuess et al. 2020), suggesting reciprocal indirect genetic effects.

In goldenrod (*Solidago*; Fig. 1B), size of galls produced by the gall fly *Eurosta* is

determined by genotypes of both fly and plant, and evolution of gall size is influenced in part

by cross-species selection imposed on *Eurosta* larvae by species at other trophic levels (Weis

and Abrahamson 1986, Weis et al. 1992, Abrahamson and Weis 1997). This type of

interaction is common across gall-forming insects and their plant hosts; in *Hormaphi*s aphids,

variation the *bicycle* gene has been linked to variation in gall size (Korgaonkar et al. 2021).

In many herbivore-plant interactions, physical damage to leaves induces upregulation of

defensive compounds to deter further herbivory, which can be countered by matching

physiological changes in the herbivore (Ohgushi 2005). For example, *Littorina* snail

herbivory changes foliar chemistry of the brown seaweed *Ascophyllum nodosa* (increased

phlorotannin concentrations), which in turn reduces snail movement and consumption rates

(Borell et al. 2004).

Host-parasite, host-parasitoid, and some plant-herbivore interactions can entail

intimate long-term associations between individuals. In contrast, predator-prey interactions

tend to be more diffuse (Brodie and Brodie 1999). Prey sense predation risk through

chemical, auditory, or visual cues and change their morphology, physiology, or behavior in

Figure 1. Examples of traits potentially mediated by IIGEs. IIGEs are a common feature of species interactions, particularly in host-parasite systems where prolonged contact between single individuals of both species determines fitness and trait expression for both. Here we highlight four examples where such IIGEs are likely contributing to trait expression and possibly patterns of among-population (co)variation. Panel A shows threespine stickleback fish (*Gasterosteus aculeatus*) and their cestode parasite *Schistocephalus*. Freshwater stickleback are the intermediate host for *Schistocephalus*, and each cestode acquires most of its lifetime resource pool while living inside a single host fish. *Schistocephalus* impose substantial reproductive and survival costs on hosts, and hosts have evolved an inducible (by the cestode) defense that suppresses cestode growth. Substantial among population-variation in this trait and in infection rates suggest these IIGEs may be mediating coevolution. Panel B shows goldenrod (*Solidago*) and a gall induced by the larvae (visible inside the gall) of the specialist gall-forming fly *Eurosta*. Gall size is induced by the genotype of the *Eurosta* larvae, and past work has shown that gall expression is a complex interaction between plant and fly genotype. Panel C shows the fungus *Ophiocordyceps unilateralis*, which manipulates behavior of its ant host prior to emergence of its fruiting body (Anderson et al. 2009). Panel D shows the water flea *Daphnia lumholtzi*, which induces growth of protective spines in response to chemical cues released by predatory fish (Agrawal 2001). Panel A and B main photos: S. De Lisle. Inset (B) photo: SriMesh / CC BY-SA (https://creativecommons.org/licenses/by-sa/3.0); C photo: "File:Ophiocordyceps unilateralis.png" by David P. Hughes, Maj-Britt Pontoppidan / CC BY 2.5; Panel D: "Water flea (Daphnia lumholtzi)" by Frupus / CC BY-NC 2.0.

ways that mitigate their risk of predator encounter (Werner and Peacor 2003, Preisser et al.

2005). Perhaps the most prominent example is the tendency of some *Daphnia* genotypes to

grow spines (Fig. 1D), or to migrate to other depths, when they detect scent cues

(kairomones) from predatory fish (Weber and Declerck 1997, Boersma et al. 1998). These

antipredator responses lead to systemic changes in gene expression and morphology (Tams et

al. 2019), which are controlled in part by fish traits (e.g., production of a scent cue). These

scents can themselves be variable and genetic, as illustrated by differences in how *Daphnia*

respond to cues from landlocked versus anadromous alewife (Walsh and Post 2011). The

Daphnia traits can thus be described resulting from IIGEs controlled by both *Daphnia* and

fish genotypes. Unlike the intimate host-parasite interactions, however, prey may be

responding to diffuse cues from a predator population as a whole.

Although IIGEs would appear to play a central role in trait interactions between many

coevolving species, little is known about how these effects influence the dynamics of

coevolution (Scheiner et al. 2015). Past approaches, which have included variance-

partitioning models of community assembly (Shuster et al. 2006, Whitham et al. 2020) and

models of "joint traits" expressed together by interacting species (Queller 2014, O'Brien et

al. 2021), are suggestive of an important role for IIGEs in species interactions. However, we

lack a general understanding of how reciprocal IIGEs may affect the coevolution of

interacting phenotypes. The fact that indirect genetic effects within a species can create

feedback loops and other complex evolutionary dynamics, including in the context of within-

species coevolution (Drown and Wade 2014), suggests that IIGEs could play a major role in

mediating trait coevolution between species. Coopting concepts from intraspecific social

evolution theory, where trait-based IGEs are well developed, thus provides a natural way to

understand the trait interactions that drive coevolution. Importantly, this trait-based approach

allows the contribution of indirect genetic effects to the coevolutionary process to be fully

explored. Our goal in this paper is to develop such models to provide a comprehensive

theoretical assessment of how IIGEs contribute to coevolution between species.

Here we adapt the trait-based theory of intraspecific social evolution (Moore et al.

1997, Wolf et al. 1999, McGlothlin et al. 2010) to the case of two interacting species. Our

model applies to both pairwise interactions between individuals (e.g., stickleback and

cestodes) as well as diffuse interactions between species mean values (e.g., predatory fish and

Daphnia). Importantly, our model accommodates both IIGEs, where genes in individuals of

one species influence trait expression in individuals of another species, and cross-species

selection, where phenotypes of individuals in one species influence fitness of individuals of

another species (Figs. 1, 2). In addition to describing the contribution of these interspecific

interactions to evolutionary change, we develop expressions for the among-population

evolutionary covariance between traits of two interacting species; that is, the expected

covariance in trait means across populations between two coevolving species. In the process,

we formalize an interspecific analog of Zeng's (1988) quantitative genetic model of among-

population trait covariation, which we expand to incorporate interacting phenotypes. Our

analysis shows that IIGEs may have a central role in driving and mediating coevolution.

Reciprocal evolutionary change in interacting species

To model coevolution in two interacting species, we first decompose trait expression into

three components: direct genetic effects, environmental effects, and indirect effects mediated

by the phenotype of an interacting species. The phenotypic interface of coevolution involves

traits with interacting effects across individuals of two species *x* and *y* and can be written as

$$\mathbf{z}_x = \mathbf{a}_x + \mathbf{e}_x + \mathbf{\Phi}_{xy}\mathbf{z}_y' \tag{1a}$$

$$\mathbf{z}_y = \mathbf{a}_y + \mathbf{e}_y + \mathbf{\Phi}_{yx}\mathbf{z}_x', \tag{1b}$$

where $\mathbf{z}_i$ is a column vector of *m* traits expressed in an individual of species $i = x$ or y (*m* is

not necessarily equal in each species), $\mathbf{a}_i$ is the corresponding column vector of direct genetic

effects, and $\mathbf{e}_i$ is an uncorrelated vector of residual environmental effects; primes denote traits

of interacting individuals of another species. The matrix $\mathbf{\Phi}_{xy}$ quantifies the effect of traits in

an interacting individual of species y ($\mathbf{z}'_y$) on the expression of traits in a focal individual of

species *x* (Fig. 2, Table 2), while the matrix $\mathbf{\Phi}_{yx}$ quantitifies such effects in the opposite

direction. Thus, $\mathbf{\Phi}_{ij}$ is an interspecific analog of the matrix of conspecific indirect genetic

effects, $\mathbf{\Psi}$ (Moore et al. 1997). The individual elements of $\mathbf{\Phi}_{ij}$, which we write as ϕ_{ij}^{kl},

represent partial regression coefficients of trait *l* in species *j* on trait *k* in species *i*. When

traits are standardized to the same scale, these coefficients will typically be limited to a range

of -1 to 1. Although we focus on interactions between pairs of individuals for simplicity of

notation, this model could be easily extended to describe interactions with multiple

individuals (cf. McGlothlin et al. 2010). For more diffuse interactions (e.g., alewife and

Daphnia), the elements of $\mathbf{\Phi}_{ij}$ could represent a weighted average of effects from integrating

across the phenotype distribution of the interacting population.

Figure 2. Path model of selection when traits and fitness are determined by interactions with heterospecifics. Individual trait values of two interacting species, *x* (blue) and *y* (red), are represented by *z*. Traits directly influence individual relative fitness (*w*) of the species that express them via natural selection (β_{xx} and β_{yy}, black arrows). Traits can also reciprocally influence expression of traits of heterospecific social partners, via interspecific indirect genetic effects Φ. Traits can also influence fitness of heterospecific social partners, via cross-species selection (β_{xy} and β_{yx}, grey arrows). Social effects are illustrated in grey arrows, direct effects in black.

Table 2. Definition of key parameters and expressions.

Expression / parameter	Biological definition
$\mathbf{z}_x, \mathbf{z}_y$	Vectors of individual trait values for species x and y, with a prime when present denoting partner traits.
$\mathbf{a}_x, \mathbf{a}_y$	Vectors of individual breeding values for species x and y.
$\mathbf{\Phi}_{xy}, \mathbf{\Phi}_{yx}$	Interspecific indirect effects (in matrix form). The effect of traits in individuals of species y on trait expression in species x, and vice-versa.
$\mathbf{\beta}_{xx}, \mathbf{\beta}_{yy}$	Vectors of within-species natural selection gradients for each species.
$\mathbf{\beta}_{xy}, \mathbf{\beta}_{yx}$	Vectors of cross-species selection gradients for each species. The effects of traits in species y on fitness of individual species x, and vice-versa.
$\mathbf{G}_{xx}, \mathbf{G}_{yy}$	Genetic covariance matrices of traits for species x and y.
$\mathbf{G}_{xy}, \mathbf{G}_{yx}$	Covariance matrices for breeding values of interacting individuals of species x and y within a population.
$\mathrm{Cov}(\Delta\bar{z}_x, \Delta\bar{z}_y)$	Coevolutionary covariance, defined as the covariance in evolutionary change between interacting species, across populations. When standardized by the variances in evolutionary response for each species, becomes a scale-free correlation.
$\mathrm{Cov}(\beta_{xx}, \beta_{yy})$,	Between-species, among-population covariance in selection gradients (shown here for within-species selection)

172   Assuming that the mean residual environmental effect is zero, the population mean

173   phenotype vector for each species is

$$\bar{\mathbf{z}}_x = \bar{\mathbf{a}}_x + \mathbf{\Phi}_{xy}\bar{\mathbf{z}}_y \tag{2a}$$

$$\bar{\mathbf{z}}_y = \bar{\mathbf{a}}_y + \mathbf{\Phi}_{yx}\bar{\mathbf{z}}_x. \tag{2b}$$

176   Noting the change in the mean additive genetic value in each species is $\Delta\bar{\mathbf{a}} = \text{Cov}(\mathbf{a}, w)$

177   (Robertson 1966, Price 1970, 1972) where w is relative fitness, we can now define

178   evolutionary change in the multivariate mean phenotype as

$$\Delta\bar{\mathbf{z}}_x = \text{Cov}(\mathbf{a}_x, w_x) + \mathbf{\Phi}_{xy}\Delta\bar{\mathbf{z}}_y \tag{3a}$$

$$\Delta\bar{\mathbf{z}}_y = \text{Cov}(\mathbf{a}_y, w_y) + \mathbf{\Phi}_{yx}\Delta\bar{\mathbf{z}}_x. \tag{3b}$$

In equation (3), the first term on each right-hand side describes direct change due to natural

selection on the focal species, and the second term describes indirect change due to the

product of IIGEs and phenotypic change in the interacting species. This second term is

analogous to transmission bias (Fisher and McAdam 2019), and in this case, specifically

describes changes in one species that are induced by evolution in an interacting species. It

can be seen from equation (3) that species will coevolve whenever there are IIGEs in both

species, because the change in phenotypic mean in species x depends upon the change of

mean in species y and vice versa whenever both $\mathbf{\Phi}_{ij} \neq \mathbf{0}$. Expanding equation (3) by

substitution results in explicit equations for evolutionary change in each species,

$$\Delta\bar{\mathbf{z}}_x = \left(\mathbf{I}_x - \mathbf{\Phi}_{xy}\mathbf{\Phi}_{yx}\right)^{-1}\left[\text{Cov}(\mathbf{a}_x, w_x) + \mathbf{\Phi}_{xy}\text{Cov}(\mathbf{a}_y, w_y)\right] \tag{4a}$$

$$\Delta \bar{\mathbf{z}}_y = \left(\mathbf{I}_y - \boldsymbol{\Phi}_{yx}\boldsymbol{\Phi}_{xy}\right)^{-1}[\text{Cov}(\mathbf{a}_y, w_y) + \boldsymbol{\Phi}_{yx}\text{Cov}(\mathbf{a}_x, w_x)]. \tag{4b}$$

where $\mathbf{I}_x$ and $\mathbf{I}_y$ are identity matrices with dimensionality equal to the number of traits in the

two species and the multiplier $\left(\mathbf{I} - \boldsymbol{\Phi}_{ij}\boldsymbol{\Phi}_{ji}\right)^{-1}$ quantifies the feedback effect of reciprocal

IIGEs. Equation 4 illustrates that total evolutionary change in a species is determined both

by change in the breeding value of that species ($\text{Cov}(\mathbf{a}_i, w_i)$) and by change in the breeding

value of the interacting species, mediated by IIGEs ($\boldsymbol{\Phi}_{ij}\text{Cov}(\mathbf{a}_j, w_j)$). Whenever IIGEs occur

in both species such that $\boldsymbol{\Phi}_{ij}\boldsymbol{\Phi}_{ji} \neq \mathbf{0}$, this multiplier alters the total amount of evolutionary

change in both species. In order for such an effect to arise, there must be a feedback loop in

phenotypic expression. The simplest of these arises when there are two traits with reciprocal

IIGEs, such that trait k in species i affects trait l in species j and trait l in turn influences the

expression of trait k. In general, when ϕ_{ij}^{kl} and ϕ_{ji}^{lk} are of the same sign, the magnitude of

evolutionary change will be enhanced, and when they are of opposite signs, evolutionary

change will be diminished (Fig. 3).

To expand equation (4), we define the fitness of each interacting species using the

linear equations

$$w_x = \alpha_x + \mathbf{z}_x^{\text{T}}\boldsymbol{\beta}_{xx} + \mathbf{z}_y'^{\text{T}}\boldsymbol{\beta}_{xy} + \varepsilon_x \tag{5a}$$

$$w_y = \alpha_y + \mathbf{z}_y^{\text{T}}\boldsymbol{\beta}_{yy} + \mathbf{z}_x'^{\text{T}}\boldsymbol{\beta}_{yx} + \varepsilon_y \tag{5b}$$

where w is individual relative fitness, α is an intercept, ε is an error term, and the superscript

T denotes transposition. Directional selection in each species is partitioned into two selection

gradients. First, the within-species selection gradients ($\boldsymbol{\beta}_{xx}$ and $\boldsymbol{\beta}_{yy}$) describe the direct

effects of an individual's traits on its own fitness (Fig. 2; Lande and Arnold 1983). The

cross-species selection gradients ($\boldsymbol{\beta}_{xy}$ and $\boldsymbol{\beta}_{yx}$) relate the fitness of a focal individual of one

species (*x* or *y*) to the traits of individuals of the other coevolving species (*y* or *x*) (Fig. 2).

Both of these are linear components of selection; we consider nonlinear terms (interactions

for fitness between focal and partner traits) below (see Incorporating Specific Fitness

Models). The cross-species selection gradient is analogous to the directional social selection

gradient in within-species models (Wolf et al. 1999) and if pairs or groups of interacting

individuals can be identified in a natural population, can be estimated in a similar way (cf.

Ridenhour 2005).

We now substitute our definitions of fitness into equation (4) and expand, yielding

predictive equations for evolutionary change in each species:

$$\Delta\bar{\mathbf{z}}_x = \left(\mathbf{I}_x - \boldsymbol{\Phi}_{xy}\boldsymbol{\Phi}_{yx}\right)^{-1}[\text{Cov}(\mathbf{a}_x, \mathbf{z}_x^{\text{T}})\boldsymbol{\beta}_{xx} + \text{Cov}(\mathbf{a}_x, \mathbf{z}_y'^{\text{T}})\boldsymbol{\beta}_{xy} +$$
$$\boldsymbol{\Phi}_{xy}\text{Cov}(\mathbf{a}_y, \mathbf{z}_y^{\text{T}})\boldsymbol{\beta}_{yy} + \boldsymbol{\Phi}_{xy}\text{Cov}(\mathbf{a}_y, \mathbf{z}_x'^{\text{T}})\boldsymbol{\beta}_{yx}] \tag{6a}$$

$$\Delta\bar{\mathbf{z}}_y = \left(\mathbf{I}_y - \boldsymbol{\Phi}_{yx}\boldsymbol{\Phi}_{xy}\right)^{-1}[\text{Cov}(\mathbf{a}_y, \mathbf{z}_y^{\text{T}})\boldsymbol{\beta}_{yy} + \text{Cov}(\mathbf{a}_x, \mathbf{z}_x^{\text{T}})\boldsymbol{\beta}_{yx} +$$
$$\boldsymbol{\Phi}_{yx}\text{Cov}(\mathbf{a}_x, \mathbf{z}_x^{\text{T}})\boldsymbol{\beta}_{xx} + \boldsymbol{\Phi}_{yx}\text{Cov}(\mathbf{a}_x, \mathbf{z}_y'^{\text{T}})\boldsymbol{\beta}_{xy}]. \tag{6b}$$

Each of these equations consists of four terms representing four components of the total

response to selection. The first term ($\text{Cov}(\mathbf{a}_x, \mathbf{z}_x^{\text{T}})\boldsymbol{\beta}_{xx}$) represents response to within-species

selection ($\boldsymbol{\beta}_{xx}$ or $\boldsymbol{\beta}_{yy}$), the second ($\text{Cov}(\mathbf{a}_x, \mathbf{z}_y'^{\text{T}})\boldsymbol{\beta}_{xy}$) represents response to cross-species

linear selection ($\boldsymbol{\beta}_{xy}$ or $\boldsymbol{\beta}_{yx}$), and the last two terms ($\boldsymbol{\Phi}_{xy}\text{Cov}(\mathbf{a}_y, \mathbf{z}_y^{\text{T}})\boldsymbol{\beta}_{yy} +$

$\boldsymbol{\Phi}_{xy}\text{Cov}(\mathbf{a}_y, \mathbf{z}_x'^{\text{T}})\boldsymbol{\beta}_{yx}$) represent the component of change caused by the change in mean of

the interacting species. Note that the third and fourth terms of the first part of equation (6a)

equal the first two terms of (6b) multiplied by the IIGE coefficient $\boldsymbol{\Phi}_{xy}$. Equation (6) also

shows that the change in response to within-species selection depends on the covariance of

additive genetic value with the phenotype of the same species, while change in response to

cross-species selection depends on the covariance of additive genetic value with the

phenotype of the opposite species.

To determine the components that give rise to these covariances, we expand our

definition of the phenotypes in equation (2) and rearrange, yielding

$$\Delta\bar{\mathbf{z}}_x = \left(\mathbf{I}_x - \boldsymbol{\Phi}_{xy}\boldsymbol{\Phi}_{yx}\right)^{-1}\left[\left(\mathbf{G}_{xx} + \mathbf{G}_{xy}\boldsymbol{\Phi}_{xy}^{\mathrm{T}}\right)\left(\mathbf{I}_x - \boldsymbol{\Phi}_{yx}^{\mathrm{T}}\boldsymbol{\Phi}_{xy}^{\mathrm{T}}\right)^{-1}\boldsymbol{\beta}_{xx} + \right. \tag{7a}$$

$$\left(\mathbf{G}_{xy} + \mathbf{G}_{xx}\boldsymbol{\Phi}_{yx}^{\mathrm{T}}\right)\left(\mathbf{I}_y - \boldsymbol{\Phi}_{xy}^{\mathrm{T}}\boldsymbol{\Phi}_{yx}^{\mathrm{T}}\right)^{-1}\boldsymbol{\beta}_{xy} + $$

$$\boldsymbol{\Phi}_{xy}\left(\mathbf{G}_{yy} + \mathbf{G}_{yx}\boldsymbol{\Phi}_{yx}^{\mathrm{T}}\right)\left(\mathbf{I}_y - \boldsymbol{\Phi}_{xy}^{\mathrm{T}}\boldsymbol{\Phi}_{yx}^{\mathrm{T}}\right)^{-1}\boldsymbol{\beta}_{yy} + $$

$$\left. \boldsymbol{\Phi}_{xy}\left(\mathbf{G}_{yx} + \mathbf{G}_{yy}\boldsymbol{\Phi}_{xy}^{\mathrm{T}}\right)\left(\mathbf{I}_x - \boldsymbol{\Phi}_{yx}^{\mathrm{T}}\boldsymbol{\Phi}_{xy}^{\mathrm{T}}\right)^{-1}\boldsymbol{\beta}_{yx}\right]$$

$$\Delta\bar{\mathbf{z}}_y = \left(\mathbf{I}_y - \boldsymbol{\Phi}_{yx}\boldsymbol{\Phi}_{xy}\right)^{-1}\left[\left(\mathbf{G}_{yy} + \mathbf{G}_{yx}\boldsymbol{\Phi}_{yx}^{\mathrm{T}}\right)\left(\mathbf{I}_y - \boldsymbol{\Phi}_{xy}^{\mathrm{T}}\boldsymbol{\Phi}_{yx}^{\mathrm{T}}\right)^{-1}\boldsymbol{\beta}_{yy} + \right. \tag{7b}$$

$$\left(\mathbf{G}_{yx} + \mathbf{G}_{yy}\boldsymbol{\Phi}_{xy}^{\mathrm{T}}\right)\left(\mathbf{I}_x - \boldsymbol{\Phi}_{yx}^{\mathrm{T}}\boldsymbol{\Phi}_{xy}^{\mathrm{T}}\right)^{-1}\boldsymbol{\beta}_{yx} + $$

$$\boldsymbol{\Phi}_{yx}\left(\mathbf{G}_{xx} + \mathbf{G}_{xy}\boldsymbol{\Phi}_{xy}^{\mathrm{T}}\right)\left(\mathbf{I}_x - \boldsymbol{\Phi}_{yx}^{\mathrm{T}}\boldsymbol{\Phi}_{xy}^{\mathrm{T}}\right)^{-1}\boldsymbol{\beta}_{xx} + $$

$$\left. \boldsymbol{\Phi}_{yx}\left(\mathbf{G}_{xy} + \mathbf{G}_{xx}\boldsymbol{\Phi}_{yx}^{\mathrm{T}}\right)\left(\mathbf{I}_y - \boldsymbol{\Phi}_{xy}^{\mathrm{T}}\boldsymbol{\Phi}_{yx}^{\mathrm{T}}\right)^{-1}\boldsymbol{\beta}_{xy}\right],$$

where $\mathbf{G}_{xx}$ and $\mathbf{G}_{yy}$ represent within-species genetic (co)variance matrices and $\mathbf{G}_{xy}$ and $\mathbf{G}_{yx}$

represent cross-species genetic covariance. Note that in a model of pairwise interactions,

$\mathbf{G}_{yx} = \mathbf{G}_{xy}^{\mathrm{T}}$.

The cross-species genetic covariance represents a covariance of breeding values

between the species within the population of interest. This covariance may arise via any

mechanism that leads to nonrandom genetic assortment between species x and y with respect

to traits at the phenotypic interface of coevolution. A variety of phenomena could lead to

such assortment including behavioral preference for certain traits in heterospecific partners,

fine-scale population structure, habitat preference, and vertical transmission of symbionts.

We elaborate on the contribution of $\mathbf{G}_{xy}$ and $\boldsymbol{\Phi}$ to phenotypic assortment between individuals

of interacting species (C_{xy}) in equation (A1). However, there is currently limited evidence of

this type of interspecific genetic assortment, as few investigators seemed to have attempted to

measure such a covariance. We therefore investigate dynamics in both the presence and

absence of such assortment in our model development below.

Response to within-species selection depends on the sum of within-species genetic

variance, $\mathbf{G}_{yy}$ or $\mathbf{G}_{xx}$, and $\mathbf{G}_{xy}\boldsymbol{\Phi}_{xy}^{\mathrm{T}}$ or $\mathbf{G}_{yx}\boldsymbol{\Phi}_{yx}^{\mathrm{T}}$, which represent an interaction between IIGEs

and cross-species genetic covariance. Response to cross-species selection depends on the

sum of cross-species genetic variance and $\mathbf{G}_{xx}\boldsymbol{\Phi}_{yx}^{\mathrm{T}}$ or $\mathbf{G}_{yy}\boldsymbol{\Phi}_{xy}^{\mathrm{T}}$, which describes the genetic

variance created by IIGEs. Each term in equation (7) also contains an additional feedback

multiplier, $\left(\mathbf{I}_y - \boldsymbol{\Phi}_{xy}^{\mathrm{T}}\boldsymbol{\Phi}_{yx}^{\mathrm{T}}\right)^{-1}$, which further enhances the response to selection when there

are reciprocal IIGEs. Because of this additional multiplier, when IIGEs are reciprocal and of

the same sign, their effects on can be massive, mirroring the effects observed for within-

species IGEs (Moore et al. 1997, McGlothlin et al. 2010). As in previous equations, the last

two terms in equation (7) represent a sort of evolutionary feedback that occurs across

generations and is only present when there are IIGEs. These effects of $\boldsymbol{\Phi}$ and cross-species

selection on evolutionary response are illustrated in Fig. 3.

Figure 3. Reciprocal IIGEs and cross-species selection change evolutionary response in a single species. Panels show the separate effects on evolutionary response in species x of indirect genetic effects (Panel A) and cross-species selection with genetic assortment (Panel B). Panel A shows the effects of reciprocal IIGEs holding all other evolutionary parameters constant, and assuming no cross-species selection. Panel B shows the effects of cross-species selection imposed by species y on species x, in combination with the genetic assortment between interactants of the different species, and assuming IIGEs are absent. In both panels, $G_x = G_y = 1$, $\beta_{yy} = \beta_{xx} = 1$.

Correlated evolution between interacting species: the coevolutionary covariance

A key feature of coevolving lineages is correlated evolution across populations subject to

varying ecological conditions. Here, we seek to understand how the equations of selection

response can be used to understand how selection and IIGEs contribute to this shared among-

population divergence. We can explore the contribution of interspecific social effects to

correlated evolution between interacting species by solving for the covariance in evolutionary

change between interacting species, $\mathrm{Cov}(\Delta\bar{z}_x, \Delta\bar{z}_y)$, which we call the coevolutionary

covariance (with the caveat that this covariance also reflects changes due in part to effects of

IIGEs). Such a covariance represents the expected pattern of reciprocal phenotypic change

through time in a single pair of populations of two species experiencing fluctuating selection,

or perhaps more importantly, across a set of populations in space under varying selection

pressures. Although mathematically equivalent, we focus on the latter scenario for its

relevance to understanding geographic variation among populations.

The coevolutionary covariance reflects the degree to which divergence among

population means in two species has occurred jointly. High absolute values of the

coevolutionary covariance indicate tightly-coupled coevolutionary change between the two

species, whereas values around zero indicate that evolutionary change occurs independently.

When scaled by the total amount of population divergence (variances in $\Delta\bar{z}_x$ and $\Delta\bar{z}_y$), this

becomes a scale-free correlation describing the proportion of total divergence shared between

the two species. Importantly, we expect the coevolutionary covariance to be often be related

to the covariance in population means, $\text{Cov}(\bar{z}_x, \bar{z}_y) \propto \text{Cov}(\Delta\bar{z}_x, \Delta\bar{z}_y)$ because, for example,

$\text{Cov}(\bar{z}_x, \bar{z}_y)^{t+1} = \text{Cov}(\bar{z}_x, \bar{z}_y)^t + \text{Cov}(\Delta\bar{z}_x, \Delta\bar{z}_y)$ (under the assumption $\text{Cov}(\Delta\bar{z}_x, \bar{z}_y) +$

$\text{Cov}(\bar{z}_x, \Delta\bar{z}_y) = 0$). Thus, expanding $\text{Cov}(\Delta\bar{z}_x, \Delta\bar{z}_y)$ allows the possibility to assess how

selection and IIGEs contribute to patterns of geographic variation in species mean

phenotypes.

To simplify our analysis, we use models of a single trait in each species. Developing

a full equation including all sources of covariance quickly becomes cumbersome, so we focus

on three instructive special cases that illustrate the explicit impacts of considering cross-

species selection and interspecific interacting phenotypes. The simplest case occurs when

there are no IIGEs ($\phi_{xy} = \phi_{yx} = 0$) and heterospecific interactions occur at random within

populations ($G_{xy} = G_{yx} = 0$). If we make the simplifying assumption that genetic variance

does not differ among populations, the only source of covariance in selection response is

covariance in direct within-species selection, or

$$\text{Cov}(\Delta\bar{z}_x, \Delta\bar{z}_y) = G_{xx}G_{yy}\text{Cov}(\beta_{xx}, \beta_{yy}). \tag{8}$$

Equation (8) represents what is usually thought of as one source of correlated evolution

between interacting species: predictable variation in the type of selection occurring across

space. Empirical data suggest that selection varies substantially in magnitude across space

(Siepielski et al. 2013) and that spatially autocorrelated biotic selection plays a substantial

role in driving divergence in trait means (Urban et al. 2011). Importantly, not just any

variance in selection will do. In order to create covariance in evolutionary change, selection

must covary between the interacting species. Cross-species selection, however, does not play

a role in equation (8) because it does not contribute to an evolutionary response to selection

in the absence of IIGEs and cross-species genetic covariance. Equation (8) is analogous to

the results of Zeng's (1988) model of correlated trait evolution under directional selection,

although for the case of traits expressed in different species.

To see an effect of cross-species selection, we first add cross-species genetic

covariance but no IIGEs. For a one-trait (per species) model, $G_{xy} = G_{yx}$. Adding this effect

leads to three new sources of covariance among populations:

$$\text{Cov}(\Delta \bar{z}_x, \Delta \bar{z}_y) = G_{xx} G_{yy} \text{Cov}(\beta_{xx}, \beta_{yy}) + G_{xy}^2 \text{Cov}(\beta_{yx}, \beta_{xy}) +$$
$$G_{xx} G_{xy} \text{Cov}(\beta_{xx}, \beta_{yx}) + G_{yy} G_{xy} \text{Cov}(\beta_{yy}, \beta_{xy}). \tag{9}$$

The second term in equation (9) $(G_{xy}^2 \text{Cov}(\beta_{yx}, \beta_{xy}))$ shows that cross-species selection will

contribute to the coevolutionary covariance only when there is cross-species genetic

covariance for the traits that affect fitness. Such covariances occur if, for example, parasite

and host genotypes associate nonrandomly for the traits that influence their partner's fitness.

The third and fourth terms $(G_{xx} G_{xy} \text{Cov}(\beta_{xx}, \beta_{yx}) + (G_{yy} G_{xy} \text{Cov}(\beta_{yy}, \beta_{xy}))$ represent a

relationship between the effect of species on itself and on its heterospecific partner. These

will be nonzero if populations with strong within-species selection also exhibit strong cross-

species selection.

The most interesting effects on the coevolutionary covariance occur when we add

IIGEs (ϕ_{xy} and ϕ_{yx}). With $G_{xy} = 0$, an appropriate simplifying assumption given that the

prevalence of G_{xy} is uncertain and is likely to be transient through time, the covariance

among populations in evolutionary response to selection is

$$
\begin{aligned}
\mathrm{Cov}(\Delta \bar{z}_x, \Delta \bar{z}_y) = \; &\mathcal{U}[G_{xx}G_{yy}\mathrm{Cov}(\beta_{xx},\beta_{yy}) + \\
&G_{xx}^2\phi_{yx}\mathrm{Var}(\beta_{xx}) + G_{yy}^2\phi_{xy}\mathrm{Var}(\beta_{yy}) + \\
&G_{xx}G_{yy}\phi_{xy}\phi_{yx}\mathrm{Cov}(\beta_{xx},\beta_{yy}) + G_{xx}G_{yy}\phi_{xy}\phi_{yx}\mathrm{Cov}(\beta_{xy},\beta_{yx}) + \\
&G_{xx}G_{yy}\phi_{xy}\mathrm{Cov}(\beta_{xx},\beta_{yx}) + G_{xx}G_{yy}\phi_{yx}\mathrm{Cov}(\beta_{yy},\beta_{xy}) + \\
&2G_{xx}^2\phi_{yx}^2\mathrm{Cov}(\beta_{xx},\beta_{xy}) + 2G_{yy}^2\phi_{xy}^2\mathrm{Cov}(\beta_{yy},\beta_{yx}) + \mathcal{V}],
\end{aligned}
\tag{10}
$$

340   where feedback effects of IIGEs are described by $\mathcal{U} = (1 - \phi_{xy}\phi_{yx})^{-4}$ and $\mathcal{V}$ collects

341   negligible third- and fourth-order ϕ terms:

$$
\begin{aligned}
\mathcal{V} = \; &G_{xx}^2\phi_{yx}^3\mathrm{Var}(\beta_{xy}) + G_{yy}^2\phi_{xy}^3\mathrm{Var}(\beta_{yx}) + \\
&G_{xx}G_{yy}\phi_{xy}\phi_{yx}^2\mathrm{Cov}(\beta_{yy},\beta_{xy}) + G_{xx}G_{yy}\phi_{xy}^2\phi_{yx}\mathrm{Cov}(\beta_{yx},\beta_{xx}) + \\
&G_{xx}G_{yy}\phi_{xy}^2\phi_{yx}^2\mathrm{Cov}(\beta_{yx},\beta_{xy}).
\end{aligned}
\tag{11}
$$

344   The first term in equation (10) is identical to equation (8) and represents covariance in direct

345   within-species selection. The second ($G_{xx}^2\phi_{yx}\mathrm{Var}(\beta_{xx})$) and third ($G_{yy}^2\phi_{xy}\mathrm{Var}(\beta_{yy})$) terms

Figure 4. IIGEs drive and mediate coevolution between interacting species. Panels show the coevolutionary correlation between two interacting species as a function of the indirect genetic effect parameters ϕ, from equation (10) standardized by the evolutionary rates. In the absence of social interaction effects, correlated evolution (coevolution) between species is driven entirely by covariance in natural selection between the species (see text). Reciprocal IIGEs can generate coevolution even when there is not covariance in natural selection (Panel A), and can even drive coevolution even when one species lacks genetic variance (Panel B). IIGEs modify observed coevolutionary patterns when natural selection does covary ($\mathrm{Cov}(\beta_{xx},\beta_{yy}) = 0.5$; Panel C). When cross-species selection and IIGEs act together, coevolutionary patterns are a complex third order polynomial (Panel D; $\mathrm{Cov}(\beta_{xy},\beta_{yx}) = 0.5$, $\mathrm{Var}(\beta_{xy}) = \mathrm{Var}(\beta_{yx}) = 1$). When natural and cross-species selection both act ($\mathrm{Var}(\beta_{xy}) = \mathrm{Var}(\beta_{yx}) = \mathrm{Var}(\beta_{yy}) = \mathrm{Var}(\beta_{xx}) = 1$) and covary positively (0.5), (Panel E), effects of IIGEs become stronger in comparison to the case of (co)variance in natural selection alone. When covariance between natural and cross-species selection is negative (-0.8) a ridge is observed (Panel F).

show that in the presence of IIGEs, simple variance in within-species selection across

populations can generate a coevolutionary covariance (or correlation, Fig. 4A; corresponding

evolutionary rates and covariances are plotted in Figs. S1 and S2, respectively). This

covariance in evolutionary response occurs as a necessary consequence of the dependence of

the trait mean of one species on that of the other. Thus, in the presence of IIGEs, a

coevolutionary covariance may occur even when selection is uncorrelated between the

species, and in extreme cases, when only one species varies in selection or when only one

species has genetic variation (Fig. 4B).

The fourth and fifth terms $(G_{xx}G_{yy}\phi_{xy}\phi_{yx}\text{Cov}(\beta_{xx},\beta_{yy}) +$

$(G_{xx}G_{yy}\phi_{xy}\phi_{yx}\text{Cov}(\beta_{xy},\beta_{yx}))$ show the effects of reciprocal IIGEs. When traits in the two

species exist in a feedback loop, the effect of covariance in within-species selection is

amplified if ϕ_{xy} and ϕ_{yx} are of the same sign and diminished if they are of opposite signs.

The combined effects of the second, third, and fourth terms lead to a complex relationship

between IIGEs and the coevolutionary covariance when within-species selection varies (Fig.

4C). In some cases, IIGEs can even reverse the sign of the correlation that would be

expected in their absence (Fig. 4C). In addition, reciprocal IIGEs may cause covariance in

cross-species selection to contribute to the coevolutionary covariance, causing an even more

complex relationship (Fig. 4D).

The last four terms in equation (10) show that in the presence of IIGEs, covariance

between within-species and cross-species selection may contribute to the coevolutionary

covariance (Fig. 4E-F). Because IIGEs inherently tie together the evolutionary responses of

the two species through their effects on the coevolutionary covariance, this can occur both

when there is covariance in gradients across species (terms 6 and 7) and when there is

covariance in gradients in the same species (terms 8 and 9). The total effect of IIGEs may be

quite complex when cross-species selection varies, with subtle changes in ϕ resulting in

dramatic changes in the expected among population correlation (Fig. 4D, E-F).

An equation incorporating both IIGEs and G_{xy} quickly becomes unwieldy, but we

present a compact form containing 10 (co)variance terms as equation (A2), which illustrates

that when present, non-random assortment and IIGEs together interact in complex ways to

influence coevolution.

378 Incorporating specific fitness models

Coevolutionary models often posit complex relationships between interacting phenotypes and

fitness (Nuismer 2017). Although the selection model we present here is linear, more

complex relationships can be incorporated by translating specific fitness functions to

selection gradients. The adaptive landscape represents the theoretical relationship between a

population's mean fitness and its phenotypic mean (Arnold et al. 2001). Selection gradients

represent the partial slope of the adaptive landscape with respect to a given phenotype, and

thus the multivariate selection gradient may be calculated using a vector of partial derivatives

if the adaptive landscape can be written as a differentiable function (Lande 1979, Lande and

Arnold 1983). In many cases, the multivariate selection gradient may be calculated using

partial derivatives of the individual fitness function as well (Lande and Arnold 1983, Abrams

et al. 1993, McGlothlin et al. 2021), evaluated over the phenotypic distribution (Phillips and

Arnold 1989). Once selection gradients have been calculated for a given model, they may be

substituted into equations (6–7) to explore the effects of a given fitness function on selection

response and the coevolutionary covariance (see also Brodie and Ridenhour 2003). This

exercise also allows us to explore the effects of adding IIGEs to an existing coevolutionary

model.

First, we consider a "phenotypic difference" model of ecological trait interaction

(Nuismer et al. 2007 , Nuismer 2017), where absolute fitness is a function of the difference

between interacting traits, $W_x \propto \exp(z_x - z_y)$ and $W_y \propto \exp(z_y - z_x)$. Using the

logarithm of the fitness function to calculate relative fitness (Lande and Arnold 1983),

$$w_x = \alpha_x + b_x(z_x - z_y) + \varepsilon_x$$
$$w_y = \alpha_y + b_y(z_y - z_x) + \varepsilon_y. \tag{12}$$

401   where b_x and b_y are constants that may vary across populations. Differentiating, the selection

402   gradients are then $\beta_{xx} = -\beta_{xy} = b_x$ and $\beta_{yy} = -\beta_{yx} = b_y$. Substituting these selection

403   gradients into equation (7) is trivial. However, is worth noting that under this fitness model,

404   the coevolutionary covariance simplifies to a function of just three (co)variance components,

$$\text{Cov}(\Delta\bar{z}_x, \Delta\bar{z}_y) = \mathcal{U}[G_{xx}G_{yy}(1 - \phi_{xy} - \phi_{yx} + 2\phi_{xy}\phi_{yx})\text{Cov}(b_x, b_y) +$$
$$G_{xx}^2\phi_{yx}(1 - 2\phi_{yx})\text{Var}(b_x) + G_{yy}^2\phi_{xy}(1 - 2\phi_{xy})\text{Var}(b_y) + \mathcal{V}], \tag{13}$$

which corresponds to a relationship with ϕ as in Figs. 4F, S1F, and S2F. This analysis

illustrates that reciprocal IIGEs have greatest impact on mediating coevolutionary outcomes

in a trait-matching models when IIGEs are similar in both sign and magnitude. Biologically,

such a situation corresponds to a scenario where, for example, trait expression is reciprocally

escalated in response to heterospecific partners.

Another important class of fitness models to consider is the case of nonlinear fitness

interactions. Nonlinearity is fundamental to many coevolutionary models that invoke

epistatic interactions across species' genomes, such as the trait matching model for

quantitative traits, or the single-locus matching-allele and gene-for-gene models of host-

parasite coevolution (Dybdahl et al. 2014). For example, in the traditional model of trait

matching, absolute fitness is assumed to be $W_x \propto \exp\left(-(z_x - z_y)^2\right)$ and so relative fitness

can be described by

$$w_x = \alpha_x - b_{xy}(z_x - z_y)^2 + \varepsilon_x$$
$$w_y = \alpha_y - b_{yx}(z_y - z_x)^2 + \varepsilon_y. \tag{14}$$

421 In this model the sign of the *b* terms correspond to different types of biological interaction;

422 for example, in a mutualism, both species may take positive values of *b* and thus have a

423 fitness peak when their trait value matches with their heterospecific partner. Taking partial

424 derivatives over the phenotypic distribution yields the selection gradients

$$\beta_{xx} = -\beta_{xy} = -2b_{xy}(\bar{z}_x - \bar{z}_y)$$
$$\beta_{yy} = -\beta_{yx} = -2b_{yx}(\bar{z}_y - \bar{z}_x) \tag{15}$$

427 In the simplest case of no IIGEs, $G_{xy} = 0$, and no variation in *b*, these gradients yield

$$\text{Cov}(\Delta\bar{z}_x, \Delta\bar{z}_y) = -4G_{xx}G_{yy}b_{xy}b_{yx}\text{Var}(\bar{z}_x - \bar{z}_y). \tag{16}$$

Thus, within the limits of genetic variation and ignoring stochastic forces (as we have done),

evolutionary response will be perfectly correlated in the phenotypic matching model. When

b_{xy} and b_{yx} share the same sign, this covariance in evolutionary response will be negative

because each species will be evolving towards (or away) from each other in trait space. This

covariance in evolutionary response occurs even without variation in b or covariance in trait

means. These results are broadly consistent with other analyses of the trait matching model,

which indicate that this model can generate relatively strong covariance in phenotypic means

across populations (Nuismer et al. 2010). Adding indirect genetic effects and assuming

$G_{xy} = 0$ and noting that $\text{Var}(\bar{z}_x - \bar{z}_y) = \text{Var}(\bar{z}_y - \bar{z}_x)$,

$$
\begin{aligned}
\text{Cov}(\Delta\bar{z}_x, \Delta\bar{z}_y) = \; & \mathcal{U}[-4b_{xy}b_{yx}G_{xx}G_{yy}\text{Var}(\bar{z}_x - \bar{z}_y) + \\
& 4b_{xy}{}^2 G_{xx}^2 \phi_{yx}\text{Var}(\bar{z}_x - \bar{z}_y) + 4b_{yx}{}^2 G_{yy}^2 \phi_{xy}\text{Var}(\bar{z}_x - \bar{z}_y) \\
& -8b_{xy}b_{yx}G_{xx}G_{yy}\phi_{xy}\phi_{yx}\text{Var}(\bar{z}_x - \bar{z}_y) + \\
& 4b_{xy}b_{yx}G_{xx}G_{yy}\phi_{xy}\text{Var}(\bar{z}_x - \bar{z}_y) + 4b_{yx}b_{xy}G_{xx}G_{yy}\phi_{yx}\text{Var}(\bar{z}_x - \bar{z}_y) \\
& -8b_{xy}{}^2 G_{xx}^2 \phi_{yx}^2 \text{Var}(\bar{z}_x - \bar{z}_y) - 8b_{yx}{}^2 G_{yy}^2 \phi_{xy}^2 \text{Var}(\bar{z}_x - \bar{z}_y) + \mathcal{V}]
\end{aligned} \tag{17}
$$

This expression again shows the covariance in evolutionary response occurs even in the

absence of (co)variance in b, and the addition of IIGEs alter the magnitude of the covariance

substantially. Analysis of more complex cases of the trait matching model, for example

where selection varies across space and/or across terms in the expanded polynomial

$(z_x - z_y)^2$, would be straightforward in this framework.

As a final example of how this approach can be used to understand simple variations

on classic coevolutionary models, we focus on a multiplicative model of trait interaction,

where fitness depends on the interaction between traits of the two species. Consider a fitness

model where relative fitness depends solely on the product of the two phenotypes:

$$w_x = \alpha_x + b_{xy}z_x z_y + \varepsilon_x$$
$$w_y = \alpha_y + b_{yx}z_y z_x + \varepsilon_y.$$

(18)

451   This model is conceptually and mathematically similar to the previous trait matching model,

452   but importantly lacks stabilizing selection terms (e.g., $b_{xy}z_x{}^2$) present in the trait matching

453   model. Because of this, directional natural selection depends only on the means of the other

454   species:

$$\beta_{xx} = b_{xy}\bar{z}_y$$
$$\beta_{xy} = b_{xy}\bar{z}_x$$
$$\beta_{yy} = b_{yx}\bar{z}_x$$
$$\beta_{yx} = b_{yx}\bar{z}_y$$

(19)

457   This relationship may cause selection to (co)vary across populations even when b_{xy} and b_{yx}

458   are homogeneous. In the absence of IIGEs, this simplest case would lead to a coevolutionary

459   covariance defined by

$$\mathrm{Cov}(\Delta\bar{z}_x, \Delta\bar{z}_y) = G_{xx}G_{yy}b_{xy}b_{yx}\mathrm{Cov}(\bar{z}_x, \bar{z}_y).$$

(20)

461 Thus, any initial covariance in the population means leads to a covariance in the response to

462 selection across species, lending a runaway aspect to the coevolutionary covariance. Adding

463 IIGEs, this becomes

$$
\begin{aligned}
\mathrm{Cov}(\Delta \bar{z}_x, \Delta \bar{z}_y) = \ & \mathcal{U}[G_{xx}G_{yy}b_{xy}b_{yx}\mathrm{Cov}(\bar{z}_x, \bar{z}_y) \\
& + G_{xx}^2 \phi_{yx} b_{xy}^2 \mathrm{Var}(\bar{z}_y) + G_{yy}^2 \phi_{xy} b_{yx}^2 \mathrm{Var}(\bar{z}_x) + \\
& 2 G_{xx} G_{yy} \phi_{xy} \phi_{yx} b_{xy} b_{yx} \mathrm{Cov}(\bar{z}_x, \bar{z}_y) \\
& + G_{xx} G_{yy} \phi_{xy} b_{xy} b_{yx} \mathrm{Var}(\bar{z}_y) + G_{xx} y G_y \phi_{yx} b_{xy} b_{yx} \mathrm{Var}(\bar{z}_x) + \\
& 2 G_{xx}^2 \phi_{yx}^2 b_{xy}^2 \mathrm{Cov}(\bar{z}_x, \bar{z}_y) + 2 G_{yy}^2 \phi_{xy}^2 b_{yx}^2 \mathrm{Cov}(\bar{z}_x, \bar{z}_y) + \mathcal{V}].
\end{aligned}
\tag{21}
$$

This equation again shows that when IIGEs are present, any variance across populations in

the trait mean of either species leads to a cross-species covariance in the response to selection

(terms 2, 3, 5, and 6). Cross species covariance in selection response is also mediated further

by covariance in trait means in this model (terms 1, 4, 7, and 8) as well as higher order

products of IIGEs captured in V (equation 11). These effects may lead trait means to become

correlated across populations in future generations. This analysis, particularly contrasting

equations 16 vs. 20 and 17 vs. 21, also reveals that coevolutionary dynamics can be

substantially different even across models that share a common polynomial relative fitness

function.

Discussion

Our model adapts the theory of trait-based intraspecific social evolution to the phenotypic

interface between two coevolving species. We show that two forms of interspecific

interaction, interspecific indirect genetic effects (IIGEs) and cross-species selection

(analogous to within-species social selection), both contribute to correlated evolution

between interacting species. Our analysis shows that reciprocal IIGEs modulate selection

response, suggesting that IIGEs may play a major role in generating and mediating patterns

of correlated evolution between species. Further, we show that constant (across space) IIGEs

can generate a coevolutionary covariance even in the absence of covariance in selection, or

even the absence of genetic variance in one species. When selection does covary between

species across populations, reciprocal IIGEs will promote changes in the magnitude of

coevolution and even reversals in the expected among-population covariance. IIGEs also

allow cross-species selection, which we model as the effect of the traits of one species on the

fitness of another, to influence evolutionary response. Such a response may also be mediated

by cross-species genetic assortment between interacting individuals. When IIGEs and cross-

species selection act together, effects on the coevolutionary covariance can be complex, with

dramatic changes in the expected sign and magnitude of correlated evolution occurring with

subtle changes in these parameters of interspecific social interaction. Our results indicate that

whenever coevolving species socially interact to modify expression of one another's

phenotypes, these interspecific social interactions are key to understanding coevolution.

IIGEs represent a scenario where phenotypes in individuals of one species influence

trait expression of individuals of another species. Thus, IIGEs are a specific type of

environmental effect (Moore et al. 1997, Drown and Wade 2014) where the environment is

the phenotypic value of the interspecific individual(s) with which an organism interacts. To

our knowledge, this type of environmental effect on between-species coevolution has been

considered in only two other theoretical studies (Scheiner et al. 2015, O'Brien et al. 2021; but

see Shuster et al. 2006, Witham et al. 2020 for a variance-partitioning approach). Scheiner et

al. (2015) consider a special case of our model, where evolvable IIGEs are present in only

one of the two interacting species. In this non-reciprocal model, they show a much more

limited role for IIGEs in coevolution. Our results are broadly consistent with this conclusion,

in that IIGEs in only a single species do not generate the reciprocal effects that lead to

massive inflation of evolutionary response. However, IIGEs in only a single species (e.g.,

$\Phi_{xy} = 0, \; \Phi_{yx} \neq 0$) still play a role in mediating response to interspecific social selection

whenever interspecific social selection is a function of individual trait values (as opposed to

the population mean, as modeled by Scheiner et al. 2015). We also note that our fully

multivariate model accommodates the possibility that reciprocal IIGEs act across different

types or numbers of traits in the two interacting species. More recently, O'Brien et al. (2021;

see also Queller 2014) developed a model of coevolution between host plant and microbial

symbionts. Their parameterization differed from ours in that they consider evolution of a

single joint trait governed by genetic variation in host and symbiont, and so is most directly

applicable to plant-microbe systems or other intimate interactions. Nonetheless, their model

shows an important role for reciprocal fitness feedbacks, consistent with the conclusions of

our trait-based model.

Within-species models of interacting phenotypes clarify the line between genetic and

environmental effects, furthering an understanding of how genotypes expressed in an

individual that act as environments for other individuals can influence phenotypic change

(Wolf et al. 1998, Wolf 2003). That is, IGEs are genetically-based environments that

influence phenotypic expression during interactions. Conceptually, this relationship is similar

to genetically based plasticity. Extending these types of models to the case of interspecific

interaction carries similar challenges and benefits. We have referred to reciprocal change in

phenotypic means between interacting species as "coevolution," even when these effects are

mediated by IIGEs. This is a broad use of the term coevolution, as IIGEs are an

environmental effect that can themselves evolve, again similar in concept to phenotypic

plasticity. However, a critical difference is that changes in phenotypic response mediated by

IIGEs represent changes driven by evolution of an interacting species (made clear in equation

4). Our point is not to broaden the definition of coevolution, but rather to highlight that

IIGEs can have substantial impact on patterns of phenotypic divergence in coevolving

species. For example, our model highlights that genetic divergence across populations of a

single species is sufficient to generate tightly-coupled patterns of correlated change in an

interacting species, a result that suggests the challenges of interpreting correlated phenotypes

as evidence of genetic response to reciprocal selection may be even greater than already

appreciated (e.g. Nuismer et al. 2010, Gomulkiewicz et al 2007, Janzen 1980).

Interspecific indirect genetic effects, or at least the potential for a prevalence of such

effects, appear to be commonplace in many biological systems. In Table 1, we provide in a

breakdown of types of biological interaction in which there is a large literature suggesting

importance of IIGE-like phenomena. These types of effects on trait expression across

species, widely appreciated in their own specific contexts (Weis and Abrahamson 1986,

Peacor and Werner 2001, Werner and Peacor 2003, Chen 2008, Thomas et al. 2012, O'Brien

et al. 2021), have taken on a variety of different forms. We suggest that these disparate

biological phenomena may nonetheless share a commonality—reciprocal effects on trait

expression across interspecific partners—that we have shown can affect the coevolutionary

process in dramatic, and in some cases predictable, ways.

Cross-species selection features prominently in verbal descriptions of the

coevolutionary process (Thompson 1982), and we show that such selection is especially

important in the presence of interspecific indirect genetic effects. When individual trait

values of one species affect individual fitness of another, focal species, this cross-species

selection can manifest evolutionary change in the focal species when there is phenotypic

assortment between interspecific interactants. This assortment, analogous to that required for

evolutionary response to social selection within species (Wolf et al. 1999, McGlothlin et al.

2010, Brodie et al. submitted), can be generated directly by a non-random genetic assortment,

or via IIGEs. Examples of processes that could generate direct genetic assortment between

interacting individuals of two different species include shared genetic structure, to the extent

that such structure manifests assortment of breeding values for the relevant traits. Such

shared genetic structure could arise through shared features of the environment that limit

gene flow and panmictic mating in both species, or alternatively, through variation in habitat

preference across individuals of both species. Direct genetic assortment could also arise

through behavioral preference for certain trait values in heterospecific partners. Such

preferences may be especially common in predator-prey interactions, where, for example,

predator body size may be expected to coevolve with behavioral preference for prey size

(Troost et al. 2008). Currently, it is unclear how common this type of cross-species genetic

assortment may be, although in part this likely reflects a lack of studies that have attempted to

measure assortment between breeding values of individuals of separate species. Moreover,

when it does occur, such assortment is likely to be transient because it does not rely on

transmission of pleiotropic alleles that may stabilize within-species genetic correlations over

multiple generations. The substantial evidence for IIGEs, but limited evidence of G_{xy}

suggests that IIGEs may play a prominent role in mediating any realized response to cross-

species selection.

Our results also indicate that nonlinear effects on cross-species selection can

contribute to coevolution even in the absence of genetic assortment or IIGEs. This form of

selection corresponds to an interaction between focal and interspecific-partner trait values for

focal individual fitness. The effect of this form of cross-species selection on evolutionary

response in a population depends on the mean genotype of the other species, and thus

represents a diffuse effect of population mean phenotype of the coevolving species. Such

interspecific interactions are potentially less intimate, for example diffuse predator chemical

cues in aquatic environments, than the individual level interactions (e.g., of host and parasite)

required to generate response from linear cross-species selection. Across populations,

nonlinear cross-species selection contributes to coevolution via covariance in mean genetic

values and/or linear selection between the species. This result is consistent with past models

of coevolution, verbal and mathematical, that indicate trait interactions for fitness are a key

feature of coevolution (Thompson 1982, 1994, 2005, Nuismer 2017), and in our model, such

interactions lead to a dependence between selection in one species and the mean trait value of

another. By defining these interaction terms in the framework of social evolution, our model

adds to past work by indicating that reciprocal IIGEs can substantially increase the degree to

which trait interactions for fitness contribute to reciprocal evolutionary change.

A key feature of our model is the development of a formal expression for the expected

covariance in evolutionary response, $\mathrm{Cov}(\Delta\bar{z}_x, \Delta\bar{z}_y)$. This coevolutionary covariance is

expected to be a key contributor to generating among-population covariation in species mean

trait values, a major focus in many empirical (Thompson 1994, 1995, Zangerl and

Berenbaum 2003, Toju and Sota 2005, Hanifin et al. 2008, Hague et al. 2020) and theoretical

(Nuismer et al. 2010, Nuismer and Week 2019, Week and Nuismer 2019) studies of

coevolution. Importantly, similar to existing within-species models of among-population

quantitative genetic variation (Zeng 1988, Chenoweth et al. 2010), defining this

coevolutionary covariance illustrates how selection, IIGEs, and genetics may contribute to

patterns of trait variation across populations.

Our model subsumes mechanistic detail into broad statistical descriptions of species

interactions and thus provides a general description of how IIGEs and cross-species selection,

when present, contribute to reciprocal evolutionary change and correlated evolution across

populations. In contrast to our approach, some models of coevolution have focused instead

on specific ecological mechanisms that generate trait-fitness relationships between interacting

species (reviewed in Nuismer 2017). By highlighting the key parameters that contribute to

coevolution—covariance in natural selection, covariance in cross-species selection, and

IIGEs—our model indicates various pathways through which specific ecological mechanisms

may affect coevolution. Our framework can be tailored to specific scenarios by substituting

different fitness models into the general equations we present here.

Our model generates quantitative predictions for the shape of coevolution that are

directly testable with empirical data because it focuses on estimable statistical effects of

underlying ecological mechanisms rather than the mechanisms themselves, which are often

unknown (Wade and Kalisz 1990). For example, using an empirical estimate of Φ (which

could be measured using methods analogous to those used to measure within-species indirect

genetic effects; Bleakley and Brodie 2009, McGlothlin and Brodie 2009), one could use

matrix comparison of covariances among population means and the covariance terms

presented here to quantitatively test the contribution of IIGEs to among-population

covariance in selection response between two interacting species (e.g., see Chenoweth et al.

2010 for a within-species test of the predictions of Zeng's 1988 model). More generally,

Week and Nuismer (2019; see also Nuismer and Week 2019) have shown how datasets of

among-population variation in trait means can be used to test for conformation to

expectations from coevolutionary models. Concomitantly, our models show how

environmental effects can be partitioned into terms describing genotypes of other species in

the ecological community, which could be useful in understanding when and why

evolutionary response fails to conform to predictions arising from the standard breeder's

equation.

Social interactions between individuals of the same species play a central role in the

evolutionary process. Within a single lineage, indirect genetic effects and social selection

fundamentally change selection response, the expression of genetic variance, and together

determine the course of social evolution (Moore et al. 1997, Wolf et al. 1998, Wolf et al.

1999, McGlothlin et al. 2010). We have shown that these effects of interactions among

individuals may transcend species boundaries and profoundly impact the dynamics of

coevolution between interacting lineages.

Literature Cited

Abrahamson, W. G., and A. E. Weis. 1997. Evolutionary Ecology Across Three Trophic Levels: Goldenrods, Gallmakers, and Natural Enemies. Princeton University Press Princeton.

Abrams, P. A., Y. Harada, and H. Matsuda. 1993. On the relationship between quantitative genetic and ESS models. Evolution 47:982-985.

Agrawal, A. A. 2001. Phenotypic plasticity in the interactions and evolution of species. Science 294:321-326.

Arnold, S. J., M. E. Pfrender, and A. G. Jones. 2001. The adaptive landscape as a conceptual bridge between micro- and macroevolution. Genetica 112/113:9-32.

Arnqvist, G., and L. Rowe. 2002. Antagonistic coevolution between the sexes in a group of insects. Nature 415:787-789.

Barribeau, S. M. 2014. Gene expression differences underlying genotype-by-genotype specificty in a host-parasite system. Proceedings of the National Academy of Sciences 111:3496-3501.

Bijma, P., and M. J. Wade. 2008. The joint effects of kin, multilevel selection and indirect genetic effects on response to genetic selection. Journal of Evolutionary Biology 21:1175-1188.

Bleakley, B. H., and E. D. Brodie III. 2009. Indirect genetic effects influence antipredator behavior in guppies: Estimates of the coefficient of interaction *psi* and the inheritance of reciprocity. Evolution 63:1796-1806.

Boersma, M., P. Spaak, and L. De Meester. 1998. Predator-mediated plasticity in morphology, life history, and behavior of *Daphnia*: the uncoupling of responses. American Naturalist 152:237-248.

Borell, E. M., A. Foggo, and R. A. Coleman. 2004. Induced resistance in intertidal macroalgae modifies feeding behavior of herbivorous snails. Oecologia 140:328-334.

Brodie, E. D., III, and E. D. Brodie, Jr. 1999. Predator-prey arms races. BioScience 49:557-568.

Brodie, E. D. Jr., B. J. Ridenhour, and E. D. Brodie III. 2002. The evolutionary response of predators to dangerous prey: hotspots and coldspots in the geographic mosaic of coevolution between garters snakes and newts Evolution 56:2067-2082.

Brodie, E. D., III and B. J. Ridenhour 2003 Reciprocal selection at the phenotypic interface of coevolution. Integrative and Comparative Biology 43:408-418

Brodie, E.D., III, P. A. Cook, R. A. Costello, V. A. Formica. Submitted. Phenotypic assortment changes the landscape of selection. Journal of Heredity.

Chen, M.-S. 2008. Inducible direct plant defense against insect herbivoes: A review. Insect Science 15:101-114.

Chenoweth, S. F., H. D. Rundle, and M. W. Blows. 2010. The contribution of selection and genetic contraints to phenotypic divergence. American Naturalist 175:186-196.

Damian, R. T. 1997. Parasite immune evasion and exploitation: reflections and projections. Parasitology 115:S169-175.

Dawkins, R. 1982. The Extended Phenotype. Oxford Univ. Press, Oxford.

Drown, D. M., and M. J. Wade. 2014. Runaway coevolution: adaptation to heritable and nonheritable environments. Evolution 68:3039-3046.

Dybdahl, M. F., C. E. Jenkins, and S. L. Nuismer. 2014. Identifying the molecular basis of host-parasite coevolution: merging models and mechanisms. American Naturalist 184:1-13.

Ehrlich, P. R., and P. H. Raven. 1964. Butterflies and plants: a study in coevolution. Evolution 18:586-608.

Fisher, D.N. and A. McAdam. 2019. Indirect genetic effects clarify how traits can evolve
even when fitness does not. Evolution Letters 3:4-14.
Foster, K. R., J. Schluter, K. Z. Coyte, and S. Rakoff-Nahoum. 2017. The evolution of the
host microbiome as an ecosystem on a leash. Nature 548:43-51.
Friesen, M. L., S. S. Porter, S. C. Stark, E. J. von Wettberg, J. L. Sachs, and E. Martinez-
Romero. 2011. Microbially mediated plant functional traits. Annual Review of
Ecology and Systematics 42:23-46.
Fuess, L., J. N. Weber, S. den Haan, N. C. Steinel, K. C. Shim, and D. I. Bolnick. 2020. A
test of the Baldwin effect: Differences in both constitutive expression and inducible
responses to parasites underlie variation in host response to a parasite. bioRxiv
https://doi.org/10.1101/2020.07.29.216531.
Gianinazzi-Pearson, V., N. Séjalon-Delmas, A. Genre, S. Jeandroz, and P. Bonfante. 2007.
Plants and arbuscular mycorrhizal fungi: cues and communication in the early steps of
symbiotic interactions. Advances in Botanical Research 46:181-219.
Gomulkiewicz, R., D.M. Drown, M.F. Dybdahl, W. Godsoe, S.L. Nuismer, K.M. Pepin, B.J.
Ridenhour, C.I. Smith, and J.B. Yoder. 2007. Dos and don'ts of testing the geographic
mosaic theory of coevolution. Heredity 98:249-258.
Hague, M. T. J., A. N. Stokes, C. R. Feldman, E. D. Brodie Jr., and E. D. Brodie III. 2020.
The geographic mosaic of arms race coevolution is closely matched to prey
population structure. Evolution Letters 164:1567–16.
Hanifin, C. T., E. D. Brodie Jr., and E. D. Brodie III. 2008. Phenotypic mismatches reveal
escape from arms-race coevolution. PLoS Biology 6:e60.
Janzen, D. H. 1980. When is it coevolution? Evolution 34:611-612.
Johnson, S. D., and B. Anderson. 2010. Coevolution between food-rewarding flowers and
their pollinators. Evolution: Eduacation and Outreach 3:32-39.
Kirkpatrick, M. and R. Lande 1989. The evolution of maternal characters. Evolution 43:485-
503.
Korgaonkar, A., C. Han, A. L. Lemire, I. Siwanowicz, D. Bennouna, R. E. Kopec, P.
Andolfatto, S. Shigenobu, and D.L. Stern. 2021. A novel family of secreted insect
proteins linked to plant gall development. Current Biology 31:1-14
Lande, R. 1979. Quantitative genetic analysis of multivariate evolution, applied to brain:
body size allometry. Evolution 33:402-416.
Lande, R., and S. J. Arnold. 1983. The measurement of selection on correlated characters.
Evolution 37:1210-1226.
McAdam, A.G., S. Boutin, D. Réale, and D. Berteaux. 2002. Maternal effects and the
potential for evolution in a natural population of animals. Evolution 56: 846-851.
McGlothlin, J. W., and E. D. Brodie III. 2009. How to measure indirect genetic effects: the
congruence of trait-based and variance-partitioning approaches. Evolution 63:1785-
1795.
McGlothlin, J. W., E. Akçay, E. D. Brodie III, A. J. Moore, and J. Van Cleve. 2021. A
synthesis of game theory and quantitative genetic models of social evolution. bioRxiv
https://doi.org/10.1101/2021.03.27.437341
McGlothlin, J. W., A. J. Moore, J. B. Wolf, and E. D. Brodie III. 2010. Interacting
phenotypes and the evolutionary process. III. Social evolution. Evolution 64:2558-
2574.
Moore, A. J., E. D. Brodie III, and J. B. Wolf. 1997. Interacting phenotypes and the
evolutionary process: I. Direct and indirect genetic effects of social interactions.
Evolution 51:1352-1362.
Mousseau, T.A. and C.W. Fox. 1998. The adaptive significance of maternal effects. Trends in
Ecology and Evolution 13:403-407

Nuismer, S. L. 2017. Introduction to Coevolutionary Theory. W.H. Freeman and Company
New York
Nuismer, S. L., B. J. Ridenhour, and B. P. Oswald. 2007. Antagonistic coevolution mediated
by phenotypic differences between quantitative traits. Evolution 61:1823-1834.
Nuismer, S.L., R. Gomulkiewicz, and B.J. Ridenhour. 2010. When is correlation coevolution.
The American Naturalist 175:525-537.
Nuismer, S.L., and B. Week. 2019. Approximate Bayesian estimation of coevolutionary arms
races. PLoS Computational Biology 15:e1006988.
Ohgushi, T. 2005. Indirect interaction webs: Herbivore-induced effects through trait change
in plants Annual Review of Ecology, Evolution, and Systematics 36:81-105.
Oladiran, A., and M. Belosevic. 2012. Immune evasion strategies of trypanosomes: a review.
Journal of Parasitology 98:284-292.
O'Brien, A. M., C. N. Jack, M. L. Friesen, and M. E. Frederickson. 2021. Whose trait is it
anyways? Coevolution of joint phenotypes and genetic architecture in mutualisms.
Proceedings of the Royal Society of London Series B. 288:20202483.
Peacor, S. D., and E. E. Werner. 2001. The contribution of trait-mediated indirect effects to
the net effects of a predator. Proceedings of the National Academy of Sciences
98:3904-3908.
Pellmyr, O. 2003. Yuccas, yucca moths, and coevolution: A review. Annals of the Missouri
Botanical Garden 90:35-55.
Phillips, P.C., and S.J. Arnold. 1989. Visualizing multivariate selection. Evolution 43:1209-
1222.
Preisser, E. L., D. I. Bolnick, and M. F. Benard. 2005. Scared to death? The effects of
intimidation and consumption in predator-prey interactions. Ecology 86:501-509.
Price, G. R. 1970. Selection and covariance. Nature 227:520-521.
Price, G. R. 1972. Extension of covariance mathematics. Annals of Human Genetics 35:485-
490.
Queller, D. C. 2014. Joint phenotypes, evolutionary conflict and the fundamental theorem of
natural selection. Philosophical Transactions of the Royal Society B. 369:20130423.
Ridenhour, B. J. 2005. Identification of selective sources: Partitioning selection based on
interactions. The American Naturalist 166:12-25.
Robertson, A. 1966. A mathematical model of the culling process in dairy cattle. Animal
Production 8:95-108.
Scharsack, J. P., A. Gossens, F. Franke, and J. Kurtz. 2013. Excretory products of the
cestode, Schistocephalus solidus, modulate in vitro responses of leukocytes from its
specific host, the three-spined stickleback (*Gasterosteus aculeatus*). Fish and
Shellfish Immunology 35:1779-1787.
Scharsack, J. P., M. Kalbe, R. Derner, J. Kurtz, and M. Milinski. 2004. Modulation of
granulocyte responses in three-spined sticlebacks Gasterosteus aculeatus infected with
the tapeworm Schistocephalus solidus. Diseases of Aquatic Organisms 59:141-150.
Scharsack, J. P., K. Koch, and K. Hammerschmidt. 2007. Who is in control of the stickleback
immune system: interactions between *Schistocephalus solidus* and its specific
vertebrate host. Proceedings of the Royal Society of London Series B. 274:3151-
3158.
Scheiner, S. M., R. Gomulkiewicz, and R. D. Holt. 2015. The genetics of phenotypic
plasticity. XIV. Coevolution. American Naturalist 185:594-609.
Schmid-Hempel, P. 2008. Parasite immune evasion: a momentous molecular war. Trends in
Ecology and Evolution 23:318-326.

Shuster, S. M., E. V. Lonsdorf, G. M. Wimp, J. K. Bailey, and T. G. Whitham. 2006.
Community heritability measures the evolutionary consequences of indirect genetic
effects on community structure. Evolution 60:991-1003.
Siepielski, A. M., K. M. Gotanda, M. B. Morrissey, S. E. Diamond, J. D. DiBattista, and S.
M. Carlson. 2013. The spatial patterns of directional phenotypic selection. Ecology
Letters 16:1382-1392.
Stappenbeck, T. S., and H. W. Virgin. 2016. Accounting for reciprocal host-microbiome
interactions in experimental science. Nature 534:191-199.
Stearns, S. C. 2012. Evolutionary routes leading to host manipulation by parasites: Afterward
in F. Thomas, J. Brodeur, and D. P. Hughes, editors. Host manipulation by parasites.
Oxford University Press, Oxford.
Tams, V., J. H. Nickel, A. Ehring, and M. Cordellier. 2019. Insights into the genetic basis of
predator-induced response in *Daphnia* - a comparative transcriptomic approach.
bioRxiv. https://doi.org/10.1101/503904
Thomas, F., J. Brodeur, and D. P. Hughes. 2012. Host manipulation by parasites Oxford
University Press, Oxford.
Thompson, J. N. 1982. Interaction and Coevolution. University of Chicago Press, Chicago.
Thompson, J. N. 1994. The Coevolutionary Process. University of Chicago Press, Chicago.
Thompson, J. N. 2005. The Geographic Mosaic of Coevolution. University of Chicago Press,
Chicago.
Toju, H., and T. Sota. 2005. Imbalance of predator and prey armament: geographic clines in
phenotypic interface and natural selection. American Naturalist 167:105–117
Troost, T. A., B. W. Kooi, and U. Dieckmann. 2008. Joint evolution of predator body size
and prey size preference. Evolutionary Ecology 22:771-799.
Urban, M.C. 2011. The evolution of species interactions across natural landscapes. Ecology
Letters 14:723-732.
Wade, M. J., and Kalisz. 1990. The causes of natural selection. Evolution 44:1947-1955.
Walsh, M. R., and D. M. Post. 2011. The impact of intraspecific vriation in a fish predator on
the evolution of phenotypic plasticity and investment in sex in *Daphnia ambigua*.
Journal of Evolutionary Biology 25:80-89.
Weber, A., and S. Declerck. 1997. Phenotypic plasticity of *Daphnia* life history traits in
response to predator kairomones: genetic variability and evolutionary potential
Hydrobiologia 360:89-99.
Weber, J. N., N. C. Steinel, K. C. Shim, and D. I. Bolnick. 2017. Recent evolution of extreme
cestode growth suppression by a vertebrate host. Proceedings of the National
Academy of Sciences 114:6575-6580.
Week, B. and S.L. Nuismer. 2019. The measurement of coevolution in the wild. Ecology
Letters 22:717-725
Weis, A. E., and W. G. Abrahamson. 1986. Evolution of host-plant manipulations by gall
makers: ecological and genetic factors in the Solidago-Eurosta system. American
Naturalist 127:681-695.
Weis, A.E., W.G. Abrahamson, and M.C. Anderson. 1992. Variable selection on Eurosta's
gall size, I: The extent and nature of variation in phenotypic selection. Evolution
46:1674-1697.
Werner, E. E., and S. D. Peacor. 2003. A review of trait-mediated indirect interactions in
ecological communities. Ecology 84:1083-1100.
West-Eberhard, M. J. 1979. Sexual selection, social competition, and evolution. Proceedings
of the American Philosophical Society 123:222-234.
West-Eberhard, M. J. 1983. Sexual selection, social competition, and speciation. Quarterly
Review of Biology 58:155-183.

831 West-Eberhard, M. J. 1984. Sexual selection, competitive communication and species-
832    specific signals in insects. Pages 283-342 *in* T. Lewis, editor. Insect communication.
833    Academic Press, New York.
834 Whitham, T. H., G. J. Allan, H. F. Cooper, and S. M. Shuster. 2020. Intraspecific genetic
835    variation and species interactions contribute to community evolution. Annual Review
836    of Ecology and Systematics 51:587-612.
837 Wolf, J. B., E. D. Brodie III, J. M. Cheverud, A. J. Moore, and M. J. Wade. 1998.
838    Evolutionary consequences of indirect genetic effects. Trends in Ecology and
839    Evolution 13:64-69.
840 Wolf, J. B., E. D. Brodie III, and A. J. Moore. 1999. Interacting phenotypes and the
841    evolutionary process. II. Selection resulting from social interactions. American
842    Naturalist 153:254-266.
843 Wolf, J.B. 2003. Genetic architecture and evolutionary constraint when the environment
844    contains genes. Proceedings of the National Academy of Sciences 100:4655-4660.
845 Zangerl, A. R., and M. R. Berenbaum. 2003. Phenotype matching in wild parsnip and parsnip
846    webworms: causes and consequences. Evolution 57:806–815
847 Zeng, Z.-B. 1988. Long-term correlated response, interpopulation covariation, and
848    interspecific allometry. Evolution 42 363-374.
849 Züst, T., S. Mou, and A.A. Agrawal. 2018. What doesn't kill you makes you stronger: The
850    burdens and benefits of toxin sequestration in a milkweed aphid. Functional Ecology
851    8:1972-1981

Figure S1. IIGEs accelerate evolutionary rate in a single species. Panels show the evolutionary rate, Var($\Delta \bar{z}_x$), of species x as a function of the indirect genetic effect parameters ϕ, under the same parameter values as in Figure 3. Reciprocal IIGEs between interacting species generally accelerate evolutionary rate. Note that in the absence of any other effects, the evolutionary rate is equal to the variance in natural selection, which is unity in Panels A, C, E, and F. In panel D, evolutionary rate is driven entirely by cross-species selection and IIGEs. In panel B, where $G_{xx} = 0$, evolutionary rate in species x is driven entirely by reciprocal IIGEs and evolutionary change in species y.

Figure S2. Among population covariances. Panels show the coevolutionary covariance between two interacting species as a function of the indirect genetic effect parameters ϕ, from equation (7) unstandardized. Parameter values are as in Figs. 4 and S1. For all panels, genetic variances were set to unity.

Appendix

A1. Phenotypic covariance between interacting species

For a single trait, we can solve for the covariance between z_y and z_x to partition the phenotypic covariance between individuals of two interacting species into terms describing the contribution of IIGEs and terms describing non-random genetic assortment. Assuming cross-species environmental covariance is zero,

$$C_{xy} = (1 - \phi_{yx}\phi_{xy})^{-2}[(1 + \phi_{yx}\phi_{xy})G_{xy} + \phi_{yx}(G_x + E_x) + \phi_{xy}(G_{yy} + E_x)]$$

where E_x and E_y represent within-species environmental variance and $G_{xy} = G_{yx}$. When IIGEs are absent, non-random genetic assortment G_{xy} is the sole contributor to the phenotypic association between individuals of coevolving species. When IIGEs are present, they can substantially change this phenotypic association.

A2. Covariance in selection response with nonzero IIGEs and genetic assortment

We expand the covariance in evolutionary response in two species when IIGEs are present and constant and when genetic assortment $G_{xy} = G_{yx}$ is present and constant,

$$\begin{aligned}
\text{Cov}(\Delta\bar{z}_x, \Delta\bar{z}_y) = \mathcal{U}[&\mathcal{A}^2\phi_{yx}\text{Var}(\beta_{xx}) + \mathcal{B}^2\phi_{yx}\text{Var}(\beta_{xy}) + \mathcal{C}^2\phi_{xy}\text{Var}(\beta_{yy}) \\
&+ \mathcal{D}^2\phi_{xy}\text{Var}(\beta_{yx}) + \mathcal{A}\mathcal{B}\mathcal{E}\text{Cov}(\beta_{xx}, \beta_{xy}) + \mathcal{A}\mathcal{C}\mathcal{E}\text{Cov}(\beta_{xx}, \beta_{yy}) \\
&+ \mathcal{A}\mathcal{D}\mathcal{E}\text{Cov}(\beta_{xx}, \beta_{yx}) + \mathcal{B}\mathcal{C}\mathcal{E}\text{Cov}(\beta_{xy}, \beta_{yy}) + \mathcal{B}\mathcal{D}\mathcal{E}\text{Cov}(\beta_{xy}, \beta_{yx}) \\
&+ \mathcal{C}\mathcal{D}\mathcal{E}\text{Cov}(\beta_{yy}, \beta_{yx})]
\end{aligned}$$

874   where

$$\mathcal{U} = (1 - \phi_{xy}\phi_{yx})^{-4}$$

$$\mathcal{A} = G_{xx} + G_{xy}\phi_{xy}$$

$$\mathcal{B} = G_{xy} + G_{xx}\phi_{yx}$$

$$\mathcal{C} = G_{yy} + G_{xy}\phi_{yx}$$

$$\mathcal{D} = G_{xy} + G_{yy}\phi_{xy}$$

$$\mathcal{E} = 1 + \phi_{xy}\phi_{xy}$$