Title: Clinicopathologic features of a feline SARS-CoV-2 infection model

parallel acute COVID-19 in humans

Authors: Jennifer M. Rudd[1], Miruthula Tamil Selvan[1], Shannon Cowan[1], Cecily C. Midkiff[2],

Jerry W. Ritchey[1], Craig A. Miller[1*]

Affiliations:

[1] Department of Veterinary Pathobiology, College of Veterinary Medicine, Oklahoma State

University; Stillwater, OK, USA

[2] Division of Comparative Pathology, National Primate Research Center, Tulane University;

Covington, LA, USA

* Corresponding author. Email: craig.miller@okstate.edu (CAM)

One Sentence Summary: Intra-tracheal SARS-CoV-2 inoculation in domestic cats induces

analogous clinicopathologic features to hospitalized patients with acute COVID-19.

Keywords: COVID-19, SARS-CoV-2, Domestic, Cats, Feline, Model, ARDS, Clinical Disease

Abstract: The emergence and ensuing dominance of COVID-19 on the world stage has

emphasized the urgency of efficient animal models for the development of therapeutics and

assessment of immune responses to SARS-CoV-2 infection. Shortcomings of current animal

models for SARS-CoV-2 include limited lower respiratory disease, divergence from clinical

COVID-19 disease, and requirements for host genetic modifications to permit infection. This

study validates a feline model for SARS-CoV-2 infection that results in clinical disease and

histopathologic lesions consistent with severe COVID-19 in humans. Intra-tracheal inoculation

of concentrated SARS-CoV-2 caused infected cats to develop clinical disease consistent with

that observed in the early exudative phase of COVID-19. A novel clinical scoring system for

feline respiratory disease was developed and utilized, documenting a significant degree of

lethargy, fever, dyspnea, and dry cough in infected cats. In addition, histopathologic pulmonary

lesions such as diffuse alveolar damage, hyaline membrane formation, fibrin deposition, and

proteinaceous exudates were observed due to SARS-CoV-2 infection, imitating lesions identified

in people hospitalized with ARDS from COVID-19. A significant correlation exists between the

degree of clinical disease identified in infected cats and pulmonary lesions. Viral loads and

ACE2 expression were quantified in nasal turbinates, distal trachea, lung, and various other

organs. Natural ACE2 expression, paired with clinicopathologic correlates between this feline

model and human COVID-19, encourage use of this model for future translational studies.

Main Text:

INTRODUCTION

Since the emergence of severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) from China in late 2019, Coronavirus Disease 2019 (COVID-19) has swept across the globe resulting in nearly 3 million deaths worldwide as of March 2021 [1]. Although a wide range of clinical symptoms are reported, mortality of COVID-19 patients is closely correlated with progression of viral infection to severe lung disease (pneumonia) and respiratory failure due to acute respiratory distress syndrome (ARDS), which is further complicated by immune cell dyscrasias and hyperinflammation (cytokine storm) in critically ill patients [2-4]. Features of pulmonary pathology that are hallmarks of severe COVID-19 (i.e., diffuse alveolar damage with hyaline membrane formation, type II pneumocyte hyperplasia, vascular thrombi, fibrin and serous exudation) have been difficult to reproduce in animal models, making it impossible to completely understand the pathophysiology of disease and to test efficacy of new therapeutics and vaccines [5, 6]. Thus, identification of a translational animal model that parallels clinical and pathologic features of disease in addition to route of infection, replication, and transmission kinetics is of paramount importance.

SARS-CoV-2 viral infection and replication within a host requires the presence and distribution of angiotensin-converting enzyme 2 (ACE2) receptors similar to humans [7]. Natural SARS-CoV-2 infections in animals are documented to occur in a diverse range of species, including domestic and exotic cats, dogs, mink, and Golden Syrian hamsters [8-11], and this diverse host range is largely due to natural expression of ACE2 receptors and host tropism of this receptor with the S protein of SARS-CoV-2 [12, 13]. Due to the natural availability of ACE2

receptors and confirmed host susceptibility and transmission [10, 14-17], domestic cats offer an

exciting advantage as experimental models for SARS-CoV-2 infection [18, 19]. Comorbidities

that exacerbate COVID-19 disease, such as hypertension, diabetes, renal disease, and obesity, are

readily adapted to feline models [20-25]. In addition, establishing a SARS-CoV-2 infected feline

model is prudent to better understand zoonotic transmission potential from domestic cats back to

their owners and others in close contact.

Previous studies have successfully infected cats with SARS-CoV-2 via intra-nasal (1-

3.05×10^5 PFU) and/or intra-oral routes (5×10^5 $TCID_{50}$/ml) and have confirmed cat-to-cat

transmission through both respiratory droplets and aerosolization [16, 26-28]. However, these

infection studies failed to produce clinical signs in infected cats, and evidence of lower

respiratory pathology mirroring severe COVID-19 in humans was not observed [16, 26-28],

potentially due to concentration of the viral inoculum and/or inoculation route. Interestingly,

pulmonary disease with diffuse alveolar damage was previously documented in cats intra-

tracheally infected with 1×10^6 $TCID_{50}$ SARS-CoV-1, which also resulted in efficient

transmission of virus to uninfected animals [29, 30].

Based on outcomes of these former studies, we hypothesized that inoculation with a

higher concentration of SARS-CoV-2 via the intra-tracheal route would result in pulmonary

pathology and clinical disease in domestic cats similar to COVID-19 in human patients. The

experiments reported in this study provide the first feline model of SARS-CoV-2 infection with

significant lower respiratory disease that displays features of diffuse alveolar damage seen in the

early exudative phase of human COVID-19. In addition, SARS-CoV-2 infected cats exhibited

clinical signs of lower respiratory disease characterized by increased respiratory effort and

coughing in addition to signs of systemic involvement such as pyrexia and lethargy. While the

role of cats in zoonotic transmission is still under investigation, the applicability of a clinically

significant SARS-CoV-2 feline model with pathological lesions that mirror severe COVID-19 is

of high impact for future studies.

RESULTS

SARS-CoV-2 infected cats exhibit clinical signs of lower respiratory disease

In order to clinically assess the feline model in Animal Biosafety Level-3 conditions, a

novel clinical scoring system for feline respiratory disease was developed by integrating features

of previously utilized systems [31-33] (Table 1). Each cat was assigned a score from 0 to 2 for

each of the following categories: body weight loss, activity levels, behavioral changes, body

temperature, respiratory effort, ocular or nasal discharge, and coughing. Scores were then

summated to assign an overall clinical score for each day.

	0 (Healthy)	1	2
Body Weight	No weight loss	0 to 5% weight loss	> 5% weight loss
Temperature	99 to 102.5° F	102.6 to 103.5° F	> 103.5° F
Respiratory Effort	Normal resting respiratory rate and normal effort	Mild tachypnea, but no overt increase in effort	Marked increase in both respiratory rate and effort; dyspnea
Activity	Normal	Reduced activity when disturbed* (lethargy)	Little to no activity disturbed*; reduced activity stimulated**
Behavior	Normal	Reduced interest in food and/or attention	Anorexia and lack of interest
Ocular/Nasal Discharge	None	Mild discharge noted	Discharge evident from both nasal and ocular regions
Coughing/Wheezing	None	Mild wheezing, but no coughing	Coughing and/or marked wheezing

*Disturbed: observer in the room, but kennel unopened
**Stimulated: kennel open

Table 1. Clinical Scoring System for Feline Respiratory Disease. A scoring system was

designed to assess clinical lower respiratory and systemic disease in the feline model. Each cat

was scored daily at the same time point (morning) by a small animal clinician (JMR). Cats were

assigned a score from 0 to 2 for each clinical parameter: body weight, temperature, respiratory

effort, activity, behavior, ocular/nasal discharge, coughing/wheezing. The parameter scores were

summed to assign an overall score per cat per day. Potential scores can range from 0 (healthy

with no signs of disease) to 14 on any given day. Resting respiratory rate was considered normal

if <36 breaths per minute. Marked increases in rate were >50 breaths per minute at rest.

Temperatures were obtained via thermal microchips, and body weights were obtained last to

limit stress affecting clinical scoring.

SARS-CoV-2-infected cats exhibited a significant increase in clinical disease scores starting on 4

days post-inoculation (dpi) and then on 5, 6, and 8 dpi when compared to sham-inoculated

controls (Fig. 1A). Clinical disease peaked on 4 dpi and continued through the study endpoint at

day 8. The most prominent clinical signs noted were lethargy and increased respiratory effort,

which were observed in 100% (12/12) of SARS-COV-2-infected cats during this study. Both

lethargy and respiratory effort increase significantly between 3 and 4 dpi (p=0.0027; p=0.0027)

and remained elevated with significantly higher scores through 8 dpi when compared with day 0

(Figure 1B). Coughing was noted in 4 of 12 infected cats with peak clinical signs occurring at 4

dpi. Pyrexia (temperature > 102.5°F) was documented in 8 of 12 SARS-CoV-2 infected cats over

the course of the study, while 7 infected cats displayed altered behavior (reduced interest in food

or attention) and 5 had measurable weight loss. No cats had ocular or nasal discharge (Table S1).

Sham-inoculated cats did not exhibit clinical signs except for one cat with mild weight loss on

day 4.

Figure 1. Intra-tracheal SARS-CoV-2 inoculation results in clinical disease. Clinical

parameters were assessed using the feline respiratory disease clinical scoring system (see Table

1). **(A)** Clinical parameters summated to provide an overall clinical score per cat per day.

Clinical disease severity peaked on 4 dpi and was significantly higher than sham-inoculated cats

on 4 dpi (p=0.0054), 5 dpi (p=0.0257), 6 dpi (p=0.0004), and 8 dpi (p=0.0453). A noticeable

trend in severity was also noted on 7 dpi as compared with sham-inoculated controls (p=0.0654).

(B) Lethargy and increased respiratory effort were the most prominent clinical signs observed in

SARS-CoV-2-infected cats; both of which were significantly increased between days 3 and 4

(p=0.0027; p=0.0027) and remained significantly elevated in infected cats after 4 dpi as

compared to day 0. Coughing was most prominent on 4 dpi and was identified in 4/12 infected

cats. Pyrexia was noted in 8/12 cats over the course of the study. Data are expressed as means ±

SEM. Statistical comparisons made via mixed effects analysis. *p<0.05; **p<0.01; ***p<0.001;

****p<0.0001.

Feline SARS-CoV-2 infection pathology mirrors acute COVID-19

Complete post-mortem evaluation was performed for all sham-inoculated control animals

(n=6) and SARS-CoV-2-infected cats euthanized on day 4 (n=6) and day 8 (n=6) post-

inoculation. Necropsy tissues from SARS-CoV-2-infected cats (including lung, trachea, nasal

turbinates, and tracheobronchial lymph node) were grossly examined and compared to those

from sham-inoculated cats (Figure 2A-C). At 4 dpi, the lungs of SARS-CoV-2-infected cats were

grossly heavy and wet, with large multifocal to coalescing regions of dark red consolidation that

exuded a moderate amount of serosanguineous fluid (edema) upon cut section (Figure 2B). Gross

lung lesions were similar at 8 dpi in SARS-CoV-2-infected cats, although the degree of

pulmonary edema was moderately more pronounced (Figure 2C). The tracheobronchial lymph

nodes (TB LN) of all SARS-CoV-2-infected cats were diffusely enlarged 4-5 times normal at

both 4 dpi (n=6) and 8 dpi (n=6). No other significant gross findings were observed.

Figure 2. Pathologic features of acute SARS-CoV-2 infection in cats are analogous to the exudative phase of COVID-19. Compared to lungs from healthy sham-inoculated cats **(A)**, the lungs of SARS-CoV-2-infected cats were diffusely consolidated, dark red and edematous at both 4 dpi **(B)** and 8 dpi **(C)**. The lungs of healthy, uninfected cats **(D)** were histologically normal, with open alveoli and minimal atelectasis. At 4 dpi, the lungs of SARS-CoV-2-infected cats **(E)** exhibited discrete foci of alveolar inflammation and necrosis with fibrin deposition (⬚), increased alveolar macrophages (arrowhead), perivascular lymphocytes (⬚), and type II pneumocyte hyperplasia (arrows). The alveoli in these cats' lungs were frequently filled with large amounts of edema and fibrin strands **(E inset)**, and there were multifocal areas of hyaline membrane formation **(F)** (⬚). The distal trachea of 1 SARS-CoV-2 infected cat **(G)** was multifocally ulcerated at 4 dpi (arrow) with diphtheritic membrane formation (arrowhead) and multifocal areas of submucosal necrosis and fibrinoid vasculitis (⬚). At 8 dpi **(H)**, fibrinoid vasculitis, vascular thrombosis (arrow), and occasional syncytial cells **(H inset)** were observed in addition to the histopathologic changes described above. Tracheal lesions observed at 8 dpi **(I)** were characterized by varying degrees of lymphoplasmacytic, histiocytic, and neutrophilic inflammation with multifocal areas of submucosal necrosis. Magnification: (D, H) 10x, scale bar = 100µm; (E) 20x, scale bar = 50µm; (E inset, G, H inset, I) 40x, scale bar = 25µm; (F) 60x, scale bar = 17µm

Microscopic evaluation of selected necropsy tissues (lung, trachea, nasal turbinates, TB LN,

and kidney) was performed for all study animals. Tissue sections from all sham-inoculated

animals (n=6) were histologically unremarkable (Figure 2D and Table S2). In contrast,

histopathologic features of feline SARS-CoV-2 infection exhibited striking similarities to

documented pathologic features of the acute (exudative) and organizing phases of human

COVID-19 [34-37]. At 4 dpi, 100% (6/6) of SARS-COV-2 -infected cats exhibited a significant

degree of lung pathology (interaction, $p<0.0001$) and prominent histologic features consistent

with diffuse alveolar damage (DAD) (Figure 2E-F). Pulmonary serous exudate/edema (5/6 cats),

multifocal alveolar damage and necrosis (5/6 cats), perivascular lymphocytic and neutrophilic

infiltrates (6/6 cats), and significantly increased numbers of intra-alveolar macrophages (5/6 cats)

were noted in SARS-CoV-2-infected cats at 4 dpi (Table S2). These changes were occasionally

accompanied by multifocal areas of hyaline membrane formation (3/6 cats), mild to moderate

amounts of intra-alveolar fibrin (2/6 cats), type II pneumocyte hyperplasia (2/6 cats), and intra-

alveolar syncytial cells (2/6 cats). One SARS-CoV-2 infected cat exhibited severe inflammation

in the distal trachea at 4 dpi characterized by multifocal areas of submucosal necrosis, fibrinoid

vasculitis admixed with degenerate neutrophils and multifocal areas of mucosal ulceration with

diphtheritic membrane formation (Figure 2G and Table S2).

Similar histologic features of DAD were also observed in the lungs of SARS-CoV-2-infected

cats at 8 dpi, however, the overall pattern of lung injury appeared to exhibit more prominent

features of vascular injury compared to day 4 (Figure 2H). A significant degree of pulmonary

edema/exudate, perivascular inflammatory infiltration, and alveolar histiocytosis was present in

100% of SARS-CoV-2 animals (6/6 cats) at 8 dpi (Table S2). Alveolar damage and necrosis (4/6

cats) and intra-alveolar fibrin (3/6 cats) were also prominent features at this time point.

Moreover, histologic evidence of fibrinoid vasculitis (2/6 cats) and vascular thrombosis (2/6

cats) was also observed at 8 dpi, in addition to occasional viral syncytia (1/6 cats) (Figure 2H).

In 2 of these cats, the tracheal submucosa was multifocally expanded and effaced by moderate to

severe lymphoplasmacytic, histiocytic, and neutrophilic inflammation with necrosis (Figure 2I

and Table S2).

Mild, multifocal lymphofollicular inflammation was observed in the nasal turbinates of all

SARS-CoV-2-infected cats (6/6) at 4 dpi and in 4/6 cats at 8 dpi, with variable neutrophilic

infiltration (Table S2). All SARS-CoV-2-infected animals (12/12) exhibited mildly increased

lymphoid hyperplasia in tracheobronchial lymph nodes at 4 and 8 dpi characterized by increased

medullary cords and extranodal proliferations (Table S2), but overall changes were not

statistically significant. No significant histopathologic findings were observed in renal tissues at

either time point. A positive linear correlation exists between peak clinical scores and

histopathology scores of the lungs in SARS-CoV-2 infected cats (p=0.0002; R^2=0.5884)

indicating that more severe clinical signs of disease correlate with worsened pulmonary

pathology (Figure S1).

Fluorescent immunohistochemistry was performed to detect SARS-CoV-2 positive cells in

lung and TB LN of 2 SARS-CoV-2 infected cats (n=1 at 4 dpi, n=1 at 8 dpi). At both time

points, low numbers of mononuclear cells positive for SARS-CoV-2 nucleoprotein were detected

within the medulla of the TB LN, however, no positive cells were observed in lungs of these

animals. (Figure 3).

Figure 3. Fluorescent immunohistochemistry for SARS-CoV-2 nucleoprotein identifies

mononuclear cells in tracheobronchial lymph node of intratracheally-infected cats. Low

numbers of SARS-CoV-2 positive cells (green, white arrows) are detected in **(A)** positive control

tissue (lung) from an African Green Monkey infected with SARS-CoV-2 [38], and within **(B)**

mononuclear cells in the TB LN of SARS-CoV-2 infected cats (green, white arrow). White =

DAPI/nuclei; green = CoV-2. Magnification (A-B) 40x, scale bar = 20 µm.

ACE2 expression and viral RNA in feline tissues during SARS-CoV-2 infection

SARS-CoV-2 viral RNA and feline ACE2 (fACE2) receptor RNA expression was

quantified in the nasal turbinates, tracheobronchial lymph nodes, distal trachea, kidneys and

lungs of all SARS-CoV-2-infected cats (n=12) and sham-inoculated controls (n=6) using droplet

digital PCR (ddPCR) technology (Table S3). Viral RNA was detected in 100% of tissues

collected on 4 dpi from SARS-CoV-2-infected cats (Figure 4A). At 8 dpi, viral RNA was also

detectable in all tissues, except 1 of 6 cats had no detectable viral RNA in the nasal turbinates

and 1 of 6 cats had none detected in the distal trachea. No SARS-CoV-2 viral RNA was detected

in tissues collected from sham-inoculated cats at either time point (Table S3). SARS-CoV-2 viral

RNA copies were elevated in the tracheobronchial lymph node at 8 dpi compared with day 4

samples, although this trend was not significant (p=0.0567). In contrast, SARS-CoV-2 viral load

in lung samples was significantly lower at 8 dpi than at 4 dpi (p=0.0007) (Figure 4A). SARS-

CoV-2 RNA was not reliably detected in nasal swabs or plasma of infected cats at either time

point. However, a positive linear correlation was observed between SARS-CoV-2 RNA in the

lung and pulmonary histopathology scores of SARS-CoV-2 infected cats (p=0.0183; R^2=0.

3012), suggesting that increased pulmonary viral loads correlate with more severe pulmonary

pathology (Fig. S1).

In sham-inoculated cats, Kruskal Wallis test revealed that fACE2 RNA in the nasal

turbinates was significantly higher than in the lung (p=0.0093) and TB LN (p=0.0049). fACE2

RNA was also higher in the kidney when compared to lung (p=0.0003), trachea (p=0.0034), and

TB LN (p=0.0001). These findings were similar in SARS-CoV-2-infected animals at 4 and 8 dpi,

with fACE2 RNA levels being significantly higher in the nasal turbinates and kidney versus

other tissues (p<0.05) (see Table S4). Overall, fACE2 RNA in the kidney was significantly

increased in SARS-CoV-2 infected cats on 4 dpi when compared to both sham-inoculated

controls, as well as SARS-CoV-2 infected cats on 8 dpi ($p<0.0001$) (Figure 4B). No other

significant changes in ACE2 RNA were observed over time.

Figure 4. fACE2 RNA and SARS-CoV-2 viral RNA quantification in feline tissues.

Extraction of SARS-CoV-2 and ACE2 RNA was performed as described from tissues samples

collected on either 4 or 8 dpi. Tissue samples included nasal turbinates, tracheobronchial lymph

node (TB LN), trachea, kidney, and lung. **(A)** SARS-CoV-2 RNA copies were detected in all

tissues collected from cats inoculated with SARS-CoV-2. No viral RNA was detected in tissues

from sham-inoculated cats. In SARS-CoV-2-infected cats, viral RNA copies were slightly

increased in the TB LN between 4 to 8 dpi ($p=0.0567$) while viral RNA load in the lungs

significantly decreased over the same period ($p=0.0007$). **(B)** fACE2 receptor RNA is

significantly increased in the kidney of SARS-CoV-2 infected cats at 4 dpi compared to sham-

inoculated controls ($p<0.0001$) SARS-CoV-3-infected cats at 8 dpi ($p<0.0001$). Data are

expressed as means ± SEM. n=6 cats per group. Statistical comparisons made via two-way

ANOVA. ***$p<0.001$; ****$p<0.0001$.

DISCUSSION

The potential of this feline model for future evaluation of COVID-19 is extensive.

Challenges with earlier feline models of SARS-CoV-2 infection included a lack of clinical

disease and/or pathology of the lower respiratory tract that resembles lesions seen in patients

with COVID-19. The differences in clinical presentation between previous feline models and the

model described here are likely attributed to modifications in routes of inoculation and

concentration of the inoculate. In this study, SARS-CoV-2 was inoculated through an intra-

tracheal route and at a higher concentration than previously reported [16, 26, 27]. Route of

inoculation is an important consideration when establishing an animal model for disease, and

previous studies in other animal models have exhibited marked differences in primary disease

severity and distribution based on route of inoculation [39]. For example, in feline and ferret

models of influenza A/H5N1, intra-tracheal inoculation of the virus resulted in more severe

lower respiratory disease as well as more prominent signs of systemic disease [39, 40].

While previous feline models offer value for study of asymptomatic infections, viral

shedding, and transmission of SARS-CoV-2, cats infected through an intra-tracheal route exhibit

clinical disease that aligns with that seen in early phases of acute COVID-19. Clinical

assessment of infectious lower respiratory disease in a cat can be challenging, and it is not

uncommon for cats with confirmed histologic infectious pneumonia to have limited clinical

respiratory signs [41]. Therefore, it is highly significant that clinical signs of respiratory disease

were induced in this model. In order to carefully assess for clinical disease in this study, a novel

clinical scoring system was designed that could be applied in the Animal Biosafety Level-3

facility. Interestingly, the disease noted in the SARS-CoV-2 infected cats was similar to that

described in hospitalized patients with COVID-19. Clinical disease of hospitalized human

COVID-19 patients is characterized by fever (70-90%), dry cough (60-86%), shortness of breath

(53-80%), and fatigue (38%) [42] while predominant clinical sings in SARS-CoV-2 infected cats

consisted of fever, cough, lethargy and increased respiratory effort, with lethargy and increased

respiratory effort being the most consistent and notable clinical signs (Fig. 1).

In addition to clinical signs of lower respiratory and systemic disease, SARS-CoV-2

infected cats also exhibited conspicuous pulmonary lesions of diffuse alveolar damage (DAD) by

4 dpi, and additional evidence of vascular damage by 8 dpi (Fig. 2). Specific histopathological

lesions align closely with those reported in human COVID-19 patients [34-36, 43-45], including

DAD resulting in hyaline membrane formation, type II pneumocyte hyperplasia, occasional

intra-alveolar syncytial cells, and the development of fibrinous exudate and vascular thrombi. To

the author's knowledge, this is the first report of hyaline membrane formation and type II

pneumocyte hyperplasia in feline SARS-CoV-2 infection, which are classic characteristics of

human SARS-CoV-2 infection. Peak clinical disease scores positively correlated with severity of

histologic lesions in the lungs (Fig. S1), which further support that cats with marked pulmonary

histologic damage also had more severe clinical signs of disease.

Surprisingly, intra-tracheal inoculation of SARS-CoV-2 did not produce high viral RNA

loads in the lungs as compared with other studies in which the inoculate was delivered via the

intranasal route [26]. However, despite by-passing the upper airway and inoculating via intra-

tracheal administration, virus was still detected in the nasal turbinates by 4 and 8 dpi, suggesting

the virus may utilize the mucociliary escalator to travel up the respiratory tree and establish

infection intra-nasally even without intra-nasal inoculation. Although seemingly lower quantities

of SARS-CoV-2 RNA were recovered from lungs of intra-tracheally inoculated cats, the damage

to lung tissues was highly evident, indicating that extensive pulmonary damage will occur even

without high levels of viral replication within the pulmonary tissue at 4 and 8 dpi. Viral

migration from lung to the TBLN occurred quickly (by 4 dpi) and this TBLN involvement is a

novel finding to the feline model of SARS-CoV-2, including detection of viral antigen within the

TBLN via fluorescent immunohistochemistry.

Similar to humans, fACE2 RNA expression varied by tissue location, but were relatively

low in the lungs of both infected and uninfected cats. It is important to note that RNA

measurements indicate an upregulation or downregulation of production of proteins, but do not

necessarily indicate an absolute number of receptors available. However, it is possible that

inefficient replication and rapid clearance of SARS-CoV-2 in the lungs is related to lower

expression of ACE2 receptors as compared with nasal turbinate ACE2. Histopathology shows

that cells regularly expressing ACE2 are damaged in the lung and this viral-induced pulmonary

epithelial pathology may contribute further. ACE2 RNA copies in the feline kidney are

significantly higher than that of other assessed tissues, and viral infection resulted in a significant

upregulation of ACE2 RNA by 4 dpi and then a subsequent reduction by 8 dpi. Hypertension and

activation of the renin-angiotensin system may have driven this rise in ACE2 in order to

counterbalance system effects on infection, and future studies should include blood pressure

evaluation in conjunction with other clinical parameters such as oxygen saturation, chemistry

panels, and imaging. Further studies are needed to fully understand the role of ACE2 in SARS-

CoV-2 viral replication kinetics and disease.

Limitations to this study include sample sizes as well as sampling time points. Further

studies are needed to try and better identify peak viral loads in various tissues as well as ACE2

expression in order to investigate when viral clearance occurs and when this feline model moves

from early exudative disease to more organized, fibrotic disease. A better delineation of these

events would add value to the model and its potential for use at other stages of disease. In

addition, transmission and viral shedding after intra-tracheal inoculation of SARS-CoV-2 should

be evaluated and compared with that of intra-nasal inoculation and spread. Expansion of in depth

diagnostics was limited due to animal biosafety level requirements and availability of resources,

but future studies will seek to evaluate other clinical parameters, such as oxygen saturation,

thoracic imaging and complete blood counts, chemistry panels and urine analysis to better assess

damage to other organ systems and further compare with human disease.

This feline model of SARS-CoV-2 infection offers an animal model that closely mirrors

both clinical disease and pathology identified in hospitalized patients with severe COVID-19,

making the model a potential option for future studies addressing novel therapeutics for COVID-

19. Therapeutic measures can be thoroughly assessed for improvement in pathology and

mitigation of clinical disease in cats before being validated in human trials, and more thorough

evaluation of the feline immune response to infection may elucidate other options for COVID-19

treatments that could mitigate disease and improve clinical outcomes. The continued emergence

of novel variants, circulating globally, ceaselessly contributes to the complexity and duration of

this pandemic. This animal model offers an ease of use, which may positively impact further

vaccination and control strategies necessary to achieve an end to the rapid spread of COVID-19.

Because cats are naturally infected with SARS-CoV-2 and have been shown to have varying

degrees of clinical disease, this model also offers utility in a One Health approach to the role of

companion animals in disease transmission, antigenic drift, and more thorough evaluation of the

potential for feline contributions to the spread of SARS-CoV-2.

MATERIALS AND METHODS

Ethics Statement. This study was approved by the Oklahoma State University Institutional

Animal Care and Use Committee; IACUC-20-48, Validation of a naturally occurring animal

model for SARS-CoV-2 infection. Oklahoma State University's animal care and use program is

licensed by the United States Department of Agriculture (USDA), and accredited by the

Association for Assessment and Accreditation of Laboratory Animal Care (AAALAC). In

accordance with the approved IACUC protocol, animals were monitored at least once daily for

evidence of morbidity and discomfort by trained animal care staff. Prior to experimental

procedures, all study animals were anesthetized to minimize animal suffering and distress.

Human euthanasia procedures were conducted by phenobarbital overdose in accordance with

IACUC protocols and American Veterinary Medical Association (AVMA) Guidelines for the

Euthanasia of Animals. Prior to euthanasia, all study animals were anesthetized by intramuscular

injection of ketamine (4 mg/kg), dexmedetomidine (0.02 mg/kg), and butorphanol (0.4 mg/kg).

No animals died without euthanasia during this study.

Virus. SARS-CoV-2 virus isolate USA-WA1/2020 was obtained from BEI Resources, passaged

up to 6 times in Vero E6 cells in Vero E6 cell growth medium. Virus stock was titrated and

quantified on Vero E6 cells using a standard MERS-CoV quantification assay [46]. $TCID_{50}$ was

calculated using the Reed and Muench method.

Animals. Eighteen adult (9 males, 9 females, all 9 months-old) specific pathogen free (SPF) cats

were obtained from Marshall BioResources (North Rose, NY). Animals intended for SARS-

CoV-2 inoculation were housed within Biosafety Level 3 (BSL-3) barrier animal rooms at

Oklahoma State University, individually housed and fed dry/wet food with access to water *ad*

libitum. Animals intended for sham-inoculation were group-housed within the AAALAC

International Accredited animal facility at Oklahoma State University. Animals were allowed 30

days for acclimation prior to initiation of the study. Temperature-sensing microchips (Bio Medic

Data Systems, Seaford, DE) were implanted subcutaneously in the dorsum after 30 days.

Baseline weights, body temperatures, clinical evaluation, and nasal swab sampling were obtained

prior to inoculation. All animals were in apparent good health at the onset of the study.

Virus Challenge. Cats were lightly anesthetized with ketamine (4 mg/kg), dexmedetomidine (20

µg/kg), and butorphanol (0.4 mg/kg) intramuscularly. Cats were then positioned in ventral

recumbency and intubated so that the end of an endotracheal tube is positioned within the distal

trachea as described [47]. In twelve cats, a 3-cc syringe was used to inoculate 1 mL 9×10^5 PFU

(1.26×10^6 $TCID_{50}$) per mL SARS-CoV-2, isolate USA-WA1/2020 in Dulbecco's Modified Eagle

Medium (DMEM), followed by 2 mL of air from an empty syringe. The remaining six cats were

sham-inoculated using sterile PBS via the same method. Viral inoculum dosage was confirmed

through virus back-titration on E6 cells immediately following inoculation.

Sampling. Blood and nasal swab samples were collected under sedation (described above) from

all study animals (n=18) at day 0 to serve as baseline for ddPCR analysis. Blood samples (6 mL)

were obtained from all cats via cephalic or medial saphenous venipuncture and processed

immediately for viral quantification. Nasal swab samples obtained from the nares of all cats

using ultrafine flocked swabs (Puritan) were placed in 2 mL microcentrifuge tubes containing

RNAlater solution (Ambio, Austin, TX) and stored at -80°C until processed. At day 4 and day 8

post-inoculation, a subset of SARS-CoV-2 infected cats (n=6 per time point) and sham-

inoculated control cats (n=3 per time point) were anesthetized for blood and nasal swab

collection then humanely euthanized (pentobarbital >80mg/kg) and necropsied to collect tissue

samples. Necropsied tissues were processed for histologic examination, immunohistochemistry

(IHC), and RNA analysis as described below.

Clinical Observations and Scoring. Animals were monitored at least once daily for evidence of

morbidity and discomfort by a licensed veterinary practitioner. Body weights and temperatures

(thermal microchips) were documented daily every morning for the duration of the study. Full

clinical scoring included evaluation of body weight, body temperature, activity levels, behavior,

respiratory effort, evidence of ocular/nasal discharge, and recognition of coughing or wheezing.

Each factor was assigned a score of 0 (normal), 1 (mild-moderate), or 2 (severe) as described in

Table 1. Each clinical factor parameter was added to assign an individual animal a summed

clinical score every 24 hours for the duration of the study. Cats were observed at rest for

respiration rates, activity levels and other notable clinical signs before stimulation.

Histopathology. Necropsy was performed on six (n=6) SARS-CoV-2-infected cats at 4 dpi and

the remaining six (n=6) SARS-CoV-2-infected cats at 8 dpi. Three (n=3) sham-inoculated cats

were necropsied at each time point (4 dpi and 8 dpi) to provide control samples. Tissue

collection included: lung, tracheobronchial lymph nodes, nasal turbinates, distal trachea, and

kidney. Necropsy tissues were halved and then placed into either 1 mL tubes and frozen at -

80°C, or into standard tissue cassettes that were then fixed in 10% neutral-buffered formaldehyde

for 96 hours prior transferring to 70% ethanol for 72 hours. Tissues were then trimmed and

processed for histology. Five μm paraffin sections were collected onto charged slides, and one

slide of each tissue was stained with hematoxylin and eosin (H & E) for microscopic evaluation.

Necropsy tissues were evaluated for evidence of inflammation and/or aberrations in lymphoid

populations as reported in human COVID-19 patients [34-37]. Lung tissues were specifically

evaluated for the following pathology: alveolar damage (pneumocyte necrosis, hyaline

membrane formation,) alveolar fibrin deposition (± organization), serous exudate/edema,

perivascular infiltrates, alveolar histiocytes, type II pneumocyte hyperplasia, syncytia,

thrombosis, and fibrinoid vasculitis. All tissues were assigned a quantitative histologic score

based on previously documented criteria [48, 49]: 0 = no apparent pathology/change; 1 =

minimal change (minimally increased numbers of inflammatory cells); 2 = mild change (mild

inflammatory infiltrates, alveolar damage/necrosis, fibrin deposition and/or exudation); 3 =

moderate change (as previously described, but more moderately extensive); 4 = marked changes

(as previously described, but with severe inflammation, alveolar damage, hyaline membrane

formation, necrosis, exudation, vasculitis and/or thrombosis). All tissues were evaluated and

scored by a board-certified veterinary pathologist blinded to study groups to ensure scientific

rigor and reproducibility.

Viral RNA Analysis. Viral RNA analysis was performed on samples from nasal swabs,

collected plasma, and necropsy tissues. Nasal swabs were immediately broken off into 1.5 mL

microcentrifuge tubes containing 200 μL RNAlater Solution (Ambion, Austin, TX) and stored at

-20°C. The nasal swabs were vortexed for 15 seconds, then inverted and centrifuged at 1500 rpm

for 10 minutes. RNA was extracted from frozen necropsy tissues using a QIAamp Viral RNA

Mini Kit (Qiagen, Germantown, MD) and tissue homogenizer. SARS-CoV-2 viral RNA was

quantified by droplet digital PCR (ddPCR) as previously described [50]. Briefly, ddPCR was

performed according to manufacturer's instructions for the 2019-nCoV CDC ddPCR Triplex

Probe Assay (Bio-Rad, Hercules, California, USA). PCR reaction mixtures were as follows: 5.5

µl One-Step RT-ddPCR Advanced Kit for Probes Supermix (no dUTP's) (Bio-Rad), 2.2 µl

reverse transcriptase, 1.1 µl 300 mM Dithiothreitol (DTT), 1.1 µl triplex probe assay (for N1,

N2, RPP30 detection), 2.2 µl RNase free water, and 9.9 µl RNA template in a final volume of 22

µl per sample. Duplicate 20 µl samples were partitioned using a QX200 droplet generator (Bio-

Rad, Hercules, California, USA) and then transferred to a 96-well plate and sealed. Samples

were processed in a C1000 touch Thermal Cycler (Bio-Rad) under the following cycling

protocol: 50 °C for 60 min for reverse transcription, 95 °C for 10 min for enzyme activation, 94

°C for 30 s for denaturation and 55 °C for 60 s for annealing/extension for 45 cycles, 98 °C 10

min for enzyme deactivation, 4 °C for 30 min for droplet stabilization followed by infinite 4 °C

hold. The amplified samples were read in the FAM and HEX channels using the QX200 reader

(Bio-Rad). Each experiment was performed with a negative control (no template control, NTC)

and a positive control (RNA extracted from SARS-CoV-2 viral stock and diluted 1:12,000). Data

were analyzed using QuantaSoft™ v1 AnalysisPro Software (Bio-Rad) and expressed as Log_{10}

(copies/mL).

Feline ACE2 Analysis. Feline angiotensin converting enzyme 2 (fACE2) RNA was quantified

by ddPCR using similar methods for CoV-2. RNA was extracted from frozen necropsy tissues

as outlined above, and cDNA was synthesized as previously published [49]. Design of primers

and probe targeting fACE2 was performed according to manufacturer's recommendation,

namely keeping GC content between 50–60 % for primers and 30–80 % for probes, and melting

temperatures between 50–65 ◦C for primers and 3–10 ◦C higher for probes. Oligoes were

synthesized by Integrated DNA Technologies (IDT, Coralville, Iowa, USA), and the sequences

are as follows: Forward: 5′- ACGGAGGCGTAAGGATTT -3′ , Reverse: 5′ -

GTGTGGTAGTGGTTGGTATTG -3′ , probe: 5′ - CGGGATCAGAAATCGAAGGAAGAA -

3′. BLAST analysis [51] of the primer and probe sequences against the domestic cat (*Felis catus*)

genome was performed to ensure no similar sequences could be amplified. ddPCR reactions

were prepared by adding 11 µl Supermix for Probes (no dUTP) (Bio-Rad), 1.1 µl of primer/probe

mix (final concentration is 500nM for primers and 250 nM for probe) and 8.8 µl of cDNA

template containing 110 ng RNA equivalent. Droplets were partitioned and PCR executed as

above using the following cycling conditions: 95 °C for 10 min, 95 °C 30 s for denaturation and

58.8 °C for 60 s for annealing/extension for 45 cycles, 98 °C 10 min for enzyme deactivation.

Droplets were read and analyzed as described above.

Immunohistochemistry. 5µm sections of formalin-fixed, paraffin-embedded lung were

mounted on charged glass slides, baked for one hour at 60°C, and passed through Xylene, graded

ethanol, and double distilled water to remove paraffin and rehydrate tissue sections. A

microwave was used for heat induced epitope retrieval. Slides were heated in a high pH solution

(Vector Labs H-3301), rinsed in hot water and transferred to a heated low pH solution (Vector

Labs H-3300) where they were allowed to cool to room temperature. Sections were washed in a

solution of phosphate-buffered saline and fish gelatin (PBS-FSG) and transferred to a humidified

chamber, for staining at room temperature. Tissues were blocked with 10% normal goat serum

(NGS) for 40 minutes, followed by a 60-minute incubation with a guinea pig anti-SARS

antibody (BEI NR-10361) diluted 1:1000 in NGS. Slides were washed and transferred to the

humidified chamber for a 40-minute incubation with a goat anti-guinea pig secondary antibody

(Invitrogen A11073) tagged with Alexa Fluor 488 and diluted 1:1000 in NGS. Following

washes, DAPI (4',6-diamidino-2-phenylindole) was used to label the nuclei of each section.

Slides were mounted using a homemade anti-quenching mounting media containing Mowiol

(Calbiochem#475904) and DABCO (Sigma #D2522) and imaged at 20X with a Zeiss Axio Slide

Scanner.

Statistical Analyses. When applicable, data were expressed as mean ± SEM and statistically

analyzed using GraphPad Prism 9.0 software (La Jolla, CA). Kruskal–Wallis test, Pearson

correlations, and ANOVA were used to compare differences in clinical score, histopathology,

SARS-CoV-2 viral load, and ACE2 RNA among uninfected and SARS-CoV-2-infected

individuals, between sample type, for each tissue individually, and between tissues. For all

significant results, pair-wise comparisons were made by post-hoc analysis. P-values < 0.05 were

considered significant.

Supplementary Materials (see attached)

Table S1. Clinical scoring data, summary, analysis

Table S2. Histopathologic scoring data and statistical analyses

Figure S1. Positive linear correlation between histopathologic score and peak clinical score

Table S3. SARS-CoV-2 and fACE2 RNA data and statistical analyses

Table S4. fACE2 statistical analysis by tissue type

References and Notes

1.   WHO. *World Health Organization Coronavirus (COVID-19) Dashboard*. 2021; Available

from: https://covid19.who.int.

2.   Wu, C., et al., *Risk factors associated with acute respiratory distress syndrome and death*

in patients with coronavirus disease 2019 pneumonia in Wuhan, China. JAMA internal

medicine, 2020.

3.   Chen, G., et al., *Clinical and immunologic features in severe and moderate Coronavirus*

Disease 2019. The Journal of Clinical Investigation, 2020.

4.   Pedersen, S.F. and Y.-C. Ho, *SARS-CoV-2: a storm is raging.* The Journal of Clinical

Investigation, 2020.

5.   Kumar, S., et al., *Selection of animal models for COVID-19 research.* Virusdisease, 2020.

31(4): p. 1-6.

6.   Gretebeck, L.M. and K. Subbarao, *Animal models for SARS and MERS coronaviruses.* Curr

Opin Virol, 2015. **13**: p. 123-9.

7.   Hikmet, F., et al., *The protein expression profile of ACE2 in human tissues.* Mol Syst Biol,

2020. **16**(7): p. e9610.

8.   Kiros, M., et al., *COVID-19 pandemic: current knowledge about the role of pets and other*

animals in disease transmission. Virol J, 2020. **17**(1): p. 143.

9.   Oreshkova, N., et al., *SARS-CoV-2 infection in farmed minks, the Netherlands, April and*

May 2020. Euro Surveill, 2020. **25**(23).

10.   Newman, A., et al., *First Reported Cases of SARS-CoV-2 Infection in Companion Animals -*

New York, March-April 2020. MMWR Morb Mortal Wkly Rep, 2020. **69**(23): p. 710-713.

11.   Molenaar, R.J., et al., *Clinical and Pathological Findings in SARS-CoV-2 Disease Outbreaks*

517   *in Farmed Mink (Neovison vison).* Vet Pathol, 2020. **57**(5): p. 653-657.

518   12.   Luan, J., et al., *Spike protein recognition of mammalian ACE2 predicts the host range and*

519   *an optimized ACE2 for SARS-CoV-2 infection.* Biochem Biophys Res Commun, 2020.

520   **526**(1): p. 165-169.

521   13.   Liu, Y., et al., *Functional and genetic analysis of viral receptor ACE2 orthologs reveals a*

522   *broad potential host range of SARS-CoV-2.* Proc Natl Acad Sci U S A, 2021. **118**(12).

523   14.   Hayashi, T., et al., *Highly conserved binding region of ACE2 as a receptor for SARS-CoV-2*

524   *between humans and mammals.* Vet Q, 2020. **40**(1): p. 243-249.

525   15.   McAloose, D., et al., *From People to Panthera: Natural SARS-CoV-2 Infection in Tigers*

526   *and Lions at the Bronx Zoo.* mBio, 2020. **11**(5).

527   16.   Shi, J., et al., *Susceptibility of ferrets, cats, dogs, and other domesticated animals to*

528   *SARS-coronavirus 2.* Science, 2020.

529   17.   Zhang, Q., et al., *A serological survey of SARS-CoV-2 in cat in Wuhan.* Emerg Microbes

530   Infect, 2020. **9**(1): p. 2013-2019.

531   18.   Lakdawala, S.S. and V.D. Menachery, *The search for a COVID-19 animal model.* Science,

532   2020. **368**(6494): p. 942-943.

533   19.   Hernández, M., et al., *Are Animals a Neglected Transmission Route of SARS-CoV-2?*

534   Pathogens, 2020. **9**(6).

535   20.   Van de Velde, H., et al., *The cat as a model for human obesity: insights into depot-*

536   *specific inflammation associated with feline obesity.* Br J Nutr, 2013. **110**(7): p. 1326-35.

21.   Nelson, R.W. and C.E. Reusch, *Animal models of disease: classification and etiology of*

diabetes in dogs and cats. J Endocrinol, 2014. **222**(3): p. T1-9.

22.   Samaha, G., et al., *The Burmese cat as a genetic model of type 2 diabetes in humans.*

Anim Genet, 2019. **50**(4): p. 319-325.

23.   Hoenig, M., *The cat as a model for human obesity and diabetes.* J Diabetes Sci Technol,

2012. **6**(3): p. 525-33.

24.   Wallner, M., et al., *A Feline HFpEF Model with Pulmonary Hypertension and*

Compromised Pulmonary Function. Sci Rep, 2017. **7**(1): p. 16587.

25.   Prat, V., et al., *Human heart failure with preserved ejection versus feline*

cardiomyopathy: what can we learn from both veterinary and human medicine? Heart

Fail Rev, 2017. **22**(6): p. 783-794.

26.   Gaudreault, N.N., et al., *SARS-CoV-2 infection, disease and transmission in domestic cats.*

Emerg Microbes Infect, 2020. **9**(1): p. 2322-2332.

27.   Bosco-Lauth, A.M., et al., *Experimental infection of domestic dogs and cats with SARS-*

CoV-2: Pathogenesis, transmission, and response to reexposure in cats. Proc Natl Acad

Sci U S A, 2020. **117**(42): p. 26382-26388.

28.   Gaudreault, N.N., et al., *Experimental re-infected cats do not transmit SARS-CoV-2.*

Emerg Microbes Infect, 2021. **10**(1): p. 638-650.

29.   van den Brand, J.M., et al., *Pathology of experimental SARS coronavirus infection in cats*

and ferrets. Vet Pathol, 2008. **45**(4): p. 551-62.

30. van den Brand, J.M., et al., *The pathology and pathogenesis of experimental severe*

acute respiratory syndrome and influenza in animal models. J Comp Pathol, 2014.

151(1): p. 83-112.

31. Huet, O., et al., *Ensuring animal welfare while meeting scientific aims using a murine*

pneumonia model of septic shock. Shock, 2013. **39**(6): p. 488-94.

32. Hartmann, A.D., et al., *Efficacy of pradofloxacin in cats with feline upper respiratory tract*

disease due to Chlamydophila felis or Mycoplasma infections. J Vet Intern Med, 2008.

22(1): p. 44-52.

33. Steagall, P.V. and B.P. Monteiro, *Acute pain in cats: Recent advances in clinical*

assessment. J Feline Med Surg, 2019. **21**(1): p. 25-34.

34. Polak, S.B., et al., *A systematic review of pathological findings in COVID-19: a*

pathophysiological timeline and possible mechanisms of disease progression. Modern

Pathology, 2020. **33**(11): p. 2128-2138.

35. Tian, S., et al., *Pathological study of the 2019 novel coronavirus disease (COVID-19)*

through postmortem core biopsies. Modern Pathology, 2020. **33**(6): p. 1007-1014.

36. von der Thüsen, J. and M. van der Eerden, *Histopathology and genetic susceptibility in*

COVID-19 pneumonia. European journal of clinical investigation, 2020. **50**(7): p. e13259.

37. Xu, Z., et al., *Pathological findings of COVID-19 associated with acute respiratory distress*

syndrome. The Lancet respiratory medicine, 2020. **8**(4): p. 420-422.

38. Blair, R.V., et al., *Acute Respiratory Distress in Aged, SARS-CoV-2–Infected African Green*

Monkeys but Not Rhesus Macaques. The American journal of pathology, 2021. **191**(2): p.

274-282.

39.   Bodewes, R., et al., *Pathogenesis of Influenza A/H5N1 virus infection in ferrets differs*

between intranasal and intratracheal routes of inoculation. Am J Pathol, 2011. **179**(1): p.

30-6.

40.   Rimmelzwaan, G.F., et al., *Influenza A virus (H5N1) infection in cats causes systemic*

disease with potential novel routes of virus spread within and between hosts. Am J

Pathol, 2006. **168**(1): p. 176-83; quiz 364.

41.   Macdonald, E.S., et al., *Clinicopathologic and radiographic features and etiologic agents*

in cats with histologically confirmed infectious pneumonia: 39 cases (1991-2000). J Am

Vet Med Assoc, 2003. **223**(8): p. 1142-50.

42.   Wiersinga, W.J., et al., *Pathophysiology, Transmission, Diagnosis, and Treatment of*

Coronavirus Disease 2019 (COVID-19): A Review. Jama, 2020. **324**(8): p. 782-793.

43.   Xu, Z., et al., *Pathological findings of COVID-19 associated with acute respiratory distress*

syndrome. Lancet Respir Med, 2020. **8**(4): p. 420-422.

44.   Fox, S.E., et al., *Pulmonary and cardiac pathology in African American patients with*

COVID-19: an autopsy series from New Orleans. Lancet Respir Med, 2020. **8**(7): p. 681-

686.

45.   Borczuk, A.C., et al., *COVID-19 pulmonary pathology: a multi-institutional autopsy cohort*

from Italy and New York City. Mod Pathol, 2020. **33**(11): p. 2156-2168.

46.   Coleman, C.M. and M.B. Frieman, *Growth and quantification of MERS-CoV infection.*

Current protocols in microbiology, 2015. **37**(1): p. 15E. 2.1-15E. 2.9.

47.   Ritchey, J.W., et al., *Constitutive expression of types 1 and 2 cytokines by alveolar*

600   *macrophages from feline immunodeficiency virus-infected cats.* Vet Immunol

601   Immunopathol, 2001. **79**(1-2): p. 83-100.

602   48.   Miller, C., et al., *Strain-specific viral distribution and neuropathology of feline*

603   *immunodeficiency virus.* Veterinary immunology and immunopathology, 2011. **143**(3): p.

604   282-291.

605   49.   Miller, C., et al., *Pathogenesis of oral FIV infection.* PloS one, 2017. **12**(9): p. e0185138.

606   50.   Deiana, M., et al., *Assessment of the direct quantitation of SARS-CoV-2 by droplet digital*

607   *PCR.* Scientific reports, 2020. **10**(1): p. 1-7.

608   51.   O'Driscoll, A., et al., *HBLAST: Parallelised sequence similarity--A Hadoop MapReducable*

609   *basic local alignment search tool.* J Biomed Inform, 2015. **54**: p. 58-64.

Acknowledgments: The authors would like to thank Girish Patil, Akhilesh Ramachandran, and

Sai Narayanan with the Oklahoma Animal Disease Diagnostic Lab for their assistance in viral

RNA extraction and technical advisement. We would also like to acknowledge Eva Kao for her

animal handling and sample collection assistance with control cats during the study and Curtis

Andrew for assistance with slide preparation and coordination of IHC from the

Immunopathology Core Laboratory at Oklahoma State College of Veterinary Medicine.

Funding: Research reported in this publication was supported by the National Institute of

General Medical Sciences of the National Institutes of Health under Award Number

P20GM103648 (CAM). The content is solely the responsibility of the authors and does not

necessarily represent the official views of the National Institutes of Health.

Author Contributions:

Conceptualization: CAM, JMR, JWR

Methodology: CAM, JMR, JWR, SC, CCM

Investigation: CAM, JMR, MTS, SC, JWR, CCM

Visualization: CAM, JMR, MTS

Funding Acquisition: CAM

Writing – original draft: JMR, CAM

Writing – review and editing: JWR, CCM, SC, MTS

Competing Interests: Authors declare that they have no competing interests.

Data and materials availability: All data associated with this study are available in the main

text or the supplementary materials.

	0 (Healthy)	1	2
Body Weight	No weight loss	0 to 5% weight loss	> 5% weight loss
Temperature	99 to 102.5° F	102.6 to 103.5° F	> 103.5° F
Respiratory Effort	Normal resting respiratory rate and normal effort	Mild tachypnea, but no overt increase in effort	Marked increase in both respiratory rate and effort; dyspnea
Activity	Normal	Reduced activity when disturbed* (lethargy)	Little to no activity disturbed*; reduced activity stimulated**
Behavior	Normal	Reduced interest in food and/or attention	Anorexia and lack of interest
Ocular/Nasal Discharge	None	Mild discharge noted	Discharge evident from both nasal and ocular regions
Coughing/Wheezing	None	Mild wheezing, but no coughing	Coughing and/or marked wheezing

*Disturbed: observer in the room, but kennel unopened
**Stimulated: kennel open