Environmental changes dictate selection and nonspecific epistasis in an empirical phenotype-environment-fitness landscape

Chen, J.Z.[1], Fowler, D.M.[2,3], and Tokuriki, N.[1] *

[1]Michael Smith Laboratories, University of British Columbia

[2]Department of Genome Sciences, University of Washington

[3]Department of Bioengineering, University of Washington

*Correspondence: tokuriki@msl.ubc.ca

Summary

The fitness landscape, a function that maps genotypic and phenotypic changes to their effects on fitness, is an invaluable concept in evolutionary biochemistry. Though widely discussed, measurements of phenotype-fitness landscapes in proteins remain scarce. Here, we quantify all single mutational effects on fitness and phenotype (antibiotic resistance level) of VIM-2 β-lactamase (5600 variants) across a 64-fold range of ampicillin concentrations by deep mutational scanning. We then construct a phenotype-fitness landscape that takes variations in environmental selection pressure into account (a phenotype-environment-fitness landscape). We found that a simple, empirical landscape accurately models the ~39,000 mutational data points, which suggests the evolution of VIM-2 can be predicted based on the selection environment. Our landscape provides new quantitative knowledge on the evolution of the β-lactamases and proteins in general, particularly their evolutionary dynamics under sub-inhibitory antibiotic concentrations, as well as the mechanisms and environmental dependence of nonspecific epistasis.

Introduction

The quantitative understanding of genotype-phenotype-fitness relationships of proteins is central in evolutionary biochemistry (Dalziel et al., 2009; Kaltenbach and Tokuriki, 2014; Yi and Dean, 2019). Genotype, a protein's sequence information, largely dictates its phenotypes, which are the measurable *in vitro* physio-chemical traits of proteins such as protein activity and stability. Nonetheless, *in vivo*, multiple factors could modulate protein function and behavior in the cell (*e.g.*, the level of protein expression). The fitness of the organism carrying a specific protein is greatly influenced by the environment where the organism lives, and the selection pressure to a particular protein's function vary significantly with environmental factors. Thus, deciphering the 'phenotype-fitness' relationship across environmental variation is essential to understanding and potentially predicting a protein's evolutionary dynamics (Bershtein et al., 2017; DePristo et al., 2005; Kemble et al., 2019; Pál et al., 2006). The phenotype-fitness landscapes typically manifest as a non-linear relationship, in the form of a 'threshold' or 'mesa-like' model (Dasmeh and Serohijos, 2018; DePristo et al., 2005; Hartl et al., 1985; Lundin et al., 2018; Lunzer et al., 2005; Sarkisyan et al., 2016; Tokuriki and Tawfik, 2009). In this model, the fitness conferred by variants with biochemical traits above a certain selection 'threshold' would saturate or plateau, while fitness rapidly decreases for variants below the threshold of the trait. Such models have been widely recognized conceptually, and provide molecular explanations for important evolutionary phenomenon such as nonspecific epistasis (Bershtein et al., 2006; Dasmeh and Serohijos, 2018; Diss and Lehner, 2018; Kemble et al., 2019; Pokusaeva et al., 2019), mutational robustness (Bershtein et al., 2013, 2006; Hartl et al., 1985; Lundin et al., 2018; Tokuriki and Tawfik, 2009) and evolvability (Dean, 1995; DePristo et al., 2005; Lunzer et al., 2005; Mehlhoff et al., 2020; Sarkisyan et al., 2016; Stiffler et al., 2015; Tokuriki and Tawfik, 2009). However, quantitative and experimental studies of the threshold model are rare, as most reports contain just a handful of data points and each landscape is confined to the scope of a single level of selection pressure (Bershtein et al., 2013; Dean, 1995; Hartl et al., 1985; Lunzer et al., 2005; Pokusaeva et al., 2019).

Consequently, detailed empirical models that can recapitulate the fitness landscapes of proteins, let alone the phenotype-fitness relationship across varying environmental conditions, have never been established. Such models will enable the quantitative evaluation of the evolutionary dynamics in proteins, and the development of predictive models for protein evolution in the response to a given environmental selection pressure.

To describe a realistic phenotype-fitness landscape, we require a quantitative mapping of the phenotype-fitness relationship with careful examinations in the context of environmental factors. Recent advances in largescale mutational characterization experiments, such as deep mutational scanning (DMS)(Fowler et al., 2014, 2010), have enabled the high-throughput measurement of protein fitness (growth of host organism carrying encoded protein variants) and phenotype (*e.g.*, antibiotic resistance level, protein binding and expression) for several thousands of mutational variants (Firnberg et al., 2014; Hietpas et al., 2011; Jacquier et al., 2013; Klesmith et al., 2017; Mavor et al., 2016; Melnikov et al., 2014; Noda-García et al., 2019; Pokusaeva et al., 2019; Starr et al., 2020, 2017; Stiffler et al., 2015; Wrenbeck et al., 2017). DMS across different conditions or genetic backgrounds can also determine higher dimensional characteristics of protein phenotype and/or fitness, such as antibiotic concentration dependent fitness(Firnberg et al., 2014; Melnikov et al., 2014; Stiffler et al., 2015), substrate specificity(Chen et al., 2020; Meer et al., 2016; Melnikov et al., 2014) and inter/intra-protein and protein-environment interactions (Diss and Lehner, 2018; Flynn et al., 2020; Horovitz et al., 2019; Kemble et al., 2020; Mavor et al., 2018,

2016; Noda-García et al., 2019; Salinas and Ranganathan, 2018; Starita et al., 2013; Thompson et
al., 2020).

In this work, we seek to obtain a global phenotype-environment-fitness landscape using
the VIM-2 β-lactamase as a model system, for which we previously generated all single point
mutants (~5600 variants) (Chen et al., 2020). To this end, we determined the fitness of an
Escherichia coli strain carrying VIM-2 variants across different environments dictated by seven
different ampicillin (AMP) concentrations (~39,000 data points). We also measured the phenotype,
specifically the antibiotic resistance level (EC_{50}), of each variant by integrating the experimental
data across all tested AMP concentrations into dose-response curves. We show that the phenotype-
fitness relationships exhibit a universal behavior according to AMP concentration. Taking
advantage of this result, we generate an empirical model for generalizing the phenotype-
environment-fitness landscape of VIM-2. We then demonstrate that this empirical and quantitative
model can explain the evolutionary behavior of the resistance gene at low antibiotic concentrations,
and the molecular mechanisms underlying the prevalence of nonspecific epistasis in mutational
datasets.

Results

Selection of VIM-2 variant libraries in a range of ampicillin concentrations

The overall workflow for our antibiotic selection experiment is presented in **Figure 1A**. We
previously generated a library containing ~5600 single amino acid variants of wild-type (*wt*) VIM-
2 in a vector that expresses VIM-2 through a constitutive ampR promoter (Chen et al., 2020). The
library was transformed into *E. coli* and subsequently grown in LB liquid media containing a range
of AMP concentrations (4.0-256 µg/mL AMP in two-fold increments), or in the absence of AMP,
for 6 hours at 37°C. The AMP concentrations used in this study covers the full dynamic range of
VIM-2's conferred resistance, from just above 2 µg/mL AMP where *E. coli* can grow without
VIM-2 expression (Chen et al., 2020), to more than double the EC_{50} of 84 µg/mL where *E. coli*
expressing *wt*VIM-2 can no longer grow. The concentration of cells in each culture after the
selection was measured via OD_{600} to track the total population size (**Figure 1B**), and the mutated
region of VIM-2 gene was amplified by PCR for deep sequencing (Chen et al., 2020). Two
replicates of the experiment with antibiotic selection and three replicates of growth without
selection were conducted.

Figure 1. Workflow for generating fitness scores and EC_{50} values using DMS. (**A**) Flowchart of the DMS process. A library of protein variants is first generated and transformed into host *E. coli*, which are then placed under selection with a range of antibiotic concentrations. The OD_{600} of the selected culture is recorded after selection and the selected culture is processed for deep sequencing. For each variant, the DNA frequency can be used to directly calculate the fitness scores, as well as the population size of each variant (N_{var}) after selection (using the OD_{600}). The population size can be used to generate a dose-response curve for each variant. (**B**) Cell density of a culture pool encompassing variants in the first 39 positions of VIM-2, measured after selection for each AMP concentration. (**C**) Example of a dose-response curve generated for the I19A variant. The solid line represents a fit to a sigmoidal curve. The red dashed line indicates the average non-selected population size of the replicate. Additional examples of dose-response curves are available in **Figure S2**. (**D**) Correlation of variant fitness scores between two replicates under selection with 64 µg/mL AMP. Correlation plots for other AMP concentrations are presented in **Figure S1**. For panels **D-F**, variants are colored by mutation type; the same color scheme applies to all panels. (**E**)

112   Correlation of EC_{50} values fitted from two replicates. (**F**) Correlation between EC_{50} determined by using
113   deep sequencing (y-axis) and or from isolated variants grown in separate liquid cultures (x-axis). See **Table**
114   **S1** for the EC_{50} values of isolated variants.

Calculation of fitness scores and EC_{50}

The output of the DMS experiments generates sequencing counts for each variant in the library,
which can be used to calculate the fitness scores (reflecting relative growth) at each AMP
concentration as well as the phenotype (EC_{50}) for each variant. The crux of our approach is that
the EC_{50} values are calculated independently from the fitness score in each condition, such that
their values are not interdependent, even when using the same set of deep sequencing data. Thus,
this allows us to examine the relationship between phenotype and fitness as separate factors under
each selection pressure.

The fitness (f) of a variant under each AMP concentration is calculated as the relative
growth of a variant compared to *wt*VIM-2 as we described previously (**Figure 1A**, **Table S2** for
all fitness scores)(Chen et al., 2020). For each variant, and at each level of AMP selection, an
enrichment score was calculated as the ratio of the frequency of a variant after selection to the
frequency without selection. Next, the fitness score of each variant was calculated by normalizing
the enrichment score of a variant to that of *wt*VIM-2, generating a measure of fitness relative to
*wt*VIM-2 using **equation (1)**.

130
$$f_{var,[AMP]} = \log_2\left(\frac{\frac{freq_{var\,sel}}{freq_{var\,nosel}}}{\frac{freq_{wt\,sel}}{freq_{wt\,nosel}}}\right) \tag{1}$$

The antibiotic resistance level of each variant was determined via a dose-response curve
by integrating the deep sequencing data from all seven different conditions (**Figure 1A-1C, Table**
S3 for all EC_{50} values). To this end, we first calculated the population size of cells expressing each
variant (N_{var}) at each antibiotic concentration by multiplying the frequency with the cell
concentration of the culture after antibiotic selection. Then we generated a dose-response curve
for each variant by plotting N_{var} against the antibiotic concentration, and calculated the EC_{50} by
fitting the sigmoidal relationship in **equation (2)**

138
$$N_{var} = N_{min} + \frac{N_{max} - N_{min}}{1 + \left(\frac{EC_{50}}{[AMP]}\right)^k} \tag{2}$$

139   Where N_{max} and N_{min} correspond to the maximum and minimum value of the variant population
140   size, N_{var}, [AMP] refers to the AMP concentration in µg/mL and k represents the Hill coefficient,
141   which determines the rate of change of the curve.

Quality of fitness scores and EC_{50} values

Next, we confirmed that our experiments yielded reliable fitness and EC_{50} values for the ~5600
variants across a range of AMP conditions. The fitness scores show a strong linear correlation
between replicates, with an R^2 of at least 0.92 for all conditions except at the highest AMP
concentration (256 µg/mL), which displayed an R^2 of 0.45 (**Figure 1D** and **S1**). The observed
noise arising at 256 µg/mL between replicates likely results from the use of an AMP concentration
much higher than the $EC_{50\text{-}wt}$ (84 µg/mL). At this high concentration, the frequency of the *wt*VIM-
2 after selection, which is the denominator in **equation (1)**, is drastically reduced and introduces
greater error in this condition. The fitness scores show distinct separation between functional and
non-functional variants. Variants with synonymous mutations are centered at a fitness score of 0

and the variants containing a stop codon mutation are centered near a fitness of -4 (**Figure 1D** and **S1**). Hence, a fitness score of -4 or lower denotes inactive variants, and scores that are below -4 are set to -4 in subsequent analyses.

We also found the EC_{50} values to strongly correlate between replicates, with a linear regression R^2 of 0.98 (**Figure 1E**). Moreover, the EC_{50} values derived from DMS are comparable with those measured on individual variants by a conventional dose-response curve resistance assay using the same selection method (87 unique variants), with a strong linear regression R^2 of 0.88 (**Figure 1F**).

Dependence of fitness on phenotype and selection pressure

The fitness scores we obtained across a range of AMP concentrations allows us to examine how the selection pressure influences the distribution of fitness effects (DFE). The fitness scores reflect relative growth of *E. coli* expressing a given VIM variant relative to *wt*VIM-2, and by extension, the resistance conferred to *E. coli* by a VIM variant in each AMP concentration. The resistance conferred by each variant is further dependent the EC_{50}, which reflects the variant's enzymatic efficiency (k_{cat}/K_M) of and/or the amount of enzyme expressed in the periplasm fraction of the cell. Thus, the DFE is inherently affected by both protein phenotype as well as the external selection environment.

Overall examination shows the DFEs and the distribution of EC_{50} values are bimodal (**Figure 2**), as typically observed in other proteins (Diss and Lehner, 2018; Firnberg et al., 2014; Fowler et al., 2010; Gray et al., 2018; Hietpas et al., 2011; Jacquier et al., 2013; Mavor et al., 2016; Pokusaeva et al., 2019; Rockah-Shmuel et al., 2015; Stiffler et al., 2015; Wrenbeck et al., 2017). In each distribution, one peak is formed by variants of near-neutral effects and a second peak is formed from variants with strongly deleterious effects, while only a small fraction of mutations exhibit higher values than *wt*. To further quantify and classify the variants in the distributions as having 'negative', 'neutral' or 'positive' effects with respect to the *wt*, we performed two-tailed z-tests between the fitness (or EC_{50}) of the non-synonymous variants against the distribution of the synonymous variants (a null model with N=251); test results are controlled for FDR by the

Benjamini-Hochberg procedure with $\alpha = 0.05$. The distribution of EC_{50} exhibits the most distinct separation between neutral and non-functional variants, where 63% are negative ($EC_{50} < 72$ µg/mL), 35% of mutations are neutral (72 µg/mL $\leq EC_{50} \geq 171$ µg/mL) and 2% are positive ($EC_{50} > 171$ µg/mL) (**Figure 2A**).

Figure 2. Distributions of fitness and EC_{50} effects. (**A**) The distribution of EC_{50} for all variants. (**B**) The DFEs of all variants under selection across all AMP selection concentrations. In all plots, the distributions are shown as a percentage of all variants within a mutation type (synonymous, missense or nonsense), with the color legend in **B**). The background colors indicate regions of positive (white), neutral (gray) and negative (hatched) effects, with the legend and a pie chart summary of the proportion of positive/neutral/negative effects below each histogram. For DFEs, the proportion of variants where positive/neutral/negative effects match the effects in the EC_{50} are displayed as percentages at the bottom. See **Table S2** for all fitness scores and **Table S3** for all EC_{50} values.

Naturally, fitness in the DFEs decline with increasing AMP concentration (**Figure 2B**), as can be inferred from the overall growth of the *E. coli* transformed with the library (**Figure 1B**). At lower AMP concentrations of 4.0-32 µg/mL, the majority (54-69%) of variants exhibit neutral effects. When the AMP concentration approaches two-fold of the $EC_{50\text{-wt}}$ of 84 µg/mL, the majority of variants become negative, up to 60%. At concentrations greater than $EC_{50\text{-wt}}$ (128-256 µg/mL), the distribution of synonymous mutations becomes broader as the *wt* resistance is also no longer sufficient to sustain high growth, and the data becomes noisier. In particular, at 256 µg/mL almost all variants are neutral again relative to *wt*VIM-2.

The shifting of the DFEs illuminates how mutational responses can be substantially altered depending on the selection pressure. For example, when comparing variant effects (negative/neutral/positive) across the DFEs and the distribution of EC_{50} values, only 52.3% of mutations exhibit the same effect across all distributions, with 26.7% consistently deleterious, 25.6% consistently neutral and none that are consistently positive. The remaining 47.7% of mutations exhibit distinct effects dependent on the selection pressure, where 39.5% switch between neutral and deleterious, 5.9% switch between positive and neutral, and 2.1% switch between all three states. We find that effects in the DFEs at 64-128 µg/mL mostly closely resemble the effects in EC_{50}, where 93% and 86% of variants have the same effect across fitness and EC_{50}, respectively (**Figure 2**); as these DFEs are measured at the closest concentrations to the $EC_{50\text{-wt}}$ of 84 µg/mL, the full extent of negative mutational effects is captured by the fitness scores without affecting the *wt* fitness.

Phenotype-fitness relationships across different AMP concentrations

Next, we seek to quantify the relationship between the EC_{50} values and the fitness scores obtained at each antibiotic concentration. We fit their relationship with a 'threshold' model (Bershtein et al., 2013; Chen et al., 2020; Dean, 1995; Hartl et al., 1985; Lunzer et al., 2005; Stiffler et al., 2015), *i.e.*, the four-parameter sigmoidal function in **equation (3)** (**Figure 3**).

$$f(EC_{50}) = f_{min} + \frac{f_{max} - f_{min}}{1 + \left(\frac{X_{50}}{EC_{50}}\right)^n} \tag{3}$$

The f_{min} and f_{max} denote the minimum and maximum fitness scores, respectively, while n represents the Hill coefficient. The inflection point of the curve, also the concentration of AMP where fitness is half-way between f_{min} and f_{max}, is denoted by X_{50}.

Figure 3. Phenotype-fitness landscapes across different AMP concentrations. Panels **A**-**G** show the comparison of DMS fitness scores with EC_{50} of all variants under selection in (**A**) 4.0 (N=5517), (**B**) 8.0 (N=5516), (**C**) 16 (N=5503), (**D**) 32 (N=5517), (**E**) 64 (N=5517), (**F**) 128 (N=5517) and (**G**) 256 (N=5517) μg/mL AMP. In each panel, variants are colored according to mutation type with the color legend in **A**). The solid black curve indicates the line of best fit for a sigmoidal curve fit (**equation 3**), while the black vertical dashed line indicates the AMP concentration used during selection. The dark grey regions extending to the x and y-axes indicates the region of linear association in the sigmoidal curve. The light grey regions extending to both axes indicate the range between 1% - 99% fitness in the fitted curve. The correlation coefficient and P-value of a Spearman rank-order correlation between the fitness score and EC_{50} is shown above each plot. (**H**) Sigmoidal fit parameters (**equation 3**) for the curves shown in panels **A**-**G**. The X_{50} values are shown in orange diamonds and the Hill coefficients (n) are shown in blue circles. The filled symbols indicate the conditions where the region of linear association in the sigmoidal curve (dark grey) was fully captured by the AMP concentrations of the experiment, while open symbols indicate the opposite. The R^2, P-value and equation for the linear regression of X_{50} with [AMP] is shown above the plot. Plots of all parameters can be found in **Figure S3**.

Overall, the threshold model recapitulates the phenotype-fitness relationship across all AMP concentrations, with Spearman rank-order correlation between fitness and EC_{50} ranging from

0.57-0.92 (**Figure 3A**-**3G**). Outside the sigmoidal transition range, fitness scores plateau at high and low fitness for variants with higher and lower EC_{50}, respectively. In particular, at AMP concentrations of 4.0-64 µg/mL (**Figure 3A**-**3E**), f_{min} and f_{max} maintain steady values in the fit at an average of -3.8 and 0.1, respectively (**Table 1**, **Figure S3A** and **S3B**). Since, wtVIM-2 can fully survive in these conditions, higher EC_{50} variants do not exhibit any fitness advantage over wtVIM-2. The existence of a fitness plateau for variants with an EC_{50} higher than the selection threshold also suggests that there are no systematic trade-offs between EC_{50} and fitness in our experimental system; higher EC_{50} variants do not exhibit low fitness in the presence of low AMP. At concentrations above EC_{50-wt} (128 and 256 µg/mL) the f_{max} increases as wtVIM-2 becomes challenged and exhibits decreased growth, while higher EC_{50} variants continue to grow (**Figure 3F** and **3G**).

The quantification of the phenotype-fitness relationships reveals some notable traits. First, the transitions of fitness between the lower and upper fitness plateaus (X_1- X_{99}, points in the curve at 1% and 99% fitness) (**Table 1**) occur in a relatively small range of EC_{50} differences (3.0-7.3 fold range) when considering conditions where the transition can be fully resolved within experimental conditions (8.0-64 µg/mL AMP). This confirms AMP selects the phenotype of VIM-2 variants in a discrete, threshold-like manner. Second, where the linear range can be fully captured (8.0-128 µg/mL AMP), the shape of the transitions occur in a very similar fashion across AMP concentrations; the Hill coefficient remaining steady with an average of 5.7 ± 1.8 S.D, leading to linear transition ranges that spans a 1.4 - 2.3 fold change in EC_{50}. (**Figure 3H, S3C** and **S3D**). This suggests that the mode of selection by AMP against VIM phenotypes is universal across a wide range of AMP concentrations. Third, the inflection points (X_{50}) in the phenotype-fitness relationship are a nearly perfect function of AMP concentration as described by the linear relationship in **equation (4)** (**Figure 3H**).

$$X_{50}([AMP]) = 0.93 \times [AMP] - 1.29 \qquad (4)$$

Taken together, the selection of VIM-2 phenotypes is strongly dictated by the concentration of AMP in the environment, and the range of phenotypic selection is largely predictable across a defined range of concentrations.

Table 1. Fitted parameters of sigmoidal phenotype-fitness relationships

[AMP] (µg/mL)	Equation fit [a]				Linear range [a]				Transition range [a]			
	f_{max}	f_{min}	n	X_{50} (µg/mL)	EC_{50} lower (µg/mL)	EC_{50} upper (µg/mL)	f lower	f upper	X_1 (µg/mL)	X_{99} (µg/mL)	$f_{1\%}$	$f_{99\%}$
4	0.0	-3.5	788	4.0	4.0	4.0	-2.9	-0.6	4.0	4.0	-3.5	0.0
8	0.0	-3.8	8.3	6.6	5.5	7.9	-3.1	-0.6	3.8	11.4	-3.8	0.0
16	0.0	-3.8	5.4	13.9	10.5	18.5	-3.1	-0.6	6.0	32.4	-3.8	0.0
32	0.1	-3.8	4.6	26.9	19.2	37.5	-3.1	-0.6	9.9	72.7	-3.8	0.1
64	0.2	-3.9	6.6	58.9	46.6	74.5	-3.2	-0.5	29.3	118	-3.9	0
128	4.3	-3.7	3.6	117	76.7	179	-2.3	2.9	33.2	415	-3.6	4
256	18.0	-2.7	2.9	256	149	439	0.9	14.4	51.4	1280	-2.5	18

[a] All parameters shown are calculated from a fit to equation (3).

A phenotype-environment-fitness landscape of VIM-2

Utilizing the universal behavior of the Hill coefficient (n) and the inflection points (X_{50}) in the phenotype-fitness relationships, we propose a simple, empirical model to describe the complete phenotype-fitness landscape of VIM-2 under AMP selection in **equation (5)**.

$$f(EC_{50}, [AMP]) = \frac{1}{1+\left(\frac{X_{50}([AMP])}{EC_{50}}\right)^{5.7}} \quad (5)$$

This equation describes the fitness (f) of a variant as a function of its EC_{50} and the AMP concentration. All parameters are functionally the same as **equation (3)**, but $n = 5.7$, and the inflection point of the curve (X_{50}) is now a linear function of [AMP] as shown in **equation (4)**. Furthermore, we normalize the observed fitness scores to the f_{min} and f_{max} determined from the sigmoidal fits at each AMP concentration in **Table 1**, such that a fitness of 0 represents the minimum fitness and a fitness of 1 represents the maximum fitness (**Figure 4A**). The model enables us to depict the entire phenotype-environment-fitness landscape, which shows agreement with ~39,000 data points across seven different selection conditions (**Figure 4B**). Thus, the universal description by a simple mathematical model suggests that the selection of VIM variants is highly dictated by the environmental AMP concentration, and that the evolutionary response of any given variant can be rather predictable.

Figure 4. The VIM-2 Phenotype-fitness landscape as a function of EC_{50} and AMP concentration. (**A**) For each variant, the observed fitness score is plotted in relation to the AMP concentration during selection, and its respective EC_{50}. The fitness scores are normalized at each concentration according to **equation (13)** in the Methods. The black dots represent the fitness of *wt*VIM-2 after normalization. (**B**) The modelled behavior of fitness scores according to **equation (5)**. The dots are the same data as panel **A**) while the surface depicts the predicted fitness scores according to the model. The black line indicates the model fitness at the $EC_{50\text{-wt}}$. Cross sections of the model at two-fold intervals of EC_{50} or AMP concentrations are marked by grey lines.

 Insight into the evolutionary dynamics of sub-MIC selection

Our global phenotype-environment-fitness landscape allows us to gain quantitative insights into
an important question regarding the evolution of antibiotic resistance: a phenomenon known as
sub-MIC selection. Antibiotic concentrations well below the 'minimal inhibitory concentration'
(MIC) of the *wt* in the population can still promote the emergence of higher resistance variants
(Fröhlich et al., 2020; Gullberg et al., 2011; Jasinska et al., 2020; Liu et al., 2011; Westhoff et al.,
2017). Typically, any resistant variant with MIC_{var} higher than MIC_{wt} will have an absolute
resistance and fitness advantage when the antibiotic concentration is over *MICwt* (*i.e.*, when MIC_{wt}
< [AMP] < MIC_{var}) (Andersson and Hughes, 2014; Baquero et al., 1998; Blondeau, 2009; Drlica
and Zhao, 2007; Gullberg et al., 2011) (**Figure 5A**). In sub-MIC selection where the antibiotic
concentration is below *MICwt*, the *wt* is still viable, but the resistance difference could still lead to
the selection of the variant over the *wt*, provided the benefit of resistance is high enough to offset
any trade-offs due to fitness costs (Andersson and Hughes, 2014; Gullberg et al., 2011). The lowest
antibiotic concentration where the variant can overtake the *wt* in fitness is designated the 'minimal
selective concentration' or MSC. Quantitative knowledge supporting existing models of sub-MIC
selection remains rare, however (Das et al., 2020; Gullberg et al., 2011).

312
Figure 5. Sub-MIC selection behavior of VIM-2 variants. (**A**) The fitness of *wt*VIM-2 and a hypothetical
variant with $2 \times EC_{50\text{-}wt}$ as a function of AMP concentration at a fixed EC_{50}. The MIC ($f = 0.01$) of both curves
and the MSC ($f_{wt} = 0.99$) are marked as vertical lines. The region in light grey is the 'mutant selection
window', the region in dark grey is the 'sub-MIC selection window' and the hatched region indicates no
selection of the variant over wt. (**B**) The difference in fitness between *wt*VIM-2 and higher resistance
variants across different AMP concentrations. Each curve indicates a variant with an EC_{50} of 1-8 fold relative
to *wt*VIM-2. The coloring of the sub-MIC selection range in the background is the same as in **A**).

Our model can bridge this gap by providing a quantitative depiction of the mechanism
underlying sub-MIC selection. For example, *wt*VIM-2, with EC_{50} of 84 µg/mL, would exhibit an
MIC (1% of f_{max}) of 203 µg/mL AMP, and an MSC (99% of f_{max}) of 42 µg/mL (see '**Calculation**
of MIC and MSC' in methods). Hence, for *wt*VIM-2, the range for sub-MIC selection can extend
to ~5-fold below the MIC*wt*. Furthermore, we can calculate the fitness difference between *wt*VIM-
2 and variants exhibiting higher resistance (**Figure 5B**). We observe that selection can still strongly
favor the higher resistance variant ($f_{var} - f_{wt} > 0.1$) until the AMP concentration drops to ~2-fold

above the MSC (**Figure 5B**). This is the case even when the variant is only 1.5 to 2-fold more
resistant than the *wt*, with the fitness difference being more pronounced for variants with higher
resistance. Thus, AMP concentrations as low as 5-fold below the *wt* MIC can still select for
variants with higher resistance than *wt*VIM-2, which is largely consistent with studies of sub-MIC
selection using other distinct systems (Fröhlich et al., 2020; Gullberg et al., 2011; Jasinska et al.,
2020; Liu et al., 2011; Westhoff et al., 2017).

The molecular basis for nonspecific epistasis

Our empirical model can also provide a quantitative understanding of the mechanisms underlying
mutational epistasis, *i.e.*, the non-additive interaction between mutational effects (Domingo et al.,
2019; Starr and Thornton, 2016). Epistasis is a key phenomenon in protein evolution that dictates
the shape and ruggedness of fitness landscapes, which subsequently affect important evolutionary
phenomenon such as robustness (Bershtein et al., 2013, 2006; Hartl et al., 1985; Lundin et al.,
2018; Tokuriki and Tawfik, 2009), evolvability (Dean, 1995; DePristo et al., 2005; Lunzer et al.,
2005; Mehlhoff et al., 2020; Sarkisyan et al., 2016; Stiffler et al., 2015; Tokuriki and Tawfik, 2009;
Yang et al., 2019) and evolutionary contingency (Baier et al., 2019; Starr et al., 2017; Starr and
Thornton, 2016). One form of epistasis, defined as 'nonspecific epistasis', is known to be highly
prevalent and strongly influences the accessibility of mutations. Nonspecific epistasis typically
stems from the non-linear relationship between fitness and protein phenotype, such as we observed
between fitness and EC_{50} in this study (**Figure 3**). The key distinction of nonspecific (as opposed
to specific) epistasis is that epistasis can still be observed at the level of fitness even if the
mutational effects of two mutations on a phenotype, *i.e.*, EC_{50} (and thus enzymatic efficiency or
protein expression) are non-epistatic (changes relative to *wt* are log-additive, or more simply
described as 'additive') (Bank et al., 2015; Bershtein et al., 2006; Dasmeh and Serohijos, 2018;
DePristo et al., 2005; Diss and Lehner, 2018; Domingo et al., 2019; Starr and Thornton, 2016).
While the existence of nonspecific epistasis is widely appreciated, we lack quantitative
understanding of the extents to which different types (magnitude/sign) of epistasis can arise
depending on the selective environment.

To this end, we simulated the prevalence and types of nonspecific epistasis in VIM-2
fitness using the empirically measured distribution of EC_{50} and the phenotype-environment-fitness
landscape (**Figure 6**). Since each variant only contains a single mutation, we can treat the variants'
effects directly as mutational effects. Specifically, we randomly sampled 200 single point missense
mutations, calculated the ΔEC_{50} values of a given mutation from *wt* (**Figure 6B**), then combined
all pairwise mutations based on a null model (**Figure 6C**)—a double mutational effect on EC_{50} is
equal to the additive effect of two single mutations. All pairwise combinations of 200 mutations
leads to $_{200}C_2 = 19900$ mutant pairs. Then, using the model in **equation (5)**, we computed the
change in fitness (Δf) for all individual and pairwise combinations at each AMP concentration
using the single and pairwise EC_{50}. We quantified epistasis by comparing the observed fitness
change (Δf_{obs}) calculated from the pairwise EC_{50}, and the expected fitness change (Δf_{exp}), which is
the sum of the Δf of both individual mutations (**Figure 6D**, See methods for '**Nonspecific epistasis**
of VIM-2 variants'). Finally, we categorized the observed epistasis into different types, based on
the magnitude and sign of the observed fitness change compared to the expected additive change:
First, magnitude epistasis, where Δf_{obs} has the same sign as Δf_{exp} but has significantly greater
(positive magnitude epistasis) or lesser (negative magnitude epistasis) magnitude. Second, sign
epistasis, where Δf_{obs} has an opposite sign to Δf_{exp} (**Figure 6A**).

Figure 6. Nonspecific epistasis varies across selection conditions. (**A**) Conceptual illustration of each type of epistasis considering an overall positive (left) or negative (right) additive effect. This given color scheme for epistasis types is maintained throughout the figure. (**B**) Distribution of the 200 randomly sampled EC_{50} values, with the change in EC_{50} relative to *wt*VIM-2 displayed on the right y-axis. (**C**) Distribution of the 19900 pairwise additive EC_{50} values, with the change in EC_{50} relative to *wt*VIM-2 displayed on the right y-axis. (**D**) The proportion of mutational pairs exhibiting epistatic effects under each condition; only proportions higher than 0.5% are labelled. Below the pie charts are scatterplots from key conditions showing the difference between expected and observed fitness changes. The backgrounds of the scatterplots are also colored according to the type of epistatic effect. A gray background indicates no significant epistasis, or additivity, and is defined as a range of 1.96 × S.D. of the synonymous mutant fitness

distribution after normalization by **Equation (13)** for each AMP concentration. In each scatterplot, points representing the combined effect of a pair of mutations are colored according to the effect of each single mutation (red when both increase EC_{50} over wtVIM-2, blue when both decrease EC_{50} and black when one increases while the other decreases EC_{50}). Histograms of all data points in each scatterplot are shown for each axis. (**E**) Visual depictions of each type of epistasis at distinct AMP concentrations, as a result of a non-linear relationship between fitness and EC_{50}.

Intriguingly, all types of epistasis can be found in our sample set, and the proportion of each type is drastically dependent on the environment (0 - 26% epistasis). At the lowest AMP concentration (4 µg/mL), no epistasis can be observed as most mutations simply exhibit neutral effects. At 8 µg/mL AMP, the greatest proportion of magnitude epistasis is observed, with 20% of mutational combinations exhibiting negative magnitude epistasis and about 4% of combinations showing positive magnitude epistasis. At 16 and 32 µg/mL AMP, smaller fractions of positive and negative magnitude epistasis are observed (7.3% and 8.3% in total, respectively). However, from 64-256 µg/mL AMP, a substantial fraction of mutational combinations exhibit sign-epistasis. A small fraction of positive sign epistasis are observed at 64 µg/mL AMP, and negative sign epistasis is observed at 128 and 256 µg/mL AMP. In particular, a striking 26% of mutational combinations exhibit negative sign epistasis at 128 µg/mL AMP. It should be emphasized that these substantially different epistasis patterns observed in different environments are based on the same set of mutational combinations within the experimentally obtained distribution of EC_{50}, even with non-epistatic effects on the phenotypic level (EC_{50}); such patterns are the results of a non-linear mapping between phenotype and fitness.

An examination of the non-linear relationship can provide the mechanistic explanation underlying the constraints (**Figure 6E**). Positive magnitude epistasis only occurs in combinations of mutants that both increase EC_{50} or decrease EC_{50} over wtVIM-2, where either mutation alone exhibits a weak effect but the combined additive change in EC_{50} can traverse the steep fitness transition in the landscape and cause a much stronger effect. In contrast, negative magnitude epistasis only occurs in mutant combinations of opposite effects (one increases while the other decreases EC_{50}), as only mutations of opposing effects can diminish the combined additive EC_{50} change. Sign epistasis is only present near AMP concentrations close to the EC_{50-wt} of 84 µg/mL, and occurs only between mutant combinations of opposite effects; sign epistasis cannot occur when the wtVIM-2 is on a high or low fitness plateau, where fitness changes can only occur in one direction (*e.g.*, fitness can only decrease when on the high fitness plateau). Positive sign epistasis is only observed when [AMP] is below EC_{50-wt} while negative sign epistasis only occurs when [AMP] is above EC_{50-wt} (**Figure 6D**).

Taken together, using the empirical distribution of mutational effects and the threshold model that we have established, we demonstrated that the selective environment strongly affects the resulting epistasis. We have also illustrated how epistasis can be prevalent in largescale mutational studies using fitness measurement (Araya et al., 2012; Bank et al., 2015; Diss and Lehner, 2018; Fowler et al., 2010; Olson et al., 2014; Palmer et al., 2015; Pokusaeva et al., 2019; Sarkisyan et al., 2016), as epistasis can still arise even when the underlying phenotype simply experience an additive change (Dasmeh and Serohijos, 2018). As the trend of apparent epistasis can be drastically different according to the environment, an understanding of the environmental factors is essential to decipher the evolutionary dynamics of proteins. Note that, in this model, we assumed additive effects on EC_{50}, while empirical mutational combinations are likely to exhibit non-additive effects with respect to protein function and stability due to specific epistasis (Domingo et al., 2019; Miton and Tokuriki, 2016; Starr and Thornton, 2016). In such cases,

428 epistasis will be even more frequently observed, and a rigorous model such as ours may help to
429 distinguish between nonspecific and specific epistasis within high-throughput mutational
430 experiments.

Discussion

In this study, we have presented a phenotype-environment-fitness landscape of VIM-2 β-lactamase,
constructed using a comprehensive set of ~39,000 data points, across a 64-fold range of AMP
concentrations. Our observations provide a long-sought, quantitative verification of an important
model for protein evolution, the 'threshold' model. With this comprehensive and empirical dataset,
we unveil how the phenotype-fitness relationship can be shaped by environmental variation,
particularly the selection strength. Importantly, we demonstrate that a simple quantitative model
can recapitulate the behavior of VIM-2 β-lactamase within the landscape. Our findings provide
evidence that the evolution of resistance in a population follows a relatively simple and predictable
rule, dictated through the selection pressure exerted by β-lactam antibiotics. Moreover, we provide
a mechanistic understanding of how different environments can cause different types of epistasis
to arise, solely from knowledge of the threshold relationship between protein phenotype and fitness.
Our observations suggest that the influence of the environment on the phenotype-fitness
relationship is extremely important to understand the evolutionary dynamics of proteins (Das et
al., 2020; Steinberg and Ostermeier, 2016). Hence, strong understanding of the effect of
environmental factors may allow us to accurately predict the evolution of antibiotic resistance
genes and other proteins.

Since we employed VIM-2, a class B β-lactamase, as a model for this study and measured
changes in EC_{50}—an important trait commonly measured trait for β-lactamases as well as other
antibiotic resistance genes—our model may be applicable to other β-lactamases, even in other
classes (A, C and D classes). Our findings could prove useful in developing a more general
understanding of β-lactamase evolution. Indeed, our observations agree with several behaviors
observed in β-lactamases, or other antibiotic resistance genes, beyond just the threshold-like
selection behavior. The ~5-fold range of sub-MIC selection estimated from our model largely
matches with other studies of the sub-MIC selection range in other systems (Fröhlich et al., 2020;
Gullberg et al., 2011; Jasinska et al., 2020; Liu et al., 2011; Westhoff et al., 2017). We do not
observe any notable, systematic trends that indicate trade-offs between high EC_{50} and fitness at
lower AMP concentrations, supporting observations that higher resistance does not necessarily
convey a large fitness cost, and that it can be maintained even under low selection pressure
(Gullberg et al., 2011; Westhoff et al., 2017). This also matches previous modelling where a flat
maximum fitness plateau is found for the mutants of TEM-1 irrespective of the selection pressure
(Stiffler et al., 2015). The lack of a fitness cost for more resistant variants is likely a result of the
non-essential nature of VIM-2 in our model system, coupled with a fairly moderate protein
expression level. Of course, when present, fitness trade-offs add greater complexity to the
dynamics of sub-MIC selection (Das et al., 2020; Gullberg et al., 2011). Regardless, our
quantitative descriptions for those phenomena are invaluable to further deepening our
understanding of the behaviors of these antibiotic resistance genes.

We note our relatively simple experimental system poses limits on the generalization of
our findings, and additional complexities may be studied in the future. *i*) We only mutated within
the VIM-2 gene, and mutations in other genetic context such as the promoter and origin of
replication of the plasmid might trigger different dynamics. *ii*) We conducted selection over 6
hours of continuous growth, up to the point where the cells are reaching the early phases of growth

saturation. Different culturing conditions may also expose different dynamics. *iii*) The results of AMP selection in *E. coli* may not reflect the behavior of VIM-2 with other classes of antibiotics, such as cephalosporins or carbapenems, as well as other clinically important bacterial hosts, *e.g.*, *P. aeruginosa* and *A. baumannii*. *iv*) These findings from a simple laboratory environment may only be partly applicable to behavior in natural environments, which will have additional variation outside of antibiotic concentration. *v*) Though we observed no systematic fitness-resistance trade-offs, higher resolution measurements of individual variants, and comparison to a condition without any selection may reveal trade-offs that are not as pronounced in our system (Gullberg et al., 2011; Jasinska et al., 2020; Mehlhoff et al., 2020; Westhoff et al., 2017; Zwart et al., 2018).

Nonetheless, we demonstrated that a quantitative mapping of the fitness landscape using deep sequencing is an extremely powerful method to study protein evolution. Similar studies for more complex conditions, and of different model proteins, will be key for further developing our understanding of protein evolution, in particular, the molecular mechanisms underlying key evolutionary phenomenon like mutational robustness, epistasis and evolvability. Ultimately, better knowledge of phenotype-fitness relationships will also improve our ability to predict the evolution of proteins, such as antibiotic resistance genes, in a complex, natural environment. Such knowledge should also lead to better design and engineering strategies to generate novel proteins and metabolic pathways.

Acknowledgements

We thank Adrian Serohijos, Pouria Dasmeh and members of the Tokuriki lab for insightful comments on the manuscript. We thank the Canadian Institute of Health Research (CIHR) Foundation Grant (FDN-148437) for the financial support. Nobuhiko Tokuriki is a Michael Smith Foundation of Health Research (MSFHR) career investigator.

Author Contributions

John Z Chen: Conceptualization, Software, Formal analysis, Investigation, Methodology, Writing - original draft, Writing - review and editing; Douglas M Fowler: Software, Validation, Methodology, Writing - review and editing; Nobuhiko Tokuriki: Conceptualization, Supervision, Funding acquisition, Project administration, Writing - review and editing

Declaration of Interest

The authors declare no competing interests.

Methods

Deep mutational scanning of VIM-2

The variant library of VIM-2 was generated in a previous study, and all selection experiments, deep sequencing and quality control for sequencing data were conducted in the same manner as before(Chen et al., 2020). Briefly, the plasmid based wild-type VIM-2 gene (*wt*VIM-2) gene was mutated into a library containing all single amino acid substitutions through PCR mutagenesis using a primer that replaces each codon position with a degenerate 'NNN' sequence. The library of all 267 codon positions was divided into 7 groups of 39 amino acids (31 in the last group), forming 117nt 'tiles' that can be sequenced with full overlapping reads using Illumina Next-seq 550. The libraries were transformed into *E. coli* E. cloni 10G electrocompetent cells (Lucigen) using the accompanying protocol, and grown in liquid culture (Miller LB broth) without selection, or selected with 4.0-256 µg/mL AMP in two-fold increments for 6 hours at 37°C. After 6 hours the cell concentration of the *E. coli* culture was measured by optical density at 600 nm (OD_{600}). The culture was then removed from selection by centrifugation and aspiration of the selection media, then resuspended in fresh LB; this procedure was repeated for a total of three washes. The washed cells were grown overnight, then the plasmid DNA was isolated using a QiaPrep 96 DNA extraction kit (Qiagen). One 'sample' in this experiment encompasses a single replicate of one library group of variants grown under a single condition. This process was conducted in two replicates on separate days, while an additional replicate (three total) was performed for the non-selected library. The third non-selected library was initially intended as a control to detect if variants in the library are lost during the 6 hours of growth without selection, and was grown overnight but was not subject to the 6 hours of growth at 37°C. However, we observed no notable differences in variants present between the three replicates of the non-selected library and the third replicate was treated the same as the other two replicates in subsequent analysis.

Following selection and DNA extraction, plasmid DNA from each library group was amplified by PCR for 15 cycles, using primers that amplify only the 117 bp region of each group and attach the universal Nextera adaptor sequence to each end of the amplicon. The amplicons are extended by PCR again to include the sample indices (i7 and i5) and flow cell binding sequence, then sequenced using a NextSeq 550 sequencing system (Illumina, Inc.). The *wt*VIM-2 plasmid was also included in the sequencing run to measure deep sequencing error rates. The raw sequencing data can be found in the NCBI Sequencing Read Archive (SRA) (BioProject accession: **PRJNA634597**). Subsequent processing of deep sequencing data and calculation of fitness scores are conducted using our previously published in-house Python scripts (https://github.com/johnchen93/DMS-FastQ-processing). Using these scripts after sequencing, the forward and reverse paired-end sequencing reads are merged and the posterior quality score of each position is calculate from the quality scores of the two reads (Edgar and Flyvbjerg, 2015). Sequences that have more than 20 mismatches between forward and reverse reads, or contains any position with a posterior quality score less than 10 were discarded. The median number of reads that pass the merging process is ~660,000. Only one of the samples from the non-selected library and 7 samples from the selected conditions had less than 100,000 reads after filtering (all limited to replicate 1), and these low read samples were handled accordingly in the subsequent applications (see sections below on '**Calculation of fitness scores**' and '**Population size estimates using deep sequencing data**').

The merged sequences were used to identify mutations, and sequences with more than one codon mutated were discarded. To further ensure the quality of the variants we identified, the

sequencing errors observed when sequencing *wt*VIM-2 was used to calculate the expected frequency of single amino acid variants arising from sequencing errors alone. Then, the non-selected library was filtered so that variants with frequencies lower than 2 × of the expected frequency from sequencing errors or have a read count less than 5 are excluded from further analysis. Only the variants that pass all of the above quality control measures are used for further calculations. The frequencies of each amino acid variant were calculated using **equation (6)** for each variant. Variants that passed quality filtering in the non-selected library but were no longer present in the selected libraries were given a dummy count of 1 to estimate the lowest possible frequency.

$$freq_{var} = read\ count_{var}\ /\ read\ count_{total} \tag{6}$$

Calculation of fitness scores

The variant frequencies were used to calculate fitness scores for selection conducted at each AMP concentration using **equation (1)**, where each selected library is matched with the non-selected replicate that was grown in the same replicate for calculations; the extra non-selected library replicate was not considered. Fitness scores were averaged by taking the arithmetic mean and averaged scores below -4.0 were set to a minimum of -4.0 for further analysis. The non-selected library in group 1 of replicate 1 contained less than 10,000 reads after merging and only ~25% of the variants were detected after filtering, meaning that most variants in positions 1-39 only have fitness scores from replicate 2. Additionally, of the 7 samples selected with AMP that contained less than 100,000 reads after FastQ merging, 2 samples were retained after scoring as they share correlation R^2 greater than 0.8 with the other replicate while the scores from the other 5 samples were excluded; this includes library group 2 (selected with 16 and 128 µg/mL AMP), 4 (8 µg/mL AMP), 6 (16 µg/mL AMP) and 7 (128 µg/mL AMP) from replicate 1.

Population size estimates using deep sequencing data

Population sizes were calculated using the variant frequencies and the OD_{600} of the library after selection. The code used to calculate the population size and EC_{50} can be found at https://github.com/johnchen93/DMS-EC50. To consolidate the 3 replicates of the non-selected library, the frequency of each variant was averaged between any replicates where the variant passed quality filtering; frequencies from selected libraries were kept as separate replicates. The frequencies of each variant were then used to estimate the population size of cells containing that variant by multiplying the frequency with the OD_{600} observed immediately after the 6 hours of selection, then multiplied again with an estimated cell count of 10^9 cells per 1 unit of OD_{600} using **equation (7)**; all cultures were selected at the same total volume of 1mL.

$$N_{var} = OD_{600} \times 10^9\ cells\ per\ OD_{600} \times freq_{var} \tag{7}$$

It is possible for the frequency to represent an unrealistic cell count, but in our case even purged variants that have been given a dummy count of 1 will still have population estimates of at least 300-500 cells. Samples with less than 100,000 reads after merging were excluded from this analysis, which consist of library groups 2 (at 16 and 128 µg/mL AMP), 4 (8 µg/mL AMP), 6 (8, 16 and 32 µg/mL AMP) and 7 (128 µg/mL AMP) in replicate 1.

Calculation of EC_{50} for each variant

The estimated population size of each variant (y-axis) is plotted against the AMP concentration (x-axis) to generate a dose-response curve. A sigmoidal fit is applied to the curve using the

scipy.optimize package's 'curve_fit' function to calculate the EC_{50} using **equation (2)**. In the initial fit, $\underline{N}_{max}$ is estimated as the average population size of the variant in the non-selected libraries or the library selected at the lowest AMP concentration, whichever is the higher value. N_{min}, k and EC_{50} were estimated as 0, -1 and 32μg/mL AMP, respectively. The fitting was performed separately for each replicate.

All dose-response curves were plotted regardless of whether a successful fit was made and each curve fit for each replicate was manually curated. Curves where the EC_{50} lies in a region with data missing in two consecutive concentrations are discarded. Some variants exhibit EC_{50} transitions outside the measured range of concentrations (4 - 256 μg/mL AMP), in which case the EC_{50} is estimated as the lowest (4 μg/mL AMP) and highest AMP concentration (256 μg/mL AMP) if the EC_{50} transition occurs past the lowest and highest AMP concentration, respectively. To merge the EC_{50} between replicates, we took the arithmetic average of the $\log_2 (EC_{50})$ (**Table S1**).

Measuring EC_{50} using conventional drug-response assays

Previously, we measured the EC_{50} for AMP of 39 amino acid variants that were isolated and assayed individually(Chen et al., 2020). In this study, we determined the EC_{50} of an additional 48 amino acid variants (63 unique codons) and measured their EC_{50} using the same method. Variant libraries at positions across the length of the VIM-2 sequence (positions 5, 16, 42, 73, 96, 153, 162, 173, 201, 209 and 251) were transformed into *E. coli*, plated on LB-agar plates and single colonies were picked, cultured and stored in glycerol stocks in a 96 well plate; for each position, we picked 8 colonies. Two colonies each of *E. coli* transformed with *wt*VIM-2 or empty vector were also isolated in the same manner and placed in the same plate as controls. The identity of each isolated variant was confirmed through Sanger sequencing.

Antibiotic selection was conducted using the same procedure as the DMS selection presented above, where a liquid culture of the E. coli containing each isolated variant is grown for 6 hrs under selection with 2.0-256 ug/mL AMP at 2-fold increments, at 37°C. The OD_{600} of each culture is measured after selection and plotted against the AMP concentration to generate a dose-response curve. The curve was fitted with the sigmoidal **equation (2)** to calculate the EC_{50}, where OD_{600} was used directly in place of N_{var}, while N_{min} and N_{max} were estimated as 0 and the OD_{600} from growth without AMP, respectively. Final EC_{50} values of the variants picked in this study can be found in **Table S3**.

Phenotype-fitness relationship at single AMP concentrations

The fitness scores (y-axis) at each AMP concentration were plotted against the EC_{50} (x-axis). The relationship between fitness scores and EC_{50} at each AMP concentration was fitted to the sigmoidal curve in **equation (3)**. The initial estimates for f_{max}, f_{min} and n are 2.0, -4.0 and 1.0 respectively for all AMP concentrations. The initial estimate for X_{50}, the inflection point of the curve, was set to be the same as the AMP concentration from which the dataset was measured.

The boundaries of the transition between the lower and upper fitness plateaus (X_1-X_{99}) were calculated as the EC_{50} values at 1% and 99% fitness relative to the range between the fitted f_{max} and f_{min} of each AMP concentration.

The linear region of the sigmoidal curve for the DMS fitness scores was calculated using **equations (8-11)**, based on the final fitted values for each AMP concentration (Sebaugh and McCray, 2003).

633 $$Y_{lower} = f_{max} + (\frac{f_{min}-f_{max}}{1+1/4.6805}) \qquad (8) \qquad Y_{upper} = f_{max} + (\frac{f_{min}-f_{max}}{1+4.6805}) \qquad (9)$$

634 $$X_{lower} = X_{50}(\frac{f_{min}-Y_{lower}}{Y_{lower}-f_{max}})^{\frac{1}{n}} \qquad (10) \qquad X_{upper} = X_{50}(\frac{f_{min}-Y_{upper}}{Y_{upper}-f_{max}})^{\frac{1}{n}} \qquad (11)$$

635 In this case, X_{lower} and X_{upper} indicate the start and the end of the linear range in terms of EC_{50} while
636 Y_{lower} and Y_{upper} indicate the start and end of the linear range in terms of fitness score.

General model for the phenotype-fitness landscape of VIM-2

To generate the model, we start by connecting the sigmoidal behavior of the phenotype-fitness
relationships across different AMP concentrations into the general form in **equation (12)**.

$$f(EC_{50}, [AMP]) = f_{min} + \frac{f_{max} - f_{min}}{1+\left(\frac{X_{50}([AMP])}{EC_{50}}\right)^n} \qquad (12)$$

All parameters are functionally the same as **equation (3)** except the inflection point of the curve
(X_{50}), which is now a linear function of AMP as shown in **equation (4)**. The Hill coefficient (n) is
5.7, the average where the linear range of the sigmoidal transition was not limited by the selection
concentration (8-128 µg/mL). Furthermore, we normalize the observed fitness scores to the f_{min}
and f_{max} determined from the sigmoidal fits at each AMP concentration in **Table 1** using **equation**
(13), such that a fitness of 0 represents the minimum fitness and a fitness of 1 represents the
maximum fitness.

$$f_{obs\ norm} = \frac{f_{obs}-f_{min}}{f_{max}-f_{min}} \qquad (13)$$

This allows us to arrive at the final form of the model presented in **equation (5)**.

Calculation of MIC and MSC from the general model

We calculated the minimal inhibitory concentration (MIC) of wtVIM-2 as the AMP concentration
at which fitness is reduced to a threshold $t = 0.01$ (1% of f_{max}), according to **equation (14)** which
is simply a rearrangement of **equation (5)**.

$$[AMP] = \frac{\left[\left(\frac{1}{t}-1\right)^{\frac{1}{5.7}} \times EC_{50}+1.29\right]}{0.93} \qquad (14)$$

Similarly, we calculated the minimal selective concentration (MSC) as the AMP concentration at
99% fitness (t = 0.99) for wtVIM-2, such that no variant can exhibit more than a 1% increase in
fitness over wtVIM-2 at this concentration of AMP or lower.

Nonspecific epistasis of VIM-2 variants

We randomly sampled 200 missense variants, extracted their EC_{50} values, and generated all
pairwise mutational combinations ($_{200}C_2 = 19900$ mutant pairs). Since the analysis was focused on
the numerical behavior rather than the physical behavior of the enzyme, mutations from the same
residue positions were allowed to be combined, though this rarely occurred as only 88 out of 19900
combinations (~0.4%) were from combining the same positions. From these mutant pairs, we used
equation (5) to calculate the expected fitness effect due to additive changes in fitness relative to
wtVIM-2. First, the relative change in fitness for each single variant in the pair is calculated
according to **equation (15)**, then added together as shown in **equation (16)**.

$$\Delta f_i = f(EC_{50-i}, [AMP]) - f(EC_{50-wt}, [AMP]) \qquad i = 1,2 \qquad (15)$$

$$\Delta f_{exp} = \Delta f_1 + \Delta f_2 \qquad (16)$$

Since the model explicitly defines 0 as the minimum fitness and 1 as the maximum fitness, $\Delta f\ exp$ is limited to the maximum change that can occur within this range (-1 to 1).

We also calculate the observed fitness effect according to **equation (5)** as an additive effect in the $\log_2(EC_{50})$ change relative to wtVIM-2. First, the relative change in EC_{50} is calculated in **equation (17)** and the additive EC_{50} effects are used to calculate the new EC_{50} in **equation (18)**.

$$\Delta \log_2(EC_{50-i}) = \log_2(EC_{50-i}) - \log_2(EC_{50-wt}) \qquad i = 1,2 \qquad (17)$$

$$EC_{50-1,2} = 2^{(\log_2(EC_{50-wt}) + \Delta\log_2(EC_{50-1}) + \Delta\log_2(EC_{50-2}))} \qquad (18)$$

We limit $EC_{50\text{-}1,2}$ to be no lower than 2.0 (4.0 µg/mL untransformed), as an [AMP] of 2.0 µg/mL is not selective and is not accounted for by **equation (5)**. The observed fitness effect is then calculated according to **equation (19)**.

$$\Delta f_{obs} = f(EC_{50-1,2}, [AMP]) - f(EC_{50-wt}, [AMP]) \qquad (19)$$

To identify significant epistatic effects, we use the synonymous fitness distribution at each [AMP] as a measure of expected measurement noise. When $|\Delta f_{obs} - \Delta f_{exp}| < 1.96 \times$ S.D. of the synonymous fitness (after normalization by **equation (13)**), the effect of the mutational pair is considered not significantly epistatic. For mutational pairs with significant epistasis, they are classified as exhibiting magnitude epstasis if the Δf_{obs} and Δf_{exp} have the same sign, or classified as exhibiting sign epistasis if Δf_{obs} and Δf_{exp} have opposite signs. For magnitude epistasis, positive magnitude epistasis is defined as when $\Delta f_{obs} > \Delta f_{exp}$ and $\Delta f_{exp} > 0$ or $\Delta f_{obs} < \Delta f_{exp}$ and $\Delta f_{exp} < 0$, while negative magnitude epistasis is defined when $\Delta f_{obs} < \Delta f_{exp}$ and $\Delta f_{exp} > 0$ or $\Delta f_{obs} > \Delta f_{exp}$ and $\Delta f_{exp} < 0$. For sign epistasis, positive sign epistasis is defined as when $\Delta f_{obs} > 0$ and $\Delta f_{exp} < 0$, while negative sign epistasis is defined as when $\Delta f_{obs} < 0$ and $\Delta f_{exp} > 0$.

References

Andersson, D.I., Hughes, D., 2014. Microbiological effects of sublethal levels of antibiotics. Nature Reviews Microbiology 12, 465–478. https://doi.org/10.1038/nrmicro3270

Araya, C.L., Fowler, D.M., Chen, W., Muniez, I., Kelly, J.W., Fields, S., 2012. A fundamental protein property, thermodynamic stability, revealed solely from large-scale measurements of protein function. Proceedings of the National Academy of Sciences 109, 16858–16863. https://doi.org/10.1073/pnas.1209751109

Baier, F., Hong, N., Yang, G., Pabis, A., Miton, C.M., Barrozo, A., Carr, P.D., Kamerlin, S.C., Jackson, C.J., Tokuriki, N., 2019. Cryptic genetic variation shapes the adaptive evolutionary potential of enzymes. Elife 8. https://doi.org/10.7554/eLife.40789

Bank, C., Hietpas, R.T., Jensen, J.D., Bolon, D.N.A., 2015. A systematic survey of an intragenic epistatic landscape. Mol Biol Evol 32, 229–238. https://doi.org/10.1093/molbev/msu301

Baquero, F., Negri, M.C., Morosini, M.I., Blázquez, J., 1998. Selection of very small differences in bacterial evolution. Int Microbiol 1, 295–300.

Bershtein, S., Mu, W., Serohijos, A.W.R., Zhou, J., Shakhnovich, E.I., 2013. Protein Quality Control Acts on Folding Intermediates to Shape the Effects of Mutations on Organismal Fitness. Molecular Cell 49, 133–144. https://doi.org/10.1016/j.molcel.2012.11.004

Bershtein, S., Segal, M., Bekerman, R., Tokuriki, N., Tawfik, D.S., 2006. Robustness–epistasis link shapes the fitness landscape of a randomly drifting protein. Nature 444, 929–932. https://doi.org/10.1038/nature05385

Bershtein, S., Serohijos, A.W., Shakhnovich, E.I., 2017. Bridging the physical scales in evolutionary biology: from protein sequence space to fitness of organisms and populations. Curr. Opin. Struct. Biol. 42, 31–40. https://doi.org/10.1016/j.sbi.2016.10.013

Blondeau, J.M., 2009. New concepts in antimicrobial susceptibility testing: the mutant prevention concentration and mutant selection window approach. Vet Dermatol 20, 383–396. https://doi.org/10.1111/j.1365-3164.2009.00856.x

Chen, J.Z., Fowler, D.M., Tokuriki, N., 2020. Comprehensive exploration of the translocation, stability and substrate recognition requirements in VIM-2 lactamase. eLife 9, e56707. https://doi.org/10.7554/eLife.56707

Dalziel, A.C., Rogers, S.M., Schulte, P.M., 2009. Linking genotypes to phenotypes and fitness: how mechanistic biology can inform molecular ecology. Mol. Ecol. 18, 4997–5017. https://doi.org/10.1111/j.1365-294X.2009.04427.x

Das, S.G., Direito, S.O., Waclaw, B., Allen, R.J., Krug, J., 2020. Predictable properties of fitness landscapes induced by adaptational tradeoffs. eLife 9, e55155. https://doi.org/10.7554/eLife.55155

Dasmeh, P., Serohijos, A.W.R., 2018. Estimating the contribution of folding stability to nonspecific epistasis in protein evolution. Proteins: Structure, Function, and Bioinformatics 86, 1242–1250. https://doi.org/10.1002/prot.25588

Dean, A.M., 1995. A molecular investigation of genotype by environment interactions. Genetics 139, 19–33.

DePristo, M.A., Weinreich, D.M., Hartl, D.L., 2005. Missense meanderings in sequence space: a biophysical view of protein evolution. Nature Reviews Genetics 6, 678. https://doi.org/10.1038/nrg1672

Diss, G., Lehner, B., 2018. The genetic landscape of a physical interaction. eLife 7, e32472. https://doi.org/10.7554/eLife.32472

Domingo, J., Baeza-Centurion, P., Lehner, B., 2019. The Causes and Consequences of Genetic
    Interactions (Epistasis). Annu Rev Genomics Hum Genet 20, 433–460.
    https://doi.org/10.1146/annurev-genom-083118-014857

Drlica, K., Zhao, X., 2007. Mutant Selection Window Hypothesis Updated. Clin Infect Dis 44,
    681–688. https://doi.org/10.1086/511642

Edgar, R.C., Flyvbjerg, H., 2015. Error filtering, pair assembly and error correction for next-
    generation sequencing reads. Bioinformatics 31, 3476–3482.
    https://doi.org/10.1093/bioinformatics/btv401

Firnberg, E., Labonte, J.W., Gray, J.J., Ostermeier, M., 2014. A Comprehensive, High-
    Resolution Map of a Gene's Fitness Landscape. Molecular Biology and Evolution 31,
    1581–1592. https://doi.org/10.1093/molbev/msu081

Flynn, J.M., Rossouw, A., Cote-Hammarlof, P., Fragata, I., Mavor, D., Hollins, C., III, Bank, C.,
    Bolon, D.N., 2020. Comprehensive fitness maps of Hsp90 show widespread
    environmental dependence. eLife 9, e53810. https://doi.org/10.7554/eLife.53810

Fowler, D.M., Araya, C.L., Fleishman, S.J., Kellogg, E.H., Stephany, J.J., Baker, D., Fields, S.,
    2010. High-resolution mapping of protein sequence-function relationships. Nature
    Methods 7, 741–746. https://doi.org/10.1038/nmeth.1492

Fowler, D.M., Stephany, J.J., Fields, S., 2014. Measuring the activity of protein variants on a
    large scale using deep mutational scanning. Nat Protoc 9, 2267–2284.
    https://doi.org/10.1038/nprot.2014.153

Fröhlich, C., Gama, J.A., Harms, K., Hirvonen, V.H.A., Lund, B.A., Kamp, M.W. van der,
    Johnsen, P.J., Samuelsen, Ø., Leiros, H.-K.S., 2020. Cryptic β-lactamase evolution is
    driven by low β-lactam concentrations. bioRxiv 2020.12.01.404343.
    https://doi.org/10.1101/2020.12.01.404343

Gray, V.E., Hause, R.J., Luebeck, J., Shendure, J., Fowler, D.M., 2018. Quantitative Missense
    Variant Effect Prediction Using Large-Scale Mutagenesis Data. Cell Syst 6, 116-124.e3.
    https://doi.org/10.1016/j.cels.2017.11.003

Gullberg, E., Cao, S., Berg, O.G., Ilbäck, C., Sandegren, L., Hughes, D., Andersson, D.I., 2011.
    Selection of resistant bacteria at very low antibiotic concentrations. PLoS Pathog. 7,
    e1002158. https://doi.org/10.1371/journal.ppat.1002158

Hartl, D.L., Dykhuizen, D.E., Dean, A.M., 1985. Limits of adaptation: the evolution of selective
    neutrality. Genetics 111, 655–674.

Hietpas, R.T., Jensen, J.D., Bolon, D.N.A., 2011. Experimental illumination of a fitness
    landscape. Proc. Natl. Acad. Sci. U.S.A. 108, 7896–7901.
    https://doi.org/10.1073/pnas.1016024108

Horovitz, A., Fleisher, R.C., Mondal, T., 2019. Double-mutant cycles: new directions and
    applications. Current Opinion in Structural Biology, Cryo electron microscopy ●
    Biophysical and computational methods ● Biophysical and computational methods - Part
    B 58, 10–17. https://doi.org/10.1016/j.sbi.2019.03.025

Jacquier, H., Birgy, A., Nagard, H.L., Mechulam, Y., Schmitt, E., Glodt, J., Bercot, B., Petit, E.,
    Poulain, J., Barnaud, G., Gros, P.-A., Tenaillon, O., 2013. Capturing the mutational
    landscape of the beta-lactamase TEM-1. PNAS 110, 13067–13072.
    https://doi.org/10.1073/pnas.1215206110

Jasinska, W., Manhart, M., Lerner, J., Gauthier, L., Serohijos, A.W.R., Bershtein, S., 2020.
    Chromosomal barcoding of E. coli populations reveals lineage diversity dynamics at high

resolution. Nature Ecology & Evolution 4, 437–452. https://doi.org/10.1038/s41559-020-1103-z

Kaltenbach, M., Tokuriki, N., 2014. Dynamics and constraints of enzyme evolution. J. Exp. Zool. B Mol. Dev. Evol. 322, 468–487. https://doi.org/10.1002/jez.b.22562

Kemble, H., Eisenhauer, C., Couce, A., Chapron, A., Magnan, M., Gautier, G., Nagard, H.L., Nghe, P., Tenaillon, O., 2020. Flux, toxicity, and expression costs generate complex genetic interactions in a metabolic pathway. Science Advances 6, eabb2236. https://doi.org/10.1126/sciadv.abb2236

Kemble, H., Nghe, P., Tenaillon, O., 2019. Recent insights into the genotype–phenotype relationship from massively parallel genetic assays. Evolutionary Applications 12, 1721–1742. https://doi.org/10.1111/eva.12846

Klesmith, J.R., Bacik, J.-P., Wrenbeck, E.E., Michalczyk, R., Whitehead, T.A., 2017. Trade-offs between enzyme fitness and solubility illuminated by deep mutational scanning. Proceedings of the National Academy of Sciences 114, 2265–2270. https://doi.org/10.1073/pnas.1614437114

Liu, A., Fong, A., Becket, E., Yuan, J., Tamae, C., Medrano, L., Maiz, M., Wahba, C., Lee, C., Lee, K., Tran, K.P., Yang, H., Hoffman, R.M., Salih, A., Miller, J.H., 2011. Selective advantage of resistant strains at trace levels of antibiotics: a simple and ultrasensitive color test for detection of antibiotics and genotoxic agents. Antimicrob. Agents Chemother. 55, 1204–1210. https://doi.org/10.1128/AAC.01182-10

Lundin, E., Tang, P.-C., Guy, L., Näsvall, J., Andersson, D.I., 2018. Experimental Determination and Prediction of the Fitness Effects of Random Point Mutations in the Biosynthetic Enzyme HisA. Mol Biol Evol 35, 704–718. https://doi.org/10.1093/molbev/msx325

Lunzer, M., Miller, S.P., Felsheim, R., Dean, A.M., 2005. The Biochemical Architecture of an Ancient Adaptive Landscape. Science 310, 499–501. https://doi.org/10.1126/science.1115649

Mavor, D., Barlow, K., Thompson, S., Barad, B.A., Bonny, A.R., Cario, C.L., Gaskins, G., Liu, Z., Deming, L., Axen, S.D., Caceres, E., Chen, W., Cuesta, A., Gate, R.E., Green, E.M., Hulce, K.R., Ji, W., Kenner, L.R., Mensa, B., Morinishi, L.S., Moss, S.M., Mravic, M., Muir, R.K., Niekamp, S., Nnadi, C.I., Palovcak, E., Poss, E.M., Ross, T.D., Salcedo, E.C., See, S.K., Subramaniam, M., Wong, A.W., Li, J., Thorn, K.S., Conchúir, S.Ó., Roscoe, B.P., Chow, E.D., DeRisi, J.L., Kortemme, T., Bolon, D.N., Fraser, J.S., 2016. Determination of ubiquitin fitness landscapes under different chemical stresses in a classroom setting. Elife 5. https://doi.org/10.7554/eLife.15802

Mavor, D., Barlow, K.A., Asarnow, D., Birman, Y., Britain, D., Chen, W., Green, E.M., Kenner, L.R., Mensa, B., Morinishi, L.S., Nelson, C.A., Poss, E.M., Suresh, P., Tian, R., Arhar, T., Ary, B.E., Bauer, D.P., Bergman, I.D., Brunetti, R.M., Chio, C.M., Dai, S.A., Dickinson, M.S., Elledge, S.K., Helsell, C.V.M., Hendel, N.L., Kang, E., Kern, N., Khoroshkin, M.S., Kirkemo, L.L., Lewis, G.R., Lou, K., Marin, W.M., Maxwell, A.M., McTigue, P.F., Myers-Turnbull, D., Nagy, T.L., Natale, A.M., Oltion, K., Pourmal, S., Reder, G.K., Rettko, N.J., Rohweder, P.J., Schwarz, D.M.C., Tan, S.K., Thomas, P.V., Tibble, R.W., Town, J.P., Tsai, M.K., Ugur, F.S., Wassarman, D.R., Wolff, A.M., Wu, T.S., Bogdanoff, D., Li, J., Thorn, K.S., O'Conchúir, S., Swaney, D.L., Chow, E.D., Madhani, H.D., Redding, S., Bolon, D.N., Kortemme, T., DeRisi, J.L., Kampmann, M., Fraser, J.S., 2018. Extending chemical perturbations of the ubiquitin fitness landscape in

a classroom setting reveals new constraints on sequence tolerance. Biol Open 7. https://doi.org/10.1242/bio.036103

Meer, J.-Y. van der, Poddar, H., Baas, B.-J., Miao, Y., Rahimi, M., Kunzendorf, A., Merkerk, R. van, Tepper, P.G., Geertsema, E.M., Thunnissen, A.-M.W.H., Quax, W.J., Poelarends, G.J., 2016. Using mutability landscapes of a promiscuous tautomerase to guide the engineering of enantioselective Michaelases. Nat Commun 7, 1–16. https://doi.org/10.1038/ncomms10911

Mehlhoff, J.D., Stearns, F.W., Rohm, D., Wang, B., Tsou, E.-Y., Dutta, N., Hsiao, M.-H., Gonzalez, C.E., Rubin, A.F., Ostermeier, M., 2020. Collateral fitness effects of mutations. PNAS 117, 11597–11607. https://doi.org/10.1073/pnas.1918680117

Melnikov, A., Rogov, P., Wang, L., Gnirke, A., Mikkelsen, T.S., 2014. Comprehensive mutational scanning of a kinase *in vivo* reveals substrate-dependent fitness landscapes. Nucleic Acids Research 42, e112–e112. https://doi.org/10.1093/nar/gku511

Miton, C.M., Tokuriki, N., 2016. How mutational epistasis impairs predictability in protein evolution and design. Protein Sci. 25, 1260–1272. https://doi.org/10.1002/pro.2876

Noda-García, L., Davidi, D., Korenblum, E., Elazar, A., Putintseva, E., Aharoni, A., Tawfik, D.S., 2019. Chance and pleiotropy dominate genetic diversity in complex bacterial environments. Nat Microbiol 4, 1221–1230. https://doi.org/10.1038/s41564-019-0412-y

Olson, C.A., Wu, N.C., Sun, R., 2014. A comprehensive biophysical description of pairwise epistasis throughout an entire protein domain. Curr Biol 24, 2643–2651. https://doi.org/10.1016/j.cub.2014.09.072

Pál, C., Papp, B., Lercher, M.J., 2006. An integrated view of protein evolution. Nat. Rev. Genet. 7, 337–348. https://doi.org/10.1038/nrg1838

Palmer, A.C., Toprak, E., Baym, M., Kim, S., Veres, A., Bershtein, S., Kishony, R., 2015. Delayed commitment to evolutionary fate in antibiotic resistance fitness landscapes. Nat Commun 6, 7385. https://doi.org/10.1038/ncomms8385

Pokusaeva, V.O., Usmanova, D.R., Putintseva, E.V., Espinar, L., Sarkisyan, K.S., Mishin, A.S., Bogatyreva, N.S., Ivankov, D.N., Akopyan, A.V., Avvakumov, S.Y., Povolotskaya, I.S., Filion, G.J., Carey, L.B., Kondrashov, F.A., 2019. An experimental assay of the interactions of amino acids from orthologous sequences shaping a complex fitness landscape. PLOS Genetics 15, e1008079. https://doi.org/10.1371/journal.pgen.1008079

Rockah-Shmuel, L., Tóth-Petróczy, Á., Tawfik, D.S., 2015. Systematic Mapping of Protein Mutational Space by Prolonged Drift Reveals the Deleterious Effects of Seemingly Neutral Mutations. PLoS Comput. Biol. 11, e1004421. https://doi.org/10.1371/journal.pcbi.1004421

Salinas, V.H., Ranganathan, R., 2018. Coevolution-based inference of amino acid interactions underlying protein function. eLife 7, e34300. https://doi.org/10.7554/eLife.34300

Sarkisyan, K.S., Bolotin, D.A., Meer, M.V., Usmanova, D.R., Mishin, A.S., Sharonov, G.V., Ivankov, D.N., Bozhanova, N.G., Baranov, M.S., Soylemez, O., Bogatyreva, N.S., Vlasov, P.K., Egorov, E.S., Logacheva, M.D., Kondrashov, A.S., Chudakov, D.M., Putintseva, E.V., Mamedov, I.Z., Tawfik, D.S., Lukyanov, K.A., Kondrashov, F.A., 2016. Local fitness landscape of the green fluorescent protein. Nature 533, 397–401. https://doi.org/10.1038/nature17995

Sebaugh, J.L., McCray, P.D., 2003. Defining the linear portion of a sigmoid-shaped curve: bend points. Pharmaceutical Statistics 2, 167–174. https://doi.org/10.1002/pst.62

Starita, L.M., Pruneda, J.N., Lo, R.S., Fowler, D.M., Kim, H.J., Hiatt, J.B., Shendure, J.,
Brzovic, P.S., Fields, S., Klevit, R.E., 2013. Activity-enhancing mutations in an E3
ubiquitin ligase identified by high-throughput mutagenesis. Proceedings of the National
Academy of Sciences 110, E1263–E1272. https://doi.org/10.1073/pnas.1303309110
Starr, T.N., Greaney, A.J., Hilton, S.K., Ellis, D., Crawford, K.H.D., Dingens, A.S., Navarro,
M.J., Bowen, J.E., Tortorici, M.A., Walls, A.C., King, N.P., Veesler, D., Bloom, J.D.,
2020. Deep Mutational Scanning of SARS-CoV-2 Receptor Binding Domain Reveals
Constraints on Folding and ACE2 Binding. Cell 182, 1295-1310.e20.
https://doi.org/10.1016/j.cell.2020.08.012
Starr, T.N., Picton, L.K., Thornton, J.W., 2017. Alternative evolutionary histories in the
sequence space of an ancient protein. Nature 549, 409–413.
https://doi.org/10.1038/nature23902
Starr, T.N., Thornton, J.W., 2016. Epistasis in protein evolution. Protein Sci. 25, 1204–1218.
https://doi.org/10.1002/pro.2897
Steinberg, B., Ostermeier, M., 2016. Environmental changes bridge evolutionary valleys. Sci.
Adv. 2, e1500921. https://doi.org/10.1126/sciadv.1500921
Stiffler, M.A., Hekstra, D.R., Ranganathan, R., 2015. Evolvability as a Function of Purifying
Selection in TEM-1 β-Lactamase. Cell 160, 882–892.
https://doi.org/10.1016/j.cell.2015.01.035
Thompson, S., Zhang, Y., Ingle, C., Reynolds, K.A., Kortemme, T., 2020. Altered expression of
a quality control protease in E. coli reshapes the in vivo mutational landscape of a model
enzyme. eLife 9, e53476. https://doi.org/10.7554/eLife.53476
Tokuriki, N., Tawfik, D.S., 2009. Stability effects of mutations and protein evolvability. Current
Opinion in Structural Biology, Carbohydradtes and glycoconjugates / Biophysical
methods 19, 596–604. https://doi.org/10.1016/j.sbi.2009.08.003
Westhoff, S., van Leeuwe, T.M., Qachach, O., Zhang, Z., van Wezel, G.P., Rozen, D.E., 2017.
The evolution of no-cost resistance at sub-MIC concentrations of streptomycin in
Streptomyces coelicolor. ISME J 11, 1168–1178. https://doi.org/10.1038/ismej.2016.194
Wrenbeck, E.E., Azouz, L.R., Whitehead, T.A., 2017. Single-mutation fitness landscapes for an
enzyme on multiple substrates reveal specificity is globally encoded. Nature
Communications 8, 15695. https://doi.org/10.1038/ncomms15695
Yang, G., Anderson, D.W., Baier, F., Dohmen, E., Hong, N., Carr, P.D., Kamerlin, S.C.L.,
Jackson, C.J., Bornberg-Bauer, E., Tokuriki, N., 2019. Higher-order epistasis shapes the
fitness landscape of a xenobiotic-degrading enzyme. Nature Chemical Biology 15, 1120–
1128. https://doi.org/10.1038/s41589-019-0386-3
Yi, X., Dean, A.M., 2019. Adaptive Landscapes in the Age of Synthetic Biology. Mol. Biol.
Evol. 36, 890–907. https://doi.org/10.1093/molbev/msz004
Zwart, M.P., Schenk, M.F., Hwang, S., Koopmanschap, B., de Lange, N., van de Pol, L., Nga,
T.T.T., Szendro, I.G., Krug, J., de Visser, J.A.G.M., 2018. Unraveling the causes of
adaptive benefits of synonymous mutations in TEM-1 β-lactamase. Heredity 121, 406–
421. https://doi.org/10.1038/s41437-018-0104-z

Supplemental Figures

Figure S1. Replicate correlation of DMS fitness scores, Related to Figure 1. Each scatter plot shows the replicate fitness scores of all variants under selection in (**A**) 4.0, (**B**) 8.0, (**C**) 16, (**D**) 32, (**E**) 64, (**F**) 128 and (**G**) 256 µg/mL AMP. In each panel, variants are colored according to mutation type with the legend in the lower right. The solid black line indicate the line of best fit for a linear regression, with the R^2 and P-value of the regression above each plot.

Figure S2. Example dose-response curves for EC_{50} calculation based on deep sequencing data, Related to Figure 1. Each pair of plots represent the two replicate dose-response curves of a single variant in the library. The solid curve indicates the results of fitting to the sigmoidal **equation (2)**. The horizontal dashed line marks the average population size of the variant in the non-selected conditions. Dose-response curves shown here were randomly sampled from the entire pool of all variants with a successful fit.

Figure S3. Fit parameters for the phenotype-fitness relationship across AMP concentrations, Related to Figure 3. Each plot shows one of the final fitted value of a parameter in the four-parameter sigmoidal curve in **equation (5)** when fitted to the fitness scores and EC_{50} values from the AMP concentration indicated in the x-axis. (**A**) Plot for maximum fitness (f_{max}) (**B**) Plot for minimum fitness (f_{min}), (**C**) Plot for Hill coefficient (n) (**D**) Plot for the inflection point (X_{50}). For all plots, solid points represent AMP concentrations where the linear portion of the fit were not limited by the AMP concentration range, while hollow points indicate the opposite. For **D**), a linear regression was conducted with the solid data points. The black line indicates the line of best fit for a linear regression, and the R^2, P-value and the equation of the fit shown at the top.