Beyond the fish-*Daphnia* paradigm: testing the potential for *Neoplea striola* (Hemiptera: Pleidae) to cause a trophic cascade in subtropical ponds

Chase J. Rakowski[1], Mathew A. Leibold[2]

[1] Department of Integrative Biology, University of Texas, Austin, TX, USA
[2] Department of Biology, University of Florida, Gainesville, FL, USA

Corresponding Author:
Chase Rakowski[1]
2415 Speedway, PAT 506B, Austin, TX, 78712, USA
Email address: chaserak@utexas.edu

Abstract

Trophic cascades, or indirect effects of predators on non-adjacent lower trophic levels, are thought to pervade diverse ecosystems, though they tend to be stronger in aquatic ecosystems. Most research on freshwater trophic cascades focused on temperate lakes where *Daphnia* tend to dominate the zooplankton community, and these studies identified that *Daphnia* plays a key role in facilitating trophic cascades by linking fish to algae with strong food web interactions. However, *Daphnia* are rare or absent in most tropical and subtropical lowland freshwaters, and many invertebrate predators have received little attention in food web research despite being common and widespread. Therefore, we aimed to test whether trophic cascades are possible in small warmwater ponds where small invertebrates are the top predators and *Daphnia* are absent. We collected naturally occurring plankton communities from small fishless water bodies in central Texas and propagated them in replicate pond mesocosms. We removed zooplankton from some mesocosms, left the plankton community intact in others, and added one of two densities of the predaceous insect *Neoplea striola* to others. Following an incubation period we then compared biomasses of plankton groups to assess food web effects between the trophic levels including whether *Neoplea* caused a trophic cascade by reducing zooplankton. The zooplankton community became dominated by copepods which prefer large phytoplankton and exhibit a fast escape response. Perhaps due to these qualities of the copepods and perhaps due to slow consumption rates by *Neoplea* on key grazers, no food web effects were found other than zooplankton marginally reducing large phytoplankton. More research is needed to understand the behavior and ecology of *Neoplea*, but trophic cascades may generally be weak or absent in subtropical and tropical lowland freshwaters where *Daphnia* is rare.

Introduction

An extensive body of literature has demonstrated the importance of direct food web effects, as well as indirect food web effects, or trophic cascades, with much of this work focusing on freshwater ecosystems from an early time. Many of the studies on trophic cascades in freshwater pelagic ecosystems have focused on large-bodied cladocerans, especially *Daphnia*, as the herbivorous prey linking predators to autotrophs (e.g. Carpenter et al. 2001). However, a relative paucity of such studies were performed in tropical and subtropical lowlands, where *Daphnia* is rare (Dumont 1994). Furthermore, the predators that most of these studies focused on were fish, with certain invertebrate taxa such as *Chaoborus* or *Notonecta* receiving some attention while the food web effects of other common and widespread invertebrate predators remain little known (Carpenter et al. 1992, 2001). Understanding the effects of these understudied predaceous invertebrate species is important for understanding their role in ecosystems and for conservation planning, as predators are more threatened with extinction than lower trophic levels (Purvis et al. 2000).

Members of the family Pleidae are small but common heteropteran insects which are closely related to the family Notonectidae which includes *Notonecta* (backswimmers). Pleids have been much less studied than their larger cousins the notonectids (especially *Notonecta* and *Buenoa*). In this paper we specifically study the pleid *Neoplea striola* (hereafter, *Neoplea*) as our manipulated zooplanktivorous predator. *Neoplea* is a widespread inhabitant of lentic freshwater in Central and North America, and is known to tolerate low oxygen conditions (Gittelman 1975). It is a small insect, with adults measuring 1.5 mm in length. *Neoplea* is an active hunter which uses sight, vibrations, and possibly chemicals to sense its prey (Papacek 2001). They have been shown to attack and consume small zooplankton such as rotifers, mesozooplankton such as *Daphnia*, and even prey as large as dipteran larvae (Gittelman 1974, Hampton and Gilbert 2001, Papacek 2001). However, the effects of *Neoplea* on pond communities are still not fully understood, including whether they can cause trophic cascades.

Here we report the results of a field mesocosm experiment in which we manipulated densities of *Neoplea* and zooplankton to test the effects of both of these trophic groups on plankton composition and biomass, including direct effects on the next trophic level and indirect (trophic cascade) effects of *Neoplea* on phytoplankton. Our plankton communities were composed of freshwater plankton collected locally in central Texas, with no *Daphnia* or other large-bodied cladocerans present. Due to the fast escape response of copepods, we expected *Neoplea* would reduce the biomass of non-copepod zooplankton, especially cladocerans and ostracods. We further predicted that zooplankton would reduce the biovolume of total phytoplankton, and that *Neoplea* would cause a trophic cascade, i.e. indirectly increase total phytoplankton biovolume via its suppression of herbivorous zooplankton. However, we found no evidence for food web effects of *Neoplea* and only weak evidence for effects of zooplankton on phytoplankton in the experiment.

Materials & Methods

Organism collection

We allowed phytoplankton communities to naturally assemble in two plastic tanks at the University of Texas' Brackenridge Field Laboratory in Austin, TX for ~3 months. We then mixed 10 L from each tank with 12 L carbonated mineral water to narcotize any zooplankton, filtered this mixture through 45 µm mesh to remove the zooplankton, and mixed well to produce a phytoplankton inoculum. To create a zooplankton inoculum, we collected and mixed zooplankton-rich water from several small fishless water bodies nearby, and concentrated the mixture with a 45 µm mesh. We collected *Neoplea* from small water bodies in Austin, TX.

Experiment setup and design

We established 20 pond communities in 200 L cattle tanks in an unshaded field at
Brackenridge Field Laboratory. We fit the tanks with float valves to maintain constant water
levels and covered them in 1 mm^2 screens to prevent insect immigration. Prior to the experiment,
we analyzed total N and P in the water following standard American Public Health Association
methods (APHA 1989). We then added NaNO$_3$ to bring total N to 14 mg/L N and added
NaH$_2$PO$_4$•H$_2$O to bring total P to 1.55 mg/L P. These are the total N and P concentrations in
COMBO medium, a eutrophic medium commonly used for culturing plankton (Kilham et al.
1998). Every five or six days thereafter for six weeks, enough of both nutrients were added to
each tank to compensate for a 5% daily loss rate from the water column (as per Hall et al. 2004);
this same amount of both nutrients was also added immediately following the first sampling
(methods described below) after a 22-day pause.
106
We inoculated each tank with 600 mL of the phytoplankton inoculum. After allowing
phytoplankton to reproduce for five days, we added an equal volume of zooplankton inoculum to
15 of the tanks. To the other five tanks we added some of the filtrate left from concentrating the
zooplankton inoculum, to ensure all tanks received the same phytoplankton and picoplankton
species, as there may have been strains in the zooplankton inoculum not represented in the
phytoplankton inoculum. Lastly, after allowing the zooplankton to reproduce for 15 days we
added 20 *Neoplea* adults to five tanks with zooplankton and 40 *Neoplea* adults to another five,
then added the same amount to the same tanks two days later to bring the totals to 40 and 80.
Thus there were four treatments: no zooplankton added ("no zoop."), zooplankton but no
Neoplea added ("no *Neoplea*"), zooplankton and 40 *Neoplea* added ("40 *Neoplea*"), and
zooplankton and 80 *Neoplea* added ("80 *Neoplea*"). Each treatment was replicated five times for
a total of 20 mesocosms arranged randomly.
119
120 ***Sampling and biomass estimation***
121
We sampled zooplankton and phytoplankton 40 days after adding the *Neoplea*, and then
again six days later. To sample zooplankton, we used tube samplers to collect ten whole water
column subsamples spread across each tank, and pooled them into a 10 L sample for each tank.
We filtered this sample through 65 µm mesh, returning any predators to the tank, and preserved
the retained organisms in 10% Lugol's solution. To sample phytoplankton, we used PVC pipes
with 1 cm diameter (one per tank) to collect three whole water column subsamples spread over
the tank and pooled them into a 50 mL sample for each tank. We preserved these phytoplankton
samples in 10% Lugol's solution. We additionally estimated surviving *Neoplea* populations after
the second plankton sampling by using a dipnet to count individuals until we returned three
successive empty sweeps.
132
133   To estimate biomass of zooplankton taxa, we identified, counted, and measured
134 zooplankton in subsamples such that for each taxon, at least 25 individuals or 10% of the sample

was counted – whichever came first – and at least 50 total individuals were counted. We used an ocular micrometer to measure the length of each crustacean and individual *Spirostomum* to the nearest half increment (0.24 mm), and to measure the length and width of each rotifer and width of *Spirostomum* to the nearest 0.05 increment (0.024 mm). We converted crustacean length to dry mass using length-mass regressions (Culver et al. [1985] for copepods, McCauley [1984] for *Scapholeberis*, and Anderson et al. [1998] for ostracods). When rotifers were identified to species, we converted rotifer dimensions to dry mass using species-specific equations from the EPA protocol (EPA 2016). Otherwise, we used biovolume equations (McCauley 1984) and converted to dry mass assuming a 10:1 biovolume:dry mass ratio. We converted *Spirostomum* dimensions to dry mass by approximating cells as cylinders and assuming the same 10:1 biovolume:dry mass ratio.

To estimate biovolume of phytoplankton taxa, we calculated phytoplankton densities with a hemocytometer. We counted 25 nL or 50 cells of the most common morphospecies – whichever came first – and we counted at least 100 nL for the other taxa. We also took several micrographs of each morphospecies from various tanks and sampling dates and used ImageJ to measure the cell dimensions (Schneider et al. 2012), measuring at least 15 cells of all but the rarest morphospecies. We then used geometric approximations to calculate the biovolume of each cell.

Data analysis

Based on previous research (Rakowski et al. *under review*), we expected *Neoplea* to have differential effects on zooplankton groups based on taxonomy/behavior and size. Copepods have been shown to have faster escape responses than other zooplankton, so we analyzed their biomass separately. Since copepods dominated the zooplankton community, we analyzed the sum of all other zooplankton biomass ("non-copepods") as we predicted *Neoplea* would primarily affect these other taxa. We also separately analyzed the major groups of non-copepod zooplankton, including cladocerans and ostracods as one group (grouped for similarity in size, morphology, and slow swimming speed), *Spirostomum* as its own group as it is quite distinct from any other taxon in the community, and lastly the rotifers. We expected zooplankton to either affect the whole phytoplankton community, or alternatively to affect only a certain size class, either large or small morphospecies. Therefore we analyzed total phytoplankton biovolume as well as the summed biovolume of larger morphospecies and of smaller morphospecies.

We tested for a difference in *Neoplea* survival between the two densities using a t test. To analyze the effects of zooplankton and *Neoplea* additions on the biomass of plankton groups, we fit a generalized linear mixed model in the gamma family (gamma GLMM) for each plankton grouping using the *lme4* package (Bates et al. 2015). We included tank as a random effect to account for the repeated measures, and fixed effects for zooplankton addition and initial *Neoplea*

175 density. We also fit the nested models excluding *Neoplea* density as well as the null models with
176 neither *Neoplea* density nor zooplankton addition, and used likelihood ratio tests to assess
177 whether including either addition significantly improved model fit. All analyses were performed
178 in R v. 3.5.3 (R Core Team 2017).
179
Results
181
182   On average *Neoplea* survived at a rate of 74%, with no significant difference in survival
183 between the two densities (d.f. = 7.98, *t* = -0.518, *P* = 0.618). No evidence was found of *Neoplea*
184 reproduction during the experiment.
185
By the sampling dates, the zooplankton community had become dominated by copepods.
However, cladocerans, *Spirostomum*, and rotifers all individually composed at least 15% of the
zooplankton mass on average in one or more treatments (Fig. 1). A small amount of zooplankton,
mostly copepods, became established in the control tanks receiving no zooplankton (Fig. 1, Fig.
2). The zooplankton additions were successful in sustaining a significant increase in biomass of
all zooplankton groups (Table 1, Fig. 2). However, *Neoplea* additions had no significant effects
on the biomass of any zooplankton group (Fig. 2, Table 1). Mean copepod mass was higher and
the mean proportion of rotifers was lower the more *Neoplea* were added (Fig. 1), but neither
copepod nor rotifer mass was significantly affected by *Neoplea* (Fig. 2, Table 1).
195
The phytoplankton community became dominated by ovoid single-celled green algae,
with pennate diatoms contributing the next most biovolume in tanks with no zooplankton added
and *Oocystis* contributing the next most biovolume in tanks with 80 *Neoplea* added (Fig. 3).
While the average total phytoplankton biovolume with zooplankton was less than half of the
average without zooplankton, there was no significant effect of zooplankton addition on total
phytoplankton biovolume due to large variation within treatments (Fig. 3, Fig. 4A, Table 3A,
Table 4A). When the largest morphospecies were taken together, including the larger ovoid
chlorophytes, pennate diatoms, and *Oocystis*, zooplankton reduced their summed biovolume
marginally significantly by 77.9% (Fig. 4B, Table 3B, Table 4B). On the other hand, summed
biovolume of the smaller morphospecies (small ovoid chlorophytes, *Chlorella*, *Selenastrum*, and
photosynthetic picoplankton) was not significantly impacted by zooplankton addition (Fig. 4C,
Table 3C, Table 4C). Similarly, *Neoplea* addition had no significant effects on the biovolume of
any phytoplankton grouping (Fig. 4, Table 3).
209
Discussion
211
Our results show that *Neoplea* had no demonstrable effect on the plankton in this
experiment. Furthermore, the copepod-dominated zooplankton only caused a marginal reduction
of larger phytoplankton taxa. Any effect of *Neoplea* on phytoplankton composition or biomass

would have likely been mediated by an effect on zooplankton composition or biomass, as
Neoplea are carnivorous. Therefore it is unsurprising there was no indirect effect of *Neoplea* on
phytoplankton, or trophic cascade, considering there were only weak or nonexistent direct effects
between the three trophic levels.
219
There are several potential reasons why *Neoplea* did not reduce the biomass of any
zooplankton groups in our experiment. First, it is possible *Neoplea*'s predation rate was similar
enough to the reproduction rate of its prey that prey populations did not significantly change over
the course of the 46-day experiment, a scenario most likely if *Neoplea* preferred prey with short
generation times such as cladocerans. While a few studies observed predation by *Neoplea* on
various prey taxa, we are aware of only one study that measured the effect of *Neoplea* on the
biomass of prey taxa over time (Rakowski et al. *under revision*). In this study *Neoplea* had strong
effects on non-copepod zooplankton, which were dominated by fast-reproducing cladocerans, on
the short term (under one cladoceran generation). In contrast, *Neoplea* had weak effects on a
longer term (~six cladoceran generations) in the absence of other predators. Interestingly
however, when the larger predator *Notonecta* was also present, *Neoplea* acted synergistically
with *Notonecta* to suppress the non-copepod zooplankton. Therefore it may be that *Neoplea*
exhibits a slow predation rate and prefers prey with short generation times, with the effect that
over the long term it can have a cryptic food web effect not apparent unless other predators are
present.
235
Other potential reasons for the lack of an effect of *Neoplea* on zooplankton relate to other
aspects of the predaceous insect's behavior and to the composition of potential prey. It is
possible that *Neoplea* consumed chironomid larvae, which were common in the mesocosms but
not effectively sampled due to their benthic nature and our sampling methods targeting plankton.
Neoplea is known to attack and consume chironomid larvae (Papacek 2001). Indeed, the *Neoplea*
were most commonly observed clinging to the sides and bottom of the mesocosms, putting them
in close proximity to the chironomids. In nature, *Neoplea* is normally found clinging to
submerged vegetation (Gittelman 1974). However, it was impossible to observe the *Neoplea*
throughout most of the experiment due to the high density of phytoplankton. This high turbidity
may have made it difficult for the *Neoplea* to hunt effectively, as they can detect prey visually
and are most commonly found in clear waters, although they can also detect prey by tactile and
possibly chemosensory methods (Gittelman 1974, Papacek 2001). When the *Neoplea* did swim
through the water column, they would have mostly encountered copepods, the largest and
dominant group of zooplankton. However, even in clear water copepods are relatively resistant
to predation due to their fast escape response. Among copepods, diaptomids have an especially
fast escape response, and the diaptomid *Arctodiaptomus* dominated our mesocosms. Therefore it
may not be surprising that the *Neoplea* were unsuccessful in suppressing the zooplankton in the
experiment. While other more easily captured prey were also present, it may have been difficult
for the *Neoplea* to encounter these rare prey among all the copepods in a turbid environment.

The dominance of copepods in the zooplankton community may also explain the weak
effects of zooplankton on phytoplankton in the experiment. Unlike *Daphnia*, copepods do not
generally impose strong top-down control on community phytoplankton biomass, due in part to
their selective grazing on larger phytoplankton (Sommer and Sommer 2006). Indeed, the
copepod-dominated zooplankton only reduced larger phytoplankton, and even this effect was
marginal, fitting the general understanding of copepods' top-down effects on phytoplankton in
ecosystems with copepod-dominated herbivore communities such as open oceans (Sommer and
Sommer 2006). Such weak top-down control of crustacean zooplankton on phytoplankton may
be more pervasive at low latitudes, as *Daphnia* is largely absent from tropical and subtropical
lowland freshwaters and copepods are instead more likely to dominate (Dumont 1994, Havens
and Beaver 2011). In combination with the ability of copepods to evade predation much more
easily than *Daphnia*, it appears likely that trophic cascades mediated by zooplankton are less
common in warm, low-latitude lakes and ponds than in their colder counterparts (Rejas et al.
2005).

The food web ecology of *Neoplea* and pleids generally will remain unclear without
further research. More behavioral research on these diminutive predators is needed to better
understand their hunting habits, such as where in the habitat they make most of their captures,
their relative dependence on different senses for prey detection, and their relative preference for
various prey. Experimental work is needed to better quantify the ecology of pleids, and these
experiments will likely benefit from better replicating their preferred habitat of clear, still water
with submerged vegetation. A higher population density may be necessary for their food web
effects to be apparent when they are the sole predator.

Conclusions

We predicted that *Neoplea* would suppress non-copepod zooplankton, and that
zooplankton would suppress phytoplankton, resulting in a trophic cascade where *Neoplea*
indirectly increased phytoplankton biomass. Instead, *Neoplea* had no effect on plankton biomass
or composition in this field mesocosm experiment. The zooplankton, dominated by copepods and
lacking *Daphnia* as is typical of lowland tropical and subtropical lakes and ponds, only weakly
reduced larger phytoplankton. While our data cannot definitively explain these weak effects, they
could have resulted from *Neoplea* consuming prey able to reproduce quickly enough to make up
for losses to predation, from *Neoplea* consuming benthic prey which was not effectively
sampled, or from the dominance of copepods which are adept at evading capture and which
selectively feed on larger phytoplankton. This study suggests that lentic ecosystems dominated
by *Neoplea* and copepods may be characterized by weak top-down control. This represents
another example of a weak or non-existent trophic cascade in lowland tropical or subtropical
lentic freshwater, which appears to be much more common in these systems than in cooler lakes

and ponds. However, more research will be needed to achieve a clearer understanding of the
ecological impacts of pleids. A better understanding of the ecology of these and other
understudied invertebrate predators will be important for conservation planning, as predators are
more threatened with extinction than lower trophic levels, a pattern that does not only apply to
charismatic megafauna.
300

Acknowledgements

303 Thanks to L. A. Sekula for help processing samples, to R. Deans for help with insect collection
304 and experiment setup, to S. Duchicela for help with experimental setup, to D. Correa for help
305 with nutrient analysis, and to D. Nobles for providing equipment. This research was supported
306 by the Department of Integrative Biology at the University of Texas at Austin and was made
307 possible by the facilities at Brackenridge Field Laboratory.

References

Anderson, D. H., S. Darring, and A. C. Benke. 1998. Growth of Crustacean Meiofauna in a
Forested Floodplain Swamp: Implications for Biomass Turnover. Journal of the North
American Benthological Society 17:21–36.
Carpenter, S., J. Cole, J. Hodgson, J. Kitchell, M. Pace, D. Bade, K. Cottingham, T. Essington, J.
Houser, and D. Schindler. 2001, May. Trophic cascades, nutrients, and lake productivity:
whole-lake experiments. WILEY, 111 RIVER ST, HOBOKEN 07030-5774, NJ USA.
Carpenter, S., C. Kraft, R. Wright, H. Xi, P. Soranno, and J. Hodgson. 1992, November.
Resilience and resistance of a lake phosphorus cycle before and after food web
manipulation. UNIV CHICAGO PRESS, 5720 S WOODLAWN AVE, CHICAGO, IL
60637.
Culver, D. A., M. M. Boucherle, D. J. Bean, and J. W. Fletcher. 1985. Biomass of Freshwater
Crustacean Zooplankton from Length–Weight Regressions. Canadian Journal of
Fisheries and Aquatic Sciences 42:1380–1390.
Dumont, H. 1994, January 7. On the diversity of Cladocera in the tropics. SPRINGER, VAN
GODEWIJCKSTRAAT 30, 3311 GZ DORDRECHT, NETHERLANDS.
Dumont, H. J. (n.d.). On the diversity of the Cladocera in the tropics:12.
Gittelman, S. H. 1974. The Habitat Preference and Immature Stages of Neoplea striola
(Hemiptera: Pleidae). Journal of the Kansas Entomological Society 47:491–503.

Gittelman, S. H. 1975. Physical Gill Efficiency and Winter Dormancy in the Pigmy

Backswimmer, ISeoplea striola (Hemiptera: Pleidae). Annals of the Entomological

Society of America 68:1011–1017.

Hampton, S. E., and J. J. Gilbert. 2001. Observations of insect predation on rotifers. Pages 115–

121 *in* L. Sanoamuang, H. Segers, R. J. Shiel, and R. D. Gulati, editors. Rotifera IX.

Springer Netherlands, Dordrecht.

Havens, K. E., and J. R. Beaver. 2011. Composition, size, and biomass of zooplankton in large

productive Florida lakes. Hydrobiologia 668:49–60.

McCauley, E. 1984. The estimation of the abundance and biomass of zooplankton in samples.

Pages 228–265 *in* J. A. Downing and F. H. Rigler, editors. A Manual for the Assessment

of Secondary Productivity in Fresh Waters. Blackwell Scientific Publishers.

Papacek, M. 2001. Small aquatic and ripicolous bugs (Heteroptera: Nepomorpha) as predators

and prey: The question of economic importance. European Journal of Entomology 98:1–

12.

Purvis, A., J. L. Gittleman, G. Cowlishaw, and G. M. Mace. 2000. Predicting extinction risk in

declining species. Proceedings of the Royal Society of London. Series B: Biological

Sciences 267:1947–1952.

Rejas, D., S. Declerck, J. Auwerkerken, P. Tak, and L. De Meester. 2005, January. Plankton

dynamics in a tropical floodplain lake: fish, nutrients, and the relative importance of

bottom-up and top-down control. WILEY, 111 RIVER ST, HOBOKEN 07030-5774, NJ

USA.

Schneider, C. A., W. S. Rasband, and K. W. Eliceiri. 2012. NIH Image to ImageJ: 25 years of

image analysis. Nature Methods 9:671–675.

Sommer, U., and F. Sommer. 2006. Cladocerans versus copepods: the cause of contrasting top–

down controls on freshwater and marine phytoplankton. Oecologia 147:183–194.