Alterations to the Gut Microbiota of a Wild Juvenile Passerine in an Urban Mosaic

Running Title: Urban-related changes in tit microbiota

Öncü Maraci[1*], Michela Corsini[2], Anna Antonatou-Papaioannou[.3,4], Sebastian Jünemann[5,6], Joanna Sudyka[2,7], Irene Di Lecce[2], Barbara A. Caspers[1], Marta Szulkin[2]

Affiliations:

[1]Department of Behavioural Ecology, Bielefeld University, Konsequenz 45, 33619 Bielefeld, Germany.

[2]Centre of New Technologies, University of Warsaw, Banacha Street 2C, 02-097 Warsaw, Poland.

[3]Evolutionary Biology, Bielefeld University, Universitätsstrasse 25, 33615, Bielefeld, Germany.

[4]Institute of Biology-Zoology, Freie Universität Berlin, Köning-Luise-Str. 1-3, 14195, Berlin.

[5]Faculty of Technology, Bielefeld University, Universitätsstrasse 25, 33615, Bielefeld, Germany.

[6]Center for Biotechnology (CeBiTec), Bielefeld University, Sequenz 1, 33615, Bielefeld, Germany.

[7]Institute of Environmental Sciences, Jagiellonian University, ul. Gronostajowa 7, 30-387 Kraków, Poland.

*Corresponding author

Öncü Maraci

E-mail: oncu.maraci@uni-bielefeld.de

Abstract

Urbanisation is a major anthropogenic perturbation that progressively alters multiple environmental parameters, thereby presenting novel evolutionary challenges to wild populations. Symbiotic microorganisms residing in the gastrointestinal tracts (gut) of vertebrates have well-established mutual connections with the physiology of their hosts and respond quickly to environmental alterations. However, the impacts of anthropogenic changes and urbanisation on gut microbiota remain poorly understood, especially in early development. To address this knowledge gap, we investigated the gut microbiota in juvenile great tits (*Parus major*) in an urban mosaic. First, we compared the microbiota of nestlings reared in artificial nest boxes or natural cavities. Next, we analysed microbiota variations employing two distinct urbanisation frameworks, (i) the classical urban/rural dichotomy and (ii) gradual changes in the amount of impervious surface area (ISA) in the urban space and identified the environmental variables that predicted the changes in the microbial communities. Alpha diversity was influenced by cavity type and by the amount of impervious surface surrounding the breeding location. Alpha diversity was also positively correlated with the tree cover density. Beta diversity differed between urban and rural sites, and these alterations covaried with sound pollution, the distance to the city centre and the distance to the closest road. Overall, the microbial communities reflect and are possibly influenced by the heterogeneous environmental modifications that are typical of the urban space. Strikingly, the choice of framework used to define the urban space can influence the outcomes of studies investigating animal-microbe symbiosis. Our results open new perspectives from which the potential function of microbial symbionts in the adaptation of their hosts to anthropogenic stress can be investigated.

Keywords: avian gut microbiota, host-microbiome interactions, birds, great tit, urbanisation, anthropogenic change, early development

1. Introduction

In today's fast-changing world, cities are expanding at an unprecedented rate to accommodate over half of the human population (World urbanization Prospects 2018: Highlights, 2019). Urban growth modifies a wide range of biotic and abiotic ecosystem properties, thereby presenting new

ecological and evolutionary challenges for wild populations (Szulkin et al., 2020). Given the

scales and magnitudes of these environmental changes, adapting to human-altered environments

is a challenging task for animals, and their responses vary among species and individuals

(Pfennig et al., 2010): while some animals are outcompeted, others can survive and reproduce in

urban habitats. Populations living in urban spaces are often found to differ from their rural

counterparts in terms of their morphological, physiological, and behavioural traits, as well as in

terms of their ecological interactions with other organisms (Ditchkoff et al., 2006; Fischer and

Lindenmayer, 2007; Slabbekoorn, 2013; Alberti et al., 2017; Christiansen et al., 2019).

The radically transformed environment present in the urban space is consequently expected to

influence the symbiotic interactions between microorganisms and animals. The gut

microbiome—defined as "the entire gut habitat, including the whole community of

microorganisms (microbiota), their genomes and the surrounding environmental conditions"

(Marchesi and Ravel, 2015)—is increasingly recognized as a key player in several aspects of

host physiology (Zilber-Rosenberg and Rosenberg, 2008; McFall-Ngai et al., 2013). Indeed, as

urbanisation alters the distribution of multiple environmental variables (Szulkin et al., 2020) as

well as various traits of individual hosts, these changes altogether shape the assembly, taxonomic

diversity, and functioning of animal-associated microorganisms via different mechanisms

(Trevelline et al., 2019). For example, urbanisation can change the environmental pool of

microorganisms; this pool is a potential reservoir for the recurrent colonization of the gut

(Afshinnekoo et al., 2015). Via the removal of native plant or animal species (Miles et al., 2019)

or the introduction of anthropogenic food sources (Auman et al., 2008; Oro et al., 2013),

urbanisation can also alter the host diet (Jarrett et al., 2020), which is the main determinant of gut

microbiota (Ley et al., 2008; Kohl et al., 2014). Alternatively, urban-driven environmental

changes can exert their effects indirectly by altering host physiology. For instance, the extensive

use of artificial lights in urban settings disturbs the circadian rhythm (Ouyang et al., 2018).

Furthermore, increased competition, human disturbances, constant exposure to toxic substances,

urban noises and fluctuating microclimates can lead to chronic stress (Powell et al., 2014;

Isaksson, 2015; Zollinger et al., 2019). Chronic stress cascades downstream and has the potential

to impact immunity and metabolism, initiate oxidative stress and inflammatory responses

(Watson et al., 2017; Gadau et al., 2019) and consequently affect microbial communities (Hieke

et al., 2019; Jiang et al., 2020). Accordingly, independent of the mechanisms involved,

urbanisation has the considerable potential to re-shape several interactive pathways in animal-
microbe symbiosis.

An adequate definition of urban space is, however, essential when addressing this potential.
Classical frameworks describing urban space often rely on a threshold-based human density
value, often captured by a simple urban-rural dichotomy that overlaps with a city's
administrative borders (Szulkin et al., 2020). Accordingly, cities consist of a core area in which
the majority of the population lives (urban site) and peri-urban sites (rural) that accommodate
relatively smaller proportions of inhabitants. This dichotomous categorization can potentially
capture some alterations that are directly related to human activities and, consequently, change
predictably from the core of a city to the periphery, such as air pollution, sound pollution and
light pollution (Sprau et al., 2017). Nonetheless, the spatial units defined by humans rarely
consist of homogeneous habitats. Rather, cities are characterised by mosaics of urban landscapes
with varying degrees of actual surface use and accompanying fine-scale differences in habitat
types and several environmental characteristics that are not reflected in a simple urban/rural
categorization. In this light, defining the urban space by quantifying the actual changes in its
physical attributes, such as the percentage of impervious surface area (ISA) that characterises the
urban space, might be a more relevant approach biologically. Consequently, the methodological
approach used to characterize the urban space can potentially influence the outcomes of studies
investigating the impacts of urbanisation on ecological interactions.

To the best of our knowledge, no comparative studies have investigated whether the choice of
framework defining the urban space influences the outcomes or interpretations in the context of
animal-microbe symbiosis. Moreover, studies investigating rapid environmental changes often
focus on individual potent environmental variables, such as temperature change (Kohl and Yahn,
2016; Bestion et al., 2017; Fontaine et al., 2018), exposure to anthropogenic pollutants (Kohl et
al., 2015; Knutie et al., 2018; Trevelline et al., 2018) or human presence (Knutie et al., 2019). In
contrast, urbanisation manifests itself as the disturbance of several environmental parameters that
often covary (Szulkin et al., 2020) or interact with each other to restructure ecological
interactions. However, analyses of urban-driven shifts in animal-microbe symbiosis inferred in
the context of multiple urban environment dimensions are largely lacking to date. The lack of an
explicit framework characterizing the urban environment may explain why the evidence for
urban-driven shifts in avian gut microbiota stipulated by a handful of studies is mixed. On one

hand, urban-driven environmental changes were associated with reduced gut microbial diversity

in species such as the house sparrow (*Passer domesticus*) (Teyssier et al., 2018b), American

white ibises (*Eudocimus albus*) (Murray et al., 2020) and herring gulls (*Larus argentatus*) (Fuirst

et al., 2018). On the other hand, microbial diversity was found to be higher in city-dwelling

white-crowned sparrows (*Zonotrichia leucophrys*) compared with their rural counterparts

(Phillips et al., 2018; Berlow et al., 2020b).

Another knowledge gap related to our understanding of animal-microbe symbiosis in the urban

space is associated with the fact that all studies published to date were performed on adult birds

and reptiles; thus, there are currently no data exploring the extent to which the gut microbiomes

of birds hatching and growing in urban environments are dissimilar to birds developing in natural

forests. This information gap is particularly striking, as the microbial colonies established in

early life are known to have vital functions in the developmental trajectory of an individual,

affecting survival and fitness (Borre et al., 2014; Cox et al., 2014). Also pertinent to the theme of

juvenile development and life-long physiological homeostasis is the fact that cavity-breeding

songbirds in urban environments are known to occupy both nest boxes and natural cavities.

However, nest boxes, whose supplementation in urban environments is essential to provide

breeding cavities for hole-nesting birds, provide different microclimatic conditions than natural

cavities (Maziarz et al., 2017; Sudyka, in prep). However, the impact of the use of artificial

breeding cavities such as nest boxes on gut microbiota has never been investigated.

Understanding these overlooked aspects of songbird biology can advance our understanding of

urban-driven responses in animal-associated microbial communities and broaden our perspective

on the evolutionary consequences of animal-microbe symbiosis.

To address these knowledge gaps, we investigated changes in the taxonomic diversity and

community composition of gut microbiota in great tit (*Parus major*) nestlings sampled in an

urban mosaic ranging from natural rural landscapes to highly modified urban habitats,

consequently exhibiting profound variations with several environmental parameters, such as the

extent of ISA, noise or light pollution, as well as the rearing environment (natural cavities and

nest boxes). Such a sampling strategy allowed us to test whether (i) community diversity and

composition are influenced by breeding cavity type (artificial nest box or natural cavity), (ii)

microbiota differ in terms of diversity and community structure as described by two different

frameworks defining the urban environment, and, finally, (iii) distinct urban environmental

variables predict changes in microbial alpha and beta diversity.

2. Materials and Methods

2.1. Study Site

Great tits are hole-nesting passerine birds known to occupy a wide range of habitats, from

primary forests (Tomiałojć and Wesołowski, 2004) to urban city centres (Corsini et al., 2020). In

this species, the nesting environment is an important driver that shapes gut microbiota during

early development (Teyssier et al., 2018a), making the great tit an ideal subject to study

microbial changes across an urban mosaic. Here, we monitored the great tit reproductive cycle in

the capital city of Warsaw in Poland, where 565 woodcrete Schwegler nest boxes (type 1b) were

set up from 2016 to 2018 at 9 sites located within and outside the city (Supplementary Material

S1). These sites aimed to cover the urban heterogeneity that is too often simplified to a

dichotomous distinction of urban and rural sampling sites. We included the following sampling

sites: a natural forest (c. 10 km from the city borders and approximately 20 km from the city

centre), a peri-urban village bordering a natural forest, 2 residential areas, 2 urban woodlands, a

large urban park and an office area. These sites are presented in greater detail by Corsini et al.

(2019, 2020). In adsdition to the 8 sites described above, woodcrete nest boxes (Schwegler,

Germany, Type 1B) and natural cavities were also monitored at a 9th site: the Bielany Forest,

which is a unique forest remnant of the Masovian Primeval forest (and retains some of its

features); this forest has an area of ca. 150 ha and is located within the urban administrative

borders, surrounded by a dense urban matrix. The nest box study site was located in the NW part

of the reserve, while the natural cavity study site was located in the SE part of the reserve; both

sites were characterised by similar vegetation structures including hornbeam and oak stands with

100-year succession. A spatial representation of all sites is presented in Figure 1A.

2.2. Capturing Urbanisation

The urban space was defined employing two different frameworks: under the first framework

used to assess the impact of urbanisation on animal-microbe symbiosis, each nest was

categorised depending on whether it was located within the administrative limits of the city

(hereafter - urban) or outside of these limits (hereafter - rural). Based on this dichotomous

categorization, the urban sites encompassed an office area, two residential areas, an urban forest,

an urban park and two urban woodlands. The rural sites comprised a natural forest and a peri-
urban village (Supplementary Material S1). Nonetheless, this often-used dichotomous
partitioning of the urban environment rarely constitutes homogeneous habitats with similar land
use characteristics (Szulkin et al., 2020); for example, urban sites within city limits are, in fact,
mosaic landscapes with varying degrees of actual surface use patterns and accompanying fine-
scale differences in habitat types and environmental characteristics. Consequently, under the
second framework used to assess the impact of urbanisation on animal-microbe symbiosis and to
capture the heterogeneity of both urban and rural sites, we defined urbanisation using the
percentage of impervious surface area (ISA) surrounding the respective location of every nest
analysed in the study. A map of the ISA in Warsaw (in a range from 0 to 100%) with a 20-m
pixel resolution was downloaded from Copernicus Land Monitoring Services
(https://land.copernicus.eu/sitemap). The ISA included all built-up areas and soil-sealing surfaces
substituting original or semi-natural surfaces. Importantly, the ISA index included all artificial
surfaces characterised by long-cover duration (for more details see the official technical
document at https://land.copernicus.eu/sitemap). Averaged ISA values at the nest level were
obtained using a 100-m-radius buffer using geographic information system tools (i.e., QGIS).
The ISA percentage was used as a continuous variable in alpha diversity analysis. For the beta
diversity analysis in which the differences between the groups were investigated, the samples
were categorized depending on whether the ISA percentage of the area with a radius of 100 m
surrounding any specific nest was lower than the median ISA percentage (hereafter, low ISA) or
higher than the median (hereafter, high ISA); the median value was 4.43.

2.3. Environmental and Spatial Variables

Ten environmental and spatial variables were collected i) on the ground and ii) using QGIS and
iii) remote sensing techniques (i.e., digital photography and satellite imagery). While these
methods are described in Szulkin et al. (2020), details on how each environmental variable was
extrapolated in this study are presented in Supplementary Material S2 and are further briefly
specified below.

Regarding the environmental variables collected on the ground, the (1) *human presence* was
derived on the ground by quantifying all humans and dogs within a 15-m radius around each nest
box. A repeatable human presence index value was derived for each nest box. This methodology

is described by Corsini et al. (2017, 2019). (2) ***Sound pollution*** was recorded on the DbC scale using hand-held sound level meters equipped with a microphone (Digital Sound Level Meter SL-200). For each nest box, the sound level was recorded over 4 days throughout the field season, three times per day. (3) The location-specific ***temperature*** was obtained from a Thermocrones ibutton DS1921G set with a 1-hour sampling frequency.

Spatial variables such as the distances from each nest box to (4) the ***closest road*** and to (5) the ***closest path*** were computed and measured in metres using the "Measure line" tool in QGIS, as described by Corsini et al. (2017). The distance from each nest box to (6) the ***city centre*** (i.e., the location of the Palace of Culture and Sciences - 52°13′54″N, 21°00′23″E) was measured in metres using the "Measure line" tool in QGIS. Furthermore, (7) ***a distance matrix of the geographical coordinates*** of the sampling sites was calculated based on the Haversine distances.

Finally, we also extracted environmental variables using digital photography and satellite imagery techniques and computed the values of these variables at the nest level in a 100-m-radius buffer. For (8) ***light pollution***, a map of light pollution in Warsaw with a 10-m pixel resolution was extrapolated from night-time digital photographic images shot on 08/10/2015 by astronauts from the International Space Station (Kyba et al., 2015). For (9) ***tree cover density***, a map of tree cover density in Warsaw (in a range from 0 to 100%) with a 20-m pixel resolution was downloaded from Copernicus Land Monitoring Services (https://land.copernicus.eu/sitemap). For a proxy of live green vegetation, (10) the normalized difference vegetation index (***NDVI***) was estimated using satellite images derived from SENTINEL2 that are available on the Earth Explorer website (https://earthexplorer.usgs.gov).

2.4. Sample Collection

As faeces can be collected using non-lethal methods and can be used to reliably capture gut microbial structure in avian species (Videvall et al., 2018; Berlow et al., 2020a), the microbial communities present in faecal samples were used in this study as a proxy for the gut microbial communities. In 2018, faecal samples were collected from 15-day-old great tit chicks. The faeces were collected from one chick per nest (from the largest chick in the brood), except for four nests in which more than one chick was sampled per nest (all originating from woodcrete nest boxes). Altogether, one to three chicks were collected per nest, with a total of 89 nests sampled from a mosaic of diverse urban environments. Of these nests, 8 nests were located in natural cavities,

238 while 81 nests were located in woodcrete nest boxes. The faeces were collected by manually

239 applying mild pressure on the cloaca during the time of ringing. The faeces were directly

240 deposited in 5-mL sterile Eppendorf tubes, each filled with 3 mL of RNAlater and further stored

241 at -20 °C.

2.5. DNA Extraction and Library Preparation

The details of the microbial analysis are described in Maraci et al. (2021). In short, microbial

DNA was extracted using the QIAamp PowerFecal DNA Kit (Qiagen), as described in the

manufacturer's protocol. The hypervariable V3–V4 region of the 16S ribosomal RNA (rRNA)

gene was targeted following the Illumina 16S Metagenomic Library Preparation Guide,

15044223-B. The final amplicon pool contained a pool of blank controls for DNA extraction and

PCR amplification and one replicate of a single sample, alongside 94 biological samples

sequenced on the Illumina MiSeq system (Illumina, Inc., San Diego, CA, USA).

2.6. Bioinformatic Processing

The bioinformatic processing was carried out as described in detail (Maraci et al., 2021). In

short, MiSeq PE reads were assembled in an iterative manner using Flash v1.2.11 (Magoč and

Salzberg, 2011). All other bioinformatic steps consisted of i) adapter clipping with cutadapt

v1.18 (Martin, 2011); ii) de-replication, alignment, filtering and de-noising with mothur v1.41.3

(Schloss et al., 2009); iii) chimaera checking and operational taxonomic unit (OTU) clustering

with USEARCH v8.0.1477 (Edgar, 2010); and iv) taxonomic classification based on the full

SILVA database v132 (Quast et al., 2013).

2.7. Statistical Analyses

All consecutive statistical analyses were conducted in R version 4.0.0 (R Core Team, 2020). As

an initial quality filtering step, all OTUs that could not be classified at the phylum level (N=41)

or that were classified as mitochondria or chloroplasts (N=121) were discarded. Next, all OTUs

that were not represented by at least 1 read in 2% of the samples (N=1035) or with fewer read

counts than 0.001% of the total number of reads (N=561) were excluded. Subsequently, samples

with poor qualities (N=12) were removed from the dataset. After these quality filtering steps

were completed, our dataset contained 82 biological samples and 1090 OTUs (Supplementary

Material S1).

To test whether the microbial communities were affected by the cavity type, we created a subset

of data (hereafter, the cavity type dataset) including the samples collected from nest boxes (N=7)

and those collected from natural cavities (N=6) located in the same urban forest. As there were

significant differences in alpha diversity between the samples collected from these different

cavity types, the six samples obtained from natural cavities were excluded from the remaining

analyses, and the remaining dataset ultimately consisted of 76 samples collected from nest boxes

to be used in the further analyses (Supplementary Material S1). Using a two-step approach, we

analysed the interactions among microbial community differences, urbanisation and

environmental change. First, the urban space was defined using two different frameworks based

on (i) the often-used administrative border delineation (urban/rural site) and (ii) the percentage of

impervious surface area (ISA) surrounding each nest (see 2.2.), irrespective of the location of the

nest within or outside of the city borders. Thus, microbial alpha diversity was compared (i)

between urban and rural sites but was also (ii) examined in relation to ISA percentage, where

ISA was fitted as a continuous variable. The differences in microbial beta diversity that existed

between urban and rural sites as well as between low- and high-ISA categories were also

examined. Next, the environmental and spatial variables that covaried most strongly with alpha

and beta diversity differences were further identified.

As a measure of alpha diversity (the species diversity within each site), Shannon's diversity

index, which accounts for both the abundance and evenness of the taxa present (Shannon, 1997),

was calculated. To analyse the impact of the cavity type on alpha diversity, a linear model was

fitted using the square-root-transformed Shannon's diversity indices of the samples in the cavity

type datasets as the response variable and the cavity type as a fixed effect, implemented using

lme4 version 1.1-25 (Bates et al., 2015). To analyse the impact of urbanisation on alpha

diversity, we employed two separate linear mixed models (LMMs) using square-root-

transformed Shannon's diversity indices of the samples collected from nest boxes as the response

variable. The fixed effects differed between the two models depending on the framework used to

define urbanisation: in the first model, the urban/rural categorisation was used as a fixed effect,

while the second LMM was constructed using ISA (as a continuous variable) as the fixed effect.

The brood ID and sampling site were included in these two models as random effects to account

for the nonindependence of the samples coming from the same brood or the same sampling site.

The taxonomic and compositional structure of the microbial communities collected from

different localities were visualized with stacked bar plots based on family-level taxonomy using

ggplot2 version 3.3.2 (Wickham, 2009). To identify differentially abundant OTUs in samples

collected from urban and rural localities as well as from low- and high-ISA territories, the

logarithmic fold changes between groups were estimated using a negative binomial Wald test

implemented in *Deseq2* version 1.12.4 extension (Love et al., 2014) of the *Phyloseq* package

version 1.32.0 (McMurdie and Holmes, 2013). The significance threshold of the p values was set

as 0.05 after a Benjamini and Hochberg false⬜discovery rate correction (Benjamini and

Hochberg, 1995).

To analyse beta diversity, the degree of community differentiation among sites (Whittaker,

1960), first, the filtered absolute abundance data were normalized by applying cumulative sum

scaling (CSS) normalization in the R package *Metagenomeseq* version 1.30.0 (Paulson, 2014) to

account for unequal sequence coverage. To address compositional variations, the data were log

$(x + 0.0001)$⬜transformed and, later, the transformed values were corrected by subtracting the

log of the pseudocount. The beta dissimilarities between *a priori* defined groups, i.e., urban and

rural sites and low- and high-ISA territories, were visualized using non-metric multidimensional

scaling (nMDS) based on Bray–Curtis dissimilarity (Bray and Curtis, 1957), implemented using

the *Vegan* package version 2.5-6 (Oksanen et al., 2019). The differences in the gut microbial

communities between the *a priori* defined groups were also statistically tested using a

nonparametric analysis of similarities (ANOSIM) with 9999 permutations in the *Vegan* package

based on Bray–Curtis and the weighted UniFrac (Lozupone et al., 2007) dissimilarities.

Employing a similar approach to that used in the alpha diversity analyses, first, the differences

between the samples collected from nest boxes and natural cavities in the cavity type dataset

were tested. Then, the dissimilarities between samples originating from urban and rural

territories, among different sampling sites and between low- and high-ISA categories were

analysed using ANOSIM.

Finally, the associations between distinct variables characterizing the urban environment and

microbial alpha and beta diversity were also inferred. As an initial step, the extent to which the

environmental parameters varied between urban and rural sites and between the low- and high-

ISA categories was tested by Welch's two-sample t-test. To analyse the interactions between

alpha diversity and environmental variables, the correlations among the environmental variables

were first analysed by using Pearson's correlation. Second, a linear mixed model was fitted using

the square-root-transformed Shannon's diversity index as the response variable, all

environmental variables fixed effects and the brood ID and sampling site as random effects. To

examine multicollinearity between the variables, the variance inflation factor (VIF) was

calculated with the *Car* package (Fox et al., 2012), and the predictors with the largest VIF values

were sequentially removed from the model and VIF values were recalculated (Zuur et al., 2010).

This procedure was repeated until all the VIF values were smaller than two. The final model

fitted using the remaining environmental variables, distance to the city centre, human presence,

temperature, tree cover density and distance to the closest path, as fixed effects. The interactions

between spatial and environmental variables and beta diversity were also evaluated by

employing different methods. First, to identify the variables that were strongly related to the first

two ordination axes, the multiple regression between each environmental variable and the

ordination axes was evaluated using the *Envfit* function, and the significance of each correlation

was tested based on 9999 permutations. Consequently, the subset of environmental variables

whose Euclidean distance matrices correlated maximally with the microbial distance matrix

based on Bray-Curtis was identified employing the *BIOENV* procedure (Clarke and Ainsworth,

1993). Finally, the correlations between the microbial dissimilarity matrices obtained based on

weighted Bray-Curtis distances and the distance matrices of the environmental variables

obtained based on Euclidean distances were then evaluated using the Mantel test. Geographic

separation, an additional non-linear parameter that could not be included in BIOENV or Envfit,

was included in the Mantel test to compare the distance matrix of the geographical coordinates of

the sampling sites with the microbial distance matrix.

3. Results

After the quality-filtering steps were completed, the resulting dataset consisted of 82 biological

samples and 1090 OTUs with a total read count of 1277151 (mean: 15575.01). These OTUs

were represented by 19 different microbial phyla, three of which constituted 96.9% of the OTUs

based on their total relative abundances: Firmicutes (69.5%), Proteobacteria (18.6%) and

Actinobacteria (8.8%). The identified taxa corresponded to 213 microbial families, among which

only eight constituted 76.7% of all obtained microbial taxa: *Catellicoccaceae* (31.1%),

Clostridiaceae (18.3%), *Mycoplasmataceae* (8.4%), *Enterobacteriaceae* (6.7), *Enterococcaceae*

(4.2%), *Yersiniaceae* (3.1%), *Lactobacillaceae* (2.7%) and *Moraxellaceae* (2.2%) (Figure 1B).

3.1. Microbial Community Diversity and Structure in Relation to Cavity Type

The microbial alpha diversity was higher in nest box-reared nestlings than in nestlings reared in natural cavities (linear model; β=-1.62 ±0.61, 95% CI [-2.96;-0.29], p=0.021). In contrast, beta diversity did not differ among nestlings originating from these two cavity types (ANOSIM Bray-Curtis, global R=-0.04, p=0.61; ANOSIM Weighted UniFrac, global R =-0.18, p = 0.97).

3.2. The Impact of Urbanization on Microbial Community Diversity and Structure

To understand the interaction between urbanisation and gut microbial communities, we analysed different gut microbial measures of nestlings growing in an urban mosaic defined by two different urbanisation frameworks: (i) first framework based on administrative boundaries (contrasting alpha and beta diversity in urban vs. rural areas); and (ii) second framework based on ISA (by examining the interaction between ISA as a continuous variable and alpha diversity and by contrasting beta diversity between areas of low- and high-level ISA).

First, we analysed the interaction between Shannon's diversity index as a measure of alpha diversity and urbanisation. When analysed based on administrative boundaries, the alpha diversity did not differ between hosts originating from urban and rural sites (LMM; β=-0.23 ±0.18, 95% CI [-0.59-0.13], p=0.215). However, when we analysed the covariation of ISA (as a continuous variable) and alpha diversity, the LMM explained considerable variation in alpha diversity (LMM; R^2-marginal=0.069, R^2-conditional=0.528) and revealed a significant negative association between the two (β=-0.14 ±0.06, 95% CI [-0.23--0.01], p=0.029).

Beta diversity also differed across the urban mosaic. In contrast to alpha diversity, we found significant differences in community structure between urban and rural sites when analysed based on administrative boundaries (ANOSIM Bray-Curtis, global R=0.180, p=0.001; ANOSIM Weighted UniFrac, global R=0.160, p=0.002) but also among sampling sites (ANOSIM Bray-Curtis, global R=0.174 p<0.001; ANOSIM Weighted UniFrac, global R=0.150, p=0.004; Figure 1C). Importantly, the microbial community structure also differed between low-ISA and high-ISA territories (ANOSIM Bray-Curtis, global R=0.090, p=0.003; ANOSIM Weighted UniFrac, global R=0.080, p =0.003); however, this change was less prominent (Figure 1D).

Importantly, the taxonomic compositions of microbial communities varied considerably in the urban mosaic (Figure 1B): when the urban and rural microbial samples were compared, the

dominating microbial phyla were the same; however, the relative abundances of these phyla differed markedly. The relative abundance of Firmicutes was higher in the urban samples than in the rural samples (72.8% vs. 59%), while the relative abundances of Proteobacteria and Actinobacteriota were higher in rural samples than in the urban samples (20.9% vs 17.9% and 12.9% vs 7.6%, respectively). The distinction between urban and rural samples was even more evident at a finer taxonomic scale: approximately 80% of the microbial taxa were represented by six and 22 families in the urban and rural samples, respectively (Figure 1B). When comparing the low- and high-ISA territories, the relative abundance of Firmicutes was higher in high-ISA territories than in low-ISA territories (70.5 vs 65.4), while Proteobacteria and Actinobacteriota were more abundant in low-ISA territories than in high-ISA territories (19.9 vs 18.3 and 11.4 vs. 8, respectively). Similarly, as observed in the urban-rural dichotomy, 80% of the microbial taxa were constituted by nine and 15 families in the high-ISA and low-ISA samples, respectively.

Differentially abundant taxa were determined with the DESeq2 method. Between the urban and rural territories, overall, we found 20 differentially abundant OTUs (Figure 2A). Of these, 10 were significantly more abundant in the rural territories, and 10 were significantly more abundant in urban territories. Urban hosts exhibited higher abundances of a potentially pathogenic microbial family, *Enterobacteriaceae* (Sackey et al., 2001; Benskin et al., 2009). Between the low- and high-ISA territories, only nine OTUs were differentially abundant. Among these, four OTUs were more abundant in hosts from high-ISA territories, and five OTUs were more abundant in hosts from low-ISA territories (Figure 2B). Overall, seven of these OTUs were detected as differentially abundant in the urban space under both urbanisation frameworks (urban/rural and ISA-based). All five of the OTUs that were detected as being more abundant in birds from low-ISA territories were also found to be more abundant in rural sites. Only one out of four OTUs that were found to be more abundant in hosts originating from high-ISA territories were also more abundant in urban sites. One OTU belonging to the family *Enterococcaceae* was found to be more abundant in high-ISA territories than in low-ISA territories and, contrastingly, more abundant in rural sites than in urban sites.

3.3. Interaction between Spatial and Environmental Parameters and Microbial Change

While nesting sites are here referred to as being located in urban or rural spaces or areas with high or low ISA, each nest box is also embedded in a complex web of specific environmental

and spatial parameters characterising the urban mosaic; these parameters are defined herein as

the distance to the city centre, light pollution, sound pollution, human presence, distance to the

closest road, temperature, tree cover density, NDVI, and distance to the closest path. All these

environmental parameters differed between the urban and rural sites (except for the distance to

the closest path, NDVI and tree cover density; Supplementary Material S3) and between the low-

and high-ISA sites (except for the distance to the closest path; Supplementary Material S3).

As multiple environmental and spatial variables were found to be correlated with each other

(Supplementary Material S4), we examined multicollinearity among all variables and

sequentially excluded predictors with the largest VIF values when analysing the predictors of

alpha diversity. Our final model included five predictors: the distance to the city centre, human

presence, temperature, tree cover density and distance to the closest path; among these

predictors, only the tree cover density was significantly associated with alpha diversity (LMM,

$\beta=0.18 \pm 0.09$, 95% CI [0.01-0.35], p=0.043).

To identify the strongest predictors of community structure, we tested the associations between

all environmental variables and beta diversity by employing different analytical methods to

obtain a coherent vision of the urban-driven variation in gut microbiota community structure.

Based on *BIOENV*, the best model with the strongest relationship with microbial dissimilarities

(R=0.209) contained sound pollution, the distance to the city centre, the distance to the closest

road and light pollution. Based on *Envfit*, the patterns observed in the ordination plot were

significantly associated with sound pollution ($R^2=0.332$; p<0.001), the distance to the city centre

($R^2=0.318$; p<0.001), the distance to the closest road ($R^2=0.216$; p<0.001), light pollution

($R^2=0.196$; p<0.001), tree cover ($R^2=0.117$; p=0.01) and the distance to the closest path

($R^2=0.089$; p=0.03) but not with temperature, NDVI or human presence. Mantel tests further

confirmed the correlation of the microbial dissimilarity matrix calculated based on Bray-Curtis

distances for three of these variables: sound pollution (R=0.188; p<0.001), the distance to the

city centre (R=0.139; p<0.001), and the distance to the closest road (R=0.110; p=0.034).

Additionally, the Mantel tests showed an association between tree cover (R=0.084; p=0.008) and

geographic separation (R=0.130; p=0.001), a non-linear parameter that could not be included in

the *BIOENV* procedure or *Envfit*. All the other parameters were found to be nonsignificant in the

Mantel tests (Supplementary Material S5). Overall, there was coherent and conclusive support

across all models for sound pollution, the distance to the city centre and the distance to the

closest road acting as possible drivers of juvenile gut microbiota variation (beta diversity) across

the urban mosaic.

4. Discussion

To understand how ecological and evolutionary drivers impact urban biodiversity, one must

understand both how biodiversity scales with city size, as well as with the various environmental

dimensions of the urban landscape (Uchida et al., 2021). As demonstrated in this study, this

statement fully pertains to biodiversity even at its finest scale, such as the gut microbiota of wild

passerines. We investigated the impacts of urbanisation inferred at different spatial scales and

environmental dimensions by comparing gut microbiota from juvenile birds reared (i) in natural

cavities and nest boxes, (ii) within the city boundaries and from rural areas, and (iii) in the

context of varying levels of sealable surfaces (ISA) in the immediate surroundings of each nest.

Furthermore, independent urban environmental variation axes were further tested against gut

microbiota variations. Based on our findings, alpha diversity was reduced as land-use intensity

increased, as measured by ISA, but this change was not evident when the data were analysed

based on a simple urban-rural dichotomy (Figure 3). This indicates that the frameworks in which

actual land-use patterns are quantified across an urban mosaic might be more efficient in

capturing the alterations in taxonomic diversity and the evenness of microbial communities than

a simple dichotomous approach (Figure 3). By contrast, the differences in community structure

(beta diversity) were more pronounced between urban and rural sites as defined by

administrative borders than when defined using ISA. We further aimed to assess the covariation

between the observed differences in alpha and beta diversity with distinct environmental and

spatial variables characterizing the urban mosaic. Alpha diversity was associated with tree cover

density, while the strongest predictors of beta diversity were sound pollution, the distance to the

city centre and the distance to the closest road (Figure 3). We also showed that the gut

microbiota of birds reared in nest boxes was more diverse than those of birds reared in natural

cavities. Overall, the present study provides new perspectives on how the gut microbiota, an

essential component of host fitness (Gould et al., 2018), differs in an urban mosaic in the early

developmental stage of a wild passerine species and how the choice of framework used to

describe the urban space can influence the outcomes of ecological studies.

4.1. Cavity Type Influences Microbial Diversity but not Community Structure

Our dataset, albeit small, clearly revealed that alpha diversity is higher in the nestlings reared in artificial cavities, nest boxes, relative to those reared in natural cavities. By contrast, community composition (beta diversity) did not differ between these two cavity types. Nest boxes are frequently used in ecological studies of whole-nesting breeding birds, as they are convenient substitutes for natural cavities. However, human-made nest boxes differ from natural cavities in various aspects by having different humidity values and providing poorer insulation, leading to prominent fluctuations in daily ambient temperatures across 24 hours (Maziarz et al., 2017). For example, at the same ambient temperature, the minimum and maximum temperature values in a nest box can be, on average, 2.8 °C and 3.6 °C higher, respectively, than those in a natural cavity (Maziarz et al., 2017). Similarly, based on microclimate data collected in 2018 and 2019 in great tit nests located in natural cavities and nest boxes in the urban forest location, woodcrete nest boxes were found to have higher temperatures relative to natural cavities (mean ± SD: 20.5±5.09 °C vs 16.2±3.08 °C) and higher absolute humidity values (mean ± SD: 25.4±4.55 hPa vs 20.1± 2.08 hPa; Sudyka et al. in prep). The gastrointestinal ecosystem is relatively well-regulated and is not expected to exhibit prominent alterations in response to moderate fluctuations in ambient conditions in endothermic animals. At the same time, it should be considered that the nestlings of altricial passerines cannot regulate their body temperatures (Hansell and Overhill, 2000). Therefore, even moderate fluctuations can alter nestling physiology and, consequently, nestling gut microbiota (Sepulveda and Moeller, 2020). Another explanation is that such fluctuations can influence the growth of microbial communities on nesting materials, which is an important reservoir for the colonisation of the gut by microbiota in early life (Chen et al., 2020; van Veelen et al., 2020). Considering that even small temperature changes can influence the microbial growth rate (Nottingham et al., 2019), most mesophilic bacteria need temperatures of at least 20 °C for optimal growth (Schiraldi and De Rosa, 2014); the moderately higher temperatures provided by the nest boxes than by natural cavities might thus be deterministic in terms of which species will grow on the nesting material and consequently colonize the guts of the developing nestlings. It is important to note that this novel finding should be confirmed with larger datasets and considered while planning and interpreting future studies.

4.2. Alterations in the Taxonomic Composition, Diversity and Community Structure of Gut Microbiota in the Urban Space

To understand the extent to which gut microbial communities differ between urban and rural spaces and whether the criteria used to define the urban space have any impact on these observations, we compared the taxonomic composition, diversity and community structure between urban and rural sites as defined by administrative borders (urban/rural) as well as by high- and low-ISA territories (Figure 3).

Microbial alpha diversity was found to be lower in highly urbanised spaces than in rural spaces. Remarkably, this decrease in alpha diversity did not follow a simple urban-rural dichotomy pattern; instead, the change was correlated with actual land-use intensity, as measured by ISA. The observed alterations in alpha diversity are probably more strongly associated with fine-scale variations in the local habitat, which shows remarkable heterogeneity within the urban mosaic, rather than with factors related to human activities or the geographical locations of the nests. These variations, in turn, can influence the diversity of gut microbiota via different mechanisms, which are discussed in the next section. Earlier studies investigating how microbial alpha diversity varies in urban spaces have revealed contradictory results: urban populations of adult house sparrows (Teyssier et al., 2018b, 2020), American white ibises (Murray et al., 2020), Darwin's finches (Knutie et al., 2019) and herring gulls (Fuirst et al., 2018) exhibited reduced diversity in urban spaces. In contrast, the gut microbial diversity was found to be higher in urban populations of white-crowned sparrows than in their rural counterparts (Phillips et al., 2018; Berlow et al., 2020b). Notably, in the study by Phillips et al. (2018), this increase was not evident when analysed based on ISA. Furthermore, alpha diversity in juvenile house sparrows did not differ between urban and rural populations (Teyssier et al., 2020). As indicated by our results, these discrepancies may be driven by differences in the criteria used to define the urban space. Alternatively, but not exclusively, biological differences originating from the taxonomy and life-history stage of the host may also play a role in these discrepancies. For example, avian species with higher tolerance to anthropogenic food might exhibit higher alpha diversity in urban areas. Another explanation is that urban growth patterns are not uniform across the world, as cities exhibit remarkable variations in their ecosystem characteristics and present distinct selection pressures that, in turn, influence the evolution of urban organisms and their microbial symbionts differently.

The taxonomic compositions of microbial communities differed between urban and rural

territories as well as between high- and low-ISA territories: the microbial communities collected

from hosts occupying highly urbanised areas were less diverse than those collected from rural

areas. Although this result was true under both frameworks (Figure 3) and some of the

differentially abundant taxa observed in the urban spaces were the same, we also observed some

differences when applying different urbanisation frameworks. Among these, the most striking

difference was noted when the urban environment was defined based on administrative borders,

as the urban hosts were enriched by a potentially pathogenic microbial family,

Enterobacteriaceae (Sackey et al., 2001; Benskin et al., 2010). However, when defined based on

ISA categories, this pattern was not evident. A higher prevalence of *Enterobacteriaceae* has been

associated with dysbiosis in mice (Garrett et al., 2010) and higher mortality rates in ostriches

(Videvall et al., 2020). Similarly, urban populations of American white ibises were found to have

lower microbial diversity and were more susceptible to *Salmonella* infections than their rural

counterparts (Murray et al., 2020). Collectively, these findings indicate that pathogen

susceptibility increases in densely populated urban sites, potentially adversely influencing the

overall health and fitness of animal hosts.

The impacts of urbanisation on microbial community structures have been previously studied in

house sparrows (Teyssier et al., 2018b), American white ibises (Murray et al., 2020) and white-

crowned sparrows (Phillips et al., 2018; Berlow et al., 2020b). We also demonstrated shifts in

beta diversity between habitats with contrasting land-use patterns. The community membership,

relative abundance and phylogenetic distance of the microbial species differed between

territories with low and high ISA percentages. Importantly, and in contrast to the results obtained

for alpha diversity, these community structure alterations were even more prominent under the

urban/rural distinction reflected by administrative borders than that obtained using ISA,

indicating that beta diversity was largely determined by the geographical location of the nest or

by whether the nestling lived in a densely populated urban site (Figure 3).

Taken together, our findings revealed that the taxonomic composition, diversity and community

structure of the gut microbial communities of juvenile great tits exhibit variations in the urban

space; using different frameworks to describe the urban space leads to the capture of different

aspects of these variations. The initial microbial colonies inhabited by animal hosts in early life

are not only critical for the establishment of healthy gut microbiota but also have crucial

functions, such as the programming of the immune system (Hooper and MacPherson, 2010), and

are involved in the maturation of the nervous system (Borre et al., 2014). Therefore, the observed

alterations might have long-term consequences on the survival and fitness of hosts (Cox et al.,

2014).

4.3. Potential Environmental Factors Underlying Gut Microbial Alterations in the Urban

Space

Urbanization is one of the strongest anthropogenic forces that modify multiple environmental

variables (McKinney, 2006; Grimm et al., 2008). In line with this, most of the environmental

variables investigated in the present study varied in the analysed urban space regardless of the

framework used to define that space. One of our primary goals was to determine whether some

of these environmental variables can predict changes in the gut microbiota of great tit nestlings.

Based on our findings, alpha diversity was positively associated with tree cover density (Figure

3). In line with our findings, white-crowned sparrows occupying territories with greater tree

cover also exhibited more diverse gut microbial communities than those occupying regions with

less tree cover (Phillips et al., 2018; Berlow et al., 2020b). This is likely to be explained by the

tree cover density influencing the diversity and abundance of invertebrates (Jones and Leather,

2012), which are the primary food source for great tits during the chick-rearing period (Wilkin et

al., 2009). Reductions in tree cover density, often observed in urban spaces, can also direct

animals to search for anthropogenic food (Pollock et al., 2017). Consistently, the camera

recordings of the nests sampled in this study revealed that the amount of anthropogenic food

brought into the nests was higher in territories characterized by high ISA percentages than in

low-ISA territories (Corsini et al., in prep.). Dietary alterations are known to be associated with

changes in alpha diversity (Davidson et al., 2020; Teyssier et al., 2020). However, it is important

to note that these assumptions should be interpreted with caution due to some shortcomings of

our study. First, as we did not analyse the diets of the studied nestlings, our predictions regarding

their diets remain speculative. Furthermore, due to multicollinearity issues, the majority of the

environmental variables could not be included in the model used to investigate alpha diversity.

Therefore, the observed patterns in alpha diversity might stem from another predictor that is

correlated with tree cover density.

In contrast to alpha diversity, beta diversity is largely influenced by whether the nestlings live in
an urban or rural habitat. The variables that showed the strongest associations with the microbial
community structure in all models—BIOENV, Envfit and Mantel tests—were the distance to the
city centre, sound pollution, and the distance to the closest road. Consequently, this result can be
interpreted by different but non-exclusive mechanisms. As the distance from the city centre was
among the strongest predictors of beta diversity and geographic separation was among the
significant explanatory variables, microbial communities might be geographically structured.
One potential explanation is that the microbial communities residing in the guts of studied
nestlings, at least to some extent, represents the environmental pool of microorganisms (Sullam
et al., 2012), which exhibits microgeographic variations (Martiny et al., 2006). In line with this,
the gut microbiota is spatially structured, even when social interactions are minimal (Antwis et
al., 2018). Alternatively, observed community changes might be driven by environmental
variables that are distributed linearly from the city centre such as sound, air or light pollution.
Consistently, another potential driver of the microbial community structure determined in this
study was sound pollution. Noise might impair parent-offspring communication by masking the
begging calls of the young, and parents might adjust their food provisioning accordingly
(Schroeder et al., 2012; Lucass et al., 2016). Therefore, the amount of food received by young
might vary depending on the background noise levels, which in turn could influence gut
microbiota. Alternatively, noise might have an impact on the host physiology: noise can activate
the hypothalamic-pituitary-adrenal axis and alter the endocrine profiles (Kight and Swaddle,
2011; Zollinger et al., 2019), causing cascading impacts on glucose metabolism (Cui et al.,
2016), immune function (Campo et al., 2005) oxidative stress (Injaian et al., 2018) and gut
physiology (Cui et al., 2018) and consequently influencing gut microbiota (Cui et al., 2016,
2018). The third predictive variable supported by all three tests was the distance to the closest
road. The observed correlations between the distance to the closest road and beta diversity might
be associated with other factors showing collinearity with the distance from the closest road as
well as the distance from the city centre, such as air pollution, a variable that was not tested in
our study, or light pollution, which was found to be a significant predictor based on the BIOENV
procedure. Indeed, both particulate matter in ambient air and artificial light at night can indirectly
affect gut microbiota by altering the host physiology (Wei et al., 2020). It is important to note
that due to the multicollinearity among these variables, it is not possible to infer the exact

correlative factors explaining changes in the microbial community structure, and each of these

assumptions requires further experimental testing to infer the exact causative factors driving the

observed differences in beta diversity.

4.4. Conclusions and Outlook

We analysed how anthropogenic modifications present in the rearing environment due to

urbanisation and the use of artificial nest boxes can reshape animal-microbe interactions in great

tit nestlings. We presented clear trends regarding urban-related shifts in taxonomic composition

and community structure, a reduction in alpha diversity, and an increase in pathogen

susceptibility. While the exact causal interactions between the environmental parameters and

observed microbial changes remain to be tested experimentally, we outlined different

mechanisms that are likely to be the driving forces of the measured microbiota variation in the

studied urban space. This study fills an important gap by providing pioneering evidence on how

anthropogenic changes may relate to gut microbiota during early nestling development, the most

critical life-history stage during which gut microbial communities are established and

programme several developmental processes of their hosts (Sommer and Bäckhed, 2013; Borre

et al., 2014; Cox et al., 2014). Another novel piece of information provided by our study is how

the use of nest boxes, the inevitable instruments of ecological studies, can affect the outcomes of

studies investigating gut microbial communities, an important point that should be considered in

future investigations. Our study further adds to the existing evidence that the frameworks used to

define urban spaces might have a prominent impact on the outcomes of ecological studies.

Our study also underscores some further questions and opportunities. First, it is pivotal to

conduct experimental testing in which the confounding variables are strictly controlled to infer

the exact causal effects of the identified environmental factors that are strongly correlated with

microbial changes. Second, whether and how gut microbes can facilitate the adaptations of their

hosts to changing environments is an important point that remains to be investigated to achieve

full comprehension of the long-term fitness consequences of host-microbe interactions. Third,

our study raised the possibility of answering a central question in microbial ecology by

leveraging the urban-rural gradient; cross-fostering experiments among contrasting habitats

would allow us to disentangle the relative importance of host-specific and environmental factors

in shaping microbial communities. Answering these questions would allow us to fully

comprehend the ecological and evolutionary dynamics and consequences of animal-microbe

symbiosis and make more informed decisions regarding urban planning, wildlife and public

health management in the face of globally increasing urbanisation.

5. Ethics Statement

The research was carried out with a permit from the Regional Directorate for Environmental

Protection (RDOŚ) in Warsaw, Poland.

6. Data Accessibility

Sequences generated for this study have been uploaded to the European Nucleotide Archive

(ENA) repository under the accession number: PRJEB44290. The scripts used for processing the

data are provided in the GitHub repository at:

https://github.com/AnnaAntonatouPap/Urban-related-changes-in-tit-microbiota

7. Authors' Contributions

Ö. M., B. A. C. and M. S. conceptualized the research idea and planned the experiments. M. C.,

M. S., I. dL. and J. S. collected the biological samples and environmental data. Ö. M. carried out

the laboratory experiments. S. J. performed the bioinformatic analyses. A. A. P. carried out the

statistical analyses with the supervision of B. A. C. and Ö. M., with input from M. C. and M. S.

Ö. M. wrote the manuscript in consultation with M. S., and all authors approved the final version

of the manuscript.

8. Competing Interests

We have no competing interests.

9. Funding Information

This study was financially supported by a small-scale research grant from the Research and
Innovation Fund of the Faculty of Biology, Bielefeld University to Ö. M., a Freigeist Fellowship
from the Volkswagen Foundation to BAC, research grants from the Polish National Science
Foundation (NCN) Sonata Bis 2014/14/E/NZ8/00386 to M. S. and M. C., and OPUS
2016/21/B/NZ8/03082 to M. S., I. D. L. and J. S. Additionally, S. J. received financial support
from the German Federal Ministry of Education and Research (BMBF) (Grant number
031A533). This research was inspired by the SFB-TRR 212.

10. Acknowledgements

We are grateful to Elke Hippauf, Barbara Fuchs, Tobias Busche, Yvonne Kutter and Katharina
Hanuschka for their technical support during the laboratory procedures and to all members of the
Wild Urban Evolution and Ecology Lab for collecting data in Warsaw and its surrounding
regions.

11. References

Afshinnekoo, E., Meydan, C., Chowdhury, S., Jaroudi, D., Boyer, C., Bernstein, N., et al. (2015).
    Geospatial Resolution of Human and Bacterial Diversity with City-Scale Metagenomics.
    Cell Syst. doi:10.1016/j.cels.2015.01.001.

Alberti, M., Correa, C., Marzluff, J. M., Hendry, A. P., Palkovacs, E. P., Gotanda, K. M., et al.
    (2017). Global urban signatures of phenotypic change in animal and plant populations.
    Proc. Natl. Acad. Sci. U. S. A. doi:10.1073/pnas.1606034114.

Antwis, R. E., Lea, J. M. D., Unwin, B., and Shultz, S. (2018). Gut microbiome composition is
    associated with spatial structuring and social interactions in semi-feral Welsh Mountain
    ponies. *Microbiome.* doi:10.1186/s40168-018-0593-2.

Auman, H. J., Meathrel, C. E., and Richardson, A. (2008). Supersize me: Does anthropogenic
    food change the body condition of silver gulls? A comparison between urbanized and

remote, non-urbanized areas. *Waterbirds*. doi:10.1675/1524-
4695(2008)31[122:SMDAFC]2.0.CO;2.

Bates, D., Mächler, M., Bolker, B. M., and Walker, S. C. (2015). Fitting linear mixed-effects
models using lme4. *J. Stat. Softw.* doi:10.18637/jss.v067.i01.

Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B*. doi:10.1111/j.2517-
6161.1995.tb02031.x.

Benskin, C. M. W. H., Rhodes, G., Pickup, R. W., Wilson, K., and Hartley, I. R. (2010).
Diversity and temporal stability of bacterial communities in a model passerine bird, the
zebra finch. *Mol. Ecol.* doi:10.1111/j.1365-294X.2010.04892.x.

Benskin, C. M. W. H., Wilson, K., Jones, K., and Hartley, I. R. (2009). Bacterial pathogens in
wild birds: A review of the frequency and effects of infection. *Biol. Rev.*
doi:10.1111/j.1469-185X.2008.00076.x.

Berlow, M., Kohl, K. D., and Derryberry, E. P. (2020a). Evaluation of non-lethal gut
microbiome sampling methods in a passerine bird. *Ibis (Lond. 1859).*
doi:10.1111/ibi.12807.

Berlow, M., Phillips, J. N., and Derryberry, E. P. (2020b). Effects of Urbanization and
Landscape on Gut Microbiomes in White-Crowned Sparrows. *Microb. Ecol.*
doi:10.1007/s00248-020-01569-8.

Bestion, E., Jacob, S., Zinger, L., Di Gesu, L., Richard, M., White, J., et al. (2017). Climate
warming reduces gut microbiota diversity in a vertebrate ectotherm. *Nat. Ecol. Evol.*
doi:10.1038/s41559-017-0161.

Borre, Y. E., O'Keeffe, G. W., Clarke, G., Stanton, C., Dinan, T. G., and Cryan, J. F. (2014).
Microbiota and neurodevelopmental windows: implications for brain disorders. *Trends Mol.*
Med. doi:10.1016/j.molmed.2014.05.002.

Bray, J. R., and Curtis, J. T. (1957). An Ordination of the Upland Forest Communities of
Southern Wisconsin. *Ecol. Monogr.* doi:10.2307/1942268.

Campo, J. L., Gil, M. G., and Dávila, S. G. (2005). Effects of specific noise and music stimuli on

stress and fear levels of laying hens of several breeds. *Appl. Anim. Behav. Sci.*

doi:10.1016/j.applanim.2004.08.028.

Chen, C. Y., Chen, C. K., Chen, Y. Y., Fang, A., Shaw, G. T. W., Hung, C. M., et al. (2020).

Maternal gut microbes shape the early-life assembly of gut microbiota in passerine chicks

via nests. *Microbiome*. doi:10.1186/s40168-020-00896-9.

Christiansen, N. A., Fryirs, K. A., Green, T. J., and Hose, G. C. (2019). The impact of

urbanisation on community structure, gene abundance and transcription rates of microbes in

upland swamps of Eastern Australia. *PLoS One*. doi:10.1371/journal.pone.0213275.

Clarke, K. R., and Ainsworth, M. (1993). A method of linking multivariate community structure

to environmental variables. *Mar. Ecol. Prog. Ser.* doi:10.3354/meps092205.

Corsini, M., Dubiec, A., Marrot, P., and Szulkin, M. (2017). Humans and tits in the city:

Quantifying the effects of human presence on great tit and blue tit reproductive trait

variation. *Front. Ecol. Evol.* doi:10.3389/fevo.2017.00082.

Corsini, M., Marrot, P., and Szulkin, M. (2019). Quantifying human presence in a heterogeneous

urban landscape. *Behav. Ecol.* doi:10.1093/beheco/arz128.

Corsini, M., Schöll, E. M., Di Lecce, I., Chatelain, M., Dubiec, A., and Szulkin, M. (2020).

Growing in the city: Urban evolutionary ecology of avian growth rates. *Evol. Appl.*

doi:10.1111/eva.13081.

Cox, L. M., Yamanishi, S., Sohn, J., Alekseyenko, A. V., Leung, J. M., Cho, I., et al. (2014).

Altering the intestinal microbiota during a critical developmental window has lasting

metabolic consequences. *Cell*. doi:10.1016/j.cell.2014.05.052.

Cui, B., Gai, Z., She, X., Wang, R., and Xi, Z. (2016). Effects of chronic noise on glucose

metabolism and gut microbiota-host inflammatory homeostasis in rats. *Sci. Rep.*

doi:10.1038/srep36693.

Cui, B., Su, D., Li, W., She, X., Zhang, M., Wang, R., et al. (2018). Effects of chronic noise

exposure on the microbiome-gut-brain axis in senescence-accelerated prone mice:

Implications for Alzheimer's disease. *J. Neuroinflammation*. doi:10.1186/s12974-018-1223-

4.

Davidson, G. L., Wiley, N., Cooke, A. C., Johnson, C. N., Fouhy, F., Reichert, M. S., et al.

(2020). Diet induces parallel changes to the gut microbiota and problem solving

performance in a wild bird. *Sci. Rep.* doi:10.1038/s41598-020-77256-y.

Ditchkoff, S. S., Saalfeld, S. T., and Gibson, C. J. (2006). Animal behavior in urban ecosystems:

Modifications due to human-induced stress. *Urban Ecosyst.* doi:10.1007/s11252-006-3262-

3.

Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST.

Bioinformatics. doi:10.1093/bioinformatics/btq461.

Fischer, J., and Lindenmayer, D. B. (2007). Landscape modification and habitat fragmentation:

A synthesis. *Glob. Ecol. Biogeogr.* doi:10.1111/j.1466-8238.2007.00287.x.

Fontaine, S. S., Novarro, A. J., and Kohl, K. D. (2018). Environmental temperature alters the

digestive performance and gut microbiota of a terrestrial amphibian. *J. Exp. Biol.*

doi:10.1242/jeb.187559.

Fox, J., Weisberg, S., Bates, D., Firth, D., Friendly, M., Gor-, G., et al. (2012). The car Package.

R.

Fuirst, M., Veit, R. R., Hahn, M., Dheilly, N., and Thorne, L. H. (2018). Effects of urbanization

on the foraging ecology and microbiota of the generalist seabird Larus argentatus. *PLoS*

One. doi:10.1371/journal.pone.0209200.

Gadau, A., Crawford, M. S., Mayek, R., Giraudeau, M., McGraw, K. J., Whisner, C. M., et al.

(2019). A comparison of the nutritional physiology and gut microbiome of urban and rural

house sparrows (Passer domesticus). *Comp. Biochem. Physiol. Part - B Biochem. Mol. Biol.*

doi:10.1016/j.cbpb.2019.110332.

Garrett, W. S., Gallini, C. A., Yatsunenko, T., Michaud, M., Dubois, A., Delaney, M. L., et al.

(2010). Enterobacteriaceae Act in concert with the gut microbiota to induce spontaneous

and maternally transmitted colitis. *Cell Host Microbe*. doi:10.1016/j.chom.2010.08.004.

Gould, A. L., Zhang, V., Lamberti, L., Jones, E. W., Obadia, B., Korasidis, N., et al. (2018).

Microbiome interactions shape host fitness. *Proc. Natl. Acad. Sci. U. S. A.*

doi:10.1073/pnas.1809349115.

Grimm, N. B., Faeth, S. H., Golubiewski, N. E., Redman, C. L., Wu, J., Bai, X., et al. (2008).
Global change and the ecology of cities. *Science.* doi:10.1126/science.1150195.

Hansell, M., and Overhill, R. (2000). *Bird Nests and Construction Behaviour.*
doi:10.1017/cbo9781139106788.

Hieke, A.-S. C. S. C., Hubert, S. M., and Athrey, G. (2019). Circadian disruption and divergent
microbiota acquisition under extended photoperiod regimens in chicken. *PeerJ.*
doi:10.7717/peerj.6592.

Hooper, L. V., and MacPherson, A. J. (2010). Immune adaptations that maintain homeostasis
with the intestinal microbiota. *Nat. Rev. Immunol.* doi:10.1038/nri2710.

Injaian, A. S., Taff, C. C., and Patricelli, G. L. (2018). Experimental anthropogenic noise impacts
avian parental behaviour, nestling growth and nestling oxidative stress. *Anim. Behav.*
doi:10.1016/j.anbehav.2017.12.003.

Isaksson, C. (2015). Urbanization, oxidative stress and inflammation: A question of evolving,
acclimatizing or coping with urban environmental stress. *Funct. Ecol.* doi:10.1111/1365-
2435.12477.

Jarrett, C., Powell, L. L., McDevitt, H., Helm, B., and Welch, A. J. (2020). Bitter fruits of hard
labour: diet metabarcoding and telemetry reveal that urban songbirds travel further for
lower-quality food. *Oecologia.* doi:10.1007/s00442-020-04678-w.

Jiang, J., He, Y., Kou, H., Ju, Z., Gao, X., and Zhao, H. (2020). The effects of artificial light at
night on Eurasian tree sparrow (Passer montanus): Behavioral rhythm disruption, melatonin
suppression and intestinal microbiota alterations. *Ecol. Indic.*
doi:10.1016/j.ecolind.2019.105702.

Jones, E. L., and Leather, S. R. (2012). Invertebrates in urban areas: A review. *Eur. J. Entomol.*
doi:10.14411/eje.2012.060.

Kight, C. R., and Swaddle, J. P. (2011). How and why environmental noise impacts animals: An
integrative, mechanistic review. *Ecol. Lett.* doi:10.1111/j.1461-0248.2011.01664.x.

Knutie, S. A., Chaves, J. A., and Gotanda, K. M. (2019). Human activity can influence the gut
microbiota of Darwin's finches in the Galapagos Islands. *Mol. Ecol.*

doi:10.1111/mec.15088.

Knutie, S. A., Gabor, C. R., Kohl, K. D., and Rohr, J. R. (2018). Do host-associated gut
microbiota mediate the effect of an herbicide on disease risk in frogs? *J. Anim. Ecol.*
doi:10.1111/1365-2656.12769.

Kohl, K. D., Amaya, J., Passement, C. A., Dearing, M. D., and Mccue, M. D. (2014). Unique and
shared responses of the gut microbiota to prolonged fasting: A comparative study across
five classes of vertebrate hosts. *FEMS Microbiol. Ecol.* doi:10.1111/1574-6941.12442.

Kohl, K. D., Cary, T. L., Karasov, W. H., and Dearing, M. D. (2015). Larval exposure to
polychlorinated biphenyl 126 (PCB-126) causes persistent alteration of the amphibian gut
microbiota. *Environ. Toxicol. Chem.* doi:10.1002/etc.2905.

Kohl, K. D., and Yahn, J. (2016). Effects of environmental temperature on the gut microbial
communities of tadpoles. *Environ. Microbiol.* doi:10.1111/1462-2920.13255.

Kyba, C. C. M., Garz, S., Kuechly, H., de Miguel, A. S., Zamorano, J., Fischer, J., et al. (2015).
High-resolution imagery of earth at night: New sources, opportunities and challenges.
Remote Sens. doi:10.3390/rs70100001.

Ley, R. E., Hamady, M., Lozupone, C., Turnbaugh, P. J., Ramey, R. R., Bircher, J. S., et al.
(2008). Evolution of mammals and their gut microbes. *Science.* 320, 1647–51.
doi:10.1126/science.1155725.

Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* doi:10.1186/s13059-014-0550-8.

Lozupone, C. A., Hamady, M., Kelley, S. T., and Knight, R. (2007). Quantitative and qualitative
β diversity measures lead to different insights into factors that structure microbial
communities. *Appl. Environ. Microbiol.* doi:10.1128/AEM.01996-06.

Lucass, C., Eens, M., and Müller, W. (2016). When ambient noise impairs parent-offspring
communication. *Environ. Pollut.* doi:10.1016/j.envpol.2016.03.015.

Maraci, Ö., Antonatou-Papaioannou, A., Jünemann, S., Castillo-Gutiérrez, O., Busche, T.,
Kalinowski, J., et al. (2021). The Gut Microbial Composition Is Species-Specific and
Individual-Specific in Two Species of Estrildid Finches, the Bengalese Finch and the Zebra

Finch. *Front. Microbiol.* doi:10.3389/fmicb.2021.619141.

Marchesi, J. R., and Ravel, J. (2015). The vocabulary of microbiome research: a proposal.
Microbiome. doi:10.1186/s40168-015-0094-5.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal. doi:10.14806/ej.17.1.200.

Martiny, J. B. H., Bohannan, B. J. M., Brown, J. H., Colwell, R. K., Fuhrman, J. A., Green, J. L.,
et al. (2006). Microbial biogeography: Putting microorganisms on the map. *Nat. Rev.*
Microbiol. doi:10.1038/nrmicro1341.

Maziarz, M., Broughton, R. K., and Wesołowski, T. (2017). Microclimate in tree cavities and
nest-boxes: Implications for hole-nesting birds. *For. Ecol. Manage.*
doi:10.1016/j.foreco.2017.01.001.

McFall-Ngai, M. M., Hadfield, M. G., Bosch, T. T. C. G. T., Carey, H. V. H., Domazet-Lošo, T.,
Douglas, A. A. E., et al. (2013). Animals in a bacterial world, a new imperative for the life
sciences. *Proc. Natl. Acad. Sci.* doi:10.1073/pnas.1218525110.

McKinney, M. L. (2006). Urbanization as a major cause of biotic homogenization. *Biol.*
Conserv. doi:10.1016/j.biocon.2005.09.005.

Miles, L. S., Breitbart, S. T., Wagner, H. H., and Johnson, M. T. J. (2019). Urbanization shapes
the ecology and evolution of plant-arthropod herbivore interactions. *Front. Ecol. Evol.*
doi:10.3389/fevo.2019.00310.

Murray, M. H., Lankau, E. W., Kidd, A. D., Welch, C. N., Ellison, T., Adams, H. C., et al.
(2020). Gut microbiome shifts with urbanization and potentially facilitates a zoonotic
pathogen in a wading bird. *PLoS One*. doi:10.1371/journal.pone.0220926.

Nottingham, A. T., Bååth, E., Reischke, S., Salinas, N., and Meir, P. (2019). Adaptation of soil
microbial growth to temperature: Using a tropical elevation gradient to predict future
changes. *Glob. Chang. Biol.* doi:10.1111/gcb.14502.

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., Mcglinn, D., et al. (2019).
Package "vegan" Title Community Ecology Package Version 2.5-6.

Oro, D., Genovart, M., Tavecchia, G., Fowler, M. S., and Martínez-Abraín, A. (2013).

Ecological and evolutionary implications of food subsidies from humans. *Ecol. Lett.*
doi:10.1111/ele.12187.

Ouyang, J. Q., Davies, S., and Dominoni, D. (2018). Hormonally mediated effects of artificial
light at night on behavior and fitness: linking endocrine mechanisms with function. *J. Exp.*
Biol. doi:10.1242/jeb.156893.

Paulson, J. (2014). metagenomeSeq: Statistical analysis for sparse high-throughput sequencing.
Bioconductor.Jp.

Pfennig, D. W., Wund, M. A., Snell-Rood, E. C., Cruickshank, T., Schlichting, C. D., and
Moczek, A. P. (2010). Phenotypic plasticity's impacts on diversification and speciation.
Trends Ecol. Evol. doi:10.1016/j.tree.2010.05.006.

Phillips, J. N., Berlow, M., and Derryberry, E. P. (2018). The Effects of Landscape Urbanization
on the Gut Microbiome: An Exploration Into the Gut of Urban and Rural White-Crowned
Sparrows. *Front. Ecol. Evol.* doi:10.3389/fevo.2018.00148.

Pollock, C. J., Capilla-Lasheras, P., McGill, R. A. R., Helm, B., and Dominoni, D. M. (2017).
Integrated behavioural and stable isotope data reveal altered diet linked to low breeding
success in urban-dwelling blue tits (Cyanistes caeruleus). *Sci. Rep.* doi:10.1038/s41598-
017-04575-y.

Powell, C., Lill, A., and Johnstone, C. P. (2014). Body Condition and Chronic stress in Urban
and Rural Noisy Miners. *Open Ornithol. J.* doi:10.2174/1874453201306010025.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., et al. (2013). The SILVA
ribosomal RNA gene database project: Improved data processing and web-based tools.
Nucleic Acids Res. doi:10.1093/nar/gks1219.

R Core Team (2020). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. Available at: http://www.r-project.org/index.html.

Sackey, B. A., Mensah, P., Collison, E., and Sakyi-Dawson, E. (2001). Campylobacter,
Salmonella, Shigella and Escherichia coli in live and dressed poultry from metropolitan
Accra. *Int. J. Food Microbiol.* doi:10.1016/S0168-1605(01)00595-5.

Schiraldi, C., and De Rosa, M. (2014). "Mesophilic Organisms," in *Encyclopedia of Membranes*

doi:10.1007/978-3-642-40872-4_1610-2.

Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., et al.
(2009). Introducing mothur: Open-source, platform-independent, community-supported
software for describing and comparing microbial communities. *Appl. Environ. Microbiol.*
doi:10.1128/AEM.01541-09.

Schroeder, J., Nakagawa, S., Cleasby, I. R., and Burke, T. (2012). Passerine birds breeding under
chronic noise experience reduced fitness. *PLoS One*. doi:10.1371/journal.pone.0039200.

Sepulveda, J., and Moeller, A. H. (2020). The Effects of Temperature on Animal Gut
Microbiomes. *Front. Microbiol.* doi:10.3389/fmicb.2020.00384.

Shannon, C. E. (1997). The Mathematical Theory of Communication. *M.D. Comput.*
doi:10.2307/410457.

Slabbekoorn, H. (2013). Songs of the city: Noise-dependent spectral plasticity in the acoustic
phenotype of urban birds. *Anim. Behav.* doi:10.1016/j.anbehav.2013.01.021.

Sommer, F., and Bäckhed, F. (2013). The gut microbiota-masters of host development and
physiology. *Nat. Rev. Microbiol.* doi:10.1038/nrmicro2974.

Sprau, P., Mouchet, A., and Dingemanse, N. J. (2017). Multidimensional environmental
predictors of variation in avian forest and city life histories. *Behav. Ecol.*
doi:10.1093/beheco/arw130.

Sullam, K. E., Essinger, S. D., Lozupone, C. A., O'Connor, M. P., Rosen, G. L., Knight, R., et al.
(2012). Environmental and ecological factors that shape the gut bacterial communities of
fish: A meta-analysis. *Mol. Ecol.* doi:10.1111/j.1365-294X.2012.05552.x.

Szulkin, M., Garroway, C. J., Corsini, M., Kotarba, A. Z., Dominoni, D., Szulkin, M., et al.
(2020). "How to Quantify Urbanization When Testing for Urban Evolution?," in *Urban
Evolutionary Biology* doi:10.1093/oso/9780198836841.003.0002.

Teyssier, A., Lens, L., Matthysen, E., and White, J. (2018a). Dynamics of gut microbiota
diversity during the early development of an avian host: Evidence from a cross-foster
experiment. *Front. Microbiol.* doi:10.3389/fmicb.2018.01524.

Teyssier, A., Matthysen, E., Hudin, N. S., de Neve, L., White, J., and Lens, L. (2020). Diet

contributes to urban-induced alterations in gut microbiota: Experimental evidence from a
wild passerine. *Proc. R. Soc. B Biol. Sci.* doi:10.1098/rspb.2019.2182.

Teyssier, A., Rouffaer, L. O., Saleh Hudin, N., Strubbe, D., Matthysen, E., Lens, L., et al.
(2018b). Inside the guts of the city: Urban-induced alterations of the gut microbiota in a
wild passerine. *Sci. Total Environ.* doi:10.1016/j.scitotenv.2017.09.035.

Tomiałojć, L., and Wesołowski, T. (2004). Diversity of the Białowieza forest avifauna in space
and time. *J. Ornithol.* doi:10.1007/s10336-003-0017-2.

Trevelline, B. K., Fontaine, S. S., Hartup, B. K., and Kohl, K. D. (2019). Conservation biology
needs a microbial renaissance: A call for the consideration of host-associated microbiota in
wildlife management practices. *Proc. R. Soc. B Biol. Sci.* doi:10.1098/rspb.2018.2448.

Trevelline, B. K., MacLeod, K. J., Knutie, S. A., Langkilde, T., and Kohl, K. D. (2018). In ovo
microbial communities: A potential mechanism for the initial acquisition of gut microbiota
among oviparous birds and lizards. *Biol. Lett.* doi:10.1098/rsbl.2018.0225.

Uchida, K., Blakey, R. V., Burger, J. R., Cooper, D. S., Niesner, C. A., and Blumstein, D. T.
(2021). Urban Biodiversity and the Importance of Scale. *Trends Ecol. Evol.*
doi:10.1016/j.tree.2020.10.011.

van Veelen, H. P. J., Falcão Salles, J., Matson, K. D., van der Velde, M., and Tieleman, B. I.
(2020). Microbial environment shapes immune function and cloacal microbiota dynamics in
zebra finches Taeniopygia guttata. *Anim. Microbiome.* doi:10.1186/s42523-020-00039-3.

Videvall, E., Song, S. J., Bensch, H. ., Strandh, M., Engelbrecht, A., Serfontein, N., et al. (2020).
Early-life gut dysbiosis linked to juvenile mortality in ostriches. *BMC Microbiome* 8.
doi:https://doi.org/10.1186/s40168-020-00925-7.

Videvall, E., Strandh, M., Engelbrecht, A., Cloete, S., and Cornwallis, C. K. (2018). Measuring
the gut microbiome in birds: Comparison of faecal and cloacal sampling. *Mol. Ecol. Resour.*
doi:10.1111/1755-0998.12744.

Watson, H., Videvall, E., Andersson, M. N., and Isaksson, C. (2017). Transcriptome analysis of
a wild bird reveals physiological responses to the urban environment. *Sci. Rep.*
doi:10.1038/srep44180.

Wei, L., Yue, F., Xing, L., Wu, S., Shi, Y., Li, J., et al. (2020). Constant Light Exposure Alters
Gut Microbiota and Promotes the Progression of Steatohepatitis in High Fat Diet Rats.
Front. Microbiol. doi:10.3389/fmicb.2020.01975.

Whittaker, R. H. (1960). Vegetation of the Siskiyou Mountains, Oregon and California. *Ecol.*
Monogr. doi:10.2307/1948435.

Wickham, H. (2009). *ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York*.

Wilkin, T. A., King, L. E., and Sheldon, B. C. (2009). Habitat quality, nestling diet, and
provisioning behaviour in great tits Parus major. *J. Avian Biol.* doi:10.1111/j.1600-
048X.2009.04362.x.

World Urbanization Prospects 2018: Highlights (2019). doi:10.18356/6255ead2-en.

Zilber-Rosenberg, I., and Rosenberg, E. (2008). Role of microorganisms in the evolution of
animals and plants: the hologenome theory of evolution. *FEMS Microbiol. Rev.*
doi:10.1111/j.1574-6976.2008.00123.x.

Zollinger, S. A., Dorado-Correa, A., Goymann, W., Forstmeier, W., Knief, U., Bastidas-Urrutia,
A. M., et al. (2019). Traffic noise exposure depresses plasma corticosterone and delays
offspring growth in breeding zebra finches. *Conserv. Physiol.* doi:10.1093/conphys/coz056.

Zuur, A. F., Ieno, E. N., and Elphick, C. S. (2010). A protocol for data exploration to avoid
common statistical problems. *Methods Ecol. Evol.* doi:10.1111/j.2041-210x.2009.00001.x.

12. Figure Captions and Legends

Figure 1. Alterations to the gut microbiota in the urban mosaic. (A) Study site locations in

the urban mosaic of the capital city of Warsaw, Poland. These locations include an office area,

two residential areas, an urban park, two urban woodlands, an urban forest, a peri-urban village

and a natural forest. The impervious surface area (ISA, in %), shown here as the original map

layer, is further described and used for analysis in section 2.2.3. A Google satellite plug-in was

used to zoom in on each study site and visualize the locations of nest boxes (dots) and natural

cavities (triangles). **(B)** The relative abundances of microbial families in gut samples from

different sampling sites located in urban and rural territories. Only the 20 families with the

highest relative abundances are shown. **(CD)** An nMDS plot of the dissimilarities of the gut

microbiota among samples collected from different sampling sites located (**C)** in urban and rural

sites and **(D**) with low (lower than median, 4.43) and high (larger than the median) ISA value**s.**

The distances were computed using the Bray–Curtis dissimilarity index. The lengths of the

vectors are proportional to their predictive strength, determined based on Envfit. All variables

that were significant based on at least one test (*BIOENV*, *Envfit* or Mantel) are represented. The

parameters considered significant in all tests are coloured in red.

Figure 2. Differentially abundant families between **(A)** urban and rural samples and between **(B)**

high and low ISA categories. OTUs with a log2-fold change larger than zero are more abundant

in urban and high-ISA territories (grey bars), while OTUs with a log2-fold change smaller than

zero are more abundant in rural and low-ISA territories (green bars).

1004

1005 **Figure 3.** Summary of the methodological framework and main findings of this study.

Figure 1

Figure 2

Figure 3

Defining urban space based on:	
A - Administrative boundaries (urban / rural)	**B – Impervious Surface Area (ISA)** (ISA fitted as: high/low, or as continuous value)

1. Do microbial communities differ across the urban mosaic?			
Alpha diversity (urban / rural) no significant difference LMM: β=-0.23 ±0.18, p=0.215	**Beta diversity** (urban / rural) Yes: stronger ANOSIM, R=0.180, p=0.001	**Alpha diversity** (ISA) Negative correlation LMM: β=-0.14 ±0.06, p=0.029	**Beta Diversity** (ISA) Yes: weaker ANOSIM, R=0.090, p=0.003

2. Independent environmental & spatial predictors of microbial communities
Predictors: sound pollution, distance to city center, distance to closest road, tree cover,
light pollution, distance to closest walking path, temperature, NDVI , human presence

Alpha diversity: LMM: tree cover density
Beta diversity: MANTEL; ENVFIT; BIOENV: sound pollution, distance to city centre, distance to closest road