Pore dynamics and asymmetric cargo loading in an encapsulin nanocompartment

Jennifer Ross[1,5†], Zak McIver[2], Thomas Lambert[1], Cecilia Piergentili[2], Jasmine Emma Bird[2], Kelly J. Gallagher[1], Faye L. Cruickshank[1], Patrick James[2], Efrain Zarazúa-Arvizu[3], Louise E. Horsfall[3], Kevin J. Waldron[5], Marcus D. Wilson[4], C. Logan Mackay[1], Arnaud Baslé[5], David J. Clarke[1]*, Jon Marles-Wright[2,5†]*

Short title: Haliangium ochraceum encapsulin structure

[1]EaStCHEM School of Chemistry, The University of Edinburgh, Joseph Black Building, David Brewster Road, Edinburgh, EH9 3FJ.
[2]School of Natural and Environmental Sciences, Newcastle University, Newcastle upon Tyne, NE1 7RU.
[3]School of Biological Sciences, The University of Edinburgh, Edinburgh, EH9 3BF.
[4] Wellcome Centre for Cell Biology, University of Edinburgh, Michael Swann Building, Kings Buildings, Mayfield Road, Edinburgh, EH9 3JR, UK
[5]Newcastle University Biosciences Institute, Medical School, Newcastle University, Newcastle upon Tyne, NE2 4HH.

†Current Address: Newcastle University Biosciences Institute, Medical School, Newcastle University, Newcastle upon Tyne, NE2 4HH

* To whom correspondence should be addressed:
Jon Marles-Wright, +44(0)191 2084855, Jon.marles-wright1@ncl.ac.uk
David J. Clarke, +44(0)131 650 4808, Dave.clarke@ed.ac.uk

Keywords:
Encapsulin, encapsulated ferritin, ferritin, ferroxidase, *Haliangium ochraceum*, cryo-EM, hydrogen-deuterium exchange (HDX), mass spectrometry (MS), metalloprotein, protein structure.

Abstract

Encapsulins are protein nanocompartments that house various cargo enzymes, including a family of decameric ferritin-like proteins. Here, we study a recombinant *Haliangium ochraceum* encapsulin:encapsulated ferritin complex using electron cryo-microscopy and hydrogen/deuterium exchange mass spectrometry to gain insight into the structural relationship between the encapsulin shell and its protein cargo. An asymmetric single particle reconstruction reveals four encapsulated ferritin decamers in a tetrahedral arrangement within the encapsulin nanocompartment. This leads to a symmetry mismatch between the protein cargo and the icosahedral encapsulin shell. The encapsulated ferritin decamers are offset from the interior face of the encapsulin shell. Using HDX-MS, we observed dynamic behavior of the major five-fold pore in the encapsulin shell and show the pore opening via the movement of the encapsulin A-domain. These data will accelerate efforts to engineer the encapsulation of heterologous cargo proteins and to alter the permeability of the encapsulin shell via pore modifications.

Teaser

Cryo-EM and HDX-MS analysis of an encapsulin nanocompartment shows that the pores at the five-fold icosahedral vertex of the shell are flexible.

Introduction

Cellular metabolism and reaction pathways can produce toxic by-products which damage proteins, DNA, and lipids, or can become involved in potentially harmful side-reactions. Eukaryotes use membrane-bound organelles, such as lysosomes, to prevent this damage by housing dangerous reactions in chemically privileged environments. In a similar manner, prokaryotes use large protein-based compartments to sequester such reactions and act as barrier from the cytosol [1, 2]. Prokaryotes have a variety of metabolic compartments including carboxysomes, which are used for carbon dioxide fixation in photosynthetic and some chemotrophic bacteria[3, 4]; the related bacterial microcompartments, which allow the catabolism of carbon sources that produce aldehyde intermediates in their breakdown, such as ethanolamine and propanediol[5]; and ferritins, used for iron oxidation and storage [6, 7].

One other key compartmentalization strategy utilized by prokaryotes and archaea is the encapsulin system [8–10]. Encapsulin (Enc) nanocompartments are hollow icosahedral complexes which range in size from 20 nm to 42 nm [9, 11, 12]. Encapsulin proteins are structurally related to the viral capsid protein (gp5) of the HK97 bacteriophage and self-assemble from a single monomer into one of three forms: 60 subunits ($T = 1$ capsid symmetry), 180 subunits ($T = 3$ capsid symmetry) or 240 subunits ($T = 4$ symmetry)[8, 9, 12, 13]. Encapsulins share a common feature of housing a cargo enzyme, such as ferritin-like proteins (encapsulated ferritins, EncFtn), iron-mineralizing encapsulin-associated firmicute (IMEF), or dye-decolorizing peroxidases[12, 14]. Cargo enzymes are directed inside the encapsulin nanocompartment by a terminal localization sequence (LS) which binds to the interior face of the encapsulin[9, 15]. Encapsulins and their cargo proteins are found throughout the bacterial and archaeal domains in species inhabiting a range of environmental niches; consequently, the proteins are stable in diverse physical conditions[8, 16–19]. For these reasons, the encapsulins have attracted considerable interest for biotechnological applications, through their ability to separate potentially hazardous heterologous reactions from the native cytosol[20, 21]. The expression of functional engineered encapsulins has also been demonstrated in eukaryotes, where they have been developed as metabolic organelles in *Saccharomyces cerevisiae*[21] and in cellular imaging applications in human cells[22].

The EncFtn cargo proteins are of particular interest, as they differ from their classical ferritin relatives. Although both proteins oxidize iron using a conserved catalytically active ferroxidase center (FOC), they have remarkably different structural architectures. Classical ferritins oxidize ferrous iron, Fe(II), into a mineral ferric form, Fe(III), which is then stored within a 24-meric 12 nm nanocage [6, 23]. In contrast to this, the EncFtn proteins have an annular structure formed from a pentamer of dimers with the FOC active sites located at a dimer interface [11, 16]. EncFtn oxidizes iron in a similar manner to other ferritins, but due to its open structure, it must be associated with an encapsulin nanocage to act as an iron store [11]. Together the encapsulin EncFtn (Enc:EncFtn) complex can perform both the oxidation and storage functions of classical ferritins. However, due to its increased size when compared to classical ferritins, the Enc:EncFtn complex has the potential to house significantly greater quantities of iron, and has been described as an iron megastore [13]. For these reasons, the encapsulins have attracted considerable interest for biotechnological applications, through their ability to separate potentially hazardous heterologous reactions from the host cytosol.

Although there have been several structural studies on encapsulins, several key
questions remain unanswered. Most notably, for the EncFtn containing encapsulin
nanocompartments, the structural relationship between the encapsulin shell and the
EncFtn cargo protein is unknown. Studies on Dyp loaded encapsulin
nanocompartments have shown loading with either one(*24*) or two hexameric
complexes(*25*). Previous models for the loading of EncFtn in encapsulin
nanocompartments have suggested a symmetric arrangement of the D5 decameric
encapsulated ferritin at the five-fold icosahedral vertices of the encapsulin shell,
giving a theoretical maximum of twelve EncFtn decamers per encapsulin
nanocage(*11*, *26*).
Herein, we investigate the structure of the Enc:EncFtn nanocompartment from
the halophilic bacterium *Haliangium ochraceum,* to gain a better understanding of
the arrangement and stoichiometry of the complex. Using the complementary
structural biology techniques of cryogenic electron microscopy (cryo-EM) and
hydrogen-deuterium exchange mass spectrometry (HDX-MS), we study the
structural relationship between Enc and EncFtn in the encapsulin nanocage.
We present the first cryo-EM structure of a mesophilic Enc:EncFtn
nanocompartment, which reveals a symmetry-breaking tetrahedral arrangement of
the EncFtn decamers within the encapsulin shell. Analysis of the encapsulin shell by
symmetry expansion of an icosahedral reconstructions and focused 3D refinement
on the pentameric vertex reveals a flexible pore in the shell. The dynamic nature of
this pore region was investigated by HDX-MS. We show that this region has a high
rate of H/D exchange, demonstrating its conformational flexibility. Our combination of
HDX-MS and cryo-EM models affords insight into the loading capacity and dynamics
of the Enc:EncFtn nanocompartment system.
138

Results

Recombinant *Haliangium ochraceum* encapsulin complexes form regular nanocompartments that recruit active EncFtn cargoes

In order to gain an understanding of the relationship between encapsulin nanocompartments and their EncFtn cargoes, the Enc:EncFtn nanocompartment from the halophilic mesophile *Haliangium ochraceum* was chosen as our model system. This was primarily due to the high yields and ease of purification of the recombinant nanocompartment. Constructs to produce empty (Empty-Enc) and EncFtn loaded (Loaded-Enc) encapsulin nanocompartments were assembled for recombinant protein expression in *Escherichia coli*. The protein complexes were purified by heat treatment, followed by anion exchange and size-exclusion chromatography (**Figure 1A and B, and Figure S1A-D**).

Size-exclusion chromatography was used to separate correctly formed encapsulin nanocompartments from contaminating proteins and partially formed sub-complexes (**Figure S1A/B**). Fractions corresponding to the major peaks were visualized by negative stain transmission electron microscopy (TEM) (**Figure S2**), with the fraction corresponding to a retention volume of 63 mL for both the Empty-Enc and Loaded-Enc containing the homogeneous encapsulin nanocompartments (**Figure 1C**). Both empty and EncFtn-loaded encapsulins assembled into regular nanocompartments, with an average diameter of approximately 21 nm, consistent with other T=1 type encapsulins (**Figure S1E and F**) (*9, 25, 27, 28*). The micrographs of the Loaded-Enc sample reveal a regular internal density visible within the nanocompartment, suggesting that the EncFtn cargo has been encapsulated in an organized manner.

The molecular masses of the protein constituents of the Empty-Enc and Loaded-Enc nanocompartments were determined by LC-MS (**Table S1**). MS analysis of the Empty-Enc assembly revealed a single charge state distribution corresponding to a monomer of the encapsulin protein. MS analysis of the Loaded-Enc revealed three charge state distributions present, with deconvoluted masses consistent with the encapsulin protein monomer, a monomer of EncFtn and a dimer of EncFtn (**Figure S1G and H**) (*29*)**.** These results indicate that the Loaded-Enc sample contained both the encapsulin and EncFtn cargo proteins, whilst Empty-Enc has only the encapsulin protein.

Ferroxidase assays were performed on the samples to confirm the ability of the EncFtn loaded encapsulin nanocompartment to convert Fe(II) to Fe(III) (**Figure 1D**). This result is consistent with our previous observations for the *Rhodospirillum rubrum* Enc:EncFtn encapsulin complex(*11*). The empty encapsulin, which lacks the EncFtn cargo is enzymatically inactive.

Taken together, these data demonstrate that functionally active EncFtn has been successfully loaded into the encapsulin nanocompartment during expression in the heterologous *E. coli* host.

The cryo-EM structure of the Loaded-Enc nanocompartment

Motivated by the apparent interior density in the Loaded-Enc sample, we performed single particle cryo-EM on the Loaded-Enc nanocompartment complex (**Table S2**). Consistent with previously published X-ray crystallographic and cryo-EM derived encapsulin models, an initial reconstruction was produced using Relion3.1 (*30*) with imposed I1 symmetry (**Figure 2, Figure S3**, **Figure S4** and **Table S3**) (*8, 9, 12, 27, 31*). This resulted in a reconstruction with a global resolution of 2.5 Å as determined by the gold-standard Fourier shell correlation at 0.143 (GS-FSC) (**Figure S3D**). The

reconstruction displays a *T* = 1 icosahedral arrangement of sixty encapsulin
monomers, with clearly resolved secondary structure elements. Small pores are
visible in the shell at, or close to, the 2-, 3-, and 5-fold symmetry axes where protein
monomers interact with each other (**Figure 2A**). Regions of the reconstruction
around the icosahedral five-fold axes displayed a lower resolution than the other
regions of the structure in a local resolution map (**Figure 2B**). This is consistent with
observations of local resolution maps for the *Quasibacillus thermotolerans*
encapsulin reconstruction(*12*). The monomer of the encapsulin nanocompartment
from the reconstruction displays a HK97-fold typical for encapsulins (**Figure 2C**),
with the position of the E-loop broadly determining the topology of the encapsulin
nanocage. In the *T* = 1 Family I encapsulins this loop is at 45° to the boundary
between the P- and A-domains (**Figure 2C**); whereas in the *T* = *3* HK97, and *T* = *3*
and *T* = *4* encapsulins, it is found in close apposition to the A-domain. The recently
described family II encapsulin from *Synechococcus elongatus*(*31*) has an E-loop
topology consistent with the *T* = *3* and *T* = *4* encapsulins, but adopts a *T* = *1*
icosahedral nanocage. The shift in E-loop position relative to other *T* = *1* encapsulins
is a result of an extended N-terminal region, that extends beneath the loop, this
forces the pentameric vertices into a raised morphology consistent with the larger
encapsulin nanocages.

The encapsulin nanocompartment recruits four EncFtn decamers to its lumen
With the imposition of I1 symmetry on the reconstruction, the EncFtn cargo is
not visible within the lumen. This suggests that the organization of the EncFtn protein
within the encapsulin shell does not conform to icosahedral symmetry and is
rotationally averaged through our symmetry-imposed processing. To gain insight into
the structural relationship between the encapsulin shell and its EncFtn cargo protein,
a reconstruction was produced with no imposed symmetry averaging (**Figure 3 and**
Figure S5). Separation of the dataset into five 3D classes revealed a highly
populated class with amorphous density in the interior **(Figure S4, panel 5B)**, and
two other main classes, both containing four distinct densities, arranged in a similar
tetrahedral fashion within the encapsulin shell. To obtain the clearest reconstruction
of both the encapsulin nanocompartment and EncFtn cargoes we took only one of
these latter classes forward for full 3D refinement in Relion3.1 to produce a final C1
reconstruction with a resolution of 3.7 Å at the 0.143 FSC threshold (**Figure 3A and**
Figure S5A). Interestingly, the EM map revealed four distinct densities within the
encapsulin nanocompartment lumen, which are consistent in size and shape with
four EncFtn decamers (**Figure 3B** and **Figure S5B**).
A local resolution map calculated for the asymmetric reconstruction indicates a
degree of flexibility at the pentameric pores compared to the trimeric pores of the
encapsulin shell (**Figure 3A**). The interior of the nanocompartment shows a
significant falloff in resolution from the inner face of the encapsulin shell to the
EncFtn densities. This is consistent with the tethering of the EncFtn to the encapsulin
nanocompartment via its localization sequence, with some degree of conformational
freedom of the EncFtn decamer with respect to the encapsulin shell.
The EncFtn decamers are located approximately 3 nm away from the encapsulin
interior wall, which corresponds to the linker region between the main EncFtn
domain and the localization sequence on the EncFtn C-terminus (**Figure S5C**). The
extended localization sequence of the EncFtn protein acts to offset it from the inner
face of the encapsulin shell, an observation consistent with previous reports of the

IMEF encapsulin complex from *Q. thermotolerans*(*12*). Due to the dynamic nature of
the EncFtn within the encapsulin it was not possible to trace the path of the
localization sequence to its binding site. Interestingly, the four EncFtn decamers are
in a tetrahedral arrangement within the encapsulin nanocompartment, with the five-
fold axes of the EncFtn decamers aligned to the three-fold tetrahedral axes (**Figure
3C and Figure S5C**). This results in a double symmetry mismatch between the
icosahedral shell and the EncFtn decamers in the complex. A concurrent structural
study of the *Thermotoga maritima* encapsulin complex(*32*) revealed five EncFtn
decamers within the encapsulin shell, with each decamer found in approximation to a
pentameric vertex. The overall arrangement of the five EncFtn complexes within the
nanocompartment is incompatible with the formation of a regular platonic solid and
results in a symmetry breaking arrangement of the EncFtn within the encapsuling
nanocompartment. To enhance the resolution of the individual EncFtn density,
symmetry expansion and local refinement focused on a single EncFtn density was
performed in CryoSPARC. This resulted in an EM map into which the *H. ochraceum*
EncFtn crystal structure (PDB ID: 5N5F)(*16*) could be docked (**Figure 3D**).
Analysis of the relationship between the EncFtn decamers and the inner face of
the *H. ochraceum* encapsulin shell reveals distinct EncFtn environments (**Figure
S6**). The first EncFtn environment is shared by two EncFtn decamers and is in line
with the five-fold pore of the encapsulin nanocompartment (EncFtn 1 and 2 in **Figure
S6**). The shared symmetry of the Enc nanocompartment five-fold pores and of the
EncFtn D5 annular structure in these positions is consistent with our previously
proposed hypothesis for the Enc:EncFtn relationship(*11*) and is also found in the *T.
maritima* encapsulin. However, the symmetry-breaking tetrahedral arrangement of
the EncFtn decamers in the *H. ochraceum* encapsulin creates a second distinct
environment shared by the remaining two EncFtn decamers, where they are offset
between five-fold and three-fold axes of the icosahedral encapsulin shell (EncFtn 3
and 4 in **Figure S6**). With the proposed route of iron entry through the 5-fold pores of
the encapsulin shell, the EncFtn decamers in proximity to the pores may have more
favorable substrate access than those found in the alternative positions. Given the
fast mass-transport of substrates across the shell(*33*), this symmetry breaking
arrangement of the EncFtn decamers is unlikely to have significant functional
consequences in terms of the speed of iron oxidation in the different EncFtn
environments.
272

Structural dynamics in the pentameric vertices of the encapsulin shell
To further investigate the apparent conformational flexibility of the encapsulin
shell at the pentameric vertices in the reconstructions, we analyzed the full data set
in CryoSPARC to take advantage of recent advances in 3D variability analysis
(3DVA) of single-particle datasets(*34*). A C1 refinement calculated with 333,945
particles performed in CryoSPARC (**Table S4**) was subjected to 3DVA with three
variability components. The three components resolve to show the five-fold pores
irising between 'open' and 'closed' states. The three components primarily differ in
which pores are open/closed and show no other major differences. Adjacent five-fold
pores show a mixture of correlated and anti-correlated opening (**Figure 4, Figure S7**
and **Supplementary Movies 1-3**). This movement gives the impression of a
pumping motion, which may enhance the transfer of substrates between the exterior
and interior lumen of the nanocompartment.
Within the encapsulin nanocompartment, the EncFtn densities show some
uncoordinated lateral motions, but their position relative to the inner face of the

encapsulin does not change. This is because they are tethered by their localization
sequences to the P-domain of the encapsulin, which does not show any significant
displacement in any of the variability components.
To capture the extreme 'open' and 'closed conformations of the five-fold pores we
performed symmetry expansion on the I1 particle set in Relion3.1, followed by
masked 3D-classification without alignment centered on the vertex. Of the five
conformations observed, the most extreme of these were subjected to 3D refinement
with local searches (**Figure 5**, **Figure S8**, and **Figure S9**). This resulted in an 'open'
pentamer conformation of 2.4 Å resolution and a 'closed' conformation of 2.3 Å,
allowing for fitting of residue side chains (**Figure S7**).
The 'open' conformation has a five-fold pore with an aperture diameter of
approximately 15 Å, while in the 'closed' conformation the aperture is reduced to 5 Å
diameter. To understand the structural changes taking place in the transition
between these conformations, an atomic model of the encapsulin protein was refined
against both maps (**Table S5**). The two models show a significant movement in the
A-domain, with the A-domain pivoting at the loops connecting it to the P-domain
(residues 121 – 131 and 201-222) (**Figure S10A/B**), opening the pore like an iris
(**Figure 5**). In the open conformation, the pore loop region (residues 182 - 189) is not
well defined in the density; while it is tightly locked in the closed conformation, with
Asp186 forming the outer boundary of the pore and Tyr188 and Lys192 forming the
inner bounds (**Figure S10C**). The tyrosine is well conserved among the family 1 T=1
encapsulins, while the lysine is substituted for a glutamine in the *R. rubrum*
encapsulin (**Figure S11**). The family 2 T=1 encapsulin from *Synechococcus*
elongatus has a five-residue sequence insertion in this region, which forms an
extended linker between secondary structure elements, rather than a distinct loop
within the pore.
Additionally, our focused refinements of the pentameric subunits allowed us to
build and sequence the $_{117}$GSLGIGSLR$_{125}$ peptide from the EncFtn protein (**Figure
6**). This region of the localization sequence forms a network of hydrophobic
interactions with the inner face of the P-domain of a single encapsulin monomer, with
further stabilization by several water-mediated backbone contacts. The core
GxLGIxxL motif found in this region of the localization sequence is conserved
between the *H. ochraceum* EncFtn and other proteins in the family and is observed
in the crystal structure of the *T. maritima* encapsulin(*9*).
In the *H. ochraceum* encapsulin the five-fold pore has a negative charge on the
exterior of the encapsulin shell and positive charge on the interior in both the open
and closed conformations (**Figure S12**). No changes are observed in the structure,
or charge of the 3- and 2-fold pores (**Figure S12**). The closed conformation of the 5-
fold pore is consistent with observations from the crystal structure of the *T. maritima*
encapsulin(*9*) and high resolution cryo-EM structures of other encapsulins(*12*, *31*,
32). However, this is the first time that an 'open' pore-conformation has been
observed in an encapsulin protein.
This observation has important implications for efforts to engineer the pores of
encapsulin nanocages. Efforts to widen the five-fold pores have demonstrated an
increase in mass-transport of model substrates and ions across the encapsulin
shell(*33*, *35*); however, the observed changes in pore diameters are limited to a
maximum of around 10 Å in the *T. maritima* encapsulin(*35*). Interestingly, the
encapsulin shell appears to interact strongly with Tb^{3+} ions in molecular dynamics
simulations, this is consistent with our previous observations of iron binding by the *R.*

rubrum encapsulin(*11*) and highlights a possible cooperative role for substrate
channelling to the EncFtn cargoes.
Our results suggest that the flexibility of the A-domain should also be considered
when engineering the pore structure for enhanced transport across the encapsulin
shell. The loops connecting the A-domain to the P-domain present ideal sites for
targeted mutagenesis to stiffen, or lock, the position of the A-domain and thus
constrain the five-fold pore loop to engineer enhanced substrate selectivity into these
systems.
345
Dynamics of the 5-fold encapsulin pore through hydrogen/deuterium exchange
mass spectrometry.
To further investigate the dynamic nature of the five-fold pore of the
encapsulin shell and the docking of the EncFtn localization sequence to the interior
of the encapsulin nanocompartment, we performed hydrogen/deuterium exchange
mass spectrometry (HDX-MS) on both Empty-Enc and Loaded-Enc
nanocompartments. The extent of backbone-amide hydrogen exchange was
determined at seven time points (0 seconds, 10 seconds, 30 seconds, 5 minutes, 30
minutes, 4 hours, and 24 hours). By calculating the rate of hydrogen exchange
throughout the protein, regions that differ in solvent exposure and/or dynamics can
be detected.
HDX-MS analysis of the encapsulin nanocompartment resulted in 40 pepsin
peptides, which constituted a protein sequence coverage of 85%, with peptide
redundancy of 2.28 (**Table S6**, **Table S7** and **Figure S13**). The encapsulin
nanocompartment displayed variable exchange rates throughout the protein
sequence and regions of the protein displaying elevated H/D exchange rates were
evident. Overlaying these local H/D exchange rates onto the cryo-EM reconstruction
revealed that the regions of highest exchange were located around pentameric
vertices. (**Figure 7**). This was most notable with the peptide spanning the region
between amino acids 180-196, which includes the five-fold pore loop (**Figure S14**).
In contrast, lower rates of HDX are observed at the 2-fold interface and the potential
3-fold pore (**Figure 7**). These findings agree with our cryo-EM structural analyses
and support the proposed conformational flexibility at the 5-fold pore.
Comparison of the H/D exchange rates of Empty-Enc and Loaded-Enc
revealed similar exchange profiles throughout the encapsulin protein sequence,
suggesting that cargo loading has little effect on the overall architecture and
dynamics of the assembled nanocompartment shell (**Table S6**, **Table S7**, and
Figure S15). However, after prolonged exchange times (4 hours), the Loaded-Enc
exhibited areas with a modest reduction in exchange when compared to the Empty-
Enc. Notably, several peptides in the N-terminal region displayed reduced exchange
rates in Loaded-Enc; for example, the peptide covering amino acids 21-37 displayed
almost twelve percent reduction. Mapping the position of this region onto our
encapsulin reconstruction highlights that this peptide is located on the interior face of
the nanocompartment and included the proposed binding site for the localization
sequence of EncFtn (**Figure 7**). A reduction in exchange across in this region is
likely a consequence of shielding by the engaged EncFtn localization sequences.

Discussion
Our cryo-EM reconstruction of an Enc:EncFtn nanocompartment complex reveals
key areas of divergence from a true icosahedral complex with important functional
consequences. The asymmetric reconstruction showed that the encapsulin

nanocompartment sequesters four decamers of EncFtn within its lumen. In our
recombinant system, with EncFtn produced in excess, this likely represents a
maximum loading capacity for the Enc:EncFtn nanocompartments. The symmetry
breaking tetrameric arrangement of the EncFtn decamers within the encapsulin shell
leads to two distinct environments for EncFtn, with two decamers aligned at the five-
fold symmetry axes, and the remaining two residing between three- and five-fold
axes (**Figure S6**). The concurrent observation of five EncFtn decamers within the *T.*
maritima encapsulin nanocompartment, aligned close to the icosahedral five-fold
symmetry axes of the encapsulin shell, highlights differences in cargo loading
between encapsulins from mesophilic and thermophilic bacteria(*32*). The enhanced
cargo loading seen in the *T. maritima* encapsulin may be a consequence of a more
compact and rigid structure adopted by thermophilic proteins, as seen in
comparisons of the crystal structures of the *H. ochraceum* and *Pyrococcus furiosus*
EncFtn proteins(*16*).
The interior volume of the *T=1* encapsulin lumen is around 4000 nm^3, while an
EncFtn decamer is only 120 nm^2; therefore, the additional cargo seen in the *T.*
maritima encapsulin should not significantly impact the iron-storage potential. The
order of magnitude discrepancy in the iron-loading capacity measured for the *R.*
rubrum(*11*) and *T. maritima*(*32*) encapsulins is likely a result of differences in
experimental conditions in different laboratories.
While the volume occupied by the EncFtn decamers represents less than 15 % of
the total lumen of the encapsulin nanocompartments, the consistent observation of a
gap between the encapsulin shell and encapsulin cargo proteins implies that the
loading and capacity of encapsulins is limited by broader steric effects. Further steric
effects would also be found from the unengaged localization sequences at the core
of the nanocompartment. These effects have implications for efforts to target
heterologous proteins to the encapsulin nanocage; effectively setting a limit on the
volume of protein that can be accommodated within, which is much lower than the
total volume of the lumen of the nanocage.
Our observations have functional implications for the oxidation and storage of
iron within the Enc:EncFtn nanocompartment. The EncFtn decamers are in non-
equivalent positions, and thus have different relationships to the pores of the
encapsulin shell. Therefore, if the pores limit the diffusion of substrates, the EncFtn
decamers would be subjected to different chemical environments. Furthermore, both
the engaged and unengaged localization sequences present a 'soft' steric barrier to
the diffusion of substrates. It is notable that the ferroxidase activity of the Enc:EncFtn
complex is significantly higher than the isolated EncFtn protein(*36*). The dynamic
behavior of the encapsulin five-fold pores, which is apparent in our 3DVA movies,
may increase EncFtn activity by enhancing diffusion of substrates into the lumen of
the nanocompartment. While it is not possible to make mechanistic conclusions from
our model, the complex interactions with the components of the encapsulin
nanocompartment clearly enhance the iron oxidation activity of the EncFtn protein.
Our data were collected on iron-free apo-Enc:EncFtn complexes, and thus it is
not possible to infer the nature of the iron mineralization pathway within the
encapsulin nanocage. Further careful work must be performed to titrate iron into the
complex prior to structural analysis to gain insight into the flow of metal ions from the
exterior to the interior of the encapsulin and to determine if metalation influences the
conformational flexibility of the EncFtn within the encapsulin nanocage, as we have
demonstrated for isolated EncFtn proteins(*11, 16, 36*). Finally, the nature of the iron
mineral and its localization within the encapsulin nanocage is still to be determined.

These cryo-EM and HDX-MS data illustrating a highly dynamic five-fold pore in the encapsulin shell, have major implications for efforts to engineer recombinant encapsulins for improved access for both native and non-native substrates. The limitations of previously published studies, where the five-fold pore is modified for altered substrate access can be explained by a highly dynamic pore structure. Our work suggests new hypotheses for engineering pore selectivity, through modifications to the hinge regions between the P- and A-domains, which are responsible for the opening of the pore.

445 Materials and Methods
446
Experimental Design
The objective of this study was to understand the structural relationship
between encapsulins and their EncFtn cargo using cryo-EM to determine the
complex structure and ferroxidase assays for validation of the complex activity. HDX
was utilized to establish differences between solvent accessibility of empty and
loaded encapsulins and garner insight into the impact of cargo on the interior of the
encapsulin nanocompartment.
454
Cloning of encapsulin expression constructs
The *Haliangium ochraceum* encapsulin and encapsulated ferritin protein
expression constructs were based on the Hoch_3836 and Hoch_3837 genes
downloaded from www.kegg.jp and were codon optimized for expression in
Escherichia coli and synthesized as CIDAR MoClo compatible gBlocks by Integrated
DNA Technologies (IDT) (**Table S8**). The gBlocks were assembled into a Level 0
CIDAR MoClo storage vector(*37*), DVA_CD, for subsequent use. The coding
sequences for the encapsulin and the EncFtn were assembled into expression
cassettes in the level 1 backbones DVK_AE and DVK_EF respectively, each with T7
promoter and transcription terminator parts. The resulting expression cassettes were
then combined into the DVA_AF backbone to produce a co-expression plasmid. All
assembled plasmids were sequence verified by Sanger sequencing by Eurofins
Genomics. The protein sequences for each construct are listed in **Table S9**.
468
Protein expression
The Empty-Enc and Loaded-Enc expression plasmids were transformed into
E. coli BL21(DE3) cells and grown overnight at 37 °C on LB-agar plates containing
appropriate selection antibiotics (kanamycin for Empty-Enc and ampicillin for
Loaded-Enc). A single colony of cells was added to 1 L of autoinduction media(*38*)
supplemented (**Table S10)** with appropriate antibiotic and grown for 38 hours at 37
°C with shaking at 200 rpm. Cells were harvested by centrifugation at $12{,}000 \times g$.
476
Encapsulin nanocompartment purification
E. coli cell pellets expressing the empty-Enc and loaded-Enc constructs were
resuspended in $10 \times$ v/w of lysis buffer (20 mM HEPES, pH 8; 2 mM $MgCl_2$; 1 mg/mL
lysozyme, and benzonase, 12.5 – 25 units/mL). Cells were lysed by sonication whilst
on ice; sonication was carried out in six 1-minute cycles (30 seconds sonication, 30
seconds rest). The lysate was clarified by centrifugation at $20{,}000 \times g$ for 1 hour, 4
°C.
The supernatant from cell lysis was heated to 85 °C for 10 minutes in a water
bath and transferred to a 4 °C ice bath for 10 minutes. The supernatant was then
collected after centrifugation at $10{,}000 \times g$ for 1 hour.
Anion exchange chromatography of the clarified supernatant was performed
using a 1 mL HiTrap Q Sepharose FF column from Cytiva on an ÄKTA™ start. The
column and ÄKTA™ start system were equilibrated with QA buffer (20 mM HEPES,
pH 8.0) and the protein sample was loaded. Unbound proteins were removed by
washing with QA buffer. Bound proteins were eluted by QB buffer (20 mM HEPES,
pH 8.0, 1 M NaCl) over a linear gradient of 0-100% QB over 15 column volumes.
Flow-through fractions containing the sample were subjected to SDS-PAGE to

identify those containing the protein of interest. These fractions were pooled and
concentrated using centrifugal concentrators with a 30 kDa nominal molecular weight
cut off (Vivaspin).
Pooled and concentrated samples from the anion exchange step were loaded
on a size exclusion chromatography column (Sephacryl 400, Cytiva) equilibrated
with SEC buffer (20 mM HEPES, pH 8.0, 150 mM NaCl). Fractions eluting from the
column containing the desired protein, as identified by SDS-PAGE were pooled and
concentrated as above. Protein aliquots were flash cooled in liquid nitrogen and
stored at -80 °C (**Figure S1Ai**, **Aii**, **Bi and Bii**). (*11, 16*).
Following purification, aliquots of Empty-Enc and Loaded-Enc were subjected
to a second round of polishing size exclusion chromatography using an Sephacryl
400 column (Cytiva) equilibrated with SEC buffer (20 mM HEPES, pH 8.0, 150 mM
NaCl) (**Figure 1B**). (Experimental data for the size-exclusion chromatography runs
are available at: doi: 10.6084/m9.figshare.16698106)
508
509

Negative stain TEM
Purified encapsulin nanocompartments were initially imaged by negative stain
TEM. Continuous carbon/formvar coated copper grids (200 mesh) were glow-
discharged for 30 seconds using a Pelco glow discharge system. 5 µL Enc was
pipetted onto the glow-discharged grids and excess liquid was removed after 30
seconds with Whatman filter paper (grade 1, diameter 24.0 cm). The grids were
washed with distilled water three times, followed by staining with 2 % uranyl acetate
for 5 seconds. Grids were left to air dry and then imaged with a JEOL JEM-1400
transmission electron microscope. Images were collected with a Gatan CCD
OneView camera and analyzed using FIJI(*39*). (Raw negative stain images are
available at: doi: 10.6084/m9.figshare.16698088)
521

Ferroxidase activity assay
The enzymatic activity of Empty-Enc and Loaded-Enc were assessed by
ferroxidase assay, as previously described (Piergentili, 2020). Fe(II) samples were
prepared by dissolving $FeSO_4.7H_2O$ in HCl 0.1 % (v/v) under anaerobic conditions.
Protein samples were diluted anaerobically in Buffer GF (20 mM HEPES, pH 8.0,
150 mM NaCl) to a final encapsulin monomer concentration of 9 µM to allow
comparison between experiments.
Iron and protein aliquots were added aerobically to a quartz cuvette (Hellma)
resulting in a final concentration of 100 µM iron and 15 µM (Loaded-Enc), or 9 µM
(Empty-Enc). The cuvette was placed in a UV-visible spectrophotometer
(PerkinElmer Lambda 35) and the reaction sample was incubated at 21°C for 50 s to
stabilise. Absorbance at 315 nm was then recorded every second for 1450 s using
the Time-Drive software. A control experiment was conducted by monitoring the
background oxidation by atmospheric oxygen of 100 µM $FeSO_4*7H_2O$ in the
absence of the enzyme. Loaded-Enc experiments were carried out with three
biological replicates. There were six technical replicates for batch one, two
relplicates using batch two and one replicate from batch three. Means and standard
deviations were calculated on the time zero-subtracted progress curves.
(Experimental data for the ferroxidase assays are available at: doi:
10.6084/m9.figshare.16698127)
542
543

Liquid Chromatography Mass Spectrometry

LC-MS experiments were performed on a Synapt G2 Q-ToF instrument
(Waters Corp., Manchester, UK) and an Acquity UPLC equipped with a reverse
phase C4 Aeris Widepore 50 × 2.1 mm HPLC column (Phenomenex, CA, USA).
Mobile phases of; A= water + 0.1% formic acid, and B=acetonitrile + 0.1% formic
acid were used on a ten-minute gradient from 5% B to 95% B. Samples were
analyzed at ~2 µM, and data analysis was performed using MassLynx v4.1 and
MaxEnt deconvolution.

Cryo-EM data collection and analysis

Sample vitrification

Holey grids (gold, 200 mesh, r 2/2 by Quantifoil) were glow-discharged for 30
seconds using a Pelco glow discharge system. The grids were then mounted into a
FEI vitrobot and 4 µL of encapsulin sample (3 mg/mL) was applied. Grids were then
blotted (100% humidity, 8 °C, blot force -5, wait time 10 seconds and blot time of 3
seconds) with Whatman filter paper (grade 1) and flash cooled in liquid ethane,
cooled with liquid nitrogen.

Cryo-EM data collection

Cryo-EM grid screening was performed on a FEI F20 microscope equipped
with a FEG electron source (200 kV) and a TVIPS F816 CMOS detector at the
University of Edinburgh. The dataset used for single particle reconstruction was
obtained at eBIC on a FEI Titan Krios microscope equipped with Gatan K3 camera
(data collection settings are shown in **Table S2**). Alignments, grid transfer and
imaging set up was performed by the eBIC local contact Dr Yun Song. (Cryo-EM
movies are available in the EMPIAR database: EMPIAR-1218).

Single particle reconstruction

Processing steps for the main reconstruction were performed with the
Relion3.1 software package (*30*). Super-resolution movies were binned at 2 × 2
pixels for both I1and C1 and motion corrected using MotionCor2(*40*). Defocus values
of images (summed movies) were determined by CTFFIND4 (*41*) and those with
poor CTF fits, or bad ice, were manually discarded. A template for autopicking was
created using 2D classes from manually picked particles. Autopicked particles were
extracted (using box a box size of 512 pixels and subjected to three rounds of 2D
classification to remove bad particles. An initial 3D model was created from particles
selected from 2D classes. 3D classes were generated both with and without
icosahedral symmetry imposed ('I1' and 'C1' symmetry). The best class from each
was taken forward for 3D refinement and then CTF refinement followed by further
rounds of 3D refinement and Bayesian polishing. After a final round of 3D
refinement, postprocessing was performed using a soft spherical mask. Local
resolution estimation was performed in Relion3.1. The data processing and
refinement pipeline is shown in **Figure S3** with data processing and refinement
statistics in **Table S3**.

3D variability analysis was performed using CryoSPARC(*32*, *41*). Super-
resolution movies were patch motion corrected and patch CTF estimated. Initally,
particles were picking using blob picker (250-300 Å) and these particles were
subjected to 2D classification followed by a round of ab initio model generation to
create an autopick template. Successive rounds of rounds of 2D classification were

performed on the autopicked particles until less than 5% of the particles were
rejected. Five volumes were created by the ab initio model generation algorithm, and
these were heterogeneously refined. The volume with the most particles (99.4 %)
was homogeneously refined and subsequently subjected to 3D variability
analysis(*34*). The homogenously refined particles were also used to enhance the
resolution of the EncFtn cargo. Particles were subjected to symmetry expansion, and
then an individual EncFtn was masked and subjected to local refinement. The data
processing and refinement performed in CryoSPARC pipeline is shown in **Figure S4**
with data processing and refinement statistics in **Table S4**. Movies were created with
ChimeraX(*43*).
Motivated by the dynamic behaviour of the 5-fold pores of the encapsulin
shell, the 5-fold pore pentamer was subjected to symmetry expansion and focused
classification. The I1 particle set was expanded using the sym_expand job in
Relion3.1 and masked 3D classification without alignment was performed focused on
the five-fold symmetry axis. Of the five classes produced the two most highly
populated and distinct were taken forward for masked 3D refinement with local
searches only, these two classes represent the 'open' and 'closed' conformation of
the five-fold pore. The mask used in these steps was produced in Chimera using the
molmap command on a docked model of a pentamer of the encapsulin protein. To
ensure the box-size and pixel-size of the mask were correct, they were resampled
onto the icosahedral map using the vop resample command.
614
Model building and refinement
An initial homology model of the encapsulin nanocompartment monomer was
generated using Phyre 2.0(*44*) based on the *T. maritima* structure. This was docked
into the open and closed maps using ChimeraX(*45*) and expanded to a full pentamer
model. The model was then fit to the map through an iterative process of automated
model refinement with phenix real-space refinement(*46*) and manual model building
in Coot(*47*), waters were added using phenix.douse and validated in in Coot. The
resulting models and maps were validated using Molprobity(*48*), phenix.mtriage(*49*),
and EM ringer(*50*) (**Table S2**). Models and maps were visualized using ChimeraX.
624
625 **Encapsulin sequence analysis**
626   Encapsulin sequences were obtained from the Kyoto Encyclopedia of Genes
627 and Genomes (www.kegg.jp). Sequence alignments were performed using Clustal
628 Omega(*51*) and visualized using ESPript(*52*).
629
Hydrogen/Deuterium Exchange Mass Spectrometry
Hydrogen/deuterium exchange mass spectrometry (HDX-MS) was performed
on a Synapt G2 MS system coupled to an ACQUITY UPLC M-Class UPLC with the
HDX manager module (Waters Corporation, Manchester, UK)(*53*). For improved
reliability and precision, a custom-built Leap automated platform was utilized in all
sample preparation and injections. Prior to HDX-MS analysis, three buffer solutions
were prepared – (i) equilibration buffer (4.7 mM K_2HPO_4, 0.3 mM KH_2PO_4 in H_2O),
adjusted to pH 8.0 with formic acid; (ii) labelling buffer (4.7 mM K_2HPO_4, 0.3 mM
KH_2PO_4 in D_2O), adjusted to pH 8.0 with DCl and quench buffer (50 mM K_2HPO_4, 50
mM KH_2PO_4 in H_2O) adjusted to pH 2.3 with formic acid. Protein samples were
diluted in equilibration buffer to final stock concentration of 42 mM. The time course
experiments consisted of 7 timepoints: T0 (0 minute; undeuterated control), T1 (20
seconds), T2 (30 seconds), T3 (2 minutes), T4 (5 minutes) and T5 (30 minutes) T6

(4 hours) and T7 (24 hours) with each timepoint being performed in triplicate.
Sample preparation consisted of 5 µL protein solution, 57 µL equilibrium buffer (T0)
or labelling buffer (T1-7). The final concentration of deuterium during the labelling
step was 91.2 %. Exchange was allowed to proceed at 4 °C. To arrest the exchange
reaction, 50 µL of quench buffer was added to this initial solution just prior to sample
injection.
After injection, samples underwent proteolytic digestion on a 2.1 x 30 mm
Waters Enzymate BEH pepsin column for 3 minutes at 200 µL/min. After digestion,
the peptide digest was loaded on to a 2.1 x 5.0 mm Acquity BEH C18 VanGuard
1.7µm C18 Trapping column to pre-concentrate the sample for 3 minutes at
200 µL/min. Following trapping, the digests were separated through a 2.1 x 5.0 mm
Acquity BEH 1.7 µm analytical column prior to MS/MS (MS^e) analysis via the Water
Synapt G2 MS system running MassLynx v4.1 software (Waters Corporation,
Manchester, UK). The separation gradient was 5-95 % acetonitrile with 0.1 % formic
acid over 12 minutes at 40 µL/min. Both the trapping and LC separation were
performed at 1 °C to minimize back exchange. Post-processing was performed using
Proteinlynx Global Server 3.0.3 and Dynamx 3.0 software to determine the average
deuterium uptake for each peptide at each time point. For comparative analyses, the
relative fractional uptake was determined by dividing the observed deuterium uptake
by the number of available amide exchangers on each peptide.
663

References

1. Y. Diekmann, J. B. Pereira-Leal, Evolution of intracellular compartmentalization. *Biochem. J.* **449**, 319–331 (2013).

2. C. A. Kerfeld, S. Heinhorst, G. C. Cannon, Bacterial Microcompartments (2010) (available at http://www.annualreviews.org/eprint/uv9wHHypNKsEb5yrt6z8/full/10.1146/annurev.micro.112408.134211).

3. W. Bonacci, P. K. Teng, B. Afonso, H. Niederholtmeyer, P. Grob, P. A. Silver, D. F. Savage, Modularity of a carbon-fixing protein organelle. *Proc. Natl. Acad. Sci.* **109**, 478–483 (2012).

4. C. V Iancu, D. M. Morris, Z. Dou, S. Heinhorst, C. Gordon, G. J. Jensen, NIH Public Access. **396**, 105–117 (2010).

5. T. O. Yeates, C. A. Kerfeld, S. Heinhorst, G. C. Cannon, J. M. Shively, Protein-based organelles in bacteria: carboxysomes and related microcompartments. *Nat. Rev. Microbiol.* **6**, 681–691 (2008).

6. S. C. Andrews, The Ferritin-like superfamily: Evolution of the biological iron storeman from a rubrerythrin-like ancestor. *Biochim. Biophys. Acta.* **1800**, 691–705 (2010).

7. E. Chiancone, P. Ceci, A. Ilari, F. Ribacchi, S. Stefanini, Iron and proteins for iron storage and detoxification. *BioMetals.* **17**, 197–202 (2004).

8. F. Akita, K. T. Chong, H. Tanaka, E. Yamashita, N. Miyazaki, Y. Nakaishi, M. Suzuki, K. Namba, Y. Ono, T. Tsukihara, A. Nakagawa, The crystal structure of a virus-like particle from the hyperthermophilic archaeon Pyrococcus furiosus provides insight into the evolution of viruses. *J. Mol. Biol.* **368**, 1469–83 (2007).

9. M. Sutter, D. Boehringer, S. Gutmann, S. Günther, D. Prangishvili, M. J. Loessner, K. O. Stetter, E. Weber-Ban, N. Ban, Structural basis of enzyme encapsulation into a bacterial nanocompartment. *Nat. Struct. Mol. Biol.* **15**, 939–947 (2008).

10. T. W. Giessen, P. A. Silver, Widespread distribution of encapsulin nanocompartments reveals functional diversity. *Nat. Microbiol.* **2**, 17029 (2017).

11. D. He, S. Hughes, S. Vanden-Hehir, A. Georgiev, K. Altenbach, E. Tarrant, C. L. Mackay, K. J. Waldron, D. J. Clarke, J. Marles-Wright, Structural characterization of encapsulated ferritin provides insight into iron storage in bacterial nanocompartments. *Elife.* **5**, e18972 (2016).

12. T. W. Giessen, B. J. Orlando, A. A. Verdegaal, M. G. Chambers, J. Gardener, D. C. Bell, G. Birrane, M. Liao, P. A. Silver, Large protein organelles form a new iron sequestration system with high storage capacity. *Elife.* **8**, 1–23 (2019).

13. C. A. McHugh, J. Fontana, D. Nemecek, N. Cheng, A. A. Aksyuk, J. B. Heymann, D. C. Winkler, A. S. Lam, J. S. Wall, A. C. Steven, E. Hoiczyk, A virus capsid-like nanocompartment that stores iron and protects bacteria from oxidative stress. *EMBO J.* **33**, 1896–1911 (2014).

14. T. W. Giessen, P. A. Silver, Widespread distribution of encapsulin nanocompartments reveals functional diversity. *Nat. Microbiol.* **2**, 17029 (2017).

15. A. Tamura, Y. Fukutani, T. Takami, M. Fujii, Y. Nakaguchi, Y. Murakami, K. Noguchi, M. Yohda, M. Odaka, Packaging guest proteins into the encapsulin nanocompartment from Rhodococcus erythropolis N771. *Biotechnol. Bioeng.*

112, 13–20 (2015).

16.   D. He, C. Piergentili, J. Ross, E. Tarrant, L. R. Tuck, C. L. Mackay, Z. McIver,
K. J. Waldron, D. J. Clarke, J. Marles-Wright, Conservation of the structural
and functional architecture of encapsulated ferritins in bacteria and archaea.
Biochem. J. **476**, 975–989 (2019).

17.   R. Rahmanpour, T. D. H. Bugg, Assembly in vitro of Rhodococcus jostii RHA1
encapsulin and peroxidase DypB to form a nanocompartment. *FEBS J.* **280**,
2097–2104 (2013).

18.   J. Snijder, O. Kononova, I. M. Barbu, C. Uetrecht, W. F. Rurup, R. J. Burnley,
M. S. T. Koay, J. J. L. M. Cornelissen, W. H. Roos, V. Barsegov, G. J. L.
Wuite, A. J. R. Heck, Assembly and Mechanical Properties of the Cargo-Free
and Cargo-Loaded Bacterial Nanocompartment Encapsulin.
Biomacromolecules. **17**, 2522–9 (2016).

19.   M. P. Andreas, T. W. Giessen, Large-scale computational discovery and
analysis of virus-derived microbial nanocompartments. *Nat. Commun.* **12**,
4748 (2021).

20.   M. C. Jenkins, S. Lutz, Encapsulin Nanocontainers as Versatile Scaffolds for
the Development of Artificial Metabolons. *ACS Synth. Biol.* **10**, 857–869
(2021).

21.   Y. H. Lau, T. W. Giessen, W. J. Altenburg, P. A. Silver, Prokaryotic
nanocompartments form synthetic organelles in a eukaryote. *Nat. Commun.* **9**,
1311 (2018).

22.   F. Sigmund, S. Pettinger, M. Kube, F. Schneider, M. Schifferer, S. Schneider,
M. V. Efremova, J. Pujol-Martí, M. Aichler, A. Walch, T. Misgeld, H. Dietz, G.
G. Westmeyer, Iron-Sequestering Nanocompartments as Multiplexed Electron
Microscopy Gene Reporters. *ACS Nano.* **13**, 8114–8123 (2019).

23.   S. Recalcati, E. Gammella, P. Buratti, G. Cairo, Molecular regulation of cellular
iron balance. *IUBMB Life.* **69**, 389–398 (2017).

24.   A. Tamura, Y. Fukutani, T. Takami, M. Fujii, Y. Nakaguchi, Y. Murakami, K.
Noguchi, M. Yohda, M. Odaka, Packaging guest proteins into the encapsulin
nanocompartment from Rhodococcus erythropolis N771. *Biotechnol. Bioeng.*
112, 13–20 (2015).

25.   Y. Tang, A. Mu, Y. Zhang, S. Zhou, W. Wang, Y. Lai, X. Zhou, F. Liu, X. Yang,
H. Gong, Q. Wang, Z. Rao, Cryo-EM structure of Mycobacterium smegmatis
DyP-loaded encapsulin. *Proc. Natl. Acad. Sci.* **118**, e2025658118 (2021).

26.   M. Sutter, D. Boehringer, S. Gutmann, S. Günther, D. Prangishvili, M. J.
Loessner, K. O. Stetter, E. Weber-Ban, N. Ban, Structural basis of enzyme
encapsulation into a bacterial nanocompartment. *Nat. Struct. Mol. Biol.* **15**,
939–947 (2008).

27.   X. Xiong, C. Sun, F. S. Vago, T. Klose, J. Zhu, W. Jiang, Cryo-EM structure of
heterologous protein complex loaded thermotoga maritima encapsulin capsid.
Biomolecules. **10**, 1–13 (2020).

28.   R. M. Putri, C. Allende-Ballestero, D. Luque, R. Klem, K.-A. Rousou, A. Liu, C.
H.-H. Traulsen, W. F. Rurup, M. S. T. Koay, J. R. Castón, J. J. L. M.
Cornelissen, Structural Characterization of Native and Modified Encapsulins as
Nanoplatforms for *in Vitro* Catalysis and Cellular Uptake. *ACS Nano.* **11**,
12796–12804 (2017).

29.   J. Ross, T. Lambert, C. Piergentili, D. He, K. J. Waldron, C. L. Mackay, J.
Marles-Wright, D. J. Clarke, Mass spectrometry reveals the assembly pathway
of encapsulated ferritins and highlights a dynamic ferroxidase interface. *Chem.*

Commun. (Camb). **56**, 3417–3420 (2020).

30.   T. Nakane, E. Lindahl, J. Zivanov, W. J. J. H. Hagen, S. H. W. H. Scheres, D.
766   Kimanius, B. B. O. Forsberg, T. Nakane, B. B. O. Forsberg, D. Kimanius, W. J.
767   J. H. Hagen, E. Lindahl, S. H. W. H. Scheres, New tools for automated high-
768   resolution cryo-EM structure determination in RELION-3. *Elife.* **7**, 1–38 (2018).

769   31.   R. J. Nichols, B. LaFrance, N. R. Phillips, D. R. Radford, L. M. Oltrogge, L. E.
770   Valentin-Alvarado, A. J. Bischoff, E. Nogales, D. F. Savage, Discovery and
771   characterization of a novel family of prokaryotic nanocompartments involved in
772   sulfur metabolism. *Elife.* **10** (2021), doi:10.7554/eLife.59288.

773   32.   B. LaFrance, C. Cassidy-Amstutz, R. J. Nichols, L. M. Oltrogge, E. Nogales, D.
774   F. Savage, *bioRxiv*, in press, doi:10.1101/2021.04.26.441214.

775   33.   E. M. Williams, S. M. Jung, J. L. Coffman, S. Lutz, Pore Engineering for
776   Enhanced Mass Transport in Encapsulin Nanocompartments. *ACS Synth. Biol.*
777   **7**, 2514–2517 (2018).

778   34.   A. Punjani, D. J. Fleet, 3D variability analysis: Resolving continuous flexibility
779   and discrete heterogeneity from single particle cryo-EM. *J. Struct. Biol.* **213**,
780   107702 (2021).

781   35.   L. Adamson, N. Tasneem, M. P. Andreas, W. Close, T. N. Szyszka, E. Jenner,
782   R. Young, L. Chen Cheah, A. Norman, F. Sainsbury, T. W. Giessen, Y. Heng
783   Lau, *bioRxiv*, in press, doi:10.1101/2021.01.27.428512.

784   36.   C. Piergentili, J. Ross, D. He, K. J. Gallagher, W. A. Stanley, L. Adam, C. L.
785   Mackay, A. Baslé, K. J. Waldron, D. J. Clarke, J. Marles-Wright, Dissecting the
786   structural and functional roles of a putative metal entry site in encapsulated
787   ferritins. *J. Biol. Chem.* (2020), doi:10.1074/jbc.RA120.014502.

788   37.   S. V. Iverson, T. L. Haddock, J. Beal, D. M. Densmore, CIDAR MoClo:
789   Improved MoClo Assembly Standard and New E. coli Part Library Enables
790   Rapid Combinatorial Design for Synthetic and Traditional Biology. *ACS Synth.*
791   *Biol.* **5**, 151019092657002 (2015).

792   38.   F. Studier, Protein production by auto-induction in high-density shaking
793   cultures. *Protein Expr. Purif.* **41**, 207–234 (2005).

794   39.   J. Schindelin, I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T.
795   Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D. J.
796   White, V. Hartenstein, K. Eliceiri, P. Tomancak, A. Cardona, Fiji: an open-
797   source platform for biological-image analysis. *Nat. Methods.* **9**, 676–682
798   (2012).

799   40.   S. Q. Zheng, E. Palovcak, J.-P. Armache, K. A. Verba, Y. Cheng, D. A. Agard,
800   MotionCor2: anisotropic correction of beam-induced motion for improved cryo-
801   electron microscopy. *Nat. Methods.* **14**, 331–332 (2017).

802   41.   A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation
803   from electron micrographs. *J. Struct. Biol.* **192**, 216–21 (2015).

804   42.   A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, No Title. *Nat.*
805   *Methods.* **14**, 290–296 (2017).

806   43.   T. D. Goddard, C. C. Huang, E. C. Meng, E. F. Pettersen, G. S. Couch, J. H.
807   Morris, T. E. Ferrin, UCSF ChimeraX: Meeting modern challenges in
808   visualization and analysis. *Protein Sci.* **27**, 14–25 (2018).

809   44.   L. A. Kelly, S. Mezulis, C. Yates, M. Wass, M. Sternberg, The Phyre2 web
810   portal for protein modelling, prediction, and analysis. *Nat. Protoc.* **10**, 845–858
811   (2015).

812   45.   E. F. Pettersen, T. D. Goddard, C. C. Huang, E. C. Meng, G. S. Couch, T. I.
813   Croll, J. H. Morris, T. E. Ferrin, UCSF ChimeraX: Structure visualization for

researchers, educators, and developers. *Protein Sci.* **30**, 70–82 (2021).

46.   P. V. Afonine, B. K. Poon, R. J. Read, O. V. Sobolev, T. C. Terwilliger, A.
Urzhumtsev, P. D. Adams, Real-space refinement in PHENIX for cryo-EM and
crystallography. *Acta Crystallogr. Sect. D Struct. Biol.* **74**, 531–544 (2018).

47.   P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of
Coot. *Acta Crystallogr. D. Biol. Crystallogr.* **66**, 486–501 (2010).

48.   V. B. Chen, W. B. Arendall, J. J. Headd, D. A. Keedy, R. M. Immormino, G. J.
Kapral, L. W. Murray, J. S. Richardson, D. C. Richardson, MolProbity: all-atom
structure validation for macromolecular crystallography. *Acta Crystallogr. D.*
Biol. Crystallogr. **66**, 12–21 (2010).

49.   P. V. Afonine, B. P. Klaholz, N. W. Moriarty, B. K. Poon, O. V. Sobolev, T. C.
Terwilliger, P. D. Adams, A. Urzhumtsev, IUCr, New tools for the analysis and
validation of cryo-EM maps and atomic models. *Acta Crystallogr. Sect. D*
Struct. Biol. **74**, 814–840 (2018).

50.   B. A. Barad, N. Echols, R. Y.-R. Wang, Y. Cheng, F. DiMaio, P. D. Adams, J.
S. Fraser, EMRinger: side chain-directed model and map validation for 3D
cryo-electron microscopy. *Nat. Methods.* **12**, 943–6 (2015).

51.   F. Sievers, D. G. Higgins, Clustal Omega, accurate alignment of very large
numbers of sequences. *Methods Mol. Biol.* **1079**, 105–16 (2014).

52.   P. Gouet, X. Robert, E. Courcelle, ESPript/ENDscript: Extracting and rendering
sequence and 3D information from atomic structures of proteins. *Nucleic Acids*
Res. **31**, 3320–3 (2003).

53.   G. R. Masson, J. E. Burke, N. G. Ahn, G. S. Anand, C. Borchers, S. Brier, G.
M. Bou-Assaf, J. R. Engen, S. W. Englander, J. Faber, R. Garlish, P. R. Griffin,
M. L. Gross, M. Guttman, Y. Hamuro, A. J. R. Heck, D. Houde, R. E. Iacob, T.
J. D. Jørgensen, I. A. Kaltashov, J. P. Klinman, L. Konermann, P. Man, L.
Mayne, B. D. Pascal, D. Reichmann, M. Skehel, J. Snijder, T. S. Strutzenberg,
E. S. Underbakke, C. Wagner, T. E. Wales, B. T. Walters, D. D. Weis, D. J.
Wilson, P. L. Wintrode, Z. Zhang, J. Zheng, D. C. Schriemer, K. D. Rand,
Recommendations for performing, interpreting and reporting hydrogen
deuterium exchange mass spectrometry (HDX-MS) experiments. *Nat.*
Methods. **16**, 595–602 (2019).

54.   M. Laguerre, M. Saux, J. P. Dubost, A. Carpy, in *Pharmaceutical Sciences*
(John Wiley & Sons, Ltd, 1997), vol. 3, pp. 217–222.

Funding

This work was supported a Royal Society Research Grant awarded to JMW [RG130585] and a BBSRC New Investigator Grant to JMW and DJC [BB/N005570/1]. CP was funded by the BBSRC New Investigator Grant [BB/N005570/1]. JMW is funded by Newcastle University. DJC, TL and JR are funded by the University of Edinburgh. JR was funded by a BBSRC EastBio DTP studentship [BB/M010996/1]. ZM is funded by a BBSRC NLD DTP studentship [BB/M011186/1]. JEB was funded by DSTL via Imperial College. EZA was funded by an IBioIC PhD studentship with Fujifilm Diosynth Biotechnologies. KJW was funded by the Biotechnology and Biological Sciences Research Council (BB/S006818/1)

Equipment for Transmission Electron Microscopy at Newcastle University was funded through the BBSRC 17ALERT call [BB/R013942/1]. Equipment for Mass spectrometry at the University of Edinburgh was funded through the BBSRC 17ALERT call [BB/R013993/1]. MDW's work is supported by the Wellcome Trust and Royal Society (210493), Medical Research Council (T029471/1), and the University of Edinburgh. The Wellcome Centre for Cell Biology is supported by core funding from the Wellcome Trust (203149).

Acknowledgements

We acknowledge Diamond Light Source for access and support of the cryo-EM facilities at the UK's national Electron Bio-imaging Centre (eBIC) [under proposal EM16637], funded by the Wellcome Trust, MRC and BBRSC. We thank Dr Yun Song for their assistance at eBIC with sample loading and data collection setup. This research made use of the Rocket High Performance Computing service at Newcastle University; we thank Dr Karen Bower for assistance with the use of the Rocket service. Negative stain TEM was performed in Edinburgh Electron Microscopy Lab and the Newcastle University Electron Microscopy Research Services, we thank Steve Mitchell, Tracey Davey, and Ross Laws for their technical support. Cryo-EM grid screening was performed in the Edinburgh cryo-EM facility in School of Biological Sciences at the University of Edinburgh. The cryo-EM facility was set up with funding from the Wellcome Trust (087658/Z/08/Z) and SULSA and is supported by the Wellcome Centre for Cell Biology. We thank Professor Martin Noble for generously donating GPU time to our CryoSPARC analyses.

Data Availability

Please see Experimental Procedures and tables for links to these. All data needed to evaluate the conclusions in the paper are present in the Supplementary Materials or have been deposited in appropriate publicly available data repositories. Please see the Materials and Methods sections and experimental tables for links to these datasets.

Competing Interests

All authors declare that they have no competing interests.

Author Contributions

Conceptualization: JR, JMW, DJC
Methodology: JR, ZM, TL, CP, AMBL, JMW, DJC
Validation: JR, ZM, TL, JMW, DJC
Formal analysis: JR, ZM, TL, AMBL, PJ, JMW, DJC
Investigation: JR, ZM, KG, JEB, CP, EZA

Resources: KG, FC, CLM, JEB, AMBL, EZA, MDW
Data curation: JR, TL, DJC, AMBL, JMW
Writing – original draft preparation: JR, DJC, JMW
Writing – review and editing: JR, ZM, CP, AMBL, MDW, KJW, JMW, DJC
Visualization: JR, ZM, TL, DJC, JMW
Supervision: JMW, DJC
Funding acquisition: MDW, LEH, KJW, DJC, JMW
903

Figures

905
Figure 1. Validation of the Assembly and Activity of Loaded-Enc and Empty-
Enc nanocompartments.
A: SDS-PAGE of purified Empty-Enc and Loaded-Enc. Proteins resolved by 15%
acrylamide SDS-PAGE and stained with Coomassie blue stain. Encapsulin bands
are near the 30 kDa marker and highlighted by a black arrow. The EncFtn cargo of
Loaded-Enc appears as both a monomer and dimer. Bands corresponding to the
monomer and dimer of EncFtn are highlighted with green arrows. **B**: Recombinant,
purified Empty-Enc (blue trace) and Loaded-Enc (orange, dotted trace) were
analyzed by size exclusion chromatography using a Sephacryl 400 column (Cytiva).
Both encapsulins elute at the same volume, suggesting the encapsulin complexes
are the same size regardless of cargo loading. **C**: Negative stain transmission
micrographs of Empty-Enc (upper micrograph) and Loaded-Enc (lower micrograph)
displaying individual particles for each complex. One nanocompartment of Empty-

Enc and Loaded-Enc is shown in the upper right corner of each micrograph, with a
hexagonal 2D geometry observed. **D**: upper panel: Ferroxidase activity of Loaded-
Enc compared to Empty-Enc. Protein samples were mixed with 100 μM
$FeSO_4.7H_2O$. Following an incubation period at room temperature of 50 seconds
absorbance at 315 nm was measured over a time-course of 1450 seconds. Control
reference established using enzyme-free reaction as a measure of background iron
oxidation. The lines represent the mean of all repeats, error bars represent the
standard deviation from the mean. Lower panel: End point ferroxidase assay
comparison. Ferroxidase activity shown by the total increase in $A_{315\ nm}$ at the end
point of the assay. Bars represent the mean of all repeats, error bars represent the
standard deviation from the mean.
930

931
932
Figure 2: Architecture of the *H. ochraceum* encapsulin nanocompartment
shell.
Visualization of the electronic potential map of the *H. ochraceum* encapsulin from an
icosahedrally averaged single particle reconstruction. **A**: The exterior of the
encapsulin shell visualized at 2.4 Å resolution. Five subunits of the encapsulin
nanocompartment shell have been colored to highlight the 5-fold axis. **B**: Icosahedral
EM map of Loaded-Enc sharpened by local resolution estimate and colored by local
resolution. The estimated resolution varies across the exterior of the encapsulin
nanocompartment with the lowest resolution at the 5-fold pores. Color key of
resolution mapping is shown on the right-hand side of the figure. **C**: The shared
phage-like fold in the HK97 bacteriophage capsid and encapsulin proteins.

Monomeric subunit of the *H. ochraceum* encapsulin protein modelled from our
reconstruction are shown (pink), with comparisons to other *T=1* encapsulins from:
T. maritima (blue, PDB ID:3DKT); *M. smegmatis* (lilac, PDB ID: 7BOJ); and
*S. elongatu*s (grey, PDB ID: 6X8M). The HK97 bacteriophage Head II *T=7* monomer
(green, PDB ID: 2FT1) is also shown.. The N-terminus of each monomer is
highlighted in blue and the C-termius in red. **Lower right:** Overlay comparison of the
encapsulin monomers showing similar A- and P-domains orientations.

951
952
Figure 3: Asymmetric reconstruction of the *H. ochraceum* encapsulin complex
reveals a tetrahedral arrangement of EncFtn within the encapsulin
nanocompartment.
A: Electronic potential map of the asymmetric reconstruction of the *H. ochraceum*
Enc:EncFtn complex. The map is colored by local resolution with the color key
shown on the right side of **A**. **B:** Radially colored cryo-EM derived map of the
Loaded-Enc nanocompartment displaying the interior EncFtn (cyan). **C**: Gaussian
smoothed C1 map showing the four discrete EncFtn densities (red, green, yellow,
and blue), consistent with the size of decameric EncFtn complexes, within the
encapsulin nanocompartment (pink). The four EncFtn are in a tetrahedral

arrangement highlighted by grey lines connecting their centers of mass. The average
distance between each EncFtn decamer is 57 Å. **D**: EM Map of a single EncFtn
density following symmetry expansion and local refinement in CryoSPARC. The *H.*
ochraceum EncFtn decamer crystal structure (PDB ID: 5N5F, red) is docked into the
density.

968
969
Figure 4: 3D variability analysis of Loaded-Enc.
The first and last frames from the trajectory of component 0 from the 3DVA are shown as density maps centered on a single five-fold pore. Pores across the encapsulin nanocompartment are seen to adopt conformations consistent with both 'closed' (green) 'open' states (purple) and move between these along the trajectory. Different pores adopt this dynamic in the two other components calculated.

976
Figure 5: Open and closed conformations of the dynamic 5-fold pore of the *H. ochraceum* encapsulin shell.
Masked 3D refinements centered around the 5-fold pore of the icosahedral reconstruction were performed after symmetry
expansion of the asymmetric units.
The 'closed' (**A**, green) and 'open' (**B**, purple) conformations of the Loaded-Enc shell pentamer are represented with transparent
solvent accessible surface rendered over cartoon secondary structure. Both 'closed' and 'open' pentamers are shown in top-down
(left-hand side) and side-on views (right-hand side). The closed conformation has a 5-fold pore diameter of 5 Å, whilst the
equivalent diameter of the open conformation is 15 Å. To allow for the widening of the pore, the open pentamer protrudes from the
nanocompartment by 5 Å.
985
986 .
987

988
989
Figure 6: The closed conformation of the *H. ochraceum* five-fold pore allows docking and sequencing of the EncFtn
localization sequence to the interior face of the encapsulin monomer.
A: Binding of localization sequence residues from the EncFtn to the interior wall of the encapsulin nanocompartment monomer
(green residues and transparent cartoon) and the localization sequence (yellow sticks). Lower panel: Hydrophobic residues from
the interior face of the encapsulin form the binding pocket for the localization sequence. The first and last residues modelled for the
LS have been labelled in black, and key residues from encapsulin have been labelled in green. Modelled water molecules are
shown as red spheres. **B:** The spatial relationship between an encapsulin monomer and the localization sequence (shown in gold
with its EM map density as in **A**). The Enc monomer has been colored by molecular lipophilicity potential (*54*) which ranges from
dark cyan (corresponding to the most hydrophilic) to white to dark gold (most lipophilic). This highlights the hydrophobic pocket on
the interior of the encapsulin nanocompartment where the localization sequence binds. **C**: The 'closed' conformation pentamer
(green) with a localization sequence (yellow) shown on each monomer.
1001

Figure 7: HDX-MS fractional uptake mapped to the symmetry axes of the icosahedral reconstruction of the *H. ochraceum* encapsulin complex.

From top to bottom: The top row depicts molecular surface models of the *H. ochraceum* encapsulin in three orientations to show the 5-fold, 3-fold and 2-fold symmetry axes. Monomers have been colored individually (pink, yellow, green, blue, and purple) to highlight the symmetry axes. The final two rows display the relative fractional uptake of each peptide from HDX experiments displayed on the 5-fold, 3-fold and 2-fold pores of the encapsulin nanocompartment. Peptide fractional uptake is show on a white-to-orange color scale with a color key shown on the right-hand side of the figure. Areas colored black correspond to no peptide coverage.