Deep mutational scanning of the plasminogen activator inhibitor-1 functional landscape

Zachary M. Huttinger[1,2,#,**], Laura M. Haynes[1,**], Andrew Yee[3], Colin A. Kretz[4], David R. Siemieniak[1,5], Daniel A. Lawrence[6,7], and David Ginsburg[1,2,5,6,8,†]

[1]Life Sciences Institute, University of Michigan, Ann Arbor, MI

[2] Cellular and Molecular Biology Program, University of Michigan Medical School, Ann Arbor, MI

[3]Department of Pediatrics, Baylor College of Medicine, Houston, TX

[4]Department of Medicine, McMaster University and the Thrombosis and Atherosclerosis Research Institute, Hamilton, ON, Canada

[5]Howard Hughes Medical Institute

[6]Department of Internal Medicine, University of Michigan Medical School, Ann Arbor, MI

[7]Department of of Molecular & Integrative Physiology, University of Michigan Medical School, Ann Arbor, MI

[8]Departments of Human Genetics and Pediatrics, University of Michigan, Ann Arbor, MI

[#]Current Address: Department of Otolaryngology, The Ohio State University College of Medicine, Columbus, OH

[**]These two authors contributed equally to this work.

[†]Corresponding author: David Ginsburg

E-mail: ginsburg@umich.edu

Running Title: PAI-1 functional landscape

Keywords: high-throughput sequencing (HTS), phage display, serpin, enzyme inhibitor, protease inhibitor, fibrinolysis

ABSTRACT

The serine protease inhibitor (SERPIN) plasminogen activator inhibitor-1 (PAI-1) is a key regulator of the fibrinolytic system, inhibiting the serine proteases tissue- and urokinase-type plasminogen activator (tPA and uPA, respectively). Missense variants may render PAI-1 non-functional through misfolding, leading to its turnover as a protease substrate, or to a more rapid transition to the latent/inactive state. Deep mutational scanning was performed to evaluate the impact of amino acid sequence variation on PAI-1 inhibition of uPA using an M13 filamentous phage display system. The effects of single amino acid substitutions on PAI-1's functional inhibition of its canonical target proteases, tPA and uPA, have been determined for only a small fraction of potential mutations. To construct a more comprehensive dataset, a mutagenized PAI-1 library, encompassing ~70% of potential single amino acid substitutions, was displayed on M13 filamentous phage. From this library, the relative effects of 27% of all possible missense variants on PAI-1 inhibition of urokinase-type plasminogen activator were determined using high-throughput DNA sequencing with 826 missense variants demonstrating conserved inhibitory activity and 1137 resulting in loss of PAI-1 function. Comparison of these deep mutational scanning results to predictions from PolyPhen-2 and SIFT demonstrate the limitations of these algorithms, consistent with similar reports for other proteins. Comparison to common human PAI-1 gene variants present in the gnomAD database is consistent with evolutionary selection against loss of PAI-1 function. These findings provide insight into structure-function relationships for PAI-1 and other members of the SERPIN superfamily.

INTRODUCTION

Plasminogen activator inhibitor-1 (PAI-1, *SERPINE1*) is a member of the serine protease inhibitor (SERPIN) protein superfamily (1). SERPINSs function as irreversible inhibitors that covalently bind the active site of their target proteases (2). All inhibitory SERPINs share a common mechanism wherein a flexible reactive center loop (RCL) extends outside the central structure of the molecule serving as a cleavable peptide "bait" for the target protease (Fig. 1A) (3). The SERPIN and protease form an acyl intermediate when the active site serine acts as a donor nucleophile to bind to the carbonyl carbon of the P1 residue within the RCL(4, 5). However, the RCL rapidly inserts into central β-sheet A before the hydrolysis reaction is completed, carrying the tethered protease to the opposite side of the molecule in a pole-to-pole transition (6). This structural transformation stabilizes the protease-SERPIN complex and renders both the SERPIN and the serine protease no longer active (2, 7, 8). If the rate of RCL insertion is slower than the resolution of the acyl intermediate, complete proteolysis of the scissile bond occurs, with the SERPIN then functioning as a substrate instead of an inhibitor (9). Either of these mechanisms is possible when a SERPIN interacts with a given protease. The specificity of a SERPIN for its target protease is in part driven by the probability that the inhibitory pathway dominates over the substrate pathway (10). This balance presents a unique challenge when engineering SERPIN variants that inhibit non-canonical target proteases (11, 12).

PAI-1 is the principal inhibitor of the serine proteases tissue-type and urokinase-type plasminogen activators (tPA and uPA, respectively)(13, 14). tPA and uPA activate the zymogen plasminogen to plasmin (15), the enzyme responsible for the proteolysis of fibrin—the structural backbone of blood clots (16). Human deficiency of PAI-1 results in excessive plasmin generation

and a mild to moderate bleeding diathesis (17-21). Likewise, PAI-1 deficient mice are viable with a mild increase in fibrinolytic activity (22). In addition to its critical role in regulating hemostasis, PAI-1has been implicated in a number of other processes, including a link to longevity (23). PAI-1 lacks native cysteine residues and remains functional in the absence of glycosylation, facilitating large-scale production of functionally active, recombinant PAI-1 in bacterial systems (24, 25).

The present work couples phage display with high-throughput DNA sequencing (HTS), to measure the effects of multiple missense mutations on PAI-1's uPA inhibitory function in a massively parallel fashion.

MATERIALS AND METHODS

Materials

Restriction enzymes, protein G magnetic beads, enterokinase, and NEBNext DNA Library Prep Master Mix Kit were purchased from New England Biolabs (Ipswitch, MA). The GeneMorph II Random Mutagenesis Kit, QuickChange II XL Site-Directed Mutagenesis Kit, and chemically and electrocompetent XL-1 Blue MRF' *E. coli* were obtained from Agilent Technologies (Santa Clara, CA). LB broth and anti-c-myc agarose gel were purchased from Invitrogen (Waltham, MA). M13KO7 helper phage were obtained from GE Healthcare (Chicago, IL). Ampicillin, kanamycin, 3X FLAG peptide, and anti-FLAG agarose beads were purchased from Sigma-Aldrich (Saint Louis, MO). High molecular weight (HMW)-uPA was obtained from Molecular Innovations (Novi, MI). EDTA-free protease inhibitor cocktail and isopropyl-β-D-thiogalactoside (IPTG) were purchased from Roche (Basel, Switzerland). A polyclonal rabbit-anti-uPA antibody (ab2412) was purchased from Abcam (Cambridge, UK). Z-Gly-Gly-Arg-

AMC is a product of Bachem (Bubendorf, Switzerland). NEXTflex multiplex adapters were purchased from Bioo Scientific (Austin, TX).

Construction of a phage display library expressing PAI-1 fusion proteins

For display of PAI-1 on the M13 filamentous bacteriophage (phPAI-1, Fig. 1B), human SERPINE1 cDNA including a N-terminal myc tag and Gly-Gly-Gly-Ser linker was cloned between the AscI and NotI restriction sites of pAY-FE (Genbank #MW464120) (27, 28). The resulting construct encodes a phage-displayed PAI-1 protein N-terminally fused to a myc tag and C-terminally fused with FLAG and E tags (Fig. 1). The PAI-1 fusion protein was randomly mutagenized using the GeneMorph II Random Mutagenesis Kit. Primers used for PCR mutagenesis (SI Table 1) maintained the AscI and NotI restriction sites for ligation of the restriction digested insert into pAY-FE. Following ligation, the library was transformed into electrocompetent XL-1 Blue MRF' *E. coli* as per manufacturer's instructions. The depth of the library was determined by quantifying the number of ampicillin resistant colonies. Mutation frequency was estimated by Sanger sequencing of the SERPINE1 inserts from randomly selected individual colonies (n=13).

Phage production and purification

Phage were prepared as previously reported (26) with modifications to minimize PAI-1 conversion to the latent conformation. Briefly, *E. coli* harboring pAY-FE PAI-1 were grown in LB Broth supplemented with 2% glucose and ampicillin (100 μg/mL) at 37°C and during mid-log phase (OD_{600} 0.3-0.4) were infected with M13KO7 helper phage at a multiplicity of infection of ~100, followed by growth for an additional 1 h at 37°C. Cells were pelleted by centrifugation

(4250xg for 10 m at 4°C), resuspended in 2xYT media (16 g/L tryptone, 10 g/L yeast extract, 5

g/L NaCl) supplemented with ampicillin (100 µg/mL), kanamycin (30 µg/mL) and IPTG (0.4

mM) to induce expression of the PAI-1 fusion protein, and grown for 2 h at 37°C (27). All

subsequent phage preparation steps were carried out at 4°C. Phage were precipitated with

polyethylene glycol-8000 (2.5% w/v) and NaCl (0.5 M) for up to 16 h (with confirmation of no

change in the number of active phPAI-1 present after purification) followed by centrifugation

(20,000xg for 20 min at 4°C). The precipitated phage pellet was resuspended in 50 mM Tris

containing 150 mM NaCl (pH 7.4; TBS). Phage titer was determined by transducing naïve XL-1

Blue MRF' *E. coli* grown to mid-log phase for 1 hour at 37°C and plating on LB-agar

supplemented with ampicillin (100 µg/mL) and 2% glucose.

Selection of uPA-bound phPAI-1

Phage displaying the A3 domain of von Willebrand Factor (phVWF-A3), in which the VWF A3

domain (Ser1681-Cys1872) was PCR amplified (SI Table 1) and cloned into the pAY-FE vector

between the AscI and NotI restriction sites (generating pAY-FE-VWF-A3), were used as a

negative control for uPA binding. phPAI-1 were diluted 9:1 with phVWF-A3 and then incubated

with uPA (1.7 nM) for 30 minutes at 37°C. Residual protease activity was inhibited by

incubating the reaction mixture with 1X EDTA-free protease inhibitor cocktail for 10 min at

37°C. uPA (free and complexed) was immunoprecipitated using magnetic protein G beads (15

µL), which were previously coupled to a polyclonal anti-uPA antibody (17 nM). Beads were

washed four times with TBS containing 5% BSA (1 mL), resuspended in Tris (20 mM) pH 8.0

containing 50 mM NaCl, 2 mM CaCl$_2$, and 5% BSA, and eluted by digestion with enterokinase

(16 U) for 16 hours at 4°C. The eluted phage pool was used to infect naive XL-1 Blue MRF' *E.*

coli. Eluted phage titers were quantified by transduction of XL-1 Blue MRF' cells as described

above. To determine the composition of the phage pools before and after selection, single

colonies of ampicillin resistant bacteria were selected, and their DNA amplified by PCR using

primers annealing outside the insertion site, to a region common to both pAY-FE:PAI-1 and

pAY-FE:VWF-A3 (SI Table 1) with three replicates of n = 24 colonies in each.

High-throughput sequencing (HTS)

Twelve overlapping amplicons (150 bp) were PCR amplified from pAY-FE PAI-1 (SI Table 1)

with overlapping regions only analyzed on one amplicon (SI Table 2). PCR amplicon products

were gel purified, pooled (100 ng DNA divided equally between 12 amplicons), dA-tailed

(NEBNext Ultra End Repair/dA-tail), and ligated to NextFlex barcodes with NEBNext Ultra

Ligation. Ligated Products were purified with AmPure beads according to the manufacturer's

directions. HTS was performed as previously described (25) using the Illumina HiSeq2500 or

HiSeqX platforms (Illumina, San Diego, CA) at the University of Michigan DNA Sequencing

Core (Ann Arbor, MI) or MedGenome, Inc (Foster City, CA). HTS data were analyzed using

DESeq2 (29) analyzing mutations at each position independent of other mutations within a given

amplicon.

Comparison of variant selection results to publicly available datasets

Results of the DESeq2 analysis of input versus selected were compared to the output from the

Sorting Intolerant from Tolerant (SIFT) algorithm that predicts the effect of an amino acid

substitution on protein function by multiple sequence alignments of related proteins (PAI-1 from

S. scrofa, B. taurus, M. vison, R. norvegicus, and *M. musculus;* glia-derived nexin from *H.*

sapiens, M. musculus, and R. norvegicus; neuroserpin from *H. sapiens, G. gallus, R. norvegicus, and M. musculus*) (30). To compare SIFT results to our data, tolerated mutations were defined as those that were able to inhibit uPA (log$_2$-fold change > 0) at 0 h, and noninhibitors (log$_2$-fold < 0) at 0 h were defined as not tolerated. Finally, a χ^2 test (SI Table 3) was used to determine if the variants identified as loss-of-function by our high-throughput screen were significantly underrepresented in the gnomAD database (31) by comparing the expected frequency of variants identified in our screen that were present in gnomAD versus those that were not present.

RESULTS AND DISCUSSION

Characterization of PAI-1 fusion protein

Selection of a 9:1 mixture of phage displayed PAI-1 (phPAI-1) and VWF A3 domain (phVWF-A3) with uPA (Fig. 2) resulted in at least a five-fold enrichment in phPAI-1 relative to phVWF-A3, indicating that the immunoprecipitation is specific for uPA and uPA:PAI-1 complexes. Given the rigorous washing of the immunoprecipitated complex, this enhancement further suggests that PAI-1 expressed as a pIII fusion protein on the phage surface retains its inhibitory activity.

Characterization of the mutant library

The phPAI-1 mutant library exhibits a depth of 8.04×10^6 independent clones with an average of 4.2 ± 1.8 amino acid substitutions/molecule as determined by Sanger sequencing of 13 randomly selected phage clones. HTS demonstrated that 5,117 of the possible 7,201 missense variants (71%) are present in the mutant phPAI-1 library, along with 269 of the 379 possible nonsense mutations (71%). The frequency of DNA sequencing reads for individual amino acid

substitutions within the starting library ranged over $>10^4$-fold. Additionally, biased amino acid

subsitutions were also observed with reduced representation of Met and Trp substitutions (both

encoded by only a single codon), compared to Arg, Leu, and Ser substitutions (each encoded by

six codons) (Fig. 3).

To limit the proportion of phPAI-1 variants transitioning to the non-reactive latent

conformation, all reactions with uPA were performed immediately following phage production.

To limit false positives within the dataset, only those variants with a basemean score (the average

of the counts in the input and selected libraries) greater than 10 and an adjusted p-value (p_{adj}) <

0.05 were included in further analyses (Fig. 4) (29). Based on these criteria, 1,963 (38%) of the

5,117 missense variants present in the starting library could be scored for uPA reactivity or lack

thereof.

Massively parallel assessment of variant impact on uPA inhibitory function

Following selection with uPA, 826 PAI-1 missense variants retained the ability to form a

complex with uPA, with a range of enrichment scores likely representing varying degrees of

inhibitory activity towards uPA (Fig. 5). Similarly, depleted variants (log_2-fold enrichment score

≤ 0, n = 1137) are broadly classified here as loss-of-function, likely include variants that retain a

low level of inhibitory activity towards uPA—again, reflecting that this approach enables the

mapping of functional variability with respect to both gain and loss of function (Fig. 5). While

WT PAI-1 was enriched 2-fold in the selected versus the input library, other variants were

enriched or depleted up to 6- or 23-fold respectively. For comparison, consider two mutations at

Ile[91], I91L and I91N, each representing approximately 0.0005% of the of the input library (Fig.

3). Following selection with uPA, I91L was enriched three-fold, consistent with previous reports

(32). In contrast, I91N was depleted three-fold—demonstrating that while the I91L mutation is well tolerated, I91N results in loss of function with respect to uPA inhibition. Of note, the selection method employed here (complex formation with uPA) does not distinguish between the three potential mechanisms for loss-of-function: PAI-1 misfolding, accelerated transition to the inactive latent state, and/or serving as a substrate for uPA.

Implications for structure-function relationships

The results of our PAI-1 functional screen in the region of the RCL (residues 331-350), are illustrated in Fig. 6. The observed enrichment and depletion at the P1 and P1' positions (residues 346 and 347) are consistent with our understanding of PAI-1 biology. The P1 position has been shown to be a key determinant of SERPIN target protease specificity (33-37), with PAI-1 inhibitory activity toward uPA requiring either a P1 Lys or the WT Arg residue (38) Consistent with these previous reports, no missense mutations were tolerated at P1 in our screen (of note, lysine at this position is absent from our library), with several substitutions significantly depleted (Fig 3 and Fig 6). Consistent with the previously reported tolerance of the P1' position for most amino acid substitutions (38), our screen identified no loss-of-function PAI-1 variants at this position (Fig 6).

At the N-terminus of the RCL, enriched or tolerated substitutions observed in our data generally consist of small aliphatic and polar amino acids. For PAI-1 to retain its inhibitory function, this region of the RCL must be able to insert into β-sheet A (9). These small amino acids allow the RCL to undergo the dramatic conformational changes that are required for this insertion. Consistent with this model, substitutions with bulky and/or charged side chains (Lys, Arg, Pro, Asp, Phe) were the most depleted residues at positions in which the RCL side chains

become oriented into the core beta sheet upon insertion (39-42). In contrast, residues C-terminal to the scissile bond (P2'-P4') are more tolerant of mutations than those at the N-terminus of the RCL, as the former region does not insert into the central β-sheet (43). Finally, the flexibility of the RCL is also important for dictating PAI-1's inhibitory behavior, and our data are concordant with a previous proline-scanning mutagenesis screen (39). Proline residues in the RCL would also be incompatible with RCL insertion into β-sheet A, which transforms it from a largely parallel β-sheet to a more stable anti-parallel β-sheet.

Correlation with predictive algorithms and human genome sequence variant data

A number of algorithms have been developed to predict the impact of single amino acid substitutions on protein function based on evolutionary conservation and/or amino acid type (44). We compared our high throughput screening data with predictions from two commonly used algorithms, SIFT (45) and PolyPhen-2 (46). SIFT predicts the effects of amino acid substitutions by comparison to homologous sequences, and PolyPhen-2 uses both sequence conservation and structural homology to predict the effects of amino acid substitutions on protein function.

The SIFT algorithm prediction was concordant for 745 of the 1137 (66 %) amino acid substitutions scored as "loss of function" in our screen and for 538 of the 826 (65%) scored as neutral. This level of concordance is similar to that previously reported for known deleterious human genetic mutations in other genes (45). PolyPhen-2 exhibited concordance with our data for 994 of 1137 (87%) "loss of function" substitutions, but only 454 of the 836 (54%) neutral PAI-1 amino acid substitutions (46).

Additionally, available human genomic sequence information provides support for the potential value of our data in interpreting the significance of human genetic variation identified by future clinical sequencing. The gnomAD database (31) catalogs human amino acid sequence variant information from ~140,000 human exomic/genomic sequences, including 202 variants scored in our mutation screening analysis. Of these 202 variants, 92 were classified by our data as "loss of function", significantly less than expected by chance (p $= 2\times10^{-4}$ SI Table 3), consistent with evolutionary selective pressure in the human population to maintain PAI-1 activity.

CONCLUSION

Deep mutational scanning has been applied to a number of proteins to analyze function, binding interactions, cellular protein abundance, cell growth/viability, and protein stability (24, 25, 47-49). In the present study, we have adapted this approach to construct a detailed map for the mutational landscape of PAI-1 with respect to the gain or loss of its capacity to inhibit uPA. We anticipate that the mutational landscape of PAI-1 for other serine proteases, including its other canonical substrate, tPA, would likely demonstrate significant differences (50).

Previous PAI-1 mutational studies (51) have been restricted to limited segments of PAI-1 (38, 50), or selected for a few variants with a unique functional impact, such as extended functional stability (32, 52). The error prone PCR approach used here to generate the phPAI-1 mutant library offers speed and ease of application with broad coverage of a significant subset of potential single amino acid substitutions. However, variant coverage is incomplete (Fig. 3), providing significant loss of function data for only a subset of mutation space (Fig 5). Advances in molecular approaches and machine learning algorithms should facilitate a comprehensive map

of the mutation landscape for PAI-1 and many other proteins (53). These data should provide a

valuable resource for the interpretation of sequence variants in PAI-1 and other genes identified

by the expanding clinical application of human whole genome sequencing. Future PAI-1 deep

mutational screening studies should provide additional insight into PAI-1 structure-function

relationships and the determinants of SERPIN target protease specificity.

Figure 1. PAI-1 displayed as a fusion protein on the surface of filamentous phage is an inhibitor of uPA. *(A)* Active PAI-1 (PDB: 3PB1) reacts with free uPA (PDB:1W0Z) to form a

covalent uPA:PAI-1 complex (54, 55). Both free and complexed uPA can be immunoprecipitated using an anti-uPA antibody. Alternatively, PAI-1 can spontaneously relax from its active, metastable state to a low-energy yet chemically inert latent conformation (PDB: 1DVN) (56). PAI-1's reactive center loop is highlighted in orange. *(B)* Schematic of phPAI-1 displayed as a fusion to the pIII coat protein of M13 filamentous phage, with the N-terminal myc- and C-terminal FLAG- and E-tags highlighted.

Figure 2. Immunoprecipitation is specific for phPAI-1:uPA complexes. Nine parts phVWF-A3 and one part phPAI-1 (9:1) were combined (*input*, n = 3), incubated with uPA (1.7 nM) for 30 min at 37°C, and selected by immunoprecipitation with an anti-uPA antibody (*selected*, n =3). For each replica, 24 colonies were genotyped by PCR using primers common to both the phVWF-A3 (677 bp) and phPAI-1 inserts (1271 bp) followed by analysis on a 1% agarose gel.

Figure 3. phPAI-1 mutant library generated by error prone PCR includes more than two-thirds of all possible missense mutations. The mutational library contains 71% of all possible missense and nonsense mutations with 27% of all missense variants present with sufficient depth (base mean score > 10, p_{adj} < 0.05) to accurately determine the effects of the mutation on PAI-1 function. The primary amino acid position within PAI-1 is indicated along the x-axis and single amino acid substitutions are listed along the y-axis. WT amino acid residues are indicated in yellow, while missense and nonsense (X) mutations not present in the input library are shown in white. Variants present within the library are shown in grey (see scale) as a percentage of the input library represented by that variant.

Figure 4. phPAI-1 uPA-selected libraries are enriched in variants that retain their inhibitory function and depleted of those without inhibitory activity. An MA plot (29) of base mean score (average of counts in the input and selected libraries) vs. log₂-fold change is shown, with missense mutations in pink and nonsense variants in blue. Variants with $p_{adj} < 0.05$ are shown as closed circles, while those that do not meet this significance threshold are shown as open circles. A base mean score greater than 10 was also set as a threshold for determining significance.

Figure 5. phPAI-1 mutational libraries contain missense variants that result in loss of function. Amino acid position is indicated on the x-axis, while amino acid substitutions are indicated on the y-axis. Loss of function missense variants, as well as those with a reduced capacity to inhibit uPA, are shown in red with shading as a function of their log_2-fold score, while variants in blue retain PAI-1 inhibitory function. The intensity of shading scales with the degree of enrichment or depletion. WT amino acid residues are shown in black, while beige indicates missense mutations that were present in the mutational library but did not meet significance thresholds. White indicates amino acid substitutions that were not present in the mutational library.

Figure 6. Deep mutational scanning of PAI-1 provides insight into the mutational landscape of PAI-1's RCL. Relative $\log_2$-fold change scores of missense mutations in PAI-1's RCL (amino acids 331-350) following selection by ability to inhibit uPA (36). Amino acids below the x-axis indicate depletion ($\log_2$-fold change < 0), while those above the x-axis indicate enrichment ($\log_2$-fold change > 0). Letter height corresponds to the relative $\log_2$-fold change. Residues are color coded by properties: acidic residues in red, basic residues in blue, polar amino acids in green, non-polar amino acids in orange, and Pro in purple (36). The P1 and P1' positions (Met[346] and Arg[347] in WT) are also indicated.

REFERENCES

1.   Irving JA, Pike RN, Lesk AM, Whisstock JC. Phylogeny of the serpin superfamily: implications of patterns of amino acid conservation for structure and function. Genome research. 2000;10(12):1845-64.
2.   Huntington JA. Serpin structure, function and dysfunction. Journal of thrombosis and haemostasis : JTH. 2011;9 Suppl 1:26-34.
3.   Stein PE, Carrell RW. What do dysfunctional serpins tell us about molecular mobility and disease? Nature structural biology. 1995;2(2):96-113.
4.   Lawrence DA, Ginsburg D, Day DE, Berkenpas MB, Verhamme IM, Kvassman JO, et al. Serpin-protease complexes are trapped as stable acyl-enzyme intermediates. J Biol Chem. 1995;270(43):25309-12.
5.   Silverman GA, Bird PI, Carrell RW, Church FC, Coughlin PB, Gettins PG, et al. The serpins are an expanding superfamily of structurally similar but functionally diverse proteins. Evolution, mechanism of inhibition, novel functions, and a revised nomenclature. J Biol Chem. 2001;276(36):33293-6.
6.   Huntington JA, Read RJ, Carrell RW. Structure of a serpin-protease complex shows inhibition by deformation. Nature. 2000;407(6806):923-6.
7.   Fredenburgh JC, Stafford AR, Weitz JI. Conformational changes in thrombin when complexed by serpins. J Biol Chem. 2001;276(48):44828-34.
8.   Peterson FC, Gettins PG. Insight into the mechanism of serpin-proteinase inhibition from 2D [1H-15N] NMR studies of the 69 kDa alpha 1-proteinase inhibitor Pittsburgh-trypsin covalent complex. Biochemistry. 2001;40(21):6284-92.
9.   Lawrence DA, Olson ST, Muhammad S, Day DE, Kvassman JO, Ginsburg D, et al. Partitioning of serpin-proteinase reactions between stable inhibition and substrate cleavage is regulated by the rate of serpin reactive center loop insertion into beta-sheet A. J Biol Chem. 2000;275(8):5839-44.
10.   Khan MS, Singh P, Azhar A, Naseem A, Rashid Q, Kabir MA, et al. Serpin Inhibition Mechanism: A Delicate Balance between Native Metastable State and Polymerization. J Amino Acids. 2011;2011:606797-.
11.   Marijanovic EM, Fodor J, Riley BT, Porebski BT, Costa MGS, Kass I, et al. Reactive centre loop dynamics and serpin specificity. Scientific Reports. 2019;9(1):3870.
12.   Scott BM, Sheffield WP. Engineering the serpin α(1) -antitrypsin: A diversity of goals and techniques. Protein Sci. 2020;29(4):856-71.
13.   Sprengers ED, Kluft C. Plasminogen activator inhibitors. Blood. 1987;69(2):381-7.
14.   Alessi MC, Declerck PJ, De Mol M, Nelles L, Collen D. Purification and characterization of natural and recombinant human plasminogen activator inhibitor-1 (PAI-1). Eur J Biochem. 1988;175(3):531-40.
15.   Vassalli JD, Sappino AP, Belin D. The plasminogen activator/plasmin system. The Journal of clinical investigation. 1991;88(4):1067-72.
16.   Cesarman-Maus G, Hajjar KA. Molecular mechanisms of fibrinolysis. British journal of haematology. 2005;129(3):307-21.
17.   Dieval J, Nguyen G, Gross S, Delobel J, Kruithof EK. A lifelong bleeding disorder associated with a deficiency of plasminogen activator inhibitor type 1. Blood. 1991;77(3):528-32.

18.   Lee MH, Vosburgh E, Anderson K, McDonagh J. Deficiency of plasma plasminogen activator inhibitor 1 results in hyperfibrinolytic bleeding. Blood. 1993;81(9):2357-62.

19.   Schleef RR, Higgins DL, Pillemer E, Levitt LJ. Bleeding diathesis due to decreased functional activity of type 1 plasminogen activator inhibitor. The Journal of clinical investigation. 1989;83(5):1747-52.

20.   Fay WP, Parker AC, Condrey LR, Shapiro AD. Human plasminogen activator inhibitor-1 (PAI-1) deficiency: characterization of a large kindred with a null mutation in the PAI-1 gene. Blood. 1997;90(1):204-8.

21.   Fay WP, Shapiro AD, Shih JL, Schleef RR, Ginsburg D. Brief report: complete deficiency of plasminogen-activator inhibitor type 1 due to a frame-shift mutation. The New England journal of medicine. 1992;327(24):1729-33.

22.   Carmeliet P, Kieckens L, Schoonjans L, Ream B, van Nuffelen A, Prendergast G, et al. Plasminogen activator inhibitor-1 gene-deficient mice. I. Generation by homologous recombination and characterization. The Journal of clinical investigation. 1993;92(6):2746-55.

23.   Khan SS, Shah SJ, Klyachko E, Baldridge AS, Eren M, Place AT, et al. A null mutation in SERPINE1 protects against biological aging in humans. Science Advances. 2017;3(11):eaao1617.

24.   Kretz CA, Tomberg K, Van Esbroeck A, Yee A, Ginsburg D. High throughput protease profiling comprehensively defines active site specificity for thrombin and ADAMTS13. Sci Rep. 2018;8(1):2788.

25.   Kretz CA, Dai M, Soylemez O, Yee A, Desch KC, Siemieniak D, et al. Massively parallel enzyme kinetics reveals the substrate recognition landscape of the metalloprotease ADAMTS13. Proc Natl Acad Sci U S A. 2015;112(30):9328-33.

26.   Yee A, Tan FL, Ginsburg D. Functional display of platelet-binding VWF fragments on filamentous bacteriophage. PloS one. 2013;8(9):e73518.

27.   Levin EG, Santell L. Conversion of the active to latent plasminogen activator inhibitor from human endothelial cells. Blood. 1987;70(4):1090-8.

28.   Yee A, Dai M, Croteau SE, Shavit JA, Pipe SW, Siemieniak D, et al. Phage Display Broadly Identifies Inhibitor-reactive Regions in von Willebrand Factor. bioRxiv. 2021:2021.03.08.434464.

29.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome biology. 2014;15(12):550.

30.   Sim N-L, Kumar P, Hu J, Henikoff S, Schneider G, Ng PC. SIFT web server: predicting effects of amino acid substitutions on proteins. Nucleic Acids Research. 2012;40(W1):W452-W7.

31.   Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alföldi J, Wang Q, et al. The mutational constraint spectrum quantified from variation in 141,456 humans. Nature. 2020;581(7809):434-43.

32.   Berkenpas MB, Lawrence DA, Ginsburg D. Molecular evolution of plasminogen activator inhibitor-1 functional stability. EMBO J. 1995;14(13):2969-77.

33.   Chuang YJ, Swanson R, Raja SM, Bock SC, Olson ST. The antithrombin P1 residue is important for target proteinase specificity but not for heparin activation of the serpin. Characterization of P1 antithrombin variants with altered proteinase specificity but normal heparin activation. Biochemistry. 2001;40(22):6670-9.

34.   Owen MC, Brennan SO, Lewis JH, Carrell RW. Mutation of antitrypsin to antithrombin. alpha 1-antitrypsin Pittsburgh (358 Met leads to Arg), a fatal bleeding disorder. The New England journal of medicine. 1983;309(12):694-8.

35.   Rashid Q, Kapil C, Singh P, Kumari V, Jairajpuri MA. Understanding the specificity of serpin-protease complexes through interface analysis. Journal of biomolecular structure & dynamics. 2015;33(6):1352-62.

36.   Thomsen MC, Nielsen M. Seq2Logo: a method for construction and visualization of amino acid binding motifs and sequence profiles including sequence weighting, pseudo counts and two-sided representation of amino acid enrichment and depletion. Nucleic Acids Res. 2012;40(Web Server issue):W281-7.

37.   Lawrence DA, Strandberg L, Ericson J, Ny T. Structure-function studies of the SERPIN plasminogen activator inhibitor type 1. Analysis of chimeric strained loop mutants. J Biol Chem. 1990;265(33):20293-301.

38.   Sherman PM, Lawrence DA, Yang AY, Vandenberg ET, Paielli D, Olson ST, et al. Saturation mutagenesis of the plasminogen activator inhibitor-1 reactive center. J Biol Chem. 1992;267(11):7588-95.

39.   Audenaert AM, Knockaert I, Collen D, Declerck PJ. Conversion of plasminogen activator inhibitor-1 from inhibitor to substrate by point mutations in the reactive-site loop. J Biol Chem. 1994;269(30):19559-64.

40.   Tucker HM, Mottonen J, Goldsmith EJ, Gerard RD. Engineering of plasminogen activator inhibitor-1 to reduce the rate of latency transition. Nature structural biology. 1995;2(6):442-5.

41.   Gils A, Knockaert I, Declerck PJ. Substrate behavior of plasminogen activator inhibitor-1 is not associated with a lack of insertion of the reactive site loop. Biochemistry. 1996;35(23):7474-81.

42.   Lawrence DA, Olson ST, Palaniappan S, Ginsburg D. Serpin reactive center loop mobility is required for inhibitor function but not for enzyme recognition. J Biol Chem. 1994;269(44):27657-62.

43.   Mottonen J, Strand A, Symersky J, Sweet RM, Danley DE, Geoghegan KF, et al. Structural basis of latency in plasminogen activator inhibitor-1. Nature. 1992;355(6357):270-3.

44.   Knecht C, Krawczak M. Molecular genetic epidemiology of human diseases: from patterns to predictions. Human Genetics. 2014;133(4):425-30.

45.   Ng PC, Henikoff S. Predicting deleterious amino acid substitutions. Genome research. 2001;11(5):863-74.

46.   Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, et al. A method and server for predicting damaging missense mutations. Nat Methods. 2010;7(4):248-9.

47.   Fowler DM, Fields S. Deep mutational scanning: a new style of protein science. Nat Methods. 2014;11(8):801-7.

48.   Fowler DM, Stephany JJ, Fields S. Measuring the activity of protein variants on a large scale using deep mutational scanning. Nat Protoc. 2014;9(9):2267-84.

49.   Stein A, Fowler DM, Hartmann-Petersen R, Lindorff-Larsen K. Biophysical and Mechanistic Models for Disease-Causing Protein Variants. Trends Biochem Sci. 2019;44(7):575-88.

50.   Sherman PM, Lawrence DA, Verhamme IM, Paielli D, Shore JD, Ginsburg D. Identification of tissue-type plasminogen activator-specific plasminogen activator inhibitor-1

mutants. Evidence that second sites of interaction contribute to target specificity. J Biol Chem. 1995;270(16):9301-6.

51.   De Taeye B, Gils A, Declerck PJ. The story of the serpin plasminogen activator inhibitor 1: is there any need for another mutant? Thrombosis and haemostasis. 2004;92(5):898-924.

52.   Stoop AA, Eldering E, Dafforn TR, Read RJ, Pannekoek H. Different structural requirements for plasminogen activator inhibitor 1 (PAI-1) during latency transition and proteinase inhibition as evidenced by phage-displayed hypermutated PAI-1 libraries. Journal of molecular biology. 2001;305(4):773-83.

53.   Fernandez-de-Cossio-Diaz J, Uguzzoni G, Pagnani A. Unsupervised Inference of Protein Fitness Landscape from Deep Mutational Scan. Molecular Biology and Evolution. 2020;38(1):318-28.

54.   Lin Z, Jiang L, Yuan C, Jensen JK, Zhang X, Luo Z, et al. Structural Basis for Recognition of Urokinase-type Plasminogen Activator by Plasminogen Activator Inhibitor-1*. Journal of Biological Chemistry. 2011;286(9):7027-32.

55.   Zeslawska E, Jacob U, Schweinitz A, Coombs G, Bode W, Madison E. Crystals of urokinase type plasminogen activator complexes reveal the binding mode of peptidomimetic inhibitors. Journal of molecular biology. 2003;328(1):109-18.

56.   Stout TJ, Graham H, Buckley DI, Matthews DJ. Structures of Active and Latent PAI-1:⬜ A Possible Stabilizing Role for Chloride Ions. Biochemistry. 2000;39(29):8460-9.

SI Table 1. Primer Sequences

Primer	5' → 3' Sequence
pAY-E FLAG tag Primers	
pAY-E FLAG For	/5Phos/GGCCGCAGGTGGTGGTGACTACAAGGACGATGACGATAAAGGA GGAGGAGGTGCG
pAY-E FLAG Rev	CCGGCGCACCTCCTCCTCCTTTATCGTCATCGTCCTTGTAGTCACCACC ACCTGC
CcdB-CmR Cassette Amplification	
CcdB-CmR For	/5Phos/TCGGCCTAGCCGGCCGGCGCGCCATTAGGCACCCCAGGCTTTAC ACTTTATGCTT
CcdB-CmR Rev	TGCGGCCGCGTCGACCTGCAGACTGGCTGTGTATAA
pAY-FE PAI-1 Cloning Primers	
pAY-FE PAI-1 For	GCCGGCGCGCCAGAACAAAAACTCATCTCAGAAGAGGATC TGGGTGGAGGTTCAGTGCACCATCCCCCATCCTAC
pAY-FE PAI-1 Rev	ACCTGCGGCCGCTGAACCACCTCCGGGTTCTATCACTTGGCCCAT
pAY-FE PAI-1 Mutagenesis Primers	
PAI-1 Mutagenesis For	TTCTCGGCCTAGCCGGCCGG
PAI-1 Mutagenesis Rev	GTCCTTGTAGTCACCACCACCTGCGGCC
Sanger Sequencing/Colony PCR Primers	
pAYE Sequencing For	TTCATGCTGCCGGCTTTCTCGG
pAYE Sequencing Rev	TCGTCATCGTCCTTGTAGTCACCACCA
HTS Sequencing Primers	
PAI-1 Amplicon 1 For	NNNNNNGAACAAAAACTCATCTCAGAAGAGGATCTG
PAI-1 Amplicon 1 Rev	NNNNNNGGGTGAGAAAACCACGTTGCG
PAI-1 Amplicon 2 For	NNNNNNTTTCAGCAGGTGGCGCAG
PAI-1 Amplicon 2 Rev	NNNNNNCTTGTCATCAATCTTGAATCCCATAGCTGC
PAI-1 Amplicon 3 For	NNNNNNATGCTCCAGCTGACAACAGGA
PAI-1 Amplicon 3 Rev	NNNNNNTGTGGTGCTGATCTCATCCTTGTT
PAI-1 Amplicon 4 For	NNNNNNGCCCTCCGGCATCTGTACAAG
PAI-1 Amplicon 4 Rev	NNNNNNTTGCTTGACCGTGCTCCGG
PAI-1 Amplicon 5 For	NNNNNNAAGCTGGTCCAGGGCTTCATG
PAI-1 Amplicon 5 Rev	NNNNNNCCCAAGCAAGTTGCTGATCATACC
PAI-1 Amplicon 6 For	NNNNNNATTCATCATCAATGACTGGGTGAAGAC
PAI-1 Amplicon 6 Rev	NNNNNNCTGGAGTCGGGGAAGGGAG
PAI-1 Amplicon 7 For	NNNNNNAATGCCCTCTACTTCAACGGC
PAI-1 Amplicon 7 Rev	NNNNNNGGGCGTGGTGAACTCAGTATAG
PAI-1 Amplicon 8 For	NNNNNNCCCATGATGGCTCAGACCAAC
PAI-1 Amplicon 8 Rev	NNNNNNGGCAGAGAGAGGCACCTCT
PAI-1 Amplicon 9 For	NNNNNNAGCATGTTCATTGCTGCCCC
PAI-1 Amplicon 9 Rev	NNNNNNTTCAGTCTCCAGGGAGAACTTGG
PAI-1 Amplicon 10 For	NNNNNNCAGGCTGCCCCGCCT
PAI-1 Amplicon 10 Rev	NNNNNNACGTGGAGAGGCTCTTGGTC
PAI-1 Amplicon 11 For	NNNNNNAGACAGTTTCAGGCTGACTTCAC
PAI-1 Amplicon 11 Rev	NNNNNNGGGGGCCATGCGGGC

PAI-1 Amplicon 12 For	NNNNNNTCATCCACAGCTGTCATA
PAI-1 Amplicon 12 Rev	NNNNNNTGCGGCCGCTGAACCACCTCC

SI Table 2. PAI-1 amino acid residues spanned by HTS amplicons

PAI-1 Amplicon	Residues Sequenced	Residues Analyzed
1	V1 – D29	V1-K28
2	A26 – A60	D29-Q56
3	G52 – W86	Q57-M83
4	E81 – L116	G84-F114
5	P111 – K145	R115-T144
6	H143 – K176	K145-W175
7	Q174 – F208	K176-F208
8	K207 – E242	N209-E243
9	Y241 – L275	K244-V274
10	L273 – S310	L275-L309
11	S308 – S344	S310-V343
12	V343 – P379	S344-P379

SI Table 3. Observed frequencies for χ^2 test comparing experimental results to gnomAD database

	Scored by phage display	Present in gnomAD	Not present in gnomAD
Functional	826	110	716
Non-functional	1137	92	1045
Total	1963	202	1761