Genomic organization of the autonomous regulatory domain of *eyeless* locus

in *Drosophila melanogaster*

Shreekant Verma, Rashmi U Pathak and Rakesh K Mishra*

Centre for Cellular and Molecular Biology, Council of Scientific and Industrial

Research, Uppal Road, Hyderabad 500007, India

* Corresponding author: email: mishra@ccmb.res.in

16 Running title: Regulatory domain of *eyeless* locus in *Drosophila*

Abstract

In *Drosophila*, expression of *eyeless* (*ey*) gene is restricted to the developing eyes and central nervous system. However, the flanking genes, *myoglianin* (*myo*), and bent (*bt*) have different temporal and spatial expression patterns as compared to the *ey*. How distinct regulation of *ey* is maintained is mostly unknown. Earlier, we have identified a boundary element intervening *myo* and *ey* genes (ME boundary) that prevents the crosstalk between the *cis*-regulatory elements of *myo* and *ey* genes. In the present study, we further searched for the *cis*-elements that define the domain of *ey* and maintain its expression pattern. We identify another boundary element between *ey* and *bt*, the EB boundary. The EB boundary separates the regulatory landscapes of *ey* and *bt* genes. The two boundaries, ME and EB, show a long-range interaction as well as interact with the nuclear architecture. This suggests functional autonomy of the *ey* locus and its insulation from differentially regulated flanking regions. We also identify a new Polycomb Response Element, the *ey*-PRE, within the *ey* domain. The expression state of the *ey* gene, once established during early development is likely to be maintained with the help of *ey*-PRE. Our study proposes a general regulatory mechanism by which a gene can be maintained in a functionally independent chromatin domain in gene-rich euchromatin.

Introduction

The *Drosophila eyeless* gene (*ey*), a *Pax*-6 homolog, is an essential regulatory gene required for development of the eye (QUIRING *et al.* 1994). It is one of the many evolutionarily conserved genes that are involved in retinal determination. Molecular genetic studies have shown that *ey* along with other genes, viz., *eyes absent* (*eya*), *twin of eyeless* (*toy*), *sine oculis* (*so*), and *dachshund* (*dac*), is involved in retinal determination network (DESPLAN 1997; HALDER *et al.* 1998; PICHAUD AND DESPLAN 2001). Although the genetic interactions of *ey* with these genes are mostly known, spatial and temporal transcriptional regulation of the *ey* is less well understood.

The *ey* gene, present on the fourth chromosome of *Drosophila*, is flanked closely by *myoglianin* (*myo*) gene at the upstream, and *bent* (*bt*) gene downstream. The three genes show a very distinct spatiotemporal pattern of expression in the fly. The expression of *ey* is restricted to eye disc primordia and central nervous system (CNS) during embryogenesis. Later during larval stages it is expressed in eye imaginal discs and CNS, where it continues to express in adults (QUIRING *et al.* 1994; HAUCK *et al.* 1999; ADACHI *et al.* 2003; CHINTAPALLI *et al.* 2007). Unlike *ey, myo* gene that codes for a TGF-β superfamily protein has a high-level of maternal transcript deposition in the early embryo. Later, it expresses in the somatic, visceral, and heart musculature where its expression persists throughout embryogenesis. In third instar larvae, *myo* expression is restricted to brain and glial cells of the ventral nerve cord (LO AND FRASCH 1999; CHINTAPALLI *et al.* 2007). Likewise, the *bt* gene encodes for a titin superfamily muscle protein 'Projectin'. During development, the *bt* transcripts first appear in the mid-embryo stage and increase steadily till the adult stage (FYRBERG *et al.* 1992; MAROTO *et al.* 1992). The expression of *bt* has been found in the embryonic/larval muscles system and all types of muscles in pupae and adults (AYME-SOUTHGATE *et al.* 1991; FRISE *et al.* 2010).

Transcriptional regulation of *ey* at various developmental stages by multiple *ey* enhancers has been reported in two earlier studies (HAUCK *et al.* 1999; ADACHI *et al.* 2003). An initial study has identified a 212-bp eye-specific enhancer at 3' end of the second intron that is essential for expression of *ey* in larval eye-disc primordia (HAUCK *et al.* 1999). A later study has described a 5 kb upstream region and a 3.6 kb

second intronic fragment that act synergistically to specify the *ey* specific expression
pattern of lacZ reporter construct in developing CNS (ADACHI *et al.* 2003). The
targeted expression of *ey* induces ectopic eyes in *Drosophila* (HALDER *et al.* 1998).
This suggests that active repression of *ey* is crucial in other tissues, wherever it is
not required. However, what maintains the repressed status of *ey* in such tissues, is
not known. The distinct spatiotemporal expression of *ey* calls for the presence of
other *cis*-regulatory elements like chromatin domain boundary elements (CBEs) or
insulators, and also a memory element like Polycomb response elements (PREs).
Such *cis*-elements at *ey* locus are required to prevent the cross-talk among the
regulatory elements of the three genes and for maintenance of tissue-specific
expression, as is the case at several well characterised loci like bithorax complex
in *Drosophila* (MAEDA AND KARCH 2006; MIHALY *et al.* 2006).

Boundary elements are an established regulatory feature of most eukaryotic
genomes. They insulate a gene from the influence of neighbouring chromatin and
actively subdivide the genome into chromatin domains of independent gene activity.
The regulatory function of these elements is dependent on their association with
several boundary interacting proteins including BEAF-32, CTCF, Su(Hw), Zw5,
GAGA factor (GAF or Trl) and CP190 (GEYER *et al.* 1986; KELLUM AND SCHEDL 1991;
GERASIMOVA AND CORCES 2001; WEST *et al.* 2002; GASZNER AND FELSENFELD 2006;
VALENZUELA AND KAMAKAKA 2006). On the other hand, PREs also known as Cellular
Memory Modules (CMMs) are specific DNA elements, recruitment sites for Polycomb
group and trithorax group (PcG/trxG) of proteins, that are required to maintain the
transcriptional states of the target genes(SIGRIST AND PIRROTTA 1997; CAVALLI AND
PARO 1998). The PcG proteins are required for stable silencing, whereas trxG
proteins promote activation of target genes. Many PREs studied in *Drosophila*
suggest their role in maintenance of repressed states. One such example is *iab7*-
PRE in the *Drosophila* bithorax complex, The *iab7*-PRE helps in appropriate
maintenance of *Abd-B* gene expression pattern in parasegement 12 (MISHRA *et al.*
2001). Studies based on techniques such as FISH and chromosome conformation
capture (3C) assay, support interactions among CBEs and other regulatory elements
like promoters, enhancers and silencers. This leads to the formation of chromatin
domains with autonomous regulatory function (DEKKER *et al.* 2002; BANTIGNIES *et al.*
2003; RONSHAUGEN AND LEVINE 2004; LANZUOLO *et al.* 2007; SEXTON *et al.* 2009; HOU

et al. 2012; ZHANG *et al.* 2013). These higher order organization of chromatin

interaction are known as topological associated domains (TADs). (HOU *et al.* 2012;

SEXTON *et al.* 2012; ZHANG *et al.* 2013; CUBENAS-POTTS *et al.* 2017). Anchoring of

regulatory elements to nuclear architecture also helps in compartmentalization of the

locus into functionally independent domain (PATHAK *et al.* 2007).

In the present study, we have analyzed the *ey* locus to identify *cis*-

regulatory elements that define its precise and distinct expression pattern. Earlier,

we have identified a CBE upstream of *ey* that separates regulatory domains of *myo*

and *ey* genes (SULTANA *et al.* 2011). Here, we have identified a CBE

between *ey* and *bt*, two closely spaced but distinctly expressed genes. We refer to

this element as EB boundary and functionally characterize it in the transgenic

context. Additionally, we have identified a PRE (*ey*-PRE) that functions as a

repressor to maintain the expression pattern of *ey*. Furthermore, to understand the

mechanism of *ey* regulation by EB, *ey*-PRE, and previously identified ME boundary,

we have explored their long-range interactions in the context of nuclear architecture.

Our findings reveal two new regulatory elements in this important locus, and the

structural framework of this genomic locus based on the higher-order chromatin

organization of regulatory elements. Such organization may have a general

implication in higher eukaryotes.

Results

Identification of potential CBEs that separate ey and bt loci

Previously, we have identified a CBE (the ME boundary) in the ~1.6 kb intergenic

region between *myo* and *ey* genes, endorsing the rationale that CBEs are required to

prevent crosstalk between the regulatory elements of closely spaced but differently

expressed genes (SULTANA *et al.* 2011). Likewise, *ey* and the downstream

gene *bt,* have a disparate pattern of expression (AYME-SOUTHGATE *et al.* 1991).

Therefore, we expected a CBE to be present in the 3.2 kb intergenic region to block

crosstalk between the adjacent regulatory domains. To identify a potential CBE

between *ey* and *bt*, we used cdBEST search tool which has been successfully

utilized to identify several new CBEs in *Drosophila* and other insects like *Anopheles*

gambie (SRINIVASAN AND MISHRA 2012; AHANGER *et al.* 2013). CdBEST predicts

CBEs based on the presence of clusters of binding motifs of many known boundary

proteins like BEAF-32, CTCF, GAF, CP190, and *Zw5,* in the queried sequence.

CdBEST analysis for the ~30 kb *ey* locus predicted two CBEs, one being the

previously reported ME boundary and a new putative CBE in the seventh intron

of *ey* gene (INT7) (Figure 1). No CBE is predicted in the intergenic region between

ey and *bt,* although the region carried multiple binding motifs of BEAF-32, CTCF,

and GAF. Therefore, we analyzed the data available for *in-vivo* binding of boundary

proteins in this region using embryonic genome-wide ChIP-chip datasets from the

modENCODE project (http://www.modencode.org/) (NEGRE *et al.* 2010). We

observed a significantly high occupancy of CTCF and CP190 in a ~1.2 Kb region

within INT7. However, a prominent binding of both of these proteins is also observed

in the ~2.2 Kb region between *ey* and *bt* (EB). Particularly, CTCF binding at EB

occurs in two separate regions, one in a ~0.8 kb 3' UTR region of *ey* (named as EB-

u), and the other in adjacent ~1.1 kb intergenic region (named as EB-i). EB-i

additionally has binding of BEAF-32. We did not find a significant binding of GAF in

these regions (Figure 1).

As the existence of DNaseI hypersensitive sites (DHS) are inherent feature of

regulatory elements, including several known CBEs we next examined the presence

of DHS in INT7 and EB using the DHS dataset of embryonic stage 9 (6 h embryo)

available at UCSC genome browser (http://genome.ucsc.edu/) (THOMAS *et al.* 2011).

Both INT7 and EB contained a prominent DHS (Figure 1 and b). These observations

altogether indicate potential CBE function of INT7 and EB (Figure 1b, Supplementary

Figure 1, Supplementary Table 1).

EB functions as an enhancer blocker in the eye

To test the boundary activity of INT7 and EB, we generated transgenic flies carrying

test fragments introduced between the *white* enhancer and *mini-white* reporter gene

in enhancer blocker assay construct (pRW+) (Figure 2a) (HAGSTROM *et al.* 1996).

The level of eye pigmentation in adult transgenic flies is a responsive indicator of the

amount of *mini-white* transcription. In case a CBE placed between the enhancer

and *mini-white* acts as an enhancer blocker, it would reduce the level of *mini-*
white expression that can be scored as light eye colour in adult flies. However, a
random DNA fragment in the same position would have no effect (HAGSTROM *et al.*
1996). The test fragments in the assay vector are flanked by *lox*P sites so that it can
be flipped out using the *cre-lox*P system to confirm the enhancer blocker effect and
rule out any position effect.

All the transgenic fly lines generated for INT7 had a high level of eye pigmentation
(red to bright red eye colour) in heterozygous condition. Moreover, the removal of
INT7 in four such initial lines (P) did not change in the level of eye pigment in their
flipped-outs (ΔP) indicating that INT7 lacked an enhancer-blocker activity (Figure 2b,
Supplementary Table 2). In contrast, seven transgenic lines carrying EB fragment
displayed varying degrees of light red eye colour in heterozygous conditions, and
four of them turned darker when EB was flipped out. We found a mild increase in eye
colour in all EB flipped-out lines, and at least one of them showed a ~2-fold increase
in eye pigment level (Figure 2b, Supplementary Table 2). Homozygous (P/P)
transgenic lines mostly showed a high level of eye pigmentation (dark red eye
colour) for both the INT7 and EB transgenes. These results suggested that the EB
acts as a weak enhancer blocker in the eye and the flipped-out lines confirmed that
light eye colour is due to the boundary function of EB and not a position effect.

As explained in the earlier section, EB can be split into two fragments based on
CTCF binding sites as EB-u and EB-i. The binding of BEAF-32, however, is detected
only in the EB-i region. Both of these regions contain distinct DHS, which is more
prominent in the EB-i (Figure 1b). We further tested the enhancer blocking activity of
both the fragments separately, to identify whether the boundary function lies in one
of these regions or the entire EB sequence is required for the activity. All the EB-u
transgenic lines had dark red eye colour and did not exhibit enhancer-blocker effect
as no change in eye colour was observed in flipped-out lines. In contrast, EB-i
transgenic lines showed light eye colour, and five out of seven lines showed a mild
(~1.5 fold) increase in eye colour after P-element flipping out, suggesting that
enhancer blocker activity of EB mainly resides in EB-i (Figure 2b, Supplementary
Table 2).

In summary, the presence of DHS with significant boundary proteins occupancy at
INT7 and EB suggested their boundary features, but only EB displayed enhancer
blocker effect in the adult eye. As the CBE features were observed using datasets
from embryonic developmental stages, we reasoned that these regions might be
functional as a boundary mostly during early development. Therefore, we tested the
boundary activity of INT7 and EB in the embryo.

EB functions as an enhancer blocker in embryo

We used a P-element based CfhL assay vector to test boundary activity in
developing embryos (Figure 3a) (HAGSTROM *et al.* 1996). The construct carries two
reporter   genes *mini-white* and *hsp70/lacZ.* Two *ftz* enhancers,   an   upstream
enhancer (UPS) active during early development and a late neurogenic enhancer
(NE), drive the *hsp70/lacZ* in the embryo. The level of eye pigmentation and x-gal
staining in the embryo both indicates the expression level of *mini-white* and *lacZ*,
respectively. A test DNA inserted between the *lacZ* promoter and the enhancers that
act as an enhancer blocker causes a reduced level of *lacZ* expression. The test
element can be flipped out in transgenes to confirm the enhancer-blocker activity
and rule out any position effect. In addition, the *mini-white* expression level in the eye
can be compared to rule out a repressor effect of the test elements (HAGSTROM *et al.*
1996; SULTANA *et al.* 2011).

In EB transgenic lines, we observed significantly reduced X-gal staining in both early
and late embryonic stages suggesting EB prevented both the enhancers from acting
on *lacZ* (Figure 3b-c and Supplementary Figure 2). EB in homozygotic condition
displayed a strong boundary activity which was similar to the known *Fab7* boundary
(HAGSTROM *et al.* 1996). In two out of three EB flipped out version of the transgenic
lines, an intense X-gal staining was restored (comparable to empty vector). At the
same time, the eye colour remained the same, suggesting that EB functions as an
enhancer blocker but not a repressor (Figure 3b-c, Supplementary Figure 2 and
Supplementary Table 3). Interestingly, INT7 transgenes also displayed a weak
enhancer blocker activity in the embryo. Two out of three transgenic lines of INT7
showed a moderate increase in the X-gal staining in both early and late stages
embryos after flipping out the test elements (Figure 3b-c, Supplementary Table 3,

Supplementary Figure 2). When EB-u and EB-i fragments were tested separately,
EB-u did not exhibit any enhancer blocker activity in the embryo as an intense X-gal
staining comparable to empty vector was observed. The EB-i region, on the other
hand, showed a strong enhancer blocker activity, comparable to EB. Like EB, flipped
out EB-i lines showed a significant increase in the X-gal staining with no change in
eye colour. These observations indicated that mainly the EB-i, which carries BEAF-
32 binding sites harbours the boundary activity in this region (Figure 3b-c,
Supplementary Figure 2 and Supplementary Table 3). Altogether these results
suggest that EB and INT7 are active boundaries during embryonic development.
Further, a weak enhancer blocker activity of INT7 in the embryo suggest its assistive
function to EB (a strong boundary) in demarcating the *ey* locus.

Identification of a putative PRE associated with ey locus

It is known that multiple enhancers positively regulate the *ey* gene in developing eye
and CNS (HAUCK *et al.* 1999; ADACHI *et al.* 2003). However, the regulatory elements
that maintain the repressed state of *ey* in other tissues are not known.
In *Drosophila,* PREs are involved in establishing and keeping the repressed state of
genes (MISHRA *et al.* 2001; AMERICO *et al.* 2002). We, therefore, searched for
putative PRE in *ey* locus using a Perl-based PRE mapper tool (SRINIVASAN AND
MISHRA 2020). This tool uses a motif cluster search of known DNA binding PcG
recruiters like Pho, GAF, Dsp1, Sp1, and Cg. PRE mapper predicted a ~1.6 kb
putative PRE region upstream of the exon 1 of *ey.* We named this element as *ey-*
PRE, and observed that it has clusters of Sp1 (3), Cg (3), Pho (1), and GAF (1)
binding sites in an internal core region of ~800 bp. Interestingly, the *ey*-PRE also
contained a second *ey* promoter region that is predicted in the eukaryotic promoter
database (DREOS *et al.* 2015) (Figure 4).

To assess whether the *ey*-PRE is a target of PcG proteins, we looked into the
presence of repressive histone modification (H3K27me3) and binding of PcG
proteins (PRC1 members - Pc, Psc, and dRing) using ChIP-seq data of 14-16h
embryo available at the modENCODE database (MOD *et al.* 2010). We observed an
H3K27me3 domain with significant enrichment of Pc in *ey* locus. Interestingly, the
repressive H3K27me3 modification is primarily confined to *ey* locus only, whereas

flanking regions toward *myo* and *bt* hold active histone modification of H3K27ac. At
the same time, boundary proteins namely BEAF-32, CP190, and CTCF (Chip-seq
data from modENCODE project) are present at both, ME and EB boundaries that
demarcate the borders of the repressive domain of *ey* (Figure 4c). Presence of a Pc-
enriched repressive domain restricted to *ey* locus indicates that a repressive *cis*-
acting element is very likely to be present within the locus. Additionally, bindings of
PcG proteins Pc, dRING, and Psc in a significantly high amount as a sharp peak at
putative *ey*-PRE further strengthens the idea. The ~800 bp internal core region also
contained a DHS. With such strong indications for the presence of a functional PRE
at the *ey* locus, we decided to test both *ey*-PRE and a smaller fragment of core DHS
region, *ey*-dPRE, for PRE activity (Figure 4b, Supplementary Table 1).

ey-PRE functions as a PRE

To determine the PRE potential of *ey*-PRE, we assessed the repressor activity of
full-length *ey*-PRE (~1.6kb) and *ey*-dPRE (831 bp) in the adult eye using a
previously described pCasper vector-based transgene assay (Figure 5a). In this
assay, the presence of a PRE upstream of the *mini-white* gene causes its repression
resulting in reduced and variegating eye colour (VASANTHI *et al.* 2013). We observed
both repression and variegation of *mini-white* in the *ey*-PREs transgenic lines. Of
the *ey*-PRE transgenic lines, 52% (16 out of 31) lines showed lighter eye colour, and
55% (18 out of 31) lines showed variegation. In the case of *ey*-dPRE, 50% (13 out of
26) lines displayed lighter eye colour while 21% (4 out of 19) lines showed
variegation (Figure 5b, Table 1, and Supplementary Table 4). Further, to confirm that
the repressive activity of *ey*-PRE and *ey*-dPRE is not a position effect, we excised
out the PREs. Excision of *ey*-PRE/*ey*-dPRE led to the de-repression of *mini-white* in
both sets of transgenic lines (Figure 5b and Table 1). Specifically, 59% and 47%
of *ey*-PRE and *ey*-dPRE lines, respectively, showed an increase in the eye colour
after flipping out of the transgene. Interestingly, we also observed a loss of
variegation in the flipped-out lines of both the transgenes confirming the repressor
function of *ey*-PRE/*ey*-dPRE.

One of the well-known characteristics of PREs is pairing-sensitive silencing (PSS),
where in homozygous conditions, a PRE transgene results in stronger repression as

compared to the heterozygous lines (KASSIS 1994; KASSIS 2002; KASSIS AND BROWN

2013). In our assay, >30% of the *ey*-PRE/*ey*-dPRE lines, showed a strong PSS

effect in homozygous condition, which is lost in flipped-out lines (Figure 5b, Table 1,

and Supplementary Table 4).

In the next step, to ascertain whether the repressor activity of *ey*-PRE/*ey*-dPRE is

PcG protein-dependent, we brought a few representative *ey*-PREs lines into

mutations background of different PcG proteins by crossing homozygous *ey*-PRE

males to the females carrying PcG mutations (Table 2). We compared the de-

repression of *mini-white* scored as an increase in the eye pigment level in the

progenies from these crosses carrying *ey*-PRE with and without PcG mutation. We

found that mutation in *Psc* gene (*Psc^1* allele) caused a notable de-repression of *mini-*

white in five of *ey*-dPRE lines tested. Mutation alleles

of *Pleiohomeotic (Pho^1)* and *Polycomb* (*Pc^1*) in heterozygous condition exhibited

very mild de-repression of *mini-white*. However, a double PcG mutation

of *Psc^1* and *Pc^1* causes cumulative de-repression effect (Figure 5c, Table 2). We

also tested GAF mutation (*Trl^{R85}*) as we observed its binding at *ey*-PRE region

(Figure 4), and interestingly loss of GAF caused a notable de-repression (Figure 5c-

d and Table 2). These observations suggest that while the function of *ey*-PRE is

dependent on PcG genes as in the case of typical PREs, it is possible that factors

other than PcG proteins, like GAF, are also involved in its repressive functions. Our

observation is supported by earlier reports where GAF has been shown to interact

with a *Drosophila* PRE (MISHRA *et al.* 2001).

ME interacts with EB and ey-PRE

As the two CBEs, ME and EB sharply demarcate *ey* locus, we wanted to investigate

whether these regions interact in a long-range to mark independent chromatin

domain of *ey* regulation. We performed 3C in the 0-16h embryo. Using all forward

primers to PCR amplify the ligated hybrids (3C DNA) we calculated the interaction

frequency of ME with several regions at *ey* locus by gel quantification method as

described in Naumova et al.(NAUMOVA *et al.* 2012). Additionally, hybrid template

controls were generated to assess the forward primer pair efficiency. The fragment

comprising ME boundary showed a high interaction frequency with *ey*-PRE and a

lesser but significant interaction frequency with EB and an internal region
of *ey* (Figure 6a-b). However, we did not detect any such interaction of ME with
immediate upstream *myo* gene, or further downstream *bt* gene.

To understand the *ey* domain organization even better and to validate our 3C results,
we explored these interactions in previously published Hi-C data from the whole
embryo and a late embryo-derived Kc167 cell, using an online web tool juicebox.js
(SEXTON *et al.* 2012; LI *et al.* 2015; CUBENAS-POTTS *et al.* 2017; EAGEN *et al.* 2017;
ROBINSON *et al.* 2018). A visual inspection of the fourth chromosome Hi-C heatmap
from the embryo (SEXTON *et al.* 2012) revealed that *ey* and *bt* genes exist in a TAD.
While *myo*, present in a distinct domain upstream region to *ey*, is separated by ME
boundary (Supplementary Figure 3). We next looked into the details of chromatin
looping within the *ey* TAD, in high-resolution Hi-C data available for Kc167 cells
(CUBENAS-POTTS *et al.* 2017; EAGEN *et al.* 2017). In these Hi-C data, *ey* and *bt* are
found to reside in an independent sub-domain within the same TAD. However, *myo*
and *MED26* are part of separate domains (Figure 6b and Supplementary Figure 3).
The ME boundary appears to be in contact with EB boundary (Figure 6b). We
compared the expression profile of these genes in Kc167 cells to further understand
the functional relevance of domain organization at *ey* locus (CHERBAS *et al.* 2011).
Interestingly, we observed that *ey* and *bt* are expressed at a very low level
suggesting their co-regulation in a transcriptionally inactive domain. In contrast, *myo*
and *MED26* show a higher level of expression (Supplementary Figure 3). We also
used the data available at 4Dgenome (https://4dgenome.research.chop.edu/), which
predicts enhancer-promoter interactions, to look into the interactions within *ey* region
(HE *et al.* 2014). This data also indicates the interaction of ME to *ey*-PRE and
downstream *ey* regions, similar to our observation (Figure 6b). These observations
together suggest the interaction of ME boundary with *ey*-PRE. Also, the interaction of
ME with EB in the embryo and Kc167 cells reflects the role of the two CBEs in
defining a functional chromatin domain.

ME, EB and ey-PRE associate differently with nuclear matrix

Nuclear Matrix (NuMat) has been proposed to provide a structural framework for
targeted tethering of chromatin domain in sub-nuclear space (MISHRA AND KARCH

1999). DNA that associates with NuMat (MARs) are proposed to be the anchoring

sequence. CBEs have been shown to associate with each other and along with

boundary proteins are shown to associate NuMat (BLANTON *et al.* 2003; BYRD AND

CORCES 2003; PATHAK *et al.* 2007). As we have observed interaction amongst ME,

EB, and *ey*-PRE, we further investigated their NuMat association potential using

an *in vivo* MAR assay (MIRKOVITCH *et al.* 1984) (Supplementary Figure 4). In the

assay, an equal amount of PCR amplified test regions was Southern hybridized with

radio-labelled NuMat DNA isolated from *Drosophila* embryos. NuMat association is

determined by comparative quantification of hybridization signal of test regions with

known MAR region in Histone gene (HIS-MAR) and a non-MAR region in exonic

sequence of the BEAF-32 gene (BEAF-CDS) (MIRKOVITCH *et al.* 1984; PATHAK *et al.*

2007).

We found that ME and EB both show a comparable level of association with NuMat.

Interestingly, the *ey*-dPRE and the (EB-u region) 3'-UTR of *ey* gene also show a

strong association with NuMat. However, the INT7 region does not show any

significant association with NuMat (Figure 7). During embryonic development,

ey expresses in limited number of cells, i.e., eye-antennal disc primordium, and

embryonic brain cells (HAUCK *et al.* 1999). Reasoning that we might be losing

important information as *ey* does not express in over-whelming majority of embryonic

cells, we also tested the NuMat association of the locus in a different tissue with

significant *ey* transcriptional activity, i.e., brain-eye-antennal imaginal discs of third

instar larvae. In third instar larvae, *ey* mainly expresses in CNS and anterior to the

morphogenetic furrow in the eye disc. As the other extreme, we also tested NuMat

association of *ey* locus in S2 cells where the gene is not transcribed (CZERNY *et al.*

1999; HAUCK *et al.* 1999; ADACHI *et al.* 2003; CHINTAPALLI *et al.* 2007). The NuMat

associations of ME, *ey*-PRE, and EB varied in the embryo, brain-eye-antennal disc,

and S2 cells. Interestingly, *ey*-PRE was found to associate with NuMat in all the

tissues examined, irrespective of *ey* gene transcription status. ME and EB

boundaries were found to associate with NuMat in embryos and brain-eye disc, while

the association was completely absent in S2 cells. EB-u association to NuMat is

stronger than EB-i, indicating that while EB-i harbours boundary activity, EB-u is

responsible for tethering of the locus to nuclear architecture (Figure 7). The

differential association of *ey cis*-regulatory elements to NuMat indicates that *ey* locus

may be organized differently in different cell types depending on the transcriptional
activity of the locus. While these observations provide an initial clue to the 3D
organization of the locus, detailed studies will be needed to fully understand the role
of MAR sequences in dynamics of the region.
440
441 **Discussion**
442
The *ey* gene in *Drosophila*, a gene crucial for eye development, is differentially
expressed in comparison to its neighbouring genes, *myo,* and *bt*. Adjacent
genes, *myo,* which is upstream to *ey,* and *bt,* which is downstream, are very close
to *ey*, ~1.6 kb, and ~3.2 kb apart, respectively. Genetic studies have mainly revealed
the interaction of *ey* to other genes involved in eye development network (DESPLAN
1997; HALDER *et al.* 1998; PICHAUD AND DESPLAN 2001), but transcriptional
regulation *of ey* is not fully understood. Earlier, we have identified ME boundary in
the *myo* and *ey* intergenic region (SULTANA *et al.* 2011). Based on the distinct
expression pattern of the three genes, we had proposed that additional *cis*-regulatory
elements like CBEs and PREs might be present at this locus. In order to identify
such elements and to understand their potential role in *ey* regulation, we used
bioinformatics prediction tools and clusters of binding sites of known CBEs and
PREs binding proteins from genome-wide localization studies. We further co-
mapped the region to DHSs that are often indicative of the presence of regulatory
elements. Taking clues from *in silico* prediction, we have identified and functionally
characterized a novel CBE and a PRE associated with *ey* locus. We show that ~1.2
kb intergenic region between *ey* and *bt*, EB, functions as boundary and ~1.6 kb
region, upstream to *ey* promoter, *ey*-PRE, functions as PcG dependent repressor.
461
Our *in silico* analysis estimates three putative CBEs in *ey* locus, i.e., ME, INT7 and
EB. Of these, ME boundary has already been characterized earlier. In the present
work, we report that only EB displays a resolute boundary activity during embryonic
development as well as a mild boundary activity in the adult stages. Whereas, INT7
shows a scant boundary activity only during early embryogenesis suggesting
boundary activity of these elements could be developmentally regulated. A few
recent studies have suggested a developmental and tissue-specific regulation of
CBEs mediated by spatially and temporally controlled expression of boundary

proteins (AOKI *et al.* 2008; LIN *et al.* 2011; AOKI *et al.* 2012; MATZAT *et al.* 2012;
BONCHUK *et al.* 2015; WOLLE *et al.* 2015; GAMBETTA AND FURLONG 2018). For
example, expression of boundary factors, Elba and a late boundary complex (LBC),
mediate boundary functions of Fab7 element in early embryogenesis and adults,
respectively (AOKI *et al.* 2008; AOKI *et al.* 2012; WOLLE *et al.* 2015). In another
example, a tissue-specific protein Shep has been shown to
modulate *gypsy* boundary activity and nuclear localization, particularly in CNS
(MATZAT *et al.* 2012). Our study of *ey* locus presents a similar scenario, where the
boundaries are spatially and temporally regulated depending on the transcriptional
activity of the locus. It would be further interesting to see if ME, EB and INT7
functions are developmentally regulated or restricted to specific cell-types.

Previous studies have shown that interaction of BEAF-32 with CP190 mediates long-
range chromosomal contact, and more than 70% of BEAF-32/CP190 bindings
demarcate domain boundaries in the *Drosophila* genome (VOGELMANN *et al.* 2014;
CUBENAS-POTTS *et al.* 2017; WANG *et al.* 2018). The EB element also harbours *in*
vivo binding sites of BEAF-32, CP190, and CTCF co-mapping with a prominent
DHS. This overall architecture of the EB element shows a remarkable similarity to
the earlier identified ME (SULTANA *et al.* 2011). These evidences suggest that the
same molecular players, particularly BEAF-32 and CP190, might be contributing to
the functional linking of the two CBEs to define the loop domain of *ey*. In our 3C
experiment, we confirm the interaction between ME and EB suggesting that the two
CBEs, flanking the *ey* locus, define functional autonomy of the chromatin loop
domain. In support to our finding, we observe the chromatin loop of ME with EB in
HiC data from previous studies in Kc167 cells which places *ey* in an independent
domain within a larger TAD (LI *et al.* 2015; CUBENAS-POTTS *et al.* 2017; EAGEN *et al.*
2017). These interactions provide necessary insulation to *ey* locus from the
neighbouring regulatory environments.

We report here the identification and functional validation of a new CMM or PRE, *ey*-
PRE, within the *ey* domain. Besides conferring a repressive feature in the transgene-
based assay, the element also exhibits PSS. The *ey*-PRE element encompasses a
core DHS region spanning ~830 bp (*ey*-dPRE) that coincides with sharp binding
peaks of PcG proteins of PRC1 members, Pc, Psc, dRing. However, in the

transgene, we could only see a prominent de-repressive effect of Psc mutation.
Other PcG proteins show a moderate and varying impact on *ey*-PRE activity, which
could be because these mutations were tested in heterozygous condition, as
homozygous mutants are lethal. In case of double PcG mutants however we could
see strong de-repression effect which suggests that *ey*-PRE maintains the
expression state of the *ey* locus and requires the activity of a subset of PcG proteins
for the purpose. This property of *ey*-PRE is similar to the earlier reported PREs
(KASSIS 2002; KASSIS AND BROWN 2013). Interestingly, GAF protein was also found to
attenuate the function of *ey*-PRE. ME boundary also has been reported to interact
with GAF protein (SULTANA *et al.* 2011). The observed 3C interaction of ME and *ey*-
PRE could be mediated by GAF protein. Recent HiC studies also support the
interaction of PREs with multiple regions within a TAD and active role of GAF
proteins in such connections (OGIYAMA *et al.* 2018). The *ey*-PRE also has predicted
promoter property, which suggest it to have a dual role of promoter as well as CMM.
However, the role of GAF remains unexplored and it will be interesting to see if GAF
is crucial to these functions.

Studies have shown that NuMat provides anchoring sites for the
compartmentalization of higher-order chromatin organization and also acts as a
platform for functional activities inside the nucleus. Work from our lab and elsewhere
have shown that boundary elements function in the context of NuMat (BYRD AND
CORCES 2003; PATHAK *et al.* 2007). In this study, we show that multiple sites of
the *ey* locus, including ME, *ey*-PRE, and EB, associate with NuMat in cell type and
transcription status dependent manner. As BEAF-32 is a bona fide NuMat
associated protein in flies, and ME and EB, both carry BEAF-32 binding sites, the
anchoring of these CBEs to the nuclear architecture is probably mediated by the
protein (PATHAK *et al.* 2007). The prominent association of ey-PRE to NuMat in all
cell types, unlike other elements, suggests the role of the memory element in all
functional states or cell types. Taken together, the dynamic nature of NuMat
association of different regulatory elements of the locus supports the hypothesis that
depending on the expression state, *ey* locus is present in separate compartments
and that NuMat might be providing the structural basis for this compartmentalization.

Based on these findings, we propose a model for *ey* gene expression where *ey*-PRE

functions as both PRE/TRE to maintain the active or repress state of *ey* in different

cell/tissues initially established by early developmental cues. The CBEs, ME and EB,

demarcate the *ey* locus, to create an independent domain of *ey* differential regulation

from the neighbouring genes *myo* and *bt,* by chromatin looping and thus prevent the

cross talk among the regulatory elements of the adjacent genes. The *ey*-PRE,

perhaps with the other unidentified regulatory elements of *ey* locus, which are bound

to PcG or TrxG proteins, takes the *ey* domain either to repressive (*Polycomb* body)

or active (transcription factory) compartments, respectively (Figure 8).

Conclusions

In conclusion, we have identified boundary elements, ME and EB, that demarcate

the *ey* domain and a cellular memory element, *ey*-PRE, within this domain. In order

to get insight into the mechanism of *ey* regulation, we have investigated how these

regulatory elements function together in their genomic context. We have learnt that

long-range interactions of ME with *ey*-PRE and EB, along with their discrete

association to the NuMat brings about the tissue-specific regulation of *ey*.

As *ey* expresses only in a discrete group of cells, while in the majority of cells in the

embryo or a tissue it remains repressed, it is challenging to compare higher-order

chromatin organizational changes at the locus during embryonic development.

However, one way to unearth such mechanisms is to carry out super-resolution

imaging of the locus in a cell-type-specific manner. Further, identification of

molecular players involved in the long-range interactions and the NuMat association

of these regulatory elements will also provide an insight into the regulation of the

locus. At present, this study offers a working model for the underlying mechanism

of *ey* spatiotemporal regulation in *Drosophila*. Our study provides clues to the

hierarchical genomic conformational features, that form the basis of gene regulation

in euchromatin.

Methods

CBEs and PRE prediction in ey locus

A ~ 30 kb *ey* region was used in cdBEST analysis to identify new CBEs at *ey* locus.
Additionally, for prediction of the CBEs, *in-vivo* binding of known boundary proteins
BEAF32, CP190, CTCF, and GAF was analyzed using whole embryo (0-12 h) ChIP-
chip data from a previous by study Negre et al., 2010(NEGRE *et al.* 2010). We co-
mapped the DNaseI hypersensitive sites(DHS) in the region using stage-9 (~ 6 h)
embryo DHS data available at UCSC browser (https://genome.ucsc.edu/) (THOMAS
et al. 2011). Two potential CBEs, a ~ 1.3 kb intronic region in a seventh intron (INT7)
and a ~ 2.2 kb intergenic region between *ey* and *bt* (EB), were identified that also co-
map with DHS (Figure 1).

A Perl based program, PRE mapper, was used to predict putative PRE (*ey*-PRE) in
the *ey* locus (SRINIVASAN AND MISHRA 2020). PRE mapper analyzes motifs of known
PRE binding proteins (Pho, GAF, Dsp1, Sp1, Zeste, Grh, Adf1, and Cg), their
binding patterns, and motif clustering to define a PRE. Presence of DHS at ey-PRE
was reviewed to correlate its regulatory feature. Additionally, a comprehensive
analysis of various chromatin features in the *ey* region was carried out using 14-16-h
whole embryo ChIP-seq data available at modENCODE by Gary Karpen.
modENCODE id of the data used are H3K27Ac (4120), H3K27Me3 (3955), Pc
(3957), dRING (5071), Psc (3960), BEAF-32 (3954), CP190 (3959), CTCF (5069),
and GAF (4149) (MOD *et al.* 2010).

Cloning of predicted CBEs and PRE

All the test fragments were PCR amplified from *CantonS* fly genomic DNA by high
fidelity *Pfu* polymerase with specific primers (Supplementary Figure 1,
Supplementary Table 1 and 5). All CBE test fragments (INT7, EB, EB-u and EB-i)
were inserted first at MCS in pLML and then sub-cloned along with *loxP* at XhoI sites
in the enhancer-blocker assay vectors, pRW+ and pCfhL, (SULTANA *et al.* 2011). Test
PREs (*ey*-PRE and *ey*-dPRE) were cloned initially at SmaI blunt site in pLML, then
sub-cloned along with *loxP* at XhoI in pCasPerX (VASANTHI *et al.* 2013).

Fly culture, transgenic flies, and genetic crosses

All fly strains of *Drosophila melanogaster* were cultured in standard cornmeal

605 medium at 25 ^{0}C. Transgenic flies were generated by micro-injection of an assay

606 vector DNA and a transposase expressing plasmid *pΔ2-3* together into pre-

607 blastoderm of w^{1118} as described in Siegal et al. (SIEGAL AND HARTL 2000). Adult flies

608 from injected embryos were backcrossed to w^{1118}, and the transgenes were

609 identified by the presence of eye pigment in the adult progenies. Transgenic flies

610 were next crossed with the double balancer (*Pin/CyO; TM2/TM6*) to know

611 chromosomes of P-element integration and making homozygous stock. We

612 established several lines for each predicted CBE and PRE (Supplementary Table 2-

613 4). The test fragment was flipped out from initial transgene, to compare the effect of

614 test fragment and rule out any position effect, by crossing the flies with *Cre-*

615 *recombinase* expressing flies and later was confirmed by PCR using specific

616 primers. The homozygous *ey*-dPRE males were crossed to virgin females of different

617 PcG proteins mutant (*Pc^1, Psc^1, Pho^1*) to investigate the effect on repressor function

618 of *ey*-dPRE (listed in Supplementary Table 6). The effect of PcG mutations effect

619 were compared in progenies with or without mutations in heterozygous condition.

621 ***Visualization and quantitative measurement of eye pigment***

623 The level of pigment in the eye of similar aged (5-day) adults were visually

624 determined to compare the level of *mini-white* expression in the enhancer-blocker

625 and PRE assay transgenic flies. In transgenic flies, pigmentation range varied

626 from bright/dark red in wild-type (*CS*) toward red, dark orange/light red, yellow,

627 and light-yellow, to white in the absence of expression (Supplementary Table 2-4).

628 We also quantitated the red pigment value in eye for some transgenic flies, to

629 compare the strength and effect of boundary and PRE after flipping-out (SULTANA *et*

630 *al.* 2011). Eye pigments from a minimum of ten adult heads were extracted in a 1:1

631 mixture of chloroform/0.1% ammonium hydroxide (200 μl) then, and centrifuged the

632 mix for 2 min at 13000 rpm. Finally, 100 μl of the supernatant was taken for

633 quantification of red pigment by spectrophotometric absorbance at 485 nm. A mean

634 relative pigmentation and standard error were calculated from a minimum of three

635 independent pigment values for each line by comparing it to the pigment value

636 of w^{1118}.

638 *β-galactosidase staining and quantification*

For enhancer blocker assay in embryo, X-gal staining was performed to determine

the beta-galactosidase (lacZ) activity (BELLEN et al. 1989; SULTANA et al. 2011). In

brief, 0-12-h old embryos were collected and dechorionated in 50% bleach (sodium

hypochlorite). The dechorionated embryos were washed and fixed for 20 min in

heptane (saturated with 25% glutaraldehyde). The fixed embryos were again

washed thoroughly with 1x PBST (0.3% Triton X-100), incubated initially for 10 min

at 37^0C in X-gal staining solution (3 mM $K_3[Fe(CN)_6]$, 3 mM $K_4[Fe(CN)_6]$, 7.2 mM

Na_2HPO_4, 2.8 mM NaH_2PO_4, 1 mM $MgCl_2$, 150 mM NaCl), and finally stained for 4-6

h in X-gal staining solution with 0.2% X-gal (Sigma). The stained embryos were

imaged in Leica stereomicroscope. We performed X-gal staining of multiple

transgenic lines along with positive and negative controls in a common grid, ensuring

reliable comparison of staining between different genotypes.

To estimate and compare the enhancer blocker activity of the test fragments in the

embryos, we quantitated X-gal staining using ImageJ-image processing and analysis

software provided by the National Institutes of Health, USA (HARTIG 2013). We used

pixels value of a fixed area size of different regions from multiple stained embryos of

each genotype to calculate relative X-gal staining and standard error and plotted as

a bar graph for comparison.

3C Assay

3C experiment was carried out in wild type *CS* embryos as described earlier

(DEKKER et al. 2002; COMET et al. 2011) with few modifications. To begin with, 0-16 h

old embryos (1 g) were dechorionated in 50% bleach for 5 min. The dechorionated

embryos were washed thoroughly PBST (PBS 1x + 0.01% Triton X 100) and fixed

for 15-20 min at room temperature (RT) with 2% formaldehyde (Sigma) in 5 ml fixing

solution (50 mM HEPES-pH-7.6, 100 mM NaCl, 0.1 mM Na-EDTA-pH-8, 0.5 mM Na-

EGTA-pH-8) added with 15 ml heptane. To stop fixation, formaldehyde was

quenched by 0.125 M glycine for 5 min at RT. The fixed embryos were resuspended

in 2.5 ml of ice-cold lysis buffer (10 mM Tris-pH-8, 10 mM NaCl, 0.2% NP40 with

Roche protease inhibitor cocktail freshly added) and homogenized in Dounce

homogenizer (10-15 strokes) to create nuclei suspension. Cellular debris was

removed from the nuclei suspension by filtration through a double layer of Mira cloth.
Later, the nuclei were washed once with 500µl 1.2xNEB EcoRI buffer then
resuspended in 500µl 1.2x NEB EcoRI buffer with 0.01% Triton X-100. A 50µl of
nuclear solution (~100mg embryos) was lysed in 362µl 1.2x NEB EcoRI buffer with
1% SDS for 2 h at 37^0C. To stop the lysis triton X-100 was added to 0.01% and
incubated for another 2 h at 37^0C. Subsequently, EcoRI digestion was carried out
with 400 units of EcoRI (NEB) for 2h at 37^0C then inactivated the EcoRI at 65^0C for
20 min by adding SDS to a final concentration of 1.5 %. Afterwards, 1% Triton X-100
was added to neutralize the SDS at 37 ^{0}C for 1 h. EcoRI digested samples were
diluted ten times in 1x NEB ligation buffer and ligated with 8µl ligase (400 U/*µl*, NEB)
for 3-4 h at 22 ^{0}C. The ligated (3C) DNA was sequentially treated with RNaseA (20
µg/ml) for 30 minutes at 37 ^{0}C and with 100 µg/ml Proteinase-K at 55 ^{0}C for 1 h, and
subsequently de-crosslinked overnight at 65 ^{0}C. Finally, 3C DNA was extracted by
phenol: chloroform: isoamyl alcohol (25:24:1) and ethanol precipitation method. DNA
from nuclei suspension and EcoRI digested samples were also prepared to compare
the digestion and ligation efficiency.

The purified 3C DNA was used for PCR to check the interaction of different regions
using all forward primers. Since we did not have a BAC clone available for *ey* region,
we used forward-reverse primer pair to amplify regions flanking to selected
restriction sites. These regions were mixed in an equimolar ratio, EcoRI restricted
digested, and ligated. The randomly ligated product used to check the efficiency of
the forward primer pairs. All PCRs amplifications were carried out with the following
parameters: 95 ^{0}C for 3min, followed by 36 cycles of 95 ^{0}C- 30 s, 56 ^{0}C-10 s, and 72
^{0}C-8 s, with a final step at 72 ^{0}C-2min. All forward primer pairs having similar
amplification efficiency were next used for PCR amplification of 3C DNA. The 3C
ligated PCR products were resolved on 1.2% agarose gel, and we confirmed the
chimeric sequences by sequencing of gel extracted PCR DNA. The signal intensity
of the 3C PCR product in the gel was quantified using ImageJ. Later, interaction
frequency and standard deviations for each forward primer pairs were calculated
from three replicates as described in Naumova et al. (NAUMOVA *et al.* 2012).

Analysis of HiC data from previous studies

To understand *ey* locus organization, we used the HiC data of embryos from Sexton
et al. (SEXTON *et al.* 2012), in Kc167 cells from Eagen et al. (EAGEN *et al.* 2017), Li et
al., and Cubenas et al. (LI *et al.* 2015; CUBENAS-POTTS *et al.* 2017). The processed
HiC matrix data were used in two online visualization tools, i.e., Juicebox (ROBINSON
et al. 2018) and HiCExplorer (WOLFF *et al.* 2018), to find out chromatin interactions.

Nuclear matrix Association Assay

The nuclear matrix (NuMat) association of *ey cis*-elements were tested in a modified
in vivo MAR assay from the original protocol of Mirkovitch, *et al.* 1984 (MIRKOVITCH *et*
al. 1984; PATHAK *et al.* 2014) (Supplementary Figure 3).

First, NuMat DNA was prepared from 0-16 h old embryos, eye-antennal discs (3rd
instar larvae), and S2 cells. In brief, nuclei were isolated from 1 g of embryos and ~
200 pairs of eye-antennal imaginal discs and 2×10^6 S2 cells. An aliquot of nuclei was
used for the isolation and estimation of total genomic DNA for quality control checks.
Isolated nuclei were treated with DNaseI in nuclei isolation buffer (20 mM Tris-pH-
7.4, 20 mM KCl, 70 mM NaCl, 10 mM $MgCl_2$, 0.12 5mM spermidine 1mM PMSF,
0.5% Triton-X 100, and 200µg/ml DNaseI) at 4 ^{0}C for 1 h to remove chromatin.
Chromatin depleted nuclei were collected by centrifugation at 3000 x g for 10
minutes. Non-matrix proteins were extracted out from the nuclei with 0.4 M NaCl for
5 min in extraction buffer (10 mM Hepes-pH-7.5, 4 mM EDTA, 0.25 mM spermidine,
0.1 mM PMSF, 0.5% (v/v) triton X-100) and another 5 min with 2 M NaCl in the
extraction buffer to get NuMat. The NuMat pellet was washed twice with wash buffer
(1 mM Tris-pH-7.4, 20 mM KCl, 1 mM EDTA, 0.25 mM spermidine, 0.1 mM PMSF)
then treated with RNaseA (20 µg/ml) for 30 minutes at 37 ^{0}C and with 100 µg/ml
Proteinase-K at 55 ^{0}C for 1 h. Finally, NuMat DNA was obtained by
phenol:chloroform:isoamylalcohol (25:24:1) and ethanol precipitation method. NuMat
DNA was dissolved in DNase free water and quantified using Nano-drop. An equal
amount of NuMat DNA isolated from different tissues was labeled with 32^P-dATP by
the Random Primer Labeling method.

Next, all the test regions were PCR amplified from fly genomic DNA using specific
primers listed in Supplementary Table 5. A known histone gene matrix-associated

region (HISMAR) was used as a positive control and an exonic sequence of BEAF-32 gene (BEAF CDS) was used as a random control. The controls were used to compare the relative NuMat associations. An equal amount (~ 400 ng) of all PCR products were resolved first on a 1.2% TAE agarose gel and then transferred to a positively charged nylon membrane using a capillary transfer method. The membrane was then subjected to a standard Southern hybridization using ^{32}P-dATP labeled NuMat DNA as described earlier (PATHAK *et al.* 2014). Finally, the hybridization signals were observed in Phosphor Molecular Imager (PMI) from (BIORAD). Relative association to NuMat and standard deviation for each region was calculated by signal intensity from three replicate blots using ImageJ (HARTIG 2013).

Competing interests

The authors declare that they have no competing interests.

Funding

Council of Scientific and Industrial Research (CSIR), India has funded this work, through JRF/SRF/RA fellowships to SV and research grants to RKM.

CRediT author contribution statement

Shreekant Verma: Investigation, Methodology, Visualization, Writing – original draft. **Rashmi U Pathak**: Investigation, Writing – review and editing. **Rakesh K Mishra**: Resources, Visualization, Supervision, Funding acquisition, Writing – review and editing.

Acknowledgements

We thank Welcome Bender for his discussion and sharing of Pc binding data in *ey* locus (not shown). We thank Yaser Payferman for his help in cloning for *ey*-PRE and A. Srinivasan for his helpful inputs in PRE predictions.

References

Adachi, Y., B. Hauck, J. Clements, H. Kawauchi, M. Kurusu *et al.*, 2003 Conserved cis-
regulatory modules mediate complex neural expression patterns of the eyeless gene
in the Drosophila brain. Mech Dev 120: 1113-1126.
Ahanger, S. H., A. Srinivasan, D. Vasanthi, Y. S. Shouche and R. K. Mishra, 2013 Conserved
boundary elements from the Hox complex of mosquito, Anopheles gambiae. Nucleic
Acids Res 41: 804-816.
Americo, J., M. Whiteley, J. L. Brown, M. Fujioka, J. B. Jaynes *et al.*, 2002 A complex array of
DNA-binding proteins required for pairing-sensitive silencing by a polycomb group
response element from the Drosophila engrailed gene. Genetics 160: 1561-1571.
Aoki, T., A. Sarkeshik, J. Yates and P. Schedl, 2012 Elba, a novel developmentally regulated
chromatin boundary factor is a hetero-tripartite DNA binding complex. Elife 1:
e00171.
Aoki, T., S. Schweinsberg, J. Manasson and P. Schedl, 2008 A stage-specific factor confers
Fab-7 boundary activity during early embryogenesis in Drosophila. Mol Cell Biol 28:
1047-1060.
Ayme-Southgate, A., J. Vigoreaux, G. Benian and M. L. Pardue, 1991 Drosophila has a
twitchin/titin-related gene that appears to encode projectin. Proc Natl Acad Sci U S A
88: 7973-7977.
Bantignies, F., C. Grimaud, S. Lavrov, M. Gabut and G. Cavalli, 2003 Inheritance of Polycomb-
dependent chromosomal interactions in Drosophila. Genes Dev 17: 2406-2420.
Bellen, H. J., C. J. O'Kane, C. Wilson, U. Grossniklaus, R. K. Pearson *et al.*, 1989 P-element-
mediated enhancer detection: a versatile method to study development in
Drosophila. Genes Dev 3: 1288-1300.
Blanton, J., M. Gaszner and P. Schedl, 2003 Protein:protein interactions and the pairing of
boundary elements in vivo. Genes Dev 17: 664-675.
Bonchuk, A., O. Maksimenko, O. Kyrchanova, T. Ivlieva, V. Mogila *et al.*, 2015 Functional role
of dimerization and CP190 interacting domains of CTCF protein in Drosophila
melanogaster. BMC Biol 13: 63.
Byrd, K., and V. G. Corces, 2003 Visualization of chromatin domains created by the gypsy
insulator of Drosophila. J Cell Biol 162: 565-574.
Cavalli, G., and R. Paro, 1998 The Drosophila Fab-7 chromosomal element conveys
epigenetic inheritance during mitosis and meiosis. Cell 93: 505-518.
Cherbas, L., A. Willingham, D. Zhang, L. Yang, Y. Zou *et al.*, 2011 The transcriptional diversity
of 25 Drosophila cell lines. Genome Res 21: 301-314.
Chintapalli, V. R., J. Wang and J. A. Dow, 2007 Using FlyAtlas to identify better Drosophila
melanogaster models of human disease. Nat Genet 39: 715-720.
Comet, I., B. Schuettengruber, T. Sexton and G. Cavalli, 2011 A chromatin insulator driving
three-dimensional Polycomb response element (PRE) contacts and Polycomb
association with the chromatin fiber. Proc Natl Acad Sci U S A 108: 2294-2299.
Cubenas-Potts, C., M. J. Rowley, X. Lyu, G. Li, E. P. Lei *et al.*, 2017 Different enhancer classes
in Drosophila bind distinct architectural proteins and mediate unique chromatin
interactions and 3D architecture. Nucleic Acids Res 45: 1714-1730.
Czerny, T., G. Halder, U. Kloter, A. Souabni, W. J. Gehring *et al.*, 1999 twin of eyeless, a
second Pax-6 gene of Drosophila, acts upstream of eyeless in the control of eye
development. Mol Cell 3: 297-307.
Dekker, J., K. Rippe, M. Dekker and N. Kleckner, 2002 Capturing chromosome conformation.
Science 295: 1306-1311.

Desplan, C., 1997 Eye development: governed by a dictator or a junta? Cell 91**:** 861-864.

Dreos, R., G. Ambrosini, R. C. Perier and P. Bucher, 2015 The Eukaryotic Promoter Database:
expansion of EPDnew and new promoter analysis tools. Nucleic Acids Res 43**:** D92-
96.

Eagen, K. P., E. L. Aiden and R. D. Kornberg, 2017 Polycomb-mediated chromatin loops
revealed by a subkilobase-resolution chromatin interaction map. Proc Natl Acad Sci
U S A 114**:** 8764-8769.

Frise, E., A. S. Hammonds and S. E. Celniker, 2010 Systematic image-driven analysis of the
spatial Drosophila embryonic expression landscape. Mol Syst Biol 6**:** 345.

Fyrberg, C. C., S. Labeit, B. Bullard, K. Leonard and E. Fyrberg, 1992 Drosophila projectin:
relatedness to titin and twitchin and correlation with lethal(4) 102 CDa and bent-
dominant mutants. Proc Biol Sci 249**:** 33-40.

Gambetta, M. C., and E. E. M. Furlong, 2018 The Insulator Protein CTCF Is Required for
Correct Hox Gene Expression, but Not for Embryonic Development in Drosophila.
Genetics 210: 129-136.

Gaszner, M., and G. Felsenfeld, 2006 Insulators: exploiting transcriptional and epigenetic
mechanisms. Nat Rev Genet 7**:** 703-713.

Gerasimova, T. I., and V. G. Corces, 2001 Chromatin insulators and boundaries: effects on
transcription and nuclear organization. Annu Rev Genet 35**:** 193-208.

Geyer, P. K., C. Spana and V. G. Corces, 1986 On the molecular mechanism of gypsy-induced
mutations at the yellow locus of Drosophila melanogaster. EMBO J 5**:** 2657-2662.

Hagstrom, K., M. Muller and P. Schedl, 1996 Fab-7 functions as a chromatin domain
boundary to ensure proper segment specification by the Drosophila bithorax
complex. Genes Dev 10: 3202-3215.

Halder, G., P. Callaerts, S. Flister, U. Walldorf, U. Kloter *et al.*, 1998 Eyeless initiates the
expression of both sine oculis and eyes absent during Drosophila compound eye
development. Development 125**:** 2181-2191.

Hartig, S. M., 2013 Basic image analysis and manipulation in ImageJ. Curr Protoc Mol Biol
Chapter 14: Unit14 15.

Hauck, B., W. J. Gehring and U. Walldorf, 1999 Functional analysis of an eye specific
enhancer of the eyeless gene in Drosophila. Proc Natl Acad Sci U S A 96**:** 564-569.

He, B., C. Chen, L. Teng and K. Tan, 2014 Global view of enhancer-promoter interactome in
human cells. Proc Natl Acad Sci U S A 111**:** E2191-2199.

Hou, C., L. Li, Z. S. Qin and V. G. Corces, 2012 Gene density, transcription, and insulators
contribute to the partition of the Drosophila genome into physical domains. Mol Cell
48**:** 471-484.

Kassis, J. A., 1994 Unusual properties of regulatory DNA from the Drosophila engrailed gene:
three "pairing-sensitive" sites within a 1.6-kb region. Genetics 136**:** 1025-1038.

Kassis, J. A., 2002 Pairing-sensitive silencing, polycomb group response elements, and
transposon homing in Drosophila. Adv Genet 46**:** 421-438.

Kassis, J. A., and J. L. Brown, 2013 Polycomb group response elements in Drosophila and
vertebrates. Adv Genet 81**:** 83-118.

Kellum, R., and P. Schedl, 1991 A position-effect assay for boundaries of higher order
chromosomal domains. Cell 64**:** 941-950.

Lanzuolo, C., V. Roure, J. Dekker, F. Bantignies and V. Orlando, 2007 Polycomb response
elements mediate the formation of chromosome higher-order structures in the
bithorax complex. Nat Cell Biol 9**:** 1167-1174.

Li, L., X. Lyu, C. Hou, N. Takenaka, H. Q. Nguyen *et al.*, 2015 Widespread rearrangement of
3D chromatin organization underlies polycomb-mediated stress-induced silencing.
Mol Cell 58**:** 216-231.
Lin, N., X. Li, K. Cui, I. Chepelev, F. Tie *et al.*, 2011 A barrier-only boundary element delimits
the formation of facultative heterochromatin in Drosophila melanogaster and
vertebrates. Mol Cell Biol 31**:** 2729-2741.
Lo, P. C., and M. Frasch, 1999 Sequence and expression of myoglianin, a novel Drosophila
gene of the TGF-beta superfamily. Mech Dev 86**:** 171-175.
Maeda, R. K., and F. Karch, 2006 The ABC of the BX-C: the bithorax complex explained.
Development 133**:** 1413-1422.
Maroto, M., J. Vinos, R. Marco and M. Cervera, 1992 Autophosphorylating protein kinase
activity in titin-like arthropod projectin. J Mol Biol 224**:** 287-291.
Matzat, L. H., R. K. Dale, N. Moshkovich and E. P. Lei, 2012 Tissue-specific regulation of
chromatin insulator function. PLoS Genet 8**:** e1003069.
Mihaly, J., S. Barges, L. Sipos, R. Maeda, F. Cleard *et al.*, 2006 Dissecting the regulatory
landscape of the Abd-B gene of the bithorax complex. Development 133**:** 2983-2993.
Mirkovitch, J., M. E. Mirault and U. K. Laemmli, 1984 Organization of the higher-order
chromatin loop: specific DNA attachment sites on nuclear scaffold. Cell 39**:** 223-232.
Mishra, R. K., and F. Karch, 1999 Boundaries that demarcate structural and functional
domains of chromatin. Journal of Biosciences 24**:** 377-399.
Mishra, R. K., J. Mihaly, S. Barges, A. Spierer, F. Karch *et al.*, 2001 The iab-7 polycomb
response element maps to a nucleosome-free region of chromatin and requires both
GAGA and pleiohomeotic for silencing activity. Mol Cell Biol 21**:** 1311-1318.
mod, E. C., S. Roy, J. Ernst, P. V. Kharchenko, P. Kheradpour *et al.*, 2010 Identification of
functional elements and regulatory circuits by Drosophila modENCODE. Science 330**:**
1787-1797.
Naumova, N., E. M. Smith, Y. Zhan and J. Dekker, 2012 Analysis of long-range chromatin
interactions using Chromosome Conformation Capture. Methods 58**:** 192-203.
Negre, N., C. D. Brown, P. K. Shah, P. Kheradpour, C. A. Morrison *et al.*, 2010 A
comprehensive map of insulator elements for the Drosophila genome. PLoS Genet 6**:**
e1000814.
Ogiyama, Y., B. Schuettengruber, G. L. Papadopoulos, J. M. Chang and G. Cavalli, 2018
Polycomb-Dependent Chromatin Looping Contributes to Gene Silencing during
Drosophila Development. Mol Cell 71**:** 73-88 e75.
Pathak, R. U., N. Rangaraj, S. Kallappagoudar, K. Mishra and R. K. Mishra, 2007 Boundary
element-associated factor 32B connects chromatin domains to the nuclear matrix.
Mol Cell Biol 27**:** 4796-4806.
Pathak, R. U., A. Srinivasan and R. K. Mishra, 2014 Genome-wide mapping of matrix
attachment regions in Drosophila melanogaster. BMC Genomics 15**:** 1022.
Pichaud, F., and C. Desplan, 2001 A new visualization approach for identifying mutations
that affect differentiation and organization of the Drosophila ommatidia.
Development 128**:** 815-826.
Quiring, R., U. Walldorf, U. Kloter and W. J. Gehring, 1994 Homology of the eyeless gene of
Drosophila to the Small eye gene in mice and Aniridia in humans. Science 265**:** 785-
789.

Robinson, J. T., D. Turner, N. C. Durand, H. Thorvaldsdottir, J. P. Mesirov *et al.*, 2018
Juicebox.js Provides a Cloud-Based Visualization System for Hi-C Data. Cell Syst 6**:**
256-258 e251.
Ronshaugen, M., and M. Levine, 2004 Visualization of trans-homolog enhancer-promoter
interactions at the Abd-B Hox locus in the Drosophila embryo. Dev Cell 7**:** 925-932.
Sexton, T., F. Bantignies and G. Cavalli, 2009 Genomic interactions: chromatin loops and
gene meeting points in transcriptional regulation. Semin Cell Dev Biol 20**:** 849-855.
Sexton, T., E. Yaffe, E. Kenigsberg, F. Bantignies, B. Leblanc *et al.*, 2012 Three-dimensional
folding and functional organization principles of the Drosophila genome. Cell 148**:**
458-472.
Siegal, M. L., and D. L. Hartl, 2000 Application of Cre/loxP in Drosophila. Site-specific
recombination and transgene coplacement. Methods Mol Biol 136**:** 487-495.
Sigrist, C. J., and V. Pirrotta, 1997 Chromatin insulator elements block the silencing of a
target gene by the Drosophila polycomb response element (PRE) but allow trans
interactions between PREs on different chromosomes. Genetics 147**:** 209-221.
Srinivasan, A., and R. K. Mishra, 2012 Chromatin domain boundary element search tool for
Drosophila. Nucleic Acids Res 40**:** 4385-4395.
Srinivasan, A., and R. K. Mishra, 2020 Genomic organization of Polycomb Response Elements
and its functional implication in Drosophila and other insects. J Biosci 45.
Sultana, H., S. Verma and R. K. Mishra, 2011 A BEAF dependent chromatin domain boundary
separates myoglianin and eyeless genes of Drosophila melanogaster. Nucleic Acids
Res 39**:** 3543-3557.
Thomas, S., X. Y. Li, P. J. Sabo, R. Sandstrom, R. E. Thurman *et al.*, 2011 Dynamic
reprogramming of chromatin accessibility during Drosophila embryo development.
Genome Biol 12**:** R43.
Valenzuela, L., and R. T. Kamakaka, 2006 Chromatin insulators. Annu Rev Genet 40**:** 107-138.
Vasanthi, D., A. Nagabhushan, N. K. Matharu and R. K. Mishra, 2013 A functionally
conserved Polycomb response element from mouse HoxD complex responds to
heterochromatin factors. Sci Rep 3**:** 3011.
Vogelmann, J., A. Le Gall, S. Dejardin, F. Allemand, A. Gamot *et al.*, 2014 Chromatin insulator
factors involved in long-range DNA interactions and their role in the folding of the
Drosophila genome. PLoS Genet 10**:** e1004544.
Wang, Q., Q. Sun, D. M. Czajkowsky and Z. Shao, 2018 Sub-kb Hi-C in D. melanogaster
reveals conserved characteristics of TADs between insect and mammalian cells. Nat
Commun 9**:** 188.
West, A. G., M. Gaszner and G. Felsenfeld, 2002 Insulators: many functions, many
mechanisms. Genes Dev 16**:** 271-288.
Wolff, J., V. Bhardwaj, S. Nothjunge, G. Richard, G. Renschler *et al.*, 2018 Galaxy HiCExplorer:
a web server for reproducible Hi-C data analysis, quality control and visualization.
Nucleic Acids Res 46**:** W11-W16.
Wolle, D., F. Cleard, T. Aoki, G. Deshpande, P. Schedl *et al.*, 2015 Functional Requirements
for Fab-7 Boundary Activity in the Bithorax Complex. Mol Cell Biol 35**:** 3739-3752.
Zhang, Y., C. H. Wong, R. Y. Birnbaum, G. Li, R. Favaro *et al.*, 2013 Chromatin connectivity
maps reveal dynamic promoter-enhancer long-range associations. Nature 504**:** 306-
310.
953
954

 Figure legends

Figure 1. CBE prediction at the *ey* locus. (a) (i) *ey* locus with flanking genes *myo* and *bt* in the fourth chromosome in *Drosophila melanogaster* (genome version dm3). **(ii)** Mapping of the boundary proteins binding sites and DHS. Embryonic (0-12 h) ChIP-chip binding data of BEAF-32, CP190, CTCF, and GAF are used from Negre et al. (NEGRE *et al.* 2010). For DHS mapping embryonic stage 9 (~6 h) DHS data from Thomas et al. is used (THOMAS *et al.* 2011). **(iii)** Predicted CBEs. CdBEST predicts previously reported ME between *ey* and *myo* and INT7 in the seventh intronic region of *ey*. Additionally, boundary protein bindings and DHS coincide with INT7(~1.2 Kb) and EB (~2.2 Kb) genomic regions (highlighted in light green). **(b)** Detailed mapping of boundary proteins binding peaks (BEAF-32, CTCF and CP190) and DHS in INT7 and EB region that are tested for enhancer blockers. CTCF binding and DHS at EB present in two sites EB-u and EB-i.

Figure 2. EB functions as Enhancer blocker adult eye (a) Enhancer blocker assay construct, pRW$^+$. The test fragment flanking with *loxP* sites is inserted between the enhancer and promoter of *mini-white*. **(b)** Enhancer blocker assay in the eye. Eye color comparisons of RW$^+$ transgenes. In each panel, the eye on the left contains the test fragment in heterozygous (P/+), whereas the eye on the right is the same line after the fragment is flipped out (ΔP/+). INT7 region does not show enhancer blocker activity while EB shows a weak enhancer blocker activity. EB-u and EB-i, when tested separately, only EB-i shows similar activity to EB suggest the boundary function of EB resides in EB-i. The graph represents the relative pigment value in the eye. An average mean and standard error in bars is calculated from a minimum of 10 eyes in triplicates.

Figure 3. EB functions as an Enhancer blocker in the embryo. (a) Enhancer blocker assay construct, pCfhL. If a test DNA, inserted between the *ftz* enhancers and *hsp70/lacZ* gene, functions as a boundary, it prevents the enhancer from driving the reporter gene, thereby, giving the low intensity of X-gal staining, which restore upon flipping out the test DNA and enhance the staining. **(b**) LacZ/X-gal staining of the embryos. The panel compares the x-gal staining during early development (UPS mediated) of the initial (P/P) and flipped-out (ΔP/ΔP) transgenes of INT7, EB, EB-u,

and EB-i. The pCfhL construct alone and with Fab7 boundary used as negative and
positive controls, respectively. INT7 in shows a weak enhancer blocker effect while
EB shows a significant enhancer blocking activity. Further, the boundary activity of
EB resides in EB-i, whereas EB-u does not show enhancer blocker effect.
(c) Quantification LacZ staining. The relative x-gal staining and standard errors are
calculated from multiple stained embryos using the ImageJ software.

Figure 4. PRE prediction in *ey* locus. (a) Regulatory elements at *ey* locus. Known
enhancers (orange) of *ey* identified in previous studies which have shown *ey* specific
enhancer activity are present at upstream, i.e., *ey_12E* and in the second intron,
i.e., *ey_3.6R* and *ey_D02* (HAUCK *et al.* 1999; ADACHI *et al.* 2003). The two
boundaries, ME and EB, demarcate the *ey* locus. Two *ey* promoters, *ey_1* and *ey_2*
predicted in *ey* locus at eukaryotic promoter database (DREOS *et al.* 2015). **(b)** A
~1.6kb *ey*-PRE and a core 831bp *ey*-dPRE in *ey* locus predicted by PRE mapper.
The *ey* promoter *ey_2* also coincide with *ey*-PRE. **(c)** Mapping of histone
modifications, binding sites of PcG and boundary proteins, and DHS at *ey* locus in
the 14-16h whole embryo. H3K27Ac (blue), H3K27Me3 (red), Pc (purple),
dRING(brown), Psc(green), boundary proteins (black), and DHS(orange). ChIP-seq
data from the modENCODE project (MOD *et al.* 2010) and DHS data of stage 9 (~6h)
embryo from Thomas et al. 2011 (THOMAS *et al.* 2011) study have been used at the
UCSC browser to generate the map. The *ey* locus has a repressed H3K27Me3
domain, whereas neighboring genes *myo* and *bt* both have active H3K27Ac marks at
their promoters. The binding of Pc spreads over the repressed domain of *ey* while
dRING, and Psc, along with Pc, show a sharp peak at upstream to the first exon.
The *ey*-PRE aligns with the strong peak of PcG proteins, and core *ey*-dPRE maps to
DHS. Boundary proteins BEAF-32, CP190, and CTCF bind at both ME and EB that
appear to delimit the repressed domain of *ey*.

1017

Figure 5. Functional validation of *ey*-PRE (a) A PRE assay vector pCasPeRX.
The test DNA is inserted upstream to a *mini-white*. If the test DNA is PRE, it causes
repression of *mini-white*. The test fragments later can be removed with the help
of *loxP* to confirm the PRE mediated repression activity and rule out any position
effect of the genomic environment. **(b)** *ey*-PREs function as a repressor and show
strong PSS. Both, *ey*-PRE (~1.6kb) and *ey*-dPRE (831bp) exhibit stronger

repression in of *mini-white* in homozygous (P/P) compared to their heterozygotes

(P/+). The flipped-out versions of the same (ΔP/+ and ΔP/ΔP) show the de-

repression of *mini-white*. More than 50% of *ey*-PREs lines show intense repression

of *mini-white* represented by very light and variegating eye color. The smaller

version *ey*-dPRE show repression activity to a similar extent suggests that smaller

fragment is sufficient for the PRE activity (see Table1 and supplementary table

4). **(c)** Effect of PcG mutations on *ey*-PRE. All eyes are heterozygous for *ey*-PRE

and PcG mutations. The graph represents a quantitative analysis of the eye color in

the heterozygous condition in wild type and different PcG proteins mutation

background. PcG mutations, Pc^1 and Pho^1 both show a mild de-repression effect

while Psc^1 attenuates the *ey*-dPRE function significantly (see Table 2). Mutation in

GAF (Trl^{R85}) also attenuates the PRE repressor effect. A double PcG mutation of Pc^1

and Psc^1 show accumulative de-repressions effect.

1037

Figure 6. ME, EB, and *ey*-PRE interact in long-range to regulate *ey* locus. **(a)**

Interaction of ME with various regions at *ey* locus. 3C was performed in the 0-16h

embryo using EcoRI restrictions digestion followed by ligation, then Ligated hybrids

were PCR amplified using all forward primer pairs and confirmed by sequencing.

Interaction frequencies mean and standard deviations (b-vi) were calculated by gel

quantification of three replicates, as described (NAUMOVA *et al.* 2012). EcoRI

fragment that contains ME (primer 1) interacts with *ey*-PRE (primer 3) and with an

internal region (primer 4) and EB (primer 6) of *ey* (red lines in b-vi). However, such

interactions are not observed in a nearby upstream region towards *myo* (primer -3, -

2 and -1), a few internal regions of *ey* (primer 2 and 5) and in farther downstream

regions of *bt* (primer 7 and 8) (grey lines in a-vi). **(b)** Comparative analysis of

chromatin interaction at *ey* locus. (i & II) A merge Hi-C heatmap of chromatin

interactions in ~250 kb region of the fourth chromosome along with genes in Kc

cells. HiC data from Li et al. & Cubenas et al. (LI *et al.* 2015; CUBENAS-POTTS *et al.*

2017) visualized using HiC-Explorer (WOLFF *et al.* 2018) and chromatin loops

inferred from the same (shaded triangles). (iii) Interaction map in *ey* region inferred

from the enhancer-promoter interaction study used in an integrated method for

predicting enhancer targets (IM-PET) available from 4Dgenome

(https://4dgenome.research.chop.edu/) (HE *et al.* 2014) (iv) Genes and regulatory

region of *ey*. (v) EcoRI sites and forward primers used for 3C. (vi) & (vii) ME

interacting regions and interaction frequencies in the embryo identified by 3C in this
study. Together 3C and HiC data suggest ME and EB interacts in long range to
demarcates the ey domain.

1061

Figure 7. Different regions of eyeless associate to NuMat. (a) A map of *ey* locus
along with NuMat association of *cis*-elements ME, *ey*-dPRE, EB-u, and EB-i. **(b)**
NuMat association of *ey cis*-elements along with known MAR (histone MAR region –
HIS MAR) as a +ve and BEAF coding sequence (BEAF CDS) as a -ve control. The
PCR products of test and controls DNA (~400ng) were electrophoresed on 1%
agarose gel, transferred to the NY+ membrane. Finally, southern hybridized with
radiolabeled NuMat DNA from the embryos (0-16h), brain-eye-antennal disc (3rd
instar larvae), and S2 cells. **(C)** Relative NuMat association of all the test region is
compared with BEAF CDS (considered as one in embryos) and calculated by
blot/gel quantification using ImageJ. *Ey*-dPRE and EB-u show a strong NuMat
association (comparable to HIS MAR) in the embryo and brain-eye-antennal disc.
ME and EB-i show similar weak interaction to the matrix in the embryos while in
brain-eye-antennal disc EB-i does not appear to interact with the matrix. In S2
cells, only e*y*-dPRE associates with the matrix whereas in all experiment INT7 does
not interact NuMat (comparable to BEAF CDS).

1077

Figure 8. A Model for regulation of *eyeless* locus. Gene regulation in cells is a
result of a spatio-temporal higher-order organization of the chromatin. *Cis*-regulatory
elements aid in tethering of a locus into a needed sub-nuclear compartment, at least
in some regions, by attaching to the nuclear matrix. In the absence of activation
signal by initial regulatory proteins or upon receiving a repressive signal during
development, PcG proteins bind to *ey*-PRE to bring and maintain transcriptional
repression of *ey*. As *ey* and *bt* are present in a common TAD (Figure 6b) in Kc cells
where both the genes are transcriptionally inactive It is likely that *ey* and *bt* are in the
same repressed domain. Further, the *ey*-PRE tethering to the matrix and the
chromatin interactions of ME with *ey*-PRE, EB, and downstream regions of the *bt*
collectively determine the higher-order chromatin organization at the repressed
domain(**i**). While upon activation signal during development, the *ey* gene moves out
into an active compartment of the nucleus likely with the help of *ey*-PRE as it
associates to NuMat in all tissues (**ii**). Moreover, *ey* promoter that coincides with *ey*-

PRE becomes a target of activators, i.e., trxG proteins and enhancer proteins that

	ey-PRE	*ey*-dPRE
Total number of lines	31	26
% lines variegating	58 (18/31)	21 (4/19)
% lines with light eye color	52 (16/31)	50 (13/26)
% lines showing PSS	32 (7/22)	36 (5/14)
% lines showing increase in eye color in flipped-outs	59 (16/27)	47(8/17)

1093 keep *ey* active. At the same time, ME and EB boundaries demarcate the

1094 independent domain of *ey* that restrict cross-regulatory influences with neighboring

1095 genes *myo* and *bt.*

1096 **Table 1:** *ey*-PRE and *ey*-dPRE transgenes repressor activity

1097

1098

1099

1100 **Table 2:** PcG mutations effect on *ey*-PRE mediated repression

1101

ey-PRE lines	PcG Mutation	Effect	Strength
13.2	Pc^1	Derepression	+
	Psc^1	Derepression	++
	Pho^1	Derepression	+
	Trl^{R85}	Derepression	++
	$Pc^1;Psc^1$	Derepression	+++
76.1	Pc^1	No effect	
	Psc^1	Derepression	++
	Pho^1	No effect	
	Trl^{R85}	Derepression	++
	$Pc^1; Psc^1$	Derepression	++
100.2	Pc^1	Derepression	+
	Psc^1	Derepression	++
	Pho^1	Derepression	+
	Trl^{R85}	Derepression	++
	$Pc^1; Psc^1$		
88.1	Pc^1	Derepression	+
	Psc^1	Derepression	++
	Pho^1	Derepression	+
	Trl^{R85}	Derepression	++

	$Pc^1; Psc^1$	Derepression	+++
10.2	Pc^1	No effect	
	Psc^1	Derepression	++
	Pho^1	Derepression	+
	Trl^{R85}	Derepression	++
	$Pc^1; Psc^1$	Derepression	+++

1102

Figure 1

a

Test fragment

Enhancer

P

loxP   *loxP*

mini-white

SCS'

P

b

P/+   ΔP/+

INT7

EB

EB-u

EB-i

□P ■ΔP

Figure 5

a

Primer 1 (ME) +

| -3 | -2 | -1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | **Primer** |

200bp ➡ **Cross-linked**

200bp ➡ **Control**

EcoRI fragments interactions

b

ME boundary   *ey* enhancers   EB boundary

*ey*_PRE

myo   *ey*   *bt*   **MED26**

NuMat associated regions

Repressed state   Active state

i   *myo*   *MED26*

ey   *bt*

Repressed domain

ii   *myo*   *bt*

ey

Active domain

trxG proteins   Enhancer factors   PcG proteins   Transcription

Boundary proteins   Nuclear Matrix Proteins   Matrix   No Transcription