Ambulacrarian insulin-related peptides and their putative receptors suggest how insulin and similar peptides may have evolved from Insulin-like Growth Factor

Jan A. Veenstra[1],

[1] INCIA UMR 5287 CNRS, Université de Bordeaux, Pessac, France.

Corresponding Author:

Jan A. Veenstra[1]

INCIA UMR 5287 CNRS, Université de Bordeaux, allée Geoffroy St Hillaire, CS 50023, 33 615

Pessac Cedex, France

Email address: jan-adrianus.veenstra@u-bordeaux.fr

Abstract

Background

Some Insulin/IGF-related peptides (irps) stimulate a receptor tyrosine kinase (RTK) that transfers the extracellular hormonal signal into an intracellular response. Other irps, such as relaxin, do not use an

RTK, but a G-protein coupled receptor (GPCR). This is unusual since evolutionarily related hormones

typically either use the same or paralogous receptors. In arthropods three different irps, *i.e.* arthropod

IGF, gonadulin and *Drosophila* insulin-like peptide 7 (dilp7), likely evolved from a gene triplication, as

in several species genes encoding these three peptides are located next to one another on the same

chromosomal fragment. These arthropod irps have homologs in vertebrates, which suggests that the

initial gene triplication was perhaps already present in the last common ancestor of deuterostomes and

protostomes. It would be interesting to know whether this is indeed so and how insulin might be related

to this trio of irps.

Methodology

33 Genes encoding irps as well as their putative receptors were identified in genomes and transcriptomes

34 from echinoderms and hemichordates.

Results

A similar triplet of genes coding for irps is also found in some ambulacrarians. Two of these are

orthologs of arthropod IGF and dilp7 and the third is likely a gonadulin ortholog. In echinoderms two

novel irps emerged, gonad stimulating substance (GSS) and multinsulin, likely from gene duplications

of the IGF and dilp7-like genes respectively. The structures of GSS diverged considerably from IGF,

which would suggest they use different receptors than IGF, but no novel irp receptors evolved. If IGF

and GSS use different receptors and the evolution of GSS from a gene duplication of IGF is not

associated with the appearance of a novel receptor while irps are known to use two different types of

receptors, it seems to suggest that the ancestor of GSS and IGF might have acted on both types of

receptors while one or both of its descendants act on only one. There are three ambulacrarian GPCRs

that have amino acid sequences suggestive of being irp GPCRs, two of these are orthologs of the

gonadulin and dilp7 receptors. This suggests that the third might be an IGF receptor, and that by

deduction GSS only acts on the RTK. The evolution of GSS from IGF may represent a pattern, where

IGF gene duplications lead to novel genes coding for shorter peptides that activate an RTK. It is likely

this is how insulin and the insect neuroendocrine irps evolved independently from IGF.

Conclusion

The local gene triplication described from arthropods that yielded three genes encoding irps was

already present in the last common ancestor of protostomes and deuterostomes. It seems plausible that

irps, such as those produced by neuroendocrine cells in the brain of insects and echinoderm GSS

evolved independently from IGF and thus are not true orthologs, but the result of convergent evolution.

Key words: insulin; relaxin; receptor tyrosine kinase; G-protein coupled receptor; evolution; gonadulin;

octinsulin; multinsulin; dilp7

Introduction

Many protein hormone and neuropeptide signaling pathways have orthologs in both protostomes and deutertostomes showing that these pathways were already present in their last common bilaterian ancestor. In some cases the orthologs of the peptide ligands show only limited sequence similarity, but their receptors contain protein domains that are sufficiently conserved to establish homology. Virtually all ligands employ either a single receptor or a number of related receptors that evolved by gene duplication. Co-evolution of peptide ligands and receptors insures that related protein hormones or neuropeptides use receptors akin to those of their orthologs (Mirabeau & Joly, 2013; Hsueh & Feng, 2020).

Insulin/IGF-related peptides (irps) are an exception to this rule. Whereas insulin and IGF act through a receptor tyrosine kinase (RTK), relaxin uses a leucine-rich repeat G-protein coupled receptor (LGR). This raises the interesting question as how this apparent jump from one type of receptor to another may have come about. In cockroaches, termites and stick insects three different irp genes, gonadulin, arthropod insulin-like growth factor (aIGF) and arthropod relaxin, are located next to one another in the genome and thus likely originated from a local gene triplication (Veenstra, 2020b). To avoid confusion with the vertebrate relaxins and related peptides, the arthropod relaxins will be referred to as *Drosophila* ilp7 (dilp7) in this manuscript. One of theirps, aIGF, is known to use an insulin RTK, while gonadulin acts through insect LGR3 (Vallejo et al., 2015; Garelli et al., 2015; Colombani et al., 2015). Bioinformatic evidence suggested that dilp7 must be the ligand for insect LGR4 and this has now been confirmed experimentally in *Drosophila* (Veenstra, Rombauts & Grbić, 2012; Imambocus et al., 2020), but dilp7 may also activate an RTK (Linneweber et al., 2014). This suggests that the archtype arthropod IGF-related peptide acted through both an RTK and an LGR and that after a likely gene triplication, some of the ligands may have lost one of the two original receptors. Although it is possible that the gene triplication of the ancestral insulin gene occurred in an early arthropod or protostomian, it may well have occurred in a bilaterian ancestor, as homologs of both aIGF and dilp7 are also present in deuterostomes.

Brain neuroendocrine insect irps are more closely related to IGF than either dilp7 or gonadulin and a gene duplication that gave rise to separate genes encoding these peptides is therefore likely to have occurred after the triplication that gave rise to the ancestor genes of gonadulin and dilp7. Yet in insect genomes irp genes are not located near the IGF gene. Thus the particular organization of these genes suggests that whereas the gonadulin and dilp7 genes likely originated by two successive local gene

duplications, the IGF gene duplication that gave rise to an initial arthropod neuroendocrine brain irp

must have materialized in a different fashion. If the earlier mentioned gene triplication was already

present in the last common ancestor of the deuterostomes then a similar argument can also be made for

the evolution of insulin. Given the importance of insulin as a human hormone and the inherent interest

of its evolutionary origin, I explored the evolution of bilaterian insulin-related peptides in more detail

and here report on the genes coding for such peptides and their receptors in the Ambulacraria that

suggest how insulin may have evolved from IGF.

Materials and Methods

Nomenclature

Hormones have often been discovered independently by different groups using different bioassays.

The vertebrate insulin-like growth factors are a good example of that. Predicted protostomian peptides

and their receptors have sometimes been given names that refer to similar deuterostomian proteins. In

some cases this is very confusing, *e.g.* vertebrate LGR-3, -4 and -5 are not the orthologs of arthropod

receptors that have been given the same names. A similar problem occurs with arthropod relaxin that is

not an ortholog of vertebrate relaxin. This peptide will therefore be called dilp7 (*Drosophila* insulin-

like peptide 7). I will refer to arthropod LGR3 as the gonadulin receptor, arthropod LGR4 as the dilp7

receptor and arthropod LGR5 as GRL101, a GPCR initially identified from the pond snail *Lymnaea*

stagnalis (Tensen et al., 1994) that is an ortholog of arthropod LGR5 (Veenstra, 2020b).

Another nomenclature problem concerns the terms, insulin-like and insulin-related that are not well

defined. Insulin and IGF are related and must share a common evolutionary origin with other peptide

ligands like vertebrate relaxin, INSL3, arthropod dilp7 and gonadulin and a large number of other

bilaterian peptides. All these peptides are often collectively called insulin-like or insulin-related without

any specification as to in which aspects these hormones are similar to insulin. The typical core

sequence of six cysteine residues and its use of an RTK are two characters that are shared by vertebrate

IGF and insulin. However, several related peptides have eight cysteine residues and others like

vertebrate relaxin use an LGR and not an RTK. Insulin and IGF are also different in that IGF is a single

chain molecules, while the insulin precursor is processed into a two chain molecule. The term insulin-

like seems more appropriate for a subset of the insulin/IGF-related peptides that look similar to insulin

and act through an RTK, yet are different from IGF. Calling IGF-related peptides like vertebrate

relaxin, INSL3 or arthropod gonadulin for which there is no evidence that they act through an RTK,

insulin-like is confusing. Unfortunately for many bilaterian peptides we can only speculate as to which

type of receptor they use. The difference between one or two chain ligands, *i.e.* IGF versus insulin, is

also useless as there is good evidence that some insect IGF-related peptides are processed into two-

chain molecules when expressed in neuroendocrine cells and produced as single chain ligands when

produced by the fat body, yet in both cases stimulate an RTK. It is for these reasons that all these

peptides will be referred to as insulin/IGF-related peptides, abbreviated irps.

Sequence analysis

Sequences for insulin related peptides and their likely receptors were identified from a number of

Ambulacraria species. This was done using using the Artemis program (Rutherford et al., 2000) and the

BLAST+ program (ftp://ftp.ncbi.nlm.nih.gov/blast/executables/blast+/) on publicly available genome

sequences from the feather star *Anneissia japonica*, the sea urchins *Lytechinus variegatus* (Davidson et

al., 2000) and *Strongylocentrus purpuratus* (Sea Urchin Genome Sequencing Consortium, 2006), the

sea cucumbers *Apostichopus japonicus* (Jo et al., 2017; Zhang et al., 2017) and *Holothuria glaberrima*,

the sea stars *Acanthaster planci* (Hall et al., 2017)*, Pisaster ochraceus* (Ruiz-Ramos et al., 2020) and

Patiria miniata, the brittle star *Ophiothrix spiculata* and the hemichordates *Saccoglossus kowalevskii*

and *Ptychodera flava* (Simakov et al., 2015). The genomes were downloaded from

https://www.ncbi.nlm.nih.gov/genome. For many of these species there are also significant amounts of

RNAseq data and these were analyzed using the sratoolkit

(https://trace.ncbi.nlm.nih.gov/Traces/sra/sra.cgi?view=software) in combination with Trinity

(Grabherr et al., 2011) using methods described in detail elsewhere (Veenstra, 2020b). Some protein

sequences were found in the NCBI database, but several of them contain errors or are incomplete.

Where possible these were corrected and/or completed using the methods described above. As there is

only a single crinoid genome assembly available, transcriptome data from the three crinoid species

Antedon mediterranea, Florometra serratissima and *Oligometra serripinna* were also included. For the

same reason transcriptome data from the brittle star *Amphiura filliformis, Ophioderma brevispina* and

the hemichordate *Schizocardium californicum* were likewise analyzed. Obviously, transcriptome data

can only demonstrate the presence of gene but not its absence and their usefulness depends largely on

the variety of tissues sampled and the expression levels of the genes of interest. Nevertheless, such data

often provide additional sequences that even if they are incomplete increase the robustness of sequence

comparisons. Genomic and transcriptomic RNAseq short read archives (SRAs) were downloaded from

NCBI (https://www.ncbi.nlm.nih.gov/sra/); a list of the SRAs analyzed is provided in the

supplementary data.

As queries for the insulin-like peptides a number of such peptides from a variety of species was

used. Insulin RTKs are easily identified in genome and transcriptome assemblies, as their kinase

domains are very well conserved. The LGRs that could function as insulin receptors are more variable.

Vertebrate RXFP1 and RXFP2 are LGRs are known receptors for relaxin and Ins3 and *Drosophila*

LGR3 and LGR4 for gonadulin, and dilp7 respectively. Other LGRs function as receptors for the

various glycoprotein hormones, GPA2/GPB5, bursicon, TSH, FSH and LH. These GPCRs cluster on

phylogenetic trees with another protostomian LGR, GRL101. This GPCR was initially identified from

the pond snail *Lymnaea stagnalis* and was the first GPCR discovered to have in addition to six leucine-

rich repeats also twelve repeats of a sequence that was known to exist in the low density lipoprotein

receptor and are now called LDLa repeats (Tensen et al., 1994). I have suggested previously (Veenstra,

2020b) that this receptor might be an IGF receptor.

Both the RTK and LGR receptors have large ectodomains. Those of the insulin RTKs are very

similar from one receptor to another, while those of the LGRs differ between different types. The latter

all contain numerous Leucine-rich repeats (LRRs) and some also have LDL-receptor class A (LDLa)

repeats. Both LRRs and LDLa's are present in many other proteins. Initial searches for orthologous

receptors were therefore done using the transmembrane regions of various insect and vertebrate LGRs

and the protein kinase domain of RTK. Once partial sequences of putative receptors were identified, the

coding sequences of these domains were then used to complete the cDNA sequences as best as

possible, using either Trinity on RNAseq SRAs or Artemis on genome sequences.

Sequence similarity and phylogenetic trees

Both phylogenetic and sequence similarity trees use Clustal omega (Sievers et al., 2011) to produce

alignments. Fasttree (Price, Dehal & Arkin, 2010), using the ./FastTreeDbl command with the -spr 4, -

mlacc 2 and -slownni options, was used to construct trees and estimate probabilities.

In order to identify putative receptors for the various irps, LGRs that show homology to various

arthropod and vertebrate LGRs were identified and a phylogenetic tree based exclusively on the

transmembrane regions of these receptors was constructed.

Precursor processing

Precursors of insulin-like peptides contain signal peptides that are removed on entry into

endoplasmatic reticulum. Signal P 5.0 (Almagro Armenteros et al., 2019) was used online

(http://www.cbs.dtu.dk/services/SignalP/) to predict where this cleavage would most likely occur.

Some, but not all precursors are further processed by convertases. Of these furin is ubiquitously present

in all cell types and can thus potentially cleave any secreted protein with appropriate cleavage site. Its

consensus cleavage site is K/R-X-K/R-R, the two human IGF precursors are processed at KSAR and

KSER respectively (Humbel, 1990). Precursors that are produced in cells with a regulated pathway,

such as neuroendocrine and enteroendocrine cells, are also exposed to other convertases like PC1/3 and

PC2. Their consensus cleavages site is KR. However, effective proteolytic processing by convertases is

strongly influenced by amino acid residues surrounding these consensus cleavage sites. For example

bulky residues immediately following the arginine residue, a proline residue before the consensus site

or disulfide bridges nearby can cause sufficient steric hindrance to inhibit cleavage. Using rules

proposed to predict cleavage by PC1/3 and PC2 in both vertebrates and insects (Devi, 1991; Rholam et

al., 1995; Veenstra, 2000) I have tried to indicate in Figs. 3-6 where the various precursors might be

cleaved. It must be noted though that there is no certainty that these site will be cleaved nor can it be

excluded that proteolytic processing occurs at sites that have not been indicated as such.

Expression

With a few notable exceptions (*e.g.* Lin et al., 2017), little is known about the expression of the

various insulin-like peptides in either echinoderms or hemichordates and except for the GSS our

knowledge of their functions is also very limited. Expression data may reveal some preliminary clues

as to where and when they are expressed and thus provide a hint as to their function. For this reason the

number of reads corresponding to the various insulin-related peptides and their putative receptors was

determined in a number of SRAs that might provide evidence as to the time and tissue specific

expression of these proteins. The analysis was performed as described previously (Veenstra, 2020b)

and the data are supplied in Spreadsheet S2.

Results

Peptides related to insulin and IGF

Some protein sequences were found in the NCBI database, but several of them contain errors or are

incomplete. Where possible these were corrected and/or completed using the methods described above.

As there is only a single crinoid genome assembly available, transcriptome data from the three crinoid

species *Antedon mediterranea, Florometra serratissima* and *Oligometra serripinna* were also included.

For the same reason transcriptome data from the brittle star *Amphiura filliformis, Ophioderma*

brevispina and the hemichordate *Schizocardium californicum* were likewise analyzed. Obviously,

transcriptome data can only demonstrate the presence of gene but not its absence and their usefulness

depends largely on the variety of tissues sampled and the expression levels of the genes of interest.

Nevertheless, such data often provide additional sequences that even if they are incomplete increase the

robustness of sequence comparisons.

Insulin-like peptide precursors are typically characterized as having A, B and C domains that

correspond to the A- and B-chains of insulin and the connecting peptide respectively. In IGF D and E

domains are also recognized, in which the D domain refers to the extension of the A chain and the E

domain to part of the precursor after the D domain that is cleaved from IGF in the Golgi apparatus. For

dilp7 orthologs it is appropriate to add an F (front) domain for the sequence in N-terminal of the B-

chain that in some peptides is not only larger, but also well conserved (Fig. 1).

Previous work on insulin-related peptides in in echinoderms have identified two different types of

insulin-like peptides, gonad-stimulating substances (GSS) and insulin-like growth factors (Mita et al.,

2009; Perillo & Arnone, 2014; Semmens et al., 2016; Smith et al., 2019). The insulin-like growth

factors, but not GSS, are also present in hemichordates. While only a single IGF gene was found in the

crinoids and hemichordates, other ambulacrarians have two such genes (Figs. 2, S1, S2; Spreadsheet

S1). These proteins have large C-terminal extension that are rich in charged amino acid residues,

especially arginine and lysine, but also aspartic and glutamic acid residues. A comparison of the protein

sequences and cDNAs from human IGFs identifies the exact separation between the D and E domains

in these proteins (Humbel, 1990). However, although the corresponding sequences of the hemichordate

and echinoderm IGFs contain numerous arginine and lysine residues (Figs. 2, S1, S2), there are no

obvious convertase cleavage sites as many potential arginine residues are succeeded by residues known

to inhibit such enzymes in vertebrates. It is thus not impossible that the D domains of these proteins are

much larger than in the vertebrate IGFs and if so likely contain numerous positively charged amino

acid residues. There are few transcriptome SRAs for specific tissues, the data that is available suggest

that the IGFs are expressed by many tissues, with the ovary showing significant expression. *Patiria*

pectinifera is the only species with follicle cell specific SRAs and IGF-1 is strongly expressed by these
cells and is probably transferred to the oocyte (Spreadsheet S2).

The GSS are known to induce oocyte maturation and ovulation in a two step process, where GSS
stimulates the follicle cells to produce 1-methyladenine which subsequently induces resumption of
meiosis in the oocyte and about 30 minutes later this is followed by ovulation (Chiba, 2020).
Interestingly, GSS was not found in either the genome nor the extensive transcriptome data from the
feather star *Anneissia japonica* and was similarly not encountered in the transcriptomes of three other
crinoids (Suppl data). Transcriptomes may miss expression of some genes and large genome assemblies
are never perfect. The short sequence reads in the genomic SRAs from *Anneissia* were therefore also
analyzed for the presence of GSS, but again no evidence for such a gene was found. This peptide is
thus likely absent from *Anneissia* and perhaps all Crinoidea. In the Holothuroidea and the Asterozoa,
but not the Echinoidea, this gene is duplicated with the two paralogous peptides showing significant
sequence variability (Figs. 3, S3, S4; Spreadsheet S1). As for all these peptides and their putative
receptors expression data is very limited, but in *Apostichopus* the two GSSs are differentially
expressed, with GSS-1 being expressed at specific stages during embryonic development as well as by
muscle and GSS-2 strongly expressed by both the ovary and the testes. Interestingly, it is the ortholog
of GSS-1 that in *Holothuria scabra* has been tested for biological activity and induces ovulation (Chieu
et al., 2019). This makes one wonder what the effects of GSS-2 on ovulation might be in this species.
However, *Apostichopus* was the only species where a significant GSS expression was found in the
gonads (Spreadsheet S2).

Two other insulin-like peptides are commonly present in both hemichordates and echinoderms,
including the Crinoidea. The first is an ortholog dilp7 which has a very characteristic F domain while
its A chain is also remarkably well conserved (Figs. 4, S5, S6; Spreadsheet S1). The precursors of this
peptide contain typical neuroendocrine KR convertase sites and seems to have its highest expression in
the nervous system, although it is also found in other tissues. During embryogenesis its expression
occurs relatively late (Spreadsheet S2). The second peptide present in all ambulacrarians has been
called octinsulin as it has eight cysteine residues and is thus predicted to have four rather than three
disulfide bridges. In echinoderms octinsulin is a single copy gene, but hemichordates have several such
genes (Fig. 5, S7, S8; Spreadsheet S1). Octinsulin expression levels are the highest in nervous tissue,
and significant expression is also found in the gut and stomach of *Strongylocentrotus* and *Patiria*

pectinifera respectively. Although virtually absent from normal gut in *Apostichopus*, it has significant

expression during gut regenerating in this species (Spreadsheet S2).

The Asterozoa have genes coding for a fifth type of insulin, that is usually present in multiple

copies and that are referred to as multinsulins. The predicted peptides share structural similarity with

the dilp7 orthologs and their genes have typically four coding exons rather than the two or three of the

other irp genes. The sprawl of these peptides is perhaps best illustrated by a phylogenetic tree that

suggest independent multiplication of these genes in several species (Fig. S10). Within a single species

the various multinsulins thus often seem more closely related to one another than to their putative

orthologs of other Asterozoa. Some of the multinsulins, like the octinsulins, have acquired two

additional cysteine residues and are thus predicted to have four disulfide bridges, but the location of

these additional cysteine residues differs from that in octinsulins (Figs. 6, 7, S9, S10; Spreadsheet S1).

Like dilp7 the multinsulins have typical neuroendocrine KR convertase cleavage sites and can thus be

expected to be expressed in neuroendocrine and/or enteroendocrine cells but expression data on *P.*

pectinifera suggest a relatively ubiquitous expression in several tissues.

The genome assemblies of *A. planci* and *Pisaster ochraceus* shows these genes to be clustered in

the genome and some RNAseq sequences suggests that at least on occasion coding exons from different

genes may be combined (Fig. S10). This and the large numbers of SNPs typically present in animals

caught in nature and used for RNAseq preparation make it impossible to reliably determine their exact

numbers.

Genome assemblies allow identification of the introns in these genes. All insulin genes have a

characteristic phase 1 intron somewhere in the conceptual C domain of these molecules. This is the

only intron in the coding sequences of the octinsulin and GSS genes. The IGF genes have a phase 0

intron near the end of the coding sequence and at least some of them have another phase 1 intron just

after the transcription start site. The genes coding for the dilp7 orthologs and multinsulins share an

additional phase 2 intron and the multinsulin genes have yet another phase 1 intron. All these introns

appear perfectly conserved (Fig. 7).

Synteny of genes producing insulin-like peptides

In the *Strongylocentrotus* genome all five genes are located on the same chromosome, with the two

IGF genes and those encoding octinsulin and dilp7 orthologs next to one another and GSS at a distance

of 6,000,000 bp (base pairs). At least the *Anneissia* octinsulin and IGF genes are likely located next to

one another on the same chromosome also, as in the current genome assembly two of the three coding

exons of IGF and one of the two octinsulin coding exons are located within about 10,000 bp. The three

missing exons of these two genes are all located on minicontigs of less than 2,000 bp, as is one of the

coding exons for the dilp7 ortholog. The contigs of the *Lytechinus variegatus* genome assembly are

smaller and this may explain why in this species the genes are located on three different scaffolds, with

the two ILGF-like peptides and the octinsulin together on a single contig. However in the recently

published genome of the closely related *L. pictus* (Warner et al., 2021) the dilp7 ortholog is also closely

associated with the other three genes. The GSS gene is on the same chromosome but at a distance of

28,000,000 bp. In the *Apostichus japonicus* genome assembly the genes encoding the octinsulin and the

two IGF genes are located on the same contig, and the other genes each on a different one. In the draft

Holothuria glaberrima genome assembly only the two IGF genes are located on the same contig,

however in a single Oxford nanopore read (SRR9125585.2851.1) from *H. scabra* the octinsulin, dilp7

and two IGF genes are located next to one another as well (Fig. 8).

Whereas the various Echinozoa genome assemblies suggest a certain degree of synteny with regard

to the various irp genes, the Asterozoa genome shows that such syntenty is disintegrating. This is most

clearly demonstrated in the genome assemblies from *Pisaster ochraceus* and *Acanthaster planci,* where

the scaffolds are much larger than from *Patiria miniata.* In these species synteny is largely lost (Fig. 8).

Interestingly the various multinsulin genes are present in small clusters on different chromosomes in

those species.

Sequence similarity tree peptides related to insulin

Peptides having the characteristic insulin signature are notoriously variable in their primary amino

acid sequences. Although the various residues allows one to align those sequences, such alignments

will not always yield reliable phylogenetic trees as the basic tenet of such analyses is often not met. As

an alternative I have proposed to use "sequence similarity trees". Such trees are constructed using the

same methods but do not pretend to illustrate phylogenetic relations, rather similarities between the

different proteins.

The structures of the multinsulins are most similar to the dilp7 orthologs (Fig. 6) and so it is not

surprising that the sequence similarity tree (Fig. 9) groups the multinsulins with the dilp7 orthologs.

The hypothesis that this structural similarity between these two types of peptides may reflect a close

evolutinary relationship is reinforced by the presence of an intron that is present in the genes encoding

these peptides but lacking in the genes encoding octinsulin, IGF and GSS (Fig. 7). The tree also

illustrates significant sequence similarity between GSS and the IGF.

Orthologs of receptors for irps: Receptor tyrosine kinase

A single insulin RTK gene was found in all species analyzed here. An alternatively spliced form is

present in *Acanthaster* and is likely commonly present in echinoderms (Spreadsheet S1). Hundreds of

ambulacrarian protein sequences were identified at NCBI using a BLAST search with the *S. kovalevskii*

protein kinase domain as a query. After aligning them with Clustal omega the protein kinase domains

were used to make a phylogenetic tree. Results revealed no other known or predicted proteins with a

similar protein kinase domain (data not shown). The insulin RTK is ubiquitously expressed

(Spreadsheet S2).

Orthologs of receptors for irps peptides: LGRs

LGR sequences were obtained using the combination of genomic sequences and, where available,

transcriptome shotgun sequences and RNAseq SRAs. The latter were used to produce contigs using

Trinity (Spreadsheet S1). Short read assemblers are good in combining sequences into larger

continuous ones, but they do produce artifacts, which are more easily obtained when very similar

sequences are present in multiple copies, such as the multinsulins, or the numerous LDLa and LRR

repeats. These repeats are usually individually coded by single exons that are sometimes skipped and

when such skipped individual reads enter in the RNAseq SRA, incorrect constructs are obtained.

Furthermore, these repeats are present in numerous proteins, and from time to time this leads to

assembled sequences that are from mRNA species from different genes. It is therefore to be expected

that not all assembled transcripts, neither those in the databank nor those produced here, will be correct.

Some errors were corrected by challenging divergent sequences that were discovered on comparing

putative orthologs with one another. Other differences could be confirmed as true differences, but it is

not impossible that some errors remain, particularly for those sequences that are incomplete. LGRs that

might function as receptors for the various irps were identified by their homology with such receptors

from vertebrates and arthropods. The transmembrane regions of the GPCRs don't have the assembly

problems of the LDLa and LRR repeats and are the most characteristic domain of the GPCRs. This

makes it easier to construct a phylogenetic trees for these receptors based on their transmembrane

regions than that it is to produce complete LGR transcripts.

Results show a surprisingly similar distribution of LGRs in the species studied. The tree resolves

two major branches, one for the glycoprotein hormone receptors, which itself is divided in two

subbranches, one for orthologs of the GPA2/GPB5 receptor - containing the receptors for human TSH,

FSH and LH - and a second one for the bursicon receptor orthologs. All species studied are represented

by one member on each of these two subbranches, except for *Ophiothrix*, where the draft genome

reveals two orthologs each for the bursicon and GPA2/GPB5 receptors (Fig. 10). These are likely

receptors for the bursicon and GPA2/GPB5 orthologs identified from various echinoderm species

(Semmens et al., 2016). It is interesting to see that whereas vertebrates have different receptors for

TSH, FSH and LH, most echinoderms have only one GPA2/GPB5-receptor ortholog (Fig. S11), even

though *A. rubens* has two GPA2 and three GPB5 orthologs (Semmens et al., 2016). The LGRs for the

glycoproteins were included in the search for putative receptors for the ambulacrarian irp LGRs in

order to be sure that no such receptors would be missed.

The lower branches of the LGR phylogenetic tree are the ones of interest as they contain receptors

with irp ligands. It consists of three subbranches, that are characterized by *Drosophila* LGR3 and

LGR4 – the receptors for gonadulin and dilp7 respectively - and *Periplaneta* LGR5, an ortholog of

Lymnaea GRL101. Here in all ambulacrarian species studied only one ortholog was found for each of

them, despite extensive attempts to find additional LGRs in the various genomes and transcriptomes.

The GRL101 transmembrane regions puts it very close to vertebrate glycoprotein hormone and

relaxin LGRs. LRRs are present in many different proteins, but when the LRR part of the *Anneissia*

GRL101 (amino acid residues 576-717) is used as query in a protein BLAST against human proteins,

the glycoprotein hormone and relaxin receptors are identified as most similar to this ectodomain of

GRL101, suggesting that similarity of the GRL101 receptors with vertebrate LGRs is not limited to the

transmembrane region of this GPCR.

Sequence alignments of these GPCRs show strong sequence similarity (Figs. S12-S14), however

the dil7 receptor ortholog varies more between species. A schematic representation of the the

ectodomains of the LGRs on this second branch is drawn in Fig. 11. The orthologs of the dilp7 and

gonadulin receptors each have a single LDLa repeat, except for the *Patiria* and *Acanthaster* orthologs

of the dilp7 receptor which both have two LDLa repeats (Fig. S13). This additional LDLa is likely due

to a relatively recent duplication of the LDLa since the two LDLa repeats have very similar amino acid

sequences (Spreadsheet S1). All three receptors are expressed in the nervous system and the gonadulin

receptor is well expressed in the gonads, both testis and ovary, and strongly so in the follicle cells of *P.*

pectinifera (Spreadsheet 2).

Discussion

The genomic and transcriptomic data from both the hemichordates and the echinoderms show that

these two groups share three irps, octinsulin, IGF and a dilp7 orthologs, that are present in both

echinoderms and hemichordates. IGF and dilp7 are orthologs of the arthropod peptides that together

with gonadulin originated from a gene triplication. The structure of gonadulin is poorly maintained,

even within insects (Veenstra, 2020b). The variable structure of gonadulin and its loss in many

arthropod lineages suggests that the evolutionary pressure on gonadulin is weak. This may explain why

the amino acid sequence of gonadulin looks significantly different from octinsulin. Nevertheless, there

are two lines of evidence that suggest that these peptides must be orthologs as well. For one, synteny of

the chromosome fragment containing these genes is conserved between the sea urchin

Strongylocentrotus purpuratus, the hemichordate *Saccoglossus kowalewski* and the cockroach *Blattella*

germanica, suggesting that these peptides are likely orthologs. More importantly, all ambulacrarians

have an ortholog of the gonadulin receptor and the only plausible ligand for such a receptor encoded by

their genomes is octinsulin. Thus the gene triplication previously reported from arthropods must have

occurred in a common bilaterian ancestor of the deuterostomes and protostomes.

Crinoids have the simplest irp signaling system, one gene each for IGF, octinsulin and the dilp7

ortholog. Their putative receptors - insulin RTK, GRL101, and the orthologs of the dilp7 and gonadulin

receptors – similarly are also each coded by a single gene. The hemichordates have a very similar

repertoire, except that the octinsulin gene is systematically amplified and in some species the dilp7

ortholog as well. It thus appears likely that the first deuterostome had a single copy of each of these

genes.

Within the echinoderms the irp genes evolved considerably, as shown both by an increase in their

numbers and the loss of synteny. Whereas the feather stars appear to have only a single IGF gene, all

other echinoderms have two such genes and two novel irps, GSS and multinsulin, appeared. The GSS

sequences are most similar to those of IGF, suggesting that they evolved from a gene duplication event

from the IGF gene. Although some GSS genes are located on the same chromosome as the other irps,

they are not close to the IGF genes, indicating that the IGF-GSS split was not a local duplication but

may have been the result of an incorrectly repaired chromosome break.

In the Asterozoa a fifth type of irp gene emerged, those that code for the multinsulins which share

significant sequence similarity with the dilp7 orthologs. The initial multinsulin gene must thus have its

origin in a gene duplication of the dilp7 ortholog gene, with which they furthermore share a

characteristic intron. Later the multinsulin gene seems to have undergone several additional gene

duplications in this respect the multinsulins resemble the insect neuroendocrine irps.

The co-evolution of ligands and receptors allows one to assign the putative receptors for gonadulin,

the dilp7 ortholog and IGF as the orthologs of the receptors of their arthropod orthologs. This allows

the identification of the ambulacrarian LGRs that are the orthologs of the gonadulin and dilp7 receptors

as likely receptors for octinsulin and the dilp7 respectively, as well as the insulin RTK as a receptor for

IGF.

The appearance of the multinsulins is not accompanied by the evolution of a novel insulin-receptor.

Some animals have multiple insulin RTKs, *e.g.* some arthropods have up to four such genes (Veenstra,

2020a,b), however, in spite of extensive searches for a second insulin RTK in ambulacrarian genomes,

none was found. Searches for an additional LGR that might function as a receptor for the GSS and/or

multinsulin were unsuccessful and this raises the question which receptors are activated by these

peptides.

I have previously argued that the close chromosomal association of the IGF, gonadulin and dilp7

ortholog genes in basal insects suggest that they derived from a gene triplication (Veenstra, 2020b).

There are three possible scenarios that can explain how IGF and gonadulin came to respectively

activate an RTK and an LGR. It is possible that the original irp activated either an RTK and that an

LGR was later acquired as a second receptor by gonadulin, alternatively the original irp activated an

LGR and IGF acquired an RTK as a second receptor. Given the importance of insulin RTKs for growth

in very basal metazoans, it is improbable that the original irp activated an LGR and that an RTK was

acquired much later during evolution (see *e.g.* Mortzfeld et al., 2019). This indicates that an irp

acquired an LGR as a second receptor and the question is whether this happened before or after the

gene triplication that yielded IGF, gonadulin and dilp7. In both *Saccoglossus* and arthropods the IGF

gene is in the middle of the three. This suggests that this represents the gene organisation after the gene

triplication and that the dilp7 and gonadulin orthologs each evolved independently from the arch irp

rather than that dilp7 evolved from gonadulin or *vice versa*. Had dilp7 originated from a gene

duplication of gonadulin or the other way round, they also might have been more similar to one another

than they are. The acquisition of a second receptor must be an extremely rare event. Since both

gonadulin and dilp7 use an LGR this would mean that such an extremely rare event of the acquisition

of a second receptor would have occurred not only twice, but even with a very similar receptor.

Furthermore, some metazoans have an LGR that is closely related to the dilp7 and gonadulin LGRs

suggesting that it could be an IGF receptor (see below). It is for these reasons that the author favors the

hypothesis that the arch irp already acted on both an LRG and an RTK, but, clearly, this remains a

hypothesis.

The binding of insulin and relaxin to their respective receptors has gotten resolved in much detail

in the last couple of years. The effective binding and stimulation of insulin RTK by the small irp from

the snail *Conus* to the RTK shows that a small irp can be an effective ligand for this receptor (Menting

et al., 2015). On the other hand the complex interaction of relaxin to its LGR makes it more difficult to

imagine a smaller peptide as an effective ligand (Hoare et al., 2019). Furthermore, considering the well

conserved F-domain of the dilp7 receptor orthologs it is likely that it is necessary for interaction with

its LGR receptor. The loss of this structure in multinsulin suggests that it is unlikely to be a dilp7

receptor agonist. On the other hand, the poor sequence conservation in the various *Drosophila* irps that

activate a single RTK is reminiscent of the large structural variability of the multinsulins. This seems to

suggest that the multinsulins are RTK ligands rather than that they activate the LGR.

The emergence of the GSS is neither accompanied with the evolution of a novel receptor for these

irps. This can also be explained by assuming that IGF acts on both the RTK and an LGR and that the

GSS have lost their affinity for the LGR. This raises the question whether an IGF LGR might exist.

If there were an IGF LGR, one would expect it to be related to the gonadulin and dilp7 receptors.

GRL101 appears a plausible candidate as its transmembrane regions are closely related to the receptors

for gonadulin and dilp7. The ectodomain of GRL101 consists of two parts, a series of LRRs and a

second series of LDLa's. In the related GPCRs the LRRs are expected to bind with the insulin core of

gonadulin and dilp7 orthologs, just like the human relaxin receptors (Hoare et al., 2019). When the

LRR part of the *Anneissia* GRL101, the most basal echinoderm, was used as query for similar human

proteins in a BLAST search, the glycoprotein hormone and relaxin receptors were identified as the

most similar proteins. This shows that the resemblance of GRL101 to the other LGRs is not limited to

the transmembrane regions and reinforces the hypothesis that the ligand of GRL101 has an insulin-like

structure. GRL101 has a large number of LDLa's, the ligands of which are typically positively charged

surfaces, which in the case of proteins consist of Lys and Arg residues (Daly et al., 1995; Prévost &

Raussens, 2004; Fisher, Beglova & Blacklow, 2006; Yasui, Nogi & Takagi, 2010; Dagil et al., 2013).

Thus the ligand of GRL101 may consist of two parts, an insulin-like structure and a piece with several

positive charges that interact with the LDLa's. The C-terminal tails of the IGFs, whether from

arthropods, echinoderms or hemichordates, are all rich in charged amino acid residues. The C-terminal

tail of IGF with its numerous positively charged amino acid residue might interact with the LDLa's of

GRL101. I therefore posit that in those species that have a GRL101 it functions as the second receptor

for IGF. The absence of such a tail in GSS would make it likely that it acts on the RTK rather than an

IGF GPCR.

The suggestion that GSS activates the RTK goes against the hypothesis that these peptides act

through GPCRs. Indeed it has recently been proposed that it is the ortholog of the dilp7 receptor that

would be activated by the gonad stimulator in *P. miniata* (Mita et al., 2020). Given the clear orthology

of both the dilp7 echinoderm orthologs with the *Drosophila* peptide and the similar orthology between

the dilp7 receptor and the echinoderm receptor, the conclusion that the two constitute a functional

ligand receptor combination seems inescapable. It was impossible to find a GSS in either the genome

assembly or the individual reads of all the genomic SRAs of *Anneissia japonica*, yet it does have a

dilp7 receptor ortholog, thus if the dilp7 receptor were to function as a GSS receptor, it most likely

would not be an exclusive receptor. *A priori* this does not exclude the possibility that GSS could

function as a ligand for the same receptor. As mentioned above, since the dilp7 orthologs have well

conserved F domains, one has to assume that it is important for binding to its receptor. Since this

domain is absent from RTK ligands, it is difficult to understand how GSS that similarly lacks this

domain would be able to bind the dilp7 receptor. It would thus seem unlikely that peptides as different

as GSS and dilp7 would be effective ligands of the same LGR. Furthermore, the GSS genes have been

duplicated and their structures have diverged considerably. Those duplicate gonad stimulators are

present in many species and have not been selected against. Hence they must be phyisologically

relevant and able to interact with a receptor. Sharing a common evolutionary origin the two gonad

stimulators should be expected to act either on the same or paralogous receptors, but the number of

putative echinoderm receptors for irps is limited, so it must be the same one. The same arguments that

were used to argue that the multinsulins are likely RTK agonists but not LGR ligands, are therefore

equally valid here and suggest that GSS is an RTK ligand.

Furhermore, the experimental evidence that GSS stimulates the ortholog of the dilp7 receptor is not

convincing. The reported response to the dilp7 receptor when expressed in Sf9 cells is very weak and

does not represent a typical response seen in this type of assay. Although the authors have shown high

affinity binding of GSS to the follicle cells, such high affinity binding should also have been present in

the Sf9 cells expressing the putative GSS receptor, but this was not reported. The follicle cell SRAs

from which the putative GSS receptor was identified contains large amounts of RNAseq reads for the

gonadulin receptor, a receptor that is more closely related to the vertebrate relaxin receptors than the

dilp7 receptor, but surprisingly the authors do not mention this receptor, which they must have found

(Mita et al., 2020).

I suggest that initially there was an IGF-like hormone that activated both a GPCR and an RTK,

after two gene duplications some of the descendant ligands either lost their C-terminal tails or one

acquired a larger one and this allowed all three ligands to activate, at least initially, the RTK while each

acquired its own LGR. Later, some of the ligands may have lost their affinity for one receptor. Since

the primary amino acid sequence of gonadulin is very different from that of the other irps, it likely lost

its capacity to activate the RTK (Fig. 12). Holometabolous insect species have lost GRL101 and hence

in those species IGF can only act on the RTK. Under this hypothesis the arginine-rich C-terminal tail

would be useless in such insect species and in higher flies, such as *Drosophila,* it was indeed lost

(Veenstra, 2020b). In vertebrates, there is no GRL101 and so IGF can only activate the two RTKs,

while the relaxin related peptides are not known to interact with RTK. The presence of a similar

arginine-rich E domain of the vertebrate IGF precursors might thus be an evolutionary relict.

This scheme raises the question as to how the functions of these two receptors activated by IGF

might differ. IGF and the drosophila irps stimulate growth, the echinoderm GSS stimulates oocyte

maturation and ovulation (Mita et al., 2009), relaxin and INSL3 affet various developmental and

reproductive processes (Ivell et al., 2020; Esteban-Lopez & Agoulnik, 2020), gonadulin is expressed by

the gonads as well as the imaginal in flies (Garelli et al., 2012; Liao & Nässel, 2020; Veenstra, 2020b;

Veenstra et al., 2021) and dilp7 is expressed in a sex specific manner (Miguel-Aliaga, Thor & Gould,

2008; Yang et al., 2008; Castellanos, Tang & Allan, 2013). These hormones stimulate growth,

development and reproduction, processes that are intimately linked; without growth and development

reproduction is impossible and growth without reproduction is useless in sexually reproducing species.

On the other hand, resources used for growth and development can not be used for reproduction or *vice*

versa.

Growth is rarely a linear process independent of development; animals are not only getting bigger,

but they also mature into adults. Metamorphosis is markedly different between hemi- and holo-

metabolous insect species. Every time a cockroach nymph molts, it becomes a little more adult,

however during the first molts of a caterpillar the insects mainly become bigger, it is only when it molts

into a pupa that it significantly changes its morphology. Cockroaches have GRL101, caterpillars don't.

This suggests that the RTK might be more directed toward linear growth, or allowing growth by

increasing uptake of resources, such as glucose and amino acids, while the LGRs might be more

important for insuring that the animal develops into an adult and becomes sexually competent. Both

holometabolous insects and vertebrates have lost GRL101 and use steroid hormones to induce sexual

maturation. Interestingly, in vertebrates the production of steroid hormones is controlled by

glycoprotein hormones, the second group of ligands for LGRs.

It is plausible that IGF in an early bilaterian was produced by the tissue that stored energy and

perhaps even protein as insects do in the form of storage proteins (Haunerland, 1996). Production and

release of IGF might have happened when the animal had sufficient resources to allow for growth

and/or reproduction. In arthropods growth has become a discontinuous process in which a new cuticle

needs to be made before molting can take place. In those species IGF produced by the fat body may

well be the essential growth hormone. However, if the animal is suddenly starved, IGF would no longer

be released. If formation of a new cuticle is too advanced to be interrupted, this become problematic. It

may have obliged the brain to take at least partial control of growth away from the fat body by

releasing one or more of the neuroendocrine irps to force growth and molting to proceed. It is possible

that this achieved by simultaneously reducing growth of organs that are needed for (sexual) maturation

but not essential for immediate survival, like the gonads. This could be how the neuroendocrine insect

irps initially evolved. In echinoderms IGF probably stimulates growth of the follicles and oocytes, but

the final growth spurt, the one that permits resumption of meiosis in the oocytes and subsequent

ovulation, is delayed until optimal conditions to do so prevail. When the time and place are right the

nervous system releases GSS likely in large amounts to finish the maturation process and induce

ovulation. In vertebrates, growth and the release of IGF has also been brought under control of the

brain but more forcefully by bringing IGF secretion by the liver under control of growth hormone.

Whereas in an early ancestor high plasma concentrations of insulin might have led to secretion of IGF,

this is no longer the case. Here insulin may have evolved to insure that plasma concentrations of

glucose are kept sufficiently low by insuring its absorption by tissues in order to avoid it loss by

excretion. In the three cases these peptides have very different functions, ovulation in echinoderms,

sparing glucose in vertebrates and rescuing interrupted growth in insects. It is plausible then that these

hormones each evolved from a non-local IGF gene duplication and that they are thus not proper

orthologs but evolved by convergent evolution. This hypothesis would explain, why there is no insulin
gene located near the IGF, octinsulin/gonadulin and dilp7 triplet in cockroaches, echinoderms and
hemichordates, even though insulin – and other peptides such as the insect neuroendocrine insulin-like
peptides and GSS - almost certainly evolved from IGF much later.
587

Conclusions

The gene triplication previously reported from arthropods must have occurred in a common
bilaterian ancestor of the deuterostomes and protostomes. The hypothesis that IGF in an ancestral
bilaterian used both a GPCR and an RTK may explain the combination of echinoderm irps and putative
insulin receptors. This hypothesis implies that insulin is not a hormone that evolved before the split
between protostomes and deuterostomes, but that insulin-like peptides evolved independently in
different metazoan clades as miniature copies of IGF capable to activate the RTK but unable to
stimulate the LGR.
596

Acknowledgements

598   This manuscript benefited from the critical contributions of an editor and two reviewers for which I
599 am most grateful. Work like this is only possible because others made their transcriptome and genomic
600 sequences publicly available. I express my sincere gratitude to all of them.

References

604 Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Søren Brunak S, von
605 Heijne G, and Nielsen H. 2019. SignalP 5.0 improves signal peptide predictions using deep neural
606 networks. Nat Biotechnol. 37:420-423, doi:10.1038/s41587-019-0036-z.
607
608 Castellanos MC, Tang JC, Allan DW. 2013. Female-biased dimorphism underlies a female-specific role
609 for post-embryonic ilp7 neurons in *Drosophila* fertility. Development. 140:3915-3926. doi:
610 10.1242/dev.094714.
611
612 Chiba K, 2020. Oocyte maturation in starfish. Cells 9, 476. doi:10.3390/cells9020476.
613
614 Chieu HD, Turner L, Smith MK, Wang T, Nocillado J, Palma P, Suwansa-Ard S, Elizur A, Cummins
615 SF. 2019. Aquaculture breeding enhancement: Maturation and spawning in sea cucumbers using a
616 recombinant relaxin-like sonad-stimulating peptide. Front Genet. 10:77. doi:
617 10.3389/fgene.2019.00077.

Colombani J, Andersen DS, Boulan L, Boone E, Romero N, Virolle V, Texada M, Léopold P. 2015. *Drosophila* Lgr3 couples organ growth with maturation and ensures developmental stability. Curr Biol. 25:2723-2729. doi: 10.1016/j.cub.2015.09.020.

Dagil R, O'shea C, Nykjær A, Bonvin AM, Kragelund BB. 2013. Gentamicin binds to the megalin receptor as a competitive inhibitor using the common ligand binding motif of complement type repeats insight from the NMR structure of the 10th complement type repeat domain alone and in complex with gentamicin. J. Biol. Chem. 288, 4424–4435. doi: 10.1074/jbc.M112.434159.

Daly NL, Scanlon MJ, Djordjevic JT, Kroon PA, Smith R. 1995. Three-dimensional structure of a cysteine-rich repeat from the low-density lipoprotein receptor. Proc Natl Acad Sci U S A. 92:6334-8. doi: 10.1073/pnas.92.14.6334.

Davidson PL, Guo H, Wang L, Berrio A, Zhang H, Chang Y, Soborowski AL, McClay DR, Fan G, Wray GA. 2020. Chromosomal-level genome assembly of the sea urchin *Lytechinus variegatus* substantially improves functional genomic analyses. Genome Biol Evol. 12:1080-1086. doi: 10.1093/gbe/evaa101.

Devi L. 1991. Consensus sequence for processing of peptide precursors at monobasic sites. FEBS Lett 280:189–194. doi: 10.1016/0014-5793(91)80290-j.

Esteban-Lopez M, Agoulnik AI, 2020. Diverse functions of insulin-like 3 peptide. J Endocrinol. 247:R1-R12. doi: 10.1530/JOE-20-0168.

Garelli A, Gontijo AM, Miguela V, Caparros E, Dominguez M. 2012. Imaginal discs secrete insulin-like peptide 8 to mediate plasticity of growth and maturation. Science. 336:579-582. doi: 10.1126/science.1216735.

Garelli A, Heredia F, Casimiro AP, Macedo A, Nunes C, Garcez M, Dias ARM, Volonte YA, Uhlmann T, Caparros E, Koyama T, Gontijo AM. 2015. Dilp8 requires the neuronal relaxin receptor Lgr3 to couple growth to developmental timing. Nat Commun. 6:8732. doi: 10.1038/ncomms9732.

Fisher C, Beglova N, Blacklow SC. 2006. Structure of an LDLR-RAP complex reveals a general mode for ligand recognition by lipoprotein receptors. Molecular Cell 22, 277–283. doi: 10.1074/jbc.M112.434159.

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury R, Zeng Q, Chen Z, Mauceli E, Hacohen N, Gnirke A, Rhind N, di Palma F, Birren BW, Nusbaum C, Lindblad-Toh K, Friedman N, Regev A. 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol. 15;29:644-652. doi: 10.1038/nbt.1883.

Hall MR, Kocot KM, Baughman KW, Fernandez-Valverde SL, Gauthier MEA, Hatleberg WL, Krishnan A, McDougall C, Motti CA, Shoguchi E, Wang T, Xiang X, Zhao M, Bose U, Shinzato C, Hisata K, Fujie M, Kanda M, Cummins SF, Satoh N, Degnan SM, Degnan BM. 2017. The crown-of-

664 thorns starfish genome as a guide for biocontrol of this coral reef pest. Nature. 544:231-234. doi:
665 10.1038/nature22033.
666
667 Haunerland NH, 1996. Insect storage proteins: gene families and receptors. Insect Biochem Mol Biol.
668 26:755-65. doi: 10.1016/s0965-1748(96)00035-5.
669
670 Hoare BL, Bruell S, Sethi A, Gooley PR, Lew MJ, Hossain MA, Inoue A, Scott DJ, Bathgate
671 RAD.2019. Multi-component mechanism of H2 Relaxin binding to RXFP1 through NanoBRET kinetic
672 analysis. iScience. 2019 Jan 25;11:93-113. doi: 10.1016/j.isci.2018.12.004.
673
674 Hsueh AJW, Feng Y, 2020. Discovery of polypeptide ligand-receptor pairs based on their co-evolution.
675 FASEB J. 34:8824-8832. doi: 10.1096/fj.202000779R.
676
677 Humbel RE, 1990. Insulin-like growth factors I and II. Eur J Biochem. 190:445-62. doi:
678 10.1111/j.1432-1033.1990.tb15595.x.
679
680 Imambocus BN, Wittich A, Tenedini F, Zhou F, Hu C, Sauter K, Varela EM. Herédia F, Casimiro AP,
681 Macedo A, Schlegel P, Yang CH, Miguel-Aliaga I, Pankratz MJ, Gontijo AM, Cardona A, Soba P,
682 2020. Discrete escape responses are generated by neuropeptide-mediated circuit logic. bioRxiv preprint
683 doi:10.1101/2020.09.22.307033.
684
685 Ivell R, Alhujaili W, Kohsaka T, Anand-Ivell R. 2020. Physiology and evolution of the INSL3/RXFP2
686 hormone/receptor system in higher vertebrates. Gen Comp Endocrinol. 299:113583. doi:
687 10.1016/j.ygcen.2020.113583.
688
689 Jo J, Oh J, Lee HG, Hong HH, Lee SG, Cheon S, Kern EMA, Jin S, Cho SJ, Park JK, Park C. 2017.
690 Draft genome of the sea cucumber *Apostichopus japonicus* and genetic polymorphism among color
691 variants. Gigascience. 6:1-6. doi: 10.1093/gigascience/giw006.
692
693 Liao S, Nässel DR. 2020. *Drosophila* insulin-like peptide 8 (DILP8) in ovarian follicle cells regulates
694 ovulation and metabolism. bioRxiv. doi: 10.1101/2020.05.02.073585.
695
696 Lin M, Mita M, Egertová M, Zampronio CG, Jones AM, Elphick MR., 2017. Cellular localization of
697 relaxin-like gonad-stimulating peptide expression in Asterias rubens: New insights into neurohormonal
698 control of spawning in starfish. J Comp Neurol. 525:1599-1617. doi: 10.1002/cne.24141.
699
700 Linneweber GA, Jacobson J, Busch KE, Hudry B, Christov CP, Dormann D, Yuan M, Otani T, Knust E,
701 de Bono M, Miguel-Aliaga I., 2014. Neuronal control of metabolism through nutrient-dependent
702 modulation of tracheal branching. Cell 16;156:69-83. doi: 10.1016/j.cell.2013.12.008.
703
704 Menting JG, Gajewiak J, MacRaild CA, Chou DH, Disotuar MM, Smith NA, Miller C, Erchegyi J,
705 Rivier JE, Olivera BM, Forbes BE, Smith BJ, Norton RS, Safavi-Hemami H, Lawrence MC., 2015. A
706 minimized human insulin-receptor-binding motif revealed in a *Conus geographus* venom insulin. Nat
707 Struct Mol Biol. 23:916-920. doi: 10.1038/nsmb.3292.
708
709 Miguel-Aliaga I, Thor S, Gould AP. 2008. Postmitotic specification of *Drosophila* insulinergic neurons
710 from pioneer neurons. PLoS Biol 6: e58. doi:10.1371/journal.pbio.0060058.

Mirabeau O, Joly JS. 2013. Molecular evolution of peptidergic signaling systems in bilaterians. Proc Natl Acad Sci U S A. 110: E2028-37. doi: 10.1073/pnas.1219956110.

Mita M, Yoshikuni M, Ohno K, Shibata Y, Paul-Prasanth B, Pitchayawasin S, Isobe M, Nagahama Y. 2009. A relaxin-like peptide purified from radial nerves induces oocyte maturation and ovulation in the starfish, *Asterina pectinifera*. Proc Natl Acad Sci U S A. 106:9507-12. doi: 10.1073/pnas.0900243106.

Mita M, Matsubara S, Osugi T, Shiraishi A, Wada A, Satake H. 2020. A novel G protein-coupled receptor for starfish gonadotropic hormone, relaxin-like gonad-stimulating peptide. PLoS One. 15:e0242877. doi: 10.1371/journal.pone.0242877.

Mortzfeld BM, Taubenheim J, Klimovich AV, Fraune S, Rosenstiel P, Bosch TCG. 2019. Temperature and insulin signaling regulate body size in Hydra by the Wnt and TGF-beta pathways. Nat Commun. 10:3257. doi: 10.1038/s41467-019-11136-6.

Perillo M, Arnone MI. 2014. Characterization of insulin-like peptides (ILPs) in the sea urchin *Strongylocentrotus purpuratus*: insights on the evolution of the insulin family. Gen Comp Endocrinol. 205:68-79. doi: 10.1016/j.ygcen.2014.06.014.

Prévost M, Raussens V. 2004. Apolipoprotein E-low density lipoprotein receptor binding: study of protein-protein interaction in rationally selected docked complexes. Proteins 55:874-84. doi: 10.1002/prot.20080.

Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-likelihood trees for large alignments. PLoS One. 5:e9490. doi: 10.1371/journal.pone.0009490.

Rholam M, Brakch N, Germain D, Thomas DY, Fahy C, Boussetta H, Boileau G, Cohen P. 1995. Role of amino acid sequences flanking dibasic cleavage sites in precursor proteolytic processing. The importance of the first residue C-terminal of the cleavage site. Eur J Biochem. 227:707–714. doi: 10.1111/j.1432-1033.1995.tb20192.x.

Ruiz-Ramos DV, Schiebelhut LM, Hoff KJ, Wares JP, Dawson MN. 2020. An initial comparative genomic autopsy of wasting disease in sea stars. Mol Ecol. 29:1087-1102. doi: 10.1111/mec.15386.

Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream MA, Barrell B. 2000. Artemis: sequence visualization and annotation. Bioinformatics 16:944-5. doi: 10.1093/bioinformatics/16.10.944.

Sea Urchin Genome Sequencing Consortium. 2006. The genome of the sea urchin *Strongylocentrotus purpuratus*. Science 314: 941–952.

Semmens DC, Mirabeau O, Moghul I, Pancholi MR, Wurm Y, Elphick MR. 2016. Transcriptomic identification of starfish neuropeptide precursors yields new insights into neuropeptide evolution. Open Biol. 6:150224. doi: 10.1098/rsob.150224.

757 Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M,
758 Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of high-quality protein multiple
759 sequence alignments using Clustal Omega. Mol Syst Biol. 7:539. doi: 10.1038/msb.2011.75.
760
Simakov O, Kawashima T, Marlétaz F, Jenkins J, Koyanagi R, Mitros T, Hisata K, Bredeson J,
Shoguchi E, Gyoja F, Yue JX, Chen YC, Freeman RM Jr, Sasaki A, Hikosaka-Katayama T, Sato A,
Fujie M, Baughman KW, Levine J, Gonzalez P, Cameron C, Fritzenwanker JH, Pani AM, Goto H,
Kanda M, Arakaki N, Yamasaki S, Qu J, Cree A, Ding Y, Dinh HH, Dugan S, Holder M, Jhangiani SN,
Kovar CL, Lee SL, Lewis LR, Morton D, Nazareth LV, Okwuonu G, Santibanez J, Chen R, Richards S,
Muzny DM, Gillis A, Peshkin L, Wu M, Humphreys T, Su YH, Putnam NH, Schmutz J, Fujiyama A,
Yu JK, Tagawa K, Worley KC, Gibbs RA, Kirschner MW, Lowe CJ, Satoh N, Rokhsar DS, Gerhart J.
2015. Hemichordate genomes and deuterostome origins. Nature. 527:459-465. doi:
10.1038/nature16150.
770
771 Smith MK, Chieu HD, Aizen J, Mos B, Motti CA, Elizur A, Cummins SF. 2019. A Crown-of-Thorns
772 Seastar recombinant relaxin-like gonad-stimulating peptide triggers oocyte maturation and ovulation.
773 Gen Comp Endocrinol. 281:41-48. doi: 10.1016/j.ygcen.2019.05.009.
774
775 Tensen CP, Van Kesteren ER, Planta RJ, Cox KJ, Burke JF, van Heerikhuizen H, Vreugdenhil E. 1994.
776 A G protein-coupled receptor with low density lipoprotein-binding motifs suggests a role for
777 lipoproteins in G-linked signal transduction. Proc Natl Acad Sci U S A. 91:4816-4820. doi:
778 10.1073/pnas.91.11.4816.
779
780 Vallejo DM, Juarez-Carreño S, Bolivar J, Morante J, Dominguez M. 2015. A brain circuit that
781 synchronizes growth and maturation revealed through dilp8 binding to Lgr3. Science. 50:aac6767. doi:
782 10.1126/science.aac6767.
783
784 Veenstra JA. 2000. Mono- and dibasic proteolytic cleavage sites in insect neuroendocrine peptide
785 precursors. Arch Insect Biochem Physiol. 43:49-63. doi: 10.1002/(SICI)1520-
786 6327(200002)43:2<49::AID-ARCH1>3.0.CO;2-M.
787
788 Veenstra JA. 2020a. Gonadulins, the fourth type of insulin-related peptides in decapods. Gen Comp
789 Endocrinol. 296:113528. doi: 10.1016/j.ygcen.2020.113528.
790
791 Veenstra JA. 2020b. Arthropod IGF, relaxin and gonadulin, putative orthologs of *Drosophila* insulin-
792 like peptides 6, 7 and 8, likely originated from an ancient gene triplication. 10: e9534. doi:
793 10.7717/peerj.9534.
794
795 Veenstra JA, Rombauts S, Grbić M. 2012. *In silico* cloning of genes encoding neuropeptides,
796 neurohormones and their putative G-protein coupled receptors in a spider mite. Insect Biochem Mol
797 Biol. 42:277-295. doi: 10.1016/j.ibmb.2011.12.009.
798
799 Veenstra JA, Leyria J, Orchard I, Lange A. 2021. Identification of gonadulin and insulin-like growth
800 factor from migratory locusts and their importance in reproduction in *Locusta migratoria*. Front
801 Endocrinol., *in press*.
802

803 Warner JF, Lord JW, Schreiter SA, Nesbit KT, Hamdoun A, Lyons DC. 2021. Chromosomal-level
804 genome assembly of the painted sea urchin *Lytechinus pictus*, a genetically enabled model system for
805 cell biology and embryonic development. Genome Biol Evol:evab061. doi: 10.1093/gbe/evab061.
806
807 Yang CH, Belawat P, Hafen E, Jan LY, Jan YN. 2008. *Drosophila* egg-laying site selection as a system
808 to study simple decision-making processes. Science 319: 1679–1683. doi: 10.1126/science.1151842.
809
810 Yasui N, Nogi T., Takagi J. 2010. Structural basis for specific recognition of reelin by its receptors.
811 Structure 18, 320–331. doi: 10.1016/j.str.2010.01.010.
812
813 Zhang X, Sun L, Yuan J, Sun Y, Gao Y, Zhang L, Li S, Dai H, Hamel JF, Liu C, Yu Y, Liu S, Lin W,
814 Guo K, Jin S, Xu P, Storey KB, Huan P, Zhang T, Zhou Y, Zhang J, Lin C, Li X, Xing L, Huo D, Sun
815 M, Wang L, Mercier A, Li F, Yang H, Xiang J. 2017. The sea cucumber genome provides insights into
816 morphological evolution and visceral regeneration. PLoS Biol. 15:e2003790. doi:
817 10.1371/journal.pbio.2003790.
818

Figure 1

Domains of insulin/IGF-related peptides.

Human insulin and IGF and Drosophila dilp7 are aligned and the different domains that are recognized in the precursors of these peptides are indicated. In insulin the domain borders are the convertase cleavage sites that are hihglighted in red. The A- and B-domains of insulin correspond to the A- and B-chains of insulina and the C-domain to the connecting peptide. Although IGF consists of a single protein chain due to its strong sequences similarity to insulin the A- and B-domains correspond to homologous regions of those domains in insulin, while the C-domain is the sequence between the A- and B-domains. In insulin there is only a single amino acid residues after the last cysteine residue, but in IGF there is a longer sequence, that has been called the D-domain. The IGF precursor is cleaved by furin in the Golgi apparatus and the sequence that is removed has been called the E-domain. Dilp7 is only known from nucleotide sequences, it is unknown how the precursor is exactly processed. Nevertheless, the presence of putative convertase cleavage sites, highlighted in red, suggests the presence of A-, B- and C-domains quite similar to those in insulin. However, unlike insulin or IGF, the putative B-chain of dilp7 has a long N-terminal extension that I propose to call the F-domain. The latter is well conserved in dilp7 orthologs from other bilaterians (Fig. 4).

Figure 2

Sequences of selected ambulacrarian IGF.

Partial IGF sequences from selected ambulacrarians are illustrated to show their sequence similarity. The A-, B- and C-domains of the insulin core are aligned, but not the putative D- and E- domains, as their amino acid sequence is only conserved in closely related species (Fig. S1). Not aligning D- and E- domains allows the visualization the context of putative convertase cleavage sites. None of the arginine or lysine residues conform to a typical arthropod or vertebrate convertase cleavage site. Although the sequence of the latter part of the IGF precursors is not well conserved, all of them are rich in positively charged amino acid residues. Conserved cysteine residues are indicated in red, conserved amino acid residues are highlighted in black and conserved substitutions in grey. The arginine and lysine residues in the D- and E- domains are highlighted in blue.

```
Anneissia              QPNHVRRYCGSELTSELERRCALKGGFNSPKKQESP--------------------
Strongylocentrotus-1   ---SFPLLCGQELVKAVAAVCNDR-GYYGQPS--------------------
Strongylocentrotus-2   ----SFRLCGRELADALAVVCKGR-GYYIDDSEI--------------------
Apostichopus-1         ---SGQRYCGEALLEAMAYVCGDR-GYFGVTSGIRGRS--------------------
Apostichopus-2         ----AQRYCGTNLADALRIVCADR-GYYTQ----KGAF--------------------
Acanthaster-1          ---AVIQVCGNDLLDALKSVCGDR-GFYSPPPGYS--------------------
Acanthaster-2          ----WQRICGEQLVETVSIVCNTR-GFYSHR-E--------------------
Ophiothrix-1           ----WQRLCGTFLVDVVSQVCGER-GTYADDSTHDLRKRSLDNFVDRGKMYSAGISKT
Ophiothrix-2           -----SYLCGSQIVQMMAVVCEGR-GYYYTEGSTGQICN--------------------
Saccoglossus           ----WDKLCGRTLVDVLALICNGR-GYNSGSPKK--------------------
Ptychodera             ----WDRLCGRSLADMLALVCHGR-GYYTDVSRQQ--------------------
Schizocardium          ----WDKLCGRTLADMLSLVCHGR-GYYTDVSKHR-IR--------------------

Anneissia              ----------------YLWNQEKRIEATS-----------TERAKEINIVYECCHSACTEAFLDS
Strongylocentrotus-1   ----KRSAGIFELETRAKTFLKSGISRGETRRSKRGARTGLIVTECCLNRCSVSHLES
Strongylocentrotus-2   ----AQKDSPIVPHHVASSFLGSSS-ASAHSRQRRRVRTGQIVNECCDKECSNNIMES
Apostichopus-1         -----VSRSPFLTEERANSFLTNE-------RTRKTRRTGRIVTECCDNPCSQRNLES
Apostichopus-2         --------PPVIPRQRASSFLTTDS---ENHHPRRLRRGTGIVTECCEKACDREVLET
Acanthaster-1          ----------------------------RRTPATQTGIATRCCISYCETSVLEK
Acanthaster-2          ----KREAELFQDERTAKSFLGTHIG------SRQRRRTGRIATECCEKVCSYDIVES
Ophiothrix-1           SDTNNAESNPFRGESEAKSFLGL---------SRQKRRTGRIVHECCNNICNYRIIES
Ophiothrix-2           IVYSFLTDPPLLTERQAKTFT-KSG----------RTRTGKIVTECCDNHCDMQIIES
Saccoglossus           ---KVKRESPFRSGMEANDFFGNIS-SKEKRRQRRRSGSGKIVDECCHQACDYTTLES
Ptychodera             -KSRRARETVFATQEEANGFFGVG---S--GRTKRRRGSGLIVVECCDKICDYSTIES
Schizocardium          ----LPRETIFQTQEDAHKFFGASV-FG--ERTQRRRGSGLIVAECCEKSCDYATIES

Anneissia              FCLSSNKDEDTTVESDTVETTTITGKKRKPTRKPKNPLKSRKKPKGSSEINSEQSASN
Strongylocentrotus-1   YCNPLPPDAVHDAEVHIRLEKSAEEDADEGRPQDGPSQLDTATGTVPETEMSETRGRV
Strongylocentrotus-2   YCNRRTPEVPPESAISENPSEEITEDSTLRTDGESTEIRTDTNPATNLEVPSPDANTP
Apostichopus-1         YCNVATTQTTEIPTELTTEGTTTEPAASPRRNSRNIEADGTAAAAGGGSGQGNGRGKG
Apostichopus-2         YCNPHVTPTLALASLVTSLMTKSPTPPSSEPSSSSSSSSSRNEDKFPMTDNALGEDY
Acanthaster-1          YCNPPSTSQSQTAAAPPRITTTPDERRANEIVVDETGQTGNTNSQMLRGGNAMGAGSR
Acanthaster-2          YCNPWPVVEDRDDPMLAPVAPGRVRQDKSADADLLLRPDIAEISEDKSSLLRQAAEKD
Ophiothrix-1           YCNPWPTTTTTATTTQSPEPLPNEQEGGYLTDEIQMKHITRGQEDNSVDLLSEGSDLR
Ophiothrix-2           YCAPLQEGQVKFTSRNLDFVNEANNVVEEPSVRVPQQVEENKIVEETVPYVSQAIVGE
Saccoglossus           YCAPLPEGVVADDSLKRFLSQSFGNDFKDTANEDKLEIVTVVRPSHDEMDGTETRIED
Ptychodera             YCAPWPKDIDPAKKIEGFKEGTWEEEDYHRKYHPESVEQPNANPEEPTPEPTTTDLDK
Schizocardium          YCAPWPKDMDPALRFAGFKYGSWEDEDYYRKYHPEEFTQPDLTTWIASTDSANHDEHF

Anneissia              TELPSQTEPTKDKNGRGDKNKKDKCNKKSRDKNNKPCRRKSRKDSRRKNRKKNKKNKP
Strongylocentrotus-1   RIDAVEKVLSERLIPTSTTGSPSPSRKKPRKDKSERRNSSREAKQARREERRRNRER
Strongylocentrotus-2   DATATSDVEQPRSDNTTAVEKPRKKDNGKGKNSSLESSTKKNRTSKGMSKEDRRRIAS
Apostichopus-1         KGRHGKGNRREQDQTVDVTSAETEGNTEPPRPNQQENNEDNRGTPDQSEERPRDRCRG
Apostichopus-2         EGPTNEGPLTSGEPTPTENRISNGPRGPSSNASSLELPTRTSTATTNSSRIVTEGAHL
Acanthaster-1          ANGTKAPPTEVVDGRSDDDDDAAGEINTSERVGSLTEPDEETGRDVATNRPHKTHSKE
Acanthaster-2          EPIDDLDTLENEYADGGNVMLQAREGVKEEGAELGKEMEEGGEKMPFPEVVPTKKRRR
Ophiothrix-1           DETPMPTGRATGGRSPQLGHDQSADVDSGFHRFGDIGAEVEESDSTIDNPSSDR----
Ophiothrix-2           TTSDGFDWDSDVSETHNILNGGGDMYINDQPRDDPSENDEEHIPKEKSLIKQINVTES
Saccoglossus           NEHVTPPTKPDVITETSSLILDDINVNKQIISSNTSVEVKSKAGNTKPKREKKDRDNS
Ptychodera             SLESESRDNVDIEESRSKDKSAESEESVTENDDLKSEETNDNQDEYSREEYLGRDKGP
Schizocardium          TTATPSEETTSSENGEHIKTVNKEDDTQVTGTHELEKDHKFAEYWMNEDKMSRKHKKA

Anneissia              TRRPKRWDSETTTLSDFLFSQRFLQRAYDSDIDVDEITTEFGVVPEEDISSSLSEEEL
Strongylocentrotus-1   SSGGRSRNGRRKDKDNDRASRAKRHGLNLWRNMFSDKFFSDIPGLENQPNLHPVNGRA
Strongylocentrotus-2   DERRASRERKKELSRERRKRLKLQQRKDKKKKKRLESAERNRGTDHMGLSEDSTLLAR
Apostichopus-1         SKKKGKCRNQDRVEEEPTQDRIEEEPTTSREVSSTDGLSRERGSGSGGGRNRGNGKGK
Apostichopus-2         AESSDQSSLEDTDSESEAPTKGASHERTRKKQKKQRTRPPKKPKKLSRERKERKKKKE
Acanthaster-1          RSKNRTSKSERRRRRTNRRRSSERRMLSSERKREDATRKLRRKEQRLSRKQPHSNKR
Acanthaster-2          VEGRRSRENSRDRNGKSEGKSKRKSGSREGGRSFRRRGKNSRRKKGRDGRERSKRWEA
Ophiothrix-1           ----------------------------------------------------
Ophiothrix-2           QTDEILDEDTDSENSESKLSIGATKGNSGRTKPTHRPTSRHSRRKNSREKKKNKTKTR
Saccoglossus           SSKKRSHPKPKKSRKKQRLQRIRKKARGTRKTKLRHVKVKSKSTPIQQIETTTTMKPF
Ptychodera             FRKHKKAPTKKLFKEKKLSEDNKKKKSRAKSKNSTKVKPTYVSSMTTSDEETLTRPQG
Schizocardium          KRLMKERRARKSRTKKGNNTKSKSILGKLVENMETESPTNWQRDGTDERWWTVEDPHF
```

Figure 3

Sequences of selected echinoderm GSS.

Sequences of selected echinoderm GSS. Sequence alignment of a few echinoderm GSS showing relatively conserved A- and B- domains of the insulin core sequence and likely KR convertase cleavage sites that can be expected to be cleaved by neuroendocrine convertase as well as a few potential furin sites. Conserved cysteine residues are indicated in red, conserved amino acid residues are highlighted in black and conserved substitutions in grey. The arginine and lysine residues that form likely – or possibly in the case of *Apostichopus* GSS-2 - part of a convertase site are highlighted in blue. For the alignment of a larger number or echinoderm GSS sequences see Fig. S3.

Figure 4

Sequences of selected dilp7 orthologs.

Sequences of selected dilp7 orthologs. Sequences of *Drosophila* dilp7 and several ambulacrarian orthologs illustrating well conserved sequences, not only in typical insulin core of the peptides, but also in the F-domain (underlined in blue). Note that the sequence conservation of these peptides is stronger than in the IGFs or GSSs (Figs. 1,2). Conserved cysteine residues are indicated in red, conserved amino acid residues are highlighted in black and conserved substitutions in grey. Likely convertase cleavage sites have been highlighted in blue. Sequences are from Spreadsheet S1 and Veenstra 2020b, a comparison of a larger number of sequences is presented in Figs. S5 and S6.

Figure 5

Sequences of selected ambulacrarian octinsulins.

Sequences of selected ambulacrarian octinsulins. Sequence alignment of a number of octinsulin sequences show that these sequences all have typical neuroendocrine convertase KR cleavage sites, suggesting these precursors are processed by enteroendocrine and/or neuroendocrine cells. Conserved cysteine residues are indicated in red, conserved amino acid residues are highlighted in black and conserved substitutions in grey. Likely convertase cleavage sites have been highlighted in blue. Sequences are from Spreadsheet S1, a comparison of a larger number of sequences is presented in Figs. S7 and S8.

```
Anneissia          ------------------ARDWYC-GNAADTLKEFCQSCYASKRAHN--ALSLP
Strongylocentrotus ------------------QSWHC-GRAAQTIMGMCNSCYASHDKRS-------
Apostichopus       ------------------SWYC-GSAPETVRAICDGCYAGGIHTR--AFKRS
Acanthaster        ------------------DSWYC-SDVYSTVQSLCDSCYAGFDKRT------
Ophiothrix         ------------------NQWFC-SPVFTMLQSMCGSCYAGVDKRS------
Saccoglossus-1     ------------------MSRNWHC-GRPVETMHEVCQGCYAGHVRPR------
Ptychodera-1       ------------------LRREWHC-GRTVETMQGICRGCYAQPSERS------
Schizocardium-1    ------------------FTKHWHC-GRIVDTMRAICDGCYASPTARD------
Saccoglossus-2     RPSGSVD------------DVLTGC-KRKLLLVDQICAGCYAPPDIINNVNFDLT
Ptychodera-2       RPRNQGD------------D--VFC-SRTYSMVESVCDGCYATTQDSSPKSESAM
Schizocardium-2    RPNRPSE------------D--VRC-RKTIHMVLKLCNGCLAPIESEVENN--TI
Saccoglossus-3     ---LPVDDTTTGVQVDVRDSRLWLC-GRLVEDLRALCRGCYAG---P--------
Ptychodera-3       -DLLPGNGDY---SNSAKGRRDLDC-THLVESMSLICRGCYATDQGV--------
Schizocardium-3    ------------QTSADRPRKWHCRKSVPEILSGVCRGCYAEPLQPP--------

Anneissia          SIKA-----KKDGMFLTKEGASGYLEAKRTRL--------FSS-LHLNHRQHETT
Strongylocentrotus ISKPSY---TPAKPFLHKRNAVHFLRTTKREIESRPSMGDTAIEVAVERRSTGNR
Apostichopus       SSDIIS---LYKDPFLKKSNALNFLLPRSHTP-------S-----SLIKRGIRRS
Acanthaster        NSITRP---I-EEPFVERKNAVDFFKRGTAR--------G------------TRR
Ophiothrix         DNSDTLSQKQSLDAFIQKEVAYSFIKRTSVG--------DT----FLRNARNTHH
Saccoglossus-1     NTRS-----VDGVQAFISRRDANMFTKGMSPD-------------V----KRAIDG
Ptychodera-1       T--N-----EAERQAFIGKEEASSFTKSVP----------------------TIVKR
Schizocardium-1    V--G-----SVKGLPFLKKHEASTFTRTRS----------------------SVQKR
Saccoglossus-2     GK-------------------------GPGY---------------EEEIVKN
Ptychodera-2       NE-------------------------SVQI---------------QDEDLRE
Schizocardium-2    SN-------------------------ETPP---------------DQDVVTA
Saccoglossus-3     --------------DISKREASKFMQFNA-----------------HTIRQSR
Ptychodera-3       --------------TVNKRLASSFIPTTTTP------------R-----NKIQR
Schizocardium-3    G-----------KRDILNKQEASLFLRSP-----------------NGGDE

Anneissia          NFVTECCYNPCSSFEMIKYCCPTRQIELHNRNPNSSEDK--------------
Strongylocentrotus GFIHECCNKFCDPGEMVLYCCEKRQIEWAQFHNLLKA--------------
Apostichopus       GFIGECCEKNCEIREMVFYCCAEKQREYASFFPEIFRNRIRHT--------
Acanthaster        GIVDECCHRQCAVSEMMLYCCEQKQREYYTFVGWLKRR------------
Ophiothrix         GLIDECCTQQCDTGEMILYCCQERQREWHIMMGLYNK--------------
Saccoglossus-1     GLIEECCYSQCSLTHMITYCCAEVQNEFQVFINILGNTDESSENDGDDGEESSSV
Ptychodera-1       GLLEDCCYRRCNLQKMMTYCCAERQRELNNFFSLLNQKDNGST-----------
Schizocardium-1    GIIEHCCNHHCSFTELLIYCCEERSEEFYSFIGLLRMDDEDTDASLEKGNGDVEA
Saccoglossus-2     QIKEACCKEYCPLPKIIEFCCDERQQEFHQFMSSFASTEE--------------
Ptychodera-2       KIRDCCNRRCTIHKMMQFCCEARRNEFHKFLALMGNTDN--------------
Schizocardium-2    KVREVCCDNYCSLDKIIEFCCEDLQQEFRQFMSFVSNSK--------------
Saccoglossus-3     GIIEDCCYHTCPTERKIQYCCFEVQAQYRLFMESAI-----------------
Ptychodera-3       GIIDECCRNRCSVERKIQYCCDEIQKEFAFFFSLFGS----------------
Schizocardium-3    LIINECCLRTCTVIEKIHYCCREKQIELYIIIQSAPWLVDNQR----------

Anneissia          ------------------------------------------------------
Strongylocentrotus ------------------------------------------------------
Apostichopus       ------------------------------------------------------
Acanthaster        ------------------------------------------------------
Ophiothrix         ------------------------------------------------------
Saccoglossus-1     HED---------------------------------------------------
Ptychodera-1       ------------------------------------------------------
Schizocardium-1    EDNTDLDQGDDDDNTQNNEGGSIDVVVERDVGDVLLKTDKSKP
Saccoglossus-2     ------------------------------------------------------
Ptychodera-2       ------------------------------------------------------
Schizocardium-2    ------------------------------------------------------
Saccoglossus-3     ------------------------------------------------------
Ptychodera-3       ------------------------------------------------------
Schizocardium-3    ------------------------------------------------------
```

Figure 6

Sequence comparison of selected ambulacrarian multinsulins and dilp7 orthologs.

Sequence comparison of selected ambulacrarian multinsulins and dilp7 orthologs. Three different sets of sequences are compared. The top five sequences are dilp7 orthologs, the next five are multinsulins having three disulfide bridges and the last five multinsulins having four disulfide bridges. Note that although the multinsulins and the dilp7 orthologs share some sequences similarity this does not include the F-domain. Like the octinsulins these sequences all have typical neuroendocrine convertase KR cleavage sites, suggesting they are processed by enteroendocrine and/or neuroendocrine cells. Conserved cysteine residues are indicated in red, conserved amino acid residues are highlighted in black and conserved substitutions in grey. Likely convertase cleavage sites have been highlighted in blue. Sequences are from Spreadsheet S1, a comparison of a larger number of sequences is presented in Figs. S9 and S10.

```
Anneissia          --------------------------------------------LRDYSDRSHNDWARVWTVE
Antedon            --------------------------------------------LQDYSDRSHHDWARVWTVE
Strongylocentrotus ------------------------------EKFCNCMVLPELTMEDYEDRTPEEWRESWNMD
Apostichopus       ---------------------------------TLQELNSRTQPSWEQLWIVE
Acanthaster        ------------------------------APHLPVEQWNSRSKADWVKLWNTE
Ophiothrix         ------------------------------KITDYSSRTKADWQRLWLTE

Amphiura-1         ------------------------------------------D----WVG-
Ophioderma-1       ---------DEIGAMKSQNEEPDEHDS----HIGTTQQPVINQRSGTMEKR----WRG-
Pisaster-2         ----------------KLNMKPDEVLSVE-SEHCSAVKEFRQVLADNPGVV---KRS-
Asterias-1         ---------------------ELRS-----QDGLMREVRQALARNGRHF----RAA-
Patiria-1          ---------APPANDGIEVLTGGKIE-----EDDFEGDDFDVKNKNQYYS----RSG-
Acanthaster-1      -----------------------HPDADAKH-GGNL-------A-

Ophioderma-5       NSLTRDADDAPVAMDDKPTSPRPIDIIKRSRHLRCAQFKDMALE-DHFRR-LVGKER-
Asterias-3         ----------------------ELN-----VASCQQLADQVKS-EGKKR----WDG-
Pisaster-1         ----------------------TELD-----EASCKQLADQVKS-EGKKR----WTG-
Acanthaster-3      ----------------------DKTPS-----TASCEALAAQIST-EGKKR----WDG-
Patiria-7          ----------------------DKAPS-----AATCEALAATVKA-EGKKR----WDG-

Anneissia          SM--RQCHENLRE--MVHVSCRNDPRKISS-----------KR-SIFIPRNEATGFLSR
Antedon            SM--RQCHEDLRE--MVHISCHNDPRKITS-----------KR-SIFIPRNEATGFLSR
Strongylocentrotus TLR-TCVGPQLQR--VGELACINDPRKTIVVKR-----SNSDRDLFLPAKLAKAFLHY
Apostichopus       NVPTVDCTVDAVQ--LHIIACSNDVYKDHEGSRRRRFSDKRRRSIFLNFSEANNFLAK
Acanthaster        RHV-NTCNEALLP--VWDVACQNDIRKITK---------RMGREFLNEWTAKNFLAG
Ophiothrix         SH--QKCNEDLLP--LWKIACTYDIRKIN ---------KRA-PEFVEDSEAKAFLIG

Amphiura-1         -FRT-ICDPPFTPSLYIEDFCGVTV---KR------E------YEATDPLGFLKM
Ophioderma-1       -YVY-ICEPQLTR--LKNVICNPASVKRS---------DTAGLEFLTEHQAKRFLMQ
Pisaster-2         -PRNYWCNTALSQ--RKTALCGCT----HH---------TLRDDDFMEEKKDATNFLE
Asterias-1         -EDHIYCGVVLEQ--NRESVCGV--IPG-----------SKRNLFVRKEAASEFLE-
Patiria-1          -PRFRVCGTTTHS--WSSFVCHPSGLI-HH---------KRDNDEFLSAGEANTFLMS
Acanthaster-1      -YRANYCGATLYE--KVRETCHA-----VR---------GVSNQEFLDSKDASTFLFG

Ophioderma-5       -VFAKYCSPTPQT--VMDNYCQCDVVPRSI----------DDKRAFVDKSSAKSFLNH
Asterias-3         -PSHKFCGETLNE--KRYAYCTCGLVPRKR---------ELELSEFLNRGKANGFLSA
Pisaster-1         -SWHTFCGETLNE--KKYAYCTCGLVPRKR---------ELDLSEFLNRGKANGFLSA
Acanthaster-3      -PSHTFCGEDLNE--RRNAYCNCQVVPRKR---------ELDLSEFLPSGKANAFLSG
Patiria-7          -PAHTFCGETLTE--KANAYCHCQVVPRKR---------ELELSEFLTPVKANSFLSG

Anneissia          FL---R-T-RRPSELHEDCCLDSRGCTWEEVAEIACINNRRRMHRPGSPVGR------
Antedon            FL---R-T-RRPSELHEDCCLDSRGCTWEEVAEIACINNRRRMHRPGSPVGR------
Strongylocentrotus RHRKDS-R-RRRVGKDEECCAEAQGCRWEELGEYCTLH-TRAYHQSGEQP--------
Apostichopus       TKRTHS-RVRRTTTFSTECCD--KLCIWEEVGEYCWH--SRVYH------------
Acanthaster        S------K-RRKRGLNEECCHEDLGCVWEEVAEYCVMH-GREKHEDGSPVRGKPGRRR
Ophiothrix         P------R-RHKRGLNEECCHESKGCVWEEIGEYCRMH-SRASHVDGRIDSR------

Amphiura-1         RGMVKRLNPNWRELMSEECCY--ESCTSEEIKELC----------------------
Ophioderma-1       NKR----IWGW-GGLSEECCN--EGCSVEEIDEVC----------------------
Pisaster-2         RGIEKR-------SLSEECCH--EGCYYEEVYEVC----------------------
Asterias-1         R--GVQ-------GLAEECCG--EGCTIEEISESC----------------------
Patiria-1          EGVGKR-------GMHEECCH--EGCWWEEMLEHC----------------------
Acanthaster-1      RGIEKR-------GLHHECCT--EGCSIEEIYESC----------------------

Ophioderma-5       RAS-TR-------SLDEECCN--EGCNLEEIVELSKTMCSSS---------------
Asterias-3         RNVQKR-------SLSEECCH--EGCYWEEIEEVC----------------------
Pisaster-1         RNLQKR-------SLSEECCS--EGCYWEEIEEVC----------------------
Acanthaster-3      RRITKR-------SLSEECCH--EGCYWEEIEEVC----------------------
Patiria-7          RNIAKR-------SLTEECCH--EGCYWEEIEEVC----------------------
```

Figure 7

Position of introns in ambulacrarian irp genes

Schematic representation of the location of the cysteine residues, indicated as purple rectangles, and introns, represented by green T's, in the coding sequences of the various types of ambulacrarian insulin-like genes. Numbers indicate the phase of each intron. All genes share the typical phase 1 intron present in insulin-like genes, whereas dilp7 and multinsulin genes also share a phase 2 intron. Signal peptides indicated as interrupted bars.

Figure 8

Synteny of ambulacrarian irp genes.

Schematic representation of the relative localization of different irp genes in several arthropod and ambulacrarian genomes. Arrow heads indicate transcription direction of the various genes, the numbers below the line indicate the number of nucleotides between the coding regions of adjacent genes in kilo base pairs. Note that the relative organization of the two insects – the cockroach *Blattella germanica* and the stick insect *Timema crisitinae* – is the same as in the hemichordate *Saccoglossus kovalewskii* and remarkably similar to that of the sea urchin *Strongylocentrotus purpuratus* and the sea cucumber *Holothuria scabra*. In the spider *Pardosa pseudoannulata* and the sea cucumber *Apostichopus japonicus* some of the genes are also next to one another. However, in the sea stars *Acanthaster planci* and *Pisaster ochraceus* synteny has been lost. Arthropod data from Veenstra, 2020b.

Pardosa	
Blattella	
Timema	
Saccoglossus	
Strongylocentrotus	
Holothuria	
Apostichopus	
Pisaster	
Acanthaster	

Legend:
- dilp7 ortholog
- GSS
- gonadulin
- octinsulin
- IGF
- multinsulin

Figure 9

Radial sequence similarity tree of ambulacrarian irps.

The five different types of irps are clearly separated from one another. Note that the GSSs are similar to IGFs and seem to be related to them, while the multinsulins are most similar to the dilp7 orthologs. Echinoderm branches are in black, hemichordate branches in red. More extensive sequence comparisons and sequence trees are the supplementary data (Figs. S1-S10). All sequences are from Spreadsheet S1.

Figure 10

Phylogenetic tree of LGRs.

Phylogenetic tree constructed from the transmembrane regions of ambulacrarian LGRs that are putative receptors for irps. A few human and insect sequences have been added for comparison. The insert at the top shows the same data to which the glycoprotein LGRs have been added and where characteristic ligands for each branch have been identified. Numbers in blue indicate the apparent probabilities as determined by Fasttree. For details of the glycoprotein LGRs see Fig. S11.

Figure 11

Ectodomains of ambulacrarian LGRs.

Ectodomains of ambulacrarian LGRs. Schematic representation of the various domains of the putative receptors for ambulacrarian insulin-related peptides. Each green circle symbolizes an LDLa repeat and each purple rectangle an LRR repeat, while the yellow oval indicates the seven transmembrane regions. The top representation corresponds to the gonadulin and dilp7 receptors (Figs. S11,S12). Note though, that the latter are somewhat variable, notably in the sea stars of two species of the *Patiria* genus and *Acanthaster planci* those receptors have two LDLa repeats (for details see Fig. S12). The bottom representation corresponds to the GRL101 receptors (Fig. S13).

Figure 12

How echinoderm irps may have evolved.

A represent an early metazoan in which an arch irp is a ligand for both an LGR and an RTK. B represents an early protostome or deuterostome that has three irps, an IGF and a dilp7 ortholog as well as gonadulin/octinsulin ortholog that evolved from local gene duplication from the arch irp. All three of these ligands each each their own LGR and at least two of them, IGF and the dilp7 ortholog, can also activate the RTK. C represents the Asterozoa where the dilp7 gene got duplicated and yielded several multinsulin genes which are represented here as one. The Asterozoa also have one or two GSS's that evolved earlier during echinoderm evolution. Both multinsulins and GSS's act exclusively through the RTK. Closed arrows indicate gene duplication events and interrupted arrows show ligand-receptor interactions. The question mark conveys uncertaintity with regard to whether or not the gonadulin/octinsulin peptides are able to activate the RTK.

LGR

IRP

RTK

A

dilp7   IGF   gonadulin/octinsulin

?

B

GSS

?

C   multinsulin