1   **Extensive transgressive gene expression in a homoploid hybrid species**

Homa Papoli Yazdi[1][*], Melissah Rowe[2], Mark Ravinet[3], Glenn-Peter Sætre[4], Caroline Øien

Guldvog[4], Fabrice Eroukhmanoff[4], Alfonso Marzal[5], Sergio Magallanes[5] & Anna Runemark[1]

[1] Department of Biology, Lund University, SE-223 62 Lund, Sweden

[2] Department of Animal Ecology, Netherlands Institute of Ecology (NIOO-KNAW), 6700

AB, Wageningen, The Netherlands

[3] School of Life Sciences, University of Nottingham, Nottingham, UK

[4] Department of Biosciences, Centre for Ecological and Evolutionary Synthesis, University of

Oslo, PO Box 1066, N-0316, Oslo, Norway

[5] Departamento de Anatomía, Biología Celular y Zoología, Universidad de Extremadura, E-

506071, Badajoz, Spain

[*]Corresponding author: E-mail: Homa.Papoli_Yazdi@biol.lu.se

Abstract

Hybridization is increasingly recognized as an important evolutionary force contributing

novel variation for selection to act on. While mis-expression in F1 hybrids is well

documented, how gene expression evolves in stabilized hybrid taxa and contributes novel

variation remains an open question, especially for hybrid species without an increase in

ploidy. As gene expression evolves in a stabilizing manner, break-up of co-evolved *cis*- and

trans-regulatory elements could lead to transgressive patterns of gene expression in hybrids.

Here, we address to what extent gonad gene expression has evolved in an old homoploid

hybrid, the Italian sparrow *Passer italiae*. Through comparing the gene expression of parental

species and F1 hybrids to that of the Italian sparrow, we find evidence for strongly

transgressive expression in the Italian sparrow, with 22% of the testis genes exhibiting

expression patterns outside the range of both parent species, compared to only 0.37% in the

F1s. In contrast, Italian sparrow ovary expression was similar to that of one parent species,

the house sparrow (*P. domesticus*). Moreover, the Italian sparrow testis transcriptome is 26

times as diverged from those of the parent species as the parental transcriptomes are from

each other, despite being genetically intermediate. This highlights the potential for regulation

of gene expression to produce novel variation following hybridization. Genes involved in

mitochondrial respiratory chain complexes and protein synthesis are enriched in the subset

that is over-dominantly expressed in Italian sparrow testis, suggesting that selection on key

functions have molded the hybrid Italian sparrow transcriptome.

Keywords: Hybridization, gene expression, transgressive expression, testis, mito-

nuclear genes

Introduction

Hybridization is an important evolutionary process, contributing to novel genetic variation, that natural selection can act on (Mallet 2005; Abbott, et al. 2013; Runemark, et al. 2019; Taylor and Larson 2019). While divergence in parental genomes brings together novel allelic combinations in the hybrid, which may lead to a reduction in hybrid fitness (Dobzhansky 1936; Muller 1942), it is now clear that hybridization is frequent (Marques, et al. 2019; Runemark, et al. 2019; Taylor and Larson 2019). Hence, we can no longer study hybridization as an isolated phenomenon, and the processes at play in hybrids should be integrated in our general understanding of how novel variation can arise. While there is ample evidence for admixed hybrid genomes (Runemark, et al. 2019) and transgressive hybrid phenotypes (Rieseberg, et al. 1999), the effect of hybridization on gene expression is not well understood. Divergence in regulation of gene expression is thought to be the basis for much of phenotypic evolution (Rest, et al. 2013; Hodgins-Davis, et al. 2015). In spite of this, we lack an understanding of how gene expression contributes to novel phenotypes and adaptation in old, stabilized hybrid taxa, in particular for homoploid hybrid species, with no increase in ploidy level.

Gene expression evolves in a stabilizing manner where regulatory elements accumulate compensatory mutations that keep gene expression at an optimum level for physiological functions (Gilad, et al. 2006; Coolon, et al. 2014; Hodgins-Davis, et al. 2015). The compensatory evolution means that as hybridization breaks up co-inheritance of regulatory elements, hybrids may experience novel combinations (Landry, et al. 2005; Renaut, et al. 2009), potentially leading to transgressive expression patterns, that transcends the range of parental expression profiles. Indeed, transgressive gene expression has been reported in F1

hybrids in several taxa, and the underlying regulatory divergence for transgressive expression

has been associated with divergence in *cis*- and *trans*-regulatory elements in the parental

species (Ranz, et al. 2004; Haerty and Singh 2006; Rockman and Kruglyak 2006; Malone and

Michalak 2008; Renaut, et al. 2009). In post-F1 generation hybrids, transcriptomes show a

higher level of transgressive expression due to uncoupling of regulatory elements in the

process of recombination. In the lake white fish *Coregonus clupeaformis*, the F2 hybrids

showed a higher level of nonadditive inheritance in gene expression compared to F1 hybrids

(Renaut, et al. 2009). An additional line of evidence of a non-linear relationship between

genetic composition and gene expression comes from recent studies on allopolyploid plant

taxa. Both preferential expression of one of the parental genomes (Edger, et al. 2017), and

tissue dependent expression of this dominance (Kryvokhyzha, et al. 2019) have been

documented in allopolyploids. Jointly, the transgressive expression in experimental hybrids

and non-linear relationship between genomic composition and gene expression similarity,

suggest that gene expression will not be proportional to the parental contributions to the

genome. Whether transgressive patterns of gene expression are found in old, stabilized

homoploid hybrid taxa is therefore important for understanding the potential of hybridization

to produce novel variation.

Gene expression studies in hybrids resulting from strongly divergent taxa has shown an

extensive level of transgressive expression arising from uncoupling of co-evolved *cis*- and

trans-regulatory elements in the hybrids (Haerty and Singh 2006; McManus, et al. 2010;

Coolon, et al. 2014). However, transgressive expression patterns in the hybrid of more

recently diverged species pairs can either arise due to polymorphic incompatible loci (Cutter

2012) or physiological response to hybrid dysfunction (Barreto, et al. 2015). In

interpopulation hybrids at F2 or later generation of copepod species, *Tigriopus californicus*,

gene expression showed transgressive inheritance only in about 1.2% of the transcriptome

and was linked to physiological response in hybrids (Barreto, et al. 2015). Additionally,

recent study on F1 hybrids between the pied and collared flycatchers with few fixed

differences and large number of shared polymorphisms showed a tissue-specific pattern of

mis-expression. Testis had the highest divergence between the parental species but the lowest

level of mis-expression in F1 hybrids (Mugal, et al. 2020). To which extent this pattern holds

up across species, and how sorting of segregating polymorphisms may result in novel patterns

of gene expression remains a challenge for researchers to address.

Here, we leverage a unique study system to address how gene expression evolves. We

investigate gene expression in the homoploid hybrid Italian sparrow (*Passer italiae*) (Fig. 1),

through combining data from the old, stabilized hybrid taxon, the parental species and

experimental F1-hybrids. The Italian sparrow has arisen from hybridization between the

house sparrow (*P. domesticus*) and Spanish sparrow (*P. hispaniolensis*) (Hermansen, et al.

2011; Hermansen, et al. 2014; Trier, et al. 2014; Elgvin, et al. 2017). The parent species

diverged approximately 0.85 million years ago (Ravinet, et al. 2018) and the Italian sparrow

originated within ca. 5800 years ago, most likely as the house sparrow expanded its range

into Europe and hybridized with the Spanish sparrow (Ravinet, et al. 2018). This system

provides a unique possibility to test to what extend an old, stabilized homoploid hybrid

species has diverged in gene expression from the parent species, and how gene expression

has evolved after hybridization. In this study, we compared the gene expression profiles of

the gonads of the Italian sparrow to gene expression of the parental species and lab bred

experimental F1 hybrids to address a) if Italian sparrows exhibit transgressive gene

expression compared to the parent species and b) whether inheritance mode and

misexpression in F1s predict which genes are transgressively expressed in Italian sparrows.

Results

Differential expression between the parental species

Gene expression divergence in gonads between the parental species was strongly asymmetric with testis showing a more conserved pattern of expression compared to ovary (Table 1). In testis, 135 genes (1.16%) and in ovary, 1382 genes (11.65%) were significantly differentially expressed. Of the differentially expressed genes, 60.7% of genes in testis and 69.3% of genes in ovary were upregulated in the Spanish sparrow. A significantly higher proportion of genes that were differentially expressed in testis were located on the Z chromosome (24 genes, hypergeometric test, P-value < 1.82-08). We found no evidence for functional enrichment among differentially expressed genes in parental species for neither testis nor ovary.

Differential expression in Italian sparrows compared to the parental species

In contrast to its genome composition, intermediate of the parent species, the Italian sparrow testis gene expression differed more in comparison to the parental species than the parent species differed to each other (Table 1). About one-third of the testis transcriptome was differentially expressed compared to both parental species, whereas the ovary expression was similar to that of the house sparrow (Table 1, Figs. 2A-3A). In testis, 3536 genes (30.45%) and 3581 genes (30.9%) were differentially expressed in comparison to house and Spanish sparrow, respectively. In ovary, only 22 genes (0.18%) differed from the house sparrow whereas 1508 genes (12.63%) were differentially expressed compared to the Spanish sparrow. Significant over-representation of Z-linked genes among the differentially expressed genes was detected in testis but only in the comparison to Spanish sparrow (196 of the 3581 genes, hypergeometric test, p value = 0.006). Most genes in both testis and ovary were up-regulated compared to house and down-regulated compared to Spanish sparrow (Chi-squared

test *p* value = 1.05e-06 in testis and 0.001 in ovary) (Figs. 2B-3B).

Genes that were differentially expressed in Italian sparrow testis were enriched for similar

biological processes in comparison to both parent species. Primarily, functions involving

protein synthesis and mitochondrial gene expression and function were overrepresented. In

the comparison to house sparrow, 27 significant GO categories were found (Table S4), with

top biological processes involved in viral transcription, SRP-dependent co-translational

protein targeting to membrane, translation, ribosome and mitochondrial respiratory chain

complex I assembly. In comparison to the Spanish sparrow, the 15 enriched GO terms

reflected similar biological processes, with ribosome as the top GO term (Table S5). In ovary,

6 GO terms were detected in the comparison to house sparrow with sodium ion

transmembrane transporter activity as the top term (Table S6) and 1 GO term, negative

regulation of protein phosphorylation detected in the comparison to Spanish sparrow (Table

S7).

Differential expression in F1 hybrids compared to the parental species

Both testis and ovary showed a more conserved pattern of expression between the F1 hybrid

and the house sparrow than between F1 hybrids and the Spanish sparrow (Table 1, Figs. 2A-

3A). The overall magnitude of differences to the parent species was much lower than for the

Italian sparrow. In testis, 106 genes (0.9%) and 263 genes (2.25%) were differentially

expressed compared to house and Spanish sparrows, respectively. In ovary, 30 genes (0.25%)

and 140 genes (1.15%) showed a significant difference in expression compared to the house

and Spanish sparrows, respectively. Similar to the Italian sparrow, the pattern of up- and

down-regulation was consistent in both testis and ovary in the comparison to each of the

parental species (Figs. 2B-3B). A larger number of genes were up-regulated in F1 hybrid in

the comparison to house sparrow and down-regulated in the comparison to Spanish sparrow

(Chi-squared test *P*-value = 1.96-05 in testis and 0.01 in ovary). As in the Italian sparrow,

there was an over-representation of Z-linked genes among those differentially expressed

between F1 hybrids and Spanish sparrows (31 of the 263 differentially expressed genes,

hypergeometric test, *P*-value = 2.42-06). The enriched functions were different to those

enriched for Italian sparrows. Testis genes differentially expressed in F1 showed functional

enrichment for primarily collagen catabolic process. In the comparison to house sparrow, 46

GO terms were significantly over-represented. Among these GO terms, 20 GO terms had an

Padj < 0.01, whereas there were few significant GO terms compared to Spanish sparrows

with top GO term common with the comparison to house sparrow (Tables S8 and S9). The

top 5 GO terms were enriched in proteinaceous extracellular matrix, extracellular matrix

structural constituent, extracellular vesicle, collagen trimer and collagen catabolic process. In

ovary, while there was no functionally significant term associated with differentially

expressed genes in F1 compared to the house sparrow, there were 28 significant GO terms in

the comparison to Spanish sparrow with the top category to be response to vitamin D (Table

S10).

Classification of inheritance patterns of gene expression

Gene expression was generally conserved in both Italian sparrow and F1 hybrid (Table 2),

but in the testis of Italian sparrow, 2611 genes (22.71%) showed a non-conserved pattern of

inheritance with transgressive expression and house-dominant being the two largest

categories (Fig. 4C). Transgressively expressed genes in Italian sparrow comprised 22% of

the testis transcriptome tested for inheritance. In contrast, in F1 testis only 0.37% of the genes

were transgressively expressed (Fig. 4D). The majority of the genes with non-conserved

pattern of inheritance in Italian sparrow ovary, over 96% of the genes, had a house-dominant

pattern of expression (Fig. 4E). In contrast to the high incidence of transgressive expression

observed in Italian sparrow testis, only 3 genes (0.028%) were transgressively expressed in

ovaries. Gene expression in F1 ovary was more conserved compared to that of the Italian

sparrow. Out of 376 genes with non-conserved pattern of inheritance, the two largest

categories of inheritance were additive and house-dominant inheritance pattern (Fig. 4F).

Overdominant genes in Italian sparrow testis were enriched for functional categories involved

in protein synthesis, mitochondrial protein complex and gene expression and binding of

sperm to zona pellucida (Table S11). We found a gene network with significant protein-

protein interaction (PPI Enrichment: 1.0E-16) among the overdominant genes in Italian

sparrow (Fig. 5, Fig. S4). A different set of genes were underdominant in Italian sparrow,

with functions including regulation of cellular component size and negative regulation of

neuron projection development. Overdominant genes in F1 testis were functionally enriched

for different GO-terms, including collagen catabolic process and proteinaceous extracellular

matrix (Table S12). The only significant GO term for underdominant genes in F1 testis was

voltage-gated potassium channel complex.

Comparison of gene expression divergence in respect to parent species between Italian

sparrows and F1

To evaluate how gene expression evolves over time asking whether mis-expression in early

generation hybrids predicts differential expression in Italian sparrows, we checked if the

same genes or gene functions that were differentially expressed in comparison to parental

species in F1 hybrids also were differentially expressed in the Italian sparrow (Figs. 2C-3C).

In testis, 1.28% of the genes that were differentially expressed in Italian sparrows compared

to house sparrow were also differentially expressed in experimental F1 hybrids, whereas in

comparison to Spanish sparrows it was 3.9%. In ovary, only 1 gene (2%) was differentially

expressed in F1 and Italian sparrow compared to house sparrow, whereas 5.9% were

differentially expressed in F1 hybrids and Italian sparrows in comparison to Spanish

sparrows. The majority of the commonly differentially expressed genes were differentially

expressed in the same direction (i.e., up- or down-regulated) in both testis and ovary, both in

F1 hybrids and Italian sparrows. Testis genes commonly differentially expressed in F1 hybrid

and Italian sparrow compared to house sparrow were enriched in extracellular vesicle, insulin

like growth factor binding, extracellular matrix structural constituent and extracellular matrix

component. In the comparison to Spanish sparrow testis, only one significant GO term,

cAMP binding was detected.

Mode of regulatory divergence and prediction of transgressive expression in Italian

sparrows

To address if the regulatory basis can predict transgressive gene expression in hybrid species,

we classified genes dependent on their regulatory basis (Table 3). We expect genes under *cis-*

trans compensatory regulation and trans-regulated genes to more easily produce transgressive

patterns of gene expression in later generation hybrids than *cis*-regulated genes, as

recombination may de-couple *cis*- and *trans*- or different trans-elements in the sorting of

parental variation in the hybrid lineage. Therefore, we checked the percentage of genes

transgressively expressed in the Italian sparrow depending on the mode of regulatory

divergence. We found that 16/130 genes (12%) of the *trans*-regulated testis genes were

transgressively expressed in Italian sparrow, and 28/159 genes (18%) of the genes under *cis-*

trans compensatory regulation were transgressively expressed in Italian sparrows. Finally, we

tested whether genes with a non-additive inheritance, that were transgressively expressed in

F1 hybrids, were more likely to be transgressively expressed in the Italian sparrow. We found

that 20 out of 45 genes that were transgressively expressed in the F1 hybrid (44.4%) also

were transgressively expressed in Italian sparrow testis.

Discussion

In this study, we aimed to document the evolution of gene expression in the gonads of a

homoploid hybrid species, the Italian sparrow. An extensive level of transgressive expression

was found in Italian sparrow testis, with an over-expression of genes involved in functions

related to mitochondrial respiratory chain and, cytosolic and mitochondrial ribosomal

proteins, as well as proteins involved in protein folding and the proteasome complex. In

contrast to testis, the gene expression pattern in the ovary was asymmetric compared to

parental species, with a house-dominant expression in the Italian sparrow. Experimental F1

hybrids showed a much more conserved pattern of gene expression in both testis and ovary.

In the following sections, we discuss the potential implications of our results and consider the

limitations of these data.

Extensive transgressive expression in Italian sparrow testis

About 22% of testis transcriptome in Italian sparrow showed transgressive expression pattern,

while this level was less than 1% in the F1 hybrid. Transgressive gene expression, despite the

fact that gene expression is conserved between parent species, is expected to occur in later

generation hybrids due to the uncoupling of co-evolved *cis*- and *trans*- regulatory elements

through recombination (Rieseberg, et al. 2003; Mallet 2007). There are, however, few studies

of gene expression in post F1-hybrids. In Lake whitefish, F2 hybrids showed higher

misexpression and percentage of genes with nonadditive inheritance compared to F1 hybrids

(Renaut, et al. 2009). Transgressive expression is commonly referred to "mis-expression" in

early generation hybrids while Italian sparrow is a successful hybrid taxon likely formed

within the last 5.800 years (Ravinet, et al. 2018). Here, we show that transgressive

expression is found not only as mis-expression in early generation hybrids but remains in

well-functioning homoploid hybrid taxa long after hybridization. Hence, evolution of the

regulation of gene expression following hybridization may produce novel expression

phenotypes from intermediate genomes. In hybrid lineages, transgressive segregation, which

leads to extreme phenotypes, has been considered as a major mechanism by which extreme or

novel adaptations observed in new hybrid ecotypes or species arise (Rieseberg, et al. 1999),

but sorting of regulatory elements have not formally been suggested to contribute to this. This

research suggests that break-up of co-evolved regulatory elements, resulting in transgressive

patterns of gene expression, partly can explain the generation of transgressive phenotypes in

hybrid taxa.

Transgressive expression arising in spite of low parental divergence

Hybrid incompatibilities arisen due to bringing together novel genetic combinations have

been mainly considered when the parental species are fixed for alleles potentially

contributing to reproductive isolation (Cutter 2012). In recently diverged species,

reproductive incompatibility may arise due to polymorphic loci within each lineage before

fixation for the incompatibility alleles (Cutter 2012). Therefore, if transgressive expression in

the Italian sparrow is driven by bringing together *cis*- and *trans*-regulatory variants not yet

co-adapted, this must either be by bringing together polymorphic loci in each of the parental

species or novel mutations that have occurred in Italian sparrow since isolation from the

parental species. Each of these hypotheses can be evaluated in future work by studying the F1

and F2 hybrids and the direct parents of the cross in order to find *cis* and *trans*-regulatory

variations. Interestingly, during ca. 5.800 years of evolution, Italian sparrows have achieved a

26 times stronger divergence in gene expression the parents have accumulated during 0.68

million years, in spite of them being genomically intermediate (Elgvin, et al. 2017)

Tissue- and parent specific patterns of differentiation in gene expression

We find that the degree of difference in gene expression is tissue- and parent specific. We

generally observed fewer differentially expressed genes relative to house sparrow than

relative to the Spanish sparrow, both for Italian sparrows and F1 hybrids, and especially for

ovary. This is likely partly explained by the fact that both Italian sparrows and F1 hybrids

have house sparrow mitochondria. Mito-nuclear interactions can strongly affect gene

expression, and even result in misexpression (Runemark, Eroukhmanoff, et al. 2018). An

additional explanation for a more conserved expression pattern in ovary could be a

contribution of the W chromosome. While we can assume that F1 hybrids examined in this

study inherit house sparrow W chromosomes, there is not yet an assembly of the W

chromosome of any of the three sparrow species. Investigating how the W chromosome

affects gene expression is an interesting future line of inquiry. Finally, we cannot exclude

that a technical artefact contributes to the house-dominant pattern of expression. More

specifically, only 3 Spanish sparrows were used in the analysis of ovary transcriptome, this

finding may be driven by large within-species variance in ovarian gene expression.

Enrichment of over-dominant genes in Italian sparrow for translation and

mitochondrial function

A large set of genes involved in protein synthesis were over-expressed in Italian sparrow

testis compared to both parents. These include a set of ribonucleoproteins present in both

cytosolic and mitochondrial ribosomes, eukaryote translation initiation factor such as *EIF1AY*

required for maximal rate of protein synthesis and SRP (signal recognition particle), a

ribonucleoprotein that recognizes and targets specific proteins to the endoplasmic reticulum

in eukaryotes. RNA binding proteins play an important role in post-translational regulation of

gene expression and spermatogenesis and appear to be key regulatory factors that ensure

male fertility (Paronetto and Sette 2010; Phillips, et al. 2019). Several genes coding for

subunits of T-complex protein Ring Complex (TRiC) (*CCT4, CCT5, CCT6A, TCP1*) were

also among the overdominant genes. The TRiC complex aids in the folding of about 10% of

the proteome. Subunits of TRiC complex has shown to be required for spermatogenesis

(Counts, et al. 2017) and have shown evidence for positive selection among the seminal fluid

genes between house and Spanish sparrows (Rowe, et al. 2020). Another category of gene

networks which were upregulated in Italian sparrow included genes encoding for components

of all mitochondrial respiratory chain enzymatic complexes. Additionally, genes encoding for

chaperone proteins involved in the assembly of these enzymatic complexes such as

TIMMDC1 were also over-expressed.

In Italian sparrow testis, mito-nuclear genes have been shown to be important in generating

reproductive isolation in the Italian sparrow (Trier, et al. 2014), and mito-nuclear sites have

been under selection for inheritance of the house sparrow allele in Italian sparrows

(Runemark, Trier, et al. 2018). Although the mitochondrion has its own genome, more than

98% of its proteins are encoded in the nuclear genes and its correct function requires a

coordinated expression of both nuclear and mitochondrial genomes (Ryan and Hoogenraad

2007). Therefore, the upregulation of genes involved in respiratory chains and protein

synthesis corroborate the evidence for a strong effect of hybridization on genes associated

with metabolism in other bird taxa (Wagner, et al. 2020). Overexpression of cytosolic

ribonucleoproteins and mitochondrial respiratory chain enzymes has also been observed in

later generation hybrids (F3+) in copepod species *Tigriopus californicus* and has been linked

to cellular responses to physiological dysfunction (Barreto, et al. 2015). Our study thus

contributes to the growing body of evidence of mito-nuclear interactions as important

reproductive barriers (Hill 2017), and provides a unique insight into altered patterns of gene

expression as a possible resolution to the conflict.

Conclusion

This study presents the first evidence for transgressive gene expression in a wild homoploid

hybrid species. The strongly divergent gene expression profile in Italian sparrow testis

highlights the potential of intermediate hybrid genomes to rapidly evolve through altered

regulation of gene expression. Comparing patterns of gene expression in wild homoploid

hybrid species to those seen in experimental F1 hybrids, we document rapid evolution of

gene expression, especially for networks of genes involved in mitochondrial respiratory chain

enzymes and protein synthesis. The strongly divergent testis transcriptome was unexpected

given the low parental divergence, highlighting that alleles contributing to reproductive

isolation may be polymorphic rather than fixed. In contrast, Italian sparrow ovary had a

transcriptome similar to that of the house sparrow. Future studies should therefore evaluate

the current results in more tissues. Using pedigrees of parents and hybrid individuals would

also improve the power to identify the regulatory underpinnings of gene expression. Finally,

examining the link between transgressive gene expression and transgressive hybrid

phenotypes is likely to significantly increase our understanding of how novel variation arise

from hybridization.

Materials and methods

Sampling, RNA extraction, library preparation, and sequencing

We compared the gene expression of Italian sparrows to its two parent species, house and

Spanish sparrow and their experimental F1-hybrids. As the species differ slightly in

phenology, birds were sampled during their peak reproductive periods at their respective

locations. Spanish sparrows and F1 hybrids were sampled near Olivenza, Spain (38º 40'

56"N, 7º 11' 17"W) in March 2016 and 2017, house sparrows in Oslo, Norway (59º 55' 2"N,

10º 46' 11"E) in May 2016 and Italian sparrows from Montanari in the Gargano peninsula

(41º 54' 36.8"N, 15º 51' 13.0"E) in May 2017 (Fig 1). Experimental F1 hybrids were bred in

outdoor cages using house sparrow females and Spanish sparrow males. All other birds were

trapped using mist nets outside or within cages. Birds were immediately euthanized by

cervical dislocation, and gonads (ovary and left testis) were harvested directly after death.

Sampling was performed between 10:00 and 16:00, with individuals from each of the species

being sampled over most of the day, in order to avoid biases in sampling time. The gonads

were stored in RNAlater, and immediately stored at -80 degrees, with extractions later

performed at the University of Oslo using the Qiagen miRNeasy micro kit. Samples were

prepped with TruSeq® Stranded mRNA library prep from Illumina on an automated Perkin

Elmer Sciclone NGSx liquid handler and sequenced in the Norwegian Sequencing Center in

Oslo, as paired-end (2x150 bp) on the Illumina Hiseq4000. The number of reads obtained per

each sample is reported in Table S1. As F1 hybrids were bred in an area where house and

Spanish sparrows are sympatric (Olivenza, Spain; see Fig 1) and correct identification of

females can be challenging, we performed a post-hoc genomic analysis based on

mitochondrial and genomic DNA to ensure that only individuals that clustered with the

expected species were included in the analyses (Supplementary materials and methods, Figs.

S1-S2).

Read mapping, quality check and counting

We mapped reads from samples of all four groups (testis sample size: house sparrow = 5,

Spanish sparrow = 5, F1 hybrid = 5, Italian sparrow = 5; ovary sample size: house sparrow =

5, Spanish sparrow = 3, F1 hybrid = 8 and Italian sparrow = 5) to the house sparrow reference

genome using STAR2 v2.7.2b (Dobin, et al. 2013) with default parameters and the house

sparrow gene annotation as reported in (Elgvin, et al. 2017). Over 90% of the reads mapped

successfully (Table S1). The house sparrow general feature format (GFF) file contained

14,734 annotated genes of which 92.52% were anchored to chromosomes and 7.48% were

located on unanchored scaffolds. Of the genes anchored to chromosomes, 12,595 genes were

annotated on autosomes and 598 genes were on Z chromosome (Table S2). To ensure

homogeneity in quality across our samples, we used RSeQC (Wang, et al. 2012) to measure

RNA integrity at the transcriptome level for each sample by calculating Transcript Integrity

Number (TIN). The median TIN scores across samples ranged between 73.7 and 85.9 (Table

S3). We counted the number of reads mapping to the house sparrow gene features using

HTSeq v0.9.1 (Anders, et al. 2015). Read counting was performed for all reads with

minimum quality score of 30 and it was configured to handle reverse-stranded sequencing

data, with the parameter controlling for overlapping gene features set to union. We created a

dendrogram to show sample relationships. The main split was clearly between tissue types,

i.e. testis and ovary (Figure S3).

Differential gene expression analysis

We analyzed the raw read counts obtained from htseq-count using the R package DESeq2

(Anders and Huber 2010; Love, et al. 2014). We first tested genes for differential expression

between house and Spanish samples. We then compared each of the parents to the

experimental F1 hybrid and the Italian sparrow. We pre-filtered the data to a minimum of

total read counts of 10 across at least half of the samples in each pairwise comparison. To

generate more accurate estimates of log2 fold change (LFC) for genes with low count number

or large dispersion, we shrank LFC estimates. We considered genes to be differentially

expressed if they showed False Discovery Rate (FDR) Padj < 0.05 and shrunken LFC > 0.32.

Classification of gene expression inheritance in experimental F1 hybrids and Italian

sparrows

The mode of inheritance for differentially expressed genes was determined following

(McManus, et al. 2010). We normalized gene expression using median of ratios method in

DESeq2 (Love, et al. 2014). Genes whose total expression in hybrids deviated significantly

more than 1.25-fold (LFC > 0.32) from that of either parent were considered to have non-

conserved inheritance. These genes were classified as having additive, dominant, under-

dominant, or over-dominant inheritance, based on the magnitude of the difference between

total expression in the Italian sparrow / experimental F1 hybrid and in each parental species.

Genes for which expression in the Italian sparrow / experimental F1 hybrid was less than for

house sparrows and greater than for Spanish sparrow (or vice versa) were classified as

additive; genes for which expression in Italian sparrow / experimental F1 hybrid was similar

to one of the parents were classified as dominant; and genes for which expression in Italian

sparrow / experimental F1 hybrid was either greater than or less than both parent species

were classified as over-dominant and under-dominant, respectively (Figs. 4A-4B).

Allele-specific expression and *cis*- and *trans*- regulatory divergence assignment

Conserved expression level between parents controlled by compensatory variation in parental

species is often associated with misregulation in their hybrids (Denver, et al. 2005; Landry, et

al. 2005; Goncalves, et al. 2012).-By using sites in experimental F_1 hybrids that contain both

parental alleles, we measured allele specific expression, and compared that to expression

levels in the parental species to infer whether the divergence is due to *cis*-, *trans*- or

compensatory evolution. A typical approach for the study of ASE is to select the sites for

which the parental species are fixed for two different alleles and study the ASE in

heterozygote hybrids (Wittkopp, et al. 2004). House and Spanish sparrows diverged only

about 0.85 MYA, with many polymorphisms still shared between the two species. We

therefore took an approach similar to the one used in other studies with species pairs

containing high number of shared polymorphisms and few fixed differences (Mugal, et al.

2020).

We performed the SNP calling and filtering as described in the section "Genetic

Characterization of Samples" (Supplementary Materials and Methods). We additionally

filtered the sites for DP of minimum 10, base quality of 20 and mapping quality of 60. We

used the R package MBASED (Mayba, et al. 2014) in the unphased mode to identify genes

with allele-specific expression. We filtered the MBASED results for the FDR < 0.05 and

major allele frequency of minimum 0.7. To balance the power, we chose 5 individuals for

each house, Spanish, F1 and Italian sparrow in testis and 3 individuals of each group for

ovary. To identify allele-specific expression at the population level, we required genes to

show significant allele-specific expression in at least two individuals. Following (McGirr and

Martin 2019), we divided the regulatory categories as follow: Cis-regulatory divergence was

attributed to genes that are differentially expressed between parents and show allele-specific

expression in hybrids. Trans-regulatory divergence was assigned to genes that are

differentially expressed between parents and do not show allele-specific expression in

hybrids. Putative compensatory regulatory variation was determined if a gene 1) did not show

differential expression between parents, 2) showed significant allele-specific expression in

hybrids, and 3) did not show allele-specific expression in parents.

Functional Annotation and Enrichment Analysis

Gene ontology (GO) analysis was done for differentially expressed genes in each pairwise

comparison. GO functional annotations and gene descriptions were obtained for protein

sequences from the entire house sparrow protein set as described in (Rowe, et al. 2020) using

PANNZER (Koskinen, et al. 2015). Functional enrichment of GO terms present in the set of

differentially expressed genes relative to the background consisting of all genes expressed

and tested for differential expression was performed using the clusterProfiler (Yu, et al. 2012)

and significant enrichment was determined at FDR < 0.05. Interactions among proteins with

significant GO terms were predicted using STRING-v11 (Szklarczyk, et al. 2019) using

"Experiment", "Databases" and "Co-expression" as interaction sources. We set the minimum

required interaction score to the highest confidence (0.9) and used only the query proteins to

build the interaction network after removing the disconnected nodes in the network. The

obtained network was then exported to Cytoscape (Shannon, et al. 2003) and major

representations of biological processes were detected using ClueGO (Bindea, et al. 2009)

with Padj< 0.01.

Animal Ethics

In Spain, all trapping and sampling of birds was conducted in accordance with Spanish

Animal Protection Regulation RD53/2013, and all methods were approved by the

Institutional Commission of Bioethics at the University of Extremadura (CBUE 49/2011).

Birds were sampled in Oslo with permission from Miljø-direktoratet (2016/2225). In Italy, all

birds were sampled with permission from ISPRA (protocol number 12404) and with

permission (no. 305) from regione Puglia. All samples taken from Italy to Oslo in accordance

with the Nagoya protocols.

Scripts for the analyses

All scripts written for the analyses presented in this manuscript are deposited under

github.com/Homap/Expression_sparrow

Author contributions

A.R. and H.P.Y. conceived of the study. A.M. raised the hybrids and M. Ro., M. Ra., A.R.,

F.E, C.O. and G-P. S. performed field work and sampling, C.O. performed the molecular lab

work, and H.P.Y. designed and performed the bioinformatical analysis based on discussions

with A.R and M. Ra. H.P.Y. drafted the manuscript, with revisions based on the input from

A.R., M.Ra and M.Ro.

Acknowledgements

We thank Angelica Cuevas for assistance with sampling. The computations were enabled by

resources in project SNIC2020-6-222 provided by the Swedish National Infrastructure for

Computing (SNIC) at UPPMAX, partially funded by the Swedish Research Council through

grant agreement no. 2018-05973. This work was financed by a Norwegian Research council

grant to G-P. S. and A. R. and a Wenner-Gren and a Swedish Research Council grant to A.R.

Figure legends

Fig. 1. Species distribution and study design. (A) Top: Illustrations of male plumage patterns in house, Italian, and Spanish sparrows modified from (Svensson, et al. 1999). Bottom: A distribution map of house, Italian, and Spanish sparrows throughout Europe and northern Africa (Summers-Smith). This picture is modified from (Elgvin, et al. 2017). (B) Pedigree of the F1 hybrids with house sparrow mother and Spanish sparrow fathers.

Fig. 2. Gene expression in Italian sparrow and experimental F1 hybrid in comparison with parental species for testis. A) Log2 fold change (LFC) as a function of mean of normalized counts. B) Proportion of up- and down-regulated genes. Dark color: up-regulated, light color: down-regulated. C) Venn diagram depicting the intersection between differentially expressed genes between the experimental F1 hybrid and Italian sparrow with each of the parental species. F1h: experimental F1 hybrid – house sparrow; Ih: Italian sparrow – house sparrow; F1S: experimental F1 hybrid – Spanish sparrow; IS: Italian sparrow – Spanish sparrow. Purple: F1 hybrids, Orange: Italian sparrows.

Fig. 3. Gene expression in Italian sparrow and experimental F1 hybrids in comparison with parental species for ovary. A) Log2 fold change (LFC) as a function of mean of normalized counts. B) Proportion of up- and down-regulated genes. Dark color: up-regulated, light color: down-regulated. C) Venn diagram depicting the intersection between differentially expressed genes between the experimental F1 hybrid and Italian sparrow with each of the parental species. F1h: experimental F1 hybrid – house sparrow; Ih: Italian sparrow – house sparrow; F1S: experimental F1 hybrid – Spanish sparrow; IS: Italian sparrow – Spanish sparrow. Purple: F1 hybrids Orange: Italian sparrows.

Fig. 4. Inheritance pattern of gene expression in testis and ovary in Italian sparrows and

experimental F1 hybrids. A and B: Schematic figures representing the classification of

inheritance patterns. C and D: Scatter plots showing shrunken log2 fold change (LFC) in

testis between the Italian sparrow and experimental F1 hybrid with the house sparrow on the

x-axis and with Spanish sparrow on the *y*-axis, respectively. E and F: Scatter plots show

shrunken log2 fold change (LFC) in ovary between the Italian sparrow and experimental F1

hybrid with the house sparrow on the *x*-axis and with Spanish sparrow on the *y*-axis,

respectively. Grey points in each graph depicts the total number of genes studied for gene

expression with those colored representing the ones significantly different from parental

species to be divided into each of the inheritance categories (Conserved: grey, additive: pink,

house dominant: blue, Spain dominant: red, transgressive (over-dominant and under-

dominant: yellow). Grey dotted lines indicate the fold-change threshold of 1.25 used for

classification of inheritance pattern. Regression line represents the major variation in log2

fold change. F1h: experimental F1 hybrid – house sparrow; Ih: Italian sparrow – house

sparrow; F1S: experimental F1 hybrid – Spanish sparrow; IS: Italian sparrow – Spanish

sparrow.

Fig. 5. Gene ontology network represented in overdominant genes in testis of Italian sparrow.

Different colors depicts different significant biological processes. Circle size represents

adjusted *P*-value for each node with all Padj < 0.01.

578 **Table 1.** Number of differentially expressed genes and log2 fold change (LFC) with Padj<

579 0.05 and LFC* > 0.32.

	Testis			Ovary		
Comparison	Significant	LFC > 0	LFC < 0	Significant	LFC > 0	LFC < 0
Spain - House	135	82	53	1382	958	424
F$_1$ - House	106	77	29	30	16	14
F$_1$ - Spain	263	125	138	140	39	101
Italy - House	3536	1962	1574	22	13	9
Italy - Spain	3581	1779	1802	1508	394	1114

580 * LFC: Log Fold Change

581 **Table 2.** Number and percentage of genes in each inheritance category.

	Testis		Ovary	
Inheritance	F$_1$	Italy	F$_1$	Italy
Conserved	11515 (99.14%)	8883 (77.30%)	11600 (96.9%)	10889 (92.26%)
Additive	6 (0.05%)	2 (0.02%)	272 (2.27%)	19 (0.16%)
house dominant	45 (0.39%)	54 (0.47%)	94 (0.78%)	895 (7.58%)
Spain dominant	4 (0.03%)	25 (0.22%)	6 (0.05%)	7 (0.06%)
Underdominant	12 (0.1%)	1205 (10.47%)	2 (0.17%)	3 (0.02%)
Overdominant	33 (0.27%)	1325 (11.53%)	2 (0.17%)	1 (0.008%)

588   **Table 3.** Counts of genes with ASE and *cis-* , *trans-* and CE[*]-evolution

Comparison	Testis				Ovary			
	ASE	cis	trans	CE	ASE	cis	trans	CE
House	261	-	-	-	72	-	-	-
Spain	456	-	-	-	118	-	-	-
F1	164	5	130	159	54	9	1373	45

589   [*]Compensatory evolution

References

Abbott R, Albach D, Ansell S, Arntzen JW, Baird SJ, Bierne N, Boughman J, Brelsford A,
Buerkle CA, Buggs R, et al. 2013. Hybridization and speciation. J Evol Biol 26:229-246.
Anders S, Huber W. 2010. Differential expression analysis for sequence count data. Genome
Biology 11.
Anders S, Pyl PT, Huber W. 2015. HTSeq-a Python framework to work with high-throughput
sequencing data. Bioinformatics 31:166-169.
Barreto FS, Pereira RJ, Burton RS. 2015. Hybrid dysfunction and physiological
compensation in gene expression. Molecular Biology and Evolution 32:613-622.
Bindea G, Mlecnik B, Hackl H, Charoentong P, Tosolini M, Kirilovsky A, Fridman WH,
Pages F, Trajanoski Z, Galon J. 2009. ClueGO: a Cytoscape plug-in to decipher functionally
grouped gene ontology and pathway annotation networks. Bioinformatics 25:1091-1093.
Coolon JD, McManus CJ, Stevenson KR, Graveley BR, Wittkopp PJ. 2014. Tempo and
mode of regulatory evolution in Drosophila. Genome Res 24:797-808.
Counts JT, Hester TM, Rouhana L. 2017. Genetic expansion of chaperonin-containing TCP-1
(CCT/TRiC) complex subunits yields testis-specific isoforms required for spermatogenesis in
planarian flatworms. Mol Reprod Dev 84:1271-1284.
Cutter AD. 2012. The polymorphic prelude to Bateson-Dobzhansky-Muller incompatibilities.
Trends in Ecology & Evolution 27:209-218.
Denver DR, Morris K, Streelman JT, Kim SK, Lynch M, Thomas WK. 2005. The
transcriptional consequences of mutation and natural selection in Caenorhabditis elegans. Nat
Genet 37:544-548.
Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29:15-21.
Dobzhansky T. 1936. Studies on Hybrid Sterility. II. Localization of Sterility Factors in
Drosophila Pseudoobscura Hybrids. Genetics 21:113-135.
Edger PP, Smith R, McKain MR, Cooley AM, Vallejo-Marin M, Yuan Y, Bewick AJ, Ji L,
Platts AE, Bowman MJ, et al. 2017. Subgenome Dominance in an Interspecific Hybrid,
Synthetic Allopolyploid, and a 140-Year-Old Naturally Established Neo-Allopolyploid
Monkeyflower. Plant Cell 29:2150-2167.
Elgvin TO, Trier CN, Torresen OK, Hagen IJ, Lien S, Nederbragt AJ, Ravinet M, Jensen H,
Saetre GP. 2017. The genomic mosaicism of hybrid speciation. Sci Adv 3:e1602996.
Gilad Y, Oshlack A, Rifkin SA. 2006. Natural selection on gene expression. Trends in
Genetics 22:456-461.
Goncalves A, Leigh-Brown S, Thybert D, Stefflova K, Turro E, Flicek P, Brazma A, Odom
DT, Marioni JC. 2012. Extensive compensatory cis-trans regulation in the evolution of mouse
gene expression. Genome Res 22:2376-2384.
Haerty W, Singh RS. 2006. Gene regulation divergence is a major contributor to the
evolution of Dobzhansky-Muller incompatibilities between species of Drosophila. Molecular
Biology and Evolution 23:1707-1714.
Hermansen JS, Haas F, Trier CN, Bailey RI, Nederbragt AJ, Marzal A, Saetre GP. 2014.
Hybrid speciation through sorting of parental incompatibilities in Italian sparrows. Mol Ecol
23:5831-5842.
Hermansen JS, Saether SA, Elgvin TO, Borge T, Hjelle E, Saetre GP. 2011. Hybrid
speciation in sparrows I: phenotypic intermediacy, genetic admixture and barriers to gene
flow. Molecular Ecology 20:3812-3822.
Hill G. 2017. The mitonuclear compatibility species concept. The auk 134:393-409.
Hodgins-Davis A, Rice DP, Townsend JP. 2015. Gene Expression Evolves under a House-of-
Cards Model of Stabilizing Selection. Molecular Biology and Evolution 32:2130-2140.

Koskinen P, Toronen P, Nokso-Koivisto J, Holm L. 2015. PANNZER: high-throughput
functional annotation of uncharacterized proteins in an error-prone environment.
Bioinformatics 31:1544-1552.
Kryvokhyzha D, Salcedo A, Eriksson MC, Duan T, Tawari N, Chen J, Guerrina M, Kreiner
JM, Kent TV, Lagercrantz U, et al. 2019. Parental legacy, demography, and admixture
influenced the evolution of the two subgenomes of the tetraploid Capsella bursa-pastoris
(Brassicaceae). Plos Genetics 15:e1007949.
Landry CR, Wittkopp PJ, Taubes CH, Ranz JM, Clark AG, Hartl DL. 2005. Compensatory
cis-trans evolution and the dysregulation of gene expression in interspecific hybrids of
Drosophila. Genetics 171:1813-1822.
Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biology 15.
Mallet J. 2007. Hybrid speciation. Nature 446:279-283.
Mallet J. 2005. Hybridization as an invasion of the genome. Trends in Ecology & Evolution
20:229-237.
Malone JH, Michalak P. 2008. Gene expression analysis of the ovary of hybrid females of
Xenopus laevis and X. muelleri. Bmc Evolutionary Biology 8:82.
Marques DA, Meier JI, Seehausen O. 2019. A Combinatorial View on Speciation and
Adaptive Radiation. Trends in Ecology & Evolution 34:531-544.
Mayba O, Gilbert HN, Liu J, Haverty PM, Jhunjhunwala S, Jiang Z, Watanabe C, Zhang Z.
2014. MBASED: allele-specific expression detection in cancer tissues and cell lines. Genome
Biol 15:405.
McGirr JA, Martin CH. 2019. Hybrid gene misregulation in multiple developing tissues
within a recent adaptive radiation of Cyprinodon pupfishes. Plos One 14:e0218899.
McManus CJ, Coolon JD, Duff MO, Eipper-Mains J, Graveley BR, Wittkopp PJ. 2010.
Regulatory divergence in Drosophila revealed by mRNA-seq. Genome Res 20:816-825.
Mugal CF, Wang M, Backstrom N, Wheatcroft D, Alund M, Semon M, McFarlane SE,
Dutoit L, Qvarnstrom A, Ellegren H. 2020. Tissue-specific patterns of regulatory changes
underlying gene expression differences among Ficedula flycatchers and their naturally
occurring F1 hybrids. Genome Res 30:1727-1739.
Muller HJ. 1942. Isolating mechanisms, evolution, and temperature. Biol Symp. 6:71-125.
Paronetto MP, Sette C. 2010. Role of RNA-binding proteins in mammalian spermatogenesis.
Int J Androl 33:2-12.
Phillips BT, Williams JG, Atchley DT, Xu X, Li JL, Adams AL, Johnson KL, Hall TMT.
2019. Mass spectrometric identification of candidate RNA-binding proteins associated with
Transition Nuclear Protein mRNA in the mouse testis. Sci Rep 9:13618.
Ranz JM, Namgyal K, Gibson G, Hartl DL. 2004. Anomalies in the expression profile of
interspecific hybrids of Drosophila melanogaster and Drosophila simulans. Genome Res
14:373-379.
Ravinet M, Elgvin TO, Trier C, Aliabadian M, Gavrilov A, Saetre GP. 2018. Signatures of
human-commensalism in the house sparrow genome. Proc Biol Sci 285.
Renaut S, Nolte AW, Bernatchez L. 2009. Gene expression divergence and hybrid
misexpression between lake whitefish species pairs (Coregonus spp. Salmonidae). Molecular
Biology and Evolution 26:925-936.
Rest JS, Morales CM, Waldron JB, Opulente DA, Fisher J, Moon S, Bullaughey K, Carey
LB, Dedousis D. 2013. Nonlinear fitness consequences of variation in expression level of a
eukaryotic gene. Molecular Biology and Evolution 30:448-456.
Rieseberg LH, Archer MA, Wayne RK. 1999. Transgressive segregation, adaptation and
speciation. Heredity (Edinb) 83 (Pt 4):363-372.

Rieseberg LH, Raymond O, Rosenthal DM, Lai Z, Livingstone K, Nakazato T, Durphy JL,
Schwarzbach AE, Donovan LA, Lexer C. 2003. Major ecological transitions in wild
sunflowers facilitated by hybridization. Science 301:1211-1216.
Rockman MV, Kruglyak L. 2006. Genetics of global gene expression. Nature Reviews
Genetics 7:862-872.
Rowe M, Whittington E, Borziak K, Ravinet M, Eroukhmanoff F, Saetre GP, Dorus S. 2020.
Molecular Diversification of the Seminal Fluid Proteome in a Recently Diverged Passerine
Species Pair. Molecular Biology and Evolution 37:488-506.
Runemark A, Eroukhmanoff F, Nava-Bolanos A, Hermansen JS, Meier JI. 2018.
Hybridization, sex-specific genomic architecture and local adaptation. Philos Trans R Soc
Lond B Biol Sci 373.
Runemark A, Trier CN, Eroukhmanoff F, Hermansen JS, Matschiner M, Ravinet M, Elgvin
TO, Saetre GP. 2018. Variation and constraints in hybrid genome formation. Nat Ecol Evol
2:549-556.
Runemark A, Vallejo-Marin M, Meier JI. 2019. Eukaryote hybrid genomes. Plos Genetics
15:e1008404.
Ryan MT, Hoogenraad NJ. 2007. Mitochondrial-nuclear communications. Annu Rev
Biochem 76:701-722.
Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B,
Ideker T. 2003. Cytoscape: a software environment for integrated models of biomolecular
interaction networks. Genome Res 13:2498-2504.
Summers-Smith JD. The Sparrows: A Study of the Genus Passer: T & AD Poyser.
Svensson L, Grant PJ, Mullarney K, Zetterstroem D. 1999. Gyldendals store fugleguide:
Europas og middelhavsområdets fugler i felt: Gyldendal.
Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, Simonovic M,
Doncheva NT, Morris JH, Bork P, et al. 2019. STRING v11: protein-protein association
networks with increased coverage, supporting functional discovery in genome-wide
experimental datasets. Nucleic Acids Res 47:D607-D613.
Taylor SA, Larson EL. 2019. Insights from genomes into the evolutionary importance and
prevalence of hybridization in nature. Nat Ecol Evol 3:170-177.
Trier CN, Hermansen JS, Saetre GP, Bailey RI. 2014. Evidence for mito-nuclear and sex-
linked reproductive barriers between the hybrid Italian sparrow and its parent species. Plos
Genetics 10:e1004075.
Wagner DN, Curry RL, Chen N, Lovette IJ, Taylor SA. 2020. Genomic regions underlying
metabolic and neuronal signaling pathways are temporally consistent in a moving avian
hybrid zone. Evolution 74:1498-1513.
Wang L, Wang S, Li W. 2012. RSeQC: quality control of RNA-seq experiments.
Bioinformatics 28:2184-2185.
Wittkopp PJ, Haerum BK, Clark AG. 2004. Evolutionary changes in cis and trans gene
regulation. Nature 430:85-88.
Yu G, Wang LG, Han Y, He QY. 2012. clusterProfiler: an R package for comparing
biological themes among gene clusters. OMICS 16:284-287.
748