Extensive transgressive gene expression in testis but not ovary in the homoploid hybrid

Italian sparrow

Homa Papoli Yazdi[1*], Mark Ravinet[2], Melissah Rowe[3], Glenn-Peter Sætre[4], Caroline Øien

Guldvog[4], Fabrice Eroukhmanoff[4], Alfonso Marzal[5], Sergio Magallanes[5,6] & Anna

Runemark[1]

[1] Department of Biology, Lund University, SE-223 62 Lund, Sweden

[2] School of Life Sciences, University of Nottingham, Nottingham, UK

[3] Department of Animal Ecology, Netherlands Institute of Ecology (NIOO-KNAW), 6700

AB, Wageningen, The Netherlands

[4] Department of Biosciences, Centre for Ecological and Evolutionary Synthesis, University of

Oslo, PO Box 1066, N-0316, Oslo, Norway

[5] Department of Anatomy, Cellular Biology and Zoology, University of Extremadura, Avda.

de Elvas s/n, Badajoz ES-06006, Spain

[6] Department of Wetland Ecology, Doñana Biological Station (EBD-CSIC), Avda. Américo

Vespucio, 41092, Seville, Spain.

[*]Corresponding author: E-mail: Homa.Papoli_Yazdi@biol.lu.se

Abstract

Hybridization can result in novel allelic combinations which can impact the hybrid phenotype

through changes in gene expression. While mis-expression in F1 hybrids is well documented,

how gene expression evolves in stabilized hybrid taxa remains an open question. As gene

expression evolves in a stabilizing manner, break-up of co-evolved *cis-* and *trans-*regulatory

elements could lead to transgressive patterns of gene expression in hybrids. Here, we address

to what extent gonad gene expression has evolved in an established and stable homoploid

hybrid, the Italian sparrow (*Passer italiae*). Through comparison of gene expression in

gonads from individuals of the two parental species (i.e., house and Spanish sparrow) to that

of Italian sparrows, we find evidence for strongly transgressive expression in male Italian

sparrows - 22% of testis genes exhibit expression patterns outside the range of both parent

species. In contrast, Italian sparrow ovary expression was similar to that of one of the parent

species, the house sparrow (*Passer domesticus*). Moreover, the Italian sparrow testis

transcriptome is 26 times as diverged from those of the parent species as the parental

transcriptomes are from each other, despite being genetically intermediate. This highlights

the potential for regulation of gene expression to produce novel variation following

hybridization. Genes involved in mitochondrial respiratory chain complexes and protein

synthesis are enriched in the subset that is over-dominantly expressed in Italian sparrow

testis, suggesting that selection on key functions have molded the hybrid Italian sparrow

transcriptome.

Keywords: Gene expression, hybridization, mito-nuclear genes, testis and transgressive

expression

Introduction

Hybridization between recently diverged species is an important evolutionary phenomenon resulting in novel gene combinations that become available to selection (Abbott et al., 2013; Mallet, 2005; Runemark, Vallejo-Marin, & Meier, 2019; Taylor & Larson, 2019). While bringing together diverged parental alleles in hybrids can lead to reduced hybrid fitness (Dobzhansky, 1936; Muller, 1942), it is now clear that hybridization is frequent and contributes to novel adaptive phenotypes (Marques, Meier, & Seehausen, 2019; Runemark et al., 2019; Taylor & Larson, 2019). Despite the growing body of evidence documenting admixed genomes, it is not clear how such intermediate genomes give rise to transgressive hybrid phenotypes (Rieseberg, Archer, & Wayne, 1999). Gene expression is the key intermediate step between genotype and phenotype, and divergence in regulation of gene expression is thought to play an important role in phenotypic evolution (Hodgins-Davis, Rice, & Townsend, 2015; Rest et al., 2013). Despite this, we lack an understanding of how gene expression evolves and contributes to novel phenotypes in hybrid taxa.

Gene expression evolves in a stabilizing manner, where regulatory elements accumulate mutations that keep gene expression at an optimum level for physiological functions (Coolon, McManus, Stevenson, Graveley, & Wittkopp, 2014; Gilad, Oshlack, & Rifkin, 2006; Hodgins-Davis et al., 2015). Specifically, changes in distal, *trans*-regulatory elements are compensated by changes in proximate, *cis*-regulatory elements (Landry et al., 2005). Such compensatory evolution means that as hybridization breaks up co-inheritance of regulatory elements, hybrids may experience novel combinations (Landry et al., 2005; Renaut, Nolte, & Bernatchez, 2009). The novel combinations of regulatory elements arising during hybridization has the potential to lead to transgressive gene expression, that transcends the range of parental expression profiles.

Hybrids resulting from strongly divergent taxa may exhibit extensive levels of transgressive

expression arising from uncoupling of co-evolved *cis*- and *trans*-regulatory elements in F1

hybrids (Coolon et al., 2014; Haerty & Singh, 2006; McManus et al., 2010). In post-F1

generation hybrids, transcriptomes show a higher level of transgressive expression due to

uncoupling of regulatory elements in the process of recombination. In the lake white fish

Coregonus clupeaformis for example, F2 hybrids showed a higher level of nonadditive

inheritance in gene expression compared to F1 hybrids (Renaut et al., 2009). An additional

line of evidence of a non-linear relationship between genetic composition and gene

expression comes from recent studies on allopolyploid plant taxa, hybrid taxa with doubled

chromosome number. Both preferential expression of one of the parental genomes (Edger et

al., 2017), and tissue dependent parental dominance in expression pattern (Kryvokhyzha et

al., 2019) have been documented in allopolyploids. Jointly, the transgressive expression in

experimental hybrids and non-linear relationship between genomic composition and gene

expression similarity, suggest that gene expression will not be proportional to the parental

contributions to the genome. Whether transgressive patterns of gene expression are found in

homoploid hybrid taxa, hybrids without increase in ploidy, is unknown.

Here, we leverage a unique study system to investigate the nature of gene expression in a

stabilized hybrid taxon, the homoploid hybrid Italian sparrow (*Passer italiae*), where

thousands of generations of selection have resulted in a stable genome composition. The

Italian sparrow arose from hybridization between the house sparrow (*Passer domesticus*) and

Spanish sparrow (*Passer hispaniolensis*) (Fig. 1) (Elgvin et al., 2017; Hermansen et al., 2014;

Hermansen et al., 2011; Trier, Hermansen, Saetre, & Bailey, 2014). The parent species

diverged approximately 0.85 million years ago (Ravinet et al., 2018) and the Italian sparrow

originated within *ca*. 5800 years, most likely as the house sparrow expanded its range into

Europe and hybridized with the Spanish sparrow (Ravinet et al., 2018). This system provides

a unique possibility to test to what extent a homoploid hybrid species has diverged in gene

expression from the parent species after thousands of generations. In this study, we compared

the gene expression profiles of the gonads of wild individuals of Italian sparrow to gene

expression of the parental species to address if gene expression in Italian sparrows is

intermediate to that of its parent species. At the same time, we opportunistically compared

gene expression profiles to experimental F1 hybrid crosses (house x Spanish) bred in

captivity. Thereby, we test whether gene expression reflects the intermediate genome

composition of the hybrid Italian sparrow, or if it is transgressive for specific gene categories.

Materials and methods

Sampling, RNA extraction, library preparation, and sequencing

We compared the gene expression of wild Italian sparrows to wild individuals of its two

parent species, the house and Spanish sparrow. Although the species differ in their

distribution and phenology, all birds were sampled during the breeding season when birds

were actively breeding (Supplementary section 1) at their respective locations. Spanish

sparrows were sampled near Olivenza, Spain (38º 40' 56"N, 7º 11' 17"W) in March 2016,

house sparrows in Oslo, Norway (59º 55' 2"N, 10º 46' 11"E) in May 2016 and Italian

sparrows from Masseria Montanari in the Gargano peninsula (41º 54' 36.8"N, 15º 51' 13.0"E)

in May 2016 (Fig. 1). All birds were trapped using mist nets and placed in a soft cotton bird

bag until processing. All birds were euthanized by cervical dislocation, and gonads (ovary

and the left testis) were harvested immediately after death. Sampling was performed between

10:00 and 16:00, with individuals from each of the species being sampled over most of the

day, in order to avoid biases in sampling time. The gonads were stored in RNAlater, and

immediately stored at -80 degrees, with extractions later performed at the University of Oslo

using the Qiagen miRNeasy micro kit. Samples were prepped with TruSeq® Stranded

mRNA library prep from Illumina on an automated Perkin Elmer Sciclone NGSx liquid

handler and sequenced in the Norwegian Sequencing Center in Oslo, as paired-end (2x150

bp) on the Illumina Hiseq4000. The number of reads obtained per each sample is reported in

Table S1.

146
Read mapping, quality check and counting

We mapped reads from samples of all three groups (testis sample size: house sparrow = 5,

Spanish sparrow = 5, Italian sparrow = 5; ovary sample size: house sparrow = 5, Spanish

sparrow = 3 and Italian sparrow = 5) to the house sparrow reference genome using STAR2

v2.7.2b (Dobin et al., 2013) with default parameters and the house sparrow gene annotation

as reported in (Elgvin et al., 2017). Over 90% of the reads mapped successfully (Table S1).

The house sparrow general feature format (GFF) file contained 14,734 annotated genes of

which 92.52% were anchored to chromosomes and 7.48% were located on unanchored

scaffolds. Of the genes anchored to chromosomes, 12,595 genes were annotated on

autosomes and 598 genes were on Z chromosome (Table S2). To ensure homogeneity in

quality across our samples, we used RSeQC (Wang, Wang, & Li, 2012) to measure RNA

integrity at the transcriptome level for each sample by calculating Transcript Integrity

Number (TIN). The median TIN scores across samples ranged between 73.7 and 85.9 (Table

S3). We counted the number of reads mapping to the house sparrow gene features using

HTSeq v0.9.1 (Anders, Pyl, & Huber, 2015). Read counting was performed for all reads with

minimum quality score of 30 and configured to handle reverse-stranded sequencing data,

with the parameter controlling for overlapping gene features set to union. We created a

dendrogram to show sample relationships. The main split was clearly between tissue types,

i.e., testis and ovary (Fig. S1).

Differential gene expression analysis

We analyzed the raw read counts obtained from htseq-count using the R package DESeq2

(Anders & Huber, 2010; Love, Huber, & Anders, 2014). We first tested genes for differential

expression between house and Spanish samples. We then compared each of the parents to the

Italian sparrow. We pre-filtered data to a minimum of total read counts of 10 across at least

half of the samples in each pairwise comparison. To generate more accurate estimates of log2

fold change (LFC) for genes with low count number or large dispersion, we shrank LFC

estimates. We considered genes to be differentially expressed if they showed False Discovery

Rate (FDR) Padj < 0.05 and shrunken LFC > 0.32. All statistical analyses were completed

using R version 4.0.2 (R Core Team 2020).

Classification of gene expression inheritance

The mode of inheritance for differentially expressed genes was determined following

(McManus et al., 2010). We normalized gene expression using median of ratios method in

DESeq2 (Love et al., 2014). Genes whose total expression in Italian sparrow deviated

significantly more than 1.25-fold (LFC > 0.32) from that of either parent were considered to

have non-conserved inheritance. These genes were classified as having additive, dominant,

under-dominant, or over-dominant inheritance, based on the magnitude of the difference

between total expression in the Italian sparrow and in each parental species. Genes for which

expression in the Italian sparrow was less than for house sparrows and greater than for

Spanish sparrow (or vice versa) were classified as additive; genes for which expression in

Italian sparrow was similar to one of the parents were classified as dominant; and genes for

which expression in Italian sparrow was either greater than or less than both parent species

were classified as over-dominant and under-dominant (transgressive), respectively (Figs. 3A-

3B).

Functional Annotation and Enrichment Analysis

Gene ontology (GO) analysis was done for differentially expressed genes in each pairwise

comparison. GO functional annotations and gene descriptions were obtained for protein

sequences from the entire house sparrow protein set as described in (Rowe et al., 2020) using

PANNZER (Koskinen, Toronen, Nokso-Koivisto, & Holm, 2015). Functional enrichment of

GO terms present in the set of differentially expressed genes relative to the background

consisting of all genes expressed and tested for differential expression was performed using

the clusterProfiler (Yu, Wang, Han, & He, 2012) and significant enrichment was determined

at FDR < 0.05. Interactions among proteins with significant GO terms were predicted using

STRING-v11 (Szklarczyk et al., 2019) using "Experiment", "Databases" and "Co-

expression" as interaction sources. We set the minimum required interaction score to the

highest confidence (0.9) and used only the query proteins to build the interaction network

after removing the disconnected nodes in the network. The obtained network was then

exported to Cytoscape (Shannon et al., 2003) and major representations of biological

processes were detected using ClueGO (Bindea et al., 2009) with Padj< 0.01.

Analysis of experimental hybrids

Although our study was designed to allow the comparison of wild Italian sparrows to wild

individuals of its two parent species, the house and Spanish sparrow, we also capitalized on

the availability of captive-bred experimental F1 hybrids produced by crossing house sparrow

females and Spanish sparrow males (Supplementary section 2, Fig. S2). As experimental

hybrids were bred in an area where house and Spanish sparrows are sympatric for at least part

of the year (Olivenza, Spain; see Fig. 1) and correct identification of females can be

challenging, we performed a post-sampling genomic analysis of all samples (Supplementary

section 3). The mtDNA of experimental F1 hybrids were correctly grouped with house

sparrow (Fig. S3). However, the F1 hybrids, did not show the expected equal ancestry ratio

from each parental species and the PCA with the Z chromosome of F1 females showed

clustering with Z chromosome from house sparrow (Figs. S4-S5). We therefore acknowledge

that the comparison between these experimental hybrids and the other species sampled in our

study is suboptimal. Nonetheless, since the comparison of the stabilized homoploid hybrid

Italian sparrow and artificial early generation hybrids between house and Spanish sparrows

provides an interesting contrast, we decided to retain the experimental hybrid samples while

clarifying the limitations to how these findings can be interpreted. All gene expression

analyses were carried out in these experimental hybrids in same way as described for the

Italian sparrow samples.

Results

228 Results

229 Differential expression between the parental species

Gene expression divergence in gonads between the parental species was strongly asymmetric

with testis showing a more conserved pattern of expression compared to ovary (Table 1). In

the testis, 135 genes (1.16%) and in ovary, 1382 genes (11.65%) were significantly

differentially expressed. Of the differentially expressed genes, 60.7% of genes in testis and

69.3% of genes in ovary were upregulated in the Spanish sparrow. A significantly higher

proportion of genes that were differentially expressed in testis were located on the Z

chromosome (24 genes, hypergeometric test, P-value < 1.82-08). We found no evidence for

functional enrichment among differentially expressed genes in parental species for either

testis or ovary.

240 Differential expression in Italian sparrows compared to the parental species

In contrast to the intermediacy of the Italian sparrow genome, Italian sparrow testis gene

expression differed more in comparison to the parental species than the parent species

differed from each other (Table 1). Specifically, approximately one-third of the testis

transcriptome was differentially expressed in Italian sparrow compared to both parental

species, whereas ovary expression was similar to that of the house sparrow (Table 1, Fig. 2).

In the Italian sparrow testis, 3536 genes (30.45%) and 3581 genes (30.9%) were differentially

expressed in comparison to house and Spanish sparrow, respectively. For Italian sparrow

ovary, only 22 genes (0.18%) differed from the house sparrow whereas 1508 genes (12.63%)

were differentially expressed compared to the Spanish sparrow. Significant over-

representation of Z-linked genes among the differentially expressed genes was detected in

testis, but only in the comparison to Spanish sparrow (196 of the 3581 genes, hypergeometric

test, p value = 0.006). Most genes in both testis and ovary were up-regulated compared to

house and down-regulated compared to Spanish sparrow (Chi-squared test p value = 1.05e-06

in testis and 0.001 in ovary) (Fig. 2).

Genes that were differentially expressed in Italian sparrow testis were enriched for similar

biological processes in comparison to both parent species. Primarily, functions involving

protein synthesis and mitochondrial gene expression and function were overrepresented.

Comparisons between the Italian sparrow and the house sparrow yielded 27 significant GO

categories (Table S5), with top biological processes involved in viral transcription, SRP-

dependent co-translational protein targeting to membrane, translation, ribosome and

mitochondrial respiratory chain complex I assembly. In comparisons between the Italian

sparrow and the Spanish sparrow, the 15 enriched GO terms reflected similar biological

processes, with ribosome as the top GO term (Table S6). In ovary, 6 GO terms were detected

in the comparison to house sparrow with sodium ion transmembrane transporter activity as

the top term (Table S7) and 1 GO term, negative regulation of protein phosphorylation

detected in the comparison to Spanish sparrow (Table S8).

Classification of inheritance patterns of gene expression

Gene expression was generally conserved in Italian sparrow (Table 2), but in the testis of

Italian sparrow, 2611 genes (22.71%) showed a non-conserved pattern of inheritance with

transgressive expression and house-dominant being the two largest categories (Fig. 3C).

Transgressively expressed genes in Italian sparrow comprised 22% of the testis transcriptome

tested for inheritance. Over 96% of genes with non-conserved pattern of inheritance in Italian

sparrow ovary, had a house-dominant pattern of expression (Fig. 3D). In contrast to the high

incidence of transgressive expression observed in Italian sparrow testis, only 3 genes

(0.028%) were transgressively expressed in ovaries. Over-dominant genes in Italian sparrow

testis were enriched for functional categories involved in protein synthesis, mitochondrial

protein complex and gene expression and binding of sperm to zona pellucida (Table S10).

We found a gene network with significant protein-protein interaction (PPI Enrichment: 1.0E-

16) among the over-dominant genes in Italian sparrow (Fig. 4). A different set of genes was

under-dominant in Italian sparrow, with functions including regulation of cellular component

size and negative regulation of neuron projection development.

Patterns of expression in experimental hybrids

The overall magnitude of differences in gene expression in the experimental hybrid compared

to both parent species was much lower than for the Italian sparrow (Figure S6). In testis, 106

genes (0.9%) and 263 genes (2.25%) were differentially expressed compared to house and

Spanish sparrows, respectively. In ovary, 30 genes (0.25%) and 140 genes (1.15%) showed a

significant difference in expression compared to the house and Spanish sparrows,

respectively. Moreover, in the experimental hybrid testis only 0.37% of the genes were

transgressively expressed (Table S11). Gene expression in experimental hybrid ovary also

was more conserved in relation to the parent species compared to that of the Italian sparrow,

with the two largest categories of inheritance being additive and house-dominant (Table S11).

Additional results from the experimental hybrids are presented in the Supplementary section

4 and Tables S12-S15.

Discussion

This study presents, to the best of our knowledge, the first evidence for transgressive gene expression in a wild homoploid hybrid species. The strongly divergent gene expression profile in Italian sparrow testis highlights the potential of intermediate hybrid genomes to rapidly evolve through altered regulation of gene expression. About 22% of testis transcriptome in Italian sparrow showed transgressive expression pattern, while this level was less than 0.03% in the ovaries. Transgressive gene expression, despite conserved parental expression, could arise in later generation hybrids due to the uncoupling of co-evolved *cis*- and *trans*- regulatory elements through recombination (Landry et al., 2005; Signor & Nuzhdin, 2018; Takahasi, Matsuo, & Takano-Shimizu-Kouno, 2011). There are, however, few studies of gene expression in post F1-hybrids and transgressive expression is commonly referred to "mis-expression" in early generation hybrids (Signor & Nuzhdin, 2018). Here, we show that the homoploid hybrid Italian sparrow has achieved a 26 times stronger divergence in testis gene expression during ca. 5800 years of evolution following hybridization than the parents have accumulated during 0.68 million years (Ravinet et al., 2018). This might imply that sorting of parental regulatory elements following hybridization can produce novel expression phenotypes from intermediate genomes.

Hybrid incompatibilities leading to reproductive isolation due to novel, untested genetic combinations are most commonly thought to occur when parental species are fixed for different alleles (Cutter, 2012). However, transgressive expression patterns in the hybrids of more recently diverged species pairs can also arise either due to polymorphic incompatible loci (Cutter, 2012) or physiological response to hybrid dysfunction (Barreto, Pereira, & Burton, 2015). In interpopulation hybrids at F2 or later generation of copepod species (*Tigriopus californicus*), transgressive gene expression mainly reflected physiological

responses in hybrids (Barreto et al., 2015). Therefore, given the relatively low divergence

between house- and Spanish sparrow, transgressive expression in the Italian sparrow could

also be due to bringing together polymorphic *cis-* and *trans-*regulatory variants from the

parental species. Alternatively, transgressive gene expression may arise from lineage specific

mutations that have been favored by selection in Italian sparrow. Each of these hypotheses

can be evaluated in future work by studying the optimal F1 and F2 hybrids and their parental

individuals to identify *cis* and *trans-*regulatory variants.

The degree of difference in gene expression was parent- and tissue specific. We generally

observed fewer differentially expressed genes relative to house sparrow than relative to the

Spanish sparrow for the Italian sparrow, and this was especially true for ovarian tissue. While

smaller sample size in Spanish sparrow might have contributed to the house-dominant pattern

of expression, it is likely that our findings are at least partly explained by the fact that the

Italian sparrow has house sparrow mitochondria (Hermansen et al., 2014; Runemark,

Eroukhmanoff, Nava-Bolanos, Hermansen, & Meier, 2018; Trier et al., 2014). Mito-nuclear

interactions can strongly affect gene expression (Sanchez-Ramirez, Weiss, Thomas, & Cutter,

2021), and even result in mis-expression (Runemark, Eroukhmanoff, et al., 2018). Mito-

nuclear genes have a role in reproductive isolation in Italian sparrows and have been under

selection for inheritance of the house sparrow allele (Hermansen et al., 2014; Runemark,

Trier, et al., 2018; Trier et al., 2014). Interestingly, we found over-expression of genes

involved in functions related to mitochondrial respiratory chain, cytosolic and mitochondrial

ribosomal proteins in the hybrid Italian sparrow. Correct mitochondrial function requires a

coordinated expression of both nuclear and mitochondrial genomes (Ryan & Hoogenraad,

2007). Therefore, the upregulation of genes involved in respiratory chains and protein

synthesis corroborate the evidence for a strong effect of hybridization on genes associated

with metabolism in other bird taxa (Wagner, Curry, Chen, Lovette, & Taylor, 2020).

Overexpression of cytosolic ribonucleoproteins and mitochondrial respiratory chain enzymes

has also been observed in later generation hybrids (F3+) in copepod species and has been

linked to cellular responses to physiological dysfunction (Barreto et al., 2015). Our study thus

contributes to the growing body of evidence of mito-nuclear interactions as important

reproductive barriers (Hill, 2017), and provides a unique insight into altered patterns of gene

expression as a possible resolution to the conflict.

In addition to mito-nuclear genes, over-dominant genes in Italian sparrow contained

ribonucleoproteins present in both cytosolic and mitochondrial ribosomes. RNA binding

proteins play an important role in post-translational regulation of gene expression and

spermatogenesis and appear to be key regulatory factors that ensure male fertility (Paronetto

& Sette, 2010; Phillips et al., 2019). Several of the over-dominant genes also coded for

subunits of T-complex protein Ring Complex (TRiC) involved in folding of about 10% of the

proteome. Subunits of TRiC complex are required for spermatogenesis (Counts, Hester, &

Rouhana, 2017) and are under positive selection among the seminal fluid genes in passerine

species (Rowe et al., 2020).

Although we suggest caution in interpretation of our experimental F1 hybrid data due to the

unusual genomic composition of F1 hybrids, our work is in agreement with findings from F1s

in two other songbird species (Davidson & Balakrishnan, 2016; Mugal et al., 2020). F1

hybrids of flycatchers also showed a tissue specific pattern of expression differences, with a

low degree of transgression in genes expressed in testis (Mugal et al., 2020), similar to the

pattern in our experimental hybrids but contrasting to the strong transgression in the testis

transcriptome of the Italian sparrow. Future studies should therefore extend sampling to non-

reproductive tissues in order to assess whether transgressive expression is consistent across

tissues. While a standardized environment and time for sampling would be ideal, this is not

easily achieved in wild species that differ in geographic distributions, phenologies and

thermal comfort zones, as sampling at the same time could capture different phenological

stages. Future studies should sample these species over time to remedy this complication.

Homoploid hybrid species are expected to have higher level of genetic variance due to

homologous recombination leading to transgressive phenotypes (Mallet, 2007; Rieseberg et

al., 2003). The extensive level of transgressive expression in the Italian sparrow testis is

therefore in agreement with this expectation. Gene expression is the key intermediate step

between genotype and phenotype, therefore examining the link between transgressive gene

expression and transgressive hybrid phenotypes is a crucial next step which will significantly

increase our understanding of how novel variation arises from hybridization and the role of

gene expression divergence in the evolution of reproductive isolation from parental species.

Acknowledgements

We thank Angelica Cuevas, Matteo Caldarella, Severino Vitulano and Giuseppe Cardone for assistance with sampling, which was done in collaboration with the Puglia region, Centre for Naturalistic Studies ONLUS (CSN, Italy) and ISPRA (Italy). We are grateful to Richard Bailey and Daniel Hooper for their comments on earlier versions of this manuscript. The computations were enabled by resources in project SNIC2020-6-222 provided by the Swedish National Infrastructure for Computing (SNIC) at UPPMAX, partially funded by the Swedish Research Council through grant agreement no. 2018-05973. This work was financed by a Norwegian Research council grant to G-P. S. and A. R. and a Wenner-Gren and a Swedish Research Council grant to A.R.

References

Abbott, R., Albach, D., Ansell, S., Arntzen, J. W., Baird, S. J., Bierne, N., . . . Zinner, D. (2013). Hybridization and speciation. *J Evol Biol, 26*(2), 229-246. doi:10.1111/j.1420-9101.2012.02599.x

Anders, S., & Huber, W. (2010). Differential expression analysis for sequence count data. *Genome Biology, 11*(10). doi:ARTN R106

10.1186/gb-2010-11-10-r106

Anders, S., Pyl, P. T., & Huber, W. (2015). HTSeq-a Python framework to work with high-throughput sequencing data. *Bioinformatics, 31*(2), 166-169. doi:10.1093/bioinformatics/btu638

Barreto, F. S., Pereira, R. J., & Burton, R. S. (2015). Hybrid dysfunction and physiological compensation in gene expression. *Molecular Biology and Evolution, 32*(3), 613-622. doi:10.1093/molbev/msu321

Bindea, G., Mlecnik, B., Hackl, H., Charoentong, P., Tosolini, M., Kirilovsky, A., . . . Galon, J. (2009). ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics, 25*(8), 1091-1093. doi:10.1093/bioinformatics/btp101

Coolon, J. D., McManus, C. J., Stevenson, K. R., Graveley, B. R., & Wittkopp, P. J. (2014). Tempo and mode of regulatory evolution in Drosophila. *Genome Res, 24*(5), 797-808. doi:10.1101/gr.163014.113

Counts, J. T., Hester, T. M., & Rouhana, L. (2017). Genetic expansion of chaperonin-containing TCP-1 (CCT/TRiC) complex subunits yields testis-specific isoforms required for spermatogenesis in planarian flatworms. *Mol Reprod Dev, 84*(12), 1271-1284. doi:10.1002/mrd.22925

Cutter, A. D. (2012). The polymorphic prelude to Bateson-Dobzhansky-Muller incompatibilities. *Trends in Ecology & Evolution, 27*(4), 209-218. doi:10.1016/j.tree.2011.11.004

Davidson, J. H., & Balakrishnan, C. N. (2016). Gene Regulatory Evolution During Speciation in a Songbird. *G3 (Bethesda), 6*(5), 1357-1364. doi:10.1534/g3.116.027946

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., . . . Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics, 29*(1), 15-21. doi:10.1093/bioinformatics/bts635

Dobzhansky, T. (1936). Studies on Hybrid Sterility. II. Localization of Sterility Factors in Drosophila Pseudoobscura Hybrids. *Genetics, 21*(2), 113-135. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/17246786

Edger, P. P., Smith, R., McKain, M. R., Cooley, A. M., Vallejo-Marin, M., Yuan, Y., . . . Puzey, J. R. (2017). Subgenome Dominance in an Interspecific Hybrid, Synthetic Allopolyploid, and a 140-Year-Old Naturally Established Neo-Allopolyploid Monkeyflower. *Plant Cell, 29*(9), 2150-2167. doi:10.1105/tpc.17.00010

Elgvin, T. O., Trier, C. N., Torresen, O. K., Hagen, I. J., Lien, S., Nederbragt, A. J., . . . Saetre, G. P. (2017). The genomic mosaicism of hybrid speciation. *Sci Adv, 3*(6), e1602996. doi:10.1126/sciadv.1602996

Gilad, Y., Oshlack, A., & Rifkin, S. A. (2006). Natural selection on gene expression. *Trends in Genetics, 22*(8), 456-461. doi:10.1016/j.tig.2006.06.002

Haerty, W., & Singh, R. S. (2006). Gene regulation divergence is a major contributor to the evolution of Dobzhansky-Muller incompatibilities between species of Drosophila. *Molecular Biology and Evolution, 23*(9), 1707-1714. doi:10.1093/molbev/msl033

Hermansen, J. S., Haas, F., Trier, C. N., Bailey, R. I., Nederbragt, A. J., Marzal, A., & Saetre,
G. P. (2014). Hybrid speciation through sorting of parental incompatibilities in Italian
sparrows. *Mol Ecol, 23*(23), 5831-5842. doi:10.1111/mec.12910
Hermansen, J. S., Saether, S. A., Elgvin, T. O., Borge, T., Hjelle, E., & Saetre, G. P. (2011).
Hybrid speciation in sparrows I: phenotypic intermediacy, genetic admixture and
barriers to gene flow. *Molecular Ecology, 20*(18), 3812-3822. doi:10.1111/j.1365-
294X.2011.05183.x
Hill, G. (2017). The mitonuclear compatibility species concept. *The auk, 134*(2), 393-409.
doi:DOI:10.1642/AUK-16-201.1
Hodgins-Davis, A., Rice, D. P., & Townsend, J. P. (2015). Gene Expression Evolves under a
House-of-Cards Model of Stabilizing Selection. *Molecular Biology and Evolution,
32*(8), 2130-2140. doi:10.1093/molbev/msv094
Koskinen, P., Toronen, P., Nokso-Koivisto, J., & Holm, L. (2015). PANNZER: high-
throughput functional annotation of uncharacterized proteins in an error-prone
environment. *Bioinformatics, 31*(10), 1544-1552. doi:10.1093/bioinformatics/btu851
Kryvokhyzha, D., Salcedo, A., Eriksson, M. C., Duan, T., Tawari, N., Chen, J., . . . Lascoux,
M. (2019). Parental legacy, demography, and admixture influenced the evolution of
the two subgenomes of the tetraploid Capsella bursa-pastoris (Brassicaceae). *Plos
Genetics, 15*(2), e1007949. doi:10.1371/journal.pgen.1007949
Landry, C. R., Wittkopp, P. J., Taubes, C. H., Ranz, J. M., Clark, A. G., & Hartl, D. L.
(2005). Compensatory cis-trans evolution and the dysregulation of gene expression in
interspecific hybrids of Drosophila. *Genetics, 171*(4), 1813-1822.
doi:10.1534/genetics.105.047449
Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology, 15*(12). doi:ARTN 550
10.1186/s13059-014-0550-8
Mallet, J. (2005). Hybridization as an invasion of the genome. *Trends in Ecology &
Evolution, 20*(5), 229-237. doi:10.1016/j.tree.2005.02.010
Mallet, J. (2007). Hybrid speciation. *Nature, 446*(7133), 279-283. doi:10.1038/nature05706
Marques, D. A., Meier, J. I., & Seehausen, O. (2019). A Combinatorial View on Speciation
and Adaptive Radiation. *Trends in Ecology & Evolution, 34*(6), 531-544.
doi:10.1016/j.tree.2019.02.008
McManus, C. J., Coolon, J. D., Duff, M. O., Eipper-Mains, J., Graveley, B. R., & Wittkopp,
P. J. (2010). Regulatory divergence in Drosophila revealed by mRNA-seq. *Genome
Res, 20*(6), 816-825. doi:10.1101/gr.102491.109
Mugal, C. F., Wang, M., Backstrom, N., Wheatcroft, D., Alund, M., Semon, M., . . . Ellegren,
H. (2020). Tissue-specific patterns of regulatory changes underlying gene expression
differences among Ficedula flycatchers and their naturally occurring F1 hybrids.
Genome Res, 30(12), 1727-1739. doi:10.1101/gr.254508.119
Muller, H. J. (1942). Isolating mechanisms, evolution, and temperature. *Biol Symp., 6*, 71-
125.
Paronetto, M. P., & Sette, C. (2010). Role of RNA-binding proteins in mammalian
spermatogenesis. *Int J Androl, 33*(1), 2-12. doi:10.1111/j.1365-2605.2009.00959.x
Phillips, B. T., Williams, J. G., Atchley, D. T., Xu, X., Li, J. L., Adams, A. L., . . . Hall, T. M.
T. (2019). Mass spectrometric identification of candidate RNA-binding proteins
associated with Transition Nuclear Protein mRNA in the mouse testis. *Sci Rep, 9*(1),
13618. doi:10.1038/s41598-019-50052-z
Ravinet, M., Elgvin, T. O., Trier, C., Aliabadian, M., Gavrilov, A., & Saetre, G. P. (2018).
Signatures of human-commensalism in the house sparrow genome. *Proc Biol Sci,
285*(1884). doi:10.1098/rspb.2018.1246

Renaut, S., Nolte, A. W., & Bernatchez, L. (2009). Gene expression divergence and hybrid
misexpression between lake whitefish species pairs (Coregonus spp. Salmonidae).
Molecular Biology and Evolution, 26(4), 925-936. doi:10.1093/molbev/msp017
Rest, J. S., Morales, C. M., Waldron, J. B., Opulente, D. A., Fisher, J., Moon, S., . . .
Dedousis, D. (2013). Nonlinear fitness consequences of variation in expression level
of a eukaryotic gene. *Molecular Biology and Evolution, 30*(2), 448-456.
doi:10.1093/molbev/mss248
Rieseberg, L. H., Archer, M. A., & Wayne, R. K. (1999). Transgressive segregation,
adaptation and speciation. *Heredity (Edinb), 83 (Pt 4)*, 363-372.
doi:10.1038/sj.hdy.6886170
Rieseberg, L. H., Raymond, O., Rosenthal, D. M., Lai, Z., Livingstone, K., Nakazato, T., . . .
Lexer, C. (2003). Major ecological transitions in wild sunflowers facilitated by
hybridization. *Science, 301*(5637), 1211-1216. doi:10.1126/science.1086949
Rowe, M., Whittington, E., Borziak, K., Ravinet, M., Eroukhmanoff, F., Saetre, G. P., &
Dorus, S. (2020). Molecular Diversification of the Seminal Fluid Proteome in a
Recently Diverged Passerine Species Pair. *Molecular Biology and Evolution, 37*(2),
488-506. doi:10.1093/molbev/msz235
Runemark, A., Eroukhmanoff, F., Nava-Bolanos, A., Hermansen, J. S., & Meier, J. I. (2018).
Hybridization, sex-specific genomic architecture and local adaptation. *Philos Trans R
Soc Lond B Biol Sci, 373*(1757). doi:10.1098/rstb.2017.0419
Runemark, A., Trier, C. N., Eroukhmanoff, F., Hermansen, J. S., Matschiner, M., Ravinet,
M., . . . Saetre, G. P. (2018). Variation and constraints in hybrid genome formation.
Nat Ecol Evol, 2(3), 549-556. doi:10.1038/s41559-017-0437-7
Runemark, A., Vallejo-Marin, M., & Meier, J. I. (2019). Eukaryote hybrid genomes. *Plos
Genetics, 15*(11), e1008404. doi:10.1371/journal.pgen.1008404
Ryan, M. T., & Hoogenraad, N. J. (2007). Mitochondrial-nuclear communications. *Annu Rev
Biochem, 76*, 701-722. doi:10.1146/annurev.biochem.76.052305.091720
Sanchez-Ramirez, S., Weiss, J. G., Thomas, C. G., & Cutter, A. D. (2021). Widespread
misregulation of inter-species hybrid transcriptomes due to sex-specific and sex-
chromosome regulatory evolution. *Plos Genetics, 17*(3), e1009409.
doi:10.1371/journal.pgen.1009409
Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., . . . Ideker, T.
(2003). Cytoscape: a software environment for integrated models of biomolecular
interaction networks. *Genome Res, 13*(11), 2498-2504. doi:10.1101/gr.1239303
Signor, S. A., & Nuzhdin, S. V. (2018). The Evolution of Gene Expression in cis and trans.
Trends in Genetics, 34(7), 532-544. doi:10.1016/j.tig.2018.03.007
Summers-Smith, J. D. (1988). *The Sparrows: A Study of the Genus Passer*: T & AD Poyser.
Svensson, L., Grant, P. J., Mullarney, K., & Zetterstroem, D. (1999). *Gyldendals store
fugleguide: Europas og middelhavsområdets fugler i felt*: Gyldendal.
Szklarczyk, D., Gable, A. L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., . . . Mering,
C. V. (2019). STRING v11: protein-protein association networks with increased
coverage, supporting functional discovery in genome-wide experimental datasets.
Nucleic Acids Res, 47(D1), D607-D613. doi:10.1093/nar/gky1131
Takahasi, K. R., Matsuo, T., & Takano-Shimizu-Kouno, T. (2011). Two types of cis-trans
compensation in the evolution of transcriptional regulation. *Proc Natl Acad Sci U S A,
108*(37), 15276-15281. doi:10.1073/pnas.1105814108
Taylor, S. A., & Larson, E. L. (2019). Insights from genomes into the evolutionary
importance and prevalence of hybridization in nature. *Nat Ecol Evol, 3*(2), 170-177.
doi:10.1038/s41559-018-0777-y

Team", R. C. (2020). R: A language and environment for statistical computing. Vienna,
575   Austria: R Foundation for Statistical Computing. Retrieved from https://www.R-
576   project.org/
577   Trier, C. N., Hermansen, J. S., Saetre, G. P., & Bailey, R. I. (2014). Evidence for mito-
578   nuclear and sex-linked reproductive barriers between the hybrid Italian sparrow and
579   its parent species. *Plos Genetics, 10*(1), e1004075. doi:10.1371/journal.pgen.1004075
580   Wagner, D. N., Curry, R. L., Chen, N., Lovette, I. J., & Taylor, S. A. (2020). Genomic
581   regions underlying metabolic and neuronal signaling pathways are temporally
582   consistent in a moving avian hybrid zone. *Evolution, 74*(7), 1498-1513.
583   doi:10.1111/evo.13970
584   Wang, L., Wang, S., & Li, W. (2012). RSeQC: quality control of RNA-seq experiments.
585   *Bioinformatics, 28*(16), 2184-2185. doi:10.1093/bioinformatics/bts356
586   Yu, G., Wang, L. G., Han, Y., & He, Q. Y. (2012). clusterProfiler: an R package for
587   comparing biological themes among gene clusters. *OMICS, 16*(5), 284-287.
588   doi:10.1089/omi.2011.0118

Data accessibility and benefit-sharing statement

Raw RNA sequencing reads will be deposited in the SRA (BioProject XXX) upon acceptance

of the manuscript for publication. R script containing code for differential gene expression

analyses and the figures presented in this manuscript is deposited under

github.com/Homap/Expression_sparrow. In this work, all samples taken from Italy and Spain

to Oslo in accordance with the Nagoya protocols. In Spain, all trapping and sampling of birds

was conducted in accordance with Spanish Animal Protection Regulation RD53/2013, and all

methods were approved by the Institutional Commission of Bioethics at the University of

Extremadura (CBUE 49/2011). Birds were sampled in Oslo with permission from Miljø-

direktoratet (2016/2225). In Italy, all birds were sampled with permission from ISPRA

(protocol number 12404) and with permission (no. 305) from Puglia region. Research was

performed in collaboration and cooperation with local researchers from University of

Extremadura. All collaborators are included as co-authors. International collaboration has

been fundamental to this work and has contributed to sharing of scientific results,

cooperation, education, and training.

Author contributions

A.R. and H.P.Y. conceived of the study. A.M. and S.M. raised the experimental F1 hybrids

and M. Ro., M. Ra., A.R., F.E, C.O.G and G-P. S. performed field work and sampling, C.O.G

performed the molecular lab work, and H.P.Y. designed and performed the bioinformatical

analysis based on discussions with A.R, M. Ra., and M.Ro. H.P.Y. drafted the manuscript,

with revisions based on the input from all co-authors.

635   **Table 1.** Number of differentially expressed genes and log2 fold change (LFC) with Padj<
636     $0.05^{\dagger}$ and LFC‡ > 0.32.

	Testis			Ovary		
Comparison	**Significant**	**LFC > 0**	**LFC < 0**	**Significant**	**LFC > 0**	**LFC < 0**
Spanish - house	135	82	53	1382	958	424
Italian - house	3536	1962	1574	22	13	9
Italian - Spanish	3581	1779	1802	1508	394	1114

637   †Results with Padj < 0.01 are reported in Table S4
638   ‡LFC: Log Fold Change
639
640
641
642   **Table 2.** Number and percentage of genes in each inheritance category†.

	Testis	Ovary
Inheritance	**Italy**	**Italy**
Conserved	8883 (77.30%)	10889 (92.26%)
Additive	2 (0.02%)	19 (0.16%)
house dominant	54 (0.47%)	895 (7.58%)
Spanish dominant	25 (0.22%)	7 (0.06%)
Under-dominant	1205 (10.47%)	3 (0.02%)
Over-dominant	1325 (11.53%)	1 (0.008%)

643   †Inheritance classification for differentially expressed genes with Padj < 0.01 is reported in
644   Table S9

653
654 **Figure 1**. Species distribution and sampling locations. Top: Illustrations of male plumage
655 patterns in house, Italian, and Spanish sparrows modified from (Svensson, Grant, Mullarney,
656 & Zetterstroem, 1999). Bottom: A distribution map of house, Italian, and Spanish sparrows
657 throughout Europe and northern Africa (Summers-Smith, 1988). This picture is modified
658 from (Elgvin et al., 2017).

661
662 **Figure 2**. Gene expression in Italian sparrow in comparison to house (Ih) and Spanish (IS)
663 for testis (top) and ovary (bottom). Log2 fold change (LFC) is plotted as a function of mean
664 of normalized counts. Bar plots summarize the proportion of up- and down-regulated genes.
665 Dark orange: up-regulated, light orange: down-regulated.
666

667
668

Figure 3. Inheritance pattern of gene expression in testis and ovary in Italian sparrows. A and
B: Schematic figures representing the classification of inheritance patterns. C and D: Scatter
plots showing shrunken log2 fold change (LFC) in testis between the Italian sparrow with the
house sparrow (Ih) on the *x*-axis and with Spanish sparrow on the *y*-axis (IS), respectively. D:
Scatter plots show shrunken log2 fold change (LFC) in ovary between the Italian sparrow
with the house sparrow on the *x*-axis and with Spanish sparrow on the *y*-axis, respectively.
Grey points in each graph depicts the total number of genes studied for gene expression with
those colored representing the ones significantly different from parental species to be divided
into each of the inheritance categories (Conserved: grey, additive: pink, house dominant:
blue, Spanish dominant: red, transgressive (over-dominant and under-dominant: yellow).
Grey dotted lines indicate the log2 fold-change threshold of 0.32 used for classification of
inheritance pattern.

689
690 **Figure 4**. Gene ontology network represented in over-dominant genes in testis of Italian
691 sparrow. Different colors depicts different significant biological processes. Circle size
692 represents adjusted *P*-value for each node with all Padj < 0.01 where node size represents the
693 term enrichment significance.
694

698