Smell-induced gamma oscillations in human olfactory cortex are required for

accurate perception of odor identity

Qiaohan Yang*[1], Guangyu Zhou*[1], Torben Noto[1], Jessica Templer[1], Stephan

Schuele[1], Joshua Rosenow[2], Gregory Lane[1], Christina Zelano[1]

[1]: Department of Neurology, Feinberg School of Medicine. Northwestern

University. 320 E Superior Ave, Chicago IL. 60611

[2]: Department of Neurosurgery, Feinberg School of Medicine. Northwestern

University. 320 E Superior Ave., Chicago IL. 60611

*, corresponding author: guangyu.zhou@northwestern.edu

This work was funded by National Institutes of Health (NIDCD) grants R01-DC-

016364 and R01-DC-018539 to C.Z.

The authors declare that they have no competing interest.

Abstract

Studies of neuronal oscillations have contributed substantial insight into the mechanisms of visual, auditory and somatosensory perception. However, progress in such research in the human olfactory system has lagged behind. As a result, the electrophysiological properties of the human olfactory system are poorly understood, and in particular, whether stimulus-driven high frequency oscillations play a role in odor processing is unknown. Here, we used direct intracranial recordings from human piriform cortex during an odor identification task to show that three key oscillatory rhythms are an integral part of the human olfactory cortical response to smell: odor induces theta, beta and gamma rhythms in human piriform cortex. We further show that these rhythms have distinct relationships with perceptual behavior. Odor-elicited gamma oscillations occur only during trials in which the odor is accurately perceived, and features of gamma oscillations predict odor identification accuracy, suggesting they are critical for odor identity perception in humans. We also found that the amplitude of high-frequency oscillations is organized by the phase of low frequency signals shortly following sniff onset, only when odor is present. Our findings reinforce previous work on theta oscillations, suggest that gamma oscillations in human piriform cortex are important for perception of odor identity, and constitute a robust identification of the characteristic electrophysiological response to smell in the human brain. Future work will determine whether the distinct oscillations we identified reflect distinct perceptual features of odor stimuli.

Introduction

Oscillations are ubiquitous across mammalian brain networks [1,2,11,3–10] and studies on their spectrotemporal dynamics have contributed important insight into the mechanisms underlying visual, auditory, and somatosensory perception [10,12–16]. Some of the earliest recordings of brain oscillations occurred in the olfactory system of hedgehogs [17], leading to decades of highly productive animal research on oscillations in the mammalian olfactory system with significant work in rabbit and cat piriform [18–22] and more recently mainly focusing on the rodent olfactory bulb [5,8,31–40,23,41–43,24–30]. However, 80 years later, we still lack understanding of the spectral and temporal dynamics of these oscillations in olfactory processing in the human brain, particularly in higher frequency ranges including beta and gamma.

Gamma oscillations, which comprise the synchronized rhythmic patterns of spiking and synaptic inhibition, have been established as an important mechanism in sensory cortical processing in the mammalian brain [44,45]. In the visual cortex, gamma oscillations are locally generated and reflect low-level stimulus attributes [46] including grating size, contrast, spatial structure and color [47–50], and are thought to support visual perception by synchronizing the processing and transfer of information within and across areas of visual cortex [51–53]. Similarly, in the auditory cortex, locally generated gamma oscillations may provide a mechanism to integrate neurons according to the similarity of their receptive fields [54], and are involved in pitch perception and sound discrimination [55,56].

In the olfactory system, odor-driven increases in beta and gamma oscillations have not

been consistently identified in direct recordings from human piriform (olfactory) cortex,

which has been found to exhibit low frequency oscillations in response to odor [57].

However, in mammals and insects, the cellular and network processes underlying

gamma oscillations in the olfactory bulb and cortex have been studied extensively

[19,29,40,58–63], and more recent work has begun to provide insight on their functional

role as well [23,26,32,33,36,37,40,64], suggesting involvement in organization of

sensory information to enable fine odor discrimination and identification [23,65–69]. In

addition to gamma oscillations, beta and theta oscillations have been extensively

studied in the mammalian olfactory system, and are involved in olfactory learning and

respiratory tracking, respectively [18,27,70,71,28,30,33,36,38,40,64,65]. Thus theta,

beta and gamma oscillations constitute three key rhythms that have functional

significance for olfactory processing in mammals [8]. However, against this background

of extensive animal work, the electrophysiological underpinnings of the human olfactory

system are vastly understudied. Only a few intracranial studies have directly measured

odor-induced oscillations in the human brain; in the amygdala [72–76] and piriform

cortex [57]. Human piriform recordings suggest an important role for low frequency (< 8

Hz) oscillations in olfactory processing [57]; theta oscillations were consistently found in

piriform cortex within 200 ms of sniff onset during an odor detection task, and their

features could be used to decode odor identity. However, the full spectral characteristics

of early and later emerging odor responses in human olfactory cortex remain unknown.

Furthermore, the functional role of high frequency oscillations in human piriform cortex

is virtually unexplored.

In this study, we used stereotactic intracranial electroencephalography (iEEG) to test

three main hypotheses about neural responses in human piriform cortex during an

olfactory identification task. First, based on the fundamental role of gamma oscillations

in human sensory perception and the established rhythms in the rodent olfactory

system, we hypothesized that odors would induce theta beta and gamma oscillations in

human piriform cortex. Second, based on the fact that olfactory oscillations are

modulated by sniffing [26,58,62,63,77–79], which is inextricably linked to odor onset

times, we hypothesized that the timing of responses relative to sniff onset would vary

across the different frequency bands. Third, based on established links between

gamma/beta oscillations and sensory coding [27,30,32,33,35,36,38,64], and the fact

that activity in rodent piriform cortex can be used to decode odor identity [80–82], we

hypothesized that human piriform beta and gamma oscillations would emerge only

during trials where odors were accurately identified, and that features of their oscillatory

rhythms would predict odor identification accuracy.

Our results show that odors elicit a stereotyped oscillatory response in human piriform

cortex that is consistent in timing and frequency composition across subjects:

Immediately following odor onset, early theta increases are quickly followed by gamma

and beta increases. The phase of early low frequency responses drives the amplitude of

high frequency rhythms only when odor is present. Finally, the distinct oscillatory

rhythms we identified were differentially related to behavioral performance, suggesting

that these rhythms could potentially give rise to non-interfering representations of

different features of odor stimuli [81,82], though future work is needed to test this

hypothesis.

Results

To examine odor-driven oscillations in human olfactory cortex, we recorded local field

potentials from 7 participants who took part in an odor identification task. Each

participant's clinical electrode coverage included piriform cortex (**Fig 1A**). The olfactory

task was performed during clinical recording of ongoing electrophysiological activity, at

sampling rates ranging between 500 to 2000 Hz, using a 256-channel clinical EEG

acquisition system (Nihon Kohden). Each trial began with an auditory cue signaling that

odor would be presented, and providing the potential identity of the odor (rose or mint).

After a jittered delay (3–7 s), odor was presented to the participant while sniff onset was

precisely measured via a pressure sensor at the nose (Salter Labs). On each trial,

participants identified the odor by indicating whether or not it matched the identity

prompt (**Fig 1B**). To isolate effects driven by odor from those driven by inhalation, which

also impacts oscillations in human piriform cortex [78,83–85], we analyzed and directly

compared LFPs during inhalations that contained odor and those that did not, resulting

in two experimental conditions: odor and no-odor (**Fig 1B**, areas shaded in pink and

blue).

Fig 1. Electrode contact locations and experimental design. (A) The location of the piriform cortex

electrode contact (red dot) is shown on each participant's (P1–P7) brain image. L, left hemisphere. (B)

Schematic illustration of the olfactory task, showing odor and no-odor conditions drawn on an illustrative

breathing signal (black line). The light blue area indicates experimental trials (starting from cue delivery to

the end of the breathing cycle in which odor was delivered), and the pink area indicates between trials.

The black boxes indicate the analysis time windows for odor and no-odor trials. The darker blue area

indicates odor delivery.

Odor elicits oscillations in theta, beta and gamma frequency bands in human piriform cortex

To test the hypothesis that odor elicits LFP oscillations in the theta, beta and gamma

frequency bands in human piriform cortex, we first conducted a time-frequency analysis

combining data from all trials and participants. We computed spectrograms aligned to

sniff onset, for odor and no-odor conditions separately. In the odor condition, we found

statistically significant increases in theta, beta and gamma frequency bands (**Fig 2A**, left

panel) (theta peak: 4.72 Hz, max z score = 14.07; beta peak: 18.98 Hz, max z score =

8.11; gamma peak: 84.94 Hz, max z score = 4.88). In the no-odor condition, we found

smaller but significant increases in theta only (**Fig 2A**, middle panel; peak: 4.02 Hz, max

z score = 3.44). To quantify the difference between odor and no-odor conditions, we

conducted a direct statistical comparison by permuting condition labels to generate a

map of z-normalized amplitude differences (**Fig 2A**, right panel). We found that theta,

beta and gamma amplitudes were significantly higher in the odor compared to the no-

odor condition (permutation test; $P < 0.05$, FDR corrected).

Fig 2. Odor induces theta, beta and gamma oscillations in human piriform cortex. (A) Results from combined analysis showing z-normalized amplitude spectrograms for no-odor and odor conditions (left and middle panels), and the difference between the two conditions (right panel). Average respiratory signal is shown on each panel (overlaid red line). Dashed line represents inhale onset. For odor and no-odor spectrograms, the black contour lines indicate statistical significance of permutation tests against pre-cue baseline ($P < 0.05$, FDR corrected). For the difference spectrogram, the black contour lines indicate statistical significance of permutation of condition labels ($P < 0.05$, FDR corrected). The far-right panel shows average z scores from 0–4 s post sniff across frequencies for odor and no-odor trials. B) Individual theta, beta and gamma summed magnitudes over 0–2 s post sniff. Each dot represents data from a single participant, and lines connect data within the same participant. Asterisk indicates statistical significance (two-tailed paired t test; $P < 0.05$). (C) Z-normalized amplitude spectrograms for each participant for no-odor (left) and odor (middle) conditions, with the average respiratory signal shown on each panel (overlaid red line). Dashed lines represent inhale onset. Black contour lines indicate statistical significance of permutation tests against pre-cue baseline ($P < 0.05$, FDR corrected). Raw broadband and beta (13–30 Hz) time series of a representative trial is shown on the right. The source data for panels (A)–(C) are available in S1 Data.

To confirm our findings at the individual level, we next computed sniff-aligned spectrograms, in each participant separately (**Fig 2B–C**). To quantify responses in each participant, we used standard human EEG frequency band definitions (theta: 4–8 Hz; beta: 13–30 Hz; gamma: > 30 Hz), and looked for responses occurring within these standard frequency ranges in each participant. We computed a direct statistical comparison between the two conditions across participants, which showed that amplitudes in these three frequency bands were significantly stronger across participants during the odor compared to the no-odor condition (**Fig 2B**) (two-tailed

paired t test, theta: T_6 = 3.19, P = 0.0187; beta: T_6 = 4.17, P = 0.0059; gamma: T_6 =

2.98, P = 0.0245). Notably, in every single participant, theta, beta and gamma

amplitudes were larger during the odor compared to the no-odor condition (**Fig 2B**,

each dot is the result from one participant and lines connect same-participant results

between conditions). Though there was some variation across individuals in the

frequency of responses, these effects were evident in the individual spectrograms of

most participants (**Fig 2C;** black outlines indicate statistical significance via permutation

test; P < 0.05, FDR corrected), and in the minimally-processed signals shown next to

each individual spectrogram (**Fig 2C**, right). In each participant, theta, beta and gamma

responses were maximal inside piriform cortex, with significantly reduced responses

outside of piriform on the same depth wire (two-tailed paired t test; theta: P = 0.0045, T_6

= 4.42; beta: P = 0.014, T_6 = 3.44; gamma: P = 0.0070, T_6 = 4.02) (**S1 Text**; **S1 Fig 3A**).

To further visualize the spatial distribution of responses in each frequency band, we

created heat maps of responses across all implanted electrodes in all participants,

projected onto an axial slice through piriform cortex (**S1 Fig 3B**). This allowed

visualization of response magnitudes in the anterior-posterior direction, with the

superior-inferior axis flattened. A hot spot of increased activity can be seen

corresponding to piriform cortex (**S1 Fig 3B**, pink boxed areas; axial slice contains

piriform cortex).

To address the possibility that the responses we observed were due to the steep slopes

of sensory-evoked potentials, we conducted an analysis to separate the spectrograms

into odor-evoked and odor-induced effects [86–88]. In this analysis, the odor-evoked

spectrogram includes only the phase-locked power, which contributes to the sensory-

evoked potential, whereas the odor-induced spectrogram includes only the non-phase-

locked power, which is oscillatory and is not related to the ERP [89,90] (see methods).

Results of this analysis showed that odor-induced oscillations were not eliminated when

the phase-locked components of the signal were removed (**S1 Fig 1**) and thus were not

due to the steep slope of sensory-evoked potentials.

In rodents, odor has been shown to elicit responses in multiple gamma subbands

[8,64,91]. To examine gamma effects more closely and to look for multiple gamma sub-

bands in humans, we re-computed individual spectrograms using a linear, as opposed

to log, frequency scale. We found that most subjects showed odor-induced oscillations

right around or just below 30 Hz. Most subjects also showed a higher frequency

broadband response, around 90–150 Hz (**S1 Fig 2**).

Together, these results suggest that odor consistently induces distinct theta, beta and

gamma oscillations in human piriform cortex.

The time course of olfactory cortical oscillations varies across frequency bands

The spectrograms in **Fig 2** showed apparent variable time courses of oscillations

relative to sniff onset across frequency bands (**Figs 2A and 2B**). Theta appeared to

emerge and dissipate soonest, with beta and gamma emerging later and persisting

longer. To quantify these differences, we conducted a series of analyses to characterize

the timing of responses across frequency bands. These included a percent-change

analysis to examine oscillatory increases over time, a bootstrapping analysis to

minimize potential impact of noisy trials, a cluster-based analysis to quantify the exact

timing and magnitude of continuously significant increases in oscillatory amplitude, and

a circular-distribution analysis to examine oscillatory peaks over respiratory phase.

Fig 3. Odor-induced theta, beta and gamma oscillations have different time courses relative to sniff onset.

(A) Percent-amplitude-change time series for odor (blue line) and no-odor (red line) conditions for each

frequency band. Shaded areas indicate one standard deviation from the mean across all trials. Thick

black lines above indicate statistically significant differences between conditions (FDR corrected *P* < 0.05,

paired t test). (B) Bootstrap colormap of z-normalized amplitude for each frequency band, with results

from each repetition stacked along the Y axis. Vertical stripes of consistent color indicate consistent

amplitudes across subsampled sets of trials. (C) Colored shaded areas represent clusters of statistical

significance of the trial-by-trial baseline-corrected amplitude time series, for each frequency band (cluster

correction test *P* < 0.05, corrected). Upper boundaries of the shaded areas represent the t statistic from

the one-sample t test of baseline-corrected amplitude at each time point against 0. Scales for t statistics

are shown to the left of each cluster. Overlaid darker dots indicate the average timing of the bootstrapped

distribution of peak amplitude occurrence for each frequency band, for each participant. Each dot

represents data from one participant. Dots from the same participant are connected by black lines. On the

right, cluster masses in relation to the permuted null distribution for each frequency band are shown, with

actual values represented by the vertical red line. (D) Respiratory phase angle distributions corresponding

to bootstrapped peak amplitudes for each frequency band. Scale of the polar histogram represents pdf of

the distribution. Darker lines overlaid on the distribution indicate the mean vector of the phase distribution.

Angle and radius represent averaged phase angle and PLV value respectively. The source data for

panels (A)–(D) are available S2 Data.

In our first analysis, we calculated the percent change in amplitude for each frequency

band at each time point, for each condition (**Fig 3A**). This allowed us to determine the

timing of the emergence of significant differences between odor and no-odor conditions

(two-tailed paired t test; $P < 0.05$, FDR corrected). This analysis showed distinct

temporal dynamics of odor responses for each frequency (**Fig 3A**, black lines above

each panel). Theta oscillations were the earliest to emerge and dissipate, beginning 126

ms prior to sniff onset and ending 516 ms after sniff onset. Gamma oscillations emerged

next, 116 ms after sniff onset, intermittently persisted through exhalation, and dissipated

3678 ms after sniff onset. Beta oscillations emerged last, 144 ms after sniff onset, and

persisted longest, ending 3786 ms after sniff onset.

In a second analysis, designed to minimize the contribution of noisy trials and to

statistically evaluate the time courses, we tested whether differences in temporal

dynamics of responses across frequency bands were stable across resampled sets of

trials. We conducted a bootstrapping analysis to create separate profiles of the timing of

emergence of responses for each frequency band. We created a z-normalized time

series of resampled mean values for each frequency band, represented as a colormap

(**Fig 3B**, y-axis is each bootstrap repetition). We found stable time courses across

repetitions, indicated by stripes of increased amplitude at particular times for each

frequency band. To quantify the timing of amplitude increases, we performed a cluster-

based statistical analysis; for each frequency band, we generated a t-statistic at each

time point for all z-normalized odor trials, and then conducted a permutation-based

analysis to correct for cluster size. We found clusters that corresponded well with the

spectrograms shown in Fig 2, with the percent-change analysis, and with the bootstrap

analysis (**Fig 3C**). We found one significant cluster for the theta band extending 266 ms

pre sniff to 810 ms post sniff (permutation test; $P < 0.0001$), one for the beta band

extending from 238 ms to 3524 ms post sniff ($P < 0.0001$), and four separate clusters

for the gamma band (326 ms to 452 ms post sniff, $P = 0.0346$; 764 ms to 892 ms post

sniff, $P = 0.0145$; 1374 ms to 1642 ms post sniff, $P < 0.0001$; 1902 ms to 2016 ms post

sniff, $P = 0.0451$). We subsequently conducted the same bootstrapping analysis at the

single subject level, and plotted the average time point of each subject's bootstrapped

peak distribution on top of the significant clusters in Fig 3C (see overlayed dots, each

representing the peak value for a single participant, with lines connecting each

participant's values across frequencies). Although single participant data was more

variable, beta and gamma peaks occurred later than theta peaks for almost all subjects.

These results suggest that the unique time courses of odor responses across

frequencies were unlikely to be caused by noise, spikes, and other artifacts.

In a third analysis, designed to account for individual respiratory differences in the time

domain, we examined the circular distributions of oscillatory peaks over the respiratory

phase. On each repetition of a bootstrapping analysis, the peak amplitude and

corresponding phase angle were calculated from the average LFP and respiratory time

series. We found that theta peaks narrowly aggregated at early stages of inhale (mean

phase angle = –0.67 rad, PLV = 0.99, Rayleigh's z = 998.05, P < 0.0001) (**Fig 3D**, left),

consistent with previous findings in humans [57]. By contrast, beta peaks consistently

aggregated at the exhale trough (mean phase angle = 2.98 rad, PLV = 0.80, Rayleigh's

z = 647.78, P < 0.0001) (**Fig 3D**, middle). Gamma peaks were more broadly distributed,

with most centered around inhale peak or the transition between inhale and exhale

(mean phase angle = 0.57 rad, PLV = 0.63, Rayleigh's z = 401.69, P < 0.0001) (**Fig 3D**,

right). Combined, these findings suggest that following presentation of odor, theta

oscillations increase earlier than beta and gamma oscillations, and that while theta and

gamma oscillations are maximal during inhalation, beta oscillations peak during exhale.

Gamma and beta oscillations are required for accurate odor identification.

Piriform cortex has been postulated to be a major driver of gamma oscillations in the brains of rodents [18,92] and cats [19–22,93]. Furthermore, numerous rodent studies suggest that beta and gamma oscillations relate to odor learning and discrimination, through both local odor coding [23,33,35,40,64,94] and recruitment of larger-scale networks [27,30,35–37,64]. We therefore hypothesized that beta and gamma oscillations in human piriform cortex would correlate more strongly with task performance than theta. We conducted three separate analyses to explore the relationship between odor-driven LFP oscillations and odor identification accuracy. These included a time-frequency analysis to look for gross differences between correct and incorrect trials, a correlation analysis to look for a relationship between accuracy and oscillations across trials, and a machine learning analysis to determine whether amplitudes of different frequency responses could predict accuracy.

Fig 4. Beta and gamma oscillations predict odor identification accuracy. (A) Z-normalized spectrograms for correct (left) and incorrect (right) trials separately. Averaged respiratory signals are overlayed on each spectrogram (red lines). Dashed lines represent sniff onset. Black contours indicate statistical significance of permutation test against pre-cue baseline ($P < 0.05$, FDR corrected). (B) Amplitudes within each frequency band, for correct and incorrect trials separately. The upper panels show the distribution of z score amplitude values for each trial for the theta, beta and gamma bands, for correct (blue) and incorrect (red) trials. The lower left panel shows box plots for the bootstrapped distributions of the difference between correct and incorrect trials. Boxes represent the 25th to 75th percentile of each distribution, the central marker indicates median, and whiskers extend to the extremes of the data excluding outliers. Colored dots represent raw difference values for each distribution. The lower right panels show individual participant's mean response amplitudes in the theta, beta and gamma bands, for correct and incorrect trials separately. Each dot represents data from a single participant, bars indicate the mean across participants and error bars are standard error across participants. Stars indicate statistical significance

(paired, two-tailed t tests). See Fig 2B for the color code of single participant dots. (C) Bar plots of

bootstrapped Pearson's correlation coefficients computed between oscillatory amplitudes and task

accuracy for each frequency band, over the inhale and exhale time periods separately. Error bars indicate

95% confidence interval of the mean. Dashed line indicates FDR threshold for significant r value. Dots

represent single correlation coefficient calculated from each bootstrap. (D) Scatter plots show

bootstrapped percent change relative to baseline for correct and incorrect trials, during inhale and exhale

time periods for each frequency band. Each dot is the average result from one bootstrap repetition. (E)

Receiver operating characteristic (ROC) curves of linear binary SVM classifier applied to each of the

scatter plots in panel C. Black dots represent optimized classifier, area under the curve (AUC) is indicated

for each ROC curve. The source data for panels (A)–(E) are available in S3 Data.

First, in a combined time-frequency analysis, we computed odor-onset aligned

spectrograms during correct and incorrect trials separately, in order to look for

differences across frequency bands. We found that during trials in which the participant

correctly identified the odor, there were statistically significant increases in theta, beta

and gamma band amplitudes (**Fig 4A**, left panel; $P < 0.05$, FDR corrected). By contrast,

during trials in which the participant failed to identify the odor, statistically significant

increases were found only in the theta band, with no such increases in the beta or

gamma bands (**Fig 4A**, right panel; $P < 0.05$, FDR corrected; **Fig 4B**, upper panels and

lower left panel show the distribution of single trial theta, beta and gamma values for

correct and incorrect trials). Since most participants performed above chance on the

task, there were a larger number of correct trials than incorrect trials (252 versus 71). To

account for this difference, we conducted a resampling analysis with 200 repetitions in

which correct and incorrect trials were resampled with replacement to include the same

number of trials on each repetition, and we computed the distribution of the differences

in amplitude (correct – incorrect) within each frequency band (**Fig 4B**, lower left panel).

We found statistically significantly larger responses during correct compared to incorrect

trials for beta and gamma bands only, with no significant differences in the theta band.

(One-sample t test against 0 for the difference between correct and incorrect trials:

theta: T_{199} = 1.21, P = 0.2247; beta: T_{199} = 35.55, P < 0.0001; gamma: T_{199} = 42.61, P <

0.0001. Paired sample t tests for differences between frequency bands: theta vs beta:

T_{199} = –41.02, P < 0.0001; beta vs gamma: T_{199} = 17.36, P < 0.0001; gamma vs theta:

T_{199} = 37.10, P < 0.0001.). This combined resampling analysis was designed to account

for differences in numbers of correct and incorrect trials, which had the advantages of

ensuring a balanced comparison across conditions and a robust number of incorrect

trials. However, the contribution of individual subjects to the observed effects was

unclear. To account for this, we quantified theta, beta and gamma responses for correct

and incorrect trials separately, in each participant. Since some participants had no, or

just a few, incorrect trials, we limited this analysis to include participants with more than

3 trials of each condition (resulting N = 5). Though numbers of incorrect trials were

relatively low for some subjects, results of this analysis confirmed the combined

findings: responses in all three frequencies were significantly above zero for correct

trials (theta: T_4 = 4.12, P = 0.014 ; beta: T_4 = 2.87, P = 0.045; gamma: T_4 = 4.71, P =

0.0092), whereas only responses in theta band were significantly above zero for

incorrect trials (theta: T_4 = 7.28, P = 0.0019; beta: T_4 = 1.77, P = 0.15; gamma: T_4 =

1.41, P = 0.23) (**Fig 4B**, lower right panels). These findings were also apparent in

individual participant spectrograms (**S4 Fig 4B**).

Second, we looked for a relationship between task performance and responses in each

frequency band on a trial-by-trial basis. To account for differences within frequency

bands over time, we conducted this analysis during the inhale and exhale periods

separately. Trials were resampled to generate a distribution of correlation coefficients

representing the relationship between response amplitudes and behavioral accuracy for

each frequency band across the resampled trial sets. Significant correlations between

accuracy and amplitude were evident in the beta and gamma bands during both inhale

and exhale (50 bootstraps, t test against 0, FDR corrected threshold for significant

correlation coefficient; beta during inhale: T_{49} = 31.25, P < 0.0001; beta during exhale:

T_{49} = 17.64, P < 0.0001; gamma during inhale: T_{49} = 23.70, P < 0.0001; gamma during

exhale: T_{49} = 28.25, P < 0.0001.), but no such correlation was found in the theta band

(theta during inhale: T_{49} = −2.04, P = 0.97; theta during exhale: T_{49} = −12.90, P = 1). In

line with this, a direct statistical comparison across frequency bands showed that

correlations were significantly stronger between behavioral accuracy and amplitude for

beta and gamma bands compared to theta band during both inhale and exhale time

periods (**Fig 4C**) **(**two way repeated measures ANOVA: main effect of time window: $F_{1,}$

$_{49}$ = 111.08, P < 0.0001; main effect of frequency band: $F_{2, 98}$ = 547.98, P < 0.0001.

Paired sample t test across frequency bands during inhale for theta vs beta: T_{49} = −

29.37, P < 0.0001; for beta vs gamma: T_{49} = 3.11, P = 0.0031; for gamma vs theta: T_{49}

= 24.95, P < 0.0001; paired sample t test across frequency bands during exhale for

theta vs beta: T_{49} = −26.51, P < 0.0001; for beta vs gamma: T_{49} = −11.13, P < 0.0001;

for gamma vs theta: T_{49} = 33.95, P < 0.0001), also see **S4 Fig 4A**.

Finally, we used machine learning to perform a classification analysis on inhale and

exhale amplitudes for each frequency band, to determine whether these features could

predict task accuracy. To this end, we applied a binary linear support vector machine

(SVM) classifier to the bootstrapped data, plotted the receiver operative characteristic

(ROC) curves and calculated the area under curve (AUC), which was used together

with optimized classifier accuracy as the predictor of data separability. We found that

beta and gamma amplitudes were highly successful at separating the trials by accuracy,

while theta amplitudes were less so (**Fig 4D**). Specifically, gamma accuracy was

maximal at 91.5%, beta accuracy was also high at 91%, whereas theta accuracy was

lower, at 66.5%. This same procedure was applied to shuffled data, which resulted in

chance performance (accuracy = 52%), validating our methods. These findings

correspond well with the results from the two previous analyses, and, taken together,

suggest that beta and gamma band amplitudes are more strongly correlated with task

performance than theta amplitudes.

Odor-elicited high-frequency oscillations are not driven by attention.

Because the odor condition occurred during experimental trials and the no-odor

condition occurred between experimental trials, responses during the odor condition

could potentially have been impacted by attentional states that were not present during

the no-odor condition. Previous human and animal work suggests that olfactory cortical

responses are modulated by attention [95,96], particularly in lower frequency LFPs

[97,98]. To control for potential effects of attention on our findings, we analyzed

odorless sniffs taken in attended states during experimental trials. Each trial began with

a cue, followed by a delay (3–7s) prior to presentation of odor. Therefore, during many

trials, sniffs of clean air occurred between the cue and odor (**Fig 5A**). These sniffs

occurred during attended states, but did not contain odor, thus providing a means to

isolate the effects of attention from the effects of odor. Because a sniff did not always

occur between the cue and the odor, the number of trials in the attended odorless

condition was lower than the odor condition. Therefore, we resampled odor trials to

match the number across conditions. We computed spectrograms aligned to attended

odorless sniffs and compared them to those aligned to odor sniffs (**Fig 5B**, left and

middle). We found that attended odorless sniffs induced oscillations in low frequencies

(below 8 Hz) only ($P < 0.05$, FDR corrected), and found no stable higher frequency

oscillations (> 8Hz). We then computed the z-normalized difference across conditions

(**Fig 5B**, right) and found significant differences between attended odor trials and

attended no-odor trials in the theta, beta and gamma bands, such that oscillations in all

three frequency bands were increased when odor was present, even after removing the

effects of attention (**Fig 5B**, right; $P < 0.05$, FDR corrected). Since theta oscillations

were still detected in the attended no-odor condition, these findings suggest that the

increases in beta and gamma bands were independent of attention, whereas theta band

increases reflected, at least to some degree, attentional states.

Fig 5. High frequency odor-elicited oscillations are not driven by attention. (A) Schematic illustration of attended and unattended conditions. Attended no-odor trials were defined as breaths taken during experimental trials, between the cue and the odor. No odor was present, but the participant was in an attended state. Blue area indicates the time of experimental trials, and pink areas indicates the time between trials. Black boxes indicate the analysis time windows for the different conditions. (B) Z-normalized spectrograms of attended odor and attended no-odor trials, and the difference between the two. Average respiratory signals are overlayed for each condition (red lines). Dashed line represents inhalation onset. Black contours indicate statistical significance (permutation tests as described elsewhere, $P < 0.05$, FDR corrected). The source data for panel (B) is available in S4 Data.

Low frequency phase modulates high frequency amplitude during inhale when odor is present.

Previous work suggests that odor-induced theta oscillations predict odor identities, even during a detection task that does not require participants to identify odors [57]. Here, we found that theta rhythms in our data appeared to have a relationship with beta responses, which appeared to oscillate at around 5 Hz during inhale. Specifically, the overall magnitude of beta oscillations increased and decreased rhythmically with a

frequency in the theta range, particularly in the first 1.5 to 2 seconds of the response

(**Fig 6A**, gray box). Based on this observation, we hypothesized that theta oscillations

might organize later-emerging higher frequency oscillations through phase modulation.

To estimate modulation of higher frequency amplitude by theta phase, we calculated the

modulation index (MI) [99] using theta phase as the modulating signal and a range of

high frequencies (13–150 Hz) as the amplitude signals, for both odor and no-odor

conditions (**Fig 6B**). We found that in the 1s time window following sniff onset, theta

phase significantly modulated beta and gamma oscillations during the odor condition (*P*

< 0.05, FDR corrected for MI in all frequency bands; peak MI at 16.16 Hz, mean z score

across all frequencies = 7.52), but not during the no-odor condition (mean z score

across all frequencies = 2.02).

Fig 6. Theta phase modulates higher frequency amplitudes during inhale, when odor is present. (A)

Apparent rhythmic beta amplitude modulations at the theta frequency. With effects of attention subtracted

out, the average Z score of the amplitude of the odor responses over time is displayed for beta (green)

and theta (blue) frequency bands. The overall magnitude of beta oscillations increased and decreased

rhythmically with a frequency in the theta range, particularly in the first 1.5 to 2 seconds of the response

(gray box). (B) Phase-amplitude coupling between theta phase and higher frequency amplitudes. Z-

normalized modulation index (MI) of theta phase and higher frequency amplitudes (13 to 150 Hz), during

0–1s post sniff for all trials in the odor and no-odor conditions. Dashed line indicates FDR threshold for

significant MI ($P < 0.05$, FDR corrected). The source data for panel (A) and (B) are available in S4 Data.

Our phase-amplitude coupling analyses was performed during a window in which a

sensory-evoked potential could have occurred, and therefore the derived phase-

amplitude coupling could have been driven by the evoked potential having a steep

slope. To control for this possibility, we conducted a permutation analysis [100] in which

we shuffled the trial-by-trial relationship between theta phase and higher frequency

amplitude to test whether the modulation effect was due to the exact trial-by-trial

relationship or rather induced by a steep slope during each trial. We normalized the

observed MI to the trial-order-shuffled null distribution and still found significant effects

for the odor condition but not the no-odor condition (**S5 Fig 5**). This finding suggests

that higher frequency oscillations were modulated by theta phase during the early

period of sniffing, when theta amplitudes were significantly increased.

Theta, beta and gamma oscillations are present across a range of olfactory tasks

Studies in rodents suggest that different olfactory tasks may elicit distinct neural

oscillatory signatures [32,91]. Whether this is the case in the human olfactory system is

unknown. It is therefore possible that the oscillations we observed in the theta, beta and

gamma ranges were due to the nature of the olfactory task (identity matching to a cue,

as opposed to detection [57] or pure identification). To explore this possibility, we

present data from two additional subjects who performed 3 different olfactory tasks,

including a detection task, an edibility assessment task and an odor naming task. This

allowed us to determine if the nature of the task was the main driver of the effects we

observed, rather than the odors. None of the tasks involved matching to a previous cue.

For each task, trials began with a countdown to a cued sniff. Subjects self-initiated odor

delivery using a manual hand-held olfactometer that allowed for precise, sniff-controlled

timing of odor delivery. Following odor presentation, during the detection task, subjects

indicated whether or not they detected an odor via button press. During the edibility

assessment task, subjects indicated whether or not the odor was edible via button

press. During the naming task, subjects indicated whether or not they could name the

smell via button press, and then they spoke the name, if possible. All three of these

tasks resulted in highly similar oscillatory responses to odor in both subjects (**Fig 7**).

Both subjects, across all three task types, showed rapid increases in theta, followed by

oscillatory increases in beta and gamma that persisted into exhalation. Detailed

analyses of the responses to these different tasks is beyond the scope of this

manuscript; results are displayed here solely for the purpose of this control analysis,

which shows that the oscillations we observed in our original data set are present

across a range of olfactory tasks, and were therefore unlikely to have been driven solely

by the nature of the task.

Fig 7. Odor-induced theta, beta and gamma oscillations are evident across a range of olfactory tasks.

Shown are spectrograms from two participants (P8 and P9) who performed three different olfactory tasks

including a detection task (top row), an edibility assessment task (middle row) and an odor naming task

(bottom row). Short-dashed vertical line indicates sniff onsets. Average respiratory signals from each

participant are shown as a red overlaid line on each spectrogram. Black contours indicate statistical

significance ($P < 0.05$, FDR corrected). Below each panel is broad-band and beta-filtered time-series from

one trial for that participant. The source data is available in S4 Data.

Discussion

We found that odor induces oscillations in human piriform cortex in the theta, beta and

gamma frequency bands, each with distinct temporal dynamics, forming the basis of a

characteristic human olfactory cortical response. While theta oscillations rapidly

emerged, peaked and dissipated within the first 500ms of inhalation, gamma and beta

oscillations emerged and peaked later, extending into the exhale period (see **Fig 2**, and

Fig 3B and 3D). These findings suggest that the spectrotemporal characteristics of

oscillations in piriform cortex in response to odor are similar in rodents and humans.

While our study cannot speak to the origin of these oscillations, it allows us to consider

the possibility that knowledge of the drivers of LFP oscillations in rodents [8] may be

applicable to humans, despite differences between species [101]; for example that beta

and gamma oscillations in piriform cortex rely on an intact olfactory tract from the bulb

[38], with important implications for our understanding of the top down or bottom up

nature of olfactory processing [102–104] . We further found that the strength of beta and

gamma oscillations was significantly more correlated with odor identification accuracy

than theta oscillations, suggesting beta and gamma rhythms are important for odor

identification in humans. Our work serves to unite results from previous intracranial

human studies that have reported odor-induced oscillations at both low and high

frequencies, with variable results [57,72–76,104].

Interestingly, we found that gamma rhythms were more tightly correlated to odor

identification accuracy than beta during exhale (see Fig 4C). Since gamma rhythms are

thought to reflect local computations [23,32,33,36,40,44,61,64], and beta oscillations

are thought to reflect long-range network interactions [27,30,35,36,61,64], local

computations in piriform cortex may have particular significance during exhale, though

our study does not directly address this and future studies on this topic are needed. Our

data also highlight an important role for theta oscillations in human piriform cortex, in

agreement with previous studies [57], as we found that theta phase modulated beta and

gamma amplitudes during inhalation, only when odor was present (see Fig 4). Rodent

work has identified functionally distinct subpopulations of piriform neurons that

differentially encode the intensity and identify of odor [82][81][105]. It is possible that the

distinct oscillations we identified in human piriform cortex represent distinct populations

of neurons that encode different features of the odor percept with unique temporal and

spectral properties, though future work is required to determine this. In particular, the

addition of single unit recordings may allow us to determine whether the different

oscillations are tied to different subpopulations of piriform neurons.

Theta, beta and gamma oscillations have been extensively studied in the olfactory

systems of rodents, providing insights into the functional roles of these oscillations in

olfactory perception [23,24,35–39,41,43,60,64,65,26,68,106,107,27–31,33,34].

Furthermore, an impactful body of rodent work has shown that the timing of olfactory

responses relative to sniffing behavior is an important mechanistic feature of odor

coding, enabling accurate olfactory percepts to form quickly [33,41,116,108–115].

Though much of this work has focused on the olfactory bulb, responses in rodent

piriform cortex also include temporal modulations [81,113,117], and its oscillations

reflect sniff phase [26,62,78]. In contrast to the extensive and elegant body of rodent

work, oscillations in human piriform cortex have been vastly understudied. Here,

dovetailing with animal work, we found that theta, beta and gamma oscillations emerged

at different times relative to sniff onset. The link between sniffing behavior and

oscillations in rodents, combined with dramatic differences in sniffing behavior between

rodents and humans, raises some intriguing questions about the timing of odor

responses in humans. When sampling odors, rodents engage in multiple sniffing

strategies, and may sniff at rates up to 12 Hz [118–120]. Humans might at times

increase their sniffing frequency a bit when smelling an odor, but often sample odors in

a single sniff, and aren't able to reach sniffing frequencies of 12 Hz (though this has not

yet been directly tested). Despite these large differences in sniffing behaviors across

species, we found surprisingly similar time courses of odor responses in piriform cortex.

For example, Frederick et al. [64] found that beta oscillations in rodent piriform cortex

emerged at around 200 ms following odor onset, similar to our findings in human

piriform cortex. We found that beta oscillations peaked and persisted longer than

gamma oscillations, also in agreement with findings in rodents [59]. In rodents, the theta

rhythm is actively shaped by sniffing behavior, which, in turn, shapes higher frequency

oscillations. In humans, sniffing behavior almost never occurs at frequencies near the

theta range, yet we found that theta oscillations shape higher frequency oscillations in

the human olfactory system during inhalation, as is the case in rodents. This may

suggest that the time scale of some odor-induced oscillations is not entirely dependent

on breathing rate, and that network properties that support these rhythms may be

conserved across species. Our findings support the idea that the theta rhythm serves as

the "internal reference clock" of odor information processing [57], though future studies

combining data from both humans and rodents, and potentially from noninvasive human

olfactory bulb recordings [121] could contribute much-needed insights to answer this

question.

An important consideration in comparing our findings to those in rodents is the nature of

the task. The two most commonly used olfactory tasks in rodent studies are the Go-No

Go (GNG) task and the 2-Alternative Forced Choice (2AFC) task [64,122–124]. The

GNG task requires a response only when a target odor is present, whereas the 2AFC

task requires a different response to each odor, depending on the identity of the odor.

Our task is more similar to the 2AFC task, as it requires subjects to respond differently

to each odor, depending on the identity of the odor. However, it is generally challenging

to compare tasks between species—both GNG and AFC tasks require training of

rodents, and include rewards during each trial. This is less common for human studies,

in which experimenters directly instruct subjects on the task, and rewards are not

required for subjects to learn the rules of the task. Several rodent studies have shown

differences in oscillatory signatures across different olfactory tasks and difficulty levels

[40,64]. It is therefore possible that differences in task-related responses could impact

differences between correct and incorrect trials between species [125–127]. A

worthwhile direction for future research includes a direct comparison between rodent

and human tasks, in which humans are trained and rewarded on a trial by trial basis,

exactly as is typically conducted with rodents.

We found that the magnitude of beta and gamma oscillations strongly predicted odor

identification accuracy, whereas this effect was weaker in the theta band, offering

support for distinct functional roles for the different rhythms in olfactory processing

[23,30,36,118]. Establishing beta rhythms in human piriform cortex may also lay the

groundwork for future work on olfactory networks and integration of olfactory information

with other cognitive processing streams and behaviors [30]. For example, integration of

olfactory information with language networks [128–130], memory networks [131–135],

sleep states [136–139], and other olfactory-guided behaviors [140,141] could potentially

involve interactions in beta oscillatory networks. Interestingly, we found that theta

oscillations, but not beta or gamma oscillations, increased during attended states when

no odor was present (**Fig 5B**), and prior to sniffing when odor was expected (**Fig 2A;**

Fig 3A and 3D). Taken together, these findings may indicate a role for theta in olfactory

attention, in line with previous studies suggesting that the phase and amplitude of lower

frequency oscillations reflect attentional states in the human brain [97]. Olfactory

attention also shapes neural responses in rodent olfactory cortex [95], and potential

future work may include a direct comparison of these oscillations between species. It

may also be interesting to explore these oscillations in other primary olfactory areas,

which may have similar neural responses to piriform cortex.

A limitation of our work is that we were unable to fully tease apart the contributions of

attention and expectation to our findings. Though we did conduct a control analysis

designed to look for effects of attention and anticipation by analyzing responses to sniffs

taken without anticipation of odor in between tasks, this analysis was limited by the fact

that there was no odor during the sniffs that we analyzed. A perfect control would

include presentation of odor in the absence of any expectation or anticipation. This

would be difficult to achieve, as multiple presentations of odor across an experimental

session almost undoubtedly would be anticipated after a couple of trials. This is an

interesting topic for future studies, once these challenging aspects of experimental

design are addressed.

Overall, our findings begin to define a characteristic human spectrotemporal odor

response, consisting of three distinct frequency bands with unique time courses relative

to sniff onset, with lower frequency phase modulating higher frequency amplitude, and

corresponding closely with cortical responses that have been found in rodents. This lays

the groundwork for future studies to further explore the functional role of the different

aspects of the human cortical olfactory response.

Methods

Ethics statement

This study was approved by the Institutional Review Board at Northwestern University,

under protocol #STU00201349, and the study adhered to the Declaration of Helsinki

and the Belmont Report. Participation was voluntary and written informed consent was

obtained from all participants.

Participants

Our study included iEEG data from nine patients with medication-resistant epilepsy. All

participants had depth electrodes implanted stereotactically for clinical pre-surgical

evaluation (**Fig 1A**). Electrode locations were determined solely based on clinical need

and, in our participant population, included piriform cortex within the medial temporal

lobe. Data were acquired at Northwestern Memorial Hospital.

Behavioral task

Participants performed a cued odor identification task in which they were asked whether

an odor matched a prior auditory cue. This task involved periodic presentation of odors

with inter-trial-intervals exceeding 6x the respiratory period of each individual, which

was at least 15 s for every participant. During this inter-trial period, participants breathed

naturally through the nose, and no odor was presented. The experimental trials were

conducted in each participant's hospital room, and were computer-controlled, presented

to participants using an Apple laptop computer running MATLAB (RRID: SCR_001622)

via the *PsychToolBox* extension (RRID: SCR_ 002881). Each trial of the task began

with an auditory cue consisting of either the word "rose" or the word "mint". The cue was

delivered by computer speaker. After a delay of 3 to 7 seconds, the odor of rose

(essential oil) or mint (methyl salicylate) was delivered through opaque squeeze bottles,

while nasal airflow was monitored in order to precisely determine sniff—and therefore

odor—onsets. We wanted participants to self-initiate odor delivery, so that they could

control the timing of the stimulus, and to ensure high quality data by allowing patients to

initiate trials at appropriate timing according to their individual situations. Note that this

design increased the jittered timing between the cue and the sniff, with patients initiating

the odor stimulus within 3 to 7 seconds following the cue. Participants were instructed

that following the auditory cue, when they were ready, they should exhale, bring the

bottle to their nose, and then sniff to sample the odor. For two patients, the

experimenter held the bottle and followed the same procedure. At the moment when the

patient sniffed, the experimenter sent a sync pulse to the clinical acquisition system via

button press. The experimenter's button press sent a signal via a data acquisition board

(USB-1208F, Measurement Computing), which translated TTL pulses from MATLAB

into the clinical EEG acquisition system (Nihon Kohden, Japan). Importantly, this button

press did not provide sniff onset information—rather, it served to mark the odor-

containing sniff in order to differentiate this sniff from other sniffs. Precise sniff onsets

were determined by analysis of the nasal airflow signals, described in the methods.

After smelling the odor, participants indicated whether the odor matched the cue via

button press, with the exception of two patients who preferred to speak their response,

which was recorded by an experimenter.

As previously described [98], participants completed between 48 and 64 trials, except

for 1 participant who completed only 16 trials due to clinical constraints. The average

inter-trial interval was 21.3 s, ranging from 14 to 28 s, across participants. The average

performance on the task was 73.3% correct (P1: 79.69%, P2: 75.56%, P3: 31.25%, P4:

91.07%, P5: 100%, P6: 87.5%, P7: 46.88%), which means the response was "yes"

when the odor matched with the sound cue or "no" when they were not matched. We

found no difference in performance between the first (mean ± standard error:

73.98% ± 9.12%) and second (72.58% ± 10.72%) half of trials (two-tailed Wilcoxon

signed-rank test; $z = 0.40$, $P = 0.69$). The performance was higher in cue-odor-matched

trials (85% ± 7.08%) than nonmatched trials (58.17% ± 15.05%) (two-tailed Wilcoxon

signed-rank test; $z = 2.20$, $P = 0.028$).

iEEG and respiratory signal recording

iEEG signals were recorded using the clinical EEG data acquisition system (Nihon

Kohden, Tokyo Japan) that is currently in use in Northwestern Memorial Hospital. The

sampling rate for each participant was determined clinically, and ranged from 500 to

2000 Hz across participants. The reference and ground consisted of an implanted

electrode strip on the surface of the brain facing the scalp. We recorded nasal airflow

using a piezoelectric pressure sensor (Salter Labs Model #5500) with a nasal cannula

placed at the patients' nostrils during the experiment. Nasal airflow signals were

recorded directly into the clinical acquisition system, and were therefore automatically

synchronized with the iEEG data. The signal was first mean centered, then key

respiratory features—including inhale and exhale onset, respiratory peak, and trough

volume and time—were detected automatically with MATLAB toolbox *BreathMetrics*,

developed by our lab [142], and then manually confirmed.

Electrode Localization

As previously described [97,98], to determine the implanted electrode locations, we

used pre-operative structural MRI scans and post-operative computed tomography (CT)

scans using the FMRIB Software Library's (FSL) registration tool *flirt* [143,144].

Individual CT images were registered to MRI images using 6 degrees of freedom and a

cost function of mutual information, which was followed by an affine registration with 12

degrees of freedom. Individual MRIs were registered to a standard Montreal

Neurological Institute (MNI) brain (MNI152_1mm_brain included in FSL) with 12

degrees of freedom. Finally, the transformation matrices generated above were

combined to create a transformation from the individual CT image to standard MNI

space.

The electrodes were localized by thresholding the raw CT image and calculating the un-

weighted mass center of each electrode. Finally, the coordinates were converted to

standard MNI space using the transformation matrix generated above.

Though we analyzed spectrograms from all electrodes on the PC depth wires and all

electrodes on the parietal grids, electrodes corresponding to those shown in Fig 2 were

selected by the following procedure. PC: For each subject, we first determined which

subset of electrodes was anatomically located inside PC using pre-drawn individual

ROIs based on the Human Brain Atlas [145]. This typically included between 1 and 3

electrodes. For the subjects who had only a single electrode in PC, we used that one.

For subjects with multiple electrodes within PC, we chose the one that was closest to

the center of PC. Notably, we also analyzed all piriform electrodes separately with

similar results.

Time-frequency analysis

For all time-frequency analyses, filtering was conducted using a two-pass, zero-phase-

lag, finite impulse response (FIR) filter, as implemented by the MATLAB toolbox *fieldtrip*

(RRID: SCR_004849), unless specified otherwise. We first low-pass filtered the iEEG

signal at 235 Hz and then removed 60 Hz noise and its harmonics with a band-stop filter

with a bandwidth of 4 Hz. We then downsampled the signal to 500Hz and re-referenced

the data to the common average. To compute spectrograms, we filtered the pre-

processed data between 1 to 200 Hz in 100 logarithmically increasing steps ranging

from 2 to 50 Hz in width. We kept only the first 95 frequency bands from 1 to 153 Hz as

our frequencies of interest for all subsequent analyses unless specified otherwise. The

analytical amplitudes of the filtered signals were calculated by taking the absolute value

of the Hilbert-transformed signal, and temporally smoothed with a moving average filter

kernel of 10 ms.

To compute spectrograms, we first created sniff-onset-aligned epochs, extending from –

2 s prior to 4 s following the sniff onset, for each trial. Sniff onsets were determined

using *BreathMetrics* [142]. Sniffs taken during experimental trials at the time of odor

presentation were used for the odor condition, sniffs taken in between trials when no

odor was present were used for the no-odor condition, and sniffs taken during

experimental trials in between the cue and odor were used for the attended-no-odor

condition. There were a larger number of no-odor trials compared to odor trials, and

therefore, no-odor trials were randomly sampled without replacement for each

participant separately to match trial numbers across conditions for further analysis. The

spectrograms were calculated by averaging the amplitude epochs across trials at each

frequency, which was further normalized by subtracting the baseline average. The time

window of [−0.55, −0.05] s prior to cue onset was used as baseline. To determine

statistical significance, we used a permutation method [97,98,146] to generate z score

values. For each repetition of the permutation process, the sniff onsets were circularly

shifted in time by a random amount while maintaining the relative distance between

them. Then, we calculate he average amplitude across these randomly shifted events.

The procedure above was repeated 10,000 times resulting a null distribution of baseline

amplitude. Finally, the baseline-corrected real spectrogram was divided by the standard

deviation of this null distribution, resulting a z score map. Raw power, baseline-

corrected power and z normalized maps can be seen in **S6 Fig 6**.

To compare the spectrograms between odor and no-odor condition, we used a

permutation method. In each permutation, the condition labels were shuffled across all

trials, and the difference of baseline-corrected spectrogram between permuted odor-

and no-odor-conditions were calculated. This process was repeated 10,000 times

resulting in a null distribution of spectrogram difference at each time-frequency point.

The mean and stand deviation of the null distribution was obtained using MATLAB's

normfit function. Finally, a z score of the real spectrogram difference was calculated by

subtracting the mean and then dividing by the stand deviation. The spectrograms

created for the attentional control analysis shown in Fig 4B were generated with the

same methods, with the number of trials adjusted to include equal numbers of odor and

attended-no-odor trials.

To compute the percentage change of the amplitude time series shown in **Fig 3A**, we

averaged the amplitude time series across frequencies within each of the following

frequency bands: theta 4–8 Hz, beta 13–30 Hz, and gamma 30–150 Hz, for each trial.

The percentage change of amplitude was calculated as (Amplitude-

Amplitude_baseline)/Amplitude_baseline for each trial. The baseline window was

defined as −500 ms to −50 ms relative to cue onset. To compare the percentage

change of amplitude between odor and no-odor conditions, we performed a paired

sample t test at each time point. Multiple comparisons across all time points were

corrected using FDR method.

In order to visualize the differences in temporal dynamics across frequency bands, we

created a z-normalized amplitude map for each frequency band using a bootstrapping

method (**Fig 3B**). First, each trial was baseline corrected and z-normalized. Then, in

each repetition, we resampled all 323 trials (with replacement), and calculated the

average across trials for each frequency band. A z-score matrix (repetition x time) was

generated for each frequency band after 2000 bootstraps (**Fig 3B**).

We then performed a cluster-based analysis to quantify the time course of significant

responses for each frequency band (**Fig 3C**). First, we performed a one sample t tests

against zero for z-normalized amplitudes for all trials at each time point for each

frequency band. To identify continuous significant clusters, We then used a cluster-

based statistical thresholding analysis to identify continuous significant clusters. The P

value threshold for initial clustering was set to $P = 0.01$, and the cluster size was defined

as the sum of the t-statistics of a given cluster. To generate a null distribution of the

cluster size, we used a permutation method. In each repetition (10000 repetition in

total), a random number of trials were circularly shifted separately, and the maximal

cluster size was calculated as described above. The 95th percentile of the distribution

was used as the cluster size significance threshold, which corresponds to cluster-based

corrected $P < 0.05$.

We next conducted an analysis to explore the timing and phase of responses in each

frequency band (**Fig 3D**). we calculated the distribution of respiratory phase values at

which the peak amplitude responses occurred using a bootstrapping method. For each

bootstrap, we resampled 323 trials (with replacement) while keeping the pairwise

relationship between LFP and breathing signals. The LFP and respiratory signals were

first averaged over resampled trials. To find the peaks of the LFP amplitude response,

we smoothed the amplitude time series using a moving average method with a kernel of

50 ms. Then, the respiratory phase of the maximal amplitude over the 4 s post-sniff time

period was extracted. The respiratory phase was obtained from the breathing signal

using Hilbert method. Finally, the distribution of the respiratory phases across trials

were qualified using PLV [147] as implemented in the MATLAB toolbox circstats (RRID:

SCR_016651) [148]. The Rayleigh test was performed to test for the non-uniformity of

phase distributions. For the peak timing analysis on the single participant level, we used

the same method.

We next conducted a series of analyses to look for relationships between LFPs and

odor identification accuracy (**Fig 4**). First, we computed amplitude spectrogram for odor

and no-odor conditions separately. The two spectrograms were z-normalized to the

same combined null distribution generated using methods described in the previous

section to rule out the effect of trial number difference across conditions. In order to

statistically compare the amplitude difference between correct and incorrect trials, we

conducted a resampling analysis. For each repetition, we resampled equal numbers of

correct and incorrect trials (71, with replacement) to ensure a fair comparison across

conditions. On each repetition, we calculated the mean of z-normalized amplitudes

across the entire time window for each frequency band for each condition (these

distributions are shown in **Fig 4B**), subtracted the mean of incorrect trials from the mean

of correct trials, and repeated this process 200 times to generate the distribution of

difference values between correct and incorrect trials. The difference values for each

frequency band was tested against 0 using one-sample t tests to generate statistics.

The distribution was plotted as scatter as well as box plot (**Fig 4B**).

We then conducted a bootstrap-correlation analysis to directly assess the correlation

between task performance and LFP amplitude of each frequency band for inhale and

exhale respectively on a trial-by-trial basis (**Fig 4C**). For each repetition (1000x) in the

analysis, we resampled 323 trials with replacement and calculated the accuracy of the

chosen trial set. We then calculated the mean amplitude during inhale and exhale

separately (inhale-exhale transition defined by phase angle of pi/2 for averaged

breathing signal) for all three frequency bands. We calculated Pearson's correlation

coefficients between task accuracy and mean amplitudes for inhale and exhale, for

each frequency band. We then repeated this entire process 50 times to generate a

distribution of correlation coefficient for each preparatory phase and frequency

condition. We used FDR to correct for multiple comparisons ($P < 0.05$) when we set the

significance threshold for the r value. We subsequently fisher-z-transformed all

Pearson's r values and conducted a repeated measures two-way ANOVA as well as

paired sample t tests to analyze the differences across frequency bands and respiratory

phases.

We next created scatter plot visualizations of amplitude responses during inhale and

exhale phases for correct and incorrect trials and fitted a linear SVM classifier to each of

the scatter plot in order to see how well the amplitude during inhale and exhale, taken

together, could predict task performance **(Fig 4D** and **4E).** To balance the discrepancy

between correct and incorrect trial numbers (252 vs. 71), we conducted a bootstrapping

analysis using equal numbers of trials across conditions on each repetition. For each

respiratory phase and frequency band, we sampled 30 trials within condition with

replacement each time, calculated percent change relative to baseline with the trial-wise

averaged amplitude time series, and calculated average amplitudes during inhale and

exhale for each trial set. Inhale and exhale periods were defined as 0–1.5s and 1.5–3 s

post sniff respectively for all trials. This process was repeated 500 times for both correct

and incorrect trials to generate a scatterplot of corresponding inhale and exhale

amplitudes for trials from both conditions. In order to test the separability of data, we

applied a linear binary SVM classifier using MATLAB function *fitcsvm* with five-fold

cross validation*. ROC curves were generated using the function *perfcurve.*

Phase-amplitude coupling analysis

We calculated the modulation index (MI) to measure coupling between theta phase and

higher frequency amplitudes (13–150 Hz) (**Fig 5**). To do so, we extracted the phase

angle time series for the theta frequency band (4–8 Hz), and then from that phase time

series, created sniff-aligned trial epochs. Data from all trials was concatenated. To

generate the higher frequency amplitude data that was used to compute the MI, the raw

time series were filtered and amplitude-extracted (as described previously) to the same

47 log-spaced frequency bands (13.05–153.04 Hz), with logarithmically increasing

bandwidth, as described in the previous section for the time-frequency analysis. We

then concatenated this data identically to the theta phase data. MI was calculated as the

Kullbach-Leibner distance divided by the logged number of bins. To normalize the MI,

we created a null distribution of MI values using permutation method. For each

repetition, and for each frequency band, we circularly shifted the amplitude time series,

and calculated the MI between the shifted amplitudes and theta phase values. This

process was repeated 1000 times for each frequency band to generate the

corresponding null distribution of MI values. We then normalized the raw MI to the null

distributions to generate the z-scores, and FDR corrected the P values for multiple

comparisons. To control for the effect of evoked potential on phase-amplitude coupling,

we used a permutation method to ensure that the significant MI we observed was

unlikely due to steep slope of evoked potential occurring in every trial. For each

repetition, we shuffled the order of theta phase but kept the order of higher frequency

amplitude to randomize the trial-by-trial relationship, and calculated MI for each

frequency band based on the shuffled trials. This process was repeated 1000 times for

each frequency band to create a null distribution for the MI value under the null

hypothesis that the phase-amplitude coupling was due to steep waveform occurring in

every trial. We then normalized the raw MI to the null distributions to generate the z-

scores, and FDR corrected the P values for multiple comparisons.

Control for nature of the task

Two participants performed three different olfactory tasks over six separate blocks.

Patients performed two blocks of each task. The three tasks required the participant to

indicate: 1) whether or not they detected any odor (Detection task), 2) whether the odor

was edible (Edibility task), or 3) the identity of the odor (Naming task). For all three

tasks, each trial began with a countdown ("3, 2, 1, Sniff") which appeared in sequential

order, with each number lasting 1 s. The participant was instructed to sniff when the

word sniff appeared, using a manual hand-held olfactometer that allowed for sniff-

controlled timing of odor delivery. The hand-held olfactometer consisted of an odor

canister connected to a nose port, constructed with one-way valves at the outlet and

inlet such that the headspace was able to be sampled only and immediately upon

sniffing. The canister was contained inside of an opaque 3D printed case. Each run

consisted of 20 trials, resulting in 40 trials of each task type. Odors were of natural

origin, and included banana, salsa, liquid smoke, blood orange tea, peanut butter,

garlic, peppermint, pine and rose, for each trial. The tasks were performed while clinical

ongoing electrophysiological activity was recorded at a sampling rate of 1000 Hz using

a 256-channel clinical EEG acquisition system (Nihon Kohden). The respiratory signals

were recorded using a pressure sensor at the nose (Salter Labs). Time-frequency

analysis of odor-induced responses in the piriform cortex was conducted using the

methods described above.

ACKNOWLEDGEMENTS

We thank the patients and families of patients who volunteered to participate in this

research; without their generous contributions of time and effort, this study would not

have been possible. We thank Navid Shadlou for his technical expertise and support,

and assistance with data collection. This work was supported by National Institutes of

Health grants R01-DC016364 (NIDCD) and R01-DC018539 (NIDCD) to C. Z.

REFERENCES:

1. Başar E, Başar-Eroglu C, Karakaş S, Schürmann M. Gamma, alpha, delta, and

theta oscillations govern cognitive processes. Int J Psychophysiol. 2001;39: 241–

248. doi:10.1016/S0167-8760(00)00145-8

2. Buzsáki G. Theta oscillations in the hippocampus. Neuron. 2002;33: 325–340.

doi:10.1016/S0896-6273(02)00586-X

3. Colgin LL, Denninger T, Fyhn M, Hafting T, Bonnevie T, Jensen O, et al.

Frequency of gamma oscillations routes flow of information in the hippocampus.

Nature. 2009;462: 353–357. doi:10.1038/nature08573

4. Engel AK, Fries P. Beta-band oscillations—signalling the status quo? Curr Opin

Neurobiol. 2010;20: 156–165. doi:10.1016/j.conb.2010.02.015

5. Fontanini A, Bower JM. Slow-waves in the olfactory system: an olfactory

perspective on cortical rhythms. Trends Neurosci. 2006;29: 429–437.

doi:10.1016/j.tins.2006.06.013

6. Fröhlich F, McCormick DA. Endogenous electric fields may guide neocortical

network activity. Neuron. 2010;67: 129–143. doi:10.1016/J.NEURON.2010.06.005

7.   Heck DH, McAfee SS, Liu Y, Babajani-Feremi A, Rezaie R, Freeman WJ, et al.

Breathing as a fundamental rhythm of brain function. Front Neural Circuits.

2017;10: 115. doi:10.3389/fncir.2016.00115

8.   Kay LM, Beshel J, Brea J, Martin C, Rojas-Líbano D, Kopell N. Olfactory

oscillations: the what, how and what for. Trends Neurosci. 2009;32: 207–214.

doi:10.1016/j.tins.2008.11.008

9.   Lisman JE, Jensen O. The theta-gamma neural code. Neuron. 2013;77: 1002–

1016. doi:10.1016/j.neuron.2013.03.007

10.   Murty DVPS, Shirhatti V, Ravishankar P, Ray S. Large visual stimuli induce two

distinct gamma oscillations in primate visual cortex. J Neurosci. 2018;38: 2730–

2744. doi:10.1523/JNEUROSCI.2270-17.2017

11.   Zoefel B, VanRullen R. Oscillatory mechanisms of stimulus processing and

selection in the visual and auditory systems: state-of-the-art, speculations and

suggestions. Front Neurosci. 2017;11: 296. doi:10.3389/fnins.2017.00296

12.   Deng Y, Choi I, Shinn-Cunningham B. Topographic specificity of alpha power

during auditory spatial attention. Neuroimage. 2020;207: 116360.

doi:10.1016/J.NEUROIMAGE.2019.116360

13.   van Ede F, de Lange F, Jensen O, Maris E. Orienting attention to an upcoming

tactile event involves a spatially and temporally specific modulation of

sensorimotor alpha- and beta-band oscillations. J Neurosci. 2011;31: 2016–2024.

doi:10.1523/JNEUROSCI.5630-10.2011

14.   Gross J, Schnitzler A, Timmermann L, Ploner M. Gamma oscillations in human

primary somatosensory cortex reflect pain perception. PLoS Biol. 2007;5: e133.

doi:10.1371/journal.pbio.0050133

15. Mayer A, Schwiedrzik CM, Wibral M, Singer W, Melloni L. Expecting to see a letter: alpha oscillations as carriers of top-down sensory predictions. Cereb Cortex. 2016;26: 3146–3160. doi:10.1093/CERCOR/BHV146

16. Todorovic A, van Ede F, Maris E, de Lange FP. Prior expectation mediates neural adaptation to repeated sounds in the auditory cortex: an MEG study. J Neurosci. 2011;31: 9118–2913. doi:10.1523/JNEUROSCI.1425-11.2011

17. Adrian ED. Olfactory reactions in the brain of the hedgehog. J Physiol. 1942;100: 459–473. doi:10.1113/jphysiol.1942.sp003955

18. Barrie JM, Freeman WJ, Lenhart MD. Spatiotemporal analysis of prepyriform, visual, auditory, and somesthetic surface EEGs in trained rabbits. J Neurophysiol. 1996;76: 520–539. doi:10.1152/jn.1996.76.1.520

19. Freeman WJ. A linear distributed feedback model for prepyriform cortex. Exp Neurol. 1964;10: 525–547. doi:10.1016/0014-4886(64)90049-4

20. Freeman WJ. Phasic and long-term excitability changes in prepyriform cortex of cats. Exp Neurol. 1962;5: 500–518. doi:10.1016/0014-4886(62)90059-6

21. Freeman WJ. Distribution in time and space of prepyriform electrical activity. J Neurophysiol. 1959;22: 644–665. doi:10.1152/JN.1959.22.6.644

22. Freeman WJ. Measurement of oscillatory responses to electrical stimulation in olfactory bulb of cat. J Neurophysiol. 1972;35: 762–779. doi:10.1152/JN.1972.35.6.762

23. Beshel J, Kopell N, Kay LM. Olfactory bulb gamma oscillations are enhanced with task demands. J Neurosci. 2007;27: 8358–8365. doi:10.1523/JNEUROSCI.1199-

984   07.2007

24.   Chapuis J, Cohen Y, He X, Zhang Z, Jin S, Xu F, et al. Lateral entorhinal

        modulation of piriform cortical activity and fine odor discrimination. J Neurosci.

        2013;33: 13449–13459. doi:10.1523/JNEUROSCI.1387-13.2013

25.   Cohen Y, Putrino D, Wilson DA. Dynamic cortical lateralization during olfactory

        discrimination learning. J Physiol. 2015;593: 1701–1714.

        doi:10.1113/jphysiol.2014.288381

26.   Courtiol E, Buonviso N, Litaudon P. Odorant features differentially modulate

        beta/gamma oscillatory patterns in anterior versus posterior piriform cortex.

        Neuroscience. 2019;409: 26–34. doi:10.1016/j.neuroscience.2019.04.025

27.   Gnaedinger A, Gurden H, Gourévitch B, Martin C. Multisensory learning between

        odor and sound enhances beta oscillations. Sci Rep. 2019;9: 11236.

        doi:10.1038/s41598-019-47503-y

28.   Kay LM. Theta oscillations and sensorimotor performance. Proc Natl Acad Sci U

        S A. 2005;102: 3863–3868. doi:10.1073/pnas.0407920102

29.   Kay LM. Circuit oscillations in odor perception and memory. Prog Brain Res.

        2014;208: 223–251. doi:10.1016/B978-0-444-63350-7.00009-7

30.   Kay LM, Beshel J. A beta oscillation network in the rat olfactory system during a

        2-alternative choice odor discrimination task. J Neurophysiol. 2010;104: 829–839.

        doi:10.1152/jn.00166.2010

31.   Kay LM, Stopfer M. Information processing in the olfactory systems of insects and

        vertebrates. Semin Cell Dev Biol. 2006;17: 433–442.

        doi:10.1016/j.semcdb.2006.04.012

1007   32.   Lepousez G, Lledo P-M. Odor discrimination requires proper olfactory fast

1008   oscillations in awake mice. Neuron. 2013;80: 1010–1024.

1009   doi:10.1016/j.neuron.2013.07.025

1010   33.   Losacco J, Ramirez-Gordillo D, Gilmer J, Restrepo D. Learning improves

1011   decoding of odor identity with phase-referenced oscillations in the olfactory bulb.

1012   Elife. 2020;9: e52583. doi:10.7554/eLife.52583

1013   34.   Lowry CA, Kay LM. Chemical factors determine olfactory system beta oscillations

1014   in waking rats. J Neurophysiol. 2007;98: 394–404. doi:10.1152/jn.00124.2007

1015   35.   Martin C, Beshel J, Kay LM. An olfacto-hippocampal network is dynamically

1016   involved in odor-discrimination learning. J Neurophysiol. 2007;98: 2196–2205.

1017   doi:10.1152/jn.00524.2007

1018   36.   Martin C, Ravel N. Beta and gamma oscillatory activities associated with olfactory

1019   memory tasks: different rhythms for different functional networks? Front Behav

1020   Neurosci. 2014;23: 218. doi:10.3389/fnbeh.2014.00218

1021   37.   Mori K, Manabe H, Narikiyo K, Onisawa N. Olfactory consciousness and gamma

1022   oscillation couplings across the olfactory bulb, olfactory cortex, and orbitofrontal

1023   cortex. Front Psychol. 2013;4: 743. doi:10.3389/fpsyg.2013.00743

1024   38.   Neville KR, Haberly LB. Beta and gamma oscillations in the olfactory system of

1025   the urethane-anesthetized rat. J Neurophysiol. 2003;90: 3921–3930.

1026   doi:10.1152/jn.00475.2003

1027   39.   Poo C, Isaacson JS. Odor representations in olfactory cortex: "sparse" coding,

1028   global inhibition, and oscillations. Neuron. 2009;62: 850–861.

1029   doi:10.1016/j.neuron.2009.05.022

1030   40.   Ravel N, Chabaud P, Martin C, Gaveau V, Hugues E, Tallon-Baudry C, et al.

1031   Olfactory learning modifies the expression of odour-induced oscillatory responses

1032   in the gamma (60-90 Hz) and beta (15-40 Hz) bands in the rat olfactory bulb. Eur

1033   J Neurosci. 2003;17: 350–358. doi:10.1046/j.1460-9568.2003.02445.x

1034   41.   Rosero MA, Aylwin ML. Sniffing shapes the dynamics of olfactory bulb gamma

1035   oscillations in awake behaving rats. Eur J Neurosci. 2011;34: 787–799.

1036   doi:10.1111/J.1460-9568.2011.07800.X

1037   42.   Wilson DA, Sullivan RM. Cortical processing of odor objects. Neuron. 2011;72:

1038   506–519. doi:10.1016/j.neuron.2011.10.027

1039   43.   Zhou Y, Fang F-H, Pan P, Liu Z-R, Ji Y-H. Visual deprivation induce cross-modal

1040   enhancement of olfactory perception. Biochem Biophys Res Commun. 2017;486:

1041   833–838. doi:10.1016/j.bbrc.2017.03.140

1042   44.   Buzsáki G, Wang X-J. Mechanisms of gamma oscillations. Annu Rev Neurosci.

1043   2012;35: 203–225. doi:10.1146/ANNUREV-NEURO-062111-150444

1044   45.   Spencer KM, Niznikiewicz MA, Shenton ME, McCarley RW. Sensory-evoked

1045   gamma oscillations in chronic schizophrenia. Biol Psychiatry. 2008;63: 744–747.

1046   doi:10.1016/J.BIOPSYCH.2007.10.017

1047   46.   Bartoli E, Bosking W, Foster BL. Seeing visual gamma oscillations in a new light.

1048   Trends Cogn Sci. 2020;24: 501–503. doi:10.1016/j.tics.2020.03.009

1049   47.   Edden RAE, Muthukumaraswamy SD, Freeman TCA, Singh KD. Orientation

1050   discrimination performance is predicted by GABA concentration and gamma

1051   oscillation frequency in human primary visual cortex. J Neurosci. 2009;29: 15721–

1052   15726. doi:10.1523/JNEUROSCI.4426-09.2009

1053   48.   Hermes D, Miller KJ, Wandell BA, Winawer J. Stimulus dependence of gamma

1054   oscillations in human visual cortex. Cereb Cortex. 2015;25: 2951–2959.

1055   doi:10.1093/cercor/bhu091

1056   49.   Honkanen R, Rouhinen S, Wang SH, Palva JM, Palva S. Gamma oscillations

1057   underlie the maintenance of feature-specific information and the contents of visual

1058   working memory. Cereb Cortex. 2015;25: 3788–3801.

1059   doi:10.1093/CERCOR/BHU263

1060   50.   Muthukumaraswamy SD, Singh KD. Visual gamma oscillations: the effects of

1061   stimulus type, visual field coverage and stimulus motion on MEG and EEG

1062   recordings. Neuroimage. 2013;69: 223–230.

1063   doi:10.1016/J.NEUROIMAGE.2012.12.038

1064   51.   Bosman CA, Schoffelen JM, Brunet N, Oostenveld R, Bastos AM, Womelsdorf T,

1065   et al. Attentional stimulus selection through selective synchronization between

1066   monkey visual areas. Neuron. 2012;75: 875–888.

1067   doi:10.1016/J.NEURON.2012.06.037

1068   52.   Rodriguez E, George N, Lachaux J-PP, Martinerie J, Renault B, Varela FJ.

1069   Perception's shadow: long-distance synchronization of human brain activity.

1070   Nature. 1999;397: 430–433. doi:10.1038/17120

1071   53.   Samonds JM, Bonds AB. Gamma oscillation maintains stimulus structure-

1072   dependent synchronization in cat visual cortex. J Neurophysiol. 2005;93: 223–

1073   236. doi:10.1152/JN.00548.2004

1074   54.   Brosch M, Budinger E, Scheich H. Stimulus-related gamma oscillations in primate

1075   auditory cortex. J Neurophysiol. 2002;87: 2715–2725.

doi:10.1152/JN.2002.87.6.2715

55.   Albouy P, Mattout J, Bouet R, Maby E, Sanchez G, Aguera P-E, et al. Impaired

pitch perception and memory in congenital amusia: the deficit starts in the

auditory cortex. Brain. 2013;136: 1639–1661. doi:10.1093/brain/awt082

56.   Ross B, Tremblay KL, Alain C. Simultaneous EEG and MEG recordings reveal

vocal pitch elicited cortical gamma oscillations in young and older adults.

Neuroimage. 2020;204: 116253. doi:10.1016/j.neuroimage.2019.116253

57.   Jiang H, Schuele S, Rosenow J, Zelano C, Parvizi J, Tao JX, et al. Theta

oscillations rapidly convey odor-specific content in human piriform cortex. Neuron.

2017;94: 207-219.e4. doi:10.1016/j.neuron.2017.03.021

58.   Cenier T, Amat C, Litaudon P, Garcia S, Lafaye de Micheaux P, Liquet B, et al.

Odor vapor pressure and quality modulate local field potential oscillatory patterns

in the olfactory bulb of the anesthetized rat. Eur J Neurosci. 2008;27: 1432–1440.

doi:10.1111/j.1460-9568.2008.06123.x

59.   David F, Courtiol E, Buonviso N, Fourcaud-Trocmé N. Competing mechanisms of

gamma and beta oscillations in the olfactory bulb based on multimodal inhibition

of mitral cells over a respiratory cycle. eNeuro. 2015;2: ENEURO.0018-15.2015.

doi:10.1523/ENEURO.0018-15.2015

60.   Fukunaga I, Herb JT, Kollo M, Boyden ES, Schaefer AT. Independent control of

gamma and theta activity by distinct interneuron networks in the olfactory bulb.

Nat Neurosci. 2014;17: 1208–1216. doi:10.1038/nn.3760

61.   Kopell N, Ermentrout GB, Whittington MA, Traub RD. Gamma rhythms and beta

rhythms have different synchronization properties. Proc Natl Acad Sci U S A.

2000;97: 1867–1872. doi:10.1073/pnas.97.4.1867

62. Litaudon P, Garcia S, Buonviso N. Strong coupling between pyramidal cell activity

and network oscillations in the olfactory cortex. Neuroscience. 2008;156: 781–

787. doi:10.1016/J.NEUROSCIENCE.2008.07.077

63. Manabe H, Mori K. Sniff rhythm-paced fast and slow gamma-oscillations in the

olfactory bulb: relation to tufted and mitral cells and behavioral states. J

Neurophysiol. 2013;110: 1593–1599. doi:10.1152/jn.00379.2013

64. Frederick DE, Brown A, Brim E, Mehta N, Vujovic M, Kay LM. Gamma and beta

oscillations define a sequence of neurocognitive modes present in odor

processing. J Neurosci. 2016;36: 7750–7767. doi:10.1523/JNEUROSCI.0569-

16.2016

65. Freeman WJ, Schneider W. Changes in spatial patterns of rabbit olfactory EEG

with conditioning to odors. Psychophysiology. 1982;19: 44–56.

doi:10.1111/J.1469-8986.1982.TB02598.X

66. MacLeod K, Bäcker A, Laurent G. Who reads temporal information contained

across synchronized and oscillatory spike trains? Nature. 1998;395: 693–698.

doi:10.1038/27201

67. Nusser Z, Kay LM, Laurent G, Homanics GE, Mody I. Disruption of GABAA

receptors on GABAergic interneurons leads to increased oscillatory power in the

olfactory bulb network. J Neurophysiol. 2001;86: 2823–2833.

doi:10.1152/JN.2001.86.6.2823

68. Stopfer M, Bhagavan S, Smith BH, Laurent G. Impaired odour discrimination on

desynchronization of odour-encoding neural assemblies. Nature. 1997;390: 70–

74. doi:10.1038/36335

69.   Viana Di Prisco G, Freeman WJ. Odor-related bulbar EEG spatial pattern analysis

during appetitive conditioning in rabbits. Behav Neurosci. 1985;99: 964–978.

doi:10.1037/0735-7044.99.5.964

70.   Rojas-Líbano D, Frederick DE, Egaña J, Kay LM. The olfactory bulb theta rhythm

follows all frequencies of diaphragmatic respiration in the freely behaving rat.

Front Behav Neurosci. 2014;8: 214. doi:10.3389/fnbeh.2014.00214

71.   Xu W, Wilson DA. Odor-evoked activity in the mouse lateral entorhinal cortex.

Neuroscience. 2012;223: 12–20. doi:10.1016/j.neuroscience.2012.07.067

72.   Hudry J, Ryvlin P, Royet J-P, Mauguière F. Odorants elicit evoked potentials in

the human amygdala. Cereb Cortex. 2001;11: 619–627.

doi:10.1093/CERCOR/11.7.619

73.   Jung J, Hudry J, Ryvlin P, Royet J-P, Bertrand O, Lachaux J-P. Functional

significance of olfactory-induced oscillations in the human amygdala. Cereb

Cortex. 2006;16: 1–8. doi:10.1093/cercor/bhi090

74.   Hughes JR, Andy OJ. The human amygdala. I. Electrophysiological responses to

odorants. Electroencephalogr Clin Neurophysiol. 1979;46: 428–443.

doi:10.1016/0013-4694(79)90144-5

75.   Hughes JR, Hendrix DE, Wetzel N, Johnston JW. Correlations between

electrophysiological activity from the human olfactory bulb and the subjective

response to odoriferous stimuli. Electroencephalogr Clin Neurophysiol. 1970;28:

97–8. Available: http://www.ncbi.nlm.nih.gov/pubmed/4188883

76.   Hudry J. Olfactory short-term memory and related amygdala recordings in

patients with temporal lobe epilepsy. Brain. 2003;126: 1851–1863.

doi:10.1093/brain/awg192

77.   Buonviso N, Amat C, Litaudon P, Roux S, Royet J-P, Farget V, et al. Rhythm

sequence through the olfactory bulb layers during the time window of a respiratory

cycle. Eur J Neurosci. 2003;17: 1811–1819. doi:10.1046/j.1460-

9568.2003.02619.x

78.   Herrero JL, Khuvis S, Yeagle E, Cerf M, Mehta AD. Breathing above the brain

stem: volitional control and attentional modulation in humans. J Neurophysiol.

2018;119: 145–159. doi:10.1152/jn.00551.2017

79.   Lockmann AL V., Laplagne DA, Tort ABL. Olfactory bulb drives respiration-

coupled beta oscillations in the rat hippocampus. Eur J Neurosci. 2018;48: 2663–

2673. doi:10.1111/ejn.13665

80.   Pashkovski SL, Iurilli G, Brann D, Chicharro D, Drummey K, Franks K, et al.

Structure and flexibility in cortical representations of odour space. Nature.

2020;583: 253–258. doi:10.1038/s41586-020-2451-1

81.   Bolding KA, Franks KM. Complementary codes for odor identity and intensity in

olfactory cortex. Elife. 2017;6: pii: e22630. doi:10.7554/eLife.22630

82.   Roland B, Deneux T, Franks KM, Bathellier B, Fleischmann A. Odor identity

coding by distributed ensembles of neurons in the mouse olfactory cortex. Elife.

2017;6: e26337. doi:10.7554/eLife.26337

83.   Sobel N, Prabhakaran V, Desmond JE, Glover GH, Goode RL, Sullivan E V, et al.

Sniffing and smelling: separate subsystems in the human olfactory cortex. Nature.

1998;392: 282–286. doi:10.1038/32654

84.   Mainland J, Sobel N. The sniff is part of the olfactory percept. Chem Senses.

2006;31: 181–196. doi:10.1093/chemse/bjj012

85.   Zelano C, Jiang H, Zhou G, Arora N, Schuele S, Rosenow J, et al. Nasal

respiration entrains human limbic oscillations and modulates cognitive function. J

Neurosci. 2016;36: 12448–12467. doi:10.1523/JNEUROSCI.2586-16.2016

86.   Kalcher J, Pfurtscheller G. Discrimination between phase-locked and non-phase-

locked event-related EEG activity. Electroencephalogr Clin Neurophysiol.

1995;94: 381–384. doi:10.1016/0013-4694(95)00040-6

87.   Donner TH, Siegel M. A framework for local cortical oscillation patterns. Trends

Cogn Sci. 2011;15: 191–199. doi:10.1016/j.tics.2011.03.007

88.   Cohen MX. Differences among total, phase-locked, and non-phase-locked power

and intertrial phase consistency. Analyzing neural time series data.

Massachusetts Institute of Technology; 2014. pp. 259–262.

89.   Pfurtscheller G, Lopes Da Silva FH. Event-related EEG/MEG synchronization and

desynchronization: basic principles. Clin Neurophysiol. 1999;110: 1842–1857.

doi:10.1016/S1388-2457(99)00141-8

90.   Tallon-Baudry C, Bertrand O. Oscillatory gamma activity in humans and its role in

object representation. Trends Cogn Sci. 1999;3: 151–162. doi:10.1016/S1364-

6613(99)01299-1

91.   Barnes DC, Chapuis J, Chaudhury D, Wilson DA. Odor fear conditioning modifies

piriform cortex local rield potentials both during conditioning and during post-

conditioning sleep. PLoS One. 2011;6: e18130.

doi:10.1371/journal.pone.0018130

92.   van der Meer M. Integrating early results on ventral striatal gamma oscillations in

1192   the rat. Front Neurosci. 2010;0: 300. doi:10.3389/fnins.2010.00300

1193   93.   Bressler SL, Freeman WJ. Frequency analysis of olfactory system EEG in cat,

1194   rabbit, and rat. Electroencephalogr Clin Neurophysiol. 1980;50: 19–24.

1195   doi:10.1016/0013-4694(80)90319-3

1196   94.   Rojas-Líbano D, Kay LM. Olfactory system gamma oscillations: the physiological

1197   dissection of a cognitive neural system. Cogn Neurodyn. 2008;2: 179–194.

1198   doi:10.1007/s11571-008-9053-1

1199   95.   Carlson KS, Gadziola MA, Dauster ES, Wesson DW. Selective attention controls

1200   olfactory decisions and the neural encoding of odors. Curr Biol. 2018;28: 2195-

1201   2205.e4. doi:10.1016/j.cub.2018.05.011

1202   96.   Zelano C, Bensafi M, Porter J, Mainland J, Johnson B, Bremner E, et al.

1203   Attentional modulation in human primary olfactory cortex. Nat Neurosci. 2005;8:

1204   114–120. doi:10.1038/nn1368

1205   97.   Arabkheradmand G, Zhou G, Noto T, Yang Q, Schuele SU, Parvizi J, et al.

1206   Anticipation-induced delta phase reset improves human olfactory perception.

1207   Hanslmayr S, editor. PLOS Biol. 2020;18: e3000724.

1208   doi:10.1371/journal.pbio.3000724

1209   98.   Zhou G, Lane G, Noto T, Arabkheradmand G, Gottfried JA, Schuele SU, et al.

1210   Human olfactory-auditory integration requires phase synchrony between sensory

1211   cortices. Nat Commun. 2019;10: 1168. doi:10.1038/s41467-019-09091-3

1212   99.   Tort ABL, Komorowski R, Eichenbaum H, Kopell N. Measuring phase-amplitude

1213   coupling between neuronal oscillations of different frequencies. J Neurophysiol.

1214   2010;104: 1195–210. doi:10.1152/jn.00106.2010

100.   Voytek B, D'Esposito M, Crone N, Knight RT. A method for event-related

        phase/amplitude coupling. Neuroimage. 2013;64: 416–424.

        doi:10.1016/j.neuroimage.2012.09.023

101.   Lane G, Zhou G, Noto T, Zelano C. Assessment of direct knowledge of the

        human olfactory system. Exp Neurol. 2020;329: 113304.

        doi:10.1016/j.expneurol.2020.113304

102.   Olofsson JK, Bowman NE, Khatibi K, Gottfried JA. A time-based account of the

        perception of odor objects and valences. Psychol Sci. 2012;23: 1224–1232.

        doi:10.1177/0956797612441951

103.   Singh AK, Touhara K, Okamoto M. Electrophysiological correlates of top-down

        attentional modulation in olfaction. Sci Rep. 2019;9: 4953. doi:10.1038/s41598-

        019-41319-6

104.   Iravani B, Arshamian A, Lundqvist M, Kay LM, Wilson DA, Lundström JN. Odor

        identity can be extracted from the reciprocal connectivity between olfactory bulb

        and piriform cortex in humans. Neuroimage. 2021;237: 118130.

        doi:10.1016/J.NEUROIMAGE.2021.118130

105.   Xia CZ, Adjei S, Wesson DW. Coding of odor stimulus features among secondary

        olfactory structures. J Neurophysiol. 2015;114: 736–745.

        doi:10.1152/JN.00902.2014/ASSET/IMAGES/LARGE/Z9K0121531780004.JPEG

106.   Ketchum K, Haberly L. Synaptic events that generate fast oscillations in piriform

        cortex. J Neurosci. 1993;13: 3980–3985. doi:10.1523/JNEUROSCI.13-09-

        03980.1993

107.   Kay LM, Freeman WJ. Bidirectional processing in the olfactory-limbic axis during

olfactory behavior. Behav Neurosci. 1998;112: 541–553. doi:10.1037/0735-

7044.112.3.541

108.   Bathellier B, Buhl DL, Accolla R, Carleton A. Dynamic ensemble odor coding in

the mammalian olfactory bulb: sensory information at different timescales.

Neuron. 2008;57: 586–598. doi:10.1016/j.neuron.2008.02.011

109.   Blauvelt DG, Sato TF, Wienisch M, Murthy VN. Distinct spatiotemporal activity in

principal neurons of the mouse olfactory bulb in anesthetized and awake states.

Front Neural Circuits. 2013;7: 46. doi:10.3389/fncir.2013.00046

110.   Kepecs A, Uchida N, Mainen ZF. The sniff as a unit of olfactory processing. Chem

Senses. 2006;31: 167–179. doi:10.1093/chemse/bjj016

111.   Phillips ME, Sachdev RNS, Willhite DC, Shepherd GM. Respiration drives

network activity and modulates synaptic and circuit processing of lateral inhibition

in the olfactory bulb. J Neurosci. 2012;32: 85–98. doi:10.1523/JNEUROSCI.4278-

11.2012

112.   Grosmaitre X, Santarelli L, Tan J, Luo M, Ma M. Dual functions of mammalian

olfactory sensory neurons as odor detectors and mechanical sensors. Nat

Neurosci. 2007;10: 348–354. doi:10.1038/NN1856

113.   Rennaker RL, Chen C-FF, Ruyle AM, Sloan AM, Wilson DA. Spatial and temporal

distribution of odorant-evoked activity in the piriform cortex. J Neurosci. 2007;27:

1534–1542. doi:10.1523/jneurosci.4072-06.2007

114.   Shusterman R, Smear MC, Koulakov AA, Rinberg D. Precise olfactory responses

tile the sniff cycle. Nat Neurosci. 2011;14: 1039–1044. doi:10.1038/nn.2877

115. Uchida N, Kepecs A, Mainen ZF. Seeing at a glance, smelling in a whiff: rapid

forms of perceptual decision making. Nat Rev Neurosci. 2006;7: 485–491.

doi:10.1038/nrn1933

116. Verhagen J V, Wesson DW, Netoff TI, White JA, Wachowiak M. Sniffing controls

an adaptive filter of sensory input to the olfactory bulb. Nat Neurosci. 2007;10:

631–639. doi:10.1038/nn1892

117. Haddad R, Lanjuin A, Madisen L, Zeng H, Murthy VNN, Uchida N. Olfactory

cortical neurons read out a relative time code in the olfactory bulb. Nat Neurosci.

2013;16: 949–957. doi:10.1038/nn.3407

118. Frederick DE, Brown A, Tacopina S, Mehta N, Vujovic M, Brim E, et al. Task-

dependent behavioral dynamics make the case for temporal integration in multiple

strategies during odor processing. J Neurosci. 2017;37: 4416–4426.

doi:10.1523/JNEUROSCI.1797-16.2017

119. Kepecs A, Uchida N, Mainen ZF. Rapid and precise control of sniffing during

olfactory discrimination in rats. J Neurophysiol. 2007;98: 205–213.

doi:10.1152/jn.00071.2007

120. Sirotin YB, Costa ME, Laplagne DA. Rodent ultrasonic vocalizations are bound to

active sniffing behavior. Front Behav Neurosci. 2014;8: 399.

doi:10.3389/FNBEH.2014.00399

121. Iravani B, Arshamian A, Ohla K, Wilson DA, Lundström JN. Non-invasive

recording from the human olfactory bulb. Nat Commun. 2020;11: 648.

doi:10.1038/s41467-020-14520-9

122. Kay LM, Beshel J, Martin C. When good enough is best. Neuron. 2006;51: 277–

1283 278. doi:10.1016/J.NEURON.2006.07.015

1284 123. Friedrich RW. Mechanisms of odor discrimination: neurophysiological and

1285 behavioral approaches. Trends Neurosci. 2006;29: 40–47.

1286 doi:10.1016/J.TINS.2005.10.004

1287 124. Frederick DE, Rojas-Líbano D, Scott M, Kay LM. Rat behavior in Go/No-Go and

1288 two-alternative choice odor discrimination: differences and similarities. Behav

1289 Neurosci. 2011;125: 588–603. doi:10.1037/A0024371

1290 125. Kay LM, Laurent G. Odor- and context-dependent modulation of mitral cell activity

1291 in behaving rats. Nat Neurosci. 1999;2: 1003–1009. doi:10.1038/14801

1292 126. Doucette W, Restrepo D. Profound context-dependent plasticity of mitral cell

1293 responses in olfactory bulb. PLOS Biol. 2008;6: e258.

1294 doi:10.1371/JOURNAL.PBIO.0060258

1295 127. Doucette W, Gire DH, Whitesell J, Carmean V, Lucero MT, Restrepo D.

1296 Associative cortex features in the first olfactory brain relay station. Neuron.

1297 2011;69: 1176–1187. doi:10.1016/J.NEURON.2011.02.024

1298 128. Majid A. Human olfaction at the intersection of language, culture, and biology.

1299 Trends Cogn Sci. 2021;25: 111–123. doi:10.1016/j.tics.2020.11.005

1300 129. Majid A, Kruspe N. Hunter-gatherer olfaction is special. Curr Biol. 2018;28: 409-

1301 413.e2. doi:10.1016/J.CUB.2017.12.014

1302 130. Olofsson JK, Gottfried JA. The muted sense: neurocognitive limitations of

1303 olfactory language. Trends Cogn Sci. 2015;19: 314–321.

1304 doi:10.1016/j.tics.2015.04.007

1305 131. Arshamian A, Iravani B, Majid A, Lundström JN. Respiration modulates olfactory

memory consolidation in humans. J Neurosci. 2018;38: 10286–10294.

doi:10.1523/JNEUROSCI.3360-17.2018

132. Arzi A, Rozenkrantz L, Holtzman Y, Secundo L, Sobel N. Sniffing patterns

uncover implicit memory for undetected odors. Curr Biol. 2014;24: R263–R264.

doi:10.1016/j.cub.2014.02.004

133. Wilson DA, Stevenson RJ. The fundamental role of memory in olfactory

perception. Trends Neurosci. 2003;26: 243–247. doi:10.1016/S0166-

2236(03)00076-6

134. Yeshurun Y, Lapid H, Dudai Y, Sobel N. The privileged brain representation of

first olfactory associations. Curr Biol. 2009;19: 1869–1874.

doi:10.1016/j.cub.2009.09.066

135. Zhou G, Olofsson JK, Koubeissi MZ, Menelaou G, Rosenow J, Schuele SU, et al.

Human hippocampal connectivity is stronger in olfaction than other sensory

systems. Prog Neurobiol. 2021;201: 102027.

doi:10.1016/j.pneurobio.2021.102027

136. Arzi A, Sela L, Green A, Givaty G, Dagan Y, Sobel N. The influence of odorants

on respiratory patterns in sleep. Chem Senses. 2010;35: 31–40.

doi:10.1093/chemse/bjp079

137. Hauner KK, Howard JD, Zelano C, Gottfried JA. Stimulus-specific enhancement

of fear extinction during slow-wave sleep. Nat Neurosci. 2013;16: 1553–1555.

doi:10.1038/nn.3527

138. Perl O, Arzi A, Sela L, Secundo L, Holtzman Y, Samnon P, et al. Odors enhance

slow-wave activity in non-rapid eye movement sleep. J Neurophysiol. 2016;115:

2294–2302. doi:10.1152/JN.01001.2015

139.   Rasch B, Buchel C, Gais S, Born J. Odor cues during slow-wave sleep prompt

declarative memory consolidation. Science (80-). 2007;315: 1426–1429.

doi:10.1126/science.1138581

140.   Bhutani S, Howard JD, Reynolds R, Zee PC, Gottfried J. Olfactory connectivity

mediates sleep-dependent food choices in humans. Elife. 2019;8: e49053.

doi:10.7554/ELIFE.49053

141.   Howard JD, Reynolds R, Smith DE, Voss JL, Schoenbaum G, Kahnt T. Targeted

stimulation of human orbitofrontal networks disrupts outcome-guided behavior.

Curr Biol. 2020;30: 490-498.e4. doi:10.1016/J.CUB.2019.12.007

142.   Noto T, Zhou G, Schuele S, Templer J, Zelano C. Automated analysis of

breathing waveforms using BreathMetrics: a respiratory signal processing toolbox.

Chem Senses. 2018;43: 583–597. doi:10.1093/chemse/bjy045

143.   Jenkinson M, Smith S. A global optimisation method for robust affine registration

of brain images. Med Image Anal. 2001;5: 143–156. doi:10.1016/S1361-

8415(01)00036-6

144.   Jenkinson M, Bannister P, Brady M, Smith S. Improved optimization for the robust

and accurate linear registration and motion correction of brain images.

Neuroimage. 2002;17: 825–841. doi:10.1006/nimg.2002.1132

145.   Mai JK, Majtanik M, Paxinos G. Atlas of the human brain. 4th ed. New York:

Academic Press; 2015.

146.   Canolty RT. Spatiotemporal dynamics of word processing in the human brain.

Front Neurosci. 2007;1: 185–196. doi:10.3389/neuro.01.1.1.014.2007

147.   Aydore S, Pantazis D, Leahy RM. A note on the phase locking value and its

properties. Neuroimage. 2013;74: 231–244.

doi:10.1016/j.neuroimage.2013.02.008

148.   Berens P. CircStat : a MATLAB toolbox for circular statistics. J Stat Softw.

2009;31: 1–21. doi:10.18637/jss.v031.i10

1357

1358

1359

1360

1361

S1 Fig. Related to Fig 2. Odor-induced and odor-evoked spectrograms. Non-phase-

locked (left) and phase-locked (right) spectrograms are shown for each participant (P1–

P7). The non-phase-locked spectrogram was obtained by subtracting the event-related

potential from each trial before time-frequency decomposition. The red solid overlay

indicates each participant's respiratory signal. Black outlines indicate statistically

significant clusters ($P < 0.05$, false-discovery-rate corrected). The phase-locked

spectrogram was calculated as the baseline-corrected time-frequency decomposition of

the event-related potential. The vertical short-dashed lines indicate sniff onset. The

source is available in S5 Data.

1371

1372

1373 S2 Fig. Related to Fig 2. Sniff-onset aligned spectrograms with linear frequency scale.

1374 The black solid overlay indicates each participant's (P1–P7) respiratory signal. Black

1375 outlines indicate statistically significant clusters (P < 0.05, false-discovery-rate

1376 corrected). The vertical short-dashed line indicates sniff onset. The source is available

1377 in S5 Data.

1378

S3 Fig. Related to Fig 2. Odor-induced responses are maximal in piriform cortex. (A)

Odor-induced amplitude (sum of z score) is larger in the depth wire located in the

piriform cortex (PC) compared to those located outside of the piriform cortex (Lateral to

PC) in theta, beta and gamma frequency bands. * indicates statistically significant

difference (two-tailed paired t test). (B) The mean z score was calculated over a time

window of 2s in the theta, beta or gamma frequency band for each electrode and each

participant. The data were collapsed over the z-axis and smoothed. The background

brain is a slice (z = –16) of the Montreal Neurological Institute standard brain. The pink

rectangles outline piriform cortex. The source is available in S5 Data.

1388

S4 Fig. Related to Fig 4. Odor-induced responses for correct and incorrect trials. (A)

Example scatter plots showing the correlations plotted in Fig 4C. The scatter plot

corresponding to the correlation value of one dot from each bar in Fig 4C is shown. (B)

Representative spectrograms from 3 participants, showing correct (top) and incorrect

trials (bottom) separately. The red solid overlay indicates each participant's (P1–P7)

respiratory signal. Black outlines indicate statistically significant clusters (P < 0.05, false-

discovery-rate corrected). The vertical short-dashed line indicates sniff onset. The

source is available in S5 Data.

1397

1398

1399   S5 Fig. Related to Fig 6. Results of modulation index computed for odor and no-odor

1400   trials, accounting for possible contribution of steep slope of sensory-evoked potentials

1401   (see supplementary Methods). The source is available in S5 Data.

1402

1403

S6 Fig. Related to Fig 2. Sniff-onset aligned raw power (left), power change relative to

baseline (middle), and z score map (right, same as Fig 2C) are shown for each

participant (P1–P7). The baseline was defined as [–0.55, –0.05] s prior to cue onset.

1407   The black overlaid line indicates the respiratory signal. The vertical short-dashed line

1408   indicates sniff onset. The source is available in S5 Data.

1409

1410   S1 Text. Supplementary methods.

1411

S1 Data. Zip file containing datasets underlying Fig 2A, 2B, and 2C.

S2 Data. Zip file containing datasets underlying Fig 3A, 3B, 3C, and 3D.

S3 Data. Zip file containing datasets underlying Fig 4A, 4B, 4C, 4D, and 4E.

S4 Data. Zip file containing datasets underlying Figs 5B, 6A, 6B, and 7.

S5 Data. Zip file containing datasets underlying S1 Fig, S2 Fig, S3 Fig, S4 Fig, S5 Fig

and S6 Fig.