1
7 **Rapalogs downmodulate intrinsic immunity and promote cell entry of SARS-CoV-2**
8
9

10 Guoli Shi[1], Abhilash I. Chiramel[2], Saliha Majdoul[1], Kin Kui Lai[1], Tirhas Dempsey[1], Adam
11 Kenney[3], Ashley Zani[3], Adrian Eddy[3], Lizhi Zhang[3], Paul A. Beare[4], Swagata Kar[5], Jacob S.
12 Yount[3], Sonja M. Best[4], Alex A. Compton[1,*]
13
14
15
16 [1]HIV Dynamics and Replication Program, Center for Cancer Research, National Cancer
17 Institute, Frederick, MD, USA
18
19 [2]Laboratory of Virology, Rocky Mountain Laboratories, National Institute of Allergy and
20 Infectious Diseases, Hamilton, MT, USA
21
22 [3]Department of Microbial Infection and Immunity, The Ohio State University, Columbus, OH,
23 USA
24
25 [4]Laboratory of Bacteriology, Rocky Mountain Laboratories, National Institute of Allergy and
26 Infectious Diseases, Hamilton, MT, USA
27
28 [5]Bioqual, Rockville, MD, USA
29
30
31 [*]Address correspondence to alex.compton@nih.gov

32
33 Running title: Rapalogs promote cell entry of SARS-CoV-2
34
35 Keywords: rapamycin, rapalog, mTOR inhibitor, IFITM, interferon, SARS-CoV-2, TFEB,
36 microautophagy, COVID-19, coronavirus, membrane fusion

Abstract

SARS-CoV-2 infection in immunocompromised individuals is associated with prolonged virus shedding and the evolution of viral variants. Rapamycin and its analogs (rapalogs, including everolimus, temsirolimus, and ridaforolimus) are FDA-approved as mTOR inhibitors in clinical settings such as cancer and autoimmunity. Rapalog use is commonly associated with increased susceptibility to infection, which has been traditionally explained by impaired adaptive immunity. Here, we show that exposure to rapalogs increases susceptibility to SARS-CoV-2 infection in tissue culture and in immunologically naïve rodents by antagonizing the cell-intrinsic immune response. By identifying one rapalog (ridaforolimus) that is less active in this regard, we demonstrate that rapalogs promote Spike-mediated entry into cells by triggering the degradation of IFITM2 and IFITM3 via an endolysosomal remodeling program known as microautophagy. Rapalogs that promote virus entry inhibit the mTOR-mediated phosphorylation of the transcription factor TFEB, which facilitates its nuclear translocation and triggers microautophagy. In rodent models of infection, injection of rapamycin prior to and after virus exposure resulted in elevated SARS-CoV-2 replication and exacerbated viral disease, while ridaforolimus had milder effects. Overall, our findings indicate that preexisting use of certain rapalogs may elevate host susceptibility to SARS-CoV-2 infection and disease by activating a lysosome-mediated suppression of intrinsic immunity.

Significance

Rapamycin is an immunosuppressant used in humans to treat cancer, autoimmunity, and other disease states. Here, we show that rapamycin and related compounds promote the first step of the SARS-CoV-2 infection cycle—entry into cells—by disarming cell-intrinsic immune defenses. We outline the molecular basis for this effect by identifying a rapamycin derivative that is inactive, laying the foundation for improved mTOR inhibitors that do not suppress intrinsic immunity. We find that rapamycin analogs that promote SARS-CoV-2 entry are those that activate TFEB, a transcription factor that triggers the degradation of antiviral membrane proteins inside of cells. Finally, rapamycin administration to rodents prior to SARS-CoV-2 challenge results in enhanced viral disease, revealing that its use in humans may increase susceptibility to infection.

Introduction

Severe acute respiratory syndrome (SARS) coronavirus (CoV)-2 emerged in humans in 2019 following a species jump from bats and a possible intermediate animal host and is the cause of COVID-19, a respiratory and multi-organ disease of variable severity [1, 2]. The characterization of virus-host interactions that dictate SARS-CoV-2 infection and COVID-19 severity is a major priority for public health [3]. Immune impairment, such as that resulting from cancer, has been associated with prolonged SARS-CoV-2 shedding and the seeding of "super-spreader" events [4-8].

One group of compounds being considered for the treatment of COVID-19-related immunopathology are rapamycin (sirolimus, Rapamune) and rapamycin analogs (rapalogs) [9-20]. As Food and Drug Administration-approved inhibitors of mammalian target of rapamycin (mTOR) kinase, these macrolide compounds are used therapeutically to inhibit the processes of

cancer, autoimmunity, graft versus host disease, atherosclerosis, and aging [21]. Rapalogs,
including everolimus (RAD-001), temsirolimus (Torisel, CCI-779), and ridaforolimus
(deforolimus, AP-23573), were developed to decrease the half-life of rapamycin *in vivo* in order
to minimize the systemic immunosuppression caused by rapamycin use, which is associated with
increased susceptibility to infections [22-26]. Differing by only a single functional group at
carbon-40 **(Figure 1)**, it is believed that rapamycin and rapalogs share the same molecular
mechanism of action to inhibit mTOR kinase—they bind to FK506-binding proteins (FKBP) and
the resulting complex physically interacts with mTOR and disrupts its signaling [25, 27].
Activation of mTOR promotes cell growth, cell proliferation, and cell survival [28]. In
addition, mTOR activation promotes pro-inflammatory T-cell differentiation and mTOR
inhibitors have been used to block lymphocyte proliferation and cytokine storm [29]. Since
respiratory virus infections like SARS-CoV-2 can cause disease by provoking hyper-
inflammatory immune responses that result in immunopathology [30-32], rapalogs are being
tested as treatments to decrease viral disease burden. At least three active clinical trials have been
designed to test the impact of rapamycin on COVID-19 severity in infected patients
(NCT04461340, NCT04341675, NCT04371640).
In addition to their potential utility for mitigating disease in individuals already infected
by SARS-CoV-2, there are also calls to use rapalogs as antiviral agents to inhibit virus infection
itself (i.e. as a prophylactic) [33]. It was recently shown that rapalogs inhibit SARS-CoV-2
replication when added to cells post-infection [34], attesting to a potential use of rapalogs as
antivirals in infected individuals. Nonetheless, rapalogs are known to induce an
immunosuppressed state in humans characterized by an increased rate of infections, including
those caused by respiratory viruses. Furthermore, rapamycin administration concurrent with
virus challenge has been shown to promote Influenza A replication in mice and to exacerbate
viral disease [35, 36], but the mechanism was unknown. We previously found that exposure of
human and murine cells to rapamycin induced the lysosomal degradation of a select group of
cellular proteins, including the interferon-inducible transmembrane (IFITM) proteins, and
rendered cells more permissive to infection by Influenza A virus and gene-delivering lentiviral
vectors [37, 38]. IFITM1, IFITM2, and IFITM3 are expressed constitutively in a variety of
tissues, are further upregulated by type-I and type-II interferons, and are important components
of cell-intrinsic immunity, the antiviral network that defends individual cells against virus
invasion [39, 40]. Nonetheless, it remained to be determined how rapamycin-mediated regulation
of intrinsic immunity impacts host susceptibility to virus infection *in vivo*.
In this report, we show that rapalogs differentially counteract the constitutive and
interferon-induced antiviral state in lung cells and increase permissiveness to SARS-CoV-2
infection. We found that the enhancing effect of rapalogs on SARS-CoV-2 infection is
functionally linked to their capacity to trigger degradation of IFITM proteins, particularly
IFITM2 and IFITM3. By identifying a rapalog that lacks this activity, we found that IFITM
protein turnover and SARS-CoV-2 infection enhancement are associated with activation of
TFEB, a master regulator of lysosome function that is regulated by mTOR. Administration of
rapamycin to naive rodents four hours prior to experimental SARS-CoV-2 infection increased
virus replication and viral disease severity, indicating for the first time that suppression of
intrinsic immunity by rapamycin contributes to its immunosuppressive properties *in vivo*.
136
137 **Results**
138

Select rapalogs promote SARS-CoV-2 infection and downmodulate IFITM proteins in lung cells

To assess how rapamycin and rapalogs impact SARS-CoV-2 infection, we took advantage of a pseudovirus system based on human immunodeficiency virus (HIV). This pseudovirus (HIV-CoV-2 S) is limited to a single round of infection, cell entry is mediated by SARS-CoV-2 Spike, and infection of target cells is measured by luciferase activity. SARS-CoV-2 can enter cells via multiple routes, and sequential proteolytic processing of Spike is essential to this process. SARS-CoV-2 Spike is cleaved at a polybasic motif (RRAR) located at the S1/S2 boundary by furin-like proteases in virus-producing cells prior to release. Subsequently, the S2' site is cleaved by the trypsin-like proteases TMPRSS2 on the target cell surface or cathepsins B and L in target cell endosomes, triggering membrane fusion at those sites [41-43].

We previously found that a four-hour pre-treatment of cells with 20 μM quantities of rapamycin triggered the degradation of human IFITM3 and enhanced cellular susceptibility to Influenza A infection [44]. Therefore, we pre-treated A549-ACE2 (transformed human lung epithelial cells that overexpress the human ACE2 receptor) with 20 μM rapamycin, everolimus, temsirolimus, ridaforolimus, or DMSO (vehicle control) for four hours and then challenged cells with HIV-CoV-2. Interestingly, we found that rapalogs promoted Spike-mediated infection to different extents: rapamycin, everolimus, and temsirolimus significantly enhanced infection (up to 5-fold) while ridaforolimus did not **(Figure 2A)**. To determine whether rapalogs promote cell permissiveness to infection by upregulating dependency factors or by downregulating restriction factors, we performed the same experiment in cells pre-treated with type-I interferon. While type-I interferon suppressed infection by approximately 90%, the addition of rapamycin, everolimus, and temsirolimus resulted in rescue of infection by up to 20-fold **(Figure 2A)**. As a result, infection levels were partially restored to those achieved in the absence of interferon, with everolimus having the greatest boosting effect and ridaforolimus, the least. Therefore, rapalogs differentially promote SARS-CoV-2 Spike-mediated infection by counteracting intrinsic antiviral defenses in lung cells to different extents.

Type-I interferon treatment of A549-ACE2 resulted in upregulation of *IFITM2* and *IFITM3*, as detected by an antibody recognizing both proteins in whole cell lysates **(Figure 2B)**. A549-ACE2 cells express low but detectable levels of IFITM2/3 in the absence of interferon treatment **(Supplemental Figure 1A)**. Consistent with our previous publication, addition of rapamycin resulted in substantial loss of IFITM2/3 protein levels from cells. In a manner that mirrored the differential effects of rapalogs on pseudovirus infection, everolimus and temsirolimus greatly diminished IFITM2/3 levels while ridaforolimus reduced IFITM2/3 to a lesser extent **(Figure 2B and Supplemental Figure 1A)**. In contrast, ACE2 levels were not affected by interferon nor by rapalog treatment. Therefore, rapamycin derivatives may facilitate infection by antagonizing constituents of intrinsic immunity, including IFITM2/3, and this activity is determined by the chemical moiety found at carbon-40 of the macrolide structure.

To extend our findings to primary lung cells, we performed similar experiments in human small airway epithelial cells (HSAEC). While these cells were not permissive to HIV-CoV-2, they were susceptible to infection by pseudovirus based on vesicular stomatitis virus (VSV-CoV-2) whereby infection is reported by GFP expression. Pre-treatment of HSAEC with rapalogs enhanced VSV-CoV-2 infection to varying extents, but as observed in A549-ACE2 cells, everolimus exhibited the greatest effect and ridaforolimus, the least. Endogenous IFITM3 was readily detected in HSAEC under basal conditions (in the absence of interferon), while IFITM1

was barely detected and IFITM2 was not detected at all, and IFITM3 levels were
downmodulated differentially by rapalogs **(Supplemental Figure 1B)**. siRNA-mediated
knockdown of IFITM3 in HSAEC resulted in enhanced VSV-CoV-2 infection, indicating that
IFITM3 restricts Spike-mediated infection in these cells **(Supplemental Figure 1C)**. We also
treated semi-transformed nasal epithelial cells known as UNCNN2TS with rapalogs in order to
assess an impact on endogenous IFITM3 levels. As observed in HSAEC, downmodulation of
IFITM3 occurred following treatment of UNCNN2TS with rapamycin, everolimus,
temsirolimus, and to a lesser extent, ridaforolimus **(Supplemental Figure 1D)**.
Since 20 μM quantities of rapalogs promoted pseudovirus infection mediated by SARS-
CoV-2 Spike, we tested how pretreatment of A549-ACE2 cells with varying amounts of
everolimus impacted infection by replication-competent SARS-CoV-2. We observed a dose-
dependent enhancement of infectious SARS-CoV-2 yield in supernatants of infected cells (up to
4-fold) **(Figure 2D)**. Therefore, everolimus boosts pseudovirus infection and SARS-CoV-2
infection to similar extents, and since Spike is the only viral component shared between the two
sources of infection, cellular entry is the infection stage inhibited by the intrinsic defenses that
are sensitive to downmodulation by rapalogs.
201
202 **Rapalogs facilitate cell entry mediated by various viral fusion proteins**
203
In order to gain a greater mechanistic understanding of the effects of rapalogs on SARS-
CoV-2 infection, we took advantage of HeLa cells overexpressing ACE2 (HeLa-ACE2). HeLa-
ACE2 were pre-treated for four hours with increasing amounts of everolimus and then
challenged with SARS-CoV-2. Everolimus increased titers of infectious virus released into
supernatants in a dose-dependent manner, and to a greater extent than was observed for A549-
ACE2 cells **(Figure 3A)**. Furthermore, we found that pre-treatment of cells with 20 μM amounts
of rapalogs enhanced SARS-CoV-2 titers to varying extents—rapamycin, everolimus, and
temsirolimus significantly boosted SARS-CoV-2 infection (up to 10-fold), while ridaforolimus
had less of an impact **(Figure 3B)**. We also performed infections of HeLa-ACE2 with HIV-CoV-
2 pseudovirus, and the results were similar: the impact of ridaforolimus was minimal while the
other three compounds significantly boosted Spike-mediated infection **(Figure 3C)**. To test the
link between infection enhancement and downmodulation of IFITM proteins by rapalogs, we
probed for levels of IFITM3, IFITM2, and IFITM1 by immunoblotting whole cell lysates using
specific antibodies. All IFITM proteins were readily detected in HeLa-ACE2 in the absence of
interferon. IFITM3, IFITM2, and IFITM1 were significantly downmodulated following
treatment with rapamycin, everolimus, and temsirolimus **(Figure 3D)**. Levels of IFITM3 were
quantified over multiple experiments and presented as an average. The results show that all
rapalogs led to significant decreases in IFITM3 protein, but ridaforolimus was least potent in this
regard **(Figure 3E)**. The loss of IFITM2/3 protein was confirmed by confocal
immunofluorescence microscopy of intact cells **(Figure 3F)**. Furthermore, prolonged treatment
(24 hours) of cells with everolimus and temsirolimus resulted in prolonged suppression of
IFITM2 and IFITM3 protein levels **(Supplemental Figure 2A)**. In contrast, ACE2 levels and
ACE2 subcellular distribution were unaffected by rapalog treatment **(Figure 3D and
Supplemental Figure 2B)**. Furthermore, rapalogs did not significantly decrease cell viability
under the conditions tested **(Supplemental Figure 2C)**.
We previously showed that lysosomal degradation of IFITM3 triggered by rapamycin
requires endosomal complexes required for transport (ESCRT) machinery and multivesicular

body (MVB)-lysosome fusion [44]. We confirmed that depletion of IFITM proteins by rapalogs occurs at the post-translational level and requires endolysosomal acidification, since bafilomycin A1 prevented their loss **(Supplemental Figure 3A-B)**. The process by which rapalogs trigger IFITM protein degradation resembles endolysosomal microautophagy, an autophagy pathway that does not require an autophagosome intermediate [45-47]. Treatment of cells with U18666A, an inhibitor of MVB formation and microautophagy, mostly prevented IFITM3 turnover in the presence of rapalogs **(Supplementary Figure 3B)**. In contrast, a selective inhibitor of vps34/PI3KC3 (essential for macroautophagy induction) did not **(Supplemental Figure 3C-D)**. Therefore, rapamycin and specific rapalogs trigger the degradation of endogenous factors mediating intrinsic resistance to SARS-CoV-2 infection, including the IFITM proteins, by promoting their turnover in lysosomes via endolysosomal microautophagy.

Enveloped virus entry into cells is a concerted process involving virus attachment to the cell surface followed by fusion of cellular and viral membranes. Since IFITM proteins are known to inhibit virus-cell membrane fusion, we quantified the terminal stage of HIV-CoV-2 entry by tracking the cytosolic delivery of beta-lactamase (BlaM) in single cells. We found that treatment of cells with rapamycin, everolimus, and temsirolimus resulted in enhanced HIV-CoV-2 entry while ridaforolimus was less impactful **(Figure 4A)**. To measure whether rapalogs promote the cell entry process driven by other coronavirus Spike proteins, we produced HIV incorporating Spike from SARS-CoV (HIV-CoV-1) or MERS-CoV (HIV-MERS-CoV). Infections by both HIV-CoV-1 and HIV-MERS-CoV were elevated by rapalog treatment in HeLa-ACE2 and HeLa-DPP4 cells, respectively, although the extent of enhancement was lower than that observed with HIV-CoV-2 **(Figure 4B-C)**. Consistently, ridaforolimus was the least active among the rapalogs tested and it did not significantly promote pseudovirus infection. Since we previously showed that rapamycin enhanced the cellular entry of Influenza A virus and VSV-G pseudotyped lentiviral vectors [44], we also assessed infection of pseudoviruses incorporating hemagglutinin (HIV-HA) or VSV G (HIV-VSV G). Rapamycin, everolimus, and especially temsirolimus boosted HA- and VSV G-mediated infections (up to 30-fold and 11-fold, respectively) **(Figure 4D-E)**. Since IFITM proteins have been previously shown to inhibit infection by SARS-CoV-1, MERS-CoV, VSV, and Influenza A virus [40], these data suggest that rapalogs promote infection, at least in part, by lowering the barrier to virus entry imposed by IFITM proteins.

IFITM2/3 mediate the rapalog-sensitive barrier to SARS-CoV-2 infection in HeLa-ACE2

To formally test the link between rapalog-mediated depletion of IFITM proteins and entry by SARS-CoV-2 Spike, we used HeLa cells in which IFITM1, IFITM2, and IFITM3 were knocked out (*IFITM1-3* KO) and introduced human ACE2 by transient transfection **(Figure 5A)**. IFITM2 alone or IFITM2 and IFITM3 were restored in *IFITM1-3* KO cells by transient overexpression **(Figure 5B)** and cells were challenged with HIV-CoV-2. Relative to WT cells, HIV-CoV-2 infection was approximately 50-fold higher in *IFITM1-3* KO cells, indicating that endogenous IFITM proteins restrict SARS-CoV-2 Spike-mediated infection in this cell type. Furthermore, while temsirolimus significantly promoted infection by 10-fold in WT cells, little to no enhancement was observed in *IFITM1-3* KO cells **(Figure 5C)**. Ectopic expression of IFITM2 inhibited infection and partially restored sensitivity to temsirolimus, while the combination of IFITM2 and IFITM3 restricted infection further and fully restored temsirolimus sensitivity. These findings indicate that temsirolimus promotes Spike-mediated infection in HeLa-ACE2 cells by lowering levels of endogenous IFITM2 and IFITM3.

Since human IFITM proteins have been reported to promote SARS-CoV-2 infection in certain cell types, including the lung epithelial cell line Calu-3 [48], we tested the impact of rapalogs on HIV-CoV-2 infection in this cell type. Here, in contrast to the enhancement observed in A549-ACE2 and HeLa-ACE2 cells, rapamycin, everolimus, and temsirolimus inhibited Spike-mediated infection in Calu-3 cells (**Supplemental Figure 4**). These results confirm that the effect of rapalog treatment on Spike-mediated infection is explained by their ability to induce the degradation of IFITM proteins.

Rapalogs differentially activate a lysosomal degradation pathway orchestrated by TFEB

Since rapamycin and rapalogs are known to inhibit mTOR signaling by binding both mTOR and FKBP12 (and other FKBP members), we sought to determine whether mTOR binding and its inhibition are required for rapalog-mediated enhancement of SARS-CoV-2 infection. To that end, we tested the effect of tacrolimus (also known as FK506), a macrolide immunosuppressant that is chemically related to rapalogs but does not bind nor inhibit mTOR. Instead, tacrolimus forms a ternary complex with FKBP12 and calcineurin to inhibit the signaling properties of the latter [49]. In HeLa-ACE2 cells, a four-hour treatment of 20 μM tacrolimus did not reduce levels of IFITM2/3 (**Supplemental Figure 5A**), nor did it boost HIV-CoV-2 infection (**Supplemental Figure 5B**). These results suggest that FKBP12 binding is not sufficient for drug-mediated enhancement of SARS-CoV-2 infection. They also suggest that the extent to which mTOR is inhibited may explain the differential degree to which infection is impacted by the immunosuppressants examined in this study. Therefore, we surveyed the phosphorylation status of TFEB, a transcription factor that controls lysosome biogenesis and degradative processes carried out by lysosomes [50]. mTOR phosphorylates TFEB at serine 211 (S211), which promotes its sequestration in the cell cytoplasm and decreases its translocation into the nucleus [50-52]. Furthermore, this phosphorylation event was previously shown to be sensitive to inhibition by rapamycin and temsirolimus [51, 53]. We found that rapamycin, everolimus, and temsirolimus significantly reduced S211 phosphorylation of endogenous TFEB in A549-ACE2 cells while ridaforolimus did so to a lesser extent (**Figure 6A-B**). Furthermore, we measured the subcellular distribution of TFEB-GFP in HeLa-ACE2 treated with different compounds and found that rapamycin, everolimus, and temsirolimus induced a significantly greater accumulation of TFEB-GFP in the nucleus (**Figure 6C-D**). These findings suggest that ridaforolimus exhibits a less potent inhibition of mTOR-mediated TFEB phosphorylation under the conditions tested. Therefore, nuclear translocation of TFEB is associated with IFITM2/3 degradation and increased cellular susceptibility to SARS-CoV-2 Spike-mediated infection. Consistent with a direct relationship between TFEB activation, IFITM2/3 turnover, and Spike-mediated cell entry, we found that ectopic expression of a constitutively active form of TFEB lacking the first 30 amino-terminal residues [50] was sufficient to trigger IFITM2/3 loss from cells (**Figure 6E**) and sufficient to increase susceptibility to HIV-CoV-2 infection (**Figure 6F**). By combining transfection of the constitutively active form of TFEB with temsirolimus treatment, we found that IFITM2/3 levels were strongly suppressed irrespective of whether TFEB was detected or not. This confirms that TFEB and rapalogs are functionally redundant and operate in the same pathway to negatively regulate IFITM2/3 levels (**Supplemental Figure 5C**). Finally, we took advantage of TFEB-deficient cells to formally address the role that TFEB activation plays during rapalog-mediated enhancement of infection (**Supplemental Figure 5D**). While rapamycin, everolimus, and temsirolimus significantly boosted HIV-CoV-2 infection in

323 HeLa WT cells transfected with ACE2, no significant enhancement was observed in HeLa *TFEB*
324 KO cells **(Figure 6G)**. In summary, our results employing functionally divergent rapalogs reveal
325 a previously unrecognized immunoregulatory role played by the mTOR-TFEB-lysosome axis
326 that affects the cell entry of SARS-CoV-2 and other viruses.
327
328 **Rapamycin enhances SARS-CoV-2 infection and viral disease *in vivo***
329
Our findings from SARS-CoV-2 and pseudovirus infection of human cells demonstrate
that rapamycin, everolimus, and temsirolimus can suppress intrinsic immunity at the post-
translational level, while ridaforolimus does so to a lesser extent. However, whether these
compounds are functionally divergent when administered *in vivo* was unclear. Since
temsirolimus is a prodrug of rapamycin (it is metabolized to rapamycin), and since rapamycin
was previously shown to promote morbidity of Influenza A infection in mice [36, 54], we tested
how intraperitoneal injection of rapamycin or ridaforolimus impacted SARS-CoV-2 replication
and disease course in naïve hamsters **(Figure 7A)**. Hamsters are a permissive model for SARS-
CoV-2 because hamster ACE2 is sufficiently similar to human ACE2 to support productive
infection. Furthermore, in contrast to transgenic mice expressing human ACE2 or mice infected
with mouse-adapted (MA) SARS-CoV-2, hamsters exhibit severe disease characterized by lung
pathology when high viral loads are achieved [55].
Eight hamsters were randomly allocated to each group (rapamycin, ridaforolimus, or
DMSO) and all received an intraperitoneal injection (3 mg/kg) 4 hours prior to intranasal
inoculation with SARS-CoV-2. Furthermore, half of the mice in each group received a second
injection on day 2 post-infection. As an indicator of infection and viral disease, we tracked
weight loss for 10 days, or less if the hamster met requirements for euthanasia (loss of 20% or
more of its body weight). We observed that hamsters receiving two injections did not exhibit
significantly different rates of weight loss compared to those receiving a single injection
(Supplemental Figure 6). As a result, we consolidated hamsters into three groups of eight
according to receipt of rapamycin, ridaforolimus, or DMSO. Relative to DMSO treatment,
hamsters injected with rapamycin or ridaforolimus exhibited significantly greater weight loss at
days 2-5 post-infection, with rapamycin-treated hamsters displaying the most weight loss
(Figure 7B). While one (1/8) of the hamsters treated with DMSO exhibited severe weight loss
necessitating euthanasia, seven (7/8) of the hamsters treated with rapamycin were euthanized
following severe weight loss between days 6 and 8 post-infection **(Figure 7C)**. Meanwhile, four
(4/8) of the hamsters treated with ridaforolimus met requirements for euthanasia on days 7 or 8
post-infection. As a result, hamsters treated with rapamycin exhibited significantly reduced
survival compared to the DMSO group **(Figure 7C)**. In contrast, survival of ridaforolimus-
treated animals did not differ significantly.
Survivors in all three groups recovered weight after day 7 post-infection and infectious
virus was not detected from the lungs of these hamsters at day 10. In contrast, the lungs of
hamsters euthanized due to severe weight loss exhibited high infectious virus titers, suggesting
that morbidity was caused by viral pathogenesis (the lungs of one hamster treated with
rapamycin were not examined because it was found dead following infection) **(Figure 7D).** To
better understand the basis for differential survival between the groups, early SARS-CoV-2
replication was measured by quantitative PCR from oral swabs. We found that hamsters injected
with rapamycin exhibited significantly higher viral RNA levels in the oral cavity at day 2 post-
infection compared to animals injected with DMSO **(Figure 7E)**. In contrast, viral RNA levels in

hamsters injected with ridaforolimus were elevated relative to the DMSO group, but they did not
differ significantly. Overall, these results suggest that rapamycin administration increases host
susceptibility to SARS-CoV-2 infection and significantly increases morbidity and mortality.
We previously found that, like its human counterpart, murine IFITM3 is sensitive to
depletion by rapamycin [44]. To determine whether rapamycin promotes host susceptibility to
SARS-CoV-2 infection in mice, we injected C57BL/6 mice with rapamycin or DMSO prior to
and after challenge with MA SARS-CoV-2 and measured infectious viral burden in lungs on day
2 post-infection (**Figure 8A**). We found that virus titers in lungs were significantly increased
(144-fold) in rapamycin-treated mice compared to DMSO-treated mice (**Figure 8B**).
Furthermore, murine IFITM3 protein levels were reduced in the lungs of mice injected with
rapamycin relative to levels found in DMSO-treated mice (**Figure 8C**). Together, these findings
support the notion that rapamycin downmodulates intrinsic barriers to infection *in vivo*.
381
382 **Discussion**
383
By assessing their impact on infection at the single-cell and whole-organism level, we
draw attention to an immunosuppressive property of rapamycin and some rapalogs that acts on
cell-intrinsic immunity and increases cellular susceptibility to infection by SARS-CoV-2 and
likely other pathogenic viruses. Side effects of rapalog use in humans, including increased risk of
respiratory tract infections, are regularly attributed to immunosuppression of adaptive immunity
[56]. Indeed, rapalogs have been used to mitigate systemic immunopathology caused by T-cell
responses, and this is one reason why they are being tested for therapeutic benefit in COVID-19
patients. However, since rapamycin was injected into immunologically naïve hosts prior to and
soon after virus challenge, it is unlikely that rapalogs modulated adaptive immunity against
SARS-CoV-2 in our experiments. While immunomodulation of adaptive immunity by rapalogs
may provide benefit for patients already suffering from COVID-19, pre-existing rapalog use may
enhance susceptibility by counteracting cell-intrinsic immunity.
The injection dose of rapamycin or ridaforolimus (3 mg/kg) that we administered once to
hamsters or daily to mice, when adjusted for body surface area and an average human weight of
60 kg [57], equates to approximately 15 mg per human. This figure is similar to those
administered to humans in clinical settings, such as the use of rapamycin for the treatment of
glioblastoma (up to 10 mg daily for multiple days), the use of temsirolimus for the treatment of
renal cell carcinoma (25 mg once weekly), or the use of everolimus for the treatment of tuberous
sclerosis (TS), a genetic disorder resulting in hyperactivation of mTOR (10 mg daily,
continuously) [23, 58-60]. Interestingly, a case report detailed the deaths of two TS patients (a
father and daughter) who, despite discontinuing everolimus upon detection of SARS-CoV-2
infection, died from severe COVID-19 in late 2020 [60]. Our findings detailing the suppression
of cell-intrinsic immunity by rapalogs raise the possibility that their use may predispose
individuals to SARS-CoV-2 infection and severe forms of COVID-19. More generally, they
provide new insight into how rapamycin and rapalogs may elicit unintended
immunocompromised states and increase human susceptibility to multiple virus infections.
By leveraging the differential functional properties of rapalogs, we reveal how the
mTOR-TFEB-lysosome axis impacts intrinsic resistance to SARS-CoV-2 infection. Specifically,
rapamycin and select rapalogs (everolimus and temsirolimus) promote infection at the stage of
cell entry, and this is functionally linked to nuclear accumulation of TFEB and the lysosomal
degradation of IFITM proteins by endolysosomal microautophagy (**Figure 9**). While mTOR

phosphorylates TFEB at S211 to promote the sequestration of TFEB in the cytoplasm, the phosphatase calcineurin dephosphorylates TFEB at this position to promote nuclear translocation [61]. Therefore, the extent to which different rapalogs promote nuclear TFEB accumulation may be a consequence of differential mTOR inhibition and/or differential calcineurin activation. Calcineurin is activated by calcium release through the lysosomal calcium channel TRPML1 (also known as mucolipin-1) [61], and interestingly, it was shown that rapamycin and temsirolimus, but not ridaforolimus, promote calcium release by TRPML1 [53]. Therefore, it is worth examining whether TRPML1 or related lysosomal calcium channels are required for the effects of rapalogs on virus infection. Overall, our findings reveal a previously unrecognized mechanism by which TFEB promotes virus infections—inhibition of cell-intrinsic defenses restricting virus entry. We show that nuclear TFEB induces the degradation of IFITM proteins, but it may also trigger the loss or relocalization of other antiviral factors that remain to be uncovered. Furthermore, TFEB-mediated induction of dependency factors, such as cathepsin L, is likely to partially contribute to the overall impact of rapalogs on SARS-CoV-2 infection. Overall, this work identifies TFEB as a therapeutic target, and inhibitors that limit levels of nuclear TFEB could be mobilized for broad-spectrum antiviral activity.

We previously demonstrated that treatment of cells with micromolar quantities of rapamycin induced the lysosomal degradation of IFITM2/3 via a pathway that is independent of macroautophagy yet dependent upon endosomal complexes required for transport (ESCRT)-mediated sorting of IFITM2/3 into intraluminal vesicles of late endosomes/MVB [37]. This MVB-mediated degradation pathway is also referred to as microautophagy, which occurs directly on endosomal or lysosomal membranes and involves membrane invagination [62]. In both yeast and mammalian cells, microautophagy is characterized by ESCRT-dependent sorting of endolysosomal membrane proteins into intraluminal vesicles followed by their degradation by lysosomal hydrolases [63]. While microautophagy selectively targets ubiquitinated endolysosomal membrane proteins, cytosolic proteins can also be non-selectively internalized into intraluminal vesicles and degraded [64, 65]. Interestingly, microautophagy is known to be regulated by mTOR [66, 67], and mTOR inhibition triggers a ubiquitin- and ESCRT-dependent turnover of vacuolar (lysosomal) membrane proteins in yeast [68, 69]. Overall, our findings suggest that select rapalogs induce a rapid, TFEB-dependent, endolysosomal membrane remodeling program known as microautophagy, and IFITM proteins are among the client proteins subjected to this pathway. The full cast of cellular factors that orchestrate this selective degradation program in mammalian cells and the other client proteins subjected to it will need to be worked out. Interestingly, the E3 ubiquitin ligase NEDD4 was previously shown to ubiquitinate IFITM2 and IFITM3 and to induce their lysosomal degradation in mammalian cells [70, 71], while Rsp5, the yeast ortholog of NEDD4, was shown to ubiquitinate vacuolar proteins turned over by microautophagy in yeast [72]. Therefore, rapamycin and select rapalogs may upregulate NEDD4 function, resulting in selective degradation of a subset of the cellular proteome that includes IFITM proteins. Indeed, NEDD4 and the related NEDD4L are among the known target genes regulated by TFEB [73].

The relationship between IFITM proteins and human coronaviruses is complex. It was previously shown that IFITM3 facilitates replication of the seasonal coronavirus hCoV-OC43 [74], while we and others recently showed that SARS-CoV-1 and SARS-CoV-2 infection is inhibited by ectopic and endogenous IFITM1, IFITM2, and IFITM3 from mice and humans [75-79]. Intriguingly, mutants of human IFITM3 that lack the capacity to internalize into endosomes lost antiviral activity and promoted SARS-CoV-2 and MERS-CoV infection, revealing that

IFITM3 can either inhibit or enhance infection depending on its subcellular localization [75, 80].
Furthermore, one study reported that endogenous human IFITM proteins promoted infection by
SARS-CoV-2 in certain human tissues [48]. Overall, the net effect of human IFITM proteins on
SARS-CoV-2 infection *in vivo* remains unclear. However, the impact of rapamycin in our
experimental SARS-CoV-2 infections of hamsters and mice suggests that rapamycin-mediated
loss of IFITM proteins favors virus infection and viral disease, consistent with IFITM proteins
performing antiviral roles against SARS-CoV-2 in those species. Accordingly, it was recently
demonstrated that mouse IFITM3 protects mice from viral pathogenesis following MA SARS-
CoV-2 infection [81].
Other lines of evidence support an antiviral role for IFITM proteins during SARS-CoV-2
infection in humans. While SARS-CoV-2 infection has been shown to cause deficiencies in
interferon synthesis and interferon response pathways, administration of type I interferon *in vivo*
promotes SARS-CoV-2 clearance in hamsters and humans [82]. Notably, IFITM3 is among the
most highly induced genes in primary human lung epithelial cells exposed to SARS-CoV-2 [83,
84], and humans experiencing mild or moderative COVID-19 showed elevated induction of
antiviral genes, including *IFITM1* and *IFITM3*, in airway epithelium compared to individuals
suffering from more severe COVID-19 [85]. Single nucleotide polymorphisms in human *IFITM3*
known as ns12252 and rs34481144, which lead to IFITM3 loss-of-function, have been associated
with severe outcomes following Influenza A virus infection as well as severe COVID-19 [86,
87]. These data suggest that cell-intrinsic immunity in airways plays a role in restricting virus
spread and constraining systemic pathology during infection. Therefore, downmodulation of
IFITM proteins by select rapalogs may contribute to the immunocompromised state that these
drugs are well known to elicit in humans. This possibility warrants the close examination of
different rapalog regimens on respiratory virus acquisition and disease in humans.
485
486 **Materials and Methods**
487
488 **Cell lines, cell culture, inhibitors, and cytokines**
489
HEK293T (CRL-3216) and Calu-3 (HTB-55) cells were obtained from ATCC. HeLa-ACE2,
HeLa-DPP4, and A549-ACE2 cell lines were produced by transducing cells with lentivirus
packaging pWPI encoding ACE2 or DPP4 and selecting with blasticidin. HeLa IFITM1/2/3
Knockout (C5-9) cells were purchased from ATCC (CRL-3452). HeLa *TFEB* KO cells were
kindly provided by Ramnik J. Xavier (Broad Institute) and were described in [88]. Primary
human small airway (lung) epithelial cells (HSAEC) were purchased from ATCC (PCS-301-
010). The partially immortalized nasal epithelial cell line (UNCNN2TS) was kindly provided by
Scott H. Randell (University of North Carolina School of Medicine). Vero E6 cells (NR-53726)
were obtained from BEI Resources. Vero-TMPRSS2 cells were a kind gift from Shan-Lu Liu
(The Ohio State University). All cells were cultured at 37°C with 5% CO_2 in Dulbecco's
Modified Eagle Medium (DMEM) supplemented with 10% fetal bovine serum (HyClone,
Cytiva), except for UNCNN2TS, which were cultured in EpiX Medium (Propagenix), and
HSAEC, which were cultured with airway epithelial cell basal medium (ATCC, PCS-300-030)
and the bronchial epithelial cell growth kit (ATCC, PCS-300-040). Rapamycin (553211) was
obtained from Sigma. Everolimus (S1120), temsirolimus (S1044), ridaforolimus (S5003),
tacrolimus (S5003), and SAR405 (S7682) were obtained from Selleckchem. U18666A (U3633)

and Bafilomycin A1 (SML1661) were obtained from Sigma. Type-I interferon (human
recombinant interferon-beta$_{ser17}$, NR-3085) was obtained from BEI Resources.
508
509 **Plasmids and RNA interference**
510
pcDNA3.1 encoding human ACE2 was kindly provided by Thomas Gallagher (Loyola
University). pcDNA3.1 encoding CoV-1 Spike or CoV-2 Spike tagged with a C9 epitope on the
C-terminus, or MERS Spike, was kindly provided by Thomas Gallagher (Loyola University).
pcDNA3.1 encoding CoV-1 Spike or CoV-2 Spike tagged with a FLAG epitope on the C-
terminus was obtained from Michael Letko and Vincent Munster (NIAID). pMD2.G encoding
VSV-G (12259) was obtained from Addgene (a generous gift from Didier Trono). pWPI was
obtained from Addgene (12254) and human ACE2 or human TMPRSS2 was introduced by
Gateway cloning (Gateway LR Clonase II Enzyme mix (11791020)) as per manufacturer's
instructions. pPolII encoding hemagglutinin (HA) or neuraminidase (NA) from Influenza
A/Turkey/1/2005 (H5N1) were kindly provided by Richard Yi Tsun Kao (The University of
Hong Kong). pCMV encoding HIV-1 Vpr fused to beta lactamase (pCMV4-BlaM-Vpr) was
obtained from Addgene (21950). A plasmid encoding replication-incompetent HIV-1 lacking *env*
and *vpr* and encoding luciferase (pNL4-3LucR-E-) was kindly provided by Vineet KewalRamani
(National Cancer Institute). A plasmid encoding replication-incompetent HIV-1 lacking *env*
(pNL4-3E-) was kindly provided by Olivier Schwartz (Institut Pasteur). pEGFP-N1-TFEB
(38119) and pEGF-N1-Δ30TFEB (44445) were obtained from Addgene (a generous gift of
Shawn M. Ferguson). pEGFP-2xFYVE (140047) was obtained from Addgene (a gift from
Harald Stenmark). Silencer Select siRNA targeting IFITM3 (s195035) and a non-targeting
control (No. 1) was obtained from Ambion. Cells were transfected with 20 nM siRNA using
Opti-MEM (Gibco) and Lipofectamine RNAiMAX (Thermo Fisher).
531
532 **Virus and pseudovirus infections**
533
SARS-CoV-2 isolate USA-WA1/2020 (MN985325.1) was provided by the Centers for Disease
Control or by BEI Resources (NR-52281). Virus propagation was performed in Vero E6 cells.
Mouse-adapted (MA) SARS-CoV-2 variant MA10 (in the USA-WA1/2020 backbone) [89] was
obtained from BEI Resources (NR-55329). Virus propagation was performed in Vero E6 cells
and subsequently in Vero-TMPRSS2 cells. Virus was sequenced to ensure lack of tissue culture
adaptations, including furin cleavage site mutations. Virus titers were calculated by plaque assay
performed in Vero E6 cells as follows: serial 10-fold dilutions were added to Vero E6
monolayers in 48-well plates for 1 hour at 37°C. Cells were overlayed with 1.5% carboxymethyl
cellulose (Sigma) in modified Eagle's medium containing 3% fetal bovine serum (Gibco), 1 mM
L-glutamine, 50 units per mL penicillin and 50 µg per mL streptomycin. Three days post-
infection, cells were fixed in 10% formalin and stained with crystal violet to visualize and count
plaques as previously described [90]. Titers were calculated as plaque forming units per mL and
normalized as described in the figure captions. HIV-based pseudovirus was produced by
transfecting HEK293T cells with 12 µg of pNL4-3LucR-E- and 4 µg of plasmid encoding viral
glycoproteins (pcDNA3.1 Spike (CoV-1, CoV-2, or MERS), pMD2.G-VSV-G, or 2 µg of
pPol1II-HA and 2 µg of pPol1II-NA) using TransIT-293 (Mirus). Virus supernatant was
harvested 72 hours post-transfection and filtered through 0.22 µm filters. Pseudovirus titers were
determined by p24 ELISA (XpressBio) and 100 ng p24 equivalent was added to target cells and

incubated for 72 hours prior to lysis with Passive Lysis Buffer (Promega). Luciferase activity was measured using the Luciferase Assay System (Promega). VSV-based pseudovirus was produced as previously described [91]. In brief, HEK293T cells were transfected with 2 µg pcDNA3.1 CoV-2 Spike using Lipofectamine2000 (Thermo Fisher). At 24 hours post-transfection, culture medium was removed from cells and 2 mL of VSV-luc/GFP + VSV-G (seed particles) was added. At 48 hours post-infection, virus supernatants were collected, clarified by centrifugation at 500xG for 5 mins, and stored. 50 µL of virus supernatants were added to target cells for a period of 24 hours prior to fixation with 4% paraformaldehyde (for measurements of GFP+ cells with flow cytometry). For infections with replication-competent SARS-CoV-2, rapamycin, everolimus, temsirolimus, or ridaforolimus (20 µM) were used to pretreat cells for 4 hours and then drugs were washed away prior to addition of virus at a multiplicity of infection (MOI) of 0.1. DMSO (Sigma) was used as a vehicle control. At one hour post-virus addition, cells were washed once with 1X PBS and overlayed with complete medium. Supernatants were harvested 24 hours later, and titers were determined on plaque assays performed in Vero E6 cells. For single-round infections using HIV- or VSV-based pseudovirus, rapamycin, everolimus, temsirolimus, ridaforolimus, or tacrolimus (20 µM) were used to pretreat cells for 4 hours and were maintained for the duration of infection and until harvest of cells for luciferase assay or flow cytometry. DMSO (Sigma) was used as a vehicle control.

FRET-based virus entry assay

HIV-based pseudovirus incorporating BlaM-Vpr and CoV-2 Spike was produced by transfecting HEK293T cells with pNL4-3E- (15 µg), pCMV4-BlaM-Vpr (5 µg), and pcDNA3.1 CoV-2 Spike (5 µg) using the calcium phosphate technique. Briefly, six million 293T cells were seeded in a T75 flask. Plasmid DNA was mixed with sterile H_2O, $CaCl_2$, and Tris-EDTA (TE) buffer, and the totality was combined with Hepes-buffered saline (HBS). The transfection volume was added dropwise, and cells were incubated at 37°C for 48 h. Supernatants were recovered and clarified by centrifugation, passed through a 0.45 µm filter, and stored. Titers were measured using an HIV-1 p24 ELISA kit (XpressBio). 50 ng p25 equivalent of virus was added to HeLa-ACE2 cells for 2 hours. Cells were washed and labeled with the CCF2-AM β-lactamase Loading Kit (Invitrogen) for 2 hours and analyzed for CCF2 cleavage by flow cytometry as described [92]. Rapamycin, everolimus, temsirolimus, or ridaforolimus (20 µM) were used to pretreat cells for 4 hours prior to virus addition and were maintained for the duration of infection. DMSO (Sigma) was used as a vehicle control.

Western blot, antibodies, and flow cytometry

Whole cell lysis was performed with RIPA buffer (Thermo Fisher) supplemented with Halt Protease Inhibitor EDTA-free (Thermo Fisher). Lysates were clarified by centrifugation and supernatants were collected and stored. Protein concentration was determined with the Protein Assay Kit II (Bio-Rad), and 10-15 µg of protein was loaded into 12% acrylamide Criterion XT Bis-Tris Precast Gels (Bio-Rad). Electrophoresis was performed with NuPage MES SDS Running Buffer (Invitrogen) and proteins were transferred to Amersham Protran Premium Nitrocellulose Membrane, pore size 0.20 µm (GE Healthcare). Membranes were blocked with Odyssey Blocking Buffer (Li-COR) and incubated with the following primary antibodies diluted in Odyssey Antibody Diluent (Li-COR): anti-IFITM1 (60074-1-Ig; Proteintech), anti-IFITM2

(66137-1-Ig; Proteintech), anti-IFITM3 (EPR5242, ab109429; Abcam), anti-Fragilis (ab15592;
Abcam (detects murine IFITM3)), anti-IFITM2/3 (66081-1-Ig; Proteintech), anti-actin (C4, sc-
47778; Santa Cruz Biotechnology), anti-hACE2 (ab15348; Abcam), anti-TFEB (4240S; Cell
Signaling Technology), and anti-pTFEB (Ser211) (37681S; Cell Signaling Technology).
Secondary antibodies conjugated to DyLight 800 or 680 (Li-Cor) and the Li-Cor Odyssey CLx
imaging system were used to reveal specific protein detection. Images were analyzed (including
signal quantification) and assembled using ImageStudioLite (Li-Cor). Cell viability was
measured using LIVE/DEAD Red Dead Cell Stain Kit (Thermo Fisher). Cells were fixed and
permeabilized with Cytofix/Cytoperm reagent (BD) for 20 minutes and washed in Perm/Wash
buffer (BD). Flow cytometry was performed on an LSRFortessa (BD).

Confocal fluorescence and immunofluorescence microscopy

HeLa-ACE2 cells were fixed with 4% paraformaldehyde, stained with anti-IFITM2/3 (66081-1-
Ig; Proteintech), goat anti-mouse IgG Alexa Fluor 647 (A21235; Thermo Fisher) and DAPI
(62248; Thermo Fisher), and imaged in a glass-bottom tissue culture plate with an Operetta CLS
High-Content Analysis System (Perkin Elmer). For measurement of TFEB-GFP
nuclear/cytoplasmic distribution, HeLa-ACE2 cells were transfected with pEGFP-N1-TFEB for
24 hours, fixed with 4% paraformaldehyde, stained with HCS CellMask Red Stain (H32712;
Thermo Fisher) and DAPI, and imaged with an Operetta CLS. Using Harmony software (Perkin
Elmer), nuclear/cytoplasmic ratios of TFEB-GFP were calculated in single cells as follows: cells
were delineated by CellMask Red Stain, nuclei were delineated by DAPI, nuclear TFEB-GFP
was designated as GFP overlapping with DAPI, and cytoplasmic TFEB-GFP was designated as
total GFP signal minus nuclear TFEB-GFP. Average ratios were calculated from 20-30 cells per
field, and the mean of averages from 10 fields was obtained (total of approximately 250 cells per
condition). For measurement of IFITM2/3 levels in cells transfected with TFEBΔ30-GFP, HeLa-
ACE2 cells were transfected with pEGF-N1-Δ30TFEB for 24 hours, fixed and permeabilized
with BD Cytofix/Cytoperm (Fisher Scientific), stained with anti-IFITM2/3 and goat anti-mouse
IgG Alexa Fluor 647, and imaged with an Operetta CLS. The IFITM2/3 fluorescence intensity
within a single, medial Z section was measured in approximately 150 GFP-negative cells and
150 GFP-positive cells using the freehand selections tool in ImageJ.

In vivo infections of hamsters and mice with SARS-CoV-2

Male Golden Syrian hamsters between the ages of 6-8 weeks were acclimated for 11 days
following receipt. Hamsters received an intraperitoneal injection (500 µL) of rapamycin (HY-
10219; MedChemExpress) or ridaforolimus (HY-50908; MedChemExpress) at 3 mg/kg or an
equivalent amount of DMSO (8 hamsters per group). Four hours later, hamsters were challenged
with 6 x 10^3 plaque forming units of SARS-CoV-2 isolate USA-WA1/2020 (amplified on Calu-3
cells) through intranasal inoculation (50 µL in each nare). Half of the hamsters in each group
received a second injection at day 2 post-infection. Clinical observations and weights were
recorded daily up until day 10 post-infection. According to Institutional Animal Care and Use
Committee human euthanasia criteria, hamsters were euthanized immediately if weight loss
exceeded 20%. Otherwise, hamsters were euthanized on day 10 post-infection. Oral swabs were
collected on day 2 post-infection for measurement of viral RNA by quantitative PCR of the viral
N (nucleocapsid) gene. Lungs were harvested following euthanasia (day 10 or earlier) and

infectious viral load was determined by TCID$_{50}$ assay in Vero-TMPRSS2 cells. C57BL/6 mice
received an intraperitoneal injection of 3 mg/kg rapamycin (NC9362949; LC-Laboratories) or an
equivalent amount of DMSO (7 and 6 mice per group, respectively). The following day, mice
were challenged intranasally with 6 x 10^4 TCID$_{50}$ equivalent of MA10 SARS-CoV-2 (USA-
WA1/2020 backbone). Mice received a second injection of rapamycin or DMSO on the day of
infection and a third on day one post-infection. Mice were euthanized for lung harvest on day
two post-infection. Infectious viral load was determined by TCID$_{50}$ assay in Vero-TMPRSS2
cells. Animal studies were conducted in compliance with all relevant local, state, and federal
regulations and were approved by the Institutional Animal Care and Use Committee of Bioqual
and of the Ohio State University.

Figure Legends

Figure 1: Rapamycin and its analogs share a macrolide structure but differ by the
functional group present at carbon-40. Violet and green bubbles indicate the FKBP- and
mTOR-binding sites, respectively.

Figure 2: Rapalogs promote SARS-CoV-2 infection in lung epithelial cells to different
extents by counteracting the intrinsic antiviral state (A) A549-ACE2 were treated with or
without type I interferon (250 U/mL) for 18 hours and then treated with 20 µM rapamycin (Rap),
everolimus (Eve), temsirolimus (Tem), ridaforolimus (Rid), or an equivalent volume of DMSO
(D) for 4 hours. HIV-CoV-2 (100 ng p24 equivalent) was added to cells and infection was
measured by luciferase activity at 48 hours post-infection. Luciferase units were normalized to
100 in the DMSO condition in the absence of interferon. (B) A549-ACE2 cells from (A) were
subjected to SDS-PAGE and Western blot analysis. Immunoblotting was performed with anti-
IFITM2/3, anti-ACE2, and anti-actin (in that order) on the same nitrocellulose membrane.
Numbers and tick marks indicate size (kilodaltons) and position of protein standards in ladder.
(C) Primary HSAEC were treated with 20 µM Rap, Eve, Tem, Rid, or an equivalent volume of
DMSO for 4 hours. VSV-CoV-2 (50 µL) was added to cells and infection was measured by GFP
expression at 24 hours post-infection using flow cytometry. (D) A549-ACE2 were treated with
varying concentrations of Eve or DMSO (equivalent to 30 µM of Eve) for 4 hours. SARS-CoV-2
(nCoV-WA1-2020; MN985325.1) was added to cells at an MOI of 0.1 and infectious titers were
measured in VeroE6 cells by calculating the TCID$_{50}$ per mL of supernatants recovered at 24
hours post-infection. TCID$_{50}$ per mL values were normalized to 100 in the DMSO condition.
Means and standard error were calculated from 3-5 experiments. Statistical analysis was
performed with one-way ANOVA and asterisks indicate significant difference from DMSO. *, p
< 0.05; **, p < 0.01. Rel.; relative.

Figure 3: Rapalogs promote SARS-CoV-2 infection in HeLa-ACE2 cells. (A) HeLa-ACE2
were treated with varying concentrations of Eve or DMSO for 4 hours. SARS-CoV-2 (nCoV-
WA1-2020; MN985325.1) was added to cells at an MOI of 0.1 and infectious titers were
measured in VeroE6 cells by calculating the TCID$_{50}$ of supernatants recovered at 24 hours post-
infection. TCID$_{50}$ per mL values were normalized to 100 in the DMSO condition. (B) HeLa-
ACE2 were treated with 20 µM Rap, Eve, Tem, Rid, or an equivalent volume of DMSO for 4
hours. SARS-CoV-2 (nCoV-WA1-2020; MN985325.1) was added to cells at an MOI of 0.1 and
infectious titers were measured in VeroE6 cells by calculating the TCID$_{50}$ per mL of

supernatants recovered at 24 hours post-infection. $TCID_{50}$ per mL values were normalized to 100
in the DMSO condition. (C) HeLa-ACE2 were treated with 20 µM Rap, Eve, Tem, Rid, or an
equivalent volume of DMSO for 4 hours. HIV-CoV-2 (100 ng p24 equivalent) was added to cells
and infection was measured by luciferase activity at 48 hours post-infection. Luciferase units
were normalized to 100 in the DMSO condition. (D) HeLa-ACE2 cells from (C) were subjected
to SDS-PAGE and Western blot analysis. Immunoblotting was performed with anti-IFITM2,
anti-IFITM1, anti-IFITM3, anti-ACE2, and anti-actin (in that order) on the same nitrocellulose
membrane. (E) IFITM3 levels from (D) were normalized to actin levels and summarized from 5
independent experiments. (F) HeLa-ACE2 were treated with 20 µM Rap, Eve, Tem, Rid, or an
equivalent volume of DMSO for 4 hours and cells were fixed, stained with DAPI and anti-
IFITM2/3, and imaged by confocal immunofluorescence microscopy. Images represent stacks of
5 Z-slices and one representative image is shown per condition. Means and standard error were
calculated from 3-5 experiments. Statistical analysis was performed with one-way ANOVA and
asterisks indicate significant difference from DMSO. *, $p < 0.05$; **, $p < 0.01$. Rel.; relative.
700
Figure 4: Rapalogs promote cell entry mediated by diverse viral fusion proteins. (A) HeLa-
ACE2 were treated with 20 µM Rap, Eve, Tem, Rid, or an equivalent volume of DMSO for 4
hours. HIV-CoV-2 S pseudovirus incorporating BlaM-Vpr (HIV-BlaM-CoV-2) was added to
cells for 2 hours and washed. Cells were incubated with CCF2-AM for an additional 2 hours and
fixed. Cleaved CCF2 was measured by flow cytometry. Dot plots visualized as density plots
from one representative experiment are shown on the left and the percentage of CCF2+ cells
which exhibit CCF2 cleavage is indicated. Summary data representing the average of four
experiments is shown on the right. (B) HIV-CoV-1, (C) HIV-MERS-CoV, (D) HIV-IAV HA, or
(E) HIV-VSV G were added to HeLa-ACE2 or HeLa-DPP4 cells as in (A) and infection was
measured by luciferase activity at 48 hours post-infection. Luciferase units were normalized to
100 in the DMSO condition. Means and standard error were calculated from 3-4 experiments.
Statistical analysis was performed with one-way ANOVA and asterisks indicate significant
difference from DMSO. *, $p < 0.05$; **, $p < 0.01$. Rel.; relative.
714
Figure 5: Select rapalogs enhance Spike-mediated infection in HeLa-ACE2 by inhibiting
IFITM2 and IFITM3. (A) HeLa WT and HeLa *IFITM1-3* KO cells were transiently transfected
with 0.150 µg pcDNA3.1-hACE2 for 24 hours. Whole cell lysates were subjected to SDS-PAGE
and Western blot analysis. Immunoblotting was performed with anti-IFITM2, anti-IFITM3, anti-
IFITM1, anti-ACE2, and anti-actin (in that order) on the same nitrocellulose membrane. (B)
HeLa *IFITM1-3* KO were transfected with IFITM2 or IFITM2 and IFITM3 and SDS-PAGE and
Western blot analysis was performed. (C) HIV-CoV-2 was added to transfected cells from (B)
and infection was measured by luciferase activity at 48 hours post-infection. Luciferase units
were normalized to 100 in HeLa WT cells treated with DMSO. Means and standard error were
calculated from 5 experiments. Statistical analysis was performed with one-way ANOVA and
asterisks indicate significant difference from nearest DMSO condition. *, $p < 0.05$; **, $p < 0.01$.
Rel.; relative.
727
728 **Figure 6: Nuclear TFEB triggers IFITM2/3 turnover, promotes Spike-mediated infection,**
729 **and is required for enhancement of infection by rapalogs.** (A) A549-ACE2 were treated with
730 20 µM Rap, Eve, Tem, Rid, or tacrolimus (Tac) for 4 hours and whole cell lysates were
731 subjected to SDS-PAGE and Western blot analysis. Immunoblotting was performed with anti-

TFEB and anti-pTFEB (S211). (B) pTFEB (S211) levels were divided by total TFEB levels and summarized as an average of 3 experiments. (C) HeLa-ACE2 were transfected with TFEB-GFP for 24 hours, treated with Rap, Eve, Tem, Rid or Tac for 4 hours, stained with DAPI and CellMask, and imaged by high-content confocal fluorescence microscopy. Representative images from each condition are shown (DAPI and CellMask channels are not shown). (D) The ratio of nuclear to cytoplasmic TFEB-GFP was calculated in individual cells and the average ratio derived from 50-100 cells per condition is shown. (E) HeLa-ACE2 were transfected with 0.5 µg TFEB∆30-GFP for 24 hours, fixed, stained with anti-IFITM2/3, and imaged by confocal immunofluorescence microscopy. A representative field is shown on the left. The average intensity of IFITM2/3 levels in approximately 150 GFP-negative and 150 GFP-positive cells were grouped and summarized from two independent transfections on the right. (F) HeLa-ACE2 were transfected with 0.5 µg TFEB∆30-GFP, or not transfected, for 24 hours and HIV-CoV-2 (100 ng p24 equivalent) was added to cells. Infection was measured by luciferase activity at 48 hours post-infection. Luciferase units were normalized to 100 in the non-transfected condition. (G) HeLa WT or HeLa *TFEB* KO were transfected with 0.3 µg pcDNA3.1-hACE2 for 24 hours and treated with 20 µM Rap, Eve, Tem, Rid, or a corresponding volume of DMSO for 4 hours. HIV-CoV-2 (100 ng p24 equivalent) was added to cells and infection was measured by luciferase activity at 48 hours post-infection. Luciferase units were normalized to 100 in the non-transfected condition. Means and standard error were calculated from 3 experiments, except for TFEB-GFP imaging experiments, for which 2 experiments (transfections) were performed. Statistical analysis was performed with one-way ANOVA and asterisks indicate significant difference from DMSO. *, $p < 0.05$; **, $p < 0.01$. Rel.; relative. A.u.; arbitrary units.

Figure 7: Rapamycin injection into hamsters intensifies viral disease during SARS-CoV-2 infection. (A) Schematic of intraperitoneal injections and intranasal SARS-CoV-2 challenge in hamsters. Golden Syrian hamsters were injected with 3 mg/kg Rap, Rid, or equivalent amounts of DMSO (4 animals per group). Four hours later, hamsters were infected intranasally with 6 x 10^3 plaque forming units of SARS-CoV-2. At 2 days post-infection, half of the animals received a second injection of Rap, Rid, or DMSO. Oral swabs were taken and used for measurement of oral viral RNA load by qPCR. At 10 days post-infection (or earlier, if more than 20% of weight loss occurred), hamsters were euthanized, and lungs were harvested for determination of infectious virus titer by TCID$_{50}$ assay in Vero-TMPRSS2 cells (B) Mean body weight and standard error for each treatment group is plotted by day post-infection. (C) Kaplan-Meier survival curves were generated according to the dates of euthanasia (or in one case, when an animal was found dead). (D) Viral RNA copy number was determined by qPCR from oral swab at 2 days post-infection. Data is depicted as box and whiskers plots. (E) Infectious virus titers in lungs were determined by TCID$_{50}$ in Vero-TMPRSS2 cells. Data is depicted as floating bars and is grouped by brackets according to hamsters that survived until 10 days post-infection and those that were euthanized at 7 days post-infection. Statistical analysis in (B) was performed by student's T test and asterisks indicate significant difference from DMSO (gray asterisks for Rap and black asterisks for Rid). Statistical analysis in (C) was performed by comparing survival curves between Rap and DMSO or Rid and DMSO using the Log-rank (Mantel-Cox) test. Illustration created with BioRender.com.

Figure 8: Rapamycin injection into mice downmodulates IFITM3 in lungs and boosts MA SARS-CoV-2 titers in lungs. (A) Schematic of intraperitoneal injections and intranasal mouse-

adapted (MA) SARS-CoV-2 challenge in mice. C57BL/6 mice were injected with 3 mg/kg of Rap or an equivalent amount of DMSO (6 or 7 mice per group, respectively). The following day, mice were infected intranasally with 6 x 10^4 TCID$_{50}$ MA SARS-CoV-2. Mice received second and third injections of Rap or DMSO on the day of infection and on day 1 post-infection, respectively. (B) Lungs were harvested from infected mice upon euthanasia at day 2 post-infection and infectious viral loads were determined by TCID$_{50}$ in Vero-TMPRSS2 cells. Geometric mean TCID$_{50}$ per gram was calculated per treatment group and data is depicted as box and whiskers plots. Statistical analysis was performed with Mann-Whitney test and asterisks indicate significant difference from DMSO. *, $p < 0.05$; **, $p < 0.01$. (C) Lung homogenates (3 µg) from mice injected with Rap or DMSO were subjected to SDS-PAGE and Western blot analysis. Immunoblotting was performed with anti-Fragilis/IFITM3 (ab15592) and anti-actin. Illustration created with BioRender.com.

Figure 9: Model for rapalog-mediated enhancement of SARS-CoV-2 infection. Rapamycin and rapalogs everolimus and temsirolimus potently inhibit the phosphorylation of TFEB by mTOR, while ridaforolimus does not. As a result, TFEB translocates into the nucleus and induces genes functioning in lysosomal activities, including autophagy-related pathways. Nuclear TFEB triggers a microautophagy pathway that results in accelerated degradation of membrane proteins IFITM2 and IFITM3. Loss of IFITM2/3 promotes SARS-CoV-2 entry into cells by facilitating fusion between viral membranes and cellular membranes. Illustration created with BioRender.com.

Supplemental Figure 1: (A) A549-ACE2 cells were treated with 20 µM Rap, Eve, Rid, Tem, or an equivalent volume of DMSO (in the absence of type-I interferon) for 4 hours and whole cell lysates were subjected to SDS-PAGE and Western blot analysis. Immunoblotting was performed with anti-IFITM2/3 and anti-actin. (B) Primary HSAEC were treated with 20 µM Rap, Eve, Tem, Rid, or an equivalent volume of DMSO for 4 hours and whole cell lysates were subjected to SDS-PAGE and Western blot analysis. Immunoblotting was performed with anti-IFITM2 (not detected), anti-IFITM3, anti-IFITM1, and anti-actin. (C) Primary HSAEC were transfected with siRNA targeting IFITM3 or control siRNA for 48 hours. VSV-CoV-2 (50 µL) was added to cells and infection was measured by GFP expression at 24 hours post-infection using flow cytometry. siRNA-transfected cells were subjected to SDS-PAGE and Western blot analysis. Immunoblotting was performed with anti-IFITM2 (not detected), anti-IFITM3, anti-IFITM1, and anti-actin. (D) Semi-transformed nasal epithelial cells (UNCNN2TS) were treated with 20 µM Rap, Eve, Tem, Rid, or an equivalent volume of DMSO for 4 hours and whole cell lysates were subjected to SDS-PAGE and Western blot analysis. Immunoblotting was performed with anti-IFITM2 (not detected), anti-IFITM3, anti-IFITM1, and anti-actin. Immunoblots are representative of 3 independent experiments. Means and standard error were calculated from 3 experiments. Statistical analysis was performed with student's T test and asterisks indicate significant difference from control siRNA. *, $p < 0.05$; **, $p < 0.01$. Rel.; relative.

Supplemental Figure 2: (A) HeLa-ACE2 were treated with 20 µM Rap, Eve, Tem, Rid, or an equivalent volume of DMSO for 24 hours and whole cell lysates were subjected to SDS-PAGE and Western blot analysis. Immunoblotting was performed with anti-IFITM2/3 and anti-actin. (B) HeLa-ACE2 were treated with 20 µM Rap, Eve, Tem, Rid, or an equivalent volume of DMSO, in the presence or absence of 1 µM Bafilomycin A1, for 4 hours and whole cell lysates

were subjected to SDS-PAGE and Western blot analysis. Immunoblotting was performed with
anti-IFITM2, anti-IFITM1, anti-IFITM3, and anti-actin (in that order) on the same nitrocellulose
membrane. (C) HeLa-ACE2 cells were transected with FYVE-GFP for 24 hours followed by
treatment with 100 nM SAR405 or an equivalent volume of ethanol (vehicle) for 3 hours. Cells
were fixed and imaged by confocal immunofluorescence microscopy. For each condition, a Z-
stack of 25 slices is shown as a maximum intensity projection. (D) HeLa-ACE2 were treated
with 20 µM Rap, Eve, Tem, Rid, or an equivalent volume of DMSO in the presence or absence
of 100 nM SAR405 for 4 hours and whole cell lysates were subjected to SDS-PAGE and
Western blot analysis. Immunoblotting was performed with anti-IFITM2/3 and anti-actin on the
same nitrocellulose membrane. (E) HeLa-ACE2 were treated with 20 µM Rap, Eve, Tem, Rid, or
an equivalent volume of DMSO in the presence of 1 µM Bafilomycin A1, 5 µg/mL U18666A, or
neither, for 4 hours. Cells were then fixed, permeabilized, and stained with anti-IFITM2/3.
IFITM2/3 protein levels were measured using flow cytometry. Means and standard error were
calculated from 3 experiments. Statistical analysis was performed with one-way ANOVA and
asterisks indicate significant difference from DMSO. *, $p < 0.05$; **, $p < 0.01$. Rel.; relative. All
immunoblots are representative of three independent experiments.

Supplemental Figure 3: (A) HeLa-ACE2 cells were transfected with 0.3 µg pcDNA3.1-hACE2
for 24 hours and treated with 20 µM Rap, Eve, Tem, Rid, or the equivalent volume of DMSO for
4 hours and whole cell lysates were subjected to SDS-PAGE and Western blot analysis. Cells
were fixed, permeabilized, stained with anti-ACE2, and imaged by confocal
immunofluorescence microscopy. Images represent a single, medial Z section. (B) HeLa-ACE2
cells were treated with 20 µM Rap, Eve, Tem, Rid, or the equivalent volume of DMSO for 4
hours and subsequently fixed and stained with LIVE/DEAD Fixable Red Dead Cell Stain Kit for
30 minutes according to manufacturer's instructions. Cells were analyzed by flow cytometry.
Means and standard error were calculated from 2 experiments. Statistical analysis was performed
with one-way ANOVA and asterisks indicate significant difference from DMSO. *, $p < 0.05$; **,
$p < 0.01$. Rel.; relative.

Supplemental Figure 4: Calu-3 cells were treated with 20 µM Rap, Eve, Tem, Rid, or the
equivalent volume of DMSO for 4 hours. HIV-CoV-2 (100 ng p24 equivalent) was added to cells
and infection was measured by luciferase activity at 48 hours post-infection. Luciferase units
were normalized to 100 in the DMSO condition. Means and standard error were calculated from
3 experiments. Statistical analysis was performed with one-way ANOVA and asterisks indicate
significant difference from DMSO. *, $p < 0.05$; **, $p < 0.01$. Rel.; relative.

Supplemental Figure 5: (A) HeLa-ACE2 cells were treated with 20 µM Rap, Eve, Tem, Rid,
Tac, or the equivalent volume of DMSO for 4 hours. Whole cell lysates were subjected to SDS-
PAGE and Western blot analysis. Immunoblotting was performed with anti-IFITM2/3 and anti-
actin on the same nitrocellulose membrane. (B) HeLa-ACE2 cells were treated with 20 µM Rap,
Eve, Tem, Rid, Tac, or the equivalent volume of DMSO for 4 hours. HIV-CoV-2 (100 ng p24
equivalent) was added to cells and infection was measured by luciferase activity at 48 hours
post-infection. Luciferase units were normalized to 100 in the DMSO condition. Means and
standard error were calculated from 3 experiments. Statistical analysis was performed with one-
way ANOVA and asterisks indicate significant difference from DMSO. *, $p < 0.05$; **, $p < 0.01$.
Rel.; relative. (C) HeLa-ACE2 were transfected with 0.5 µg TFEBΔ30-GFP for 24 hours and

treated with 20 µM Tem for four hours. Cells were then fixed, permeabilized, stained with anti-IFITM2/3, and imaged by confocal immunofluorescence microscopy. Representative images are shown and anti-IFITM2/3 staining in untreated HeLa-ACE2 are shown for comparison. (D) Whole cell lysates from HeLa WT and HeLa *TFEB* KO cells were subjected to SDS-PAGE and Western blot analysis. Immunoblotting was performed with anti-TFEB and anti-actin on the same nitrocellulose membrane.

Supplemental Figure 6: Body weight measurements for individual hamsters following injections with DMSO (A), Rap (B), or Rid (C) are plotted by day post-infection and presented as % body weight change relative to Day 0. Hamsters receiving one injection of 3 mg/kg DMSO, Rap, or Rid prior to infection (n=4, 1 injection) are indicated by black squares, while hamsters receiving one injection prior to infection as well as a second injection of 3 mg/kg DMSO, Rap, or Rid at Day 2 post-infection (n=4, 2 injections) are indicated by white squares. The average daily weight change for each group is indicated by grey and black lines, respectively. If and when a hamster lost 20% or more of its body weight, it was euthanized and body weight measurements were stopped.

Acknowledgements

We thank Michael Letko and Vincent Munster for providing VSV-luc/GFP seed particles and CoV and CoV-2 Spike plasmids, Thomas Gallagher for providing CoV, CoV-2, and MERS Spike plasmids, Alan Rein for facilitating lentiviral pseudovirus production, Scott H. Randell for providing UNCNN2TS, Ramnik J. Xavier for providing HeLa *TFEB* KO cells, and Eric O. Freed for providing primary HSAEC.

Funding sources

Work in the lab of AAC was funded by the Intramural Research Program, National Institutes of Health, National Cancer Institute, Center for Cancer Research and an Intramural Targeted Anti-COVID-19 award from the National Institute of Allergy and Infectious Diseases. Work in the lab of SMB was funded by the Division of Intramural Research, National Institutes of Health, National Institute of Allergy and Infectious Diseases. The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. Government. Work in the lab of JSY was funded by National Institutes of Health grants AI130110, AI151230, AI142256, and HL154001.

References

1. Zhu, N., et al., *A Novel Coronavirus from Patients with Pneumonia in China, 2019.* New England Journal of Medicine, 2020. **382**(8): p. 727-733.
2. Wu, F., et al., *A new coronavirus associated with human respiratory disease in China.* Nature, 2020: p. 1-20.
3. Zhang, X., et al., *Viral and host factors related to the clinical outcome of COVID-19.* Nature, 2020: p. 1-18.

4.   Aydillo, T., N.E. Babady, and M. Kamboj, *Shedding of Viable SARS-CoV-2 after*
Immunosuppressive Therapy for Cancer. New England Journal of Medicine, 2020.
383(26): p. 2586-2588.
5.   Wei, J., et al., *SARS-CoV-2 infection in immunocompromised patients: humoral versus*
cell-mediated immunity. Journal for immunotherapy of cancer, 2020. **8**(2): p. e000862.
6.   Tarhini, H., et al., *Long term SARS-CoV-2 infectiousness among three*
immunocompromised patients: from prolonged viral shedding to SARS-CoV-2
superinfection. The Journal of Infectious Diseases, 2021.
7.   Baang, J.H., et al., *Prolonged Severe Acute Respiratory Syndrome Coronavirus 2*
Replication in an Immunocompromised Patient. The Journal of Infectious Diseases,
2021. **223**(1): p. 23-27.
8.   Avanzato, V.A., et al., *Case Study: Prolonged Infectious SARS-CoV-2 Shedding from an*
Asymptomatic Immunocompromised Individual with Cancer. Cell, 2020. **183**(7): p. 1901-
1912.e9.
9.   Gordon, D.E., et al., *A SARS-CoV-2 protein interaction map reveals targets for drug*
repurposing. Nature, 2020: p. 1-30.
10.   Zhou, Y., et al., *Network-based drug repurposing for novel coronavirus 2019-*
nCoV/SARS-CoV-2. Cell Discovery, 2020: p. 1-18.
11.   Appelberg, S., et al., *Dysregulation in Akt/mTOR/HIF-1 signaling identified by proteo-*
transcriptomics of SARS-CoV-2 infected cells. Emerging Microbes & Infections, 2020.
9(1): p. 1748-1760.
12.   Blagosklonny, M.V., *From causes of aging to death from COVID-19.* Aging, 2020.
12(11): p. 10004-10021.
13.   Ciliberto, G., R. Mancini, and M.G. Paggi, *Drug repurposing against COVID-19: focus*
on anticancer agents. 2020: p. 1-9.
14.   Husain, A. and S.N. Byrareddy, *Rapamycin as a potential repurpose drug candidate for*
the treatment of COVID-19. Chemico-Biological Interactions, 2020. **331**: p. 109282.
15.   Zheng, Y., R. Li, and S. Liu, *Immunoregulation with mTOR inhibitors to prevent*
COVID-19 severity: A novel intervention strategy beyond vaccines and specific antiviral
medicines. Journal of medical virology, 2020. **92**(9): p. 1495-1500.
16.   Terrazzano, G., et al., *An Open Question: Is It Rational to Inhibit the mTor-Dependent*
Pathway as COVID-19 Therapy? 2020: p. 1-5.
17.   Ramaiah, M.J., *mTOR inhibition and p53 activation, microRNAs: The possible therapy*
against pandemic COVID-19. Gene reports, 2020. **20**: p. 100765.
18.   Zhavoronkov, A., *Geroprotective and senoremediative strategies to reduce the*
comorbidity, infection rates, severity, and lethality in gerophilic and gerolavic infections.
Aging, 2020. **12**(8): p. 6492-6510.
19.   Willyard, C., *Ageing and Covid Vaccines.* Nature, 2020. **586**(7829): p. 352-354.
20.   Omarjee, L., et al., *Targeting T-cell senescence and cytokine storm with rapamycin to*
prevent severe progression in COVID-19. Clinical immunology (Orlando, Fla.), 2020.
216: p. 108464.
21.   Laplante, M. and D.M. Sabatini, *mTOR signaling in growth control and disease.* Cell,
2012. **149**(2): p. 274-293.
22.   Bertram, P.G., et al., *The 14-3-3 proteins positively regulate rapamycin-sensitive*
signaling. Current biology : CB, 1998. **8**(23): p. 1259-1267.

23. Vignot, S., et al., *mTOR-targeted therapy of cancer with rapamycin derivatives.* Annals
of Oncology, 2005. **16**(4): p. 525-537.
24. Lamming, D.W., et al., *Rapalogs and mTOR inhibitors as anti-aging therapeutics.*
Journal of Clinical Investigation, 2013. **123**(3): p. 980-989.
25. Abdel-Magid, A.F., *Rapalogs Potential as Practical Alternatives to Rapamycin.* ACS
medicinal chemistry letters, 2019. **10**(6): p. 843-845.
26. Benjamin, D., et al., *Rapamycin passes the torch: a new generation of mTOR inhibitors.*
Nature Reviews Drug Discovery, 2011: p. 1-13.
27. Marz, A.M., et al., *Large FK506-Binding Proteins Shape the Pharmacology of*
Rapamycin. Molecular and Cellular Biology, 2013. **33**(7): p. 1357-1367.
28. Zoncu, R., A. Efeyan, and D.M. Sabatini, *mTOR: from growth signal integration to*
cancer, diabetes and ageing. Nature Reviews in Molecular Cell Biology, 2011. **12**(1): p.
21-35.
29. Chi, H., *Regulation and function of mTOR signalling in T cell fate decisions.* Nature
Reviews Immunology, 2012: p. 1-14.
30. Mangalmurti, N. and C.A. Hunter, *Cytokine Storms: Understanding COVID-19.*
Immunity, 2020. **53**(1): p. 19-25.
31. Mehta, P., et al., *COVID-19: consider cytokine storm syndromes and*
immunosuppression. Lancet (London, England), 2020. **395**(10229): p. 1033-1034.
32. Manjili, R.H., et al., *COVID-19 as an Acute Inflammatory Disease.* The Journal of
Immunology, 2020. **205**(1): p. 12-19.
33. Bischof, E., et al., *The potential of rapalogs to enhance resilience against SARS-CoV-2*
infection and reduce the severity of COVID-19. The Lancet. Healthy longevity, 2021.
2(2): p. e105-e111.
34. Mullen, P.J., et al., *SARS-CoV-2 infection rewires host cell metabolism and is potentially*
susceptible to mTORC1 inhibition. Nature communications, 2021: p. 1-10.
35. Alsuwaidi, A.R., et al., *Sirolimus alters lung pathology and viral load following influenza*
A virus infection. Respiratory research, 2017. **18**(1): p. 136.
36. Huang, C.-T., et al., *Rapamycin adjuvant and exacerbation of severe influenza in an*
experimental mouse model. Scientific Reports, 2017. **7**(1): p. 873-8.
37. Shi, G., et al., *mTOR inhibitors lower an intrinsic barrier to virus infection mediated by*
IFITM3. Proceedings of the National Academy of Sciences of the United States of
America, 2018. **115**(43): p. E10069-E10078.
38. Ozog, S., et al., *Resveratrol trimer enhances gene delivery to hematopoietic stem cells by*
reducing antiviral restriction at endosomes. Blood, 2019. **134**(16): p. 1298-1311.
39. Shi, G., O. Schwartz, and A.A. Compton, *More than meets the I: the diverse antiviral*
and cellular functions of interferon-induced transmembrane proteins. Retrovirology,
2017. **14**(1): p. 1-11.
40. Majdoul, S. and A.A. Compton, *Lessons in self-defence: inhibition of virus entry by*
intrinsic immunity. Nature Reviews Immunology, 2021.
41. Bestle, D., et al., *TMPRSS2 and furin are both essential for proteolytic activation of*
SARS-CoV-2 in human airway cells. Life science alliance, 2020. **3**(9): p. e202000786.
42. Hoffmann, M., H. Kleine-Weber, and S. Pöhlmann, *A Multibasic Cleavage Site in the*
Spike Protein of SARS-CoV-2 Is Essential for Infection of Human Lung Cells. Molecular
Cell, 2020. **78**(4): p. 779-784.e5.

43. Hoffmann, M., et al., *SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor.* Cell, 2020: p. 1-19.

44. Shi, G., et al., *mTOR inhibitors lower an intrinsic barrier to virus infection mediated by IFITM3.* Proceedings of the National Academy of Sciences of the United States of America, 2018. **115**(43): p. E10069-E10078.

45. Mejlvang, J., et al., *Starvation induces rapid degradation of selective autophagy receptors by endosomal microautophagy.* J Cell Biol, 2018. **217**(10): p. 3640-3655.

46. Olsvik, H.L., et al., *Endosomal microautophagy is an integrated part of the autophagic response to amino acid starvation.* Autophagy, 2019. **15**(1): p. 182-183.

47. Sahu, R., et al., *Microautophagy of cytosolic proteins by late endosomes.* Dev Cell, 2011. **20**(1): p. 131-9.

48. Bozzo, C.P., et al., *IFITM proteins promote SARS-CoV-2 infection of human lung cells.* bioRxiv, 2020. **1**: p. 261-47.

49. Kolos, J.M., et al., *FKBP Ligands-Where We Are and Where to Go?* Frontiers in pharmacology, 2018. **9**: p. 1425.

50. Roczniak-Ferguson, A., et al., *The transcription factor TFEB links mTORC1 signaling to transcriptional control of lysosome homeostasis.* Science signaling, 2012. **5**(228): p. ra42.

51. Martina, J.A., et al., *MTORC1 functions as a transcriptional regulator of autophagy by preventing nuclear transport of TFEB.* Autophagy, 2012. **8**(6): p. 903-14.

52. Settembre, C., et al., *A lysosome-to-nucleus signalling mechanism senses and regulates the lysosome via mTOR and TFEB.* The EMBO journal, 2012. **31**(5): p. 1095-1108.

53. Zhang, X., et al., *Rapamycin directly activates lysosomal mucolipin TRP channels independent of mTOR.* PLoS Biology, 2019. **17**(5): p. e3000252-24.

54. Alsuwaidi, A.R., et al., *Sirolimus alters lung pathology and viral load following influenza A virus infection.* Respiratory research, 2017. **18**(1): p. 1-8.

55. Tostanoski, L.H., et al., *Ad26 vaccine protects against SARS-CoV-2 severe clinical disease in hamsters.* Nat Med, 2020. **26**(11): p. 1694-1700.

56. Sasongko, T.H., et al., *Rapamycin and its analogues (rapalogs) for Tuberous Sclerosis Complex-associated tumors: a systematic review on non-randomized studies using meta-analysis.* Orphanet Journal of Rare Diseases, 2015: p. 1-11.

57. Reagan-Shaw, S., M. Nihal, and N. Ahmad, *Dose translation from animal to human studies revisited.* FASEB J, 2008. **22**(3): p. 659-61.

58. Cloughesy, T.F., et al., *Antitumor activity of rapamycin in a Phase I trial for patients with recurrent PTEN-deficient glioblastoma.* PLoS Med, 2008. **5**(1): p. e8.

59. Sun, Y., et al., *Phase II study of the safety and efficacy of temsirolimus in East Asian patients with advanced renal cell carcinoma.* Jpn J Clin Oncol, 2012. **42**(9): p. 836-44.

60. Martynov, M.Y., V.A. Kutashov, and O.V. Ulyanova, *COVID 19 in a family with rare genetic disease of the nervous system.* Neurology, Neuropsychiatry, Psychosomatics, 2022. **14**(1): p. 108-114.

61. Medina, D.L., et al., *Lysosomal calcium signalling regulates autophagy through calcineurin and TFEB.* Nature Cell Biology, 2015. **17**(3): p. 288-299.

62. Schuck, S., *Microautophagy - distinct molecular mechanisms handle cargoes of many sizes.* Journal of Cell Science, 2020. **133**(17).

63. Oku, M. and Y. Sakai, *Three Distinct Types of Microautophagy Based on Membrane Dynamics and Molecular Machineries.* BioEssays, 2018. **40**(6): p. 1800008-6.

1048   64.   Santambrogio, L. and A.M. Cuervo, *Chasing the elusive mammalian microautophagy.*
1049   Autophagy, 2011. **7**(6): p. 652-4.

1050   65.   Tekirdag, K. and A.M. Cuervo, *Chaperone-mediated autophagy and endosomal*
1051   *microautophagy: Jointed by a chaperone.* Journal of Biological Chemistry, 2018.
1052   **293**(15): p. 5414-5424.

1053   66.   Mijaljica, D., M. Prescott, and R.J. Devenish, *Microautophagy in mammalian cells:*
1054   *revisiting a 40-year-old conundrum.* Autophagy, 2011. **7**(7): p. 673-682.

1055   67.   Sato, M., et al., *Rapamycin activates mammalian microautophagy.* Journal of
1056   Pharmacological Science, 2019. **140**(2): p. 201-204.

1057   68.   Yang, X., et al., *TORC1 regulates vacuole membrane composition through ubiquitin- and*
1058   *ESCRT-dependent microautophagy.* The Journal of Cell Biology, 2020. **219**(3).

1059   69.   Li, M., et al., *Ubiquitin-Dependent Lysosomal Membrane Protein Sorting and*
1060   *Degradation.* Molecular Cell, 2015. **57**(3): p. 467-478.

1061   70.   Chesarino, N.M., T.M. McMichael, and J.S. Yount, *E3 Ubiquitin Ligase NEDD4*
1062   *Promotes Influenza Virus Infection by Decreasing Levels of the Antiviral Protein*
1063   *IFITM3.* PLoS Pathogens, 2015. **11**(8): p. e1005095.

1064   71.   Compton, A.A., et al., *Natural mutations in IFITM3 modulate post-translational*
1065   *regulation and toggle antiviral specificity.* EMBO reports, 2016. **17**(11): p. 1657-1671.

1066   72.   Yang, X., et al., *TORC1 regulates vacuole membrane composition through ubiquitin- and*
1067   *ESCRT-dependent microautophagy.* 2019. **8**: p. 1835-53.

1068   73.   Di Fruscio, G., et al., *Lysoplex: An efficient toolkit to detect DNA sequence variations in*
1069   *the autophagy-lysosomal pathway.* Autophagy, 2015. **11**(6): p. 928-38.

1070   74.   Zhao, X., et al., *Interferon induction of IFITM proteins promotes infection by human*
1071   *coronavirus OC43.* Proceedings of the National Academy of Sciences, 2014. **111**(18): p.
1072   6756-6761.

1073   75.   Shi, G., et al., *Opposing activities of IFITM proteins in SARS☐CoV☐2 infection.* The
1074   EMBO journal, 2020. **3**: p. e201900542-12.

1075   76.   Winstone, H., et al., *The polybasic cleavage site in the SARS-CoV-2 spike modulates viral*
1076   *sensitivity to Type I interferon and IFITM2.* Journal of Virology, 2021.

1077   77.   Zhao, X., et al., *LY6E Restricts Entry of Human Coronaviruses, Including Currently*
1078   *Pandemic SARS-CoV-2.* Journal of Virology, 2020. **94**(18).

1079   78.   Zang, R., et al., *Cholesterol 25-hydroxylase suppresses SARS-CoV-2 replication by*
1080   *blocking membrane fusion.* Proceedings of the National Academy of Sciences of the
1081   United States of America, 2020. **32**: p. 202012197.

1082   79.   Huang, I.C., et al., *Distinct patterns of IFITM-mediated restriction of filoviruses, SARS*
1083   *coronavirus, and influenza A virus.* PLoS Pathogens, 2011. **7**(1): p. e1001258.

1084   80.   Zhao, X., et al., *Identification of Residues Controlling Restriction versus Enhancing*
1085   *Activities of IFITM Proteins on Entry of Human Coronaviruses.* Journal of Virology,
1086   2018. **92**(6): p. 374-17.

1087   81.   Zani, A., et al., *Interferon-induced transmembrane protein 3 (IFITM3) limits lethality of*
1088   *SARS-CoV-2 in mice.* bioRxiv, 2021.

1089   82.   Beyer, D.K. and A. Forero, *Mechanisms of Antiviral Immune Evasion of SARS-CoV-2.* J
1090   Mol Biol, 2021: p. 167265.

1091   83.   Hachim, M.Y., et al., *Interferon-Induced Transmembrane Protein (IFITM3) Is*
1092   *Upregulated Explicitly in SARS-CoV-2 Infected Lung Epithelial Cells.* Frontiers in
1093   Immunology, 2020. **11**: p. 1372.

84.   Blanco-Melo, D., et al., *Imbalanced Host Response to SARS-CoV-2 Drives Development*
of COVID-19. Cell, 2020. **181**(5): p. 1036-1045.e9.
85.   Ziegler, C.G.K., et al., *Impaired local intrinsic immunity to SARS-CoV-2 infection in*
severe COVID-19. bioRxiv, 2021: p. 2021.02.20.431155.
86.   Zhang, Y., et al., *Interferon-induced transmembrane protein-3 genetic variant rs12252-*
C is associated with disease severity in COVID-19. The Journal of Infectious Diseases,
2020.
87.   Nikoloudis, D., D. Kountouras, and A. Hiona, *The frequency of combined IFITM3*
haplotype involving the reference alleles of both rs12252 and rs34481144 is in line with
COVID-19 standardized mortality ratio of ethnic groups in England. PeerJ, 2020. **8**: p.
e10402.
88.   Carey, K.L., et al., *TFEB Transcriptional Responses Reveal Negative Feedback by*
BHLHE40 and BHLHE41. Cell Rep, 2020. **33**(6): p. 108371.
89.   Leist, S.R., et al., *A Mouse-Adapted SARS-CoV-2 Induces Acute Lung Injury and*
Mortality in Standard Laboratory Mice. Cell, 2020. **183**(4): p. 1070-1085 e12.
90.   Chiramel, A.I., et al., *TRIM5α Restricts Flavivirus Replication by Targeting the Viral*
Protease for Proteasomal Degradation. CellReports, 2019. **27**(11): p. 3269-3283.e6.
91.   Letko, M., A. Marzi, and V. Munster, *Functional assessment of cell entry and receptor*
usage for SARS-CoV-2 and other lineage B betacoronaviruses. Nature Microbiology,
2020. **11**: p. 1-17.
92.   Compton, A.A., et al., *IFITM Proteins Incorporated into HIV-1 Virions Impair Viral*
Fusion and Spread. Cell Host & Microbe, 2014. **16**(6): p. 736-747.
1116

Figure 1

Figure 2

Figure 3

A SARS-CoV-2
   HeLa-ACE2

B SARS-CoV-2
   HeLa-ACE2

C HIV-CoV-2 Spike
   HeLa-ACE2

D HeLa-ACE2 – 4 hr

E

F

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

A

Rapalog

MA SARS-CoV-2
+
Rapalog

1 d

2 d

Infectious
virus titer

Lung harvest

B

n = 6   n = 7

Lung TCID$_{50}$ per gram

10^{11}
10^{10}
10^{9}
10^{8}
10^{7}
10^{6}
10^{5}
10^{4}
10^{3}

**

D   Rap

C

n = 6   n = 7
DMSO   Rapamycin

15 —   IFITM3

50 —
37 —   Actin

Figure 9