# Potential transmission chains of variant B.1.1.7 and co-mutations of SARS-CoV-2

Jingsong Zhang[1#], Yang Zhang[2#], Junyan Kang[3,4], Shuiye Chen[2], Yongqun He[5], Benhao Han[1], Mofang Liu[3,4], Lina Lu[1], Li Li[6], Zhigang Yi[2,7*] and Luonan Chen[1,8,9,10*]

[1]Key Laboratory of Systems Biology, State Key Laboratory of Cell Biology, Shanghai Institute of Biochemistry and Cell Biology, Center for Excellence in Molecular Cell Science, Chinese Academy of Sciences, Shanghai 200031, China.

[2]Key Laboratory of Medical Molecular Virology (MOE/NHC/CAMS), School of Basic Medical Sciences, Shanghai Medical College, Fudan University, Shanghai 200032, China.

[3]State Key Laboratory of Molecular Biology, Shanghai Key Laboratory of Molecular Andrology, Shanghai Institute of Biochemistry and Cell Biology, Center for Excellence in Molecular Cell Science, Chinese Academy of Sciences, Shanghai 200031, China.

[4]University of Chinese Academy of Sciences, Shanghai 200031, China.

[5]Department of Computational Medicine and Bioinformatics, University of Michigan Medical School, Ann Arbor, MI 48109, USA.

[6]Department of Genetics, Harvard Medical School, Boston, MA 02115, USA.

[7]Shanghai Public Health Clinical Center, Fudan University, Shanghai 201508, China.

[8]School of Life Science and Technology, ShanghaiTech University, Shanghai 201210, China.

[9]Key Laboratory of Systems Biology, Hangzhou Institute for Advanced Study, University of Chinese Academy of Sciences, Hangzhou 310024, China.

[10]Center for Excellence in Animal Evolution and Genetics, Chinese Academy of Sciences, Kunming 650223, China.

*Correspondence: Luonan Chen (lnchen@sibs.ac.cn) or Zhigang Yi (zgyi@fudan.edu.cn)
#These authors contributed equally: Jingsong Zhang, Yang Zhang

# Abstract

The presence of SARS-CoV-2 mutants, including the emerging variant B.1.1.7, has raised great concerns in terms of pathogenesis, transmission, and immune escape. Characterizing SARS-CoV-2 mutations, evolution, and effects on infectivity and pathogenicity is crucial to the design of antibody therapies and surveillance strategies. Here we analyzed 454,443 SARS-CoV-2 spike genes/proteins and 14,427 whole-genome sequences. We demonstrated that the early variant B.1.1.7 may not have evolved spontaneously in the United Kingdom or within human populations. Our extensive analyses suggested that Canidae, Mustelidae or Felidae, especially the Canidae family (for example, dog) could be a possible host of the direct progenitor of variant B.1.1.7. An alternative hypothesis is that the variant was simply yet to be sampled. Notably, the SARS-CoV-2 whole genome represents a large number of potential co-mutations with very strong statistical significances (p value<E–44). In addition, we used an experimental SARS-CoV-2 reporter replicon system to introduce the dominant co-mutations NSP12_c14408t, 5'UTR_c241t, and NSP3_c3037t into the viral genome, and to monitor the effect of the mutations on viral replication. Our experimental results demonstrated that the co-mutations significantly enhanced the viral replication. The study provides valuable clues for discovering the transmission chains of variant B.1.1.7 and understanding the evolutionary process of SARS-CoV-2.

**Key words:** SARS-CoV-2, variant B.1.1.7, transmission chains, co-mutations, viral replication.

## Introduction

Since the outbreak in December 2019, COVID-19 has been pandemic in over 200 countries. Cases of infection and mortalities have been surging and are an ongoing threat to public health[1,2]. COVID-19 is caused by infection with the novel coronavirus SARS-CoV-2[3-5]. Although as a coronavirus, SARS-CoV-2 has genetic proofreading mechanisms[6-8], the persistent natural selection pressure in the population drives the virus to gradually accumulate favorable mutations[6,9,10]. Much attention has been paid to the mutations and evolution of SARS-CoV-2[11-15], since mutations are related to the infectivity and pathogenicity of viruses[16-21]. Beneficial mutants of the virus can better evolve and adapt to the host[9], either strengthening or weakening the infectivity and pathogenicity. In addition, certain variants may generate drug resistance and reduce the efficacy of vaccines and therapeutics[22-26]. In short, studying mutations and evolution in detail is vital to understand the transformations of viral properties and to control the pandemic.

A new variant of SARS-CoV-2 named VOC-202012/01 (Variant of Concern 202012/01) or lineage B.1.1.7 was first detected in the United Kingdom last December[27]. It appears to be substantially more transmissible than other variants[28]. The variant has been growing exponentially in the United Kingdom and rapidly spreading to other countries[29,30]. However, it is not yet clear if it evolved spontaneously in the United Kingdom or was imported from other countries. Studying how the variant B.1.1.7 mutates can enable researchers to track its spread over time and to understand the evolution of SARS-CoV-2.

In this study, large-scale SARS-CoV-2 sequences, consisting of more than 454,000 spike

genes/proteins and 14,000 whole-genome sequences were analyzed. Our extensive sequence analysis showed that many mutations always co-occur not only in the spike protein of B.1.1.7, but in the whole genome of SARS-CoV-2. The mutation trajectories of the spike protein indicate that the early variant B.1.1.7 did not evolve spontaneously in the United Kingdom or even within human populations. We also investigated possible SARS-CoV-2 transmission chains of the variant B.1.1.7 based on the mutation analysis of large-scale spike proteins and the cluster analysis of spike genes. Over the whole genome, the top 25 high-frequency mutations of SARS-CoV-2 converged into several potential co-mutation patterns, each of which showed a strong correlation with a very strong statistical significance (p value<E–44). The potential co-mutations depicted the evolutionary trajectory of SARS-CoV-2 virus in the population, shaping variable replication of SARS-CoV-2. In addition, we further explored the effect of the dominant (co-)mutations 5'UTR_c241t, NSP3_c3037t, and NSP12_c14408t on viral replication using a SARS-CoV-2 replicon based on a four plasmid *in-vitro* ligation system. The results suggest that such mutations significantly attenuate the replication of SARS-CoV-2.

## Results

### Evolutionary trajectories of variant B.1.1.7

The variant B.1.1.7 was generally defined by multiple amino acid changes including 3 deletions (69-70del and 145del) and 7 mutations (N501Y, A570D, D614G, P681H, T716I, S982A, and D1118H) in the spike protein[31]. The number of non-adjacent co-occurrent changes indicates that they resulted from accumulated mutations. We therefore explored the

evolutionary trajectories of B.1.1.7 by tracing the incremental mutations (Fig. 1a). All routes

along the directions of the arrows are possible evolutionary trajectories of lineage B.1.1.7.

Among all the mutation routes, the green one was the most probable mutation trajectory

based on the number of variant strains. However, it was unlikely that the earliest variant

B.1.1.7 (GISAID: EPI_ISL_601443, 2020-09-20, England) with 9 mutations evolved from the

existing variants with 3–8 mutations, because the former arose much earlier than the latter.

More than 454,000 SARS-CoV-2 strains have been collected and extensive sequenced from

infected humans without finding intermediate variants with 3–9 mutations. It is therefore

unlikely that the intermediate variants with 3–8 mutations have infected humans. Thus, the

early variant B.1.1.7 might not have arisen spontaneously in the UK or within human

populations. An alternative hypothesis is that spillover likely occurred from susceptible

animals.

The co-appearance rates (see Materials and Methods) of all nine mutations are shown in

Fig. 1b. We found that at least five mutations (145del, A570D, T716I, S982A, and D1118H)

of variant B.1.1.7 significantly co-occurred (rate>95%), which indicates a potential

co-mutation pattern in the spike protein, causing us to wonder what selection pressure drove

such co-occurrences of mutations and rapid evolution in the population of SARS-CoV-2.

Note that coronaviruses generally tend to exhibit rapid evolution when they jump to a

different species[32]. We therefore analyzed the key spike genes and proteins of existing

SARS-CoV-2 strains collected from animals to find a possible direct progenitor of variant

B.1.1.7. The variant with mutations "56" (labeled by "*" in Fig. 1a, termed star variant) had

the minimum phylogenetic distance with EPI_ISL_699508, which was collected from a dog

on 2020-07-28 (Fig. 2) using MEGA[33,34] (see Materials and Methods). The strains collected

from tigers, minks, and cats were also close to the star variant. Our extensive analyses

including mutations, phylogeny (Fig. 2), collection date/location and the number of

sequences (Tables S1-S3) suggested that Canidae, Mustelidae or Felidae, especially the

Canidae family (for example, dog) could be a possible host of the direct progenitor of variant

B.1.1.7. The possible transmission chains of variant B.1.1.7 are shown in Fig. 1c. This star

variant strains in humans could not have evolved into the early variant B.1.1.7, but they

might have infected high-density yet susceptible animals (such as dogs) and adapted to

these species through rapid mutation. Such progenitor variants comprised most or all of the

mutations of the early variant B.1.1.7 within the Canidae family populations, and they may

have spilled back to humans after the rapid mutation period.

## High-frequency mutations converge into potential co-mutations

Based on sequence alignment and mutation analysis, we found that 7,441 nucleotide

alterations in the viral 29903-letter RNA code occurred at least once in the samples from

COVID-19 patients. These mutations were dispersed in the 14,427 SARS-CoV-2 strains

collected from all around the world. As shown in the heatmap of the top 1% high-frequency

mutations (Table S4), some sites show very similar mutation rates on most days in samples

isolated globally (Fig. S1), including 8,898 and 815 samples isolated from the U.S. (Fig. S2)

and Australia (Fig. S3). Therefore, these mutations shown in Fig. S4a were selected and

clustered into co-occurrences, which we called potential co-mutation patterns. From the

landscape of the mutation rates (Fig. S4a), 25 nucleotide sites were clearly clustered into

several potential co-mutation patterns. Among these patterns, there was one consisting of the top 4 high-frequency mutations (i.e., 5'UTR_c241t, NSP3_c3037t, NSP12_c14408t, and S_a23403g), which converged into a dominant potential co-mutation pattern. Such co-occurrence lineage has been found in almost all sequenced samples of SARS-CoV-2. Within this co-occurrence pattern, mutation S_ a23403g resulted in the amino acid change (D614G) that apparently enhances viral infectivity[6,35], albeit debate exists[16]. Notably, there were three successive sites at the 28881st to 28883rd positions of the virus (N_g28881a, N_g28882a, and N_g28883c) that strictly co-occurred. Comparing Fig. S4a-c and Table S4, we found that the top 14 high-frequency mutations formed five common co-occurrence patterns.

To assess the above co-occurrence patterns, we analyzed the correlations and statistical significance levels of the high-frequency co-occurrence mutations. The heatmap of the paired Pearson-correlation-coefficients (Fig. 3a) shows that the top 25 high-frequency mutations clearly cluster into several potential co-mutation groups/patterns with very strong correlation ($\geq 0.8$). By regression analyses, the above co-occurrence patterns have statistical significance levels with p values less than $10^{-44}$ (Fig. 3b). The detailed mutation transitions (Fig. 3c–k, Figs. S5–7) provide further evidence that the above mutations form co-mutation patterns.

## Dominant mutations attenuate viral replication

We further explored the effect of the dominant mutations 5'UTR_c241t, NSP3_c3037t, and NSP12_c14408t on viral replication using a SARS-CoV-2 replicon based on a

four-plasmid *in-vitro* ligation system. This replicon is devoid of the viral structural proteins

while undergoing viral replication, and the viral replication is sensitive to the antiviral agent

remdesivir[36]. The 5'UTR_c241t mutation resides in a highly conserved region in the 5'UTR

(Fig. 4a). The NSP3_c3037t mutation is synonymous. The NSP12_c14408t mutation is

nonsynonymous with an amino acid change of a conserved amino acid P323 in the viral

RNA-dependent RNA polymerase (Fig. 4b). We introduced the NSP12_c14408t mutation or

the NSP12_c14408t mutation with the other two mutations 5'UTR_c241t and NSP3_c3037t

into the replicon plasmids. The fragments were released from the plasmids by BsaI digestion,

and then assembled by *in-vitro* ligation with T4 ligase (Fig. 4c). Replicon RNA transcribed

from the ligation products was co-transfected with N mRNA into Huh7 cells. RNA replication

was monitored by measuring the secreted *Gaussia* luciferase activity in the supernatants.

Enzymatic dead mutants (759-SAA-761) of the RNA-dependent RNA polymerase NSP12

were introduced, and the mutated replicon served as a non-replication control. As shown in

Fig. 4d, transfection of WT replicon RNA resulted in an obvious increase of luciferase activity,

and SAA RNA did not replicate as expected. Introduction of NSP12_c14408t mutation

resulted in a significant reduction of viral replication. The combination of NSP12_c14408t

mutation with the other two mutations further significantly reduced viral replication. These

results demonstrate that the P323L mutation in the viral RNA-dependent RNA polymerase

reduces viral replication, and the synonymous mutations further attenuate viral replication.

## Discussion

172 A well-resolved phylogeny of variant B.1.1.7 spike genes provides an opportunity to

understand the evolutionary process and transmission chains of variant B.1.1.7. Our incremental mutation and phylogenetic analyses on large-scale SARS-CoV-2 spike proteins/genes revealed that the early variant B.1.1.7 might not have evolved spontaneously in the United Kingdom or within human populations. In this case the spillover likely occurred from susceptible animals. Current evidence[37-39] indicates that SARS-CoV-2 can effectively infect both domestic animals (for example, dog, cat, pig and bovine) and wild animals (for example, mink, rabbit and fox) by binding their angiotensin converting enzyme 2 (ACE2). Our further analyses including mutations, phylogeny, collection date/location and the number of sequences suggested that the earliest variant B.1.1.7 possibly originated from Canidae, Mustelidae or Felidae, especially the Canidae family (for example, dog). The cases[40] that the variant B.1.1.7 can easily infect dogs and cats indicated that both are susceptible to B.1.1.7. Still, due to the limited information available to date, an alternative hypothesis is that the direct progenitor of variant B.1.1.7 is yet to be sampled.

By tracing the mutation trajectories, we found that at least five mutations of the spike proteins always co-occurred, and a large number of potential co-mutations appeared in the top 1% high-frequency mutations of SARS-CoV-2 whole genome. In addition, we used a SARS-CoV-2 reporter replicon system to study the dominant co-mutations (NSP12_c14408t, 5'UTR_c241t, and NSP3_c3037t) and investigated whether they affected the viral replication. We found that such co-mutations significantly enhanced the replication of SARS-CoV-2. In line with the findings that the mutation S_ a23403g results in the amino acid change of the spike protein D614G that enhances viral infectivity[19,41-44], our findings elucidate that SARS-CoV-2 undergoes an evolution toward enhancing viral infectivity while attenuating

viral replication.

So far, SARS-CoV-2 has exhibited significant mutations and co-mutations. We just
evaluated the replication of a co-mutation pattern including three dominant mutations.
Determining the functional effects of the other co-mutations is still a long-term work.
Together, this study provides insights into not only the transmission chains of variant B.1.1.7
but also our understanding of SARS-CoV-2 evolution.

## 201 Materials and Methods

### 202 Data selection and pre-processing

The 454,443 spike gene/protein sequences of SARS-CoV-2 were obtained at
https://www.gisaid.org/. The NCBI website at https://www.ncbi.nlm.nih.gov/sars-cov-2/ has
released more than 1.7 thousand sequences of SARS-CoV-2 viruses before July 31, 2020.
We selected 14,427 sequences that satisfied two criteria: (1) having specific collection dates;
(2) sequence-lengths being no less than 29,305 nt (29903*0.98). It is inevitable that some
sites of sequences are equivocal owing to the limitation of sequencing depth. For instance,
many sites were labeled as letter N in genome sequences. The noise of indeterminate
nucleic-acids was taken into consideration in our experiments so as to boost accuracy. The
co-mutation rate of multi-site co-mutations was calculated by $\mathrm{co-mutation\ rate} =$
$\frac{\text{number of sequences containing co−mutaions}}{\text{number of all sequences}}$. Moreover, the co-appearance rate of a mutation in
B.1.1.7 variant was defined by $\mathrm{co-appearance\ rate} = \frac{\text{number of B.1.1.7 sequences}}{\text{number of sequences containing a mutation}}$.

### 214 Possible animal host analyses

In addition to the phylogenetic analysis, we further explored the possible animal hosts of

the direct progenitor of variant B.1.1.7 by collection time/space of strains, the number of

sequences and the edit distance[45,46] of mutations (Table S1-2). Due the late lockdown

policies of some governmental agencies, the spread of SARS-CoV-2 has not been

prevented well in Europe, America, and Australia. We could ignore the impact of policies for

studying the origin of variant B.1.1.7. We quantified the multiple impact factors of viral

transmission as shown in Table S3 based on the criterion that the smaller the value, the

more similar. The results still supported that the Canidae family is a possible host of the

direct progenitor of variant B.1.1.7.

**MEGA version and parameter settings**

Version: MEGA-X

Statistical Method: Maximum Likelihood

Test of Phylogeny: None

Model/Method: Jones-Taylor-Thornton (JTT) model

Rates among Sites: Uniform Rates

Gaps/Missing Data Treatment: Use all sites

ML Heuristic Method: Nearest-Neighbor-Interchange (NNT)

Initial Tree for ML: Make initial tree automatically (Default - NJ/BioNJ)

Branch Swap Filter: None

Number of Threads: 7

**Statistical analysis**

The Pearson-correlation-coefficient (PCC) is a classic statistic that measures linear

correlation between two variables. Its value ranges from -1.0 to 1.0. Normally, the two

variables meet a strong correlation or a very strong correlation when the absolutes value of PCC is between 0.6 and 0.8 or between 0.8 and 1.0. Linear regression is a linear approach to model the relationship between a scalar response and one or more variables. We used PCC and significance level (p value) of regression analysis to evaluate the relationships of the co-occurrence mutations in large-scale SARS-CoV-2 examples.

**Plasmids**

Four plasmids encompassing the viral genome (pLC-nCoV-A-BsaI, pLC-nCoV-B-BsaI, pLC-nCoV-C-BsaI, and pnCoV-D-sGluc-BsaI) were described previously[36]. The 5'UTR_c-241-t and NSP3_c-3037-t mutations were introduced into the pLC-nCoV-A-BsaI by fusion PCR. The NSP12_c-14408-t mutation was introduced into the pLC-nCoV-B-BsaI by fusion PCR.

**Cell lines**

The human hepatoma cells Huh 7 were purchased from the Cell Bank of the Chinese Academy of Sciences (www.cellbank.org.cn) and routinely maintained in Dulbecco's modified medium supplemented with 10% FBS (Gibco) and 25 mM HEPES (Gibco).

***In-vitro* ligation**

BsaI digested fragments were gel purified using Gel Extraction Kit (OMEGA) and ligated with T4 ligase (New England Biolabs) at room temperature for 1 h. The ligation products were phenol/chloroform extracted, precipitated by absolute ethanol, and resuspended in

257   nuclease-free water, quantified by determining the A260 absorbance.

**_In-vitro_ transcription**

Purified _in-vitro_ ligated product was used as template for the _in-vitro_ transcription by

mMESSAGE mMACHINE T7 Transcription Kit (Ambion) according to the manufacturer's

protocol. For N mRNA production, we amplified the N coding region by PCR (sense: _GGC_

_ACA CCC CTT TGG CTC T_; antisense: _TTT TTT TTT TTT TTT TTT TTT TTT TTT TTT TTT_

_TTT TTT TCT AGG CCT GAG TTG AGT CAG CAC_) with phCMV-N as template. Then the

purified PCR product was used as a template for _in-vitro_ transcription by mMESSAGE

mMACHINE T7 Transcription Kit as described above. RNA was purified by RNeasy mini

Elute (Qiagen), eluted in nuclease-free water, and quantified by UV absorbance (260 nm).

**Transfection**

268   Cells were seeded onto 48-well plates at a density of $7.5 \times 10^4$ per well and then

269   transfected with 0.3 µg _in-vitro_ transcribed RNA using a TransIT-mRNA transfection kit

270   (Mirus) according to the manufacturer's protocol.

**Luciferase activity**

272   Supernatants were taken from cell medium and mixed with equal volumes of 2×lysis

273   buffer (Promega). Luciferase activity was measured with Renilla luciferase substrate

274   (Promega) according to the manufacturer's protocol.

## Acknowledgments

We thank associate prof. T.Z. and Dr. S.T.H. for useful comments on the manuscript. This work is supported by the National Key Research and Development Program of China (2017YFA0505500 to L.N.C., 2017YFC0909502 to J.S.Z.); the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB38040400 to L.N.C.); National Science Foundation of China (31771476 and 31930022 to L.N.C, 61602460 and 11701379 to J.S.Z.); Shanghai Municipal Science and Technology Major Project (2017SHZDZX01 to L.N.C.); National Science and Technology Major Project of China (2017ZX10103009 to Z.G.Y.); Emergency Project of Shanghai Science and Technology Committee (20411950103 to Z.G.Y.); National Postdoctoral Program for Innovative Talent (BX20180331 to J.Y.K.); and China Postdoctoral Science Foundation (2018M642018 to J.Y.K.).

## Author contributions

L.N.C. and J.S.Z. designed the study. Z.G.Y. and J.S.Z. designed the experiments. J.S.Z. analyzed data. Y. Z. performed the experiments of viral replication. J.S.Z., Z.G.Y., and J.Y.K. designed the figures. S.C. repeated and checked the experiments of viral replication. H.B.H. checked the computational analyses. J.S.Z. and Z.G.Y. wrote the manuscript. Y.Q.H, M.F.L., L.N.L, and L.L. polished the manuscript. All authors participated in result interpretation and discussion.

## Data availability

The raw sequence data reported in this paper have been deposited in the GISAID and NCBI

295    websites at https://www.gisaid.org/ and https://www.ncbi.nlm.nih.gov/sars-cov-2/,

296    respectively. Code is available from the corresponding author on reasonable request.

## Conflict of interest

298    The authors declare that they have no conflict of interest.

## References

1    Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* **367**,
1260-1263 (2020).

2    Chinazzi, M. *et al.* The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19)
outbreak. *Science*, 395-400 (2020).

3    Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *The New England journal of*
*medicine* **382**, 727-733, doi:10.1056/NEJMoa2001017 (2020).

4    Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**,
270-273, doi:10.1038/s41586-020-2012-7 (2020).

5    Chen, L. *et al.* RNA based mNGS approach identifies a novel human coronavirus from two individual pneumonia
cases in 2019 Wuhan outbreak. *Emerging microbes & infections* **9**, 313-319,
doi:10.1080/22221751.2020.1725399 (2020).

6    Korber, B. *et al.* Tracking changes in SARS-CoV-2 Spike: evidence that D614G increases infectivity of the
COVID-19 virus. *Cell*, doi:doi.org/10.1016/j.cell.2020.06.043 (2020).

7    Smith, E. C., Blanc, H., Vignuzzi, M. & Denison, M. R. Coronaviruses Lacking Exoribonuclease Activity Are
Susceptible to Lethal Mutagenesis: Evidence for Proofreading and Potential Therapeutics. *Plos Pathog* **9** (2013).

8    Sevajol, M., Subissi, L., Decroly, E., Canard, B. & Imbert, I. Insights into RNA synthesis, capping, and
proofreading mechanisms of SARS-coronavirus. *Virus Res* **194**, 90-99 (2014).

9    Vignuzzi, M., Stone, J. K., Arnold, J. J., Cameron, C. E. & Andino, R. J. N. Quasispecies diversity determines
pathogenesis through cooperative interactions in a viral population.  **439**, 344-348 (2006).

10    Garvin, M. R. *et al.* Potentially adaptive SARS-CoV-2 mutations discovered with novel spatiotemporal and
explainable AI models. *Genome Biology* **21**, 304, doi:10.1186/s13059-020-02191-0 (2020).

11    Tang, X. *et al.* On the origin and continuing evolution of SARS-CoV-2. *National Science Review* **7**, 1012-1023,
doi:10.1093/nsr/nwaa036 %J National Science Review (2020).

12    Acter, T. *et al.* Evolution of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) as coronavirus
disease 2019 (COVID-19) pandemic: A global health emergency. *Science of The Total Environment* **730**, 138996,

doi:https://doi.org/10.1016/j.scitotenv.2020.138996 (2020).

13    Plante, J. A. *et al.* Spike mutation D614G alters SARS-CoV-2 fitness. *Nature*, doi:10.1038/s41586-020-2895-3
(2020).

14    Zohar, T. *et al.* Compromised Humoral Functional Evolution Tracks with SARS-CoV-2 Mortality. *Cell* **183**,
1508-1519.e1512, doi:10.1016/j.cell.2020.10.052 (2020).

15    Li, T. *et al.* The use of SARS-CoV-2-related coronaviruses from bats and pangolins to polarize mutations in
SARS-Cov-2. *Science China Life Sciences* **63**, 1608-1611, doi:10.1007/s11427-020-1764-2 (2020).

16    Grubaugh, N. D., Hanage, W. P. & Rasmussen, A. L. Making sense of mutation: what D614G means for the
COVID-19 pandemic remains unclear. *Cell*, doi:doi.org/10.1016/j.cell.2020.06.040 (2020).

17    Liu, Z. *et al.* Identification of Common Deletions in the Spike Protein of Severe Acute Respiratory Syndrome
Coronavirus 2. *J Virol* **94** (2020).

18    Li, Q. Q. *et al.* The Impact of Mutations in SARS-CoV-2 Spike on Viral Infectivity and Antigenicity. *Cell* **182**,
1284-+ (2020).

19    Plante, J. A. *et al.* Spike mutation D614G alters SARS-CoV-2 fitness. *Nature* (2020).

20    Blanco, J. D., Hernandez-Alias, X., Cianferoni, D. & Serrano, L. In silico mutagenesis of human ACE2 with S
protein and translational efficiency explain SARS-CoV-2 infectivity in different species. *Plos Comput Biol* **16**
(2020).

21    Zhang, L. Z. *et al.* SARS-CoV-2 spike-protein D614G mutation increases virion spike density and infectivity. *Nat*
*Commun* **11** (2020).

22    A., B. *et al.* Antibody cocktail to SARS-CoV-2 spike protein prevents rapid mutational escape seen with
individual antibodies. *Science*, doi:10.1126/science.abd0831 (2020).

23    Hansen, J. *et al.* Studies in humanized mice and convalescent humans yield a SARS-CoV-2 antibody cocktail.
*Science*, doi:10.1126/science.abd0827 (2020).

24    Sheahan, T. P. *et al.* An orally bioavailable broad-spectrum antiviral inhibits SARS-CoV-2 in human airway
epithelial cell cultures and multiple coronaviruses in mice. *Sci Transl Med* **12** (2020).

25    Nunes-Santos, C. J., Kuehn, H. S. & Rosenzweig, S. D. N-Glycan Modification in Covid-19 Pathophysiology: In
vitro Structural Changes with Limited Functional Effects. *J Clin Immunol* (2020).

26    Lo, M. K. *et al.* Remdesivir targets a structurally analogous region of the Ebola virus and SARS-CoV-2
polymerases. *P Natl Acad Sci USA* **117**, 26946-26954 (2020).

27    Editorial. Evolution goes viral. *Nature Ecology & Evolution* **5**, 143-143, doi:10.1038/s41559-021-01395-2 (2021).

28    Editorial. COVID-19 vaccines: acting on the evidence. *Nature Medicine*, doi:10.1038/s41591-021-01261-5
(2021).

29    Muik, A. *et al.* Neutralization of SARS-CoV-2 lineage B.1.1.7 pseudovirus by BNT162b2 vaccine–elicited human
sera. eabg6105, doi:10.1126/science.abg6105 %J Science (2021).

30    Rice, B. L. *et al.* Variation in SARS-CoV-2 outbreaks across sub-Saharan Africa. *Nature Medicine*,

doi:10.1038/s41591-021-01234-8 (2021).

31   Muik, A. *et al.* Neutralization of SARS-CoV-2 lineage B.1.1.7 pseudovirus by BNT162b2 vaccine–elicited human
sera. *Science*, eabg6105, doi:10.1126/science.abg6105 %J Science (2021).

32   Zhou, P. & Shi, Z.-L. SARS-CoV-2 spillover events. *Science* **371**, 120-122, doi:10.1126/science.abf6097 %J Science
(2021).

33   Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular Evolutionary Genetics Analysis across
Computing Platforms. *Molecular Biology and Evolution* **35**, 1547-1549 (2018).

34   Tamura, K. & Nei, M. Estimation of the Number of Nucleotide Substitutions in the Control Region of
Mitochondrial-DNA in Humans and Chimpanzees. *Molecular Biology and Evolution* **10**, 512-526 (1993).

35   Li, Q. *et al.* The impact of mutations in SARS-CoV-2 spike on viral infectivity and antigenicity. *Cell*,
doi:doi.org/10.1016/j.cell.2020.07.012 (2020).

36   Zhang, Y., Song, W., Chen, S., Yuan, Z. & Yi, Z. A bacterial artificial chromosome (BAC)-vectored noninfectious
replicon of SARS-CoV-2. *bioRxiv*, doi:doi.org/10.1101/2020.09.11.294330 (2020).

37   Shi, J. *et al.* Susceptibility of ferrets, cats, dogs, and other domesticated animals to SARS–coronavirus 2. *Science*
**368**, 1016-1020, doi:10.1126/science.abb7015 %J Science (2020).

38   Wu, L. *et al.* Broad host range of SARS-CoV-2 and the molecular basis for SARS-CoV-2 binding to cat ACE2. *Cell*
*Discovery* **6**, 68, doi:10.1038/s41421-020-00210-9 (2020).

39   Patterson, E. I. *et al.* Evidence of exposure to SARS-CoV-2 in cats and dogs from households in Italy. *Nat*
*Commun* **11**, 6231, doi:10.1038/s41467-020-20097-0 (2020).

40   Ferasin, L. *et al.* Myocarditis in naturally infected pets with the British variant of COVID-19. *bioRxiv*,
2021.2003.2018.435945, doi:10.1101/2021.03.18.435945 %J bioRxiv (2021).

41   Raghav, S. *et al.* Analysis of Indian SARS-CoV-2 Genomes Reveals Prevalence of D614G Mutation in Spike
Protein Predicting an Increase in Interaction With TMPRSS2 and Virus Infectivity. *Front Microbiol* **11** (2020).

42   Johnson, M. C. *et al.* Optimized Pseudotyping Conditions for the SARS-COV-2 Spike Glycoprotein. *J Virol* **94**
(2020).

43   Jiang, X. Y. *et al.* Bimodular effects of D614G mutation on the spike glycoprotein of SARS-CoV-2 enhance
protein processing, membrane fusion, and viral infectivity. *Signal Transduct Tar* **5** (2020).

44   Fernandez, A. Structural Impact of Mutation D614G in SARS-CoV-2 Spike Protein: Enhanced Infectivity and
Therapeutic Opportunity. *Acs Medicinal Chemistry Letters* **11**, 1667-1670 (2020).

45   Ristad, E. S., Yianilos, P. N. J. I. T. o. P. A. & Intelligence, M. Learning string-edit distance. *IEEE Transactions on*
*Pattern Analysis and Machine Intelligence* **20**, 522-532 (1998).

46   Bille, P. J. T. c. s. A survey on tree edit distance and related problems. *Pattern Analysis and applications* **337**,
217-239 (2005).

## Figures and Figure legends

**Fig. 1 Evolutionary trajectories of variant B.1.1.7. a** Incremental mutations of variant B.1.1.7. The digits in the upper-right-corner rectangle with dotted line indicate the labels of mutations. For simplicity, the 69–70 deletions were labeled as "1", and the other mutations "2"-"9" respectively. The bottom nodes (rectangles) represent the variants with one mutation and the top one was the early variant B.1.1.7. Each rectangle with solid line consists of lineage (e.g., B.1.243), number of strains (e.g., N:2382), mutation sites (e.g., M:----56---), the earliest collection date (e.g., 20-03-29, i.e., 2020-03-29), and collection location (e.g., USA). In the labels of the mutation sites, sign "-" indicated the corresponding site did not mutate. All routes along the directions of the arrows are possible evolutionary trajectories of lineage B.1.1.7, where the green one was the most probable mutation trajectory. Large-scale SARS-CoV-2 analysis demonstrates that the early variant B.1.1.7 might not have arisen spontaneously in the UK or within human hosts. **b** Coappearances of variant B.1.1.7 mutations. At least five mutations form a potential co-mutation pattern (coappearance rate > 95%). **c** Possible transmission chains of variant B.1.1.7. Canidae, Mustelidae or Felidae, especially the Canidae family (for example, dog) could be a possible host of the direct progenitor of variant B.1.1.7.

**Fig. 2 The Canidae family could be a possible host of the direct progenitor of variant B.1.1.7.** The digits on the left of the figure indicate the labels of mutations, which correspond with the mutation labels in Fig. 1a. The strains shown in the center of the figure contain at least one spike mutation of variant B.1.1.7. And these strain examples cover all existing SARS-CoV-2 viruses that collected from animal hosts. The strain labeled by orange star corresponds with the star variant in Fig. 1a. The strain with orange solid-round label was collected from a dog on 2020-07-28. Such two strains share the same mutations "56" and have the minimum phylogenetic distance by MEGA tool. Canidae, Mustelidae or Felidae, especially the Canidae family (for example, dog) could be a possible host of the direct progenitor of variant B.1.1.7 based on existing stains collected before the end of Jan. 2021.

**Fig. 3 The strong correlations suggest that the top 25 mutations form eight potential co-mutation patterns. a** The correlation heatmap of the top 25 mutations. These mutations could be grouped into several clusters with high Pearson-correlation-coefficient (PCC). **b** Regression analysis of mutations shows that eight clusters all denote the statistical significance level: ***p value < E-44. c to k show the transitions of the high-frequency mutations. The sky-blue represents the rate per day of initial residue in population and the golden the rate per day of substitution/mutant. These mutation transitions provide further evidence that the above mutations potentially form co-mutation patterns.

**Fig. 4 Dominant co-mutation attenuates viral replication. a** Predicted RNA structure of the SARS-CoV-2 5'UTR. RNA structure of the 400-nt 5'UTR was predicted by "RNAstructure" (http://rna.urmc.rochester.edu/RNAstructureWeb). The start codon for nsp1 is grey, the TRS-L is orange, and the mutated nucleotides are red. The bottom panel shows the alignment of the 5'UTR of SARS-CoV-2 with 5'UTRs of related viruses, with c241 highlighted. **b** Structure of SARS-CoV-2 RdRp/RNA complex. The structure of SARS-CoV-2 RdRp/RNA complex (PDB, 6X2G) was visualized by Chimera (UCSF). The P323 mutation is highlighted in red, with the alignment of the amino acid sequences of SARS-CoV-2 and related viruses near the P323 position. **c** Schematic of the *in-vitro* ligation system for SARS-CoV-2 replicon. Four plasmids encompassing the viral genome were digested by BsaI to release the four fragments. After gel purification, the fragments were ligated by T4 ligase. The ligation products were purified and used as template for RNA *in-vitro* transcription. sGluc, secreted *Gaussia* luciferase; 2A, foot-and-mouth disease virus (FMDV) 2A peptide; BSD, blasticidin. **d** Huh7 cells were co-transfected with *in-vitro* transcribed replicon RNA (WT or the indicated mutants) and an mRNA encoding the SARS-CoV-2 N protein. The luciferase activity in the supernatants was measured at the time points indicated. Medium was changed at 8 hours post-transfection. Data are shown as mean±SEM (n=8). SAA, the NSP12 polymerase active-site mutant. Unpaired Student's t-test was performed between the mutants and wild type (WT) and between the mutants as indicated (statistical significance level: *p value<0.05, **p value<0.01, ***p value<0.001).